3D Mapping Reveals Network-specific Amyloid Progression and Subcortical Susceptibility

Canter RG*[1], Choi H*[4], Wang J[1], Watson LA[1], Yao CG[1], Abdurrob F[1], Bousleiman SM[1], Delalle I[2], Chung K[+1,3,4], Tsai L-H[+1]

*These authors contributed equally to this work.
+Co-corresponding authors.

[1]The Picower Institute for Learning and Memory, Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology. Cambridge, MA
[2]Department of Pathology and Laboratory Medicine, Boston University School of Medicine, Boston, MA
[3]Department of Chemical Engineering, MIT, Cambridge, MA
[4]Institute for Medial Engineering and Science (IMES), MIT, Cambridge, MA

Abstract
Alzheimer's disease is a progressive, neurodegenerative condition for which there is no cure. Prominent hypotheses posit that accumulation of beta-amyloid (Aβ) peptides drives the neurodegeneration that underlies memory loss, however the spatial origins of the lesions remain elusive. Using SWITCH, we created a spatiotemporal map of Aβ deposition in a mouse model of amyloidosis. We report that structures connected by the fornix show primary susceptibility to Aβ accumulation and demonstrate that aggregates develop in increasingly complex networks with age. Notably, the densest early Aβ aggregates occur in the mammillary body coincident with electrophysiological alterations. In later stages, the fornix itself also develops overt Aβ burden. Finally, we confirm Aβ in the mammillary body of postmortem patient specimens. Together, our data suggest that subcortical memory structures are particularly vulnerable to Aβ deposition and that functional alterations within and physical propagation from these regions may underlie the affliction of increasingly complex networks.

Author Contributions:
RGC, KC, L-HT, ID conceived of the work and planned the experiments.
RGC, HC, JW, LAW, CGY, FA, SMB performed experiments and analyzed data.
HC built the custom microscope.
RGC, L-HT, KC, ID wrote the manuscript.

Introduction

Cognitive impairments attributable to Alzheimer's disease (AD) will affect millions of individuals in the next decade, however the etiology of the disease remains unknown.[1] The results from decades of research support the hypothesis that the accumulation of toxic amyloid-beta peptides (Aβ) in the brain contributes to the onset and progression of AD.[2,3] The hypothesis was initially based on the discovery that mutations in the Aβ precursor protein (APP) and its processing enzymes cause autosomal dominant, inherited, familial AD (FAD).[4] Subsequent preclinical studies demonstrating that the Aβ peptides resulting from these mutations induces synaptic loss and neuronal death *in vitro* and *in vivo* have further established the acute toxicity of aggregate forms of the peptide on neural substrate, which suggests they directly influence the neurodegeneration observed in different subtypes of AD.[5–7] In addition to cellular harm, amyloidosis in murine models of AD contributes to AD-like memory impairments, hippocampal synaptic loss, and electrophysiological alterations -- changes that are also observed in human patients.[8–10] This data not only links lesions to dementia in FAD, but also provides a basis for understanding the etiology of the sporadic form of the disease. Despite the link between amyloid toxicity and neurodegeneration, the precipitating events that trigger Aβ accumulation, deposition, and propagation which initiate AD remain unclear.

One impediment to understanding Aβ deposition has been identifying the brain regions most vulnerable to Aβ plaques. The first studies using postmortem brain sections from AD patients' brains suggested that primary accumulation occurs in the neocortex with subsequent spread of aggregates to deeper structures implicated in learning and memory.[11] However, due to the use of post-mortem specimens, these reports could not definitively link evidence of amyloid deposition to the eventual development of AD. Although recent advances in positron emission tomography (PET) imaging have enabled longitudinal patient imaging studies that confirm the importance of cortical Aβ in the diagnosis and prediction of AD progression,[12,13] they have largely not reported the contributions of deeper structures to Aβ load and spread.[13,14] Intriguingly, while cortical Aβ does correlate with disease status in patients,[15,16] many cognitively healthy individuals also have high levels of cortical Aβ.[17] The discrepancy between cortical accumulation and cognitive impairment suggests that unexplored Aβ-induced changes in the brain contribute to memory-related alterations in AD.[18]

Increasing evidence suggests that distributed memory networks are particularly affected by Aβ in AD. Some lines of investigation demonstrate alterations in high-level cortical networks like the

default mode network (DMN),[14,19] while other recent studies show core alterations in
fundamental deep memory structures that are part of the limbic system.[10,20–22] Though the
cognitive and memory networks implicated by patient imaging somewhat overlap, there are
distinct brain areas and functions that the networks do not share, leaving the hypotheses at
odds. Whether the divergence arises from technical considerations or has unaddressed
biological significance remains unknown, and still unexplored is the question of whether specific
areas show differential susceptibility that is not seen with bulk detection. We hypothesized that
some of the discrepancies in observed network susceptibility may be due to the difficulties in
staging patient AD progression, which can be affected by variable pathological load,[18] as well as
by socioeconomic, lifestyle, and genetic factors.[23]

To overcome the variability in human disease, in this work, we capitalize on the temporal
precision of murine models harboring FAD mutations to map amyloid progression at high-
resolution throughout the brain.[24,25] We use optimized techniques for whole-brain SWITCH
immunolabeling[26] to create an unbiased, spatiotemporal map of Aβ deposition. Our
observations reveal early subcortical susceptibility to amyloidosis in regions connected by the
fornix, such as the mammillary body, which shows concurrent functional alterations alongside
aggregate formation. The fur-dimensional (4D) nature of the data also reveal potential
mechanisms of propagation along white matter tracts and suggests that pathological changes
within long-range projections that connect memory and cognitive networks may contribute to
disease progression.

Results
Using optimized system-wide control of interaction time and kinetics of chemicals (SWITCH)
techniques (Figure 1a) that enable homogenous, whole-brain immunolabeling,[26] we created a
spatially unbiased, temporally precise map of amyloidosis in a murine model of AD. We chose
the 5XFAD transgenic mouse line because it is a widely-used model of AD-like pathology that
shows an age-dependent increase in Aβ aggregates and alterations in hippocampal
physiology.[8,25] Although the SWITCH techniques are widely applicable to many antibodies and
target proteins,[26] we discovered that we needed to further refine the buffer systems to achieve
homogenous whole brain labeling with our chosen amyloid antibody. In our optimized SWITCH
system, we switch pH and ionic strength of the buffers to actively modulate antibody-antigen
binding kinetics [27] In the first step, we use high pH and high ionic strength buffer to slow the
binding reaction so that antibodies can penetrate deep into the sample. Next, the buffer is

titrated to a physiological pH and ionic strength to allow antibodies to rapidly bind to targets. By

employing this system, we were able to achieve homogenous labeling throughout thick tissue

sections (Supplemental Figure 1). To further optimize the conditions for our chosen amyloid

antibody, we used aged animals with the expectation that they would show wide-spread

aggregate accumulation based on previous reports.[25,28] Using the optimized SWITCH protocols

to label a whole brain, we observed homogenous immunolabeling of Aβ aggregates throughout

the entire tissue in aged mice (Supplemental Figure 2, Video 1). These large tissue volume

techniques allow analyses of intact specimens without the loss of critical information to the size

limitations and directional segmentation that are inherent in sectioning.[29] To our surprise, careful

examination of the entire brain revealed that, in addition to deposition in cortical and

hippocampal regions known to be affected in AD, significant accumulation of Aβ was also

evident in subcortical regions.

To determine the reproducibility of the protocols and replicate the unexpected labeling, we

repeated the procedure in a cohort of 5XFAD animals aged 12 month (12M) (Figure 1B, Video

2d). In the 12M animals, we observed substantial plaque burden throughout the brain. Region

specific quantification from white matter-based hand annotation of three-dimensional (3D)

images (Supplemental Figure 3) demonstrated area-specific accumulation in AD-associated

areas (Figure 1C). The regions harboring Aβ included the retrosplenial cortex (RSP),

hippocampus (HPC), subiculum (SUB), and pre-frontal cortex (PFC), and importantly,

aggregates were largely absent from areas thought to be relatively spared from amyloidosis in

AD like the caudoputamen (CP) and midbrain (MD) (D'Agostino and Pearson normality test; K2

= 94.61, p < 0.0001, passed normality = No; Friedman test; Q = 114.2, p < 0.0001). This

suggests that our labeling techniques are able to homogenously and specifically label Aβ

aggregates, and that accumulation of the peptide is restricted within particular brain areas.

Unexpectedly, the highest aggregate density by rank in the 12M animals appeared within

specific subcortical regions, namely the mammillary body (MB) and septum (SEPT), which

confirmed our initial observations.

Although the MB and SEPT had the densest Aβ, with data from aged animals we could not

determine whether this pattern emerged because aggregates in the 5XFAD animals develop

uniformly throughout the brain over time, thus making these smaller structures appear more

dense; or if these high-density regions show unique vulnerability to elevated levels of Aβ,

leading to earlier or more significant aggregation. To observe how the pattern emerged, we

used our optimized SWITCH protocol to analyze a time course of 5XFAD cohorts including mice

aged for 6, 4, 2, and 1 months (6M, 4M, 2M, 1M respectively) (Video 2a-c, Figures 1D-G).

Quantifying Aβ density in specific regions across multiple animals at each time point, we

observed increasingly precise aggregation with decreasing age in a pattern that was consistent

across individuals. At 6M, 5XFAD animals had region-specific aggregation, with areas harboring

significantly different densities of amyloid. (Figures 1D, 1G; D'Agostino and Pearson normality

test; K2 = 70.12, p < 0.0001, passed normality = No; Friedman test; Q = 68.56, p = 0.0171). To

better understand the brain areas that contribute to region-specific aggregation in our data, we

performed hierarchical clustering analysis, which revealed that at 6M, areas associated with the

olfactory and cortico-limbic system (e.g. basolateral amygdalar complex [BLA], piriform cortex

[PIR], entorhinal cortex [EC]) show most similar levels of Aβ (Supplementary Figure 4B;

hierarchical clustering, single linkage- nearest distance). This is particularly interesting because

it suggests that network-level distribution of Aβ is relatively uniform within functional networks

and also identifies circuits that underlie behaviors known to show alterations in early AD.[30]

Looking at younger animals to determine the progression in the predictable, genetic murine

models, we next saw that at 4M the data continues to demonstrate area-specific aggregation

and our analyses show non-normal, non-linear pattern to the regional distribution (Figure 1E,

1G; D'Agostino and Pearson normality test; K2 = 51.67, p < 0.0001, passed normality = No;

Non-linear Regression; y = -354.1*ln(x)+1200.9, R^2 = 0.82465). The skewed data suggest that

fewer regions have high burden, consistent with increasingly specific patterns of aggregation at

the earliest disease stages. To determine what regions are affected at this 4M age, we looked at

our clustering analyses (Supplementary Figure 4C), which revealed that regions homologous to

the those in the DMN (e.g. restroplenial [RDP], anterior cingulate [ACC], and parietal cortices)[31]

are better correlated to each other, with the exclusion of olfactory and limbic areas from the

cluster. To understand whether the DMN is the earliest affected network as suggested by many

human functional studies,[32,33] we next looked at 2M animals. At 2M, animals failed to

demonstrate significant differences across brain areas (Figure 1F; D'Agostino and Pearson

normality test; K2 = 105.2, p < 0.0001, passed normality = No; Friedman test; Q = 58.18, p =

0.1073), suggesting little to no specific aggregation patterns. However, our analyses revealed

that while many areas did not cluster, there was correlated deposition in the 2M animals within a

few areas (Supplementary Figure 4D). These key nodes of early aggregation were the MB,

SEPT, and SUB – regions that connect the HPC to the rest of the Papez memory circuit.[34] At

1M animals did not show discernable amyloid deposits (data not shown), suggesting that the

Figure 1

Figure 1: SWITCH labeling reveals network propagation of amyloid pathology. A) Diagram of protocol for our optimized SWITCH labeling. B) Representative images of the 12 month 5XFAD brains. C) Amyloid deposit density rank-ordered by region in 12 month aged 5XFAD mice. D-F) Representative images of amyloid labeling from D) 6 month, E) 4 month, and F) 2 month 5XFAD mice. G) Clustergram of the hierarchical clustering by density across the four time points examined. Coloring of the dendrogram represents the fifth level from the top and these clades contain regions with similar amyloid deposition patterns. H) Lists of regions contained in colored groups from panel G. *p ≤ 0.05; **p ≤ 0.01; ****p≤ 0.0001; Graphs report Mean ± Standard Deviation unless otherwise noted in the legend text. *Abbreviations: ACC, anterior cingulate cortex; RSP, retrosplenial cortex; vSub, ventral subiculu; LS, lateral septum; dSub, dorsal subiculum; MC, motor cortex; DB, diagonal band; MS, medial septum; ACB, nucleus accumbens; PIR, piriform cortex; MD, midbrain; Thal, thalamus; BLA, basolateral amygdala; OT, olfactory tubercle; PERI, perirhinal and ectorhinal corticies; GC, gustatory cortex; IC, insular cortex; VIS, visual cortex; PTL, posterial parietal association areas; SS, somatosensory cortex; DP, dorsal peduncular area; AON, anterior olfactory nucleus; EC, entorhinal cortex; PFC, prefrontal cortex; OFC, orbitofrontal cortex; TR, postpiriform transition area; TT, taenia tecta; dDG, dorsal dentate gyrus; Visc, visceral cortex; TEA, temporal association areas; AUD, auditory cortex; dCA1, dorsal CA1, vDG, ventral dentate gyrus; vCA1, ventral CA1; vCA3, ventral CA3; dCA3, dorsal CA3; CP, caudoputamen; BNST, bed nucleus of the stria terminalis; HYP, hypothalamus; CeA, centromedial amygdalar nuclei; SI, substantia innominata; PAA, piriform amygdalar area; COA, cortical amygdalar area.*

deposition is not developmental and that the SUB, MB, and SEPT are uniquely susceptible to

Aβ aggregation at the earliest stages of disease pathology.

One concern using transgenic models is whether the pattern is reflective of biologically

interesting results, or a byproduct of the transgenic expression. To ascertain whether the

subcortical aggregation pattern may be an artifact of the mouse transgene, we performed dual

immunofluorescence for Aβ peptides and *in situ* hybridization for the transgenic mRNA to

correlate the location of Aβ deposits with expression patterns of the transgene (Supplementary

Figure 5A). Patterns of expression in 2 and 4-month animals looked similar and were combined,

and the correlation analyses demonstrate that across the entire brain, transgenic RNA

expression does not correlate with deposit location (Supplementary Figure 5B; ρ = -0.0728, p =

0.3588). The insignificance of this result remains when split by region (Supplementary Figure

5C). This data suggests that 5XFAD animals show biologically relevant, region-specific amyloid

deposition. Combined with our clustering results (Figure 1H), our data show that the MB, SEPT,

and SUB are particularly vulnerable to Aβ deposition. Over time, the aggregates appear in

regions homologous to the DMN. Following the DMN, lesions are detected throughout the

extended limbic system, before finally developing throughout the entire forebrain. Thus we

propose that Aβ deposits start in susceptible subcortical structures, and spread to increasingly

complex memory and cognitive networks with age.

The detection of aggregates alone is not sufficient to demonstrate that the Aβ we observe in the

MB, SEPT, and SUB may contribute to Aβ-related cognitive decline because it is unknown

whether subcortical Aβ, especially within diencephalic structures, induces the functional

alterations thought to underlie early memory deficits in AD.[9,22,35,36] In our data, the MB shows the

densest Aβ at the earliest stages. The MB is downstream of both the SEPT and SUB,[37] and

previous reports suggest that it is related to recall memory in humans[38] and has calcium-

sensitive firing properties[39] that may render it particularly vulnerable to Aβ-induced dysfunction.

Thus, to test whether the observed aggregation leads to functional impairments within the

subcortical structures, we performed *ex vivo* whole cell patch-clamp recordings in the MB in

slices from 2M 5XFAD mice and littermate controls. 5XFAD MB neurons showed significantly

increased excitability (Figures 2A-B, 2-way Repeated Measures ANOVA; Interaction: $F_{(10,450)}$=

8.786, p < 0.0001; Genotype: $F_{(1,45)}$= 9.902, p = 0.0029), which is consistent with previous

reports of neuronal hyperexcitability in the hippocampus in response to Aβ.[9,35] Further

examination showed that this excitability was not due to increased action potential threshold

Figure 2.

Figure 2: **The MB shows significant functional impairments** and loses its synaptic input. A) Representative action potential traces following current injection of 10pA (top row) and 100pA (bottom row) into MB neurons of 5XFAD animals (right column) compared to littermate controls (left column). B) Excitability curves showing increased firing rate of FAD+ across multiple steps of current injection. C) The action potential threshold of the cells does not change, neither does the D) after-hyperpolarization. E) The action potential amplitude is significantly decreased in 5XFAD animals and F) the resting membrane potential is also significantly decreased, both of which could underlie the excitability phenotype. Graphs represent Mean ± Standard Deviation.

(Figure 2C, Unpaired Student's t-test; $t_{(46)}$ = 1.146, p = 0.2577) or altered hyperpolarlization

dynamics (Figure 2D, Unpaired Student's t-test; $t_{(32)}$ = 0.7107, p = 0.4824), but may be attributed

to slightly increased action potential amplitude (Figure 2E, D'Agostino and Pearson Normality

Test, FAD-: K2 = 11.09, p = 0.0039, passed normality = No; FAD+: K2 = 4.217, p = .1214,

passed normality = Yes; Mann Whitney U; U = 87.5, p = 0.0497) and a reduced resting

membrane potential (Figure 2F, Unpaired Student's t-test; $t_{(40)}$ = 4.807, p < 0.0001) in 5XFAD

MB neurons compared to the WT controls. This is consistent with previous findings of altered

resting membrane potential in non-pyramidal neurons of the HPC in a different murine model of

AD,[40] revealing further consistencies in amyloid-driven alterations, despite differing cell-types

and circuit structures.[41] Thus, our results suggest that early accumulation of Aβ the MB of the

5XFAD drives neuron-intrinsic hyperexcitability, which is characteristic of AD-related changes in

neuronal activity and, in this early stage, may contribute to the onset of network dysfunction and

memory impairment.[9,35,40]

Our results demonstrating that the MB is particularly susceptible to early Aβ deposition adds to

the mounting evidence that implicates subcortical Papez circuitry in AD-related memory

impairment in human patients,[21,42–44] however the relevance of these findings is tempered by

mixed reports about the involvement of MB in human mild cognitive impairment (MCI) and AD

patients.[42,45,46] Thus, to understand whether MB demonstrates Aβ lesions over the course of AD,

we acquired a small cohort of post-mortem mammillary bodies embedded in paraffin blocks

from the Netherlands Brain Bank. The blocks contained mammillary bodies from individuals that

died at each Braak and Braak stage[47] (Figure 3A). We cut traditional histology sections from the

blocks (Figure 3B) that revealed numerous Aβ deposits in the MB of AD patients at each stage,

with none present in the healthy individual (Figure 3C, 3D). This finding is consistent with older

reports of amyloid pathology in the region.[43] However, because the sections were thin and the

deposits did not label strongly, we could not determine whether there was a stage-dependent

increase in Aβ within the MB. To test this, we used iSWITCH methods to clear and label the

entire banked MB specimen from each individual. 3D reconstruction of the labeled specimens

(Figure 3E, Video 3) revealed substantial Aβ burden in the MB of AD patients, but none in the

MB from a healthy individual. Although the density of Aβ deposits did not correlate with NFT

accumulation reflected in advancing Braak stage (Figure 3F), individual aggregate volume did

(Figure 3G). This suggests substantial early Aβ deposition precedes overt dementia and that

aggregates continue to expand as the largest ones in our small cohort were seen in the Braak V

– AD patient. One additional interesting observation in the pathology sections was that there

Figure 3

A

Patient	Gender	Age	Braak Score	Diagnosis
1	male	70	0	Not Impaired
2	male	80	2	Not Impaired
3	female	91	3	Not Impaired
4	male	74	5	AD

B 5X Overview   C 40X   D Histology Count

E BB-0 BB-II BB-III BB-IV

F Plaque Number   G Mean Volume   H.

Figure 3: **Human mammillary body and associated white matter show AD-related amyloid burden**. A) Information on the patient samples used for the experiments in this figure. B) Representative images of 5X tile scans showing the overall MB structure and C) representative high-magnification image of the plaque types used for quantification (right). D) Quantification of histologically identified plaques from pathology slide sections. E) Representative images of the large-volume MB samples labeled with D54D2. F) The plaque count does not reflect the severity of the disease, however the G) mean plaque volumes increase with disease severity in this small cohort. H) White matter tracts identified anatomically and pathologically by presence of oligodendrocyte nuclei show amyloid burden too.

were numerous Aβ deposits in oligodendrocyte-rich regions of the MB (Figure 3H), suggesting a

relationship between white matter tracts and Aβ deposition in this region.

To check whether this is reflected in the 5XFAD model, we investigated the Aβ burden in major

forebrain white matter tracts. We did not observe Aβ in the white matter of 2M animals, but did

observe consistent deposits in the fornix of the 6M cohort (Figure 4A, Video 2c). The fornix,

which connects the SUB to the SEPT and MB, is the major output structure of the hippocampal

formation. To determine whether this burden was unique to the fornix, we took 100um coronal

sections from 6M 5XFAD brains (Figure 4B). Surprisingly, white matter tracts connecting major

limbic structures showed significantly more accumulation than tracts connecting other regions

(Figure 4C, Friedman Test, Q = 40.29, p <.0001), suggesting that the limbic white matter is

uniquely vulnerable. To determine whether these aggregates formed along the outside

fasciculated bundles, we performed higher-magnification imaging of the 100um sections (Figure

4D, Video 4). Interestingly, the 3D reconstruction demonstrates amyloid build-up within the

myelin sheath, along the axonal path (Figure 4D, Z-plane). This pattern was specific to the

limbic structures like the fornix, fimbria, and cingulum bundle (Figure 4E), excluded white matter

tracts like the corpus callosum (Figure 4E) and was not present in the wild-type littermate

control animals (Figure 4F). This data is particularly intriguing because in the 6M FAD there is

considerable Aβ burden in the cortical regions connected by the corpus callosum (Figure 1D).

Together, it suggests that limbic white matter tracts are uniquely affected by axonal Aβ in AD,

and may explain structural alterations in these regions in human MRI[21] and the progression of

connectivity impairments in individuals at risk or harboring significant brain Aβ.[48–50]

Discussion

Using optimized iSWITCH whole-brain clearing and immunolabeling technologies, we have

created a spatially unbiased map of the progression of Aβ deposits that revealed novel

subcortical hubs of early-disease susceptibility in a mouse model of AD. Our findings

demonstrate that areas connected by the fornix develop aggregates at the earliest phases of

disease and that, of the identified vulnerable regions, the MB shows the most significant density

of deposits across time. Although the MB is a critical part of the Papez long-term memory circuit

connecting the hippocampus to the anterior thalamus,[34,51] it has not been strongly implicated in

AD.[34,42,52,53] This is somewhat surprising because its major inputs, the subiculum and prefrontal

cortex[54], have been shown to demonstrate significant synaptic loss that correlates with memory

performance.[55–58] Furthermore, the MB is part of the head-direction system, a group of critical

navigation regions including hippocampus, thalamus, and retrosplenial cortex that are affected

in AD, and the pathology in which is thought to underlie spatial memory deficits seen in the

disease.[59,60] Integrating our findings into this framework, the present data indicate that the

subcortical and diencephalic projecting components of the Papez circuit are particularly

vulnerable in AD. Interestingly, some of the regions of the Papez circuit and head direction

systems, like the septum, were initially implicated in early AD by the cholinergic hypothesis,

which posited that loss of those neurons was causative in disease onset and progression[61].

Although the idea was overshadowed by the amyloid cascade hypothesis,[2] our data, along with

other recent evidence for Papez circuit dysfunction[50] suggests that these subcortical structures

and their white matter tracts are associated with amyloid pathology early in AD.[62] In the present

manuscript, we take this observation one step further to show that the onset of Aβ aggregation

in the MB of the 5XFAD animals correlates with functional alterations. These data are consistent

with network-level changes previously observed in the Papez circuit in AD patients and other

amyloid models,[9,22,36,40] and the significant findings at such an early pathological stage highlight

the importance of continued investigation of the network alterations and functional changes in

these structures at the initial disease phases. Because the regions in the basal forebrain are

linked in our data, and other studies, to the hypothalamic diencephalic limbic structures, we

suggest in-depth studies of these regions and their connections may lead to a better

understanding of the mechanisms of dysfunction underlying AD onset and the ensuing

progressive memory loss.

It has become increasingly important to understand the translatability of observations in animal

models.[63,64] Thus to determine whether the MB is also affected early in

human AD, we used SWITCH techniques to confirm that Aβ aggregates are present in

postmortem MB from Braak staged individuals with tau pathology and an AD patient. Our data

investigating both traditional histology sections alongside large-volume labeled samples show

significant Aβ burden that increases with stage, which is most evident using our novel

technique. The data we obtained with traditional labeling methods was consistent with the

literature[65], however the large-volume images revealed that the histology sections did not show

the full burden in the region. This observation highlights the power of these tools to enable a

better understanding of pathological affliction through molecular interrogations of intact,

precious samples. These observations of MB lesion and dysfunction in AD are of particular note

because of the amnesia associated with other MB-related conditions. For example in

Wernicke's encephalopathy and Korsakoff's syndrome, MB structural alterations were one of

Figure 4

Figure 4: White matter amyloid deposits are specific to limbic tracts. Six month 5XFAD animals develop amyloid deposits within the white matter. A) Representative images of amyloid within the fornix visualized in the 3D rendered whole brain amyloid labeling in the six month 5XFAD cohort. B) Representative image of the white matter amyloid labeling in 100um sections from a second cohort of six month 5XFAD animals. C) Quantification of white matter deposits within several posterior white matter tracts. D) Representative images demonstrating labeling of amyloid within white matter bundles in the fornix. E) Representative images showing robust amyloid deposition within the fimbria, cingulum, and alveus, and less in the corpus callosum. F) There is no amyloid labeling within the fornix of non-transgenic 6 month aged littermates. *p ≤ 0.05; ****p ≤ 0.0001; Green and white arrows point to white matter regions with observed amyloid. Graphs report Mean ± Standard Deviation. *Abbreviations: fx, fornix; cing, cingulum bundle; cc, corpus callosum; alv, alveus.*

the first anatomical abnormalities correlated with worsened memory performance.[66–68] More

recent studies utilizing functional readouts and dietary interventions suggest that while the MB

may be a primary site of sensitivity, thalamic, hippocampal, and cortical alterations also coincide

with worsening memory.[67,69,70] Interestingly, the interventions restore hippocampal connectivity

to the diencephalic mammillary body and thalamus[69,70], which are strikingly similar results to the

experimental trials using deep brain stimulation of the fornix near the MB to enhance memory in

AD patients.[71] Together, this data suggest that dysfunction within the MB correlates with HPC

abnormalities and memory impairment, like those seen in AD. Importantly, recent human

studies correlating MB abnormalities to memory performance specifically link MB aberrations to

impaired recall memory,[38,70] the type of memory thought to be most affected in AD.[72,73]

In addition to a better understanding of the regions that show primary Aβ aggregation and

promote the initial network destabilization, our data suggest mechanisms for Aβ propagation

between connected brain regions.[74] By demonstrating enhanced Aβ deposition within limbic

white matter tracts, and unique axonal patterns of Aβ within the Papez and limbic circuit tract,

we propose that specific disconnection of the core memory network underlies cognitive

impairment in AD. This may be due to particular mechanisms of Aβ handling within primary

neurons of the memory network, or the particular vulnerability of the cells to overproduction or

mishandling of APP/ Aβ. Additionally, by showing that distinct networks are burdened by Aβ,

with increasingly complex networks affected across age, we present a unifying view of early

lesions and can provide a framework in which to better interpret human pathogenesis.

Investigators have suggested that the DMN, limbic system, attentional systems, and brain stem

may all be involved at the earliest stages of prodromal AD. The data presented in this

manuscript suggest that these do not need to be disparate hypotheses, but instead represent

different stages of network affliction that occur as the disease progresses. In the mouse model,

2M 5XFAD animals show core limbic affliction, followed by the DMN at 4M and typical limbic

system at 6M. These observations suggest that early AD may be best staged by the network

aberrations detected by functional MRI techniques. These alterations occur before overt

memory loss,[33] and with increasing evidence for restoration of network function as a successful

treatment in AD models,[9,72,75] our data may lay the ground work for network-progression staging

to guide early-disease circuit interventions.

Methods

Animals

All mouse work was approved by the Committee for Animal Care of the Division of Comparative

Medicine at the Massachusetts Institute of Technology. 5XFAD (Tg 6799) breeding pairs were

acquired from the Mutant Mouse Resource and Research Center (MMRRC) (Stock No. 034848-

JAX) and maintained as hemizygous on the BL6 background. Animals were group housed on a

12h light/dark cycle with Nestlet enrichment and sacrificed at the ages noted in the text.

Mouse Tissue Fixation

Mice were deeply anesthetized with isoflurane (Isoflurane, USP, Piramal Healthcare, Andhra

Pradesh, India) and underwent transcardial perfusion with ice cold 1X PBS (10X stock, Gibco,

#70011-044) followed by ice-cold fixative made of 4% paraformaldehyde (32% stock, Electron

Microscopy Sciences (EMS), Hatfield, PA, #15714) and 1% glutaraldehyde (10% stock, EMS

#16110) in 1X PBS. Brains were removed from the skull and post-fixed in the same fixative for 3

days shaking at 4C.

Whole Mouse Brain Processing

After washing in 1X PBS, brains incubated in inactivation solution of 1% acrylamide (40% stock,

Biorad #161-0140), 1M glycine (Sigma-Aldrich, St. Louis, MO, #G7126) in 1X PBS. After

washing in 1X PBS, brains were put into clearing solution of 200mM sodium dodecyl sulfate

(SDS) (Sigma-Aldrich, #L3771), 20mM lithium hydroxide monohydrate (Sigma, #254274),

40mM boric acid (Sigma-Aldrich, #7901), pH 8.5-9.0 and left shaking at 55C for 4 weeks until

white matter tracts were translucent to the eye in SDS. Brains were washed in 1X PBS or Weak

Binding Solution (WBS) for up to 1 week and immunolabeled with the 'Whole Brain iSWITCH'

protocol attached.

Mouse Brain Section Processing

Where sections were analyzed, brains were sliced to 100um on a vibratome (Leica VT100S)

and stored at 4C in 1X PBS + 0.02% sodium azide (Sigma-ALdrich, #S2002). For iSWITCH

labeling, individual sections were incubated in clearing solution shaking at 55C for 2 hours and

were washed in 1X PBS. Sections were immunolabeled with the 'Mouse Brain Section

iSWITCH' protocol attached.

Human Tissue Processing

Human tissue blocks were deparaffinized by sequential immersion in xylene, ethanol, and water

(details in Human iSWITCH protocol). Then, blocks were incubated in 1% GA in 1X PBS for 10

days at 4C. Brains were incubated in clearing solution shaking at 55C until the tissue appeared

translucent (4-8 weeks). Following clearing, tissue was labeled using the Human iSWITCH

protocol.

Antibodies & Dyes

The primary antibodies used are as follows:

Target	Host	Company	Product #	Dilution
Aβ (D54D2)	Rabbit	Cell Signaling Technologies	8243	1:100 –1000
Myelin Basic Protein* SMI-99	Mouse	BioLegend	808401	1:50 –500
Myelin Basic Protein* SMI-94	Mouse	BioLegend	836502	1:50 –500

*These antibodies were used concurrently as per the manufacturer's recommendation.

Additional labeling reagents:

Hoechst 33528 was used for nuclear labeling (Sigma #14530).

All secondary antibodies were Pre-adsorbed F(ab)2' AlexaFluor-conjugated from AbCam.

Whole Brain Image Acquisition

Whole brain images were acquired on a custom SPIM microscope built by H.C. During imaging,

samples were illuminated with a sheet of light generated by scanning a focused beam from a

light from a broad spectrum laser unit (SOLE -6 with 405, 488, 561, 647 nm, Omicron) through a

low NA objective (Macro 4X/ 0.28 NA, Olympus) with a galvo-scanner (6215H, Cambridge

Technology). Collection of emitted light on the microscope occurs through a long working

distance high NA objective (10x/0.6 NA WD 8mm CLARITY, Olympus). The microscope is

outfitted with four sCMOS cameras (Orca Flash4.0 V2, Hamamatsu) for simultaneous

multichannel detection and signal recording. During acquisition, the samples were illuminated

simultaneously from dual illumination arms (one on each side) to minimize the shading effects of

light scattering elements in the brain. Dynamic confocal mode of detection is implemented by

synchronizing the scanning speed of the illumination beam with the read-out speed of the rolling

shutter mode of sCMOS camera, which improves the signal to background ratio by filtering out

background noise from out-of-focal planes. Sample is mounted on a motorized stage with x, y, z

translation and theta rotation (M-112.2DG, M-111K028, M-116.DG, Physik Instrumente) for

mosaic imaging. Z-stack imaging by sample scanning alone is slow due to communication

overhead between the host computer and the stage controller. We achieved high-speed volume

imaging by scanning the light sheet along the depth direction with a galvo-scanner and

synchronizing the position of the light sheet with the detection objective's focal plane by moving

the detection objective with a piezo actuator (P-628.1CL, Physik Instrumente). To maintain light

sheet position on the focal plane of the objective across the entire sample volume, we

implemented an image based autofocusing algorithm.[76] The laser settings are determined such

that about 5% of the images are saturated to its maximum gray level for high signal to

background ratio. The sample chamber is filled with the refactive index matching solution

(RIMS)[77]. Depending on the refractive index of the medium, the beam waist position of the

illumination light sheet shifts along the illumination beam direction. Each illumination objective is

mounted on the piezo actuator (P-628.1CL, Physik Instrumente) to allow the beam waist

position to be adjusted to the center of the detection objective. Sample chamber is specially

designed to allow for free motion of the detection objective while preventing leakage of the

immersion medium.

<u>Whole Brain Image Processing</u>

Each tile is first corrected for the non-uniform illumination pattern using a modified algorithm

from Smith and colleages[78]. Multiple stacks of acquired images are stitched with Terastitcher.[79]

Each tile has 15% overlapping area with the neighboring tiles for calculating stitching

parameters. The voxel size of raw data is 0.58 x 0.58 x 5.0um. After stitching, the data set is

down-sampled in X and Y dimension for further analysis with Imaris software (Bitplane).

<u>Section Image Acquisition</u>

For section and human tissue imaging, tissue was mounted onto microscope slides (VWR

VistaVision, VWR International, LLC, Radnor, PA) with either Fluromount G Mounting Medium

(Electron Microscopy Sciences, Hatfield, PA, USA) or RIMS solution[77].

Slice images were acquired on a Zeiss LSM Inverted 710 microscope using Zen 2012 software

(Carl Zeiss Microscopy, Jena, Germany). Images with cellular resolution were taken using a C-

apochromat 40X, water immersion objective, NA 1.20. Section overview images used a Plan-

apochromat 5X, air objective, NA 0.16. Pinhole, optical sectioning, and laser settings were

determined for each experiment and kept consistent for all images within an experiment or that
were included within one graph.
444
Human Brain Image Acquisition
Human brain images were acquired on a Leica TCS SP8 Confocal Microscope using LASAF
software (Leica Microsystems, Wetzlar, Germany). Images were taken using a 20X 1.0NA
CLARITY optimized objective with 6mm working distance. The pinhole, optical sectioning, and
resolution and laser settings were empirically determined for one brain, and kept constant for
imaging subsequent samples.
451
3D Image Quantification
Images were analyzed using Imaris (Bitplane, Zurich, Switzerland). All quantification steps were
performed on raw images by blinded investigators. For whole brain analyses (Figure 1), each
whole brain file was segmented by hand using white matter tract and regional guidelines from
the Allen Brain Atlas (Allen Mouse Brain Atlas, Coronal) to delineate boundaries for each major
brain region (Video 5). After segmentation, a spots object was created on a 12 month brain. The
parameters were fixed over the entire brain, and spots were separated into the bounded brain
regions using the Spots into Surfaces tool in Imaris Xtensions. A spots object was created on
each brain and these objects were similarly split into brain regions using the Xtension. Data was
exported to CSV and analyzed GraphPad Prism 7.0a for Mac. For human brain analyses
(Figure 3), the 3D rendering was randomly segmented into five, $0.5mm^3$ sub-sections to account
for differences in block size. An identical spots object was created within each of the five sub-
sections, and the data was exported to CSV and analyzed GraphPad Prism 7.0a for Mac.
465
2D Image Quantification
For 2D analyses, images were imported into FIJI[80] as TIFF files. Relevant values were exported
to GraphPad for statistical analyses. For human amyloid analyses (Figure 3), two blinded
observers counted plaques within images using the Multi-point tool. Data was saved as a CSV
and analyzed in GraphPad Prism 7.0a for Mac. For white matter quantification (Figure 4),
blinded observers analyzing only the myelin image channel of a 5X section-overview image
segmented white matter tracts. Then the segmentation was overlaid on the Aβ channel as a
selection, within which a threshold was applied to the images. Finally, the Analyze Particles tool
was used to count individual deposits. Data was saved in a spreadsheet and analyzed in
GraphPad Prism 7.0a for Mac.

<u>Representative Images and Videos from Whole Brain Data</u>

Representative images from the whole-brain datasets are either 2D images of the 3D rendered

whole dataset or digitally sectioned at 5-100um in Imaris using the Orthoslicer tool. Videos are

created using the Key Frame Animation tool in Imaris. The brightness of the images has been

individually adjusted for each brain to enhance 2D/3D viewing of specific objects. Because of

these alterations, no direct comparisons of labeling intensity should be made between images.

<u>*In situ* hybridization probe design</u>

RNA antisense probes were generated by PCR-amplifying human cDNA with human-specific

APP primers with a T7 RNA polymerase recognition sequence (TAATACGACTCACTATAGGG)

fused to the reverse primer (Table 1). The resulting PCR product was gel extracted and *in vitro*

transcribed using a DIG-RNA labeling kit (Roche).

<u>Immuno-*in situ* hybridization (Immuno-ISH)</u>

Mice were anesthetized by isoflurane in an open system and perfused with RNase-free PBS

followed by RNase-free 4% formaldehyde. Brains were dissected, drop fixed in RNase-free 4%

formaldehyde for 12 hours, equilibrated in 30% sucrose-PBS, and frozen in O.C.T. (TissueTek).

Cryosections (10μm) were incubated with a DIG-labeled RNA antisense probe (1:1000 in

hybridization buffer) overnight at 65^{o}C, washed in 1X SSC/50% formamide/0.1% Tween-20 3X

30 minutes at 65^{o}C followed by 1X MABT for 30 minutes at room temperature. Sections were

blocked with 20% heat-inactivated sheep serum/2% blocking reagent (Roche)/1X MABT for 1

hour and then incubated with mouse anti-DIG antibody (Roche; 1:2000) and rabbit anti-amyloid

β (Cell Signaling; 1:500) diluted in blocking solution overnight. Sections were washed with

1XMABT 2X 20 minutes, incubated with donkey anti-rabbit Alexa-488 (Invitrogen; 1:1000)

diluted in blocking solution for 1 hour, and washed with 1XMABT 5X 20 minutes. Sections were

then prestained with 100mM NaCl/50mM MgCl$_2$/100mM Tris pH 9.5/0.1% Tween-20 2X 10

minutes, followed by staining with NBT/BCIP (Roche; 4.5 μl/ml and 3.5 μl/ml, respectively, in

prestaining buffer) for 2 hours. Sections were washed with 1X PBS 3X 15 minutes, incubated in

xylene 3X 5 minutes, and mounted with VectaMount (Vector Laboratories).

Gene	Forward Primer	Reverse Primer
h*APP*	GAGACACCTGGGGAT GAGAA	TAATACGACTCACTATAGGGACAGAGTCAGCC CCAAAAGA

Table 1. Primer sequences (5'-3') for *in situ* probe preparation.

Ex vivo electrophysiology

Acute hippocampal slices were prepared from C57bl/6 mice aged2-2.5 months old, which are

wild type group and 5xFAD group. The mice were anesthetized with isoflurane and decapitated.

The experimenter was blinded to the group of animal. Transverse brain slices (250 μm thick)

were prepared in ice-cold dissection buffer (in mM: 211 sucrose, 3.3 KCl, 1.3 NaH2PO4, 0.5

CaCl2, 10 MgCl2, 26 NaHCO3 and 11 glucose) using a Leica VT1000S vibratome (Leica).

Slices were recovered in a submerged chamber with 95% O2/5% CO2-saturated artificial

cerebrospinal fluid (ACSF) consisting of (in mM) 124 NaCl, 3.3 KCl, 1.3 NaH2PO4, 2.5 CaCl2,

1.5 MgCl2, 26 NaHCO3 and 11 glucose for 1 h at 28–30 °C. Whole cell patch clamp with

current steps from 0 pA to +130 pA at 10 pA increments for 800ms were used to verify the

ability to form action potential (AP) on HEKA ECP 10 Amplifier (Germany) with Pulse software.

Number of APs at each current step; Resting membrane potential (RMP); Threshold of APs;

Amplitude of APs from RMP and threshould as well as fast afterhyperpolarization(fAHP) were

measured and analyzed.

Statistics

All statistics were performed in MatLab or GraphPad Prism. Individual statistical tests are

indicated in the text and/or figure legends for the appropriate experiments. Graphs were created

in the respective analytical software packages and exported as .TIFF, .PDF, or .JPEG for

inclusion in the document.

Acknowledgements

The authors would like to acknowledge Michiel Kooreman at the Netherlands Brain Bank for

help procuring the PPFE human tissue and Teresa Lima for her expert advice in preparing the

human PPFE brain tissue for the optimized CLARITY protocol. We would also like to thank Nina

Dedic for her advice and helpful comments on the project and manuscript. Additionally, we

would like to acknowledge Naveed Bakh, Sung-Yon Kim, and Kamilla Tekiela for helping lay the

groundwork for the experiments presented in this work.

Funding

This work was funded by NIH grant RF1AG047661 to L.-H.T. and by the Norman B. Leventhal

and Barbara Weedon fellowships to R.G.C. Additionally, K.C. was supported by Burroughs

Wellcome Fund Career Awards at the Scientific Interface, the Searle Scholars Program,

544
545 **Citations**

1. Hurd, M. D., Martorell, P., Delavande, A., Mullen, K. J. & Langa, K. M. Monetary costs of
dementia in the United States. *N. Engl. J. Med.* **368,** 1326–34 (2013).
2. Hardy, J. & Higgins, G. Alzheimer's Disease□: The Amyloid Cascade Hypothesis.
Science (80-.). **256,** 184–185 (1992).
3. Musiek, E. S. & Holtzman, D. M. Three dimensions of the amyloid hypothesis□: time ,
space and ' wingmen '. *Nat. Neurosci.* **18,** (2015).
4. Mutations | AlzForum. Available at: http://www.alzforum.org/mutations/. (Accessed: 20th
October 2016)
5. Hong, S. *et al.* Complement and microglia mediate early synapse loss in Alzheimer
mouse models. *Science (80-.).* **8373,** 1–9 (2016).
6. Wei, W. *et al.* Amyloid beta from axons and dendrites reduces local spine number and
plasticity. *Nat. Neurosci.* **13,** 190–196 (2009).
7. Scheuner, D. *et al.* Secreted amyloid beta-protein similar to that in the senile plaques of
Alzheimer's disease is increased in vivo by the presenilin 1 and 2 and APP mutations
linked to familial Alzheimer's disease. *Nat. Med.* **2,** 864–870 (1996).
8. Kimura, R. & Ohno, M. Impairments in remote memory stabilization precede hippocampal
synaptic and cognitive failures in 5XFAD Alzheimer mouse model. *Neurobiol. Dis.* **33,**
229–235 (2009).
9. Palop, J. J. *et al.* Aberrant Excitatory Neuronal Activity and Compensatory Remodeling of
Inhibitory Hippocampal Circuits in Mouse Models of Alzheimer's Disease. *Neuron* **55,**
697–711 (2007).
10. Huijbers, W. *et al.* Amyloid-beta deposition in mild cognitive impairment is associated with
increased hippocampal activity, atrophy and clinical progression. *Brain* **138,** 1023–1035
(2015).
11. Thal, D. R., Rüb, U., Orantes, M. & Braak, H. Phases of A beta-deposition in the human
brain and its relevance for the development of AD. *Neurology* **58,** 1791–1800 (2002).
12. Mormino, E. C. *et al.* Synergistic effect of β-amyloid and neurodegeneration on cognitive
decline in clinically normal individuals. *JAMA Neurol.* **71,** 1379–85 (2014).
13. Jack, C. R. *et al.* Brain beta-amyloid measures and magnetic resonance imaging atrophy
both predict time-to-progression from mild cognitive impairment to Alzheimer's disease.
Brain **133,** 3336–3348 (2010).
14. Buckner, R. L., Andrews-Hanna, J. R. & Schacter, D. L. The brain's default network:
Anatomy, function, and relevance to disease. *Ann. N. Y. Acad. Sci.* **1124,** 1–38 (2008).
15. Villemagne, V. L. *et al.* Amyloid Imaging with 18F-Florbetaben in Alzheimer Disease and
Other Dementias. *J. Nucl. Med.* **52,** 1210–1217 (2011).
16. Klunk, W. E. *et al.* Imaging brain amyloid in Alzheimer's disease with Pittsburgh
Compound-B. *Ann. Neurol.* **55,** 306–319 (2004).
17. Villemagne, V. L. *et al.* Longitudinal assessment of Abeta and cognition in aging and
Alzheimer disease. *Ann. Neurol.* **69,** 181–192 (2011).
18. Jack, C. R. *et al.* Rates of beta-amyloid accumulation are independent of hippocampal
neurodegeneration. *Neurology* **82,** 1605–1612 (2014).

19.   Greicius, M. D., Srivastava, G., Reiss, A. L. & Menon, V. Default-mode network activity
588   distinguishes Alzheimer's disease from healthy aging: evidence from functional MRI.
589   *Proc. Natl. Acad. Sci. U. S. A.* **101,** 4637–42 (2004).

590   20.   Fletcher, E. *et al.* Beta-amyloid, hippocampal atrophy and their relation to longitudinal
591   brain change in cognitively normal individuals. *Neurobiol. Aging* **40,** 173–180 (2016).

592   21.   Fletcher, E., Carmichael, O., Pasternak, O., Maier-Hein, K. H. & DeCarli, C. Early Brain
593   Loss in Circuits Affected by Alzheimer's Disease is Predicted by Fornix Microstructure but
594   may be Independent of Gray Matter. *Front. Aging Neurosci.* **6,** 106 (2014).

595   22.   Bakker, A., Albert, M. S., Krauss, G., Speck, C. L. & Gallagher, M. Response of the
596   medial temporal lobe network in amnestic mild cognitive impairment to therapeutic
597   intervention assessed by fMRI and memory task performance. *NeuroImage Clin.* **7,** 688–
598   698 (2015).

599   23.   Reitz, C. & Mayeux, R. Alzheimer disease: Epidemiology, diagnostic criteria, risk factors
600   and biomarkers. *Biochem. Pharmacol.* **88,** 640–651 (2014).

601   24.   De Strooper, B. & Karran, E. The Cellular Phase of Alzheimer's Disease. *Cell* **164,** 603–
602   615 (2016).

603   25.   Oakley, H. *et al.* Intraneuronal β-Amyloid Aggregates, Neurodegeneration, and Neuron
604   Loss in Transgenic Mice with Five Familial Alzheimer's Disease Mutations: Potential
605   Factors in Amyloid Plaque Formation. *J. Neurosci.* **26,** 10129–10140 (2006).

606   26.   Murray, E. *et al.* Simple, Scalable Proteomic Imaging for High-Dimensional Profiling of
607   Intact Systems. *Cell* **163,** 1500–1514 (2015).

608   27.   Barnes, A. The specificity of pH and ionic strength effects on the kinetics of the Rh (D)-
609   anti-Rh (D) system. *J. Immunol.* **96,** 854–864 (1966).

610   28.   Eimer, W. A. & Vassar, R. Neuron loss in the 5XFAD mouse model of Alzheimer's
611   disease correlates with intraneuronal Aβ42 accumulation and Caspase-3 activation. *Mol.*
612   *Neurodegener.* **8,** 2 (2013).

613   29.   Richardson, D. S. & Lichtman, J. W. Clarifying Tissue Clearing. *Cell* **162,** 246–257
614   (2015).

615   30.   Jost, B. C. & Grossberg, G. T. The evolution of psychiatric symptoms in Alzheimer's
616   disease: A natural history study. *J. Am. Geriatr. Soc.* **44,** 1078–1081 (1996).

617   31.   Stafford, J. M. *et al.* Large-scale topology and the default mode network in the mouse
618   connectome. *Proc. Natl. Acad. Sci. U. S. A.* **111,** 18745–50 (2014).

619   32.   Hedden, T. *et al.* Disruption of functional connectivity in clinically normal older adults
620   harboring amyloid burden. *J Neurosci* **29,** 12686–12694 (2009).

621   33.   Sperling, R., LaViolette, P. & O'Keefe, K. Amyloid deposition is associated with impaired
622   default network function in older persons without dementia. *Neuron* **63,** 178–188 (2009).

623   34.   Aggleton, J. P., Pralus, A., Nelson, A. J. D. & Hornberger, M. Thalamic pathology and
624   memory loss in early Alzheimer's disease: moving the focus from the medial temporal
625   lobe to Papez circuit. *Brain* aww083 (2016). doi:10.1093/brain/aww083

626   35.   Siskova, Z. *et al.* Dendritic structural degeneration is functionally linked to cellular
627   hyperexcitability in a mouse model of alzheimer's disease. *Neuron* **84,** 1023–1033
628   (2014).

629   36.   Vossel, K. A. *et al.* Seizures and epileptiform activity in the early stages of Alzheimer
630   disease. *JAMA Neurol.* **70,** 1158–66 (2013).

631   37.   Dillingham, C. M., Frizzati, A., Nelson, A. J. D. & Vann, S. D. How do mammillary body
632   inputs contribute to anterior thalamic function? *Neurosci. Biobehav. Rev.* **54,** 108–119
633   (2015).

634   38.   Tsivilis, D. *et al.* A disproportionate role for the fornix and mammillary bodies in recall
635   versus recognition memory. *Nat. Neurosci.* **11,** 834–842 (2008).

636   39.   Llinás, R. R. & Alonso, A. Electrophysiology of the mammillary complex in vitro. II. Medial
637   mammillary neurons. *J. Neurophysiol.* **68,** 1321–31 (1992).

40. Verret, L. *et al.* Inhibitory interneuron deficit links altered network activity and cognitive
dysfunction in alzheimer model. *Cell* **149,** 708–721 (2012).
41. Vann, S. D. & Nelson, A. J. D. in *Progress in Brain Research* **219,** 163–185 (2015).
42. Copenhaver, B. R. *et al.* The fornix and mammillary bodies in older adults with
Alzheimer's disease, mild cognitive impairment, and cognitive complaints: A volumetric
MRI study. *Psychiatry Res. - Neuroimaging* **147,** 93–103 (2006).
43. McDuff, T. & Sumi, S. M. Subcortical degeneration in Alzheimer's disease. *Neurology* **35,**
123–126 (1985).
44. Callen, D. J. A., Black, S. E., Caldwell, C. B. & Grady, C. L. The influence of sex on limbic
volume and perfusion in AD. *Neurobiol. Aging* **25,** 761–770 (2004).
45. Hornberger, M. *et al.* In vivo and post-mortem memory circuit integrity in frontotemporal
dementia and Alzheimer's disease. *Brain* **135,** 3015–3025 (2012).
46. Nestor, P. J., Fryer, T. D., Smielewski, P. & Hodges, J. R. Limbic hypometabolism in
Alzheimer's disease and mild cognitive impairment. *Ann. Neurol.* **54,** 343–351 (2003).
47. Braak, H. & Braak, E. Neuropathological stageing of Alzheimer-related changes. *Acta*
Neuropathol. **82,** 239–259 (1991).
48. Sheline, Y. I. *et al.* Amyloid Plaques Disrupt Resting State Default Mode Network
Connectivity in Cognitively Normal Elderly. *Biol. Psychiatry* **67,** 584–587 (2010).
49. Lim, H. K. *et al.* Regional amyloid burden and intrinsic connectivity networks in cognitively
normal elderly subjects. *Brain* **137,** 3327–3338 (2014).
50. Li, W. *et al.* Aberrant functional connectivity in Papez circuit correlates with memory
performance in cognitively intact middle-aged APOE4 carriers. *Cortex* **57,** 167–176
(2014).
51. Vann, S. D. Dismantling the papez circuit for memory in rats. *Elife* **2013,** 1–21 (2013).
52. Jucker, M. & Walker, L. C. Self-propagation of pathogenic protein aggregates in
neurodegenerative diseases. *Nature* **501,** 45–51 (2013).
53. Avila, J., Perry, G., Strange, B. A. & Hernandez, F. Alternative neural circuitry that might
be impaired in the development of Alzheimer disease. *Front. Neurosci.* **9,** 1–5 (2015).
54. Allen, G. V & Hopkins, D. a. Mamillary body in the rat: topography and synaptology of
projections from the subicular complex, prefrontal cortex, and midbrain tegmentum. *J.*
Comp. Neurol. **286,** 311–336 (1989).
55. Terry, R. D. *et al.* Physical basis of cognitive alterations in Alzheimer's disease: Synapse
loss is the major correlate of cognitive impairment. *Ann. Neurol.* **30,** 572–580 (1991).
56. DeKosky, S. T. & Scheff, S. W. Synapse loss in frontal cortex biopsies in Alzheimer's
disease: Correlation with cognitive severity. *Ann. Neurol.* **27,** 457–464 (1990).
57. Scheff, S. W. & Price, D. A. Synapse Loss in the Temporal Lobe in Alzheimer 's Disease.
Ann. Neurol. (1993).
58. Hyman, B. T., Van Hoesen, G. W., Damasio, A. R. & Barnes, C. L. Alzheimer's disease:
cell-specific pathology isolates the hippocampal formation. *Science* **225,** 1168–70 (1984).
59. Yoder, R. M., Peck, J. R. & Taube, J. S. Visual landmark information gains control of the
head direction signal at the lateral mammillary nuclei. *J. Neurosci.* **35,** 1354–67 (2015).
60. Sharp, P. E. & Koester, K. Lesions of the mammillary body region severely disrupt the
cortical head direction, but not place cell signal. *Hippocampus* **18,** 766–784 (2008).
61. Whitehouse, P. J., Price, D. L., Clark, a W., Coyle, J. T. & DeLong, M. R. Alzheimer
disease: evidence for selective loss of cholinergic neurons in the nucleus basalis. *Ann*
Neurol **10,** 122–126 (1981).
62. Dean 3rd, D. *et al.* Association of Amyloid Pathology With Myelin Alteration in Preclinical
Alzheimer Disease. *JAMA Neurol.* **Epub ahead,** (2016).
63. Hargis, K. E. & Blalock, E. M. Transcriptional signatures of brain aging and Alzheimer's
disease: What are our rodent models telling us? *Behavioural Brain Research* (2016).
doi:10.1016/j.bbr.2016.05.007

64.   Sabbagh, J. J., Kinney, J. W. & Cummings, J. L. Alzheimer's disease biomarkers:
690   Correspondence between human studies and animal models. *Neurobiology of Disease*
691   **56,** 116–130 (2013).
692   65.   Grossi, D., Lopez, O. L. & Martinez, A. J. Mamillary bodies in Alzheimer's disease. *Acta*
693   *Neurol.Scand.* **80,** 41–45 (1989).
694   66.   Sullivan, E. V *et al.* In vivo mammillary body volume deficits in amnesic and nonamnesic
695   alcoholics. *Alcohol. Clin. Exp. Res.* **23,** 1629–36 (1999).
696   67.   Sullivan, E. V. & Pfefferbaum, A. Neuroimaging of the Wernicke-Korsakoff syndrome.
697   *Alcohol Alcohol.* **44,** 155–165 (2009).
698   68.   Beh, S. C., Frohman, T. C. & Frohman, E. M. Isolated mammillary body involvement on
699   MRI in Wernicke's encephalopathy. *J. Neurol. Sci.* **334,** 172–175 (2013).
700   69.   Nahum, L. *et al.* Neural Correlate of Anterograde Amnesia in Wernicke???Korsakoff
701   Syndrome. *Brain Topogr.* **28,** 760–770 (2015).
702   70.   Kim, E. *et al.* Restoration of mammillothalamic functional connectivity through thiamine
703   replacement therapy in Wernicke's encephalopathy. *Neurosci. Lett.* **479,** 257–261 (2010).
704   71.   Laxton, A. W. *et al.* A phase I trial of deep brain stimulation of memory circuits in
705   Alzheimer's disease. *Ann. Neurol.* **68,** 521–534 (2010).
706   72.   Roy, D. S. *et al.* Memory retrieval by activating engram cells in mouse models of early
707   Alzheimer's disease. *Nature* (2016). doi:10.1038/nature17172
708   73.   Fischer, A., Sananbenesi, F., Wang, X., Dobbin, M. & Tsai, L.-H. Recovery of learning
709   and memory is associated with chromatin remodelling. *Nature* **447,** 178–82 (2007).
710   74.   Ye, L. *et al.* Progression of Seed-Induced Abeta Deposition within the Limbic
711   Connectome. *Brain Pathol.* **25,** 743–752 (2015).
712   75.   Iaccarino, H. F. *et al.* Gamma frequency entrainment attenuates amyloid load and
713   modifies microglia. *Nature* **540,** 230–235 (2016).
714   76.   Tomer, R., Ye, L., Hsueh, B. & Deisseroth, K. Advanced CLARITY for rapid and high-
715   resolution imaging of intact tissues. *Nat. Protoc.* **9,** 1682–97 (2014).
716   77.   Yang, B. *et al.* Single-cell phenotyping within transparent intact tissue through whole-
717   body clearing. *Cell* **158,** 945–958 (2014).
718   78.   Smith, K. *et al.* CIDRE: an illumination-correction method for optical microscopy. *Nat.*
719   *Methods* **l,** 1–7 (2015).
720   79.   Bria, A. & Iannello, G. TeraStitcher - a tool for fast automatic 3D-stitching of teravoxel-
721   sized microscopy images. *BMC Bioinformatics* **13,** 316 (2012).
722   80.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*
723   *Methods* **9,** 676–82 (2012).
724

725 Supplementary Figure 1 – supporting Figure 1

Supplemental Figure 1: Optimized iSWITCH enhances immunolabeling homogeneity in 100um sections. A) Representative optical sections from the first (left), middle (center), and last (right) slices of z-stack image files taken from tissue labeled for myelin basic protein (MBP, red) glial fibrillary acidic protein (GFAP, green), and with Hoechst dye (blue) using either iSWITCH buffers (top) or tradititional labeling buffer phosphate buffered saline with triton x-100 (PBST, bottom). B) Quantification of MBP labeling demonstrating signal inensity consistency between the first and last sections under both conditions (left), but enhanced middle section immunolabeling signal when compared to either the intensity of the first (center) or last (right) section. C) Immunofluorescent signal loss throughout the depth of the sections is first order decay using iSWITCH buffers, indicating relatively homogenous signal with intensity that decays with depth due to laser attenuation, but using PBST the loss is a second order polynomial, indicating non-homogenous signal intensity through the thickness of the tissue. *p ≤ 0.05; **p ≤ 0.01; ****p≤ 0.0001; Graphs report Mean ± Standard Deviation.

726 Supplementary Figure 2 – supporting Figure 1

Supplemental Figure 2: iSWITCH buffer immunolabeling whole brain for β-amyloid achieves homogenous signal. Aged 5XFAD brain was split into two hemispheres and labeled with either iSWITCH buffers or PBST. A) Representative images from the center section of each hemisphere. B) Demonstration of the consistency of overall aggregate counts between animals and no consistent effect of PBST or iSWITCH on aggregate detection. C) Aggregates labeled by iSWITCH show overall lower intensity, which may be driven by more antibody availability for surface antigens. D) Plot of the intensity of aggregates detected through sample (left axis) with regression (right axis) showing non-homogenous signal intensities in PBST labeled brains, which is rectified using iSWITCH buffers. Graphs report Mean ± Standard Deviation unless otherwise noted in the text.

Supplementary Figure 3 – Supporting Figure 1

Supplemental Figure 3: White-matter based hand segmentation of brain areas provides robust identification of specific regions. A) 3D rendering of white matter tracts in the whole mouse brain. B) Optical slice-wise visualization of white matter tracts. C) Representative examples of hand-segmentation (left) next to the Allen Mouse Brain Atlas section (right, anterior section #43, posterior section #85) from which they were drawn. Website: © 2015 Allen Institute for Brain Science. Allen Mouse Brain Atlas [Internet]. Available from: http://mouse.brain-map.org[311] D) Representative images showing the whole-brain segmentation. E) Representative images of the volume-rendered segmentation.

Supplementary Figure 4 – supporting Figure 1

Supplemental Figure 5: **Hierarchical clustering at each age reveals amyloid deposition within networks appears with age**. A-D) Dendrograms of hierarchical clustering of amyloid density at each age examined in the 5XFAD mice. A) 12 months. B) 6 months C) 4 months and D) 2 months aged animals.

Supplementary Figure 5 – supporting Figure 1

Supplemental Figure 4: Imaging reveals no correlation between transgenic mRNA expression and amyloid aggregates . A) Representative images of the double *in situ* hybridization and immunofluorescence. B) Brain-wide correlation graph demonstration lack of correlation between signals. C) Correlation coefficients split by region and displayed with p-value.