1   **Title**: Decomposing leaf mass into photosynthetic and structural components explains
2   divergent patterns of trait variation within and among plant species
3
4   **Running title**: Photosynthetic and structural leaf mass
5

6   **Authors**: Masatoshi Katabuchi[1,2 *], Kaoru Kitajima[1,3,4], S. Joseph Wright[4], Sunshine A. Van
7   Bael[4,5], Jeanne L. D. Osnas[1] and Jeremy W. Lichstein[1]
8
[1] Department of Biology, University of Florida, Gainesville, FL 32611, USA
[2] Department of Agricultural and Biological Engineering, University of Florida, Gainesville,
FL 32611, USA
[3] Graduate School of Agriculture, Kyoto University, Kitashirakawa Oiwake-Cho, Kyoto 606-
8502 Japan
[4] Smithsonian Tropical Research Institute, 9100 Panama City Pl., Washington, DC 20521
[5] Department of Ecology and Evolutionary Biology, Tulane University, New Orleans, LA
70118 USA

17
18 **Keywords:** Bayesian inference, functional diversity, functional traits, global vegetation
19 models, interspecific variation, intraspecific variability, latent variables, null models, optimal
20 leaf lifespan
21
22 **Type of article:** Letter
23
Number of words in abstract: 150
Number of words in main text: 4919
Number of references: 42
Number of figures: 5
Number of tables: 0
Number of text boxes: 1 (750 words)
30

31 [*] **Corresponding Author:** Masatoshi Katabuchi
32 **Mailing address:** Department of Biology, University of Florida, Gainesville, FL 32611,
33 USA
34 **Email:** mattocci27@gmail.com
35 **Telephone**: 352-392-1540

36
37 **Statement of authorship:**
38 MK, JWL and JLDO conceived of the study; KK, SJW and SAVB contributed data; MK
39 devised the analytical approach and performed analyses; MK and JWL wrote the first draft of
40 the manuscript, and all authors contributed to revisions.
41
42 **Data accessibility:**
43 Should the manuscript be accepted, the data supporting the results will be archived in an
44 appropriate public repository such as Dryad or Figshare, the model will be archived in
45 GitHub, and the data DOI will be included at the end of the article.
46
47

Abstract
Across the global flora, photosynthetic and metabolic rates depend more strongly on leaf area
than leaf mass. In contrast, intraspecific variation in these rates is strongly mass-dependent.
We show that decomposing leaf mass per area (LMA) into two components –
'photosynthetic' LMAp (which determines photosynthetic capacity and metabolic rates, and
also affects optimal leaf lifespan) and 'structural' LMAs (which determines leaf toughness
and potential leaf lifespan) – provides improved predictions of trait variation (photosynthesis,
respiration, and lifespan) across the global flora, and within and among tropical plant species
in Panama. Our analysis shows that most interspecific LMA variation is due to LMAs (which
explains why traits are area-dependent across species) and that intraspecific LMA variation is
due to changes in both LMAp and LMAs (which explains why traits are mass-dependent
within species). Our results suggest that leaf trait variation is multi-dimensional, thereby
questioning the validity of the one-dimensional worldwide leaf economics spectrum.

61
Introduction
Over the past two decades, the 'leaf economics spectrum' (LES) (Reich *et al.* 1997;
Wright *et al.* 2004; Reich 2014) has become a cornerstone of plant ecology. The LES
captures tight relationships among leaf mass per area (LMA), leaf lifespan (LL), and mass-
normalized leaf traits related to carbon fixation and nutrient use, suggesting the presence of a
single dominant axis of leaf functional variation, ranging from short-lived leaves with high
photosynthetic potential and fast returns on investment to long-lived leaves with low
photosynthetic potential and slow returns (Wright *et al.* 2004; Westoby & Wright 2006;
Reich 2014). Because the mass-based LES constrains variation in green leaf traits to a single
axis, it has been proposed as a simple framework for incorporating more realistic levels of
leaf trait diversity in global carbon-climate models (Bonan *et al.* 2002; Wright *et al.* 2004).
However, recent analyses suggest that the one-dimensional LES is largely the result of high
interspecific variation in LMA combined with mass-normalization of area-dependent traits
(Lloyd *et al.* 2013; Osnas *et al.* 2013).
The essence of the problem follows (see Osnas *et al.* 2013 and Supporting
Information Section 1 for details). Let X represent the whole-leaf value of a purely area-
dependent trait; i.e., X is proportional to leaf area, where X could be the photosynthetic
capacity of the entire leaf (units = moles CO_2 fixed per-unit time), the amount of nitrogen in
the entire leaf (units = grams of nitrogen), etc. The mass-normalized trait value would then be
X/Mass, which (assuming X is proportional to leaf area) is proportional to Area/Mass =
LMA^{-1}. This simplistic example shows that if traits are actually area-dependent, mass
normalization can introduce spurious correlations; i.e., in this example, X/Mass is
proportional to LMA^{-1}, even though X is assumed to depend only on leaf area. Thus, the
strong correlations observed among LMA and mass-normalized traits (Reich et al. 1997;
Wright et al. 2004) should not be taken as evidence for a single dominant axis of leaf
functional variation (Lloyd *et al.* 2013; Osnas *et al.* 2013).
An important observation suggesting the presence of multiple axes of leaf functional
variation is that trait relationships across species are sometimes inconsistent with those
observed within species. For example, across species in the global flora, LMA has a strong,
positive relationship with LL but a weak relationship with net photosynthetic capacity per-
unit leaf area (A_{area}) (Wright *et al.* 2004). In contrast, intraspecific variation in these traits
across vertical canopy (light) gradients shows the opposite pattern: within species, LMA has
a weak and/or negative relationship with LL (Osada *et al.* 2001; Lusk *et al.* 2008; Russo &
Kitajima 2016) but a strong, positive relationship with A_{area} (Ellsworth & Reich 1993; Kenzo
et al. 2006; Niinemets *et al.* 2014). These contrasting patterns suggest that the causes of LMA
variation may be fundamentally different within vs. among species. Furthermore, variation in

four traits related to photosynthesis and metabolism – net photosynthetic capacity, dark
respiration rate, and concentrations of nitrogen and phosphorus – are primarily mass-
dependent among species within some plant functional groups, and primarily area-dependent
within others (Osnas *et al*., in review). Thus, leaf trait relationships diverge in a number of
intra- and interspecific comparisons, suggesting the presence of multiple axes of functional
variation that can combine in different ways to create a variety of patterns.
A conceptual model that might explain these divergent patterns is to view LMA as
being comprised of two primary components – 'photosynthetic' leaf mass (LMAp, which
determines photosynthetic capacity and associated metabolic activity, such as phloem-
loading) and structural leaf mass (LMAs, which determines toughness and thus potential LL)
(Osnas *et al*., in review). This conceptual model can potentially explain the above divergent
trait patterns, because variation in LMAp leads to mass-dependence of photosynthetic and
metabolic traits, whereas variation in LMAs leads to area-dependence of these same traits
(Supporting Information, Section 2). Thus, the degree of area- vs. mass-dependence in a
given leaf assemblage should, according to this conceptual model, depend on the relative
contributions of LMAp and LMAs to total LMA variation in the assemblage.
A challenge to implementing the above conceptual model is the absence of direct
measurements of LMAp and LMAs. Although some leaf mass components are clearly
associated with certain functions (e.g., chloroplasts perform photosynthesis), it is difficult to
neatly partition some leaf mass components to specific functions. For example, thick cell
walls provide structural toughness and allow for long LL (Kitajima *et al.* 2012), but a leaf
without any cell wall mass would lack the biomechanical support needed to intercept light
and perform photosynthesis. Thus, it is difficult to determine how much of the cell wall mass
present in a leaf was constructed for photosynthesis and associated metabolic functions, vs.
how much additional cell wall mass was constructed for the purpose of prolonging LL.
Despite this challenge, it should be possible to use multivariate trait datasets to statistically
partition LMA into LMAp and LMAs, because statistical relationships among net
photosynthetic capacity (A_{max}), dark respiration rate (R_{dark}), LMA, and LL should hold clues
regarding the allocation of leaf mass to photosynthetic vs. structural functions.
In this paper, we explore the above conceptual model using leaf trait data from two
tropical forest sites (sun and shade leaves from wet and dry sites in Panama) and the
GLOPNET global leaf traits database (Wright *et al.* 2004). For simplicity, our analysis
assume that LMA is equal to the sum of separate additive components, LMAp and LMAs.
The goal of our analysis is to evaluate if the conceptual model described above can explain
divergent patterns in leaf trait data, and if so, to use the model to elucidate the causes of these
divergent patterns. First, we describe a statistical modeling framework to estimate LMAp and
LMAs, and we use simulations to show that the model yields robust results that are not prone
to statistical artifacts. Then, we ask the following questions: (1) Does decomposing LMA into
photosynthetic and structural components lead to improved predictions of A_{max}, R_{dark}, and
LL? (2) What is the relative importance of LMAp vs. LMAs in explaining variation in LMA
within and among species? (3) Do LMAp and LMAs differ between evergreen and deciduous
species, and between sun and shade leaves? and (4) How are measurable leaf photosynthetic
and structural traits (e.g., concentrations of nitrogen and cellulose) related to LMAp and
LMAs?
Our model is conceptually similar to that of Shipley *et al.* (2006), which explained the
LES in terms of a tradeoff between the volume of a leaf allocated to cell wall vs. protoplasm.
An important difference between our analysis and that of Shipley *et al.* (2006) is that our goal
is to understand divergent trait patterns within vs. among species, whereas Shipley *et al.*
(2006) focused only on interspecific variation. Because we treat LMA as the sum of LMAp
and LMAs without considering how these mass components are packaged within a leaf, our

model ignores the effects of leaf anatomy. For example, a given cell wall mass per leaf area
may be packaged into a large surface area with thin walls (high mesophyll conductance and
A_{max}, but low structural toughness) or a small surface area with thick walls (low mesophyll
conductance and A_{max}, but high structural toughness) (Onoda *et al.* 2017). Despite its lack of
anatomical detail, our model explains much of the inter- and intraspecific variation in A_{max},
R_{dark}, and LL (see Results) and provides novel insights into the causes of divergent patterns of
trait variation within vs. among species (see Discussion).
155
Material and Methods
Model overview
We developed a statistical modeling framework to partition LMA into additive LMAs and
LMAp components (Box 1 and Supplementary Section 3). For sample i (where 'sample'
refers to a species, or a species × canopy position combination; see Datasets below), we
partition LMA_i into $LMAp_i = f_i \times LMA_i$ and $LMAs_i = (1 - f_i) \times LMA_i$ by estimating a latent
variable f_i. The latent variables f_i are not directly observed, but they can be constrained by
available data using Bayesian methods (Bishop 2006; Gelman & Hill 2006). For example,
posterior distributions for $LMAp_i$ should tend to converge on high values for leaves with high
A_{area}, and posterior distributions for $LMAs_i$ should tend to converge on high values for leaves
with high LL. Given the large number of free parameters (i.e., one latent variable per leaf
sample), it is possible for the model to over-fit the data, which could lead to spurious
inferences. Therefore, we performed tests with randomized data (see below) and with
simulated data to evaluate model performance under a range of conditions (Supporting
Information, Section 4). Tests with simulated data suggest that our modeling approach is
robust and not prone to producing artefactual results.
172
Datasets
To fit the models described in Box 1, we used LMA (g m^{-2}), net photosynthetic capacity per
unit leaf area (A_{area}; μmol s^{-1} m^{-2}), dark respiration rate per unit leaf area (R_{area}; μmol s^{-1} m^{-2})
and LL (months) from the GLOPNET global leaf traits database (Wright *et al.* 2004) and
from two tropical forest sites in Panama: Monumental Natural Metropolitano (MNM, "dry
site") and Bosque Protector San Lorenzo (SL, "wet site"). The Panama data include leaves
sampled at two canopy positions ("sun": full sun at the top of the canopy; and "shade": well
shaded, sampled within 2 m of the forest floor) from trees within reach of a canopy crane at
each site. The dry MNM site is a semi-deciduous coastal Pacific forest with a 5-month dry
season from December-April and 1740 mm of annual rainfall (Wright *et al.* 2003). The MNM
crane is 40 m tall with a 51 m long boom. The wet SL site is an evergreen Caribbean coastal
forest with 3100 mm of annual rainfall (Wright *et al.* 2003). The SL crane is 52 m tall with a
54 m long boom.
After deleting leaf samples (i.e., database records, which typically average over
multiple individual leaves) that lacked one of the four traits (LMA, A_{area}, R_{area}, or LL), 198
samples for 198 unique species were available for GLOPNET, and 132 samples for 106
unique species were available for Panama (dry and wet sites combined; 26 species sampled in
both sun and shade; no species with all four traits available at both sites). Both datasets
include additional traits that we used to interpret model results, but which were not used to fit
models. These traits include nitrogen and phosphorus content per leaf unit area (N_{area} and
P_{area}; g m^{-2}) in both datasets, leaf habit in GLOPNET (deciduous or evergreen), and cellulose
content per unit area (CL_{area}; g m^{-2}) in Panama.
We fit the Potential LL Model (Box 1, Eq. 5) to both GLOPNET and Panama data.
We fit the Optimal LL Model (Box 1, Eq. 13; see also Supporting Information, Section 3.1)
only to the Panama data because GLOPNET primarily represents interspecific variation

among sun leaves, whereas the Optimal LL Model was motivated by the negative
intraspecific LL-LMA relationship observed in Panama (Osnas *et al*. *in review*) and
elsewhere (Lusk *et al.* 2008).
201
Parameter estimation
We used independent and non-informative prior distributions for all parameters, including
latent variables f_i for each leaf sample. Details on the prior distributions are provided in
Supporting Information, Section 3.2, and an evaluation of alternative priors for the latent
variables is presented in Supporting Information, Section 4. Models were fit using the
Hamiltonian Monte Carlo algorithm (HMC) implemented in Stan (Carpenter *et al.* 2016).
Posterior estimates were obtained from three independent chains of 20,000 iterations after a
burn-in of 10,000 iterations, thinning at intervals of 20. The Stan code use to fit models can
be accessed and downloaded from the *Ecology Letters* website. Convergence of the posterior
distribution was assessed with the Gelman-Rubin statistic with a convergence threshold of
1.1 for all diagnostics (Gelman *et al.* 2014a).
213
Model selection
Alternative LL models (Eqs. 5 and 13; see also Supporting Information, Section 3.1) fit to
Panama data were compared using the WAIC (Watanabe-Akaike information criterion)
(Watanabe 2010; Gelman *et al.* 2014b). WAIC is a predictive information criterion for
Bayesian models, with lower values indicating a more parsimonious model (Gelman *et al.*
2014b). Compared to AIC (Akaike 1973) and DIC (Spiegelhalter *et al.* 2002), WAIC has the
desirable property of averaging over the posterior distribution rather than conditioning on a
point estimate (Gelman *et al.* 2014b).
222
Variance partitioning
We used the following identity to estimate the relative contributions of LMAp and LMAs to
LMA variance, where again LMA = LMAp + LMAs:
$$\mathrm{Var}\,(Y = X1 + X1) = \mathrm{Cov}(Y, X1 + X2) = \mathrm{Cov}(Y, X1) + \mathrm{Cov}(Y, X2) \qquad (15)$$
Thus, the fractions of total LMA variance due to variance in LMAp and LMAs were
determined by the covariances Cov(LMA, LMAp) and Cov(LMA, LMAs), respectively,
taken as proportions of the total variance Var(LMA). Tests with simulated data confirm that
our modeling approach can effectively partition LMA variance into LMAp and LMAs
components (Supporting Information, Section 4).
232
Randomized LMA datasets
Because our statistical approach includes one latent variable f_i to partition LMA into LMAp
and LMAs for each leaf sample, one might expect a good match between predictions and
observations simply due to the large number of free parameters, whether or not the model
captured important biological mechanisms. To evaluate model performance while controlling
for the number of free parameters, we compared the fit of our model to observed and
randomized datasets. Specifically, we generated randomized datasets by randomizing LMA
values across species, while maintaining the original (non-randomized) data for A_{area}, R_{area}
and LL. Thus, the randomized LMA datasets had zero expected covariance between LMA
and other traits, but maintained the observed covariances among A_{area}, R_{area} and LL. If the fit
between model and data depended primarily on the number of parameters or other model
assumptions, then we would expect similar fits for the original (non-randomized) and
randomized LMA datasets. For the Panama dataset, which includes two sites (wet and dry)
and two canopy strata (sun and shade), we also considered constrained randomizations in
which LMA was randomized across canopy strata within each site. We generated 10

randomized datasets for each analysis. Analyzing a larger number of randomized datasets
(e.g., 1000) was impractical given our computationally intensive Bayesian modeling
approach.
To compare model results obtained from the observed (non-randomized) dataset to
those obtained from the randomized datasets, we calculated the standardized effect size (SES)
for correlations of interest (e.g., between LMAp and A_{area}, or between LMAs and LL) as SES
= $(r_{obs} - r_{rand})/SD_{rand}$, where r_{obs} is the Pearson correlation coefficient of interest for the
observed dataset, and r_{rand} and SD_{rand} are the mean and standard deviation, respectively, of the
corresponding correlation coefficients for the randomized datasets. Under the null hypothesis
of equal correlations for the observed and randomized data, the distribution of SES is
approximately standard normal (Gotelli & Rohde 2002), which we assumed when calculating
P-values. A significant difference ($P < 0.05$) indicates that the correlation inferred from the
observed dataset is different from what would be expected to emerge from model
assumptions and free parameters.
262
Results
1. Photosynthetic and structural LMA components (LMAp and LMAs, respectively)
can be accurately estimated.
Tests with simulated data show that our modeling approach can accurately partition LMA
into LMAp and LMAs, except in cases where the assumed relationships (e.g., between
LMAp and A_{area}, and between LMAs and LL) are weak (Supporting Information, Section 4).
However, tests with simulated data also show that the LMA-randomization test we
implemented can effectively diagnose these weak-correlation cases (Supporting Information,
Section 4). Therefore, in all cases presented below where results based on observed data are
significantly different ($P < 0.05$) than results based on randomized data, we assume that
estimates of LMAp and LMAs (Box 1, Eqs. 1-3) are accurate and that the results reflect
meaningful patterns in the data.
275
2. Nearly all leaf dark respiration is associated with photosynthetic leaf mass.
Photosynthetic mass (LMAp) accounted for nearly all leaf dark respiration; i.e., estimated
dark respiration rate per-unit structural mass (r_s) was close to zero in analyses of both
GLOPNET and Panama data (Table S5.1). Thus, although building costs are likely similar for
different leaf chemical components and tissues (Williams *et al.* 1989; Villar & Merino 2001),
our results suggest that leaf mass associated with photosynthetic function accounts for nearly
all leaf maintenance respiration.
283
3. Decomposing LMA into photosynthetic and structural components leads to improved
predictions of A_{area}, R_{area} and LL.
For the GLOPNET global dataset, A_{area} had a strong positive correlation with LMAp, a weak
negative correlation with LMAs, and a non-significant correlation with total LMA (Figs. 1a-
c; Table S5.2). A_{area} was also positively correlated with LMAp in the randomized datasets (in
which LMA was randomized among leaves in GLOPNET; see gray symbols in Fig. 1), but
these correlations were weaker than those obtained from the observed data ($P < 0.001$ for the
null hypothesis of equal correlation in observed and randomized datasets). R_{area} in GLOPNET
also had a strong positive correlation with LMAp, which was stronger than the correlation
between R_{area} and LMAp in the randomized datasets ($P < 0.001$), or between R_{area} and either
LMA or LMAs (Figs. 1d-f). Finally, LL in GLOPNET had a strong positive correlation with
LMAs, which was stronger than the correlation between LL and LMAs from the randomized
datasets ($P < 0.001$) or between LL and either LMA or LMAp (Figs. 1g-i).

For the Panama dataset, we evaluated multiple models due to the availability of both
sun and shade leaves. The Optimal LL Model (Box 1, Eq. 13) fit the data better than the
Potential LL Model (Box 1, Eq. 5) according to the Watanabe-Akaike information criterion
(WAIC; Table S5.1). Including site effects in the Optimal LL Model further improved the
WAIC but led to similar parameter estimates and inferences (Table S5.1-S5.3). All Panama
results we report are for the Optimal LL Model with site effects unless stated otherwise. A_{area}
and R_{area} had stronger and more positive correlations with LMAp than with LMA or LMAs
(Figs. 2a-f and Table S5.3). LL was not significantly correlated with LMA when all leaves
were combined, but was strongly correlated with LMA for shade leaves at the dry site (Fig.
2g). The correlation between LL and LMAp was similar for observed and randomized
datasets (Fig. 2h; Table S5.3), implying a lack of meaningful correlation between LL and
LMAp in the Panama data. The correlation between LL and LMAs was weaker in observed
than in randomized data (Fig. 2i) because the LL vs. LMAs relationship ignores important
factors in the Optimal LL Model (e.g., effects of light on realized photosynthetic rates and
thus optimal LL). The correlation between predicted and observed LL was higher in the
Optimal LL Model ($r = 0.81$) than for randomized datasets ($P < 0.001$), and also stronger
than for the Potential LL Model ($r = 0.38$) (Fig. 3; Table S5.4).

4. Most interspecific LMA variation is due to variation in structural leaf mass, not
photosynthetic leaf mass.
LMAs accounted for the majority of interspecific variation in LMA among all leaves in
GLOPNET (74.7%), among sun leaves in Panama (57.8%), and among shade leaves in
Panama (86.0%) (Fig. S5.1).

5. Evergreen leaves have greater LMAs than deciduous leaves, and sun leaves have both
greater LMAp and LMAs than shade leaves.
In the GLOPNET dataset, evergreen leaves had significantly higher LMAs than deciduous
leaves, but the two groups had similar LMAp (Fig. 4). Thus, the higher total LMA in
evergreen leaves in GLOPNET was primarily due to differences in LMAs (which comprised
a greater fraction of LMA in evergreen than in deciduous leaves; Fig. S5.2a). Similar results
were obtained for the Panama dataset (evergreen leaves have higher LMAs but similar LMAp
compared to deciduous leaves; Fig. S5.2b). In the Panama dataset, both LMAp and LMAs
were significantly higher in sun leaves than in shade leaves (Fig. 4). Thus, in contrast to
interspecific variation in LMA (which is primarily driven by variation in LMAs; Fig. S5.1),
intraspecific variation in LMA reflects changes in both LMAs and LMAp.

6. Nitrogen and phosphorus per-unit leaf area are strongly correlated with LMAp, and
cellulose per-unit leaf area is strongly correlated with LMAs. In the GLOPNET dataset,
N_{area} and P_{area} had strong positive correlations with LMAp, but only weak correlations with
LMAs (Fig. S5.4). Similarly, in the Panama dataset, N_{area} and P_{area} had strong positive
correlations with LMAp, but were not correlated with LMAs (Fig. 5). In contrast, cellulose
per-unit leaf area (CL_{area}), which was available for the Panama dataset but not for GLOPNET,
had a strong positive correlation with LMAs, and a weak positive correlation with LMAp
(Fig. 5). CL_{area} was more strongly correlated with LMA than with LMAp or LMAs, but sun
and shade leaves aligned along a common CL_{area}-LMAs relationship, as opposed to being
offset for LMA and LMAp (Figs. 5g-i). The above correlations (N_{area} and P_{area} with LMAp,
and CL_{area} with LMAs) were stronger than those based on randomized datasets ($P < 0.001$ in
all cases, Fig. 5 and Fig. S5.4).

Discussion

Our analyses demonstrate that decomposing LMA variation into separate photosynthetic and
structural components (LMAp and LMAs, respectively) leads to improved predictions of
photosynthetic capacity (A_{max}), dark respiration rate (R_{dark}), and leaf lifespan (LL), as well as
clear relationships with traits used for independent model evaluation (nitrogen, phosphorus,
and cellulose concentrations). Tests with simulated data suggest that our results are robust
and reflect meaningful, previously unreported patterns in leaf trait data. Below, we elaborate
on the insights gained from our analysis and the implications of our results for the
representation of leaf functional diversity in global ecosystem models.
Decomposing LMA into LMAp and LMAs provides insights into why interspecific
variation in leaf traits related to photosynthesis and metabolism are primarily area-dependent
(i.e., primarily independent of LMA when expressed per-unit area) rather than mass-
dependent (see Supporting Information, Section 1 and 2 and Osnas *et al.* 2013 for explanation
of trait mass- vs. area-dependence). Specifically, our results suggest that most interspecific
LMA variation across the global flora and across tropical tree species in Panama is due to
variation in leaf mass components that contribute to LL but not to A_{max} (Fig. S5.1). An
illustrative case is the comparison of evergreen and deciduous leaves in the global flora:
according to our model results, evergreen leaves have greater LMAs (and thus greater LL)
but similar LMAp (and thus similar A_{area}) compared to deciduous leaves (Figs. 1 and 4). This
comparison illustrates why interspecific LMA variation is largely unrelated to area-based
A_{max}, which in turn explains trait area-dependence (Supporting Information, Section 2).
Consistent with this explanation, the assemblage we examined where LMAs accounted for
the highest fraction of total LMA variation (86% for Panama shade leaves; Fig. S5.1) is also
the assemblage with the highest degree of trait area-dependence (Osnas *et al.* in review).
Decomposing LMA also provides insights as to why intraspecific patterns of trait
variation differ from those observed across species. In contrast to interspecific LMA
variation, our analysis suggests that LMAp contributes half or more of the intraspecific
increase in LMA from shade to sun (Figs. 4 and S5.2). The increase in LMAp from shade to
sun – which likely reflects an increase in the size and number of palisade mesophyll cells
with increasing light availability (Onoda *et al.* 2008; Terashima *et al.* 2011) – is also
associated with an increase in LMAs from shade to sun (Fig. 4). This positive covariance
between LMAp and LMAs within species means that per-area values of LMAp-dependent
traits (e.g., A_{area}) have a strong, positive relationship with total LMA, which implies trait
mass-dependence (Supporting Information, Section 1-2). In contrast, because interspecific
variation in LMA is due primarily to variation in LMAs rather than LMAp, LMAp-dependent
traits (when expressed per-unit area) are largely independent of LMA across species, which
implies trait area-dependence (Supporting Information, Sections 1-2).
Finally, decomposing LMA provides insights into the intraspecific LL 'counter-
gradient' (Osada *et al.* 2001; Lusk *et al.* 2008; Russo & Kitajima 2016), whereby LL
decreases as LMA increases from shade to sun within species, which contrasts with positive
interspecific LL-LMA relationships within a single light environment. Decomposing LMA
into LMAp and LMAs suggests that the decline in photosynthetic rate from sun to shade
(which increases optimal LL from a carbon-gain perspective; Kikuzawa 1991) is sufficient to
produce an intraspecific counter-gradient, because a counter-gradient occurs at the dry
Panama site (6 vs. 8 months for sun vs. shade LL means) despite no change in the structural
leaf mass fraction (LMAs/LMA) from sun to shade (see LMAp/LMA values in Fig. S5.2c,
with LMAs/LMA = 1 – LMAp/LMA). However, the counter-gradient is stronger at the wet
Panama site (11 vs. 31 months for sun vs. shade LL means), where LMAs/LMA is
significantly higher for shade leaves than for sun leaves (see LMAp/LMA in Fig. S5.2c).
Thus, at the wet Panama site, changes in LMAs/LMA from sun to shade appear to strengthen
the counter-gradient. The potential for LMAs/LMA to enhance the counter-gradient is

corroborated by data on cellulose percent mass, which is similar in shade and sun at the
Panama dry site (Fig. S5.5b; as observed by Lusk *et al.* 2010), but higher in shade than sun at
the wet site (Fig. S5.5b; as observed by Kitajima *et al.* 2016). In summary, the decrease in
optimal LL with increasing light (Kikuzawa 1991) appears sufficient by itself to create a LL
counter-gradient, but the counter-gradient appears to be enhanced in some assemblages by
greater structural mass fractions in shade vs. sun leaves.
The improved predictions and understanding provided by decomposing LMA into
photosynthetic and structural components challenge the view that leaf functional diversity
can be accurately represented by a single leaf economics spectrum (LES) axis (Wright *et al.*
2004). Lloyd *et al.* (2013) argued that the apparent dominance of a single LES axis is an
artifact of expressing area-dependent leaf traits on a per-mass basis, and Osnas *et al.* (2013)
demonstrated that across the global flora, traits related to photosynthesis and metabolism are
indeed area-dependent. Intraspecific patterns in trait variation, which contrast with
interspecific patterns, pose additional challenges for a one-dimensional view of leaf
functional diversity. Our analysis shows that considering two primary axes of leaf trait
variation (photosynthesis and structure) provides improved quantitative predictions and
insights compared to a single LES. Our results point to a simple two-dimensional framework
for representing leaf functional diversity in global ecosystem models: a LMAp axis that
determines A_{max}, accounts for nearly all R_{dark} (see r_p and r_s estimates in Table S5.1), and
affects LL as predicted by optimality theory (Kikuzawa 1992); and a LMAs axis that
determines potential LL through its effects on leaf toughness (Kitajima *et al.* 2012). In the
datasets we analyzed (the global flora and tropical trees in Panama), these two axes are only
weakly correlated with each other (Fig. S5.6), which suggests that trait-based approaches to
global ecosystem modeling (Scheiter *et al.* 2013; Wullschleger *et al.* 2014) could consider
these as independent axes.
Our statistical decomposition of LMA into photosynthetic and structural components
provides important insights (summarized above), but additional insights and accuracy could
be gained by a more mechanistic modeling approach. For example, John *et al.* (2017)
decomposed interspecific LMA variation into anatomical components such as the size,
number of layers, and mass density of cells in different leaf tissues. If such detailed
information became available for a large number of leaves, representing both intra- and
interspecific variation, it should be possible to quantify how these anatomical traits scale up
to leaf-level A_{max}, R_{dark}, and LL. A simpler alternative would be to modify our model (Box 1)
to account for variation in cell wall thickness (T_{CW}): for a given cell size, increasing T_{CW}
would lead to an increase in lamina density, cellulose per volume, toughness, and LL
(Kitajima and Poorter 2010; Kitajima *et al.* 2012, 2016), and a decrease in mesophyll
conductance and A_{max} (Evans *et al.* 2009; Terashima *et al.* 2011; Onoda *et al.* 2017).
434
Conclusions
It is widely recognized that LMA variation is associated with multiple tissues and functions,
including metabolically active mesophyll that largely determines photosynthetic capacity, as
well as structural and chemical components that contribute primarily to leaf toughness and
defense (Roderick *et al.* 1999; Shipley *et al.* 2006; Lusk *et al.* 2010). It should not be
surprising then, that partitioning LMA into photosynthetic and structural components yields
enhanced predictions and improved understanding of patterns of leaf trait variation both
within and among species. Yet for over a decade, the vast literature on leaf traits has been
strongly influenced by the view that leaf trait variation can usefully be represented by a single
LES axis. Our results provide quantitative evidence that this one-dimensional view of leaf
trait variation is insufficient, and our model provides a biological explanation for previous
statistical analyses that have demonstrated area-dependence of leaf traits across species

(Osnas *et al*. 2013), while also explaining mass-dependence within species. Our results suggest that most of the interspecific variation in LMA concerns structural mass that contributes to LL but not to photosynthetic capacity, and that structural and photosynthetic LMA components are only weakly correlated across species. Thus, strong interspecific correlations between LMA and mass-normalized photosynthetic capacity (and related traits, such as respiration rate, and nitrogen and phosphorus concentrations) are likely driven by mass-normalization itself, rather than any functional dependence of these photosynthetic and metabolic traits on LMA (Lloyd *et al*. 2013, Osnas *et al*. 2013). In contrast, intraspecific variation in LMA is driven by coordinated changes in structural and photosynthetic mass components, which explains why mass-normalized photosynthetic and metabolic traits vary little from sun to shade within species (Aranda *et al.* 2004; Poorter *et al.* 2006; Niinemets *et al.* 2014).

Acknowledgments
We thank Jonathan Dushoff for statistical advice and Martijn Slot for helpful comments that improved the paper. Mirna Samaniego and Milton Garica provided indispensable assistance in data collection. We thank the Smithsonian Tropical Research Institute (STRI), the Tropical Canopy Biology Program at STRI and the Andrew W. Mellon Foundation for supporting this work. MK was supported by a Postdoctoral Fellowship for Research Abroad from the Japan Society for the Promotion of Science.

References
1.Akaike, H. (1973). Information theory and an extension of the maximum likelihood principle. In: (eds. Petrov, B.N. & Csaki, F.). Presented at the Second International Symposium on Information Theory, Akadémiai Kiado, pp. 267–281.
2.Aranda, I., Pardo, F., Gil, L. & Pardos, J.A. (2004). Anatomical basis of the change in leaf mass per area and nitrogen investment with relative irradiance within the canopy of eight temperate tree species. *Acta Oecologica*, 25, 187–195.
3.Bishop, C.M. (2006). Pattern Recognition and Machine Learning.
4.Bonan, G.B., Levis, S., Kergoat, L. & Oleson, K.W. (2002). Landscapes as patches of plant functional types: An integrating concept for climate and ecosystem models. *Global Biogeochem. Cycles*, 16, 5–1–5–23.
5.Carpenter, B., Gelman, A., Hoffman, M., Lee, D. & Goodrich, B. (2016). Stan: A probabilistic programming language. *J Stat Softw*.
6.Ellsworth, D.S. & Reich, P.B. (1993). Canopy structure and vertical patterns of photosynthesis and related leaf traits in a deciduous forest. *Oecologia*.
7.Evans, J.R., Kaldenhoff, R., Genty, B. & Terashima, I. (2009). Resistances along the CO2 diffusion pathway inside leaves. *Journal of Experimental Botany*, 60, 2235–2248.
8.Gelman, A. & Hill, J. (2006). Data analysis using regression and multilevel/hierarchical models.
9.Gelman, A., Carlin, J.B., Stern, H.S., Dunson, D.B., Vehtari, A. & Rubin, D.B. (2014a). *Bayesian Data Analysis, Third Edition*. Taylor & Francis.
10.Gelman, A., Hwang, J. & Vehtari, A. (2014b). Understanding predictive information criteria for Bayesian models. *Statistics and Computing*, 24, 997–1016.
11.Gotelli, N. & Rohde, K. (2002). Co-occurrence of ectoparasites of marine fishes: a null model analysis. *Ecol Letters*, 5, 86–94.
12.John, G.P., Scoffoni, C., Buckley, T.N., Villar, R., Poorter, H. & Sack, L. (2017). The anatomical and compositional basis of leaf mass per area. *Ecol Letters*.
13.Kenzo, T., Ichie, T., Watanabe, Y., Yoneda, R., Ninomiya, I. & Koike, T. (2006). Changes in photosynthesis and leaf characteristics with tree height in five dipterocarp species in

a tropical rain forest. *Tree Physiol.*, 26, 865–873.

14.Kikuzawa, K. (1991). A cost-benefit analysis of leaf habit and leaf longevity of trees and
their geographical pattern. *Am Nat*, 1250–1263.

15.Kikuzawa, K., Onoda, Y., Wright, I.J. & Reich, P.B. (2013). Mechanisms underlying
global temperature-related patterns in leaf longevity. *Global Ecol Biogeogr*.

16.Kikuzawa, K., Shirakawa, H., Suzuki, M. & Umeki, K. (2004). Mean labor time of a leaf.
Ecol Res, 19, 365–374.

17.Kitajima, K., Llorens, A.-M., Stefanescu, C., Timchenko, M.V., Lucas, P.W. & Wright,
S.J. (2012). How cellulose-based leaf toughness and lamina density contribute to long
leaf lifespans of shade-tolerant species. *New Phytol*, 195, 640–652.

18.Kitajima, K., Wright, S.J. & Westbrook, J.W. (2016). Leaf cellulose density as the key
determinant of inter- and intra-specific variation in leaf fracture toughness in a species-
rich tropical forest. *Interface Focus*, 6, 20150100–9.

19.Lloyd, J., Bloomfield, K., Domingues, T.F. & Farquhar, G.D. (2013). Photosynthetically
relevant foliar traits correlating better on a mass vs an area basis: of ecophysiological
relevance or just a case of mathematical imperatives and statistical quicksand? *New*
Phytol, 199, 311–321.

20.Lusk, C., Onoda, Y., Kooyman, R. & Gutiérrez-Girón, A. (2010). Reconciling species-
level vs plastic responses of evergreen leaf structure to light gradients: shade leaves
punch above their weight. *New Phytol*, 186, 429–438.

21.Lusk, C.H., Reich, P.B., Montgomery, R.A., Ackerly, D.D. & Cavender-Bares, J. (2008).
Why are evergreen leaves so contrary about shade? *Trends Ecol Evol*, 23, 299–303.

22.Niinemets, Ü., Keenan, T.F. & Hallik, L. (2014). A worldwide analysis of within-canopy
variations in leaf structural, chemical and physiological traits across plant functional
types. *New Phytol*, 205, 973–993.

23.Onoda, Y., Schieving, F. & Anten, N.P.R. (2008). Effects of light and nutrient availability
on leaf mechanical properties of Plantago major: a conceptual approach. *Ann Bot*, 101,
727–736.

24.Onoda, Y., Wright, I.J., Evans, J.R., Hikosaka, K., Kitajima, K., Niinemets, Ü., *et al.*
(2017). Physiological and structural tradeoffs underlying the leaf economics spectrum**.**
New Phytol. doi:10.1111/nph.14496

25.Osada, N., Takeda, H., Furukawa, A. & Awang, M. (2001). Leaf dynamics and
maintenance of tree crowns in a Malaysian rain forest stand. *J Ecol*, 89, 774–782.

26.Osnas, J.L.D., Lichstein, J.W., Reich, P.B. & Pacala, S.W. (2013). Global Leaf Trait
Relationships: Mass, Area, and the Leaf Economics Spectrum. *Science*, 340, 741–744.

27.Poorter, H., Pepin, S., Rijkers, T., de Jong, Y., Evans, J.R. & Körner, C. (2006).
Construction costs, chemical composition and payback time of high- and low-irradiance
leaves. *Journal of Experimental Botany*, 57, 355–371.

28.Reich, P.B. (2014). The world-wide "fast-slow" plant economics spectrum: a traits
manifesto. *J Ecol*, 102, 275–301.

29.Reich, P.B., Walters, M.B. & Ellsworth, D.S. (1997). From tropics to tundra: global
convergence in plant functioning. *Proceedings of the National Academy of Sciences*, 94,
13730–13734.

30.Roderick, M.L., Berry, S.L., Noble, I.R. & Farquhar, G.D. (1999). A theoretical approach
to linking the composition and morphology with the function of leaves. *Functional*
Ecology, 13, 683–695.

31.Russo, S.E. & Kitajima, K. (2016). The Ecophysiology of Leaf Lifespan in Tropical
Forests: Adaptive and Plastic Responses to Environmental Heterogeneity. In: *Tropical*
Tree Physiology, Tropical Tree Physiology: Adaptations and Responses in a Changing
Environment (eds. Goldstein, G. & Santiago, S.L.). Springer International Publishing,

Cham, pp. 357–383.

32.Scheiter, S., Langan, L. & Higgins, S.I. (2013). Next-generation dynamic global
vegetation models: learning from community ecology. *New Phytol*, 198, 957–969.

33.Shipley, B., Lechowicz, M., Wright, I. & Reich, P. (2006). Fundamental trade-offs
generating the worldwide leaf economics spectrum. *Ecology*, 87, 535–541.

34.Spiegelhalter, D.J., Best, N.G., Carlin, B.P. & Van Der Linde, A. (2002). Bayesian
measures of model complexity and fit. *Journal of the Royal Statistical Society: Series B*
(Statistical Methodology), 64, 583–639.

35.Terashima, I., Hanba, Y.T., Tholen, D. & Niinemets, Ü. (2011). Leaf functional anatomy
in relation to photosynthesis. *Plant Physiology*, 155, 108–116.

36.Villar, R. & Merino, J. (2001). Comparison of leaf construction costs in woody species
with differing leaf life-spans in contrasting ecosystems. *New Phytol*.

37.Watanabe, S. (2010). Asymptotic Equivalence of Bayes Cross Validation and Widely
Applicable Information Criterion in Singular Learning Theory. *The Journal of Machine*
Learning Research, 11, 3571–3594.

38.Westoby, M. & Wright, I.J. (2006). Land-plant ecology on the basis of functional traits.
Trends Ecol Evol, 21, 261–268.

39.Williams, K., Field, C.B. & Mooney, H.A. (1989). Relationships among leaf construction
cost, leaf longevity, and light environment in rain-forest plants of the genus Piper. *Am*
Nat.

40.Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F., *et al.*
(2004). The worldwide leaf economics spectrum. *Nature*, 428, 821–827.

41.Wright, S.J., Horlyck, V., Basset, Y., Barrios, H., Bethancourt, A., Bohlman, S.A., *et al.*
(2003). Tropical Canopy Biology Program, Republic of Panama. In: *Studying Forest*
Canopies from Above: The International Canopy Crane Network (eds. Basset, Y.,
Horlyck, V. & Wright, S.J.). Panama, pp. 137–155.

42.Wullschleger, S.D., Epstein, H.E., Box, E.O., Euskirchen, E.S., Goswami, S., Iversen,
C.M., *et al.* (2014). Plant functional types in Earth system models: past experiences and
future directions for application of dynamic vegetation models in high-latitude
ecosystems. *Ann Bot*, 114, 1–16.

Supporting information
Additional Supporting Information may be found online in the supporting information tab for
this article.
582

583
Figure 1. Observed and estimated leaf-trait relationships in the global GLOPNET dataset.
Leaf life span (LL), net photosynthetic rate per unit leaf area (A_{area}), and dark respiration rate
per unit leaf area (R_{area}) are plotted against observed LMA and estimates (posterior means) of
photosynthetic and structural LMA components (LMAp and LMAs, respectively). Gray
symbols show 1 of 10 randomized datasets, in which LMA was randomized among all leaves
in the GLOPNET dataset we analyzed (which includes all leaves for which LMA, A_{area}, R_{area},
and LL were available). Pearson correlation coefficients are for observed LMA (left column)
and posterior means of LMAp (middle column) and LMAs (right column). *P*-values (** *P* <
0.01; *** *P* < 0.001) for LMA test the null hypothesis of zero correlation, and for LMAp and
LMAs test the null hypothesis of equal correlation in observed and randomized datasets (see
'Randomized LMA Datasets' in Methods for details). In some cases, r_{obs} is near zero but
significantly greater than r_{rand} because r_{rand} is negative (see Table S5.2 for details).

596
597 Figure 2. Observed and estimated leaf-trait relationships in the Panama dataset. Estimates are
598 from the Optimal LL Model with site effects (Eq. 13 and Supporting Information Section
599 3.1). Details as in Fig. 1, except that in the randomized dataset shown here (gray symbols),
600 LMA was randomized within sites (wet and dry) across canopy strata (sun and shade).
601 Results for other LL models are summarized in Table S5.3.
602

603
Figure 3. Observed vs. predicted leaf lifespan (LL) in the (a) Potential LL Model (Eq. 5) and
(b) Optimal LL Model with site effects (Eq. 13 and Supporting Information Section 3.1).
Predicted LL values are posterior means. The dashed line indicates the 1:1 relationship. Gray
symbols show predicted LL values from 1 of 10 randomized LMA datasets for each model.
Pearson correlation coefficients are for predictions derived from the observed datasets, and *P*-
values (*** $P < 0.001$) are for the null hypothesis of equal correlation in observed and
randomized datasets (see 'Randomized LMA Datasets' in Methods for details). Results for
other LL models are reported in Table S5.4.

Figure 4. Boxplots comparing leaf mass per area (LMA) and posterior means of
photosynthetic and structural LMA components (LMAp and LMAs, respectively) across
deciduous (D) and evergreen (E) leaves in the GLOPNET dataset (top), and across sites (wet
and dry) and canopy strata (sun and shade) in the Panama dataset (bottom). The center line in
each box indicates the median, upper and lower box edges indicate the interquartile range,
whiskers show 1.5 times the interquartile range, and points are outliers. Groups sharing the
same letters are not significantly different ($P > 0.05$; t-tests). To isolate the effects of
intraspecific variation (i.e., plastic responses to light), the Panama results shown here only
include species for which both sun and shade leaves were available. Qualitatively similar
results were obtained when all Panama species were included (Fig. S5.3). Note that the
vertical axis is on a $\log_{10}$ scale.

Figure 5. Measured traits related to photosynthesis and metabolism traits (nitrogen and phosphorus per-unit leaf area; N_{area} and P_{area}) are strongly correlated with estimates (posterior means) of the photosynthetic LMA component (LMAp), and a measured structural trait (cellulose per-unit leaf area; CL_{area}) is strongly correlated with estimates of the structural LMA component (LMAs) for the Panama dataset. Note that sun and shade leaves align along a single relationship for CL_{area} vs. LMAs, but not for CL_{area} vs. LMA or LMAp. N_{area}, P_{area}, and CL_{area} data were not used to fit the models, and are presented here as independent support for the model results. Analogous results were obtained for N_{area} and P_{area} vs. LMAp for GLOPNET (Fig. S5.4). Gray symbols show 1 of 10 datasets in which LMA was randomized within sites (wet and dry) across canopy strata (sun and shade). Others details as in Fig. 1. Results for other LL models are reported in Table S5.3.

Box 1 Modeling leaf lifespan, photosynthetic capacity, and dark respiration in relation to
photosynthetic and structural leaf mass components
We assume that the sum of photosynthetic leaf mass per area (LMAp) and structural leaf
mass per area (LMAs) is equal to total observed LMA for leaf sample i (where a 'sample' is a
species in the GLOPNET dataset, or a species × canopy position combination in the Panama
dataset):

$$LMA_i = LMAp_i + LMAs_i \qquad (1)$$
$$LMAp_i = f_i LMA_i \qquad (2)$$
$$LMAs_i = (1 - f_i) LMA_i \qquad (3)$$

where, f_i is the fraction of LMA_i that is comprised of $LMAp_i$. The f_i terms are not directly
observed but can be estimated as latent variables in a Bayesian modeling framework (see
details below). In our model, gross photosynthetic capacity (approximated here as the sum of
A_{area} and R_{area}) is determined by LMAp, LL is determined by LMAs, and total leaf dark
respiration is the sum of respiration components due to photosynthetic and structural tissues:

$$E[\text{gross photosynthesis}] = E[A_{area\,i} + R_{area\,i}] = \alpha LMAp_i = \alpha f_i LMA_i \qquad (4)$$
$$E[LL_i] = \beta_1 LMAs_i = \beta_1 (1 - f_i) LMA_i \qquad (5, \text{Potential LL Model})$$
$$E[R_{area\,i}] = r_p LMAp_i + r_s LMAs_i = r_p f_i LMA_i + r_s (1 - f_i) LMA_i \qquad (6)$$

where, $E[\cdot]$ is the expected value of the variable in brackets, $A_{area\,i}$, $R_{area\,i}$, and LL_i, are,
respectively, the net photosynthetic rate (A_{max}) per unit area, dark respiration rate (R_{dark}) per
unit area, and leaf life span of leaf i; α is net photosynthetic rate per unit photosynthetic
mass; β_1 is leaf lifespan per unit structural mass; and r_p and r_s, are, respiration rates per unit
photosynthetic and structural leaf mass, respectively. Eq. 5 is labeled "Potential LL Model",
because this model assumes that LL is affected only by structural toughness (which we
assume is proportional to LMAs), in contrast to an alternative model based on optimal LL
theory (see Eqs. 12-14 below). The logarithms of gross photosynthesis, LL, and R_{area} are
assumed to have a multivariate normal distribution (MVN):

$$\begin{pmatrix} \ln [A_{area\,i} + R_{area\,i}] \\ \ln [LL_i] \\ \ln [R_{area\,i}] \end{pmatrix} \sim MVN \begin{pmatrix} \mu_{1i} \\ \mu_{2i}, \Sigma \\ \mu_{3i} \end{pmatrix} \qquad (7)$$

$$\mu_{1i} = \ln \alpha + \ln f_i + \ln LMA_i - \frac{\sigma_1^2}{2} \qquad (8)$$

$$\mu_{2i} = \ln \beta_1 + \ln(1 - f_i) + \ln LMA_i - \frac{\sigma_2^2}{2} \qquad (9)$$

$$\mu_{3i} = \ln[r_p f_i LMA_i + r_s (1 - f_i) LMA_i] - \frac{\sigma_3^2}{2} \qquad (10)$$

$$\Sigma = \begin{pmatrix} \sigma_1^2 & \rho_{12}\sigma_1\sigma_2 & \rho_{13}\sigma_1\sigma_3 \\ \rho_{12}\sigma_1\sigma_2 & \sigma_2^2 & \rho_{23}\sigma_2\sigma_3 \\ \rho_{13}\sigma_1\sigma_3 & \rho_{23}\sigma_2\sigma_3 & \sigma_3^2 \end{pmatrix} \qquad (11)$$

where μ_{1-3} are, respectively, the expected values of $\ln[A_{area\,i} + R_{area\,i}]$, $\ln[LL_i]$, and $\ln[R_{area\,i}]$
(from Eqs. 4-6); Σ is the covariance matrix; σ_1, σ_2, and σ_3 are standard deviations of μ_{1-3}; and
ρ_{12}, ρ_{13}, and ρ_{23} are correlation coefficients among μ_{1-3} (e.g., we expect positive covariance
between μ_1 and μ_3 because both include $R_{area\,i}$).

In addition to Eq. 5 (Potential LL Model), we considered an alternative assumption
for the relationship between LL and other traits based on optimal LL theory (Kikuzawa 1991;
Kikuzawa *et al.* 2004; 2013). According to this theory, the optimal leaf lifespan (LL_{opt})
maximizes a leaf's lifetime carbon gain per-unit time, which is determined by potential LL,
net photosynthetic rate, and construction cost per unit leaf area:

$$LL_{opt} = \sqrt{2bC/(a - m)} \qquad (12)$$

where a is the realized (i.e., light-dependent) gross photosynthetic rate per unit leaf area, m is
the realized daily respiration rate per unit leaf area, b is the rate of decline in photosynthetic

capacity with leaf age (which determines potential LL in the Kikuzawa model), and C is the
construction cost per unit leaf area. To implement this Optimal LL Model, we assumed that
(i) potential LL is proportional to LMAs; and (ii) leaf construction cost per unit area (g
glucose per unit leaf area) is proportional to LMA. Assumption (ii) is justified because leaf
construction cost per unit mass (g glucose per unit leaf mass) is strongly conserved (Williams
et al. 1989; Villar & Merino 2001). Thus, Eq. 12 can be written in terms of the traits
considered in our analysis as:

$$\mathrm{E}[\mathrm{LL}_i] = \beta_2\sqrt{\mathrm{LMAs}_i\mathrm{LMA}_i/(\theta_{\mathrm{L}}A_{\mathrm{area}\,i} - R_{\mathrm{area}\,i})}$$

687
$$= \beta_2\sqrt{(1 - f_i)\mathrm{LMA}_i^2/(\theta_{\mathrm{L}}A_{\mathrm{area}\,i} - R_{\mathrm{area}\,i})} \qquad\qquad (13,\ \text{Optimal LL Model})$$

688   where β_2 is a constant, and θ_{L} $(0 < \theta_{\mathrm{L}} < 1)$ is a scaling parameter that accounts for the effects
689   of light availability on the realized photosynthetic rate (Kikuzawa *et al.* 2004) (see
690   Supporting Information Section 3.1.2 for more details). The expected value of the logarithm
691   of LL (μ_{2i} in Eq. 7) for the Optimal LL Model is:

692
$$\mu'_{2i} = \ln\beta_2 + \ln\mathrm{LMA}_i + \frac{1}{2}\left[\ln(1 - f_i) - \ln(\theta_{\mathrm{L}}A_{\mathrm{area}\,i} - R_{\mathrm{area}\,i})\right] - \frac{\sigma'^2_2}{2}. \qquad\qquad (14)$$

We fit the Optimal LL Model to the Panama data (see Methods/Datasets in main text),
which includes both sun and shade leaves for 26 out of 106 total species. We set $\theta_{\mathrm{L}} = 1$ for
sun leaves, and we fit θ_{L} as a single free parameter assigned to all shade leaves. We
considered versions of the Optimal LL Model both with and without an additional parameter
to account for differences in LL between the dry and wet sites due to mechanisms not
included in the model (Supporting Information, Section 3.1).