Aggressiveness as a latent personality trait of domestic dogs: testing local independence and measurement invariance

4   Conor Goold[1*], Ruth C. Newberry[1]

5   [1] Department of Animal and Aquacultural Sciences, Faculty of Biosciences, Norwegian

6   University of Life Sciences, Ås, Akershus, Norway

7   [*] Corresponding author: conor.goold@nmbu.no (CG)

Abstract

Studies of animal personality attempt to uncover underlying or 'latent' personality traits that explain broad patterns of behaviour, often by applying latent variable statistical models (e.g. factor analysis) to multivariate data sets. Two integral, but infrequently confirmed, assumptions of latent variable models in animal personality are: i) behavioural variables are independent (i.e. uncorrelated) conditional on the latent personality traits they reflect (*local independence*), and ii) personality traits are associated with behavioural variables in the same way across individuals or groups of individuals (*measurement invariance*). We tested these assumptions using observations of aggression in four age classes (4 - 10 months, 10 months - 3 years, 3 - 6 years, over 6 years) of male and female shelter dogs (N = 4,743) in 11 different contexts. A structural equation model supported the hypothesis of two positively correlated personality traits underlying aggression across contexts: aggressiveness towards people and aggressiveness towards dogs (comparative fit index: 0.96; Tucker-Lewis index: 0.95; root mean square error of approximation: 0.03). Aggression across contexts was moderately repeatable (towards people: intraclass correlation coefficient (ICC) = 0.479; towards dogs: ICC = 0.303). However, certain contexts related to aggressiveness towards people (but not dogs) shared significant residual relationships unaccounted for by latent levels of aggressiveness. Furthermore, aggressiveness towards people and dogs in different contexts interacted with sex and age. Thus, sex and age differences in displays of aggression were not simple functions of underlying aggressiveness. Our results illustrate that the robustness of traits in latent variable models must be critically assessed before making conclusions about the effects of, or factors influencing, animal personality. Our findings are of concern because

31   inaccurate 'aggressive personality' trait attributions can be costly to dogs, recipients of

32   aggression and society in general.

34   *Key words*: animal personality assessment; agonistic behaviour; shelter dogs;

35   measurement bias; behavioural phenotyping

Introduction

Studies of non-human animal personality demonstrate that animals show relatively

consistent between-individual differences in behaviour, and that the behavioural

phenotype is organised hierarchically into broad behavioural dimensions or personality

traits (e.g. sociability, aggressiveness or boldness) that further exhibit inter-correlations to

form behavioural syndromes (e.g. boldness with aggression; [1–5]). To interpret the

complexity inherent in behavioural phenotypes, personality traits and behavioural

syndromes are frequently inferred using latent variable statistical models [6], which

reduce two or more measured variables (the *manifest* variables) into one or more lower-

dimensional variables (the *latent* variables), following work in human psychology [7–10].

Many animal personality studies use *formative* models, such as principal components

analysis, that construct composite variables comprised of linear combinations of manifest

variables. However, formative models impose only weak assumptions about the

relationships between latent variables and manifest variables [6,11]. For instance,

formative models do not require manifest variables to be correlated with one another or

illustrate internal consistency [11]. Because behavioural variables comprising personality

traits are expected to correlate with each other [4], the utility of formative models to

revealing underlying personality traits has been criticised in both animals [12,13] and

humans [10,11,14,15]. Instead, researchers are increasingly using *reflective* models, such

as factor analysis, including confirmatory approaches such as structural equation

modelling (see [1,16–18]). Reflective models regress measured behaviours on one or

more latent variables, incorporating measurement error and possibilities to compare *a*

priori competing hypotheses [1,16,19].

Whilst reflective models offer a powerful framework to examine the latent variable

structure of animal behaviour [19], they impose certain assumptions on the interpretation

and modelling of latent variables that have received scrutiny in human psychology but

are rarely discussed in studies of animal personality. Two foundational assumptions are

local independence and *measurement invariance*. Local independence implies that

manifest variables should be independent of each other conditional on the latent variables

[20,21]. For example, given a continuous latent variable θ (e.g. boldness) and two binary

manifest variables Y_1 and Y_2 that can take the values 0 and 1, the item response theory

model asserts that $P(Y_1 = 1, Y_2 = 1|\theta) = P(Y_1 = 1|\theta)P(Y_2 = 1|\theta)$. As such, the latent

variables should 'screen off' any covariance between manifest variables. Measurement

invariance implies that the latent variables function the same (i.e. are invariant or

equivalent) in different subsets of a population or in the same individuals through time

[21–25]. In the previous example, this means that the expected values of the manifest

variables Y_1 and Y_2 should remain the same across different groups, π (e.g. sex or

different populations), for any fixed value of the latent variable θ_x e.g. $E(Y_1 \mid \theta_x) =$

$E(Y_1 \mid \theta_x, \pi)$. For studies of personality, violations of local independence or measurement

invariance highlight instances where the personality traits do not completely explain

variation in the manifest variables, which may lead to misleading conclusions about the

differences between individuals as a function of trait scores [25–27].

The goal of this study was evaluate local independence and measurement invariance in

behavioural data on domestic dogs (*Canis lupus familiaris*). Dog personality has been of

scientific interest for decades [28–30], both to predict the behaviour of dogs at future

time points [31] and to elucidate behavioural traits pertinent to dogs' domestication

history [32–35]. Research on personality in dogs has led to different numbers and

composition of hypothesised personality traits with little consensus on how such traits

should be compared within and between studies [36–38]. Dog personality studies

frequently involve collection of data on a wide range of behaviours and, as a result, latent

variable models are popular to reduce behavioural data into personality traits or

dimensions [29]. Importantly, the predictive value of personality assessments in dogs has

been inconsistent [31,39–43], perhaps most prominently in shelter dog personality

assessments (e.g. see [31] for a review). Assessments of aggression are of particular

concern, where aggression has been divided into different aggressiveness traits, such as

owner-, stranger-, dog- or animal-directed factors [29,37,44,45]. Improving inference

about aggressiveness in dogs is important because dog bites are a serious public health

concern [46], especially for animal shelters rehoming dogs to new owners, and aggressive

behaviour is undesirable to many organisations using dogs for various working roles [47].

Evaluating local independence and measurement invariance could help refine applied

personality assessments on dogs. Local independence may be violated in standardised test

batteries (a common assessment method; [48–50]) because the sequential administration

of different behavioural subtests means that how dogs responds to one sub-test may

influence their subsequent behavioural responses, as well as the responses of the dog

handlers [31]. Identifying local independence could, thus, highlight which sub-tests can

be interpreted as providing independent information. Local independence is also relevant

to the development and analysis of dog personality questionnaires completed by dog

owners, because the order in which the questions are presented or redundancy in the

content of questions can lead to dependencies between participant responses not

explained by the questionnaire's intended focus on the dog's behaviour [51].

Scientists are also concerned with understanding personality differences in dogs across a

variety of conditions, including ontogeny, age, sex, breed and neuter status (e.g. [37, 42,

52–54]). Evaluating measurement invariance in personality assessments would allow

researchers to confirm whether differences between individuals or groups of individuals

in personality assessments reflect credible differences in personality trait scores or

whether additional, unaccounted for factors are driving the differences. While it may be

unrealistic for measurement invariance to hold in all instances, it is important to establish

whether it holds for personality traits across basic biological variables such as age and

sex, which are generally applicable to dog populations undergoing personality assessment

and have previously been found to show interactions with personality traits, including

playfulness, sociability, curiosity and aggressiveness [33, 55]. However, apart from van

den Berg *et al*. [18] who assessed measurement invariance across breed groups, no

studies have confirmed measurement invariance or local independence for personality

traits.

In this paper, we assessed local independence and measurement invariance of

aggressiveness in shelter dogs using a large sample of data on inter-context aggressive

behaviour. First, we decomposed observations of aggression towards people and dogs

across contexts into separate aggressiveness traits. Secondly, we assessed whether

aggression in different contexts remained associated beyond that explained by latent

levels of aggressiveness, testing local independence. Thirdly, we investigated whether the

probability of aggression in different contexts assumed to be underpinned by the same

aggressiveness trait was measurement invariant with respect to sex and age groups.

Materials & Methods

Subjects

Observational data on the occurrence of aggression in 4,743 dogs were gathered from

Battersea Dogs and Cats Home's (UK) observational and longitudinal dog behaviour

assessment records (Table 1). The data were from a sample of dogs (N = 4,990) at the

shelter's three rehoming centres during 2014 (including dogs that arrived during 2013 or

left in 2015). We selected the records from all dogs that were at least 4 months old,

144 excluding younger dogs because they were more likely to be unvaccinated, more limited

145 in their interactions at the shelter and may have been kennelled in different areas to older

146 dogs. Although dogs were from a variety of heritages (including purebreds and

147 mongrels), the analyses here did not explore breed differences because the accurate visual

148 assessment of breed in dogs with unknown heritage has been refuted [56–58].

Table 1. Demographic characteristics of the studied dogs.

Variable	Mean ± SD / *N*
Average age at shelter (years; all ≥ 4 months of age)	3.75 ± 3.03
Total days at the shelter	25.13 ± 41.53
Weight (average weight if multiple measurements; kg)	19.06 ± 10.26
Rehoming centre: London / Old Windsor / Brands Hatch	2897 / 1280 / 566
Males / females	2749 / 1994
Neutered[1] before arrival / neutered at shelter / not neutered	1218 / 1665 / 1502
Relinquished by owners / returned to shelter / strays	2892 / 260 / 1591

[1]358 dogs had unknown neuter status

Shelter environment

The shelter was composed of three different UK rehoming centres: a high-throughput,

urban centre based at Battersea, London with capacity for approximately 150-200 dogs; a

semi-rural/rural centre based at Old Windsor with capacity for approximately 100-150

dogs; and a rural centre based at Brands Hatch with capacity for approximately 50 dogs.

All dogs arrived in an intake area of their respective rehoming centre and, when

considered suitable for adoption, were moved to a 'rehoming' area that was partially open

to the public between 1000 h and 1600 h. All kennels were indoors. Kennels varied in

size, but were usually approximately 4m x 2m and included either a shelf and bedding

alcove area, or a more secluded bedding area at the back of the kennel (see [59] for more

details). At different times throughout the day, dogs had access to indoor runs behind

their kennels. In each kennel block area, dogs were cared for (e.g. fed, exercised, kennel

cleaned) by a relatively stable group of staff members, allowing the development of

familiarity with staff members and offering some predictability for dogs after arrival at

the shelter. Although data on the number of dogs in each kennel were incomplete, in the

majority of cases dogs were kennelled singly for safety reasons. The shelter mainly

operated between 0800 h and 1700 h each day. All dogs were socialised with staff and/or

volunteers each day (often multiple times) except on rare occasions when it was deemed

unsafe to handle a dog (when training/behavioural modification proceeded without

physical contact). Dogs were provided water ad libitum and fed commercial complete dry

and/or wet tinned food twice daily (depending on recommendations by veterinary staff).

Dogs received daily tactile, olfactory and/or auditory enrichment/variety (e.g. toys,

essential oils, classical music, time in a quiet 'chill-out' room).

Data collection

In the observational assessment procedure, trained shelter employees recorded

observations of dog behaviour in a variety of contexts as part of normal shelter

procedures. Behavioural observations pertaining to each context were completed using an

ethogram specific to that context and recorded in a custom computer system. Multiple

observations could be completed each day, although we retained only one observation in

each context per day (the least desirable behaviour on that day; see below). The ethogram

code that best described a dog's behaviour in a particular context during an observation

was recorded by selecting it from a series of drop-down boxes (one for each context).

Although staff could also add additional information in character fields, a full analysis of

those comments was beyond the scope of this study. The ethogram for each context

represented a scale of behaviours ranging from desirable to undesirable considered by the

shelter to be relevant to dog welfare and ease of adoption. Contexts had between 10 and

16 possible behaviours to choose from, some of which overlapped between different

contexts. Among the least desirable behaviours in each context was aggression towards

either people or dogs (depending on context). Aggression was formally defined as

"*Growls, snarls, shows teeth and/or snaps when seeing/meeting other people/dogs,*

potentially pulling or lunging towards them", distinguished from non-aggressive but

reactive responses, defined as "*Barks, whines, howls and/or play growls when*

seeing/meeting other people/dogs, potentially pulling or lunging towards them".

Observation contexts included both onsite (at the shelter) and offsite (e.g. out in public

parks) settings. For the analyses here, we excluded offsite contexts (which had separate

observation categories) because these were less frequently recorded and offsite records

were more likely to be completed using second-hand information (e.g. from volunteers

taking the dog offsite). We focused on observations of aggression in nine core onsite

contexts that were most frequently completed by trained staff members: i) *Handling*, ii)

In kennel, iii) *Out of kennel*, iv) *Interactions with familiar people*, v) *Interactions with*

unfamiliar people, vi) *Eating food*, vii) *Interactions with toys*, viii) *Interactions with*

female dogs, ix) *Interactions with male dogs*. For the *In kennel* and *Out of kennel*

contexts, recording of aggression towards both people and dogs was possible. If both

occurred at the same time, aggression towards people was recorded. Therefore, *In kennel*

and *Out of kennel* were each divided to reflect aggression shown towards people and

towards dogs only, respectively. This resulted in 11 aggression contexts (Table 2) used as

manifest variables in structural equation models to investigate latent aggressiveness traits.

The average number of days between successive observations across these contexts and

across dogs was 3.27 (SD = 2.08), and dogs had an average of 9.77 (SD = 13.41)

observations within each context (N = 416,860 observations in total across dogs, contexts

and days). Observations were recorded in the category that best described the scenario.

Nonetheless, certain contexts could occur closely in space and time, which were

investigated for violations of local independence, as explained below.

Table 2. Behavioural observation contexts in which each dog's reactions were analysed for the presence or absence of aggression.

Context	Definition
Handling	Informal handling by people (e.g. stroking non-sensitive areas, touching the collar, fitting a harness or lead).
In kennel towards people	People approaching or walking past the kennel.
In kennel towards dogs	Dogs in neighbouring kennels or dogs walking past the kennel.
Interactions with familiar people	When outside the kennel and familiar people (interacted with at least once before) approach, make eye contact, speak to or attempt to make physical contact with the dog.
Interactions with unfamiliar people	When outside the kennel and unfamiliar people (never interacted with before) approach, make eye contact, speak to or attempt to make physical contact with the dog.
Out of kennel towards people	When around people outside the kennel who may be a long distance away and who make no attempt to engage with the dog.
Out of kennel towards dogs	When around dogs outside the kennel that may be a long distance away and that are not encouraged to interact with the focal dog.
Eating food	When eating food (e.g. from a food bowl, or toy filled with food) and people approach within close proximity or attempt to touch the food container.
Interactions with toys	When interacting with toys and people approach within close proximity or attempt to touch the toy.
Interactions with female dogs	During structured interaction with a female dog, including approaching each other, walking in parallel, and interacting off-lead. Both dogs are aware of each other's presence and are in close enough proximity to engage in a physical interaction.
Interactions with male dogs	During structured interaction with a male dog, including approaching each other, walking in parallel, and interacting off-lead. Both dogs are aware of each other's presence and are in close enough proximity to engage in a physical interaction.

We aggregated behavioural observations across time for each dog into a dichotomous

variable indicating whether a dog had or had not shown aggression in a particular context

at any time while at the shelter (Table S1). This was performed because the overall

prevalence of aggression was low, with only 1.06% of all observations across days

involving aggression towards people and 1.13% towards dogs. Thus, the main difference

between individuals was whether they had or had not shown aggression in a particular

context during their time at the shelter. We interpret aggressiveness here as a between-

individual difference variable.

Validity of behaviour recordings

Validity of the recording of behaviour was assessed separately from the main data

collection as part of a wider project investigating the use of the observational assessment

method. Ninety-three shelter employees trained in conducting behavioural observations

each watched (in groups of 5 – 10 people) 14 videos, approximately 30 seconds each,

presenting exemplars of 2 different behaviours from seven contexts (to keep the sessions

concise and maximise the number of participants). For each context, behaviours were

chosen pseudo-randomly by numbering each behaviour and selecting two numbers using

a random number generator. Experienced behaviourists working at the shelter filmed the

videos demonstrating the behaviours. Videos were shown to participants once in a

pseudo-random order. After each video, participants recorded on a paper answer sheet the

behaviour they thought most accurately described the dog's behaviour based on the

ethogram specific to the context depicted. Two of the videos illustrated aggression: one in

a combined *Interactions with new* and *familiar people* context (combined because

familiarity between specific people and dogs was not universally known) and one in the

In kennel towards dogs context. The first video had an ethogram of 13 possible

behaviours to choose from, and the second had 11 behaviours. The authors were blind to

the selection of videos shown and to the video coding sessions with shelter employees.

Data analysis

246 All data analysis was conducted in R version 3.3.2 [60].

Validity of behaviour recordings

249 The degree to which shelter employees could recognise and correctly record aggressive

250 behaviour from the videos (chosen by experienced behaviourists at the shelter) was

251 determined by the percentage of participants who correctly identified the 2 videos as

252 showing examples of aggression.

Missing data

Data were missing when dogs did not experience particular contexts while at the shelter.

The missing data rate was between 0.06% and 5% for each context, except for the

Interactions with female dogs and *Interactions with male dogs* categories which had 17%

and 18% of missing values, respectively (because structured interactions with other dogs

did not arise as frequently). Moreover, 16% and 8% of dogs were missing weight

measurement and neuter status data, respectively, which were independent variables

statistically controlled for in subsequent analyses. We created 5 multiply imputed data

sets (using the *Amelia* package; [61]), upon which all following analyses in the sections

below were conducted and results pooled. The multiple imputation took into account the

hierarchical structure of the data (observations within dogs), all independent variables

reported below, and the data types (ordered binary variables for the context data,

positive-continuous for weight measurements, nominal for neuter status; see the R script).

The data were assumed to be missing at random, that is, dependent only on other

variables in the analyses.

Structural equation models

We used structural equation modelling to assess whether aggression towards people

(contexts: *Handling*, *In kennel towards people*, *Out of kennel towards people*,

Interactions with familiar people, *Interactions with unfamiliar people*, *Eating food*,

Interactions with toys) and towards dogs (contexts: *In kennel towards dogs*, *Out of kennel*

towards dogs, *Interactions with female dogs*, *Interactions with male dogs*) could be

explained by two latent aggressiveness traits: aggressiveness towards people and dogs,

respectively. Since positive correlations between different aggressiveness traits have been

reported in dogs [55], we compared a model where the latent variables were orthogonal

to a model where variables were allowed to covary. Models were fit using the *lavaan*

package [62], with the weighted least squares mean and variance adjusted (WLSMV)

estimator and theta/conditional parameterisation, as recommended for categorical

dependent variables [8,63,64]. The latent variables were standardised to have mean 0 and

variance 1. The results were combined across imputed data sets using the 'runMI'

function in the *semTools* package [65]. The fit of each model was ascertained using the

comparative fit index (CFI) and Tucker Lewis index (TLI), where values > 0.95 indicate

excellent fit, as well as the root mean squared error of approximation (RMSEA) where

values < 0.06 indicate good fit [7]. Parameter estimates were summarised by test statistics

and 95% confidence intervals (CI).

Local independence

We tested the assumption of local independence by re-fitting the best-fitting structural

equation model with residual covariances specified between context variables. To

maintain a theoretically driven approach (see [66] regarding the best practice of including

residual covariances in structural equation models) and model identifiability, we only

tested a predefined set of covariances based on which contexts shared close temporal-

spatial relationships. First, we allowed covariances between *Handling* with *In kennel*

towards people, *Interactions with familiar people*, *Interactions with unfamiliar people*

and *Interactions with toys*, respectively, since the *Handling* context could directly

succeed these other contexts. The residual covariance between *Handling* and *Eating food*

was not estimated because shelter employees would be unlikely to handle a dog while the

dog ate its daily meals. The residual covariance between *Handling* and *Out of kennel*

towards people was not estimated because any association between *Handling* and *Out of*

kennel towards people would be mediated by either the *Interactions with familiar people*

or *Interactions with unfamiliar people* context. Therefore, secondly, we estimated the

three-way covariances between *Out of kennel towards people*, *Interactions with familiar*

people and *Interactions with unfamiliar people*. Similarly, and lastly, we estimated the

three-way covariances between *Out of kennel towards dogs, Interactions with female*

dogs and *Interactions with male dogs*. No covariances were inspected between *In kennel*

towards dogs and other aggressiveness towards dogs contexts since large time gaps were

more likely to separate observations between those contexts.

Measurement invariance

To test for measurement invariance in each of the latent traits derived from the best

fitting structural equation model, we investigated the response patterns across aggression

contexts related to the same latent aggressiveness trait using Bayesian hierarchical

logistic regression models. These models were analogous to the 1-parameter item

response theory model, which represents the probability that an individual responds

correctly to a particular test item as a logistic function of i) each individual's latent ability

and ii) the item's difficulty level. This model can be expressed as a hierarchical logistic

regression model [67,68], whereby individual latent abilities are modelled as individual-

specific intercepts (i.e. 'random intercepts'), the propensity for a correct answer to an

item i is its regression coefficient β_i, and credible interactions between items and relevant

independent variables (e.g. group status) indicate a violation of measurement invariance.

Here, the dependent variable was the binary score for whether or not dogs had shown

aggression in each context and the average probability of aggression across contexts

varied by dog, representing latent levels of aggressiveness. Context type, dog age, dog

sex and their interactions were included as categorical independent variables. Age was

treated as a categorical variable, with categories reflecting general developmental

periods: i) 4 months to 10 months (juvenile dogs before puberty), ii) 10 months to 3 years

(dogs maturing from juveniles to adults), iii) 3 years to 6 years (adults), and iv) 6 years +

(older dogs). Broad age categories were chosen due to potentially large differences in

developmental timing between individuals. Age was categorised because we predicted

that aggression would be dependent on these developmental periods.

Models included additional demographic variables (Table 1) that may mediate the

probability of aggression: body weight (average weight if multiple measurements were

taken), total number of days spent at the shelter, the rehoming centre at which dogs were

based (London, Old Windsor, Brands Hatch), neuter status (neutered before arrival,

neutered at the shelter, not neutered) and source type (relinquished by owner, returned to

the shelter after adoption, stray). Categorical variables were represented as sum-to-zero

deflections from the group-level intercept to ensure that the intercept represented the

average probability of aggression across the levels of each categorical predictor. Weight

and total days at the shelter were mean-centered and standardised by 2 standard

deviations. Due to the potentially complex relationships between these variables and

aggression (e.g. interactive effects between neuter status and sex; [52]), which could also

include violations of measurement invariance, we decided not to interpret their effects

inferentially. Instead, they were included to make the assessment of measurement

invariance between sexes and age groups conditional on variance explained by

potentially important factors.

For comparability to other studies in animal personality, behavioural repeatability was

calculated across contexts in each model using the intraclass correlation coefficient

(ICC), calculated as $\frac{\sigma_\beta^2}{\sigma_\beta^2 + \sigma_\epsilon^2}$, where σ_β^2 represented the between-individual variance of the

probability of aggression (i.e. the variance of the random intercepts), and σ_ϵ^2 was $\pi^2/3$,

the residual variance of the standard logistic distribution [69].

Computation

Models were computed using the probabilistic programming language Stan version

2.15.1 [70], using Hamiltonian Monte Carlo, a type of Markov Chain Monte Carlo

(MCMC) algorithm, to sample from the posterior distribution. Prior distributions for all

independent variables were normal distributions with mean 0 and standard deviation 1,

attenuating regression coefficients towards zero for conservative inference. The prior on

the overall intercept parameter was normally distributed with mean 0 and standard

deviation 5. The standard deviation of dog-specific intercept parameters was given a half-

Cauchy prior distribution with mean 0 and shape 2. Each model was run with 4 chains of

2,000 iterations with a 1,000 step warm-up period. The Gelman-Rubin statistic (ideally <

1.05) and visual assessment of traceplots were used to assess MCMC convergence. We

checked the accuracy of the model predictions against the raw data using graphical

posterior predictive checks. For plotting purposes, predicted probabilities of aggression

were obtained by marginalising over the random effects (explained in the Supporting

Information). Regression coefficients were expressed as odds ratios and were

summarised by their mean and 95% Bayesian highest density interval (HDI), representing

the 95% most probable parameter values. To compare levels of categorical variables and

their interactions, we computed the 95% HDI of the differences between the respective

posterior distributions.

Model selection & parameter inference

Models were run on each imputed data set and their respective posterior distributions

were averaged to attain a single posterior distribution for inference. Adopting a Bayesian

approach allowed the estimation of interaction parameters (i.e. testing measurement

invariance) without requiring corrections for multiple comparisons as in null hypothesis

significance testing [71]. Nonetheless, models included a large number of estimated

parameters. Two strategies were employed to guard against over-fitting of models to data.

First, we selected the model with the best out-of-sample predictive accuracy given the

number of parameters based on the Widely Applicable Information Criterion (WAIC;

using the R package *loo* [72]). Four variants of each model were computed: two-way

interactions between contexts and age and contexts and sex, respectively (model 1), a

single interaction with sex but not with age (model 2), a single interaction with age but

not with sex (model 3), and no interactions (model 4). All models included the mediating

independent variables above. Second, to avoid testing point-estimate null hypotheses, the

effect of a parameter was only considered credibly different from zero if the odds ratio

exceeded the region of practical equivalence (ROPE; see [73]) around an odds ratio of 1

from 0.80 to 1.25. An odds ratio of 0.80 or 1.25 indicates a 20% decrease or increase (i.e.

4/5 or 5/4 odds), respectively, in the odds of an outcome, frequently used in areas of

bioequivalence testing (e.g. [74]), which we here considered to be small enough to

demonstrate a negligible effect in the absence of additional information. If a 95% HDI

fell completely within the ROPE, the null hypothesis of no credible influence of that

parameter was accepted; if a 95% HDI included part of the ROPE, then the parameter's

influence was left undecided [73].

Ethics statement

402   Permission to use and publish the data was received from the shelter. Approval from an

403   ethical review board was not required for this study.

Data accessibility

406   Supporting Information (data, R script, Stan model code, Tables S1-4) can be found at:

407   https://github.com/ConorGoold/GooldNewberry_aggression_shelter_dogs.

Results

Validity of behaviour recordings

For the video showing aggression towards people, 52% of participants identified the behaviour correctly as aggression and 42% identified the behaviour as non-aggressive but (similarly) reactive behaviour (see definitions above). For the video showing aggression towards dogs, 53% identified the behaviour correctly and 44% identified the behaviour as non-aggressive but reactive behaviour. For the 12 other videos not showing aggression, only 1 person incorrectly coded a video as aggression towards people and 3 people incorrectly coded videos as aggression towards dogs.

Structural equation models

The raw tetrachoric correlations between the aggression contexts were all positive, particularly between contexts recording aggression towards people and dogs, respectively, supporting their convergent validity (Table S2). The model with correlated latent variables fit marginally better (CFI: 0.96; TLI: 0.95; RMSEA: 0.03) than the model with uncorrelated variables (CFI: 0.94; TLI: 0.92; RMSEA: 0.04). All regression coefficients of the model with correlated latent variables were positive and significant (i.e. the 95% CI did not include zero), and the latent variables shared a significant positive covariance (Table 3).

Table 3. Parameter estimates from the best-fitting structural equation model.

Parameter	Estimate	SE	*t* value	95% CI
Handling[a]	0.81	0.06	14.25	[0.70, 0.92]
In kennel towards people[a]	1.29	0.09	14.17	[1.12, 1.46]
Out of kennel towards people[a]	0.83	0.07	11.99	[0.69, 0.96]
Interactions with familiar people[a]	0.96	0.07	14.23	[0.83, 1.09]
Interactions with unfamiliar people[a]	1.54	0.12	12.46	[1.23, 1.78]
Eating food[a]	0.70	0.06	12.33	[0.59, 0.81]
Interactions with toys[a]	0.51	0.06	8.32	[0.39, 0.63]
In kennel towards dogs[b]	0.70	0.06	11.94	[0.59, 0.82]
Out of kennel towards dogs[b]	0.47	0.04	10.80	[0.38, 0.55]
Interactions with female dogs[b]	0.87	0.07	12.05	[0.72, 1.02]
Interactions with male dogs[b]	0.88	0.07	12.23	[0.74, 1.03]
Covariance: People ~ Dogs	0.26	0.03	7.94	[0.19, 0.33]

[a] Contexts reflecting aggressiveness towards people

[b] Contexts reflecting aggressiveness towards dogs

Local independence

Allowing the pre-defined residuals to co-vary in the best-fitting structural equation model

resulted in a better fit (CFI = 0.98; TLI = 0.97; RMSEA: 0.03). Significant negative

covariances were observed between the *Handling* and *In kennel towards people* contexts

(Table 4) and the *Handling* and *Interactions with unfamiliar people* contexts. A

significant positive covariance was observed between *Out of kennel towards people* and

Interactions with unfamiliar people contexts. No significant residual covariances between

contexts reflecting aggressiveness towards dogs were observed.

Table 4. Estimated residual covariances between contexts.

Residual covariances	Estimate	SE	t value	95% CI
Handling ~ In kennel towards people[a]	-0.60	0.21	-2.86	[-1.01, -0.19]
Handling ~ Interactions with familiar people[a]	0.16	0.09	1.84	[-0.01, 0.33]
Handling ~ Interactions with unfamiliar people[a]	-0.48	0.19	-2.49	[-0.86, -0.10]
Handling ~ Interactions with toys[a]	0.14	0.07	1.85	[-0.01, 0.28]
Out of kennel towards people ~ Interactions with familiar people[a]	0.04	0.08	0.49	[-0.12, 0.20]
Out of kennel towards people ~ Interactions with unfamiliar people[a]	0.24	0.09	2.56	[0.06, 0.42]
Interactions with familiar people ~ Interactions with unfamiliar people[a]	-0.02	0.12	-0.16	[-0.25, 0.21]
Out of kennel towards dogs ~ Interactions with female dogs[b]	-0.55	0.48	-1.15	[-1.50, 0.40]
Out of kennel towards dogs ~ Interactions with male dogs[b]	-0.45	0.40	-1.13	[-1.22, 0.33]
Interactions with female dogs ~ Interactions with male dogs[b]	-0.24	0.50	-0.49	[-1.23, 0.74]

[a] Contexts reflecting aggressiveness towards people

[b] Contexts reflecting aggressiveness towards dogs

Measurement invariance

Separate models were run for contexts reflecting aggressiveness towards people and

aggressiveness towards dogs. All models converged. Posterior predictive checks of model

estimates reflected the raw data (Figs 1 and 2). The full measurement invariance model

(model 1) including interactions between contexts and sex and contexts and age groups

had the best out-of-sample predictive accuracy for both the aggressiveness towards

people and aggressiveness towards dogs models, respectively, illustrated by the lowest

WAIC values (Table 5). Since some models included numerous interactions, we provide

an overall summary of the main results below (Figs 1 and 2) with full parameter

estimates provided in Tables S3 and S4.

Table 5. Mean ± standard error of the Widely Applicable Information Criteria (WAIC) values (lower is better) per model and aggressiveness variable.

Model	Aggressiveness towards people	Aggressiveness towards dogs
Model 1	13405.6 ± 179.0	15257.2 ± 133.1
Model 2	13506.3 ± 179.6	15381.4 ± 133.4
Model 3	13426.3 ± 179.1	15285.3 ± 133.0
Model 4	13521.7 ± 179.5	15407.6 ± 133.4

Aggressiveness towards people

The odds of aggression towards people, across categorical predictors and for an average

dog of mean weight and length of stay at the shelter, were 0.022 (HDI: 0.021 to 0.024), a

probability of approximately 2%. On average, aggression was most likely in the *In kennel*

towards people context (OR = 0.054; HDI: 0.049 to 0.058) and least probable in the

Interactions with toys context (OR = 0.008; HDI: 0.007 to 0.009).

Aggression was less likely across contexts for females than males (OR = 0.719; HDI:

0.668 to 0.770), although there were also credible interactions between sex and contexts

(Fig 1A; Table S3). Whereas males and females had similar odds of aggression in the *Out*

of kennel towards people context, smaller differences were observed between *Out of*

kennel towards people and *Handling* (OR = 0.578; HDI: 0.481 to 0.682), *Eating food*

(OR = 1.812; HDI: 1.495 to 2.152) and *Interactions with familiar people* (OR = 1.798;

HDI: 1.488 to 2.126) contexts in females compared to males. Additionally, whereas

aggression in the *Interactions with unfamiliar people* context was similar between males

and females, larger differences were observed between *Interactions with unfamiliar*

people and *Handling* (OR = 0.616; HDI: 0.530 to 0.702), *Eating food* (OR = 0.594; HDI:

0.506 to 0.686) and *Interactions with familiar people* (OR = 0.598; HDI: 0.513 to 0.687)

contexts in females compared to males.

Apart from lower odds of aggression in 4 to 10 month olds compared to 10 month to 3

year old dogs (OR = 0.638; HDI: 0.565 to 0.705), there was no simple influence of age

group on aggressiveness. Between the 4 to 10 months old and 3 to 6 years old groups,

differences between the odds of aggression across contexts varied due to an increase of

aggression in certain contexts but not others (Fig 1B; Table S4). Aggression in *In kennel*

towards people and *Interactions with unfamiliar people* contexts particularly increased,

leading to larger differences between, for example, *In kennel towards people* and *Eating*

food (OR = 0.524; HDI: 0.400 to 0.642) and *Eating food* and *Interactions with unfamiliar*

people (OR = 1.721; HDI: 1.403 to 2.059) contexts for 10 month to 3 year olds compared

to 4 to 10 month olds, and between *In kennel towards people* and *Out of kennel towards*

people (OR = 0.470; HDI: 0.355 to 0.606) and *Out of kennel towards people* and

Interactions with unfamiliar people (OR = 2.051; HDI: 1.608 to 2.543) contexts in 3 to 6

year olds compared to 4 to 10 month olds. In 3 to 6 year old compared to 10 month to 3

year old dogs, aggression increased in the *Handling* and *Eating food* contexts but

decreased in the *Out of kennel towards people* context, resulting in larger differences

between, for instance, *Handling* and *Out of kennel towards people* (OR = 0.526; HDI:

0.409 to 0.631) and *Out of kennel towards people* and *Interactions with unfamiliar people*

(OR = 2.349; HDI: 1.891 to 2.925), and smaller differences between *Eating food* and

Interactions with familiar people (OR = 0.576; HDI: 0.468 to 0.687).

Dogs over 6 years old demonstrated qualitatively different response patterns across

certain contexts than all other age groups. While aggression was most probable in *In*

kennel towards people and *Interactions with unfamiliar people* contexts for dogs aged 4

months through 6 years, dogs over 6 years old were most likely to show aggression in the

Handling context, leading to interactions between, for example, *Handling* and *In kennel*

towards people, and between *Handling* and *Interactions with unfamiliar people* contexts

compared to the other age groups (Fig 1B; Table S3). Aggression when *Eating food* and

in *Interactions with toys* contexts also increased compared to that expressed by younger

dogs, resulting in credible differences between, for instance, *Eating food* and *Interactions*

with familiar people contexts between dogs aged 10 months to 3 years and over 6 years

(OR = 0.379; HDI: 0.300 to 0.465) and between *Out of kennel towards people* and

Interactions with toys contexts between over 6 year olds and all other age groups (Table

S3).

Fig 1. Predicted probabilities of aggression towards people in different contexts by

sex (panel A) and age groups (panel B). Black points and vertical lines show mean and

95% highest density intervals of model parameter estimates; blue triangles show raw

sample data. Model estimates were obtained by marginalising over the random effects

(see the Supporting Information). Abbreviations used in the figure: HND (*Handling*); KP

(*In kennel towards people*); OKP (*Out of kennel towards people*); FPL (*Interactions with*

familiar people); UPL (*Interactions with unfamiliar people*); EAT (*Eating food*); TOY

(*Interactions with toys*).

Aggressiveness towards dogs

The odds of aggression towards dogs, across categorical predictors and for an average

dog of mean weight and length of stay at the shelter, was 0.176 (HDI: 0.168 to 0.184),

corresponding to a probability of approximately 15%. Dogs were most likely to show

aggression in the *Interactions with male dogs* context (OR = 0.297; HDI: 0.198 to 0.217)

and least likely in the *In kennel towards dogs* context (OR = 0.099; HDI: 0.094 to 0.104;

Fig 2; Table S4).

No credible mean-level differences existed between females and males (OR = 1.187;

HDI: 1.128 to 1.250). However, the difference in aggression between the *Interactions*

with female dogs and *Interactions with male dogs* contexts was smaller for females (OR

= 1.542; HDI: 1.400 to 1.704; Fig 2A; Table S4), as were the differences between

Interactions with male dogs and *In kennel towards dogs* (OR = 0.661; HDI: 0.590 to

0.732) and *In kennel towards dogs* and *Out of kennel towards dogs* (OR = 1.420; HDI:

1.269 to 1.587). Females were also more likely to show aggression in *Interactions with*

female dogs than *Out of kennel towards dogs* compared to males (OR = 1.444; HDI:

1.301 to 1.603).

Dogs aged 4 to 10 months old had credibly lower odds of aggression towards dogs than

older dogs across contexts (Fig 2B; Table S4). However, contexts and age also showed

interactive effects. In particular, aggression in *Interactions with female dogs* and

Interactions with male dogs contexts tended to increase relative to other contexts. For

instance, the relationship between *Interactions with female dogs* and *Out of kennel*

towards dogs contexts reversed in direction between 4 to 10 month and 10 month to 3

year olds (OR = 0.595; HDI: 0.495 to 0.688) as did the relationship between *Interactions*

with male dogs and *Out of kennel towards dogs* contexts (OR = 0.499; HDI: 0.422 to

0.575). The relationship between *In kennel towards dogs* and *Out of kennel towards dogs*

contexts also changed across age groups (Fig 2B; Table S4). Four to 10 months old were

more likely to show aggression in *Out of kennel towards dogs* than *In kennel towards*

dogs contexts, but the difference was smaller in 10 months to 3 year olds (OR = 0.608;

HDI: 0.505 to 0.728) and in over 6 year olds (OR = 0.396; HDI: 0.316 to 0.481). The

latter relationship was reversed in 3 to 6 year olds compared to 4 to 10 month old dogs

(OR = 0.277; HDI: 0.227 to 0.331) and 10 month to 3 year old dogs (OR = 0.456; HDI:

0.396 to 0.516).

Fig 2. Predicted probabilities of aggression towards dogs in different contexts by sex

(panel A) and age groups (panel B). Black points and vertical lines show mean and

95% highest density intervals of model parameter estimates; blue triangles show raw

sample data. Model estimates were obtained by marginalising over the random effects

(see the Supporting Information). Abbreviations used in the figure: KD (*In kennel*

towards dogs); OKD (*Out of kennel towards dogs*); DF (*Interactions with female dogs*);

DM (*Interactions with male dogs*).

Repeatability

564 Both aggressiveness towards people and dogs showed moderate repeatability across

565 contexts ($ICC_{people} = 0.479$; HDI: 0.466 to 0.491; $ICC_{dogs} = 0.303$; HDI: 0.291 to

566   0.315), although aggressiveness towards people was more repeatable than aggressiveness

567   towards dogs ($ICC_{difference} = 0.176$; HDI: 0.158 to 0.192).

Discussion

In this study, we have examined local independence and measurement invariance of aggressiveness traits in shelter dogs. Observational recordings of aggression directed towards people and dogs across different shelter contexts were explained by two positively correlated latent variables, and behaviour across contexts was moderately repeatable. These results are consistent with the concept of animal personality, which is used to describe behaviour that shows moderately consistent between-individual differences across time or contexts, and is characterised by multiple observed behaviours being decomposed into lower-dimensional behavioural traits [4]. However, we found violations of local independence between contexts with close temporal-spatial relationships and measurement invariance with respect to sex and age groups, highlighting potential measurement biases.

Local independence implies that the association between manifest variables is greater than that explained by the latent variable. For aggressiveness towards people, aggression in the *Handling* context was negatively related with the *In kennel towards people* and *Interactions with unfamiliar people* contexts, while positive covariances were present between *Out of kennel towards people* and *Interactions with unfamiliar people* contexts. Violations of local independence may arise through shared method variance [75–78] or unmodelled latent variables influencing manifest variables [79,80]. If a dog showed aggression when an unfamiliar person approached, it may be less likely to be handled by that person, which may explain the negative residual covariations between the *Handling*

and *In kennel towards people* and *Interactions with unfamiliar people* contexts,

respectively. These contexts were, in fact, positively correlated when latent levels of

aggressiveness were not accounted for (Table S4). In addition, the positive residual

correlation between *Out of kennel towards people* and *Interactions with unfamiliar*

people may be mediated by additional traits of interest to personality researchers, such as

fearfulness or anxiety [29,81], if dogs who are fearful of interacting with unfamiliar

people are more likely to show aggression beyond that described by a latent

aggressiveness trait.

While authors have argued that greater standardisation and validation of personality

assessments is key to ensuring the accurate measurement of underlying traits [36,48,49],

it may be untenable to avoid dependencies between testing contexts. Displays of

aggression in one sub-test will likely change how people conduct future sub-tests with the

same dog, regardless of test standardisation. Human psychologists have argued that

violations of local independence are a natural consequence of the organisation of

behaviour as a complex dynamic system [82,83], which unfolds with respect to time- and

context-dependent constraints [84]. Thus, awareness of local independence and its

violation could facilitate closer understanding of the dynamics driving personality test

responses beyond explanations purely based on personality traits.

While different subsets of a population may differ in mean levels of trait expression,

interactions between behavioural responses and those subsets indicate that the same

phenomenon is not under measurement across groups [23,24]. We found that the

probability of aggression across contexts was dependent on sex and age conditional on

latent levels of aggressiveness (Figs 1 and 2; Tables S3 and S4). Female dogs, for

example, were more likely than males to show aggression in *Out of kennel towards*

people and *Interactions with unfamiliar people* contexts relative to other contexts (Fig

1A). Females also demonstrated similar odds of aggression during *Interactions with*

female dogs and *Interactions with male dogs*, whereas males were more likely to show

aggression towards male than female dogs (Fig 2a). As with local independence, different

behavioural variables unaccounted for in this study may result in violations of

measurement invariance. While dogs up to 6 years old were most likely to show

aggression in *In kennel towards people* and *Interactions with unfamiliar people contexts*,

dogs over 6 years old demonstrated aggression most commonly in the *Handling* context.

Dogs over 6 years old also showed an increase in aggression in the *Eating food* and

Interactions with toys contexts relative to other age groups. These results suggest that

older dogs in shelter populations may be less tolerant during close interactions with

people (i.e. handling, people in the vicinity of their food and toys) compared to other

contexts, which may driven by other quantifiable factors such as pain or sensitivity (e.g.

[29]).

Although we have identified violations of both local independence and measurement

invariance, we remain cautious about hypothesising *a posteriori* about their causes.

Personality traits in animal behaviour are typically defined operationally, based on the

statistical repeatability of quantifiable behaviour [77,85,86]. As discussed in human

personality psychology, operational definitions can be ontologically ambiguous [87,88].

That is, while operational definitions facilitate experimentation in animal personality [4],

they do not necessarily designate biological mechanisms underlying trait expression. For

example, Budaev and Brown remark that boldness, defined as a propensity to take risks,

could encompass a range of distinct personality traits, each with a different biological

basis [75]. Whilst reflective latent variable models allow researchers to test hypotheses

about the relatedness of measured behaviours via one or more underlying traits, they have

also been criticised as ambiguous [82]. For example, it is uncertain what reflective latent

variables may represent in biological organisation [87] or even whether they are features

individuals possess or simply emergent features of between-individual differences

[89,90]. Such considerations highlight the importance of research on the proximate

mechanisms of personality [85] and longitudinal data analyses to separate between- from

within-individual behavioural variation [91,92].

A number of authors have emphasised the poor predictive value of aggression tests in

shelter dogs [39–41,50] and that low occurrence of aggression specifically can make its

accurate measurement difficult [40]. The probability of observing aggression on any

particular day was low in this study (approximately 1%), and the number of dogs who, on

average, showed aggression to people at least once while at the shelter was much lower

than the number that showed aggression towards dogs, on average (Figs 1 and 2).

Nonetheless, our evaluations of validity indicated that between 40 and 45% of the shelter

employees mistook observations of aggression for non-aggressive responses (e.g. over-

excitement and frustration when seeing people/dogs), meaning that the true probability of

aggression was potentially under-estimated (although incorrectly coding other behaviours

as aggression also occurred, albeit rarely). Moreover, our assessments of validity were

based on shelter staff evaluations of brief video recordings that may be less reliable than

the live, spontaneous behavioural recordings upon which our main analyses were based,

resulting in a lower percentage of correctly identified instances of aggression. For the two

videos being evaluated, the shelter employees had 13 and 11 different behavioural codes,

respectively, to choose from to describe the behaviours observed. Thus, while employees

as a whole were undecided about whether the motivation for the behaviour was

aggressive or non-aggressive, the vast majority of employees described the behaviour as

reactive, despite potentially erring on the side of caution by labelling aggressive

behaviours as non-aggressive. Comparable estimates of validity are not present in the

literature on dog personality, but are particularly important in shelter settings where

accurate recording of aggression is paramount. It is also worth noting that how to assess

validity has received much debate (e.g. [87,92]). In this study, we used expert judgement

as a benchmark to which shelter employees' responses were compared, but validity is

frequently assessed in dog personality by inspecting patterns of correlation coefficients

between similar and dissimilar behaviours (e.g. convergent or divergent validity; [29]).

This is less directly interpretable than reporting the percentage of answers that were

correct, as used here. Moreover, the predictive validity of personality assessments in dogs

have been inconsistent (e.g. [40-42]). More discussion of the concept of validity, and how

best to assess it, is warranted in studies of dog personality.

Infrequent occurrence and/or recording of aggression may also limit accurate predictions

of future behaviour. Patronek and Bradley [50] demonstrate using simulation that the low

prevalence of aggression inflates the chance that aggression shown in a shelter

assessment represents a false positive. In general, our results support this conclusion in

the sense that aggression may be shown differentially across contexts not explained by

latent levels of aggressiveness. Violations of local independence and measurement

invariance as found here indicate, further, that it is not only the difference between false

and true positives and negatives, but the validity of inferring homogeneous personality

traits by which to compare individual dogs, that needs careful consideration.

Consequently, we agree with recommendations to establish the efficacy of longitudinal,

observational assessments rather than relying on a single assessment made using a

traditional test battery [31,40,50]. This approach will prioritise the cumulative

understanding of a dog's context-dependent behaviour and help to guide decisions about

the potential risk a dog poses to humans and other animals.

Conclusion

This study has tested the assumptions of local independence and measurement invariance

of personality traits in shelter dogs. Using structural equation modelling, aggression

across behavioural contexts was explained by two correlated latent variables and

demonstrated repeatability. Nevertheless, significant residual covariances remained

between certain behavioural contexts related to aggressiveness towards people, violating

the assumption of local independence. In addition, aggression in different contexts

showed differential patterns of response across sex and age, indicating a lack of

measurement invariance. Violations of local independence and measurement invariance

imply that the aggressiveness towards people and dogs traits did not completely explain

patterns of aggression in different contexts, or that inferences based on these

hypothesised personality traits may in fact be misleading. We encourage researchers to

more closely assess the measurement assumptions underlying reflective latent variable

models before making conclusions about the effects of, or factors influencing,

personality.

Acknowledgements

713 The authors are extremely grateful to Battersea Dogs and Cats Home for allowing us to

714 access the data for this study.

References

1. Dochtermann NA, Jenkins SH. Behavioural syndromes in Merriam's kangaroo rats

(*Dipodomys merriami*): a test of competing hypotheses. Proc R Soc B. 2007;274:

2343–2349. doi:10.1098/rspb.2007.0622

2. Dochtermann NA, Dingemanse NJ. Behavioral syndromes as evolutionary

constraints. Behav Ecol. 2013; art002. doi:10.1093/beheco/art002

3. Westneat DF, Wright J, Dingemanse NJ. The biology hidden inside residual within-

individual phenotypic variation. Biol Rev Camb Philos Soc. 2015;90: 729–743.

doi:10.1111/brv.12131

4. Réale D, Reader SM, Sol D, McDougall PT, Dingemanse NJ. Integrating animal

temperament within ecology and evolution. Biol Rev Camb Philos Soc. 2007;82:

291–318. doi:10.1111/j.1469-185X.2007.00010.x

5. Sih A, Bell AM, Johnson JC, Ziemba RE. Behavioral syndromes: an integrative

overview. Q Rev Biol. 2004;79: 241–277. doi:10.1086/422893

6. Budaev DS. How many dimensions are needed to describe temperament in animals: a

factor reanalysis of two data sets. Int J Comp Psychol. http://cogprints.org/5478/;

1998.

7. Spearman C. "General intelligence," objectively determined and measured. Am J

Psychol. 1904;15: 201–292. doi:10.2307/1412107

8. Beaujean AA. Latent variable modeling using R: A step-by-step guide. New York:

Routledge; 2014.

9. Bollen KA. Latent variables in psychology and the social sciences. Annu Rev

Psychol. 2002;53: 605–634. doi:10.1146/annurev.psych.53.100901.135239

10. Borsboom D. The attack of the psychometricians. Psychometrika. 2006;71: 425–440.

doi:10.1007/s11336-006-1447-6

11. Bollen K, Lennox R. Conventional wisdom on measurement: a structural equation

perspective. Psychol Bull. 1991;110: 305–314. doi:10.1037/0033-2909.110.2.305

12. Budaev SV. Using principal components and factor analysis in animal behaviour

research: caveats and guidelines. Ethology. 2010;116: 472–480. doi:10.1111/j.1439-

0310.2010.01758.x

13. Dingemanse NJ, Dochtermann N, Wright J. A method for exploring the structure of

behavioural syndromes to allow formal comparison within and between data sets.

Anim Behav. 2010;79: 439–450. doi:10.1016/j.anbehav.2009.11.024

14. Fabrigar LR, Wegener DT, MacCallum RC, Strahan EJ. Evaluating the use of

exploratory factor analysis in psychological research. Psychol Methods. 1999;4: 272–

299. doi:10.1037/1082-989X.4.3.272

15. Preacher KJ, MacCallum RC. Repairing Tom Swift's electric factor analysis

machine. Understanding Statistics. 2003;2: 13–43.

doi:10.1207/S15328031US0201_02

16. Araya-Ajoy YG, Dingemanse NJ. Characterizing behavioural "characters": An

evolutionary framework. Proc R Soc B. 2014;281: 20132645.

doi:10.1098/rspb.2013.2645

17. Arden R, Adams MJ. A general intelligence factor in dogs. Intelligence. 2016;55: 79–

85. doi:10.1016/j.intell.2016.01.008

18. van den Berg SM, Heuven HCM, van den Berg L, Duffy DL, Serpell JA. Evaluation

of the C-BARQ as a measure of stranger-directed aggression in three common dog

breeds. Appl Anim Behav Sc. 2010;124: 136–141.

doi:10.1016/j.applanim.2010.02.005

19. Martin JS, Suarez SA. Personality assessment and model comparison with behavioral

data: a statistical framework and empirical demonstration with bonobos (*Pan*

paniscus). Am J Primatol. 2017;9999: e22670. doi: 10.1002/ajp.22670

20. Bartholomew DJ. Factor analysis for categorical data. J R Stat Soc Series B Stat

Methodol. 1980;42: 293–321. http://www.jstor.org/stable/2985165

21. Markus KA, Borsboom D. Frontiers of test validity theory: Measurement, causation,

and meaning. 1st ed. New York, N.Y: Routledge; 2013.

22. Drasgow F. Study of the measurement bias of two standardized psychological tests.

Journal of Applied Psychology. 1987;72: 19–29. doi:10.1037/0021-9010.72.1.19

23. Mellenbergh GJ. Item bias and item response theory. Int J Educ Res. 1989;13: 127–

143. doi:10.1016/0883-0355(89)90002-5

24. Meredith W. Measurement invariance, factor analysis and factorial invariance.

Psychometrika. 1993;58: 525–543. doi:10.1007/BF02294825

25. Reise SP, Widaman KF, Pugh RH. Confirmatory factor analysis and item response

theory: two approaches for exploring measurement invariance. Psychol Bull.

1993;114: 552–566. doi:10.1037/0033-2909.114.3.552

26. Jansen BRJ, van der Maas HLJ. Statistical test of the rule assessment methodology by

latent class analysis. Dev Rev. 1997;17: 321–357. doi:10.1006/drev.1997.0437

27. Wicherts JM, Dolan CV. Measurement invariance in confirmatory factor analysis: an

illustration using IQ test performance of minorities. Educ Meas Issues Pract. 2010;29:

39–47. doi:10.1111/j.1745-3992.2010.00182.x

28. Goddard ME, Beilharz RG. Genetic and environmental factors affecting the

suitability of dogs as Guide Dogs for the Blind. Theor Appl Genet. 1982;62: 97–102.

doi:10.1007/BF00293339

29. Hsu Y, Serpell JA. Development and validation of a questionnaire for measuring

behavior and temperament traits in pet dogs. J Am Vet Med Assoc. 2003;223: 1293–

1300. doi:10.2460/javma.2003.223.1293

30. Scott JP, Fuller JL. Genetics and the social behavior of the dog. University of

Chicago Press; 2012.

31. Rayment DJ, Groef BD, Peters RA, Marston LC. Applied personality assessment in

domestic dogs: limitations and caveats. Appl Anim Behav Sci. 2015;163: 1–18.

doi:10.1016/j.applanim.2014.11.020

32. Persson ME, Wright D, Roth LSV, Batakis P, Jensen P. Genomic regions associated

with interspecies communication in dogs contain genes related to human social

disorders. Sci Rep. 2016;6. doi:10.1038/srep33439

33. Sundman A-S, Johnsson M, Wright D, Jensen P. Similar recent selection criteria

associated with different behavioural effects in two dog breeds. Genes Brain Behav.

2016;15: 750–756. doi:10.1111/gbb.12317

34. Barnard S, Marshall-Pescini S, Pelosi A, Passalacqua C, Prato-Previde E, Valsecchi

P. Breed, sex, and litter effects in 2-month old puppies' behaviour in a standardised

open-field test. Sci Rep. 2017;7: 1 – 8. doi:10.1038/s41598-017-01992-x.

35. Svartberg K. Breed-typical behaviour in dogs – historical remnants or recent

constructs? Appl Anim Behav Sci. 2006;96: 293 – 313. doi:

10.1016/j.applanim.2005.06.014

36. Jones AC, Gosling SD. Temperament and personality in dogs (*Canis familiaris*): a

review and evaluation of past research. Appl Anim Behav Sci. 2005;95: 1–53.

doi:10.1016/j.applanim.2005.04.008

37. Fratkin JL, Sinn DL, Patall EA, Gosling SD. Personality consistency in dogs: a meta-

analysis. PLOS ONE. 2013;8: e54907. doi:10.1371/journal.pone.0054907

38. Posluns JA, Anderson RE, Walsh CJ. Comparing two canine personality assessments:

convergence of the MCPQ-R and DPQ and consensus between dog owners and dog

walkers. Appl Anim Behav Sci. doi:10.1016/j.applanim.2016.12.013

39. Bennett SL, Litster A, Weng H-Y, Walker SL, Luescher AU. Investigating behavior

assessment instruments to predict aggression in dogs. Appl Anim Behav Sci.

2012;141: 139–148. doi:10.1016/j.applanim.2012.08.005

40. Mohan-Gibbons H, Weiss E, Slater M. Preliminary investigation of food guarding

behavior in shelter dogs in the United States. Animals. 2012;2: 331–346.

doi:10.3390/ani2030331

41. Mornement KM, Coleman GJ, Toukhsati SR, Bennett PC. Evaluation of the

predictive validity of the Behavioural Assessment for Re-homing K9's (B.A.R.K.)

protocol and owner satisfaction with adopted dogs. Appl Anim Behav Sci. 2015;167:

35–42. doi:10.1016/j.applanim.2015.03.013

42. Riemer S, Müller C, Virányi Z, Huber L, Range F. The predictive value of early

behavioural assessments in pet dogs: a longitudinal study from neonates to adults.

PLOS ONE. 2014;9: e101237. doi:10.1371/journal.pone.0101237

43. Wilsson E, Sundgren P-E. Behaviour test for eight-week old puppies: heritabilities of

tested behaviour traits and its correspondence to later behaviour. Appl Anim Behav

Sci. 1998;58: 151–162. doi:10.1016/S0168-1591(97)00093-2

44. Goodloe LP, Borchelt PL. Companion dog temperament traits. J Appl Anim Welf

Sci. 1998;1: 303–338. doi:10.1207/s15327604jaws0104_1

45. Jones AC. Development and validation of a dog personality questionnaire. Doctoral

Thesis. University of Texas at Austin. 2008. http://gosling.psy.utexas.edu/wp-

content/uploads/2014/10/Amanda-Claire-Jones-Diss-2008.pdf

46. Orritt R. Dog bites: a complex public health issue. Vet Rec. 2015;176: 640–641.

doi:10.1136/vr.h3215

47. Svartberg K. Shyness-boldness predicts performance in working dogs. Appl Anim

Behav Sci. 2002;79: 157–174. doi:10.1016/S0168-1591(02)00120-X

48. Taylor KD, Mills DS. The development and assessment of temperament tests for

adult companion dogs. J Vet Behav. 2006;1: 94–108. doi:10.1016/j.jveb.2006.09.002

49. Haverbeke A, Plujimakers J, Diederich C. Behavioral evaluations of shelter dogs:

literature review, perspectives, and follow-up within the European member state's

legislation with emphasis on the Belgian situation. J Vet Behav. 2015;10: 5 – 11. doi:

10.1016/j.jveb.2014.07.004

50. Patronek GJ, Bradley J. No better than flipping a coin: reconsidering canine behavior

evaluations in animal shelters. J Vet Behav. 2016;15: 66–77.

doi:10.1016/j.jveb.2016.08.001

51. Edelen MOE, Reeve BB. Applying item response theory (IRT) modeling to

questionnaire development, evaluation, and refinement. Qual Life Res. 2007;16: 5 –

18. doi: 10.1007/s11136-007-9198-0

52. Casey RA, Loftus B, Bolster C, Richards GJ, Blackwell EJ. Human directed

aggression in domestic dogs (*Canis familiaris*): occurrence in different contexts and

risk factors. Appl Anim Behav Sci. 2014;152: 52–63.

doi:10.1016/j.applanim.2013.12.003

53. Hsu Y, Sun L. Factors associated with aggressive responses in pet dogs. Appl Anim

Behav Sci. 2010;123: 108–123. doi:10.1016/j.applanim.2010.01.013

54. Sherman CK, Reisner IR, Taliaferro LA, Houpt KA. Characteristics, treatment, and

outcome of 99 cases of aggression between dogs. Appl Anim Behav Sci. 1996;47:

91–108. doi:10.1016/0168-1591(95)01013-0

55. Asp HE, Fikse WF, Nilsson K, Strandberg E. Breed differences in everyday

behaviour of dogs. Appl Anim Behav Sci. 2015;169: 69–77.

doi:10.1016/j.applanim.2015.04.010

56. Voith VL, Trevejo R, Dowling-Guyer S, Chadik C, Marder A, Johnson V, et al.

Comparison of visual and DNA breed identification of dogs and inter-observer

reliability. Sociology. 2013;3: 17–29. doi: 10.5923/j.sociology.20130302.02

57. Olson KR, Levy JK, Norby B, Crandall MM, Broadhurst JE, Jacks S, Barton RC,

Zimmerman MS. Inconsistent identification of pit bill-type dogs by shelter staff. Vet

J. 2015;206: 197 – 202. doi:10.1016/j.tvjl.2015.07.019

58. Simpson RJ, Simpson K, VanKavage L. Rethinking dog breed identification in

veterinary practice. J Am Vet Med Assoc. 2012;241: 1163 – 1166. doi:

10.2460/javma.241.9.1163

59. Owczarczak-Garstecka, SC, Burman OHP. Can sleep and resting behaviour be used

as indicators of welfare in shelter dogs (*Canis lupus familiaris*)? PLOS ONE.

2016;11: e0163620. doi:10.1371/journal.pone.0163620

60. R Development Core Team. R: a language and environment for statistical computing.

Vienna, Austria: R Foundation for Statistical Computing; 2016. https://www.r-

project.org/

61. Honaker J, King G, Blackwell M. Amelia: A program for missing data. 2015. Version

1.7.4. https://cran.r-project.org/web/packages/Amelia/vignettes/amelia.pdf

62. Rosseel, Y., Oberski, D., Byrnes, J., Vanbrabant, L., Savalei, V., Merkle, E., et al,

Lavaan: Latent Variable Analysis. 2016. Version 0.5-22. https://cran.r-

project.org/web/packages/lavaan/

63. Muthén B, Christoffersson A. Simultaneous factor analysis of dichotomous variables

in several groups. Psychometrika. 1981;46: 407–419. doi:10.1007/BF02293798

64. Muthén B. A general structural equation model with dichotomous, ordered

categorical, and continuous latent variable indicators. Psychometrika. 1984;49: 115–

132. doi:10.1007/BF02294210

65. Jorgensen TD, Pornprasertmanit S, Miller P, Schoemann A, Rosseel Y, Quick C, et

al. semTools: useful tools for structural equation modeling. 2016.

https://rdrr.io/cran/semTools/

66. Hermida R. The problem of allowing correlated errors in structural equation

modeling: concerns and considerations. Comp Method Soc Sci. 2015;3: 05–17.

67. Kamata A. Item analysis by the hierarchical generalized linear model. J Educ Meas.

2001;38: 79–93. doi:10.1111/j.1745-3984.2001.tb01117.x

68. Van den Noortgate W, De Boeck P. Assessing and explaining differential item

functioning using logistic mixed models. J Educ Behav Stat. 2005;30: 443–464.

69. Nakagawa S, Schielzeth H. Repeatability for Gaussian and non-Gaussian data: a

practical guide for biologists. Biol Rev Camb Philos Sci. 2010;85: 935–956.

doi:10.1111/j.1469-185X.2010.00141.x

70. Stan Development Team. Stan modeling language users guide and reference manual.

2016. Version 2.15.0. https://github.com/stan-

dev/stan/releases/download/v2.14.0/stan-reference-2.14.0.pdf

71. Kruschke JK. Bayesian data analysis. Wiley Interdiscip Rev: Cogn Sci. 2010;1: 658–

676. doi:10.1002/wcs.72

72. Vehtari A, Gelman A, Gabry J, Piironen J, Goodrich B. Loo: Efficient leave-one-out

cross-validation and WAIC for Bayesian models. 2016. Version 1.0.0. https://cran.r-

project.org/web/packages/loo/loo.pdf

73. Kruschke J. Doing Bayesian data analysis: A tutorial with R, JAGS, and Stan. 2nd ed.

Academic Press; 2014.

74. Chen JJ, Tsong Y, Kang S-H. Tests for equivalence or noninferiority between two

proportions. Drug Inf J. 2000;34: 569–578. doi:10.1177/009286150003400225

75. Budaev S, Brown C. Personality traits and behaviour. In: Brown C, Laland K, Krause

J, editors. Fish cognition and behavior. Wiley-Blackwell; 2011. pp. 135–165.

76. Campbell DT, Fiske DW. Convergent and discriminant validation by the multitrait-

multimethod matrix. Psychol Bull. 1959;56: 81–105. doi:10.1037/h0046016

77. Carter AJ, Feeney WE, Marshall HH, Cowlishaw G, Heinsohn R. Animal personality:

what are behavioural ecologists measuring? Biol Rev Camb Philos Soc. 2013;88:

465–475. doi:10.1111/brv.12007

78. Podsakoff PM, MacKenzie SB, Lee J-Y, Podsakoff NP. Common method biases in

behavioral research: a critical review of the literature and recommended remedies. J

Appl Psychol. 2003;88: 879–903. doi:10.1037/0021-9010.88.5.879

79. Cole DA, Ciesla JA, Steiger JH. The insidious effects of failing to include design-

driven correlated residuals in latent-variable covariance structure analysis. Psychol

Methods. 2007;12: 381–398. doi:10.1037/1082-989X.12.4.381

80. Yen WM. Scaling performance assessments: strategies for managing local item

dependence. J Educ Meas. 1993;30: 187–213. doi:10.1111/j.1745-

3984.1993.tb00423.x

81. Carter AJ, Marshall HH, Heinsohn R, Cowlishaw G. How not to measure boldness:

novel object and antipredator responses are not the same in wild baboons. Anim

Behav. 2012;84: 603–609. doi:10.1016/j.anbehav.2012.06.015

82. Cramer AOJ, van der Sluis S, Noordhof A, Wichers M, Geschwind N, Aggen SH, et

al. Dimensions of normal personality as networks in search of equilibrium: you can't

like parties if you don't like people. Eur J Pers. 2012;26: 414–431.

doi:10.1002/per.1866

83. Schmittmann VD, Cramer AOJ, Waldorp LJ, Epskamp S, Kievit RA, Borsboom D.

Deconstructing the construct: a network perspective on psychological phenomena.

New Ideas Psychol. 2013;31: 43–53. doi:10.1016/j.newideapsych.2011.02.007

84. Hamaker EL, Nesselroade JR, Molenaar PCM. The integrated trait-state model. J Res

Pers. 2007;41: 295–315. doi:10.1016/j.jrp.2006.04.003

85. Duckworth RA. Neuroendocrine mechanisms underlying behavioral stability:

implications for the evolutionary origin of personality. Ann N Y Acad Sci.

2015;1360: 54–74. doi:10.1111/nyas.12797

86. Koski SE. How to measure animal personality and why does It matter? Integrating the

psychological and biological approaches to animal personality. In: Inoue-Murayama

M, Kawamura S, Weiss A, editors. From Genes to Animal Behavior. Springer Japan;

2011. pp. 115–136. doi:10.1007/978-4-431-53892-9_5

87. Borsboom D. Measuring the mind: Conceptual issues in contemporary

psychometrics. Cambridge University Press; 2005.

88. Maul A, Torres Irribarra D, Wilson M. On the philosophical foundations of

psychological measurement. Measurement. 2016;79: 311–320.

doi:10.1016/j.measurement.2015.11.001

89. Molenaar PCM. A manifesto on psychology as idiographic science: bringing the

person back into scientific psychology, this time forever. Measurement. 2004;2: 201–

218. doi:10.1207/s15366359mea0204_1

90. Adolf J, Schuurman NK, Borkenau P, Borsboom D, Dolan CV. Measurement

invariance within and between individuals: a distinct problem in testing the

equivalence of intra- and inter-individual model structures. Front Psychol. 2014;5.

doi:10.3389/fpsyg.2014.00883

91. Stamps JA, Briffa M, Biro PA. Unpredictable animals: individual differences in

intraindividual variability (IIV). Anim Behav. 2012;83: 1325–1334.

doi:10.1016/j.anbehav.2012.02.017

92. Biro PA, Stamps JA. Using repeatability to study physiological and behavioural

traits: ignore time-related change at your peril. Anim Behav. 2015;105: 223–230.

doi:10.1016/j.anbehav.2015.04.008

93. Borsboom D, Cramer AO, Kievit RA, Zand Scholten A, Franic S. The end of

construct validity. In Lissitz, R, editor. The concept of validity. Information Age

Publishers; 2010. pp. 135–170.

Supporting Information

Table S1. Counts of aggression per context. The number of dogs who had 0, 1, and > 1 observations of aggression while at the shelter.

Table S2. Tetrachoric correlations between aggression contexts. Tetrachoric correlations between aggression contexts on the raw binary data, before the multiple imputation. Abbreviations used: HND (*Handling*); FPL (*Interactions with familiar people*); UPL (*Interactions with unfamiliar people*); KD (*In kennel towards dogs*); KP (*In kennel towards people*); OKD (*Out of kennel towards dogs*); OKP (*Out of kennel towards people*); EAT (*Eating food*); TOY (*Interactions with toys*); DM (*Interactions with male dogs*); DF (*Interactions with female dogs*).

Table S3. Bayesian hierarchical model parameter estimates for aggression towards people in different contexts. Mean and 95% highest density interval (HDI) estimates for all parameters from the Bayesian hierarchical logistic model assessing measurement invariance for contexts reflecting aggressiveness towards people. Differences between levels of categorical variables are indicated by '.v.' in the parameter name; interactions are denoted with '*' in the parameter name. The decision rule for each parameter is given except for those variables not interpreted inferentially: YES = 95% HDI falls completely outside the region of practical equivalence (ROPE); NULL = 95% HDI falls completely inside the ROPE; ROPE = 95% HDI partly covers the ROPE.

Table S4. Bayesian hierarchical model parameter estimates for aggression towards

dogs in different contexts. Mean and 95% highest density interval (HDI) estimates for

all parameters from the Bayesian hierarchical logistic model assessing measurement

invariance for contexts reflecting aggressiveness towards dogs. Differences between

levels of categorical variables are indicated by '.v.' in the parameter name; interactions

are denoted with '*' in the parameter name. The decision rule for each parameter is given

except for those variables not interpreted inferentially: YES = 95% HDI falls completely

outside the region of practical equivalence (ROPE); NULL = 95% HDI falls completely

inside the ROPE; ROPE = 95% HDI partly covers the ROPE.

1005