***Lysinibacillus fusiformis* M5 induces increased complexity in *Bacillus subtilis* 168**

colony biofilms via hypoxanthine

Ramses Gallegos-Monterrosa[1], Stefanie Kankel[1], Sebastian Götze[2], Robert Barnett[2],

Pierre Stallforth[2,*], Ákos T. Kovács[1,*,#]

[1] Terrestrial Biofilms Group, Institute of Microbiology, Friedrich Schiller University Jena,

Jena, Germany.

[2] Junior Research Group Chemistry of Microbial Communication, Leibniz Institute for

Natural Product Research and Infection Biology, Hans Knöll Institute – HKI, Jena,

Germany.

Running Title: *L. fusiformis* M5 interaction with *B. subtilis* 168

* Address correspondence to: Ákos T. Kovács <atkovacs@dtu.dk> (on microbiology and

strain requests) or Pierre Stallforth < pierre.stallforth@leibniz-hki.de> (on isolation and

identification of metabolites)

[#] Affiliation from July 1st 2017: Department of Biotechnology and Biomedicine, Technical

University of Denmark, Denmark

ABSTRACT

In recent years, biofilms have become a central subject of research in the fields of

microbiology, medicine, agriculture, or systems biology amongst others. The

sociomicrobiology of multispecies biofilms, however, is still poorly understood. Here, we

report a screening system that allowed us to identify soil bacteria, which induce

architectural changes in biofilm colonies when cocultured with *B. subtilis*. We identified

the soil bacterium *Lysinibacillus fusiformis* M5 as inducer of wrinkle-formation in *B.*

subtilis colonies mediated by a diffusible signaling molecule. This compound was

isolated by bioassay-guided chromatographic fractionation. The elicitor was identified to

be the purine hypoxanthine using mass spectrometry and nuclear magnetic resonance

(NMR) spectroscopy. We show that the induction of wrinkle formation by hypoxanthine is

not dependent on signal recognition by the histidine kinases KinA, KinB, KinC, and KinD,

which are generally involved in phosphorylation of the master regulator Spo0A.

Likewise, we show that hypoxanthine signaling does not induce the expression of

biofilm-matrix related operons *epsA-O* and *tasA-sipW-tapA*. Finally, we demonstrate that

the purine permease PbuO, but not PbuG, is necessary for hypoxanthine to induce an

increase in wrinkleality of *B. subtilis* biofilm colonies. Our results suggest that

hypoxanthine-stimulated wrinkle development is not due to a direct induction of biofilm-

related gene expression, but rather caused by the metabolic changes derived from the

excess of hypoxanthine in *B. subtilis* cells.

IMPORTANCE

Biofilms are a bacterial lifestyle with high relevance regarding diverse human activities. Biofilms can be favorable, for instance in crop protection. In nature, biofilms are commonly found as multispecies communities displaying complex social behaviors and characteristics. The study of interspecies interactions will thus lead to a better understanding and use of biofilms as they occur outside laboratory conditions. Here, we present a screening method suitable for the identification of multispecies interactions, and showcase *L. fusiformis* as a soil bacterium that is able to live alongside *B. subtilis* and modify the architecture of its biofilms.

INTRODUCTION

Biofilms are microbial populations formed by cells living in high density communities attached to biotic or abiotic surfaces. These cells are often encased in a matrix of polymeric substances that provide the whole population with an increased resistance against environmental stress (1, 2). Furthermore, these communities exhibit highly complex structural organization and social behavior. Thus, biofilms have become an increasingly studied research subject by microbiologists, especially when it became apparent that this lifestyle is widely spread among bacteria and involved in a multitude of biological processes (3, 4). Although much attention has been given to medically-relevant biofilms (5, 6), scientists have also studied biofilms in the context of industrial applications (7), bioremediation (8), and crop protection (9).

In nature, biofilms rarely occur as single-species populations, but rather as mixed communities of diverse bacteria and other microorganisms. This leads to complex interactions between the different members of the community, usually involving communication networks based on chemical signals (10). Additionally, microorganisms need to sense and efficiently adapt to a wide array of environmental cues in order to efficiently regulate biofilm formation (11).

Bacillus subtilis is a soil-dwelling Gram-positive bacterium that has become a model for biofilm research. On agar plates, *B. subtilis* can form large colonies with remarkably complex architecture, while in liquid medium it forms robust floating biofilms known as pellicles. Both forms of biofilms are characterized by a wrinkled surface, which has been associated to the production of exopolysaccharides, biofilm maturation, and mechanical

forces concomitant with an increased population complexity (12–14). Moreover, these

biofilms display intricate cell heterogeneity, i.e. some cells become matrix producers,

while others either produce exoenzymes to harvest nutrients, or form resistant structures

known as spores (15, 16). The development of this population heterogeneity is regulated

by a complex gene regulatory network involving various sensing kinases i.e. Kin

kinases, DegS, and ComP and their concomitant response regulators Spo0A, DegU,

and ComA (15, 17), respectively.

Biofilms produced by *B. subtilis* are not only a good research model, they are currently

applied in crop protection (18, 19), and spores of this organism are readily

commercialized as a biocontrol agent for agriculture. *B. subtilis* is a prolific producer of

secondary metabolites and many potent antimicrobial compounds inhibiting both

bacteria and fungi have been identified (20–22). In addition, it has also been shown that

B. subtilis activates biofilm-related gene expression in response to chemicals produced

by other bacteria closely related to it, for instance by other members of the *Bacillus*

genus (23). Interestingly, the signaling role of the molecules can be independent from

other effects of the compounds, as in the case of antimicrobial thiazolyl peptides, which

can induce biofilm-matrix production in *B. subtilis* even when separated from their

antibiotic activity (24). Moreover, other organisms such as *Pseudomonas protegens*, are

able to inhibit cell differentiation and biofilm gene expression in *B. subtilis*, possibly as a

competition strategy during root colonization (25).

B. subtilis successfully inhabits a congested and competitive ecological niche (26–28),

and it is to be expected that this organism has a finely tuned regulatory network

governing community behavior. Therefore, further study of the signaling mechanisms
that influence *B. subtilis* biofilm formation may enhance the use of this organism, both in
biotechnological applications as well as a research model. However, the identification of
signals that induce biofilm formation is a poorly investigated field of study, possibly due
to the greater general interest in the removal of biofilms in various medical and industrial
settings (5, 29–31). Thus, we have established a co-cultivation-based screening method
to identify signaling molecules that induce biofilm formation in *B. subtilis*. Using this
system, we identified ecologically relevant soil bacteria that are able to induce the
formation of large wrinkles in colony biofilms of *B. subtilis*. The majority of these bacteria
are members of the family *Bacillaceae*, to which *B. subtilis* belongs. The strain with the
clearest wrinkle-inducing effect was identified as *Lysinibacillus fusiformis* M5. The
observed effect on *B. subtilis* is dependent on a diffusible signaling molecule, which was
identified as hypoxanthine using bioassay-guided fractionation and subsequent structure
elucidation using various spectroscopic and spectrometric methods. The induction of
wrinkles by hypoxanthine was not dependent on Kin kinases signal transduction, and
the expression levels of operons responsible for the production of biofilm matrix
components, *epsA-O* and *tapA-sipW-tasA*, remained unaffected. We show that uptake
of hypoxanthine by permease PbuO is necessary for the increased wrinkleality of *B.
subtilis* biofilms colonies, where wrinkleality refers to the degree of wrinkle formation as
defined by Spiers (32). We therefore suggest that hypoxanthine induces the formation of
wrinkles by introducing a metabolic change in *B. subtilis* cells, rather than by direct
stimulation of biofilm-related gene expression.

RESULTS

Screening of soil bacteria that induce structural changes in colonies of *B. subtilis.*

We screened a collection of 242 strains isolated from two distinct soil sampling sites in

Mexico, in order to identify bacteria that are able to induce biofilm formation or increased

complex colony architecture of *B. subtilis*. Importantly, our assay aimed to discover

alterations in biofilm colony architecture that was different from the previously described

method that identified soil-derived microbes that induce gene expression related to

biofilm formation of *B. subtilis* (23). While some *B. subtilis* strains, such as *B. subtilis*

NCIB 3610, easily and spontaneously form biofilms, we used a strain that would form

architecturally complex colonies only in the presence of specific inducers or nutrient rich

conditions. Thus, even weak biofilm-inducing effects would not be overseen. We used

B. subtilis 168 (Jena stock), a strain that can only form architecturally complex colonies

when grown on glucose-rich medium or exposed to signaling molecules as present in

plant root exudates (33). The strain of *B. subtilis* used for the assay (TB48) carried a

$P_{hyperspank}$-mKATE reporter fusion in order to facilitate the identification of *B. subtilis* from

the soil strains in mixed colonies. The reporter strain was mixed in different ratios with

the bacterial isolates and allowed to form colonies on 2×SG medium for 72 hours. Single

strain colonies of *B. subtilis* and the soil-derived isolates were also grown as neighboring

colonies under the same conditions, inoculated with a spatial distance of 5 mm between

each other to examine their interactions. The majority of these interactions resulted in

the apparent killing of one strain by the other, producing a colony identical to the pure

culture colony of the surviving partner (data not shown). However, 36 soil strains were

able to grow alongside *B. subtilis*, mainly by creating a colony where the strains

segregate in sectors. Interestingly, the *B. subtilis* sectors of these mixed colonies

showed a marked increase in the formation of wrinkles compared to its pure colonies

(Fig. 1). Furthermore, when single strain colonies of these soil-derived isolates were

grown close to *B. subtilis* colonies, the induction of wrinkle formation could be observed

in the areas of the *B. subtilis* colony that are closest to the soil strain, but not in the other

regions of the colony (Fig. 1). These results suggest that the aforementioned bacteria

produce signals that can induce an increased architectural complexity in *B. subtilis*

colony biofilms.

Structural changes in *B. subtilis* colonies are induced by diffusible signals

produced by soil bacteria.

In order to elucidate if the observed induction of wrinkle formation is caused by a

diffusible signal molecule or due to direct cell-cell interactions, we designed an assay to

test the cell-free supernatants of the selected soil strains. In this assay, we used cotton

discs infused with cell-free supernatant to simulate colonies of the tested soil strains.

B. subtilis was inoculated at a distance of 5 mm next to the cotton discs and allowed to

grow for 72 hours. Over this period, the growing colony surrounded the cotton discs,

getting into contact with the freely diffusing compounds in the supernatants.

Using this assay, we observed that the cell-free supernatants of four bacterial isolates

were able to induce efficiently the formation of wrinkles in the adjacent *B. subtilis* colony.

Importantly, this phenomenon was observed in the periphery of the cotton discs, but not

in the areas of the colony farther away from it (Fig. 2). We note that neither the cell-free

supernatant of *B. subtilis* itself nor the medium used to obtain the supernatants showed

the capacity to induce increased wrinkleality in *B. subtilis* colonies under our tested

conditions (Fig. 2). We concluded that the induction of wrinkle formation was due to a

diffusible signaling molecule produced by these soil organisms.

Using 16S rRNA locus sequencing, we characterized those soil strains whose

supernatant could best stimulate wrinkle formation in *B. subtilis*. The majority of the

sequenced strains were found to be members of the same phylogenetic family as *B.*

subtilis, such as *Bacillus pumilus* or *L. fusiformis*. The only exception was a strain

identified as the γ-proteobacterium *Acinetobacter variabilis.*

Hypoxanthine was identified in the supernatant to be responsible for wrinkle

induction.

The clearest and strongest induction of wrinkle formation was observed in the

supernatant of the soil derived strain identified as *L. fusiformis* M5. For this reason, we

decided to further investigate the respective signaling molecule produced by this

bacterium using a colony wrinkleality assay. Bio-assay-guided fractionation allowed then

to identify a compound from *L. fusiformis* M5 that induced a similar phenotype as

observed when *B. subtilis* and *L. fusiformis* M5 were co-cultured. To this end,

supernatant of *L. fusiformis* M5 was lyophilized and fractionated using Sephadex G20 as

stationary phase. Each fraction was applied to cotton discs and placed on an agar plate

in the vicinity of *B. subtilis.* The fraction that induced wrinkle formation was sub-

fractionated by high-performance liquid chromatography (HPLC) using a hypercarb

column as stationary phase. Repeating this procedure led to the isolation of a

homogeneous compound whose structure was subsequently elucidated via a

combination of nuclear magnetic resonance (NMR) spectroscopy and high-resolution

mass spectrometry (HR-MS). Finally, hypoxanthine was identified as the inducer of

wrinkle formation. We further validated our findings using commercial hypoxanthine,

which showed the same properties as the isolated hypoxanthine (Fig. 3).

Hypoxanthine signaling is not mediated by Kin kinases

In *B. subtilis*, the transcriptional regulator Spo0A controls the expression of several

biofilm-related operons, including those responsible for the production of the

exopolysaccharide and protein components of the biofilm matrix (*epsA-O* and *tapA-*

sipW-tasA, respectively) (17). Five sensor kinases (KinA, KinB, KinC, KinD, and KinE)

have been identified in *B. subtilis*, while four of them (KinA-D) are known to activate

Spo0A via a phosphorelay depending on the environmental signals (11, 15, 34). We

wanted to determine if any of these kinases is involved in hypoxanthine-mediated

induction of wrinkles. Therefore, we compared the effect that the supernatant of *L.*

fusiformis M5 has on kinase-mutant strains of *B. subtilis* using our colony wrinkleality

assay. We expected that, should one of these kinases be responsible for sensing

hypoxanthine, the corresponding mutant strain would no longer show increased colony

wrinkleality. Interestingly, all the mutant strains were still able to develop highly wrinkled

colonies when exposed to the supernatant (Fig. 4). This result indicates that

hypoxanthine-mediated induction of increased architectural complexity is not mediated

by the activation of Spo0A via the Kin kinases.

Expression levels of genes responsible for biofilm matrix production are not

affected by hypoxanthine signaling.

The *epsA-O* and *tapA-sipW-tasA* operons are related to the production of the

exopolysaccharide and protein components of the *B. subtilis* biofilm matrix, respectively.

Changes in the expression levels of these operons are associated to a maturating

biofilm, and show spatiotemporal variation during its development (16, 35, 36).

To further examine if biofilm matrix-related genes are involved in the induction of

wrinkles by hypoxanthine, we monitored the expression of P_{epsA}-*gfp* and P_{tapA}-*gfp*

fluorescent reporter fusions in colonies of *B. subtilis* using our colony wrinkleality assay.

Fluorescence emission was examined only in the sections of the colonies directly

adjacent to the cotton discs at 3 time points: (i) when *B. subtilis* has encircled the discs

(40 hours), (ii) when the colony started to expand from the disc and showed the onset of

wrinkle formation (50 hours), and (iii) when the colony has developed wrinkles and

expanded (65 hours). The examined strains also carried a $P_{hyperspank}$-*mKATE* reporter

fusion to adjust for colony growth. Under these conditions, the expression from P_{epsA} and

P_{tapA} in these colonies showed no statistical differences when exposed to cotton discs

infused with 2×SG medium or supernatant of *B. subtilis,* as compared to those infused

with supernatant of *L. fusiformis* M5 (one-way ANOVA: $P<0.05$, n=4-8 independent

colonies) (Fig. 5).

Taken together, these results suggest that the increased colony wrinkleality induced by

hypoxanthine is not directly associated to biofilm matrix-related gene expression.

The permease PbuO is necessary for hypoxanthine-induced development of

wrinkles.

We hypothesized that the observed induction of wrinkle formation may be due to

metabolic effects on *B. subtilis* cells derived from an excess of available hypoxanthine

provided by the culture supernatant of *L. fusiformis* M5, rather than to direct signal-

dependent expression of biofilm-related genes. In this case, hypoxanthine alone would

be sufficient to induce increased wrinkleality, and its uptake by *B. subtilis* would be

necessary for this phenomenon.

Different concentrations of hypoxanthine (as solution in 0.05 N NaOH) were able to

induce the formation of wrinkles in *B. subtilis* colonies as efficiently as the supernatant of

L. fusiformis M5. To test whether hypoxanthine internalization is required for the

observed wrinkle induction in *B. subtilis*, we analyzed mutant strains of *B. subtilis* that

lack *pbuG* or *pbuO*. PbuG is a previously described hypoxanthine/guanine permease

(37), while PbuO is a protein paralogous to PbuG annotated as a putative purine

permease (see SubtiWiki: http://subtiwiki.uni-goettingen.de/index.php). These strains

were studied with our colony wrinkleality assay. The hypoxanthine inducing effect

disappeared when *pbuO* alone, or in combination with *pbuG,* was deleted (Fig. 6).

These final results demonstrated that hypoxanthine uptake is important for induction of

wrinkle formation and PbuO is mainly responsible for the observed effect under our

experimental conditions.

DISCUSSION

In this work, we identified a chemical sensing mechanism between *B. subtilis* and other soil bacteria that promotes architectural complexity in colony biofilms. We devised a screening system that allowed us to analyze a collection of soil bacteria, selecting those that could form stable multispecies communities with *B. subtilis*. Using this screening system, we identified *L. fusiformis* M5 as a bacterium inducing an increase of colony wrinkleality in *B. subtilis* via hypoxanthine as chemical cue.

Hypoxanthine is a purine that plays an important role in the pentose phosphate salvage pathway, which is a mechanism for cells to interconvert nucleosides and nucleobases according to their metabolic needs (38, 39). In *B. subtilis*, hypoxanthine is particularly relevant in this pathway because it can be taken up by cells and used as a substrate by phosphoribosyl-transferases in order to produce inosine monophosphate (IMP), which in turn is converted to adenosine monophosphate or guanine monophosphate (37) (also see SubtiWiki Pathways: http://subtiwiki.uni-goettingen.de/apps/pathway.php?pathway=2). The role of hypoxanthine in eukaryotic cell metabolism has been extensively investigated. In humans, it has been studied in the context of diseases such as gout, Lesch–Nyhan disease, and endothelial cell injury of cardiovascular diseases. Although these conditions have different etiologies and clinical evolution, they have in common an excessive accumulation of hypoxanthine and uric acid that leads to oxidative-stress-induced apoptosis (40–42). In bacteria, hypoxanthine has been mainly studied in relation to DNA damage and mutagenesis due to spontaneous deamination of adenine, which yields hypoxanthine and leads to AT-to-GC transitions after DNA replication (43). In *B. subtilis*, hypoxanthine has been studied both

related to purine metabolism (37), and DNA damage and repair (44). To the best of our

knowledge, this is the first time that hypoxanthine has been reported as a mediator of

interactions in *B. subtilis* biofilms.

L. fusiformis is a free-living bacterium that can be isolated from soil and has been

studied due to its production of interesting secondary metabolites and bioremediation

potential (45–47). Here, we have identified a strain of *L. fusiformis* able to produce and

excrete hypoxanthine in sufficient levels to induce the formation of wrinkles in biofilm

colonies of *B. subtilis*. We found no evidence that this phenomenon is dependent on Kin

kinases signal transduction, and the expression levels of the matrix-component-related

operons *epsA-O* and *tapA-sipW-tasA* remained equal when *B. subtilis* was exposed to

the supernatant of *L. fusiformis* M5. However, a knock-out mutant of the

hypoxanthine/guanine permease, PbuO, lost the ability to form heavily wrinkled colonies.

A *pbuG*⁻ mutant strain showed comparable or in rare cases slightly reduced wrinkle

formation, while a *pbuO*⁻ strain completely lost the ability to form wrinkled colonies when

exposed to hypoxanthine. Based on these results, we conclude that hypoxanthine

induces the formation of wrinkles in colony biofilms of *B. subtilis* not by inducing the

expression of biofilm-related genes, but rather by metabolic effects derived from the

excess of available hypoxanthine. In this regard, we note that an excess of hypoxanthine

can cause oxidative stress and cell death in eukaryotic cells (40, 41), while localized cell

death can lead to the formation of wrinkles in colonies of *B. subtilis* by providing an

outlet for mechanical forces that shape the developing biofilm (13). Thus, we

hypothesize that the hypoxanthine provided by *L. fusiformis* induces cell stress and

possibly cell death in *B. subtilis*, which leads to the formation of wrinkles as a

mechanical consequence. This is in accordance with the fact that the observed

development of wrinkles only occurs in the interaction zone between colonies of these

organisms (Fig. 1), or in the proximity of the cotton discs during our colony wrinkleality

assays (Figs. 4 and 6). This development of wrinkles does not happen in the rest of the

B. subtilis colony, presumably due to a lower concentration of diffused hypoxanthine.

Importantly, this induction of increased wrinkleality in *B. subtilis* may be a consequence

of regular metabolic processes of *L. fusiformis*, rather than a canonical signaling

mechanism intended to elicit a response in the receiver. Regarding hypoxanthine

production by *L. fusiformis* M5, we have previously sequenced this strain, finding genes

homologous to those known in other organisms to be responsible for hypoxanthine

synthesis and export (48), although more research is needed to establish its production

yield. However, we note that hypoxanthine can be produced by spontaneous

deamination of adenine, such as that present in the surroundings of decaying cells (49).

Unfortunately, our efforts to transform *L. fusiformis* M5 failed, preventing us to construct

mutants with altered hypoxanthine production.

In recent years, biofilm research has grown from an incipient field to a major area of

microbiological interest. Due to the high cell density of biofilms, social interactions are an

inherent characteristic of these microbial populations, regardless of whether they are

formed as single or multi-species communities (10, 27, 50, 51). The interactions

between the organisms forming a biofilm are therefore an important aspect of this

research field, since they shape the development of these communities, be it by

intraspecies signaling, interspecies communications, or chemical cues derived from the

metabolism of community members, such as the case presented here. Further study of

the sociomicrobiology of biofilms will lead to an increased understanding of these

communities as they form in nature, better enabling us to eliminate them when they are

noxious to human activities, or to promote them when needed for biotechnological

applications.

MATERIAL AND METHODS

Strains, media, and general culture conditions.

All strains used in this study are listed in Table 1. When fresh cultures were needed, these strains were pre-grown overnight in Lysogeny broth medium (LB-Lennox, Carl Roth; 10 g L^{-1} tryptone, 5 g L^{-1} yeast extract, and 5 g L^{-1} NaCl) at 37°C and shaken at 225 r.p.m. LB medium was used for all *B. subtilis* and *Escherichia coli* transformations, and to screen soil samples. 2×SG medium (16 g L^{-1} nutrient broth (Difco), 2 g L^{-1} KCl, 0.5 g L^{-1} MgSO$_4$·7H$_2$O, 1mM Ca(NO$_3$)$_2$, 0.1mM MnCl$_2$·4H$_2$O, 1 μM FeSO$_4$, and 0.1% glucose) (52) was used to grow cultures intended for supernatant production. This medium was also used for all strain interaction assays and wrinkle-induction assays. Tryptone Soya broth (CASO-Bouillon, AppliChem; 2.5 g L^{-1} glucose, 5 g L^{-1} NaCl, 2.5 g L^{-1} buffers (pH 7.3), 3 g L^{-1} soya peptone, and 17 g L^{-1} tryptone) was used for screening soil samples. GCHE medium (1% glucose, 0.2% glutamate, 100mM potassium phosphate buffer (pH: 7), 3mM trisodium citrate, 3mM MgSO$_4$, 22 mg L^{-1} ferric ammonium citrate, 50 mg L^{-1} L-tryptophan, and 0.1% casein hydrolysate) was used to induce natural competence in *B. subtilis* (53). Our developed Gallegos Rich medium was used to grow *Lactococcus lactis* MG1363, in order to purify pMH66: 21 g L^{-1} tryptone, 5 g L^{-1} yeast extract, 8.3 g L^{-1} NaCl, 3 g L^{-1} soya peptone, 2.6 g L^{-1} glucose, and 2.5 g L^{-1} MgSO$_4$·7H$_2$O. Overnight cultures of *L. lactis* were incubated at 30°C without shaking. Media were supplemented with Bacto agar 1.5 % when solid plates were needed. Antibiotics were used at the following final concentrations: kanamycin, 10 μg mL^{-1}; chloramphenicol, 5 μg mL^{-1}; erythromycin-lincomycin, 0.5 μg mL^{-1} and 12.5 μg mL^{-1} respectively; ampicillin, 100 μg mL^{-1}; spectinomycin, 100 μg mL^{-1}; tetracycline, 10 μg mL^{-1}. Specific growth conditions are described in the corresponding methods section.

Importantly, all 2×SG plates used in this study were prepared with 25 mL of medium,

and dried for a minimum of 20 minutes before use. Insufficient drying resulted in

excessive colony expansion without development of architecturally complex colonies. To

dry the plates, they were first allowed to solidify at room temperature for 1 hour,

afterwards, they were kept completely open in a laminar flow sterile bench for the

duration of the drying period. These drying conditions were followed for all assays that

examined changes in colony architecture.

Strain construction.

All *B. subtilis* strains generated in this work were obtained via natural competence

transformation (53) using genomic or plasmid DNA from donor strains as indicated in

Table 1. Briefly, overnight cultures of the receiver strains were diluted to a 1:50 ratio with

GCHE medium, these cultures were incubated at 37°C for 4 h with shaking at 225 r.p.m.

After this incubation period, 5–10 µg of genomic or plasmid DNA were mixed with 500

µL of competent cells and further incubated for 2h before plating on LB plates added

with selection antibiotics. Strain TB822 was obtained by using the Cre recombinase

expressed from plasmid pMH66 to eliminate the Erm^R cassette of TB813, and

subsequently curating pMH66 via thermal loss of the plasmid (54). Briefly, TB813 was

transformed with 10 µg of pMH66, selecting transformants via incubation at 37°C on LB

plates added with tetracycline. Candidates were then screened for their capacity to grow

at 37°C on LB plates added with macrolide antibiotics (erythromycin-lincomycin), those

that were not able to grow were further incubated on LB plates at 43°C for 18 h to induce

the loss of pMH66. Candidates that were then unable to grow at 37°C on LB plates

added with tetracycline were considered to have lost pMH66.

Successful construction of all used strains and plasmids was validated via PCR and

restriction pattern analysis using standard molecular biology techniques. All PCR

primers used in this study are listed in Table 2. Primer pairs were used to amplify the

indicated loci (see Table 2), in order to confirm the proper mutation of the corresponding

gene. To confirm the correct construction of strains TB869 and TB870, primer oGFPrev2

was used in combination with oRGM38 (for $sacA$::P$_{tapA}$-gfp) or oRGM40 (for $sacA$::P$_{epsA}$-

gfp).

Isolation of bacteria from soil samples.

Two independent Mexican soil samples were screened to isolate bacteria able to grow

on LB or tryptone soya broth media. The first sample was collected from Tepoztlán,

Morelos (18° 59′ 7″ N, 99° 5′ 59″ W), a humid and verdant region of central Mexico. The

second sample was collected from Tehuacán, Puebla (18° 25′ 47.71″ N, 97° 27′ 58.1″

W), a semidesertic dry region in east central Mexico. Both samples were collected with a

clean metal spatula 5 cm below surface level, and at 15 to 20 cm of the roots of local

trees.

1 g of each soil sample was suspended in 9 mL of a sterile 0.85% NaCl solution, and 50

µL of Tween 80 were added. The resulting suspensions were vortexed for 5 minutes at

maximum speed. The bigger soil particles were allowed to sediment by keeping the

suspensions still for 10 minutes. The supernatants were then diluted with sterile 0.85%

NaCl to 1:1000, 1:10 000 and 1:100 000 ratios. 50 and 100 µL of these dilutions were

spread on LB and tryptone soya broth agar plates. These plates were incubated at 30°C

for a maximum of 5 days. Bacterial colonies that grew during the incubation period were

further isolated by cross-streaking them on LB or tryptone soya broth plates and

incubating them at 30°C for 48 hours. Single isolated colonies obtained from this

secondary cultivation were used to prepare liquid cultures on 3 mL of LB media. These

cultures were incubated at 30°C with shaking for a maximum of 48 hours. Bacteria that

grew efficiently during this incubation period were used to prepare glycerol stock

solutions (20% v/v) and stored at -80°C for further use. In total, 242 soil strains were

obtained and subsequently tested.

Soil strains interaction screening.

Overnight liquid cultures of *B. subtilis* strain TB48 and the obtained soil strains were

adjusted to OD_{600} 0.2 using LB medium. These diluted cultures were then mixed in 1:1,

10:1, and 1:10 ratios, and 2 µL of the mixed and pure cultures were inoculated on 2×SG

plates. For neighbor colonies interaction assay, 2 µL of the pure cultures were

inoculated at a distance of 5 mm from each other. Plates were incubated at 30°C for 72

h. The obtained colonies were used for microscopy analysis without further treatment.

Identification of soil strains.

Genomic DNA of selected soil strains was extracted with the GeneMATRIX Bacterial

and Yeast Genomic DNA Purification Kit, according to the manufacturer's instructions

(EURx, Poland). This DNA was used to PCR amplify a fragment of the 16S rRNA gene

using primers 27F and 1492R (55). Amplicons were purified with the High Pure PCR

Product Purification Kit according to the manufacturer's instructions (Roche Diagnostics,

Switzerland) and sequenced with primers 27F and 1492R (GATC Biotech, Germany).

Sequencing results were then compared with sequences in the National Center for

Biotechnology Information (16S ribosomal RNA Bacteria and Archaea database) using

the BLASTn algorithm (56). Soil strains' identities were established using minimum

query coverage of 98% and minimum identity values of 99%.

Purification and treatment conditions of soil strain supernatant.

To obtain cell-free supernatants of selected soil strains, 10 mL cultures on 2×SG

medium of the corresponding strains were incubated at 30°C for 24 hours with shaking

at 100 r.p.m. These cultures were then centrifuged at 7000 r.c.f. for 15 min. The

supernatants were collected and filter-sterilized using a 0.22 µm pore-size filter (Carl

Roth, Germany).

Colony wrinkleality assay.

We developed the following assay to assess the effect that the supernatants of soil

strains, and compound solutions, may have upon the architectural complexity of *B.*

subtilis colony biofilms. Sterile 12 mm-diameter cotton discs were placed on 90 mm-

diameter 2×SG agar plates, in such a way that there is a maximum amount of available

space among the discs themselves and between the discs and the border of the plates.

3 cotton discs were used per plate. 50 µL of the tested supernatant or compound

solution were deposited on the center of the cotton discs, and the plates were dried for 3

min by keeping them completely open in a laminar flow sterile bench. This drying period

was done in addition to the regular 20 min drying previously described. 2 µL of an

overnight culture of the tested strains were then inoculated at 5 mm from the edge of the

cotton discs. The plates were incubated at 30°C for a total of 72 hours. Every 24 hours

the cotton discs were reimpregnated with 25 μL of the corresponding supernatant or

compound solution. After the incubation period, the plates were used for microscopy

analysis without further treatment.

Bioassay-guided fractionation.

A 50 mL culture of M5 isolate was grown for 24 h under standard conditions. Bacterial

cells were pelleted by centrifugation for 10 min at 6.000 g and the supernatant was

filtered through a 0.2 μm filter. 25 mL of the supernatant were freeze-dried and the

remaining foam was dissolved in 1 mL water. The solution was applied to a Sephadex

G20 column (3 cm × 40 cm) and eluted with water collecting 3 mL fractions (50

fractions). Fractions were further analyzed using the colony wrinkleality assay. The

fraction with the largest activity was analyzed using an LCMS (Shimadzu Deutschland,

Germany) equipped with a Hypercarb column (100 × 3 mm, 3 μM, Thermo Fisher

Scientific, flow rate = 0.6 mL min^{-1}, method: 0-10 min: 100% (v/v) water). The main

compound of this fraction was purified using a semi-preparative HPLC (Shimadzu

Deutschland, Germany) equipped with Hypercarb column (100 × 10 mm, 5 μM, Thermo

Fisher Scientific, flow rate = 5 mL min^{-1}, method = 0-20 min: 100% (v/v) water). The pure

compound was analyzed using ^{1}H-NMR spectroscopy, LCMS, and HR-ESIMS. Obtained

analytical data were in good agreement with a hypoxanthine standard from Sigma

Aldrich.

Comparison of hypoxanthine production.

Supernatant of different isolates were heated to 80°C for 15 min and filtered through a

0.2 μm syringe filter. The samples were analyzed using an LCMS (Shimadzu

Deutschland, Germany) equipped with a Hypercarb column (100 × 3 mm, 3 μM, Thermo

Fisher Scientific, flow rate = 0.6 mL min^{-1}, method: 0-10 min: 100% (v/v) water).

Stereomicroscopy and Image Analysis.

All bright-field and fluorescence images of colonies were obtained with an Axio Zoom

V16 stereomicroscope (Carl Zeiss, Germany) equipped with a Zeiss CL 9000 LED light

source, a PlanApo Z 0.5× objective, HE 38 eGFP filter set (excitation at 470/40 nm and

emission at 525/50 nm), HE 63 mRFP filter set (excitation at 572/25 nm and emission at

629/62 nm), and AxioCam MRm monochrome camera (Carl Zeiss, Germany).

Images were obtained with exposure times of 20 ms for bright-field, and 2500 ms for red

fluorescence or 3000 ms for green fluorescence when needed. For clarity purposes, the

images of colonies are presented here with adjusted contrast and the background has

been removed, so that the colony structures can be easily appreciated. The modified

pictures were not used for any fluorescence measurements.

To assess the expression levels of P$_{epsA}$-*gfp*, P$_{tapA}$-*gfp,* and P$_{hyperspank}$-*mKATE*

fluorescent reporter fusions in colonies we used ImageJ (National Institutes of Health,

USA). Briefly, the average fluorescence emission intensities were measured in the green

fluorescence channel for GFP and red fluorescence channel for mKATE by using a

region of interest (ROI) that surrounds the cotton discs in the pictures as a partial ring,

taking care to avoid the disc area itself. The ROI was drawn in such a way that it avoids

the region of the colony that first makes contact with the cotton discs. This ROI had a

width of 0.5 mm for measurements done at 40 and 50 hours, and a width of 1 mm for

measurements done at 65 hours. The ROI was positioned in each colony image using

the bright-field channel, and the average fluorescence intensity was then measured on

the corresponding green and red fluorescence images.

Nucleotide sequence accession numbers.

Sequences used in this study have been deposited in GenBank under accession

numbers KY705015 (*Lysinibacillus* sp. M2c), KY698015 (*Bacillus pumilus* P22a), and

KY703395 (*Acinetobacter variabilis* T7a). Further, the draft genome sequence of *L.*

fusiformis M5 is available in GenBank under accession number MECQ00000000, and

the strain was deposited in the Jena Microbial Resource Collection (ST-Number:

ST036146, see http://www.leibniz-hki.de/en/jena-microbial-resource-collection.html).

AUTHORS CONTRIBUTIONS

R.G.-M., Á.T.K. conceived the project; R.G.-M., S.K. performed the screen and the microbiology assays; R.G.-M. constructed bacterial strains; S.G., purified hypoxanthine; S.G., R.B., P.S. analyzed the analytical data; R.G.-M., P.S., Á.T.K., wrote and revised the manuscript.

ACKNOWLEDGEMENTS

We would like to thank Heike Heineke (HKI Jena) for NMR measurements and Andrea Perner (HKI Jena) for HR-MS measurements.

Ákos T. Kovács was funded by a Marie Curie Career Integration Grant (PheHetBacBiofilm), Deutsche Forschungsgemeinschaft (DFG) (KO4741/2-1 and KO4741/3-1), and DFG Graduate School Jena School of Microbial Communication (JSMC) (Start-up Grant). Ramses Gallegos-Monterrosa was funded by Consejo Nacional de Ciencia y Tecnología, German Academic Exchange Service. Sebastian Götze, Robert Barnett, and Pierre Stallforth was funded by Leibniz Association (Junior Research Group of Pierre Stallforth). Robert Barnett was funded by DFG Graduate School Jena School of Microbial Communication (JSMC) (PhD fellowship to Robert Barnett).

REFERENCES

1. **Hall-Stoodley L, Costerton JW, Stoodley P.** 2004. Bacterial biofilms: from the natural environment to infectious diseases. Nat Rev Microbiol **2**:95–108.

2. **Flemming H-C, Wingender J, Szewzyk U, Steinberg P, Rice SA, Kjelleberg S.** 2016. Biofilms: an emergent form of bacterial life. Nat Rev Microbiol **14**:563–575.

3. **West SA, Diggle SP, Buckling A, Gardner A, Griffin AS.** 2007. The social lives of microbes. Annu Rev Ecol Evol Syst **38**:53–77.

4. **Stewart PS, Franklin MJ.** 2008. Physiological heterogeneity in biofilms. Nat Rev Microbiol **6**:199–210.

5. **Miquel S, Lagrafeuille R, Souweine B, Forestier C.** 2016. Anti-biofilm activity as a health issue. Front Microbiol **7**:592. doi: 10.3389/fmicb.2016.00592

6. **Bjarnsholt T, Alhede M, Alhede M, Eickhardt-Sørensen SR, Moser C, Kühl M, Jensen PØ, Høiby N.** 2013. The *in vivo* biofilm. Trends Microbiol **21**:466–474.

7. **Valderrama WB, Cutter CN.** 2013. An ecological perspective of *Listeria monocytogenes* biofilms in food processing facilities. Crit Rev Food Sci Nutr **53**:801–817.

8. **Mangwani N, Kumari S, Das S.** 2016. Bacterial biofilms and quorum sensing: fidelity in bioremediation technology. Biotechnol Genet Eng Rev 8725. doi: 10.1080/02648725.2016.1196554

9. **Ramey BE, Koutsoudis M, Bodman SB Von, Fuqua C.** 2004. Biofilm formation in plant-microbe associations. Curr Opin Microbiol **7**:602–609.

10. **Parsek MR, Greenberg EP.** 2005. Sociomicrobiology: the connections between quorum sensing and biofilms. Trends Microbiol **13**:27–33.

11. **Mhatre E, Monterrosa RG, Kovács ÁT.** 2014. From environmental signals to

regulators: Modulation of biofilm development in Gram-positive bacteria. J Basic

Microbiol **54**:616–632.

12.   **Espeso DR, Carpio A, Einarsson B.** 2015. Differential growth of wrinkled

biofilms. Phys Rev E **91**:22710. doi: 10.1080/02648725.2016.1196554

13.   **Asally M, Kittisopikul M, Rué P, Du Y, Hu Z, Çağatay T, Robinson AB, Lu H,**

Garcia-Ojalvo J, Süel GM. 2012. Localized cell death focuses mechanical forces

during 3D patterning in a biofilm. Proc Natl Acad Sci U S A **109**:18891–18896.

14.   **Wilking JN, Zaburdaev V, De Volder M, Losick R, Brenner MP, Weitz DA.**

2013. Liquid transport facilitated by channels in *Bacillus subtilis* biofilms. Proc Natl

Acad Sci U S A **110**:848–852.

15.   **Vlamakis H, Chai Y, Beauregard P, Losick R, Kolter R.** 2013. Sticking together:

building a biofilm the *Bacillus subtilis way*. Nat Rev Microbiol **11**:157–168.

16.   **Lopez D, Vlamakis H, Kolter R.** 2009. Generation of multiple cell types in

Bacillus subtilis. FEMS Microbiol Rev **33**:152–163.

17.   **Mielich-Süss B, Lopez D.** 2015. Molecular mechanisms involved in *Bacillus*

subtilis biofilm formation. Environ Microbiol **17**:555–565.

18.   **Bais HP, Fall R, Vivanco JM.** 2004. Biocontrol of *Bacillus subtilis* against

infection of Arabidopsis Roots by *Pseudomonas syringae* is facilitated by biofilm

formation and surfactin production. Plant Physiol **134**:307–319.

19.   **Chen Y, Yan F, Chai Y, Liu H, Kolter R, Losick R, Guo JH.** 2013. Biocontrol of

tomato wilt disease by *Bacillus subtilis* isolates from natural environments

depends on conserved genes mediating biofilm formation. Environ Microbiol

15:848–864.

20.   **Stein T.** 2005. *Bacillus subtilis* antibiotics: structures, syntheses and specific

functions. Mol Microbiol **56**:845–857.

21.   **Todorova S, Kozhuharova L.** 2010. Characteristics and antimicrobial activity of

Bacillus subtilis strains isolated from soil. World J Microbiol Biotechnol **26**:1207–

1216.

22.   **Fernandes PAV, De Arruda IR, Dos Santos AFAB, De Araújo AA, Maior AMS,**

Ximenes EA. 2007. Antimicrobial activity of surfactants produced by *Bacillus*

subtilis R14 against multidrug-resistant bacteria. Brazilian J Microbiol **38**:704–709.

23.   **Shank EA, Klepac-Ceraj V, Collado-Torres L, Powers GE, Losick R, Kolter R.**

2011. Interspecies interactions that result in *Bacillus subtilis* forming biofilms are

mediated mainly by members of its own genus. Proc Natl Acad Sci U S A

108:E1236-1243.

24.   **Bleich R, Watrous JD, Dorrestein PC, Bowers AA, Shank EA.** 2015.

Thiopeptide antibiotics stimulate biofilm formation in *Bacillus subtilis*. Proc Natl

Acad Sci U S A **112**:3086–3091.

25.   **Powers MJ, Sanabria-Valentín E, Bowers AA, Shank EA.** 2015. Inhibition of cell

differentiation in *Bacillus subtilis* by *Pseudomonas protegens*. J Bacteriol

197:2129–2138.

26.   **Roesch L, Fulthorpe R, Riva A, Casella G, Hadwin A, Kent A, Daroub S,**

Camargo F, Farmerie W, Triplett E. 2007. Pyrosequencing enumerates and

contrasts soil microbial diversity. ISME J **1**:283–290.

27.   **Fierer N, Lennon JT.** 2011. The generation and maintenance of diversity in

microbial communities. Am J Bot **98**:439–448.

28.   **Kraemer SA, Kassen R.** 2015. Patterns of local adaptation in space and time

among soil bacteria. Am Nat **185**:317–331.

29. **Algburi A, Comito N, Kashtanov D, Dicks LMT, Chikindas ML.** 2017. Control of

biofilm formation: antibiotics and beyond. Appl Environ Microbiol **83**:e02508-

02516.

30. **Sultana ST, Babauta JT, Beyenal H.** 2015. Electrochemical biofilm control: a

review. Biofouling **31**:745–758.

31. **Winkelströter LK, Teixeira FB dos R, Silva EP, Alves VF, De Martinis ECP.**

2014. Unraveling microbial biofilms of importance for food microbiology. Microb

Ecol **68**:35–46.

32. **Spiers AJ.** 2014. A mechanistic explanation linking adaptive mutation, niche

change, and fitness advantage for the wrinkly spreader. Int J Evol Biol

2014:675432. doi: 10.1155/2014/675432

33. **Gallegos-Monterrosa R, Mhatre E, Kovács ÁT.** 2016. Specific *Bacillus subtilis*

168 variants form biofilms on nutrient-rich medium. Microbiology **162**:1922–1932.

34. **Jiang M, Shao W, Perego M, Hoch JA.** 2000. Multiple histidine kinases regulate

entry into stationary phase and sporulation in *Bacillus subtilis*. Mol Microbiol

38:535–542.

35. **Vlamakis H, Aguilar C, Losick R, Kolter R.** 2008. Control of cell fate by the

formation of an architecturally complex bacterial community. Genes Dev **22**:945–

953.

36. **López D, Kolter R.** 2010. Extracellular signals that define distinct and coexisting

cell fates in *Bacillus subtilis*. FEMS Microbiol Rev **34**:134–149.

37. **Saxild HE, Nygaard P.** 1987. Genetic and physiological characterization of

Bacillus subtilis mutants resistant to purine analogs. J Bacteriol **169**:2977–2983.

38. **Murray AW.** 1971. The biological significance of purine salvage. Annu Rev

Biochem **40**:811–826.

39. **Dandanell G, Hove-Jensen B, Willemoës M, Jensen KF.** 2008. Nucleotides,

nucleosides, and nucleobases. EcoSal Plus 3. doi: 10.1128/ecosalplus.3.6.2

40. **Torres RJ, Prior C, Garcia MG, Puig JG.** 2016. A review of the implication of

hypoxanthine excess in the physiopathology of Lesch–Nyhan disease.

Nucleosides Nucleotides Nucleic Acids **35**:507–516.

41. **Kim Y-J, Ryu H-M, Choi J-Y, Cho J-H, Kim C-D, Park S-H, Kim Y-L.** 2017.

Hypoxanthine causes endothelial dysfunction through oxidative stress-induced

apoptosis. Biochem Biophys Res Commun **482**:821–827.

42. **Day RO, Kamel B, Kannangara DRW, Williams KM, Graham GG.** 2016.

Xanthine oxidoreductase and its inhibitors: relevance for gout. Clin Sci **130**:2167–

2180.

43. **Hill-Perkins M, Jones MD, Karran P.** 1986. Site-specific mutagenesis in vivo by

single methylated or deaminated purine bases. Mutat Res **162**:153–163.

44. **Ayala-García VM, Valenzuela-García LI, Setlow P, Pedraza-Reyes M**. 2016.

Aag hypoxanthine-DNA glycosylase is synthesized in the forespore compartment

and involved in counteracting the genotoxic and mutagenic effects of

hypoxanthine and alkylated bases in DNA during *Bacillus subtilis* sporulation. J

Bacteriol **198**:3345–3354.

45. **He M, Li X, Liu H, Miller SJ, Wang G, Rensing C.** 2011. Characterization and

genomic analysis of a highly chromate resistant and reducing bacterial strain

Lysinibacillus fusiformis ZC1. J Hazard Mater **185**:682–688.

46. **Zhao L, Bao G, Geng B, Song J, Li Y.** 2015. Draft genome sequence of

Lysinibacillus fusiformis strain SW-B9, a novel strain for biotransformation of

isoeugenol to vanillin. Genome Announc 3:e00289-15.

doi:10.1128/genomeA.00289-15

47.   **Gupta S, Goyal R, Nirwan J, Cameotra SS, Tejoprakash N.** 2012.

Biosequestration, transformation, and volatilization of mercury by *Lysinibacillus*

fusiformis isolated from industrial effluent. J Microbiol Biotechnol **22**:684–689.

48.   **Gallegos-Monterrosa R, Maróti G, Bálint B, Kovács ÁT.** 2016. Draft genome

sequence of the soil isolate *Lysinibacillus fusiformis* M5, a potential hypoxanthine

producer. Genome Announc 4:e01272-16. doi: 10.1128/genomeA.01272-16

49.   **Wang S, Hu A.** 2016. Comparative study of spontaneous deamination of adenine

and cytosine in unbuffered aqueous solution at room temperature. Chem Phys

Lett **653**:207–211.

50.   **Stefanic P, Decorosi F, Viti C, Petito J, Cohan FM, Mandic-Mulec I.** 2012. The

quorum sensing diversity within and between ecotypes of *Bacillus subtilis*. Environ

Microbiol **14**:1378–1389.

51.   **Burmølle M, Ren D, Bjarnsholt T, Sørensen SJ.** 2014. Interactions in

multispecies biofilms: Do they actually matter? Trends Microbiol **22**:84–91.

52.   **Kobayashi K.** 2007. *Bacillus subtilis* pellicle formation proceeds through

genetically defined morphological changes. J Bacteriol **189**:4920–4931.

53.   **Kunst F, Rapoport G.** 1995. Salt stress is an environmental signal affecting

degradative enzyme synthesis in *Bacillus subtilis*. J Bacteriol **177**:2403–2407.

54.   **Yan X, Yu H-JJ, Hong Q, Li S-PP.** 2008. Cre/lox system and PCR-based genome

engineering in *Bacillus subtilis*. Appl Environ Microbiol **74**:5556–5562.

55.   **Devereux R, Wilkinson SS.** 2004. Amplification of ribosomal RNA sequences, p.

509–522. *In* Kowalchuk, GA, Bruijn, F de, Head, IM, Zijpp, AJ Van der, Elsas, JD

van (eds.), Molecular Microbial Ecology Manual, 2nd edition. Kluwer Academic

Publishers, Dordrecht, The Netherlands.

56. **Zhang Z, Schwartz S, Wagner L, Miller W.** 2000. A greedy algorithm for aligning

DNA sequences. J Comput Biol **7**:203–214.

57. **van Gestel J, Weissing FJ, Kuipers OP, Kovács ÁT.** 2014. Density of founder

cells affects spatial pattern formation and cooperation in *Bacillus subtilis* biofilms.

ISME J **8**:2069–2079.

58. **Perego M, Cole SP, Burbulys D, Trach K, Hoch JA.** 1989. Characterization of

the gene for a protein kinase which phosphorylates the sporulation-regulatory

proteins Spo0A and Spo0F of *Bacillus subtilis*. J Bacteriol **171**:6187–6196.

59. **Wang L, Grau R, Perego M, Hoch JA.** 1997. A novel histidine kinase inhibitor

regulating development in *Bacillus subtilis*. Genes Dev **11**:2569–2579.

60. **Grau RR, de Oña P, Kunert M, Leñini C, Gallegos-Monterrosa R, Mhatre E,**

Vileta D, Donato V, Hölscher T, Boland W, Kuipers OP, Kovács ÁT. 2015. A

duo of potassium-responsive histidine kinases govern the multicellular destiny of

Bacillus subtilis. MBio **6**:e00581-15. doi: 10.1128/mBio.00581-15

61. **López D, Fischbach MA, Chu F, Losick R, Kolter R.** 2009. Structurally diverse

natural products that cause potassium leakage trigger multicellularity in *Bacillus*

subtilis. Proc Natl Acad Sci U S A **106**:280–285.

62. **Koo B-M, Kritikos G, Farelli JD, Todor H, Tong K, Kimsey H, Wapinski I,**

Galardini M, Cabal A, Peters JM, Hachmann A-B, Rudner DZ, Allen KN,

Typas A, Gross CA. In press. Construction and analysis of two genome-scale

deletion libraries for *Bacillus subtilis*. Cell Syst. doi: 10.1016/j.cels.2016.12.013

63.   **Murray EJ, Strauch MA, Stanley-Wall NR.** 2009. SigmaX is involved in

696   controlling *Bacillus subtilis* biofilm architecture through the AbrB homologue Abh. J

697   Bacteriol **191**:6822–6832.

698   64.   **Casadaban MJ, Cohen SN.** 1980. Analysis of gene control signals by DNA fusion

699   and cloning in *Escherichia coli*. J Mol Biol **138**:179–207.

700   65.   **Kovács ÁT, van Hartskamp M, Kuipers OP, van Kranenburg R.** 2010. Genetic

701   tool development for a new host for biotechnology, the thermotolerant bacterium

702   *Bacillus coagulans*. Appl Environ Microbiol **76**:4085–4088.

TABLES AND FIGURES

Table 1. Strains and plasmids used in this study.

Strain	Characteristics	Reference
B. subtilis		
168	168 1A700 *trpC*. Jena Stock	(33)
TB48	168 *trpC2 amyE*::P*hyperspank*-*mKATE* (*cat*)	(57)
JH12638	JH642 *trpC2 phe*-1 *kinA*::Tn917(*cat*)	(58)
JH19980	JH642 *trpC2 phe*-1 *kinB*::*tet*	(59)
RGP0203-4	JH642 *kinC*::Sp^R	(60)
DL153	NCIB 3610 *kinD*::*tet*	(61)
BKE06370	168 *trpC2 pbuG*::Erm^R	(62)
BKE29990	168 *trpC2 pbuO*::Erm^R	(62)
NRS2243	NCIB 3610 *sacA*::P*epsA*-*gfp* (*neo*), *hag::cat*	(63)
NRS2394	NCIB 3610 *sacA*::P*tapA*-*gfp* (*neo*)	(63)
TB812	168 *trpC2 amyE*::P*hyperspank*-*mKATE* (*cat*) *pbuG*::Erm^R	This study, BKE06370→TB48
TB813	168 *trpC2 amyE*::P*hyperspank*-*mKATE* (*cat*) *pbuO*::Erm^R	This study, BKE29990→TB48
TB822	168 *trpC2 amyE*::P*hyperspank*-*mKATE* (*cat*) Δ*pbuO*	This study
TB823	168 *trpC2 amyE*::P*hyperspank*-*mKATE* (*cat*) Δ*pbuO, pbuG*::Erm^R	This study, BKE06370→TB822
TB833	168 *trpC2 amyE*::P*hyperspank*-*mKATE* (*cat*) *kinA*::Tn917(*cat*)	This study, JH12638→TB48
TB834	168 *trpC2 amyE*::P*hyperspank*-*mKATE* (*cat*) *kinB*::*tet*	This study, JH19980→TB48
TB835	168 *trpC2 amyE*::P*hyperspank*-*mKATE* (*cat*) *kinC*::Sp^R	This study, RGP0203-4→TB48
TB836	168 *trpC2 amyE*::P*hyperspank*-*mKATE* (*cat*) *kinD*::*tet*	This study, DL153→TB48
TB869	168 *trpC2 amyE*::P*hyperspank*-*mKATE* (*cat*) *sacA*::P*epsA*-*gfp* (*neo*)	This study, NRS2243→TB48
TB870	168 *trpC2 amyE*::P*hyperspank*-*mKATE* (*cat*) *sacA*::P*tapA*-*gfp* (*neo*)	This study, NRS2394→TB48
E. coli		
MC1061	Cloning host; K-12 F⁻ λ⁻ Δ(*ara-leu*)7697 [*araD*139]B/r Δ*(codB-lacI)*3 *galK16 galE15* e14⁻ *mcrA0 relA1 rpsL150*(Str^R) *spoT1 mcrB1 hsdR2*(r⁻ m⁺)	(64)

Soil Isolates		
Lysinibacillus sp. M2c		This study
Lysinibacillus fusiformis M5		This study
Bacillus pumilus P22a		This study
Acinetobacter variabilis T7a		This study
Plasmid	**Characteristics**	**Reference**
pMH66	pNZ124-based Cre-encoding plasmid, Tet^R Ts	(65)

Table 2. PCR primers used in this study.

Primer	Sequence (5´ → 3´)	Target locus
oGFPrev2	TTGTGCCCATTAACATCACC	*gfp*
oRGM110	GGAATCCGCGCCGTTACATC	*pbuG*
oRGM111	CAGCCCATATAGCAAAGACC	*pbuG*
oRGM116	GCGGTGCGGAATAAGTAAAG	*pbuO*
oRGM117	TACTGAGCGGCACTTGCTTG	*pbuO*
oRGM130	TATCCACGCCTACGCAGAGC	*kinA*
oRGM131	CTCAATGGACACGCTGAGAG	*kinA*
oRGM132	GAAGACCAGCAAGCAAATCG	*kinD*
oRGM133	GCGGCTGATCGCCTTTATGG	*kinD*
oRGM38	GAGAATTCGTGGTGCCAAAGACGAGAAG	*tapA* promotor
oRGM40	GAGAATTCCCAGCTGATTAATAGAATAG	*epsA* promotor
oTB55	CATGGGATCCTGGCGGAGAAGGATTTATG	*kinB*
oTB56	CACGGAATTCTGTCTCAAACGTGCTCATC	*kinB*
oTB61	CATGGGATCCATTACGCTAAGCCCTGAG	*kinC*
oTB62	CACGGAATTCTTGTGCCAGCAAATGATG	*kinC*
27F	AGAGTTTGATCMTGGCTCAG	16S rRNA
1492R	TACGGYTACCTTGTTACGACTT	16S rRNA

Figure 1. Single strain and mixed colonies of *B. subtilis* and selected soil strains. *B. subtilis* was differentiated from the soil strains using fluorescence emission (false-colored orange) from a reporter expressed by the P*hyperspank*-mKATE construct. Neighbor colonies were inoculated with 2 μL of cultures at 5 mm from each other. Scale bars represent 5 mm.

Figure 2. Effect of cell-free supernatants of soil strains on the development of *B. subtilis* 168 biofilm colonies. Bright-field images of colonies are shown. The scale bar represents 5 mm.

Figure 3. Characterization of hypoxanthine standard and cell-free supernatants of *B. subtilis* 168 (168), and soil isolates *L. fusiformis* M5 (M5) and *Lysinibacillus* sp. M2c (M2). a) HPLC chromatograms of cell-free supernatants compared to a standard solution of hypoxanthine. b) HPLC chromatogram of *L. fusiformis* M5 cell-free supernatant compared with a standard solution of hypoxanthine, and the *L. fusiformis* M5 cell-free supernatant spiked with a standard of hypoxanthine (M5+).

Figure 4. Effect of cell-free supernatant of *L. fusiformis* M5 on the development of biofilm colonies of *B. subtilis* 168 and knock-out mutants of kin-kinase genes. Bright-field images of colonies are shown. The scale bar represents 5 mm.

B. subtilis TB869 (P*epsA*-*gfp*, P*hyperspank*-*mKATE*)

B. subtilis TB870 (P*tapA*-*gfp*, P*hyperspank*-*mKATE*)

Figure 5. Comparison of fluorescence emission of *B. subtilis* strains carrying the constitutive P*hyperspank*-*mKATE*, and the P*epsA*-*gfp* (a-d) or P*tapA*-*gfp* (e-h), reporter fusions. Bright field (a and e), red fluorescence (b and f), and green fluorescence images (c and g) of representative *B. subtilis* biofilm colonies exposed to cell-free supernatants of *B. subtilis* 168, *L. fusiformis* M5 and 2×SG medium. Box plots on the ratio of green and red fluorescence emission of biofilm colonies of *B. subtilis* TB869 (d) and TB870 (h) exposed to cell-free supernatants of *B. subtilis* 168, *L. fusiformis* M5 and 2×SG medium at different time points. Scale bars represent 5 mm. Box plots (d and h) represent fluorescence ratios of at least 4 independent colonies, processed as described in material and methods. (one-way ANOVA: P<0.05, n=4-8 independent colonies).

Figure 6. Effect of hypoxanthine and cell-free supernatant of *L. fusiformis* M5 on the

development of biofilm colonies of *B. subtilis* 168 and knock-out mutants of *pbuO* and

pbuG permease genes. Bright-field images of colonies are shown. The scale bar

represents 5 mm.