1 **Cheating on cheaters dramatically affects social interactions in**
2 ***Pseudomonas aeruginosa***
3

4 Özhan Özkaya, Roberto Balbontín, Isabel Gordo, and Karina B. Xavier#

6 Instituto Gulbenkian de Ciência, Oeiras, Portugal

9 Running Head: Cheating on cheaters prevents drastic population collapse

12 #Address correspondence to Karina B. Xavier, kxavier@igc.gulbenkian.pt

Abstract

Bacterial cooperation can be disrupted by non-producers, which can profit from public goods without paying their production cost. A cheater can increase in frequency, exhausting the public good and causing a population collapse. Here we investigate how interactions among two cheaters for distinct social traits influence the short and long-term dynamics of polymorphic populations. Using as a model *Pseudomonas aeruginosa* and its extensively studied social traits, production of the siderophore pyoverdine and the quorum sensing regulated elastase, we analyzed the social dynamics of polymorphic populations under conditions where the two traits are required for optimal growth. We show that cheaters for either trait compete with both the wild type and each other, and that mutants for pyoverdine production can prevent a drastic population collapse caused by quorum sensing cheaters. A simple mathematical model suggests that the observed social dynamics are determined by the ratio of the costs of each social trait, such that the mutant which avoids paying the highest cost dominates the population. Finally, we demonstrate how quorum sensing regulation can avoid the full loss of cooperation.

Introduction

Bacteria are unicellular organisms, but can engage in diverse group behaviors, including biofilm formation, swarming motility, and production of extracellular proteases or iron-chelating siderophores [1–4]. The production of compounds that can benefit both producers and non-producers (public goods) can be considered as one of these cooperative behaviors. Cooperation is frequently under the threat of exploitation by cheaters: individuals that benefit from the cooperative action but contribute little or nothing to the production of the public goods. This situation, where both cooperators and cheaters can access a resource produced by the formers, is referred to as public goods dilemma [5,6]. If cheaters emerge, by mutation or migration, they can increase in frequency and cause loss of cooperation. As they rise to dominance, the public goods get exhausted and a population collapse, characterized by a strong decrease in the growth yield of the entire population, can occur; this population collapse is also referred in sociomicrobiology as 'the tragedy of the commons' [7–11].

Although often theorized [5,8], population collapse due to cheater expansion is hard to observe in natural populations even under conditions where cheaters spreading has been observed [12]. This raises the question of how invasion by cheaters is prevented and cooperative behaviors are maintained in microbial populations in nature. Mechanisms such as spatial structure and diffusion [13–22], pleiotropy [10,23–30], restricted migration [31], social and non-social adaptations [11,32–34], policing mechanisms [9], molecular properties of public goods [35], and metabolic strategies [36], have been proposed to play significant roles in maintaining cooperation by preventing cheater invasions and avoiding population collapse [2]. However, cheating behavior is observed *in vitro* [9–11,25], *in vivo* [37,38], and in natural populations, including clinically relevant environments such as the lungs of cystic fibrosis (CF) patients chronically infected with *Pseudomonas aeruginosa* [12,39–45].

Importantly, cheater invasion leading to the loss of the cooperative trait and the collapse of the population have been observed in laboratory studies focusing on a single trait [9–11]. However, in environments where more than

one social trait is required, the roles among mutants for these traits are likely to be more complex, since a cheater for one trait can be a cooperator for another, making 'cheater' and 'cooperator' relative terms [46,47]. We hypothesize that, in environments requiring bacteria to express multiple cooperative traits simultaneously, competition among mutants for these traits can influence their social interactions and, therefore, dictate the fate of the population. To test this hypothesis, we examine the consequences of ecological interactions among two social cheaters (mutants for two different traits) and the full cooperator (the wild type) in *P. aeruginosa* populations under conditions where the two cooperative traits are required for optimal growth.

Both *lasR* and *pvdS* mutants have been studied individually in a large number of sociomicrobiology studies [10,25,35,48–52], and are among the most common mutants recurrently isolated from the sputum samples of CF patients [12,39,41].

LasR is the master regulator of quorum sensing and, among many other genes, controls the production of extracellular elastase [53–55], which is essential for *P. aeruginosa* to use complex sources of amino acids, such as casein, as a carbon and nitrogen source [56–59]. Previous studies showed that *lasR* mutants grow poorly in media containing casein as the only carbon source, but increase in frequency when mixed with wild-type (WT) bacteria. Such increase leads to a population collapse where total cell numbers are drastically reduced due to depletion of producers of the public good [9–11].

PvdS is an alternative sigma factor, that among other genes, controls the transcription of genes responsible for pyoverdine biosynthesis [60,61]. In iron-limited environments, *P. aeruginosa* can secrete pyoverdine, which binds iron from the environment and is subsequently retrieved, providing iron to the cell [49]. Mutants in pyoverdine synthesis (e.g. *pvdS*) do not pay the cost of its production but are still able to retrieve the iron-bound pyoverdine produced by others, gaining a fitness advantage and increasing in frequency in mixed populations [35,62,63].

We analyzed the cheating of a *lasR* knock-out (KO) mutant over wild-type bacteria in an environment where production of elastase is required (medium with casein as the sole carbon source with iron supplementation). Then, we

modified the conditions (medium with casein as the sole carbon source with iron
depletion by human apo-transferrin) to cause pyoverdine production to be also
required and added a third social player (a *pvdS* KO mutant), to study the
interactions in these two different scenarios in short and long-term competition
experiments. We found that the fitness advantage of the *lasR* mutant
disappears when the *pvdS* mutant is in the culture and the two traits are
necessary. The long-term consequence of the interaction between these two
mutants is the prevention of the drastic population collapse, which occurs (*i*)
irrespectively of the presence of *pvdS* under conditions where only elastase is
required, or (*ii*) in the absence of the *pvdS* in conditions where the two traits are
required. The observed dynamics can be explained by a simple mathematical
model of multiple public good competition, which predicts the dominance of the
mutant that avoids expressing the trait with the highest cost, eventually causing
the corresponding population collapse associated with the loss of that trait.

Results

Cheating capacity of* lasR *mutant depends on abiotic and biotic factors.

We first investigated the growth yields of the WT and *lasR* and *pvdS*
mutants in monocultures under environmental conditions where either trait,
both, or none are required. In medium where casein is the sole carbon source,
iron-supplemented casein medium (Casein + Fe), and therefore elastase is
required, the *lasR* mutant has a lower growth yield than the WT and the *pvdS*
mutant (Figure 1A). In medium strongly depleted in iron, as it has been
repeatedly shown in the literature, the growth yields of all the three strains,
including the WT, are severely reduced [34,35,63–67]. However, the growth
yield of the *pvdS* mutant is significantly lower than those of the WT and the *lasR*
mutant in iron depleted media with casamino acids (CAA + Transferrin), where
only pyoverdine is necessary (Figure 1B). In a medium with casein as the sole
carbon source and low iron, namely, iron-depleted casein medium (Casein +
Transferrin), where both traits are required, both *lasR* and *pvdS* mutants have a
lower growth yield than the WT (Figure 1C). In this medium, the growth yield of
the *lasR* mutant is smaller than that of the *pvdS* mutant, indicating that the traits
have different benefit/cost ratios in this environment. Importantly, the WT and

the two mutants have similar growth yields in a medium where none of the traits are required, iron-supplemented CAA medium (CAA + Fe) (Figure 1D), in accordance with the expectation that the differences in growth yields of the mutants across media are due to the lack of expression of each social trait in the corresponding mutant. Moreover, the observation that the *lasR* mutant only has a growth yield significantly lower than the WT in the media with casein as the sole carbon source indicates that, even though LasR regulates many genes besides those responsible for elastase production [53–55], most do not significantly affect fitness under the conditions tested (For more details about the media used in this study and the growth yield differences of the strains, see Supplemental Information).

We then determined the relative fitness of each mutant in competition with the WT and each other in the different media (Figure 2). When co-cultured, at a ratio WT+*lasR* of 9:1, in conditions where only elastase production is required, the *lasR* mutant can cheat on the WT, since *lasR* increases in frequency with respect to it (Figure 2A-left), whereas such increase does not occur when elastase production is not required (Figure 2G). The introduction of the *pvdS* mutant (at ratio WT+*lasR*+*pvdS* of 8:1:1) does not significantly affect the cheating behavior of the *lasR* mutant, since *lasR* can also increase in frequency in the triple co-culture (Figure 2A-right). The incapability of the *pvdS* mutant to affect cheating by *lasR* is consistent with the fact that *pvdS* does not cheat under these conditions (Figure 2B). As expected, *pvdS* can cheat in medium where only pyoverdine is required, whereas *lasR* cannot (Figure 2, panels D and C, respectively). Next, we studied the interaction between these social players in conditions where the two traits are necessary. In these conditions, the *lasR* mutant also increases in frequency in co-culture with the WT (Figure 2E-left). Remarkably, the introduction of the *pvdS* mutant under these conditions causes the cessation of cheating by *lasR* (Figure 2E-right), which is consistent with the fact that *pvdS* can cheat on the WT in co-culture (Figure 2F-left) and both on the WT and the *lasR* mutant in triple co-culture (Figure 2F-right, Figure S1F). Importantly, in conditions where neither of the social traits are necessary, no cheating can be observed (Figure 2G-H), further ratifying that the effects observed are due to social interactions.

Notably, *lasR* mutants have been reported to produce lower amounts of

pyoverdine than the WT in iron-limited succinate minimal medium [68].

However, we found no significant difference in pyoverdine production between

WT and *lasR* across the different media used in this study (Figure S2). The

difference between our results and those of Stintzi and colleagues might be due

to differences in the media used in the two studies [64], or potential differences

in the strains used.

Altogether, these results demonstrate that the cheating capacities of the

two social mutants studied here are context-dependent, varying not only with

the environment, but also with the level of polymorphism in the population.

Invasion of* lasR *mutant leads to a drastic collapse of the population

We next asked what are the long-term consequences of the different

cheating capacities of *lasR* for the overall fitness of the population by

performing long-term propagations (Figure 3). We started co-cultures of

WT+*lasR* or WT+*lasR*+*pvdS* (at 9:1 and 8:1:1 initial ratios, respectively), either

in medium requiring only elastase production (Figure 3A and B), or in medium

where the two traits are needed (Figure 3C and D). Propagations were

performed by transferring an aliquot of each culture to fresh media every 48

hours. Before each passage, growth yields and frequencies of WT, *pvdS,* and

lasR cells were determined (Figure 3).

We observed that, in five out of six replicates of WT+*lasR* co-cultures in

the medium where only elastase is required, the *lasR* mutant quickly increased

in frequency throughout the first 8 days (4 passages), reaching up to 90% of the

population (red bars in Figure 3A). The total cell numbers of the populations

(black lines) rapidly decreased by day 12, and no recovery was observed in

subsequent passages (Figure 3A). We defined this drastic decrease in density

as the population collapse caused by the domination of the *lasR* mutant. One

replicate, out of six, did not follow this trend; in this case, no population collapse

was observed, and the total cell numbers remained high throughout the

experiment (Figure 3A). The fact that this only occurred in one of the six

replicates suggests that, in this particular replicate, the WT may have acquired

non-social beneficial mutation(s) that could prevent invasion of the *lasR* mutant,
as it was described in a recent study [11].

Next, we analyzed long-term competitions in triple co-cultures (WT, *pvdS*,
and *lasR*; with initial frequencies of 8:1:1, respectively) in the medium where
only elastase is required (Figure 3B). In this case, we observed an increase in
lasR frequency similar to that of seen in WT+*lasR* co-cultures (Figure 3A),
which was also accompanied by a drastic decrease in the overall population
size. At day 12 of the propagation, all 6 populations collapsed (Figure 3B). The
frequencies of the *pvdS* mutant varied between 4% and 15% throughout the
experiment, with no indication of any sustained increase (blue bars in Figure
3B). This result is consistent with the predictions from the relative fitness
measurements, which shows no cheating of *pvdS* in these conditions (Figure
2B).

Then we propagated WT+*lasR* co-cultures in the medium where the two
traits are necessary (Figure 3C). In these propagations, the *lasR* mutant also
increases in frequency throughout the first days, but at a slower pace than when
only elastase is required (compare panels A and C in Figure 3). The total cell
numbers remain high until days 10-12, but, as the *lasR* frequencies increase to
about 80%, the density of the population decreases, collapsing by day 18.
Hence, in all the three scenarios described above, the dominance of the *lasR*
mutant, which presumably resulted in the exhaustion of the public good
elastase, caused a drastic population collapse (Figure 3A–C).

**pvdS *prevents the drastic population collapse caused by the invasion of*
the lasR *mutant***

Our short-term competitions revealed that the cheating capacity of *lasR* is
influenced not only by abiotic, but also by biotic conditions, as the presence of
pvdS under conditions where both traits are required reduces the relative fitness
of the *lasR* mutant (Figure 2C). Therefore, we investigated if*,* in propagations in
the medium where both traits are needed, *pvdS* could prevent the drastic
population collapse caused by *lasR* invasion. Indeed, Figure 3D shows that
lasR does not increase in frequency, staying at approximately 3% throughout
the experiment. In contrast, *pvdS* rapidly expands to an average frequency of

96% at day 18. As the *lasR* mutant does not increase in frequency, cell
densities of the overall populations do not decrease and collapse of the
population is not observed.

Given that *pvdS* dominated the populations, a reduction in cell numbers
due to its invasion could be expected. Indeed, full fixation of the *pvdS* mutant
should result in a small population decrease close to the levels of the *pvdS*
mono-cultures (Figure 1C). However, at the end of the propagation experiments
(day 18), complete fixation of *pvdS* had not yet been reached, and an average
of 4% of WT and *lasR* cells were detected in the populations (Figure 3D). We
hypothesized that the presence of only 4% of pyoverdine producers in the
population could be enough to sustain the growth of the entire populations to
levels similar to the WT mono-cultures similarly to what has been reported
recently for cultures in chemostat [69]. The results shown in Figure S3A support
this hypothesis, since the growth yields of WT+*pvdS* mixed cultures at different
initial frequencies of *pvdS* significantly decrease only when the initial frequency
of *pvdS* is 98% or less, whereas mixtures with 3-4% of WT cells (or WT and
lasR cells) have growth yields similar to that of WT monocultures. These results
demonstrate that a small proportion of pyoverdine producer cells (WT and/or
lasR cells) are sufficient to produce enough pyoverdine to sustain the entire
population. This justifies why, in the propagation shown in Figure 3D, where at
day 18, *pvdS* reached an average frequency of 96%, no significant drop in cell
numbers was observed. Moreover, these results indicate that, if the
propagations were to continue, *pvdS* fixation and the subsequent decrease in
cell density could be expected. In fact, new WT+*pvdS* and WT+*pvdS*+*lasR*
propagations with much higher initial frequencies of *pvdS* (75-85%), allowed to
observe this population collapse (Figure S3B and C). The reason why *lasR* or
pvdS domination lead to a stronger or milder population collapse, respectively,
is related with the different characteristics of these two mutants, which have
different fitness in mono-culture in the medium requiring both traits (Figure 1C),
presumably as a consequence of the differences in cost and benefits of the
traits involved.

Remarkably, the presence of the *pvdS* mutant in the 3-way competition
under conditions where the two traits are required has a strong effect on the

outcome of the propagations in terms of growth yields, which is dramatically
different from those of the other three scenarios tested, since *pvdS* domination
prevents the drastic population collapse (Figure 3D) caused by the expansion of
lasR in the other three conditions (Figure 3A–C). This occurs because, in this
environment, although the *lasR* mutant is still being able to cheat on the WT
(Figure S1C), it is being cheated by *pvdS* (Figure S1D).

Importantly, long-term propagation experiments of WT+*lasR* and
WT+*lasR*+*pvdS* in medium where neither of the traits are required showed no
significant change in the population densities (Figure S4).

Alterations in carbon or iron source availability can prevent or induce the collapse

We reasoned that, if social interactions dominate over *de novo* adaptive
mutations in long-term dynamics, alterations of the abiotic factors in the triple
cultures should modify the social role of each mutant (by changing the costs
and benefits of the cooperative traits) and therefore affect the outcome for the
populations. Indeed, in the propagation of WT+*lasR* co-cultures in the medium
where only elastase is required, changing the carbon source from casein to
CAA during the course of the propagation (making elastase unnecessary)
eliminates the advantage of the *lasR* mutant, and this environmental change is
sufficient to prevent the population collapse (Figure 4A). Conversely, addition of
iron to the medium where both traits are required (thus making pyoverdine
unneeded) reverts the expansion of the *pvdS* mutant, favoring *lasR* cheating,
and ultimately causing the collapse of the populations at day 18 (Figure 4B). We
confirmed that changes in final frequencies observed in Figure 4B were not due
to the high initial frequencies of *pvdS,* because even though the selective
advantage of *pvdS* is frequency dependent, this mutant is capable of cheating
even at frequencies higher than 90% (Figure S5).

Overall, these results confirm the cheating role of the two mutants, and
also demonstrate the preponderance of social interactions over evolutionary
adaptation by *de novo* mutation in the propagation experiments shown in Figure
3.

A mathematical model of a 3-way public goods game explains the dynamics of the cheating mutants

To further investigate the general factors determining the dynamics of competitions among cooperators and cheaters we built a simple mathematical model (see Supplemental Information, Mathematical Model 1 and 2). The model assumes that the cost (c) of a cooperative trait is lower than the benefit (b) associated with this trait ($b>c>0$), and also that the benefit provided by the cooperative trait is equal for the entire population, as it would be expected in the case of an equally accessible public good in a well-mixed environment. Spatial structure, diffusion, or privatization, which would alter the benefit gained from the public good for cooperators and cheaters asymmetrically, were not considered in the model. The parameters used are described in Supplementary Table S1. As can be seen from the fitness definitions of the three players involved in our simple 3-way public model (Supplemental Information, Mathematical Model 1 (equations 1 to 3)), the cheaters always have a higher fitness than the cooperator due to the costs (c_1 or c_2) saved. Figure 5A shows the predicted mean fitness ($\bar{\omega}$) and final frequencies of the different strains in the population assuming different c_1/c_2 ratios. It can be easily seen that cooperators will always go extinct, and that the two cheaters can only co-exist when $c_1 = c_2$. Whenever $c_1 \neq c_2$, then the cheater that produces the trait with highest cost will lose. Therefore, the relation between c_1 and c_2 determines which cheater will dominate the population, independently of the benefits (b_1 and b_2) of these cooperative traits. On the other hand, the mean fitness, $\bar{\omega}$, is affected by the difference between b and c values of each trait.

We simulated the four scenarios corresponding to the conditions in Figure 3. As shown in Figure S6 in panels A and C, the cooperator for both traits and the cheater of the 1st cooperative trait compete, while the cheater of the 2nd cooperative trait is absent, whereas in panels B and D all three strains compete. In panels A and B, only the 1st cooperative trait is necessary, while in panels C and D both traits are required. The c_1/c_2 ratios defined in the model are estimated from the ratios of the relative fitnesses determined in the competitions shown in Figure 2E and F. The results of the model for the four scenarios resemble the experimental data, explaining changes in frequencies reasonably

well (Figure S6A-D). However, this simple model predicts complete fixation of the expanding mutant (Figure S6A-D), and cannot explain the lack of fixation of *lasR* (Figure 3A–C) or *pvdS* (Figure 3D) observed experimentally. As discussed above, as long as *de novo* mutations are not acquired, *pvdS* can reach fixation when co-cultured either with WT, or with WT and *lasR* under conditions where the two traits are needed (Figure S3B and C), which is in accordance to the model. However, that was not the case when *lasR* expansion was observed. We tested experimentally whether fixation of the *lasR* mutant could occur if the propagations were continued under conditions where *lasR* was expanding. Our results show that, when we initiate WT+*lasR* competitions at initial *lasR* frequencies similar to those at day 18 in Figure 3A, *lasR* still fails to reach fixation (Figure 6A). This dynamical behavior of *lasR* is not predicted under the assumptions of the model and suggests that other processes are taking place in the experiment.

Given that the *lasR* gene and elastase production are regulated by quorum sensing, we hypothesized that quorum sensing could be responsible for the lack of fixation of *lasR* mutant observed experimentally. Quorum sensing regulation should reduce both the cost and the benefit of elastase production when the cooperators are below the quorum sensing threshold, as cells will not produce elastase in that phase. We therefore modelled the effect of quorum sensing on fitness equations by assuming a Hill function where the cost and benefit of the 1st cooperative trait are sharply reduced when the frequency of the cheater for the 1st cooperative trait reaches a given threshold value (see Supplemental Information, Mathematical Model 2 for the model including quorum sensing). In this case, fixation of the mutant for the 1st cooperative trait can only happen if $c_1 < c_2$. When $c_1 \geq c_2$, both cheaters can co-exist in the population (Figure 5B). As shown in Figure 7, the simulations of the modified model including quorum sensing for the four experimental conditions predict accurately their frequency dynamics. Moreover, it also predicts that mutants for traits regulated by quorum sensing, like *lasR*, will not reach fixation.

To test experimentally if quorum sensing regulation could indeed be the mechanism responsible for preventing fixation of *lasR* in the WT+*lasR* competitions, we repeated the propagation experiment shown in Figure 6A

adding the quorum sensing autoinducer Acyl-homoserine lactone (AHL) N-3-oxododecanoyl-homoserine lactone ($3OC_{12}$-HSL) to the culture medium. Addition of AHL abolishes the quorum sensing-dependent regulation of elastase, locking elastase production constitutively in the ON state. Remarkably, the addition of AHL allows the *lasR* mutant to expand throughout the competitions, and eventually reach fixation (Figure 6B), as the model without quorum sensing predicted (Figure 5A). Thus, regulation of the production of a public good by quorum sensing prevents full domination of the quorum sensing cheater, maintaining cooperation in populations. However, if the expanding cheater is affected in the production of a public good not regulated via quorum sensing (e.g. *pvdS*), this mutant can dominate the entire population.

In summary, the results obtained with our 3-way public goods model including quorum sensing (Figure 7) show that the dynamics observed in our propagation experiments (Figure 3) can be explained by the relationship between the cost of the different cooperative traits involved and a quorum threshold that regulates both costs and benefits of one of these traits. Figures S6E-I represent predictions, according to our model, for other possible scenarios with different relationships between the costs, which can be tested experimentally in the future (see Supplemental Information).

Discussion

The classical experimental approach in sociomicrobiology has been to study one trait at a time. The simplicity of such an approach has allowed to substantially increase our understanding of the dynamics of cooperative interactions and revealed several mechanisms involved in the maintenance of cooperation [2,4,47]. In particular, the ability of *lasR* or *pvdS* mutants to behave as cheaters individually has been extensively documented [9,11,25–27,34,35,64,67,70–74], and these mutants are commonly isolated from bacterial populations colonizing CF lungs [41,45].

We established an experimental setup where WT cooperates in more than one trait: production of elastase via quorum sensing regulation and production of siderophore pyoverdine. Under conditions where the two traits are required,

the *lasR* mutant can act a cheater for elastase but a cooperator for pyoverdine, whereas the *pvdS* mutant can cooperate for elastase production and cheats for pyoverdine production. Our results showed that, in this environment, the 3-way competitions result in a dominance of *pvdS* over both the WT and the *lasR* mutant. Presumably, this occurs because the advantage of the *pvdS* mutant (caused by not producing pyoverdine) is higher than that of *lasR* (for not producing elastase, and the other quorum sensing regulated goods) under conditions where the two traits are necessary, causing *pvdS* to be more fit than *lasR* in this environment (Figure 1C). As a consequence, the *pvdS* mutant can cheat on the *lasR* mutant (and on the WT) (Figure 2, Figure S1), dramatically affecting the outcome of the long-term competitions (Figure 3D).

The expansion of *pvdS* under conditions where the two traits are required prevents the drastic population collapse caused by invasion of *lasR* mutants observed in absence of *pvdS* (compare panels D and C of Figure 3). Even though the domination of *pvdS* mutant can also lead to a drop in the density of the population caused by the exhaustion of the public good, (Figure S3B and C), the decrease in cell density due *pvdS* domination is much less drastic than that of observed upon domination of *lasR* mutant (Figure 3A – C).

Interestingly, both *lasR* and *pvdS* mutants are stronger cheaters in the medium where either of their affected trait is required than in the medium where the two traits are necessary (Figure 2, panels A and D versus E and F). In the case of *lasR*, this difference is coherent with the lower growth yields reached under conditions where the two traits are required (Figure 1, A versus C), which allow fewer cell divisions and therefore milder cheating. The difference in cheating of *pvdS* cannot be ascribed to higher growth yields when only pyoverdine is required (Figure 1B and C). However, pyoverdine production per cell is significantly higher in the medium where only pyoverdine is required (Figure S2G-H), and this could possibly explain the boosting in the cheating by *pvdS* in this medium. A stronger iron depletion in the CAA + Transferrin medium is coherent with both the low iron content of the CAA mixture used in our experiments [75] and the iron-chelating capacity of casein [76,77], which could allow carryover of casein-bound iron to the medium and thus result in higher iron availability in the media with casein.

Our simple model assuming that the difference in costs and benefits of the
cooperative traits involved proved to be sufficient to reasonably explain our
experimental results. Therefore, the model allowed us to infer the general
parameters governing social interactions beyond the particularities of the two
mutants used in this study. Specifically, the mathematical model suggests that,
in competitions among more than one social cheater under conditions where
more than one trait is required (a scenario likely to be closer to the conditions in
nature), the mutant for the trait with the highest cost is expected to dominate.
Moreover, the degree of the decrease in population density caused by loss of
cooperation due to exhaustion of the public good is determined by the benefit
minus the cost difference of the trait affected. In case of a trait with high benefit-
cost difference, a drastic collapse on the density of the population caused by
the cheater in that trait is expected. In contrast, if the mutant for the trait with a
low benefit-cost difference (as inferred for *pvdS*) dominates, a weak drop
occurs. These scenarios that lead to different degrees of decrease in population
densities could have very different consequences for the host in the context of
infections.

Importantly, our results provide support for a dynamic view of cooperation
and cheating that is dependent on both the genotypes present in the population
and the environmental conditions. We demonstrated how changes in the abiotic
environment can cause a social mutant to start or stop cheating or being
cheated. Additionally, as shown here for the *lasR* mutant*,* quorum sensing
regulation can also favor the maintenance of polymorphism, since such
regulation alters the values of the cost and benefits of the traits as a function of
the population density.

A better understanding of the interactions in polymorphic bacterial
populations in complex environments not only helps to gain insights into key
aspects of sociomicrobiology, but also can provide a theoretical framework for
the development of new therapeutic strategies against bacterial populations
where social mutants can invade [41,45]. In particular, our study provides
relevant information about the biotic and abiotic conditions that favor the
expansion of these mutants, which should be taken into account when

considering strategies aiming to manipulate populations where this type of social interactions is taking place.

The potential effects of the appearance of *pvdS lasR* double mutants in settings similar to ours should also be considered since double mutants have the potential to occur *in vivo* [41]. Although we found no evidence for emergence of *pvdS lasR* double mutants within the period of the experiments reported here, in the course of longer propagations *pvdS lasR* double mutants generated by *de novo* mutations were identified (data not shown). The effects of these double mutants on the interactions described here should be investigated in the future. However, based on our results, we can speculate that double mutants, as full cheaters, should cause an accelerated collapse of the population.

A non-social explanation for the advantage of the *pvdS* mutant in triple co-cultures under conditions where the two traits are required was also considered given that, at least in *Pseudomonas fluorescens,* certain mutants defective in pyoverdine production have been reported to be better adapted even in environments where iron concentration is not low, and thus can be considered non-social mutations [78]. However, the fact that our *pvdS* mutant has a lower fitness than the WT in the low iron media and does not show any advantage in conditions where pyoverdine production is not necessary (Figure 1B and C, Figure 2B and H) rules out non-social adaptation as the reason for its advantage.

Collectively, our findings underline the need for including polymorphism in social phenotypes and multiple environmental conditions in experimental studies and mathematical models pertaining to cooperation in microbial populations. This need is further supported by recent theoretical and experimental studies showing that interactions between genetically and functionally interlinked cooperative traits can significantly affect the course of their social evolution [26,79]. Our results demonstrate that using experimental conditions that include more than one social trait can reveal complex and dynamic social roles in bacterial populations as well as their dependence on the environment. Understanding the dynamics of polymorphic populations in these complex environments provides insights into social interaction processes,

expanding their relevance beyond sociomicrobiology, in addition to providing important knowledge for the development of novel therapeutic tools.

Materials and Methods

Bacterial strains. The strains used in this study were *Pseudomonas aeruginosa* WT strain PA01, PA01 *lasR* mutant harboring a gentamycin resistant gene inserted in *lasR* (*lasR*::*GmR*) [80], and PA01 *pvdS* mutant harboring a gentamycin resistance gene resistance gene replacing the *pvdS* coding sequence (Δ*pvdS*::*GmR*) [81]. For more detailed information, see Supplemental Information, supplementary methods section.

Media and culture conditions. The medium where only elastase is required (iron-supplemented casein medium) contains casein (Sigma, Ref: C8654) (1% w/v) as the sole carbon and nitrogen source salts (1.18 g $K_2HPO_4.3H_2O$ and 0.25 g $MgSO_4.7H_2O$ per liter of dH_2O) and 50 µM of $FeCl_3$. The medium where only pyoverdine production is required (iron-depleted CAA medium) contains the same salt solutions indicated above, low iron CAA (BD, Ref: 223050) (1% w/v) as the sole carbon source and 100 µg/ml of human apo-transferrin (Sigma, T2036) and 20 mM sodium bicarbonate to deplete available iron and induce pyoverdine production. The medium where both traits are needed (iron-depleted casein medium) is identical to the iron-supplied casein medium but instead of $FeCl_3$ supplementation, this medium contains 100 µg/ml of human apo-transferrin (Sigma, T2036) and 20 mM sodium bicarbonate to deplete available iron and induce pyoverdine production. The medium where none of the traits is necessary (iron-supplemented CAA medium) contains the same salt solutions as the other media, low iron CAA (1% w/v) as the sole carbon source and 50 µM of $FeCl_3$. All cultures were incubated in 15 ml falcons at $37^{o}C$ with aeration (240 rpm, New Brunswick E25/E25R Shaker) for the incubation times indicated. Cell densities were estimated by measuring absorbance (Abs) at 600 nm (OD_{600}) in a Thermo Spectronic Helios δ spectrophotometer.

Determination of genotypic frequencies. Estimation of the frequencies of each strain in the co-cultures was performed by scoring fluorescence and colony morphology of colonies obtained from plating serial PBS dilutions of the cultures. For each individual sample, three aliquots (of 50µl - 200µl, as

appropriated) were plated into LB agar plates, which were used as technical replicates. Then, CFU/ml were calculated by scoring different colony morphologies on each plate (with three technical replicate for each biological replicate). A stereoscope (Zeiss Stereo Lumar V12) with a CFP filter was used to distinguish pyoverdine producers, which are fluorescent, from the non-fluorescent *pvdS* mutants [72,82]. *lasR* mutant colonies have distinct colony morphology: smaller with smooth edges whereas elastase producers are larger with rugged edges [82]. To validate the phenotypic scoring all colonies used to determine the frequency from day 18 of the propagation experiments (Figure 3D) were tested by PCR with primers for the *lasR* and *pvdS* genes. The PCR data confirmed the phenotypic scoring with 100% accuracy.

Measurement of relative mutant fitness. Relative fitness was used to determine the cheating capacity of each mutant as commonly used [72,83,84]. For both mutants (*lasR* and *pvdS*), the relative fitness of each mutant (v) was calculated as the change in frequency of the mutant over a period of 48 hours relative to the rest of the strains in the mixture, *i. e.*, $v = fm_{final}.fr_{initial} / fm_{initial}.fr_{final}$ [72,83,84]. Where *fm* is the proportion of the mutant measured at the beginning of the competitions for $fm_{initial}$, or after 48 hours of competition for fm_{final}, and *fr* is the final proportions of the rest of the strains in the competitions at time = 0 ($fr_{initial}$) or after 48 hours (fr_{final}). As $fr = (1 - fm)$, the relative fitness was determined using the following formula $v = fm_{final} (1 - fm_{initial}) / fm_{initial} (1 - fm_{final})$.

Competition experiments. We propagated six replicates under four different conditions (Figure 3). Prior to start the competition experiments, all strains were inoculated, from frozen stocks, in medium containing 1% (w/v) casein and 1% (w/v) CAA in salts solution (same as in iron-supplied casein medium, described above) for 36 hours at 37^{o}C temperature with shaking (240 rpm). Cells were then washed with PBS four times, to remove any residual extracellular factor. Next after measuring cell densities (OD_{600}), cultures were normalized to $OD_{600} = 1.0$ and used to inoculate the various media as described in the text and figures. The different strains were diluted into fresh media, at different ratios as specified, to a starting initial $OD_{600} = 0.05$. For short term competitions the relative frequencies were determined by plating an aliquot of each culture at the beginning of the experiment (t = 0), and after 48 hours of incubation. For long-

term competitions, the relative frequencies were determined at t = 0, and thereafter every 48 hours before each passage. At the end of every 48 hours 1.5 µl of each culture was transferred to 1.5ml of fresh medium (bottle-neck of 1/1000).

Statistical analysis. Independent biological replicates were separately grown from the frozen stocks of each strain. Each figure (or figure panel) includes data from at least 6 biological replicates. The sample size (N), corresponds to the total numbers of independent biological replicates in each figure panel and is provided in the corresponding figure legends. The Mann-Whitney test which is a non-parametric test, was used because it does not account for normality and it is more suitable for the sample size used in each experiment ($5 < N < 20$). For multiple corrections, Kruskal-Wallis test with Dunn's correction was used. For all statistical analyses we used GraphPad Prism 6 software (http://www.graphpad.com/scientific-software/prism).

Acknowledgements

We thank Joana Amaro for technical assistance; João B. Xavier (Memorial Sloan Kettering), João Barroso-Batista, Rita Valente, and Jessica Thompson for suggestions and helpful comments on the manuscript, Jan Engelstädter (The University of Queensland, Australia) for his help while building the mathematical model, and Kevin Foster (University of Oxford, UK) for sending some of the strains used in this work. This work is supported by the Howard Hughes Medical Institute (International Early Career Scientist grant, HHMI 55007436). R.B. is supported by European Research Council (ERC-2010-StG_20091118) and by Fundação para a Ciência e Tecnologia (FCT) with a postdoctoral fellowship SFRH/BDP/109517/2015, Ö.Ö. is supported by Fundação Calouste Gulbenkian with a Doctoral Fellowship 01/BD/13, K.X. and I.G. are supported by FCT Investigator Program.

Author Contributions

Conceptualization, Ö.Ö., K.X., I.G., and R.B.; Methodology, Ö.Ö., K.X., I.G., and R.B.; Investigation, Ö.Ö.; Writing – Original Draft, Ö.Ö. and R.B.;

Writing – Review & Editing, Ö.Ö., K.X., I.G., and R.B.; Funding Acquisition, K.X.
and I.G.; Resources, K.X. and I.G.; Supervision, K.X. and I.G.

Declaration of Interest

The authors declare no competing interests.

References

1. Foster, K.R. (2010). Social behaviour in microorganisms. In Social
Behaviour: Genes, Ecology and Evolution, A. J. M. and J. K. Tamás
Székely, ed. (Cambridge: Cambridge University Press), pp. 331–356.

2. Bruger, E., and Waters, C. (2015). Sharing the sandbox: Evolutionary
mechanisms that maintain bacterial cooperation. F1000Research 4, 2–9.

3. Parsek, M.R., and Greenberg, E.P. (2005). Sociomicrobiology: The
connections between quorum sensing and biofilms. Trends Microbiol. 13,
27–33.

4. Xavier, J.B. (2016). Sociomicrobiology and pathogenic bacteria. Microbiol
Spectr. 4(3), 1–10.

5. Dionisio, F., and Gordo, I. (2006). The tragedy of the commons, the public
goods dilemma, and the meaning of rivalry and excludability in
evolutionary biology. Evol. Ecol. 8, 321–332.

6. Frank, S. A. (2010). A general model of the public goods dilemma. J.
Evol. Biol. 23, 1245–50.

7. Rankin, D.J., Bargum, K., and Kokko, H. (2007). The tragedy of the
commons in evolutionary biology. Trends Ecol. Evol. 22, 643–651.

8. MacLean, R.C. (2008). The tragedy of the commons in microbial
populations: insights from theoretical, comparative and experimental
studies. Heredity (Edinb). 100, 471–477.

9. Wang, M., Schaefer, A.L., Dandekar, A.A., and Greenberg, E.P. (2015).
Quorum sensing and policing of *Pseudomonas aeruginosa* social
cheaters. Proc. Natl. Acad. Sci. 112, 2187–2191.

608 10. Dandekar, A.A., Chugani, S., and Greenberg, E.P. (2012). Bacterial
609   quorum sensing and metabolic incentives to cooperate. Science 338,
610   264–266.

611 11. Asfahl, K.L., Walsh, J., Gilbert, K., and Schuster, M. (2015). Non-social
612   adaptation defers a tragedy of the commons in *Pseudomonas aeruginosa*
613   quorum sensing. ISME J. 9, 1734–1746.

614 12. Andersen, S.B., Marvig, R.L., Molin, S., Krogh Johansen, H., and Griffin,
615   A.S. (2015). Long-term social dynamics drive loss of function in
616   pathogenic bacteria. Proc. Natl. Acad. Sci. 112, 10756–10761.

617 13. Krakauer, D.C., and Pagel, M. (1995). Spatial structure and the evolution
618   of honest cost-free signalling. Proc. R. Soc. B Biol. Sci. 260, 365–372.

619 14. Foster, K.R., Shaulsky, G., Strassmann, J.E., Queller, D.C., and
620   Thompson, C.R.L. (2004). Pleiotropy as a mechanism to stabilize
621   cooperation. Nature 431, 693–6.

622 15. Kreft, J.U. (2004). Biofilms promote altruism. Microbiology 150, 2751–
623   2760.

624 16. Lion, S., and Baalen, M. Van (2008). Self-structuring in spatial
625   evolutionary ecology. Ecol. Lett. 11, 277–295.

626 17. Kümmerli, R., Griffin, A.S., West, S. A, Buckling, A., and Harrison, F.
627   (2009). Viscous medium promotes cooperation in the pathogenic
628   bacterium *Pseudomonas aeruginosa*. Proc. Biol. Sci. 276, 3531–3538.

629 18. Nadell, C.D., Foster, K.R., and Xavier, J.B. (2010). Emergence of spatial
630   structure in cell groups and the evolution of cooperation. PLoS Comput.
631   Biol. 6, e1000716.

632 19. West, S.A., Winzer, K., Gardner, A., and Diggle, S.P. (2012). Quorum
633   sensing and the confusion about diffusion. Trends Microbiol. 20, 586–594.

634 20. Dobay, A., Bagheri, H.C., Messina, A., Rankin, D.J., Kümmerli, R., and
635   Rankin, D.J. (2014). Interaction effects of cell diffusion, cell density and
636   public goods properties on the evolution of cooperation in digital

microbes. J. Evol. Biol. 27, 1869–1877.

21.   Drescher, K., Nadell, C.D., Stone, H.A., Wingreen, N.S., and Bassler, B.L.
(2014). Solutions to the public goods dilemma in bacterial biofilms. Curr.
Biol. 24, 50–55.

22.   Persat, A., Nadell, C.D., Kim, M.K., Ingremeau, F., Siryaporn, A.,
Drescher, K., Wingreen, N.S., Bassler, B.L., Gitai, Z., and Stone, H.A.
(2015). The mechanical world of bacteria. Cell 161, 988–997.

23.   Banin, E., Vasil, M.L., and Greenberg, E.P. (2005). Iron and
Pseudomonas aeruginosa biofilm formation. Proc. Natl. Acad. Sci. U. S.
A. 102, 11076–81.

24.   Harrison, F., and Buckling, A. (2009). Siderophore production and biofilm
formation as linked social traits. ISME J. 3, 632–634.

25.   Sandoz, K.M., Mitzimberg, S.M., and Schuster, M. (2007). Social cheating
in *Pseudomonas aeruginosa* quorum sensing. Proc. Natl. Acad. Sci. U. S.
A. 104, 15876–15881.

26.   Ross-Gillespie, A., Dumas, Z., and Kümmerli, R. (2015). Evolutionary
dynamics of interlinked public goods traits: An experimental study of
siderophore production in *Pseudomonas aeruginosa*. J. Evol. Biol. 28,
29–39.

27.   Wilder, C.N., Diggle, S.P., and Schuster, M. (2011). Cooperation and
cheating in *Pseudomonas aeruginosa*: the roles of the *las*, *rhl* and *pqs*
quorum-sensing systems. ISME J. 5, 1332–43.

28.   Friman, V.P., Diggle, S.P., and Buckling, A. (2013). Protist predation can
favour cooperation within bacterial species. Biol. Lett. 9, 20130548.

29.   Wagner, V.E., and Iglewski, B.H. (2008). *P. aeruginosa* biofilms in CF
infection. Clin. Rev. Allergy Immunol. 35, 124–34.

30.   Bachmann, H., Fischlechner, M., Rabbers, I., Barfa, N., Branco dos
Santos, F., Molenaar, D., and Teusink, B. (2013). Availability of public
goods shapes the evolution of competing metabolic strategies. Proc. Natl.

Acad. Sci. U. S. A. 110, 14302–7.

31. Kerr, B., Neuhauser, C., Bohannan, B.J.M., and Dean, A.M. (2006). Local
migration promotes competitive restraint in a host–pathogen "tragedy of
the commons." Nature 442, 75–78.

32. Waite, A.J., and Shou, W.Y. (2012). Adaptation to a new environment
allows cooperators to purge cheaters stochastically. Proc. Natl. Acad. Sci.
U. S. A. 109, 19079–19086.

33. Hammarlund, S.P., Connelly, B.D., Dickinson, K.J., and Kerr, B. (2016).
The evolution of cooperation by the Hankshaw effect. Evolution 70, 1376–
85.

34. Kümmerli, R., Santorelli, L.A., Granato, E.T., Dumas, Z., Dobay, A.,
Griffin, A.S., and West, S.A. (2015). Co-evolutionary dynamics between
public good producers and cheats in the bacterium *Pseudomonas*
aeruginosa. J. Evol. Biol. 28, 2264–74.

35. Kümmerli, R., and Brown, S.P. (2010). Molecular and regulatory
properties of a public good shape the evolution of cooperation. Proc. Natl.
Acad. Sci. U. S. A. 107, 18921–18926.

36. Xavier, J.B., Kim, W., and Foster, K.R. (2011). A molecular mechanism
that stabilizes cooperative secretions in *Pseudomonas aeruginosa*. Mol.
Microbiol. 79, 166–179.

37. Rumbaugh, K.P., Diggle, S.P., Watters, C.M., Ross-Gillespie, A., Griffin,
A.S., and West, S.A. (2009). Quorum sensing and the social evolution of
bacterial virulence. Curr. Biol. 19, 341–345.

38. Czechowska, K., McKeithen-Mead, S., Al Moussawi, K., and
Kazmierczak, B.I. (2014). Cheating by type 3 secretion system-negative
Pseudomonas aeruginosa during pulmonary infection. Proc. Natl. Acad.
Sci. U. S. A. 111, 7801–7806.

39. Cordero, O.X., and Polz, M.F. (2014). Explaining microbial genomic
diversity in light of evolutionary ecology. Nat Rev Microbiol. 12, 263–273.

40.   Katzianer, D.S., Wang, H., Carey, R.M., and Zhu, J. (2015). Quorum non-
sensing: social cheating and deception in *Vibrio cholerae*. Appl. Environ.
Microbiol. 81, 3856–3862.

41.   Smith, E.E., Buckley, D.G., Wu, Z., Saenphimmachak, C., Hoffman, L.R.,
D'Argenio, D.A., Miller, S.I., Ramsey, B.W., Speert, D.P., Moskowitz,
S.M., *et al.* (2006). Genetic adaptation by *Pseudomonas aeruginosa* to
the airways of cystic fibrosis patients. Proc. Natl. Acad. Sci. 103, 8487–
8492.

42.   Sommer, L.M., Molin, S., Johansen, H.K., and Marvig, R.L. (2015).
Convergent evolution and adaptation of *Pseudomonas aeruginosa* within
patients with cystic fibrosis. Nat. Genet. 47, 57–65.

43.   Nguyen, A.T., O'Neill, M.J., Watts, A.M., Robson, C.L., Lamont, I.L.,
Wilks, A., and Oglesby-Sherrouse, A.G. (2014). Adaptation of iron
homeostasis pathways by a *Pseudomonas aeruginosa* pyoverdine mutant
in the cystic fibrosis lung. J. Bacteriol. 196, 2265–2276.

44.   Folkesson, A., Jelsbak, L., Yang, L., Johansen, H.K., Ciofu, O., Hoiby, N.,
and Molin, S. (2012). Adaptation of *Pseudomonas aeruginosa* to the
cystic fibrosis airway: an evolutionary perspective. Nat Rev Microbiol. 10,
841–851.

45.   Winstanley, C., O'Brien, S., and Brockhurst, M.A. (2016). *Pseudomonas*
aeruginosa evolutionary adaptation and diversification in cystic fibrosis
chronic lung infections. Trends Microbiol. 24, 327–337.

46.   Brown, S.P., and Taylor, P.D. (2010). Joint evolution of multiple social
traits: a kin selection analysis. Proc. R. Soc. B-Biological Sci. 277, 415–
422.

47.   Özkaya, Ö., Xavier, K.B., Dionisio, F., and Balbontín, R. (2017).
Maintenance of microbial cooperation mediated by public goods in single-
and multiple-traits scenarios. J. Bacteriol. 199:e00297-17.

48.   De Vos, D., De Chial, M., Cochez, C., Jansen, S., Tümmler, B., Meyer,
J.M., and Cornelis, P. (2001). Study of pyoverdine type and production by

Pseudomonas aeruginosa isolated from cystic fibrosis patients: Prevalence of type II pyoverdine isolates and accumulation of pyoverdine-negative mutations. Arch. Microbiol. 175, 384–388.

49. Visca, P., Imperi, F., and Lamont, I.L. (2007). Pyoverdine siderophores: from biogenesis to biosignificance. Trends Microbiol. 15, 22–30.

50. Cox, C.D., and Adams, P. (1985). Siderophore activity of pyoverdin for *Pseudomonas aeruginosa*. Infect. Immun. 48, 130–138.

51. Lamont, I.L., Beare, P.A., Ochsner, U., Vasil, A.I., Vasil, M.L., and Kustu, S. (2002). Siderophore-mediated signaling regulates virulence factor production in *Pseudomonas aeruginosa*. Proc Natl Acad Sci U. S. A. 99, 7072–7077.

52. Griffin, A.S., West, S. A, and Buckling, A. (2004). Cooperation and competition in pathogenic bacteria. Nature 430, 1024–1027.

53. Whiteley, M., Lee, K.M., and Greenberg, E.P. (1999). Identification of genes controlled by quorum sensing in *Pseudomonas aeruginosa*. Proc. Natl. Acad. Sci. U. S. A. 96, 13904–13909.

54. Wagner, V.E., Bushnell, D., Passador, L., Brooks, A.I., and Iglewski, B.H. (2003). Microarray analysis of *Pseudomonas aeruginosa* quorum-sensing regulons: effects of growth phase and environment. J. Bacteriol. 185, 2080–2095.

55. Schuster, M., Lostroh, C.P., Ogi, T., and Greenberg, E.P. (2003). Identification, timing, and signal specificity of *Pseudomonas aeruginosa* quorum-controlled genes: a transcriptome analysis. J. Bacteriol. 185, 2066–79.

56. Morihara, K. (1964). Production of elastase and proteinase by *Pseudomonas aeruginosa*. J. Bacteriol. 88, 745–57.

57. Gambello, M.J., and Iglewski, B.H. (1991). Cloning and characterization of the *Pseudomonas aeruginosa lasR* gene, a transcriptional activator of elastase expression. J. Bacteriol. 173, 3000–9.

754   58.   Passador, L., Cook, J.M., Gambello, M.J., Rust, L., and Iglewski, B.H.
755   (1993). Expression of *Pseudomonas aeruginosa* virulence genes requires
756   cell-to-cell communication. Science 260, 1127–30.

757   59.   Schad, P.A., Bever, R.A., Nicas, T.I., Leduc, F., Hanne, L.F., and
758   Iglewskil, B.H. (1987). Cloning and characterization of elastase genes
759   from *Pseudomonas aeruginosa*. J. Bacteriol. 169, 2691–26.

760   60.   Cunliffe, H.E., Merriman, T.R., and Lamont, I.L. (1995). Cloning and
761   characterization of *pvdS*, a gene required for pyoverdine synthesis in
762   *Pseudomonas aeruginosa*: PvdS is probably an alternative sigma factor.
763   J. Bacteriol. 177, 2744–50.

764   61.   Miyazaki, H., Kato, H., Nakazawa, T., and Tsuda, M. (1995). A positive
765   regulatory gene, *pvdS*, for expression of pyoverdin biosynthetic genes in
766   *Pseudomonas aeruginosa* PAO. Mol. Gen. Genet. 248, 17–24.

767   62.   Dumas, Z., and Kümmerli, R. (2012). Cost of cooperation rules selection
768   for cheats in bacterial metapopulations. J. Evol. Biol. 25, 473–484.

769   63.   Dumas, Z., Ross-Gillespie, A., and Kümmerli, R. (2013). Switching
770   between apparently redundant iron-uptake mechanisms benefits bacteria
771   in changeable environments. Proc Biol Sci 280, 20131055.

772   64.   Kümmerli, R., Jiricny, N., Clarke, L.S., West, S.A., and Griffin, A.S.
773   (2009). Phenotypic plasticity of a cooperative behaviour in bacteria. J.
774   Evol. Biol. 22, 589–98.

775   65.   Ghysels, B., Dieu, B.T.M., Beatson, S.A., Pirnay, J.-P., Ochsner, U.A.,
776   Vasil, M.L., and Cornelis, P. (2004). FpvB, an alternative type I
777   ferripyoverdine receptor of *Pseudomonas aeruginosa*. Microbiology 150,
778   1671–1680.

779   66.   Popat, R., Harrison, F., da Silva, A.C., Easton, S.A.S., McNally, L.,
780   Williams, P., and Diggle, S.P. (2017). Environmental modification via a
781   quorum sensing molecule influences the social landscape of siderophore
782   production. Proc. R. Soc. London B Biol. Sci. 284, 20170200.

67. Butaitė, E., Baumgartner, M., Wyder, S., and Kümmerli, R. (2017).
Siderophore cheating and cheating resistance shape competition for iron
in soil and freshwater *Pseudomonas* communities. Nature Comm. 8, 414.

68. Stintzi, A., Evans, K., Meyer, J.M., and Poole, K. (1998). Quorum-sensing
and siderophore biosynthesis in *Pseudomonas aeruginosa*: *lasR/lasI*
mutants exhibit reduced pyoverdine biosynthesis. FEMS Microbiol. Lett.
166, 341–345.

69. Sexton, D.J., and Schuster, M. (2017). Nutrient limitation determines the
fitness of cheaters in bacterial siderophore cooperation. Nature Comm. 8,
230.

70. Cabeen, M.T. (2014). Stationary phase-specific virulence factor
overproduction by a *lasR* mutant of *Pseudomonas aeruginosa*. PLoS One
9, e88743.

71. Ghoul, M., West, S.A., McCorkell, F.A., Lee, Z.-B.B., Bruce, J.B., and
Griffin, A.S. (2016). Pyoverdin cheats fail to invade bacterial populations
in stationary phase. J. Evol. Biol. 29, 1728–1736.

72. Jiricny, N., Diggle, S.P., West, S.A., Evans, B.A., Ballantyne, G., Ross-
Gillespie, A., and Griffin, A.S. (2010). Fitness correlates with the extent of
cheating in a bacterium. J. Evol. Biol. 23, 738–747.

73. Buckling, A., Harrison, F., Vos, M., Brockhurst, M. A, Gardner, A., West,
S.A., and Griffin, A.S. (2007). Siderophore-mediated cooperation and
virulence in *Pseudomonas aeruginosa*. FEMS Microbiol. Ecol. 62, 135–
41.

74. Harrison, F., McNally, A., da Silva, A.C., Heeb, S., and Diggle, S.P.
(2017). Optimised chronic infection models demonstrate that siderophore
"cheating" in *Pseudomonas aeruginosa* is context specific. ISME J. 11,
2492–2509.

75. Kümmerli, R., and Ross-Gillespie, A. (2013). Explaining the sociobiology
of pyoverdin producing *Pseudomonas*: A comment on Zhang and Rainey
(2013). Evolution 11., 3337–3343.

76. Demott, B.J., and Dincer, B. (1976). Binding Added Iron to Various Milk
Proteins. J. Dairy Sci. 59, 1557–1559.

77. Smialowska, A., Matia-Merino, L., and Carr, A.J. (2017). Oxidative
stability of iron fortified goat and cow milk and their peptide isolates. Food
Chem. 237, 1021–1024.

78. Zhang, X., and Rainey, P.B. (2013). Exploring the sociobiology of
pyoverdin-producing Pseudomonas. Evolution 67, 3161–74.

79. Mellbye, B., and Schuster, M. (2014). Physiological framework for the
regulation of quorum sensing-dependent public goods in *Pseudomonas*
aeruginosa. J. Bacteriol. 196, 1155–1164.

80. Popat, R., Crusz, S.A., Messina, M., Williams, P., West, S.A., and Diggle,
S.P. (2012). Quorum-sensing and cheating in bacterial biofilms. Proc.
Biol. Sci. 279, 4765–4771.

81. Ochsner, U.A., Johnson, Z., Lamont, I.L., Cunliffe, H.E., and Vasil, M.L.
(1996). Exotoxin A production in *Pseudomonas aeruginosa* requires the
iron-regulated *pvdS* gene encoding an alternative sigma factor. Mol.
Microbiol. 21, 1019–1028.

82. Ghoul, M., West, S.A., Diggle, S.P., and Griffin, A.S. (2014). An
experimental test of whether cheating is context dependent. J. Evol. Biol.
27, 551–556.

83. Ross-Gillespie, A., Gardner, A., West, S. A, and Griffin, A.S. (2007).
Frequency dependence and cooperation: theory and a test with bacteria.
Am. Nat. 170, 331–42.

84. Ross-Gillespie, A., Gardner, A., Buckling, A., West, S. A, and Griffin, A.S.
(2009). Density dependence and cooperation: theory and a test with
bacteria. Evolution 63, 2315–25.

Figure 1. *P. aeruginosa lasR* **and** *pvdS* **mutants have lower growth yields than WT in media where elastase and/or pyoverdine are required, respectively.** Growth yields (CFU/ml) of WT (purple triangles), *pvdS* (blue squares), and *lasR* (red circles) strains of *P. aeruginosa* monocultures after 48 hours of incubation in **(A)** iron-supplemented casein medium (Casein + Fe), **(B)** iron-depleted casamino acids medium (CAA + Transferrin), **(C)** iron-depleted casein medium (Casein + Transferrin) and **(D)** iron-supplemented casamino acids medium (CAA + Fe). Each data point represents an individual biological replicate (N≥6) and the horizontal bars indicate the means of each group. For comparisons, Kruskal-Wallis test with Dunn's correction was used, ns=not significant P>0.05, * P≤0.05, ** P≤0.01, *** P≤0.001, **** P≤0.0001.

Figure 2. Relative fitness of *lasR* or *pvdS* in various media in double or triple co-cultures. Relative fitness (v) of *lasR* (red circles in **A, C, E,** and **G**) or *pvdS* (blue squares in **B, D, F,** and **H**) were calculated as the change in frequency of each mutant relative to the rest of the strains in each culture after 48 hours of incubation in **(A** and **B)** iron-supplemented casein medium (Casein + Fe), **(C** and **D)** iron-depleted casamino acids medium (CAA + Transferrin), **(E** and **F)** iron-depleted casein medium (Casein + Transferrin), and **(G** and **H)** iron-supplemented casamino acids medium (CAA + Fe). Relative fitness of *lasR* (**A, C, E,** and **G**) was calculated in co-cultures with WT, or with WT and *pvdS.* Relative fitness of *pvdS* (**B, D, F,** and **H**) was calculated in co-cultures with WT, or with WT and *lasR.* Initial ratios of the strains in each co-culture are 9:1 for WT+*lasR* and WT+*pvdS*, and 8:1:1 for WT+*pvdS*+*lasR*. Mann-Whitney two-tailed test was used to compare the relative fitness values of each mutant in double and triple co-cultures (significance symbols are located at the middle-top of each plot above the brackets). Dotted lines indicate v=1. Relative fitness values above the dotted lines (v>1) indicate that the strain is cheating and below the dotted lines (v<1) indicate that the strain is being cheated. One-sample t-test was used to determine whether each dataset is significantly different than 1 (significance symbols are located above the dotted line when v>1 and below the dotted line when v≤1). Each data point indicates an individual biological replicate (N≥6) and horizontal lines indicate the means of each group. ns=not significant P>0.05, * P≤0.05, ** P≤0.01, *** P≤0.001, **** P≤0.0001.

Figure 3. Effects of abiotic and biotic factors on growth yields and strain composition of the population in long-term propagations. Left Y-axes represent CFU/ml values prior to subculture; black symbols correspond to the CFU/ml values of each of the 6 biological replicates tested for each condition. Right Y-axes show the frequencies of WT (purple), *lasR* (red), and *pvdS* (blue) at each time point; data are shown as bars and represent the means of 6 biological replicates, error bars indicate SD. X-axes show the days of propagations. **(A)** WT and *lasR* co-cultures mixed at an initial frequency of 9:1 in iron-supplemented casein media. **(B)** WT, *lasR,* and *pvdS* triple co-cultures mixed at initial an initial frequency of 8:1:1 in iron-supplemented casein media. **(C)** and **(D)** same as in (A) and (B) but in iron-depleted casein media.

Figure 4. Effects of alterations of abiotic conditions in long-term propagations. (A) After the 6[th] day of the competitions of WT+*lasR* co-cultures in iron-supplemented casein medium (Casein + Fe) (Figure 3A), aliquots were transferred into iron-supplemented CAA medium (CAA + Fe) to relieve the requirement for digesting casein by elastase production (N=6, data from the first 6 days are from Figure 3A). **(B)** After the 6[th] day of the competitions of WT+*lasR*+*pvdS* triple co-cultures in iron-depleted casein medium (Casein + Transferrin) (Figure 3D), aliquots were transferred into iron-supplemented casein medium (Casein + Fe) to relieve the requirement for pyoverdine production (N=6, data from the first 6 days are from Figure 3D). Legends as in Figure 3.

Figure 5. Mathematical model for the final frequencies of the three strains in relation to the ratio of c_1/c_2. In Left-Y axes, the mean fitness, $\bar{\omega}$, is shown in black. In Right-Y axes, frequencies of cheater of the 1st cooperative trait (red), cheater of the 2nd cooperative trait (blue), and cooperator of both cooperative traits (purple) are shown in relation to the ratio of c_1/c_2 (X-axes) either without **(A) - mathematical model 1** or with the influence of quorum sensing (QS) regulation on the 1st cooperative trait **(B) - mathematical model 2**. The values given to the parameters of the simulations are: $p_{coop}(0)$=0.8, $p_{ch1}(0)$=0.1, $p_{ch2}(0)$=0.1, 0.001≤c_1<0.199, b_1=1.5, c_2=0.1, b_2=0.25, ω_0=0.1, time (as arbitrary units of cumulative numbers of cell divisions)=1800. In (B) the 1st cooperative trait is regulated by QS with n=30 and th=0.8.

Figure 6. Frequencies of *lasR* in propagations of WT+*lasR* co-cultures in iron-supplemented casein media in the absence or presence of exogenously added quorum sensing signal AHL. Initial frequencies of *lasR* of 80-90% were used, these frequencies were similar to those of the 18[th] day in Figure 3A. Cultures were propagated throughout 6 days by passing the fresh media each 48 hours. **(A)** Frequency changes of *lasR* in WT+*lasR* co-cultures (red). **(B)** is the same as (A) but with 5μM AHL ($3OC_{12}$-HSL) added to the media. Red lines indicate linear regressions. Dotted lines represent 100% domination of *lasR*.

Figure 7. Results of the mathematical model simulating the four scenarios in Figure 3. Model includes quorum sensing regulation of the 1st cooperative trait (b_1 and c_1 are negatively regulated via a Hill equation as a function of the frequency of the mutant of this trait, p_{ch1}). Left Y-axes show $\bar{\omega}$, the mean fitness of the entire population which is a function of b and c values; these values correspond to the biomass gain due to benefiting from the cooperative action (b), and the energy spent to the cooperative action instead of biomass increase (c). Right Y-axes show the frequencies of p_{coop} (e.g. WT, purple), p_{ch1} (e.g. *lasR*, red) and p_{ch2} (e.g. *pvdS,* blue). X-axes show the cumulative numbers of cell divisions as arbitrary units. In panels **(A)** and **(B)**, only the 1st cooperative trait is necessary ($b_1 > c_1 > 0$, whereas $b_2 = c_2 = 0$), while in panels **(C)** and **(D)** both traits are required ($c_2 > c_1 > 0$ and $b_1 > b_2 > 0$). In panels **(A)** and **(C)**, the cooperator for both traits (WT) and the cheater of the 1st cooperative trait compete ($p_{coop}(0) = 0.9$ and $p_{ch1}(0) = 0.1$), while the cheater of the 2nd cooperative trait is absent ($p_{ch2}(0)=0$), whereas in panels **(B)** and **(D)** all three strains compete ($p_{coop}(0) = 0.8$ and $p_{ch1}(0) = p_{ch2}(0) = 0.1$). The values that are given to the parameters of the simulations are: **(A)** $p_{coop}(0)= 0.9$, $p_{ch1}(0)=0.1$, $p_{ch2}(0)=0$, $c_1=0.01$, $b_1=3.4$, $c_2=0$, $b_2=0$, $\omega_0=0.2$, $th=0.8$, $n=30$; **(B)** $p_{coop}(0)= 0.8$, $p_{ch1}(0)=0.1$, $p_{ch2}(0)=0.1$, $c_1=0.01$, $b_1=3.4$, $c_2=0$, $b_2=0$, $\omega_0=0.2$, $th=0.8$, $n=30$; **(C)** $p_{coop}(0)= 0.9$, $p_{ch1}(0)=0.1$, $p_{ch2}(0)=0$, $c_1=0.01$, $b_1=1.5$, $c_2=0.025$, $b_2=0.25$, $\omega_0=0.1$, $th=0.8$, $n=30$; **(D)** $p_{coop}(0)= 0.8$, $p_{ch1}(0)=0.1$, $p_{ch2}(0)=0.1$, $c_1=0.01$, $b_1=1.5$, $c_2=0.025$, $b_2=0.25$, $\omega_0=0.1$, $th=0.8$, $n=30$. Note that the values of parameters used in these simulations are chosen to reflect approximately the relation between the values observed in Figure 1, Figure 2. (For more detailed description please see Supplemental Information, Mathematical Model 2).