Characterization of nitrogen deposition by means of atmospheric biomonitors

E. A. Díaz-Álvarez[1,2] and E. de la Barrera[2]

[1]Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México, Ciudad

Universitaria, México, Ciudad de México 04510, México.

[2]Instituto de Investigaciones en Ecosistemas y Sustentabilidad, Universidad Nacional

Autónoma de México, Morelia, Michoacán 58190, México.

Corresponding author: Erick de la Barrera, tel.: + 52 (443) 3223810, e-mail:

delabarrera@unam.mx

Abstract

An increase of nitrogen deposition resulting from human activities is not only a major threat for global biodiversity, but also for human health, especially in highly populated regions. It is thus important and in some instances legally mandated to monitor reactive nitrogen species in the atmosphere. The utilization of widely distributed biological species suitable for biomonitoring may be a good alternative. We assessed the suitability of an ensemble of atmospheric biomonitors of nitrogen deposition by means of an extensive sampling of a lichen, two mosses, and a bromeliad throughout the Valley of Mexico, whose population reaches 30 million, and subsequent measurements of nitrogen metabolism parameters. In all cases we found significant responses of nitrogen content, C:N ratio and the δ^{15}N to season and site. In turn, the δ^{15}N for the mosses responded linearly to the wet deposition. Also, the nitrogen content ($R^2 = 0.7$), the C:N ratio ($R^2 = 0.6$), and δ^{15}N ($R^2 = 0.5$) for the bromeliad had a linear response to NOx. However, the bromeliad was not found in sites with NOx concentrations exceeding 80 ppb, apparently of as a consequence of exceeding nitrogen. These biomonitors can be utilized in tandem to determine the status of atmospheric nitrogenous pollution in regions without monitoring networks for avoiding health problems for ecosystems and humans.

Keywords: Biogeochemistry; epiphytic plants; global change; megacities; stable isotopes; urban ecology

Introduction

Nitrogen deposition is one the most predominant forms of atmospheric pollution[1]. This phenomenon results from the release of nitrogenous compounds to the atmosphere, both in cities, and in the country, including oxidized (NOx) and reduced (NHx) species, which are highly reactive. Nitrogen deposition causes acidification and eutrophication of aquatic and terrestrial ecosystems, contributes to the proliferation of invasive species leading to changes in ecosystem structure; it is also an important factor for climate change[1-6]. Nitrogenous pollution is also an issue for human health worldwide. Indeed, in traffic-jammed motorways, high accumulations of NOx at ground level are harmful by direct inhalation[7]. NOx emissions also contribute to the formation of secondary compounds including tropospheric ozone and particulate matter which are responsible in part for 9 million annual deaths worldwide[8]. For the case of Mexico City, a megalopolis with more than 20 million habitants, at least 9600 deaths can be attributed annually to atmospheric pollution, resulting from respiratory and heart diseases, and pollutant build up in the nervous system and the brain [9,10]. In turn, ammonia emissions from agricultural sources quickly react with other atmospheric pollutants to form airbone particulate matter that can be readily inhaled[7]. Yet nitrogen is mostly neglected in public policy despite its importance in the prevalent discourse of global environmental change.

While monitoring nitrogenous pollution is an important issue, the deployment of air quality monitoring networks can be cost prohibitive in regions with developing economies. To fill this gap, an affordable alternative is the use of biomonitors[11]. For instance, throughfall deposition has been determined by the nitrogen content of lichen thalli[12]. Mosses are also widely used as biomonitors that allow covering vast areas, given that their

nitrogen content reflects the rates of deposition and their isotopic composition can reflect

the possible sources of pollution[13,14]. A group of special promise for biomonitoring in the

neo-tropical region are atmospheric bromeliads, whose succulent tissues allow for yearlong

physiologically activity, taking up pollution regardless of the seasonal environmental

conditions. This is the case for the genus *Tillandsia* (Bromeliaceae), amply distributed in

the Americas, which has been utilized for monitoring NOx pollution given that its nutrition

depends exclusively from atmospheric sources[15,16]. However, physiological limitations of

these atmospheric organisms can confuse the observed deposition, therefore it has been

proposed that an assemble of various biomonitors for tracking this type of atmospheric

pollution improves biomonitoring[11].

To assess whether the combined use of various biomonitors can provide reliable

information on nitrogen deposition we: 1) determined the spatial distribution of nitrogen

content and the isotopic composition of the lichen *Anaptychia* sp., the mosses *Grimmia* sp.,

and *Fabronia* sp., and the bromeliad *Tillandsia recurvata* throughout the Valley of Mexico;

and 2) evaluated the suitability as biomonitors of these organisms by comparing their

nitrogen with an existing automated atmospheric monitoring network.

Results

Spatial distribution of nitrogen deposition

The total wet deposition in Mexico City increased from east to west, it ranged from 23 to

45 kg ha^{-1} year^{-1} during 2013, and between 24 and 50 kg ha^{-1} year^{-1} during 2014 (Figure

2a). We found the highest rates of wet deposition in northwestern of Mexico City. The rest

of the valley lacks monitoring stations. The only existing record is from a natural protected

area at the north of the Valley, whose wet deposition is always below 5 kg ha^{-1} year$^{-1, 17}$.

For NOx concentrations, the northern part of Mexico City is the zone with the highest

concentrations reaching a cumulative of 78.8 ppb during the dry season and 40.6 ppb

during the rainy season (Figure 2b). The lowest concentrations of NOx were found in the

south and the east part of Mexico City which reached 21.8 ppb and 9.2 ppb during dry and

rainy season respectively (Figure 2b).

Nitrogen relations of biomonitors

Anaptychia sp.

The nitrogen content of *Anaptychia sp*. ranged from 1.4% (dry mass) in a natural

protected area at the north of the Valley to 5.0% in an urban site in Mexico City during the

dry season (Figure 3a). We observed the same pattern during the rainy season, in which the

nitrogen content ranged from 1.3 to 4.6%. The interaction between site and season was

significant for all parameters measured (Table 1). A notable finding is that the highest

nitrogen deposition of 50 kg N ha^{-1} year^{-1} did not produce the highest nitrogen content in

this lichen, which was observed in a site whose nitrogen deposition reached 31.5 kg N ha^{-1}

year^{-1}.

The C:N ratio ranged from 7.5 to 39.1, increasing from the urban to the rural sites

during the dry season; it ranged from 9.4 to 34.1 during the rainy season. Similarly, the

δ^{15}N values ranged between –9.4 and 5.2‰, being more positive in northeastern and

downtown Mexico City and in southeastern Pachuca, contrasting with negative values from

rural areas (Figure 3b). A linear regression showed no direct relationship between nitrogen

deposition and the parameters measured for this lichen (R^2=0.14 for nitrogen content,

R^2=0.07 for the C:N ratio and R^2=0.16 for the δ^{15}N).

Grimmia sp.

We found significant differences between the nitrogen content of mosses from a

natural protected area at the north of the valley that reached 1.3% where the deposition was

5.0 Kg N ha^{-1} year^{-1}, with those of a site at the north of Mexico City of 3.8% where the

deposition reached 40 Kg N ha^{-1} year^{-1} (Figure 3c). We observed no differences in the

nitrogen between for the dry and rainy season (Table 1). A linear regression showed that

the atmospheric concentration of NOx had no effect on the nitrogen content ($R^2 < 0.01$).

The C:N ratio was higher in rural areas and decreased significantly in the urban ones, it

ranged between 9.5 and 33.6. On the contrary, NOx concentration had no effect on the C:N

($R^2 < 0.01$).

We found the most negative $\delta^{15}N$ values of –7.0‰ in a rural area and positive ones

of 6.6‰ in urban sites where wet deposition was lower than 35 Kg N ha^{-1} year^{-1},

exceeding this point $\delta^{15}N$ became negative reaching –5.3‰, as occurs with mosses growing

at 40 Kg N ha^{-1} year^{-1} (Figure 3d; Figure 4). We also observed positive values in the city

of Pachuca at the north of the Valley. Season and NOx concentration showed no effect on

the $\delta^{15}N$ of this moss (Table 1).

***Fabronia* sp.**

The nitrogen content ranged between 1.7 and 3.4% (dry weight) across the Valley.

We found no differences between the seasons, but significant differences between the rural

areas and some sites in Mexico City (Table 1; Figure 3e). Wet deposition had no effect on

the nitrogen content for the mosses from Mexico City. For example, it only increased by

0.3% when nitrogen deposition when from 24.1 to 48.3 Kg N ha^{-1} year^{-1}. The site and the

season were statistical significant (Table 1). The NOx concentration had no effect on the

nitrogen content of this moss. In addition, wet deposition, nor the NOx concentration

resulted in alterations on C:N ratio, which ranged from 9.0 and 29.5.

The δ^{15}N ranged from –6.6 to 7.8‰. We found the most negative value in a semi-

rural site at the north of the Mexico City (Figure 3f), while a δ^{15}N of –6.0 ± 0.2‰ was

observed at the south, where the deposition reached 37.6 Kg N ha^{-1} year^{-1}. The most

positive value was found in the eastern border of Mexico City, where the nitrogen

deposition was 33.7 and 27.1 Kg N ha^{-1} year^{-1} during 2013 and 2014 respectively. The

urban environment in the city of Pachuca also had a considerable effect on the isotopic

composition of this moss, as positive values were the rule. Significant differences were

found between rural areas and the urban ones. In contrast, the season showed no effect on

the isotopic composition of this moss (Table 1). We found a weak relationship between the

rate of nitrogen deposition and the δ^{15}N values (R^2 = 0.2; Figure 5).

Tillandsia recurvata

The nitrogen content for *Tillandsia recurvata* ranged from 0.8 to 3.6% during the

dry season and from 1.0 to 2.2% (dry weight) during wet season (Figure 3g). The highest

content was observed in the northern part of Mexico City and was significantly different

from the lowest value found in a semirural site in the central part of the Valley. At the same

time, bromeliads collected in the city of Pachuca had a similar nitrogen content as those

from semirural areas of the Valley (Figure 3g). However, this variation was not due to wet

deposition. Instead, it responded positively to the NOx concentration (Figure 6a). For

example, the nitrogen content reached 3.6% in an urban site where the NOx reached 57.4

ppb, contrasting with the lowest nitrogen content that reached 0.8% in a rural site where

NOx presumably was lower to 5 ppb, based on the lowest recorded concentration in

Mexico City. Wet deposition had no effect on the C:N ratio of this bromeliad, but it was

strongly affected by the concentration of NOx during both seasons (Figure 6b). For

example, we found the lowest C:N ratio of 15.9 where the NOx reached 50.3 ppb, and the

highest C:N of 40 where NOx concentrations presumably were lower than 5 ppb. The $\delta^{15}N$

values for *T. recurvata* were negative in rural areas and became positive in the cities,

varying from □5.0 to 4.4‰ during the dry season and □7.7 to 5.1‰ during the wet season

(Figure 3h). The $\delta^{15}N$ values responded positively to the NOx concentration (R^2=0.49;

Figure 6c), while the wet deposition had no effect on $\delta^{15}N$ values (R^2=0.01). the data from

two sites (Figure 6c open circles) were excluded from the analysis because topographic and

exposure to pollutants were non-representative, clearly skewed the isotopic signature of

negative values. In particular, one of the sites is a zoological park with large mammals that

is crossed by a stream that was visible polluted. In turn, the second site were from a small

ravine with dense vegetation, where the diffusion of NOx was probably reduced and where

biogenic emissions were negative.

174   **Discussion**

The distribution of nitrogenous pollution and deposition in Mexico City generally

matched that of the sources of emission. For instance, the highest values were measured in

the northern part of the city, where the majority of the manufacturing industries are located

along numerous important and busy motorways[18]. Indeed, motor vehicles, followed by

industrial emissions, are the most important source of NOx in México City, whose

concentration is directly measured by the city government from the air[18,19]. In addition,

nitrogenous emissions from the agricultural zone between Mexico City and Pachuca

substantially contribute to the observed pattern, given predominant north-to-south winds[18].

In this case, NHx originated from both agricultural and industrial activities are measured

from wet deposition.

The nitrogen content of lichens has been utilized as an effective indicator of

nitrogen deposition. For example, three species of lichens can record nitrogen inputs of wet

deposition below 10 Kg N ha^{-1} year^{-1}, point at which they saturate [12,20]. We observed a

threshold of nitrogen tolerance for lichens from the Valley, when the deposition reached

31.5 kg N ha^{-1} year^{-1} above which they could not take up additional nitrogen. We observed

a similar pattern for the $\delta^{15}N$, while the negative values from rural areas were the result of

low rates of deposition, in urban areas high rates of deposition resulted in positive isotopic

values, however, above the threshold of saturation the isotopic values turned negative, as

has been already observed in other species of lichen [21]. Their isotopic composition

responded more to wet than dry deposition as was observed for mosses. Additionally,

different studies show a point of saturation when studying throughfall deposition [20, 21].

The nitrogen content of moss tissues is affected by the rate of deposition and

responds to the distance to urban centers [14,22-24]. We also observed an important influence

of urban centers for the nitrogen content of mosses from the Valley. Their $\delta^{15}N$ values are

affected by the rates of deposition and the prevalent pollutant (e.g. wet NH_4^+ or NO_3^-; dry

NHx or NOx,), being more commonly negative in mosses from rural areas [25-28]. The

observed negative $\delta^{15}N$ for mosses from rural areas resulted from uptake of NH_4^+ derived

from fertilizers and livestock emissions which have characteristic negative $\delta^{15}N$ values

[27,29,30]. Mosses from urban areas of the Valley such as Pachuca, as well as some areas of

Mexico City where the wet deposition was below 35 Kg N ha^{-1} year^{-1} had positive $\delta^{15}N$,

suggesting that they more likely take up NO_3^- derived from NO_x of fossil fuel burning and

industrial activities which typically display positive $\delta^{15}N$ [29]. Another possible mechanism

influencing the isotopic composition of these mosses is the gas-particle conversion process,

in which isotopic enrichment occurs for the aerosol nitrogen resulted, producing positive

$\delta^{15}N$ [31].

Mosses take up NH_4^+ preferentially over NO_3^- because less energy is needed in its

assimilation [32]. Additionally, high deposition rates can cause the inhibition of nitrate

reductase, reducing NO_3^- assimilation [27,33,34]. This occurs to some degree when the

deposition reaches 10 Kg N ha^{-1} $year^{-1}$, but when it exceeds 30 Kg N ha^{-1} $year^{-1}$ the nitrate

reductase become completely inhibited not only precluding for the NO_3^- be uptake leading

to its loss by leaching [27,35,36]. Nitrate reductase inhibition also has an important effect on the

isotopic composition of mosses because it favors the take up of NH_4^+ that typically has

negative $\delta^{15}N$ values [30]. In the Valley of Mexico, mosses $\delta^{15}N$ became negative when the

rates of wet deposition exceeded 35 Kg N ha^{-1} $year^{-1}$, which suggests that nitrate reductase

was indeed inhibited, allowing NH_4^+ uptake, a pollutant that represented 35% of the total

wet deposition in Mexico City during 2014 [18,36,37]. The atmospheric concentration of NOx

had no effect on the nitrogen content nor the isotopic composition of either moss

considered here because when NH_3 is the prevalent nitrogenous pollutant in the

atmosphere, the nitrogen content of the mosses can increase more than from wet deposition.

However, in Mexico City the prevalent nitrogenous gas pollutant is NOx, suggesting that

nitrogen content and the $\delta^{15}N$ of the mosses responded more to wet deposition than gaseous

pollutants [27,38].

Despite the high rates of wet deposition recorded in Mexico City which exceeded 50

Kg N ha^{-1} $year^{-1}$ in some areas, neither the nitrogen content nor the C:N ratio of *Tillandsia*

recurvata were directly affected. This occurs because raindrops cannot be absorbed by the

non-absorptive roots of these plants [39]. Instead, the nitrogen was taken up as NOx by the

leaves of this bromeliad because it can absorb particles and gasses from the air thanks to

stomatal gas exchange and the trichomes present in the surface [39]. This is evident from the

close relationship found here between the nitrogen content and the NOx concentration in

the Valley. This is similar to what occurs for other atmospheric bromeliads, whose nitrogen

content is higher in the vicinity of highways than further away [40-42]. The seasonal

differences found for the nitrogen content appeared to respond to the phenology of this

bromeliad which grows after the rainy season. Indeed, the NOx were dragged from the

atmosphere to the ground surface during rainy season reducing its biological availability

[18,43].

Both the nitrogen content and the isotopic composition of *T. recurvata* were

determined by the predominant anthropogenic activity of the sites where the plant was

collected. Bromeliads take up NOx with positive $\delta^{15}N$ from industrialized and densely

populated areas, contrasting with the negative emissions originated from biogenic sources,

the soil and livestock waste [44,45]. The positive $\delta^{15}N$ found for *T. recurvata* suggests no

nitrate reductase inhibition, but it is likely that NOx concentrations higher than 80 ppb may

result in some inhibition of this enzyme. This could be one of the reasons why *T. recurvata*

was not found in sites with concentrations of NOx higher to 80 ppb.

The distinct responses observed for each biomonitor evaluated here are a result of

the ecophysiological traits of each functional type. Indeed, the lack of a cuticle and a

vascular system for mosses, allows a rapid/ready assimilation of incoming pollution,

mainly from wet deposition. However, their weak response to NOx levels could be an

effect of plant phenology, as the mosses remain dormant during the dry season, so the

uptake of the gas is not physiologically possible. For the case of the tillandsias, the fact that

they are adapted to a water and nutrient-limiting environment was the cause for its

suitability for monitoring NOx, as all their nutrition is atmospheric and their water is

intercepted from the atmosphere [46]. In addition, their potential uptake of wet deposition is

inherently limited by their small area to volume ratio that in turn reduces water loss [47].

Also, their rates for taking up water are low, so exposure to more benign environments does

not lead to increased water uptake as we have found experimentally [48]. Of course, caution

must be taken when utilizing a biomonitoring method utilizing spontaneously occurring

plants, as their nitrogen parameters can also respond to various factors other than

nitrogenous pollution. For instance, if dewfall is substantial during the dry season, mosses

can remain physiologically active, and thus be able to assimilate NOx. Also, water

availability and extreme temperatures can have an effect on the N relations of plants. For

instance, high daytime temperatures and very low relative humidity during the dry season

or a very cloudy sky during the rainy season can both lead to stomatal closure during the

night for the CAM bromeliads. In either case, uptake of NOx will not occur. Other factors

to consider are pathogens and herbivores [49]. For these reasons, specific biomonitors need to

be developed for each region of interest, taking in consideration the particular

environmental conditions and the ecophysiological traits of potential biomonitors [11]. For

the case of the Mexico valley, and by possible extension to the Trans-Mexican Volcanic

Belt, *T. recurvata* can be utilized for monitoring NOx along with a moss that records wet

deposition. This approach can become a useful tool for determining the status of

nitrogenous pollution in regions where air quality monitoring networks are not available.

Especially in mid-sized cities and surrounding areas where the saturation thresholds of

these organisms has not been reached. Finally, the utilization in tandem of these organisms

can inform an early alert for avoiding health problems for both ecosystems and humans.

Methods

Study area

   The Valley of Mexico is located in central Mexico spanning 7500 km^2, with an average elevation of 2240 m [50]. Annual precipitation ranges from 600 mm at the center of the valley to 1300 mm in the surrounding mountains. The predominant winds blow from the northeast and northwest [51]. Mexico City sits at the southern edge of the Valley with a population of 20 million. At the northern edge, Pachuca, the capital of the state of Hidalgo, has a population of 3 million. Additional, settlements interspersed in the valley with industrial or agricultural activities comprise the rest of the 30 million inhabitants of this basin [10].

Nitrogen environment

The Mexico City environmental authority has deployed an air quality network of 16 monitoring stations for wet deposition (Fig. 1). This network collects data during the rainy season from May to November. The nitrogen collected consists of dissolved NO_3^- and NH_4^+, so that the total nitrogen is the sum of both forms of deposition ($NO_3^- + NH_4^+$). This is the dissolved inorganic nitrogen (DIN), biologically available [27]. For this study, the rates of wet deposition measured by this monitoring network during 2014 were utilized for the analyses described below. This air quality network also has 27 monitoring stations for measuring the atmospheric NOx concentration ($NO + NO_2$) year-round, 24 hours a day. We utilized the mean seasonal concentration in ppb during dry season (November 2013 to April 2014) and during the rainy season (from May 2014 to October 2014). Data were obtained from the website of the environmental authority of the Mexico City (http://www.aire.cdmx.gob.mx/default.php).

Biomonitoring

We selected four species of three types of atmospheric organisms, the lichen *Anaptychia*

sp., the mosses *Grimmia* sp. and *Fabronia* sp., and the bromeliad *Tillandsia recurvata* (L).

These organisms have been utilized as monitors of nitrogen deposition and are abundant in

the Valley of Mexico [14,15,21]. We collected tissue samples from 5 individuals of each

species growing on different substrates from 36 sites throughout the valley, including urban

parks, agricultural sites, and natural protected areas, for the dry season on (8-20 May) and,

the wet season 3-15 November of 2014. The distribution of the sites within the basin was

determined by the occurrence of the biomonitoring species and by the complex nature of

the landscape, which precluded the collection of samples from a regular grid [52].

The tissue samples were dried at 60°C in a gravity convection oven until reaching

constant weight. The dried tissues were ground to a fine powder in a ball mill (Retsch

MM300; Retsch, Vienna, Austria), wrapped into tin capsules (Costech Analytical, Inc.

Valencia, California, USA), and weighed with a microbalance (0.01 mg, Sartorius,

Göttingen, Germany). For each sample, both the nitrogen content and their isotopic

signature were determined at the Stable Isotope Facility, University of Wyoming (Laramie,

Wyoming, USA), with a Carlo Erba EA 1110 elemental analyzer (Costech Analytical Inc.,

Valencia, CA, USA) attached to a continuous flow isotope ratio mass spectrometer

(Finnigan Delta Plus XP, Thermo Electron Corp, Waltham, MA). Nitrogen isotope ratios,

reported in parts per thousand, were calculated relative to atmospheric air standards.

The analytical precision for the $\delta^{15}N$ was 0.3 ± 0.07‰ (SD). The natural abundances of ^{15}N

were calculated as:

$$\delta^{15}N\ (‰\ _{versus\ air}) = (R_{sample}/R_{standard}\ \square\ 1) \times 1000$$

where, R is the ratio of $^{15}N/^{14}N$ for nitrogen isotope abundance for a given sample [53,54].

Statistical analyses

We utilized linear regressions to determine the relationship between total wet

nitrogen deposition and the nitrogen content (% dry weight), the C:N ratio as well as the

$\delta^{15}N$ of the organisms considered. The same was done for the concentration of atmospheric

NOx. Data accomplished normality in both cases [30]. We performed two way ANOVAs

(factors were site and season) followed by a post hoc Holm–Sidak tests ($p \leq 0.05$) to

determine differences in the nitrogen content (% dry weight), the C:N ratio, as well as the

$\delta^{15}N$ values of the tissue samples. All statistical analyses were conducted with Sigmaplot

12 (Systat Software Inc. USA).

Geostatistical analyses

We utilized the ordinary Kriging method (a geostatistical gridding tool for

irregularly spaced data [55]) to determine the rates of wet deposition, and the concentration of

NOx in the sites where no monitoring station was available for the area covered by the

monitoring network. Likewise, we modelled the geographical distribution of the nitrogen

content, C:N ratio, and the $\delta^{15}N$ values for the biomonitors, as well as the data from the

monitoring network with ArcGIS 10 (Esri, Redlands, California, USA).

Data availability

347 The datasets generated and/or analyzed during the current study are available from the

348 corresponding author on reasonable request.

Acknowledgments

We thank funding by the Dirección General de Asuntos del Personal Académico, (PAPIIT

IN205616), and institutional funds of the Instituto de Investigaciones en Ecosistemas y

Sustentabilidad, Universidad Nacional Autónoma de México. EADA held a generous

graduate research fellowship from Consejo Nacional de Ciencia y Tecnología, México. We

also thank the generousity of the personnel of Zona Arqueológica de Teotihuacán,

especially to Ms. V Ortega.

References

1. Galloway, J. N. *et al.* Nitrogen Cycles: Past, Present, and Future. *Biogeochemistry*. **70**, 153–226 (2004).

2. Sala, O.E. *et al*. Global Biodiversity Scenarios for the Year 2100. *Science*. 287: 1770–1774 (2000).

3. Phoenix, G.K. *et al.* Atmospheric nitrogen deposition in world biodiversity hotspots: the need for a greater global perspective in assessing N deposition impacts. *Glob Chang Biol*. **12**, 470–476 (2006).

4. Duprè, C. *et al*. Changes in species richness and composition in European acidic grasslands over the past 70 years: the contribution of cumulative atmospheric nitrogen deposition *Glob Chang Biol*. **16**, 344–357 (2010).

5. Scherer-Lorenzen, M., Venterink, H. O., Buschmann, H. Nitrogen enrichment and plant invasions: the importance of nitrogen-fixing plants and anthropogenic eutrophication. *Biological invasions* (ed. Netwig, W.). 163-180 (Springer, 2007).

6. Silva, L. C. R., Gómez-Guerrero, A., Doane, T. A., Horwath, W. R. Isotopic and

nutritional evidence for species and site-specific responses to N deposition and elevated

CO_2 in temperate forests. *J. Geophys. Res. Biogeosci.* **120**, 1110–1123 (2015).

7. Reay, D. Nitrogen and Climate Change: An Explosive Story**.** (Palgrave Macmillan

2015).

8. Landrigan, P. J., *et al.* The Lancet Commission on pollution and health. *Lancet* 391

462-512 (2018).

9. Stevens. G. *et al*. Characterizing the Epidemiological Transition in Mexico: National

and Subnational Burden of Diseases, Injuries, and Risk Factors. *PLoS Medicine*, **5**,

e125. (2018).

10. Instituto Nacional de Estadística y Geografía. 2011. Censo de población y vivienda

2010.

http://web.archive.org/web/20120505010044/http://www.censo2010.org.mx/doc/cpv10

p_pres.pdf. (2011).

11. Díaz-Álvarez, E. A., Lindig-Cisneros, R., de la Barrera, E. Biomonitors of atmospheric

nitrogen deposition: potential uses and limitations. *Conserv. Physiol.* Conserv 6(1):

(2018).

12. Root, H. T., Geiser, L. H., Fenn, M. E. A simple tool for estimating throughfall nitrogen

deposition in foresta of western North América using lichens. *For. Ecol. Manage.* **306**,

1☐8 (2013).

13. Pitcairn, C. E. R. *et al*. Foliar Nitrogen as an Indicator of Nitrogen Deposition and

Critical Loads Exceedance on a European Scale. *Water Air Soil Pollut.* **130**, 1037–1042

(2001).

14. Harmens, H. *et al.* Nitrogen concentrations in mosses indicate the spatial distribution of

atmospheric nitrogen deposition in Europe. *Environ pollut.* **159**, 2852–60 (2011).

15. Zambrano García, A. *et al.* Distribution and sources of bioaccumulative air pollutants at

Mezquital Valley, Mexico, as reflected by the atmospheric plant *Tillandsia recurvata* L.

Atmos Chem Phys. **9**, 6479–6494 (2009).

16. Felix, J. D., Avery, G. B., Mead, R. N., Kieber, R. J., Willey, J. D. Nitrogen Content

and Isotopic Composition of Spanish Moss (*Tillandsia usneoides* L.): Reactive

Nitrogen Variations and Source Implications Across an Urban Coastal Air Shed.

Environ Process. **3**, 711–722 (2006).

17. Arciga, P. A. Evaluación del potencial bioindicador de la depositación de nitrógeno de

dos musgos comunes en México. (PS thesis). Universidad Michoacana de San Nicolás

de Hidalgo. (2010).

18. Secretaría del Medio Ambiente del Gobierno del Distrito Federal. Calidad del aire en la

Ciudad de México 2014, 120 pp. Dirección general de gestión de la calidad del aire,

Dirección de monitoreo atmosférico. (2015).

19. Castro, T. *et al.* Water-soluble inorganic ions of size-differentiated atmospheric

particles from a suburban site of Mexico City. *J. Atmos Chem.*

https://doi.org/10.1007/s10874-017-9369-5 (2017).

20. McMurray, J. A., Roberts, D. W., Fenn, M. E., Geiser, L. H., Jovan, S. Using Epiphytic

Lichens to Monitor Nitrogen Deposition Near Natural Gas Drilling Operations in the

Wind River Range, WY, USA. *Water Air Soil Pollut.* **224**, 1487 (2013).

21. Jovan, S., Riddell, J., Padgett, P. E., Nash III, T. H. Eutrophic lichens respond to

multiple forms of N: implications for critical levels and critical loads research. *Ecol*

Appl. **22**, 1910–1922 (2012).

22. Zechmeister, H. G. *et al*. Total Nitrogen Content and δ^{15}N Signatures in Moss Tissue:

Indicative Value for Nitrogen Deposition Patterns and Source Allocation on a

Nationwide Scale. *Environ. Sci. Technol.* **42**, 8661–8667 (2008).

23. Liu. X-Y., Xiao, H-Y., Liu, C-Q., Li, Y-Y., Xiao, H-W. Atmospheric transport of

urban-derived NH(x): Evidence from nitrogen concentration and ^{15}N in epilithic mosses

at Guiyang, SW China. *Environ. Pollut.* **156**, 715–22 (2008).

24. Liu, X-Y., Xiao, H-Y., Liu, C-Q., Li, Y-Y., Xiao, H-W. Stable carbon and nitrogen

isotopes of the moss *Haplocladium microphyllum* in an urban and a background area

(SW China): The role of environmental conditions and atmospheric nitrogen deposition.

Atmos. Environ. **42**, 5413–5423 (2008).

25. Stewart, G., Aidar, M., Joly, C., Schmidt, S. Impact of point source pollution on

nitrogen isotope signatures (δ^{15}N) of vegetation in SE Brazil. *Oecologia*, **131**, 468–472

(2002).

26. Xiao, H. Y., Tang, C. G., Xiao, H. W., Liu, X. Y., Liu, C. Q. Mosses Indicating

Atmospheric Nitrogen Deposition and Sources in the Yangtze River Drainage Basin,

China. *J. Geophys. Res.* **115**, D14301 (2010).

27. Liu X-Y, Koba K, Liu C-Q, Li X-D, Yoh M. Pitfalls and new mechanisms in moss

isotope biomonitoring of atmospheric nitrogen deposition. *Environ. Sci. technol.* **46**,

12557–66 (2012).

28. Schröder, W. *et al*. Mapping correlations between nitrogen concentrations in

atmospheric deposition and mosses for natural landscapes in Europe. *Ecol. Indic.* **36**,

563–571 (2014).

29. Felix, J. D., Elliott, E. M. Isotopic composition of passively collected nitrogen dioxide

emissions: Vehicle, soil and livestock source signatures. *Atmos. Environ.* **92**, 359–366

(2014).

30. Skudnik, M., Jeran, Z., Batič, F., Simončič, P., Kastelec, D. Potential environmental

factors that influence the nitrogen concentration and δ^{15}N values in the moss Hypnum

cupressiforme collected inside and outside canopy drip lines. *Environ. Pollut.* **198**, 78–

85 (2015).

31. Pavuluri, C. M., Kawamura, K., Tachibana, E., Swaminathan T. Elevated nitrogen

isotope ratios of tropical Indian aerosols from Chennai: Implication for the origins of

aerosol nitrogen in South and Southeast Asia. *Atmos Environ* **44**, 3597–3604 (2010).

32. Heldt, H. W., Piechulla, B. Plant Biochemistry, Fourth edition. (Academic Press 2011).

33. Paulissen, M. P. C. P., Van Der Ven, P. J. M., Dees, A. J., Bobbink, R. Differential

effects of nitrate and ammonium on three fen bryophyte species in relation to pollutant

nitrogen input. *New Phytol.* **164**, 451–458 (2004).

34. Liu, X-Y. *et al*. Ammonium first: natural mosses prefer atmospheric ammonium but

vary utilization of dissolved organic nitrogen depending on habitat and nitrogen

deposition. *New phytol.* **199**, 407–19 (2013).

35. Gordon, C., Wynn, J. M., Woodin, S. J. Impacts of increased nitrogen supply on high

Arctic heath: the importance of bryophytes and phosphorus availability. *New Phytol.*

149, 461–471 (2002).

36. Pitcairn, C. *et al*. Diagnostic indicators of elevated nitrogen deposition. *Environ. Pollut.*

144, 941–50 (2006).

37. Wiedermann, M. M., Gunnarsson, U., Ericson, L., Nordin, A. Ecophysiological

adjustment of two *Sphagnum* species in response to anthropogenic nitrogen deposition.

New Phytol. **181**, 208–217 (2009).

38. Redling, K., Elliott, E., Bain, D., Sherwell, J. Highway contributions to reactive nitrogen deposition: tracing the fate of vehicular NOx using stable isotopes and plant biomonitors. Biogeochemistry **116**, 261–274 (2013).

39. Schmitt, D-BAK., Martin, C. E., Lüttge, U. E. Gas Exchange and Water Vapor Uptake in the Atmospheric CAM Bromeliad *Tillandsia recurvata* L.: The Influence of Trichomes. *Plant Biol.* 102, 80–84 (1989).

40. Figuereido, A. M. G., Alcalá, A. L., Ticianelli, R. B. The use of *Tillandsia usneoides* L. as bioindicator of air pollution in Sao Paulo, Brazil. *J. Radioanal. Nucl. Chem.* **259**, 59–63 (2004).

41. Abril, A. B., Bucher, E. H. A comparison of nutrient sources of the epiphyte *Tillandsia capillaris* attached to trees and cables in Cordoba, Argentina. *J. Arid Environ.* **73**, 393□395 (2009).

42. Santos, J. J., *et al*. Modelagem da dispersão de poluentes de origem móvel (veicular) em curitiba e a ocorrência de bromélias em cabos da rede elétrica. *Blucher Engineering Proceedings*, **1**, 189–204 (2014).

43. Taiz, L., Zeiger, E. Plant Physiology, Fifth edition. (Sinauer Associates 2010).

44. Felix, J. D. *et al.* Examining the transport of ammonia emissions across landscapes using nitrogen isotope ratios. *Atmos. Environ.* **95**, 563–570 (2014).

45. Díaz-Álvarez, E. A., Reyes-García, C., de la Barrera, E. A δ^{15}N assessment of nitrogen deposition for the endangered epiphytic orchid *Laelia speciosa* from a city and an oak forest in Mexico. *J. Plant Res.* **129**, 863–872 (2016).

46. Guevara-Escobar, A. M., *et al.* Fog interception by Ball moss (*Tillandsia recurvata*). *Hydrol. Earth. Syst. Sci.* **15,** 2509–2518 (2011).

47. Park, N. Physicochemical and environmental plant physiology, fourth edition.

(Academic Press 2009).

48. Barrios-Hernández, E. Y. Organismos potenciales para estudios de bioindicación de la

depositación de nitrógeno en México. (PS thesis). Universidad Michoacana de San

Nicolás de Hidalgo. (2010).

49. Maron, J. L., Crone, E. Herbivory: effects on plant abundance, distribution and

population growth. *Proc. Biol. Sci.* **273**, 2575–2584 (2006).

50. Calderon, R. G., Rzedowski, J. Flora fanerogamica del Valle de México. pp.1406.

(Instituto de Ecologia y CONABIO, Xalapa, Veracruz, México 2001).

51. Servicio Meteorologíco Nacional. Normales climatológicas por estado 1981-2010.

http://smn.cna.gob.mx/es/informacion-climatologica-ver-estado?estado=df.

http://smn.cna.gob.mx/es/informacion-climatologica-ver-estado?estado=gto.

http://smn.cna.gob.mx/es/informacion-climatologica-ver-estado?estado=mex (2016).

52. Wang, W., Pataki, D. E. Spatial patterns of plant isotope tracers in the Los Angeles

urban region. *Landsc. Ecol.***25**, 35–52 (2010).

53. Ehleringer, J. R., Osmond, B. O. Stable isotopes. *Plant physiological ecology.* (ed.

Pearcy, R. W., Ehleringer, J. R., Mooney, H. A., Rundel, P. W.) 281-300 (Chapman &

Hall, 1989).

54. Evans, R. Physiological mechanisms influencing plant nitrogen isotope composition.

Trends Plant Sci. **6**, 121–126 (2001).

55. Cressie, N. Spatial prediction and ordinary Kriging. *Math. Geol* **20**, 405-421 (1988).

Author information

Affiliations

Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México, Mexico

City 04510, *Mexico*

Edison A. Díaz-Álvarez

Instituto de Investigaciones en Ecosistemas y Sustentabilidad, Universidad Nacional

Autónoma de México, Morelia, Michoacán 58190, Mexico

Edison A. Díaz-Álvarez & Erick de la Barrera

Contributions

Both authors planned the study, conducted field work, took part in the preparation of the manuscript, have reviewed the results, and have approved the final version of this manuscript. E.A.D.A. analyzed the data, prepared all of the figures, and wrote the initial version of the manuscript. E.d.l.B. procured funding, revised and approved the manuscript, and drove in the field and planned a gastronomic tour in the Mexico Valley.

Competing interests

The authors declare that they have no competing interests.

Corresponding author

Correspondence to Erick de la Barrera.

555   Table 1. Two-way ANOVA for responses of potential biomonitoring organisms growing in

556   the Valley of Mexico.

		d.f.	%N (dry weight)		C:N ratio		δ^{15}N (‰)	
			F	P	F	P	F	P
Anaptychia sp.	Site	33	26.16	<0.001	34.78	<0.001	29.04	<0.001
	Season	1	7.83	0.005	10.14	0.002	0.99	<0.001
	Site × Season	33	3.18	<0.001	3.22	<0.001	2.34	<0.001
Grimmia sp.	Site	31	34.33	<0.001	40.73	<0.001	123.04	<0.001
	Season	1	1.47	0.226	30.02	<0.001	2.95	0.087
	Site × Season	31	5.27	<0.001	3.25	<0.001	4.95	<0.001

Fabronia sp.	Site	29	24.92	<0.001	15.50	<0.001	46.87	<0.001
	Season	1	2.57	0.11	6.14	0.014	1.44	0.232
	Site × Season	29	5.80	<0.001	3.93	<0.001	6.62	<0.001
Tillandsia recurvata	Site	21	34.82	<0.001	33.92	<0.001	57.47	<0.001
	Season	1	96.05	<0.001	6.53	0.011	136.29	<0.001
	Site × Season	21	16.02	<0.001	3.97	<0.001	7.56	<0.001

Figure legends

Figure 1. Localization of the Valley of Mexico. Red and yellow dots represent the spatial

distribution of the air quality network stations of the wet deposition and the

automatic monitoring network for NOx, respectively. This network is located

mainly in the Mexico City and its metropolitan area. Green triangles represent the

sites where biomonitors were collected throughout the Valley. The red line delimits

the basin, the white line indicates state division, and the blue line shows Mexico

City limits. The map was created with ArcGIS 10 (Esri, Redlands, California,

USA). Image data: Google Earth; image date: 5 September 2016.

Figure 2. Spatial distribution of the total wet deposition in Kg N ha^{-1} year^{-1} during 2014

(A), and atmospheric concentration of NOx in ppb (B). Data is available for public

access in the website of the Mexico City government

(http://www.aire.cdmx.gob.mx/default.php). The map was created with ArcGIS 10

(Esri, Redlands, California, USA). Image data: Google Earth; image date: 5

September 2016.

Figure 3. Spatial distribution of the nitrogen content (A, C, E, G) and δ^{15}N values (B, D, F,

H) for *Anaptychia* sp. (A, B), *Grimmia* sp. (C, D), *Fabronia* sp. (E, F) and

Tillandsia recurvata (G, H). The map was created with ArcGIS 10 (Esri, Redlands,

California, USA). Image data: Google Earth; image date: 5 September 2016.

Figure 4. Relationship between wet deposition of ammonium (green circles), nitrate (blue

squares), and total deposition, $NH_4^+ + NO_3^-$ (red triangles) during 2014 and the

δ^{15}N values of the moss *Grimmia* sp.

Figure 5. Relationship between wet deposition, of ammonium (green circles), nitrate (blue

592   squares) and total deposition, $NH_4^+ + NO_3^-$ (red triangles) during 2014 and the $\delta^{15}N$

593   values of the moss *Fabronia* sp.

594   **Figure 6**. Relationship between NOx concentration during the 2014 dry season and the

595   nitrogen content (A), C:N ratio (B), and the $\delta^{15}N$ values (C) of the bromeliad

596   *Tillandsia recurvata.* Open circles were excluded from the regression analysis

597   because they were collected from non-typical environmental conditions that skewed

598   the isotopic signatures of *T. recurvata* to very negative values.