A transformation from latency to ensemble coding

in a model of piriform cortex

Merav Stern[1,2,4], Kevin A. Bolding[3], L.F. Abbott[2], Kevin M. Franks[3] *

[1] Edmond and Lily Safra Center for Brain Sciences, Hebrew University, Jerusalem 9190401 Israel

[2] Department of Neuroscience, Zuckerman Mind Brain Behavior Institute, Columbia University, New York, New York 10027, USA

[3] Department of Neurobiology, Duke University School of Medicine, Durham, North Carolina 27705-4010, USA

[4] Present address: The Raymond and Beverly Sackler Scholars Program in Integrative Biophysics at the University of Washington, Seattle, Washington, 98195-3925, USA

* Correspondence: Kevin Franks

Bryan Research Building, Rm. 401D

311 Research Dr.

Durham, NC, 27705

(919) 684-3487

franks@neuro.duke.edu

Author contributions:

M.S., L.F.A. and K.M.F. conceived the project, performed and analyzed modeling studies, and wrote the manuscript with input from K.A.B.

Competing interests:

We have no financial or non-financial competing interests.

ABSTRACT

We present a spiking network model that transforms odor-dependent variable-latency olfactory

bulb responses into a cortical ensemble code. In the model, which captures basic circuit properties

of piriform cortex, the impact of the earliest-activated bulb inputs on the cortical response is am-

plified by diffuse recurrent collateral excitation, which then recruits strong feedback inhibition that

stabilizes cortical activity and decreases the impact of later-responding glomeruli. Because the

sequence of olfactory bulb activity for a particular odor is preserved across concentration, the en-

semble of activated cortical neurons is robust to concentration changes. Nevertheless, odor con-

centration is represented by the latency and synchrony of the ensemble response. Using decoding

techniques, we show that the ensemble-based coding scheme that arises in the cortical model sup-

ports concentration-invariant odor recognition.

INTRODUCTION

The coding schemes used to represent odors in the olfactory bulb and piriform cortex (PCx) are

different. In the bulb, odorants sequentially activate distinct subsets of glomeruli causing mitral

and tufted cells (MTCs) to begin firing at various odor- and cell-specific latencies following the

onset of inhalation (Spors and Grinvald 2002, Bozza et al. 2004, Bathellier et al. 2008, Carey et

al. 2009, Cury and Uchida 2010, Shusterman et al. 2011). In the PCx, this latency code is trans-

formed so that odors are represented by ensembles of neurons distributed across the piriform (Illig

and Haberly 2003, Rennaker et al. 2007, Poo and Isaacson 2009, Stettler and Axel 2009) with

spike timing providing little information about odor identity in the PCx (Miura et al. 2012, Uchida

et al. 2014) (see also Bolding & Franks, *submitted*). Here we construct a model, based on the circuit

architecture of PCx to explore the mechanisms that support the transformation from a bulbar la-

tency to a cortical ensemble odor code.

In mammals, odor perception begins when volatile molecules are inhaled and bind to odorant re-

ceptors expressed by olfactory sensory neurons (OSNs) in the nasal epithelium. Each OSN ex-

presses just one of ~1000 different odorant receptor genes, and all OSNs expressing a given re-

ceptor converge on a unique pair of glomeruli in the olfactory bulb (Wilson and Mainen 2006).

MTCs receive excitatory inputs from a single glomerulus, and their projections to PCx are diffuse

and overlapping (Ghosh et al. 2011, Miyamichi et al. 2011, Sosulski et al. 2011), allowing indi-

vidual pyramidal piriform neurons to receive inputs from different combinations of co-activated

glomeruli (Franks and Isaacson 2006, Suzuki and Bekkers 2006, Apicella et al. 2010, Davison and

Ehlers 2011). In addition to their connections onto pyramidal cells, MTCs make excitatory con-

nections onto layer 1 inhibitory interneurons that provide feedforward inhibition to pyramidal cells

(Luna and Schoppa 2008, Stokes and Isaacson 2010, Suzuki and Bekkers 2012). Pyramidal cells

form a widespread recurrent collateral excitatory plexus that may amplify the effect of the earliest

MTCs by recruiting additional pyramidal cells that initially receive slightly subthreshold bulb in-

put (Franks et al. 2011). Recurrent excitation also recruits strong feedback inhibition from layer

3 interneurons that should, along with the feedforward inhibition, lessen the impact of later MTCs

inputs (Franks et al. 2011, Suzuki and Bekkers 2012, Large et al. 2016). Piriform circuits are there-

fore poised to dramatically reshape odor representations (Davison and Ehlers 2011, Poo and

Isaacson 2011, Haddad et al. 2013).

Although MTCs are activated at different times across the full respiration cycle (Bathellier et al.

2008, Cury and Uchida 2010, Shusterman et al. 2011), rodents can rapidly discriminate different

odors within ~100 ms (Uchida and Mainen 2003, Abraham et al. 2004, Rinberg et al. 2006), well

before most responsive glomeruli are activated (Wesson et al. 2008). Thus, the earliest activated

glomeruli may provide most information about odor identity (Hopfield 1995, Schaefer and Margrie

2007, Schaefer and Margrie 2012, Wilson et al. 2015). Interestingly, most odors retain their per-

ceptual identity across a range of concentrations (Krone et al. 2001, Laing et al. 2003, Homma et

al. 2009). Note, however, that while increasing odor concentration systematically decreases MTCs

onset latencies (Cang and Isaacson 2003, Margrie and Schaefer 2003, Spors et al. 2006, Junek et

al. 2010, Gschwend et al. 2012) and can increase the total number of activated glomeruli, the

sequences in which glomeruli are activated it preserved (Spors and Grinvald 2002, Junek et al.

2010). Thus, decoding the sequence in which glomeruli, and their associated MTCs, are activated

has been hypothesized to provide a basis for concentration-invariant odor identification (Hopfield

1995, Junek et al. 2010, Schaefer and Margrie 2012, Wilson et al. 2015). Our model supports this

hypothesis by providing a mechanism for its implementation in piriform cortex.

RESULTS

We use a modeling approach to provide mechanistic insight into how the transformation from the

latency code in the bulb to the ensemble code in PCx is implemented. We developed a spiking

network model that simulates patterns of odor-evoked mitral cell activity, and this provides input

to a spiking network mode that captures the specific circuit properties of PCx. We simulated bulb

and PCx spiking activity over the course of a single respiration cycle consisting of a 100 ms exha-

lation followed by a 200 ms inhalation.

<u>Odors activate distinct ensembles of piriform neurons</u>

Our model bulb consisted of 900 glomeruli that are each innervated by a unique family of 25 mitral

cells. Odor identities are defined by sets of glomerular onset latencies so that different odors acti-

vate specific subsets of glomeruli and their associated mitral cells with odor-specific latencies after

the onset of inhalation (Figure 1A, Supplemental Figure S1, and Methods). At our reference con-

centration, 10% of the glomeruli are activated during the 200 ms sniff.

The PCx model contains 10,000 excitatory pyramidal cells, each of which receives 50 excitatory

inputs from a random subset of the mitral cells and 1,000 recurrent excitatory inputs from a random

subset of other pyramidal cells (Figure 1B). Our model also includes 1,225 feedforward inhibitory

neurons that receive input from mitral cells and provide synaptic inhibition onto the pyramidal

cells and other feedforward interneurons, and a separate population of 1,225 feedback inhibitory

neurons that each receive inputs from a random subset of pyramidal cells and provide inhibitory

input locally onto pyramidal cells. We model all three classes of PCx neurons as leaky integrate-

and-fire neurons with current-based synaptic inputs (details and model parameter values are given

A

Figure 1. Odors activate distributed ensembles of PCx neurons.
(A) Raster plots showing spiking of 1,000 mitral cells (out of 22,500 total; 25 mitral cells/glomerulus, 40 glomeruli shown) in response to 3 different odors. Each row represents a single mitral cell and all mitral cells belonging to each glomerulus are clustered. Tick marks indicate spike times. Inhalation begins at 0 ms and is indicated by the grey shaded region. The red curve shows the cumulative number of glomeruli activated across the sniff, and the blue curve is the firing rate averaged across all mitral cells. (B) Schematic of the piriform cortex model. (C) Voltage traces for three sequential sniffs in 4 model pyramidal cells. Time of inhalation is indicated by the dashed line. (D) Single-trial population activity map for all 10,000 pyramidal cells. Each pixel represents a single cell, and pixel color indicates the number of spikes fired during the 200 ms inhalation. Approximately 13% of cells fired at least 1 action potential, with activated cells randomly distributed across the cortex. (E) Response vectors shown for 20 cells in response to different odors presented on 4 sequential trials. Spiking levels are low for no-odor controls. Note the trial-to-trial variability and that individual cells can be activated by different odors.

in the Methods). Most of our analyses focus on pyramidal cell activity because these cells receive

bulb input and provide cortical output and thus carry the cortical odor code.

Low levels of spontaneous PCx spiking in the model are driven by baseline activity in mitral cells

(1-2 Hz), with 2.8 ± 0.4 % (mean $\pm$ st. dev) of pyramidal cells spiking during the 200 ms inhalation

in the absence of odor, consistent with experimental findings in awake (Zhan and Luo 2010, Miura

et al. 2012) and anesthetized (Poo and Isaacson 2009) animals. Given this low level of spontaneous

activity, cells that fire at least one action potential during the 200 ms inhalation are defined as

"activated". Because each piriform cell receives input from a random subset of mitral cells, differ-

ent odors selectively and specifically activate distinct, sparse subsets of cortical pyramidal cells

(Figure 1C) so that each cell is responsive to multiple odors and each odor activates distinct en-

sembles of neurons distributed across PCx (Figure 1D,E). At our reference concentration, for

which 10% of glo-

meruli are activated,

14.1 ± 0.59 % (mean

$\pm$ st. dev., n = 6

odors) of piriform

pyramidal cells fire

at least one action

potential during a

sniff, which is also

consistent with ex-

perimental data (Poo

Supplementary Figure 1. Modeling odor responses in olfactory bulb
(A) Raster for all 22,500 mitral cells for one odor trial. Grey shading indicates inhalation phase. Inset shows expanded view of 1,000 mitral cells belonging to 40 different glomeruli. (B) Raster showing trial-to-trial variability for a single activated glomerulus. Each box represents a different sister mitral cell, with trials 1-4 represented by the lines within each box.

and Isaacson 2009, Stettler and Axel 2009, Miura et al. 2012). In summary, we have simulated

odor-evoked activity in olfactory bulb that we use as input to a model piriform cortex. Population

odor responses in the model broadly recapitulate experimental observations. We can therefore use

the model to probe how these population responses emerge mechanistically.

<u>Evolution of cortical odor ensembles</u>

We next illustrate how the model PCx ensembles evolve over time by comparing the spiking ac-

tivity of its four different classes of neurons (mitral cell, pyramidal cell, feedforward and feedback

inhibitory) over the course of a single sniff (Figure 2A). Preceding inhalation, baseline activity in

mitral cells drives low levels of spiking in both pyramidal cells and feedforward inhibitory neu-

rons. Feedback inhibitory neurons, which do not receive mitral cell input, show no baseline activ-

ity. Shortly after inhalation, inputs from the earliest activated glomeruli initiate a dynamic cascade

of cortical activity, characterized by a transient and rapid burst of spiking in a small subset of

pyramidal cells that is sharply truncated by the strong and synchronous recruitment of feedback

inhibitory neurons. The feedback inhibitory response suppresses pyramidal cell firing for a brief

period before the network settles into a sustained state with somewhat elevated pyramidal cell

activity that both drives and is held in check by feedback inhibition (Figure 2A). This activity

remains steady for the remainder of the inhalation, even though additional mitral cells respond,

because the activity of feedforward inhibitory neurons ramps up to cancel the increase in total

mitral cell input.

What triggers the rapid transient pyramidal cell response? Each odor initially activates a subset of

glomeruli that project randomly onto different cortical pyramidal cells. A small subset of pyrami-

dal cells receives enough input from short-latency mitral cells to reach threshold and start spiking

Figure 2. Evolution of a cortical odor response.
(A) Raster for a single sniff showing spiking activity of a subset of mitral cells (2,250 out of 22,500), all 1,225 feedforward interneurons (FFIs), all 10,000 pyramidal cells, and all 1,225 feedforward interneurons (FBIs). Spiking rate for the population of pyramidal cells is shown at the bottom (average of 6 trials). Note that the earliest activated glomeruli initiate a cascade of pyramidal cell spiking that peaks after ~50 ms and is abruptly truncated by synchronous spiking of FBIs. (B) Single-trial voltage traces (black) for 3 pyramidal cells in response to the same odor. Inhalation onset is indicated by the dashed line. The red traces show the bulb input and the green traces the recurrent input received by each cell. Cell 1 receives strong bulb input and spikes soon after odor presentation. Cell 2 receives subthreshold input from the bulb and only spikes after receiving addition recurrent input from other pyramidal cells. Cell 3 receives no early odor-evoked input from the bulb, and its recurrent input is subthreshold, so it does not spike over the time period shown.

early in the sniff (Figure 2B, cell 1). This activity produces a small amount of recurrent excitation

that is dispersed across the cortex via the long-range recurrent collateral connections. The resulting

recurrent input is not strong enough to drive spiking by itself (Figure 2B, cell 3), but it can recruit

other pyramidal cells that receive moderate but subthreshold bulb input (Figure 2B, cell 2). Be-

cause more cells received subthreshold than suprathreshold bulb input, more pyramidal cells are

activated by the recurrent input, resulting in even stronger recurrent excitation that, in turn, can

help activate even more pyramidal cells receiving even less bulb input. The result is an explosive

increase in total pyramidal cell activity, and therefore a steady increase in the strength of recurrent

excitation. However, recurrent excitation onto feedback inhibitory neurons is stronger than onto

other pyramidal cells so that feedback inhibitory neurons are recruited before pyramidal cells that

have not received any direct bulb input. Thus, feedback inhibition quickly halts the explosive

growth of pyramidal cell firing because pure recurrent input always remains subthreshold for py-

ramidal cells, thereby maintaining the odor-specificity of the cortical ensemble.

<u>Specific roles for different circuit elements in shaping cortical responses</u>

To reveal the specific roles that different circuit elements play in shaping piriform output we com-

pared responses in the full circuit with those obtained after removing different circuit elements

(Figure 3). The same odor stimulus was used in all cases, so input from the olfactory bulb is iden-

tical except for the trial-to-trial stochasticity of mitral cell spiking. We first compared responses in

the full circuit (Figure 3A) with those in a purely feedforward network in which pyramidal cells

only receive mitral cell inputs (Figure 3B). In this highly reduced, feedforward circuit, pyramidal

cell spiking activity grows more slowly, without any strong initial transient, and it tracks the num-

ber of spiking mitral cells. In the absence of either feedforward or feedback inhibition, the cortical

response continues to grow over the course of the sniff as more glomeruli are activated (Figure

1A), so that a large fraction of cells respond at some point during the course of the entire sniff.

We next examined networks without feedforward or feedback inhibition, or without recurrent ex-

citation and excitatory input to feedback interneurons. Eliminating only feedforward inhibition

increases the amplitude of the peak pyramidal response by about 50%, although the general shape

Figure 3. Distinct roles for different inputs in shaping ensemble response.
(A-D) Raster plots for presentations of the same odor using different circuit configurations. The percentages of active pyramidal cells are shown in the insets. (A) As in Figure 2A. (B) Network activity when pyramidal cells get excitatory input from mitral cells but without FFI, recurrent excitation or FBI. Pyramidal cell spiking tracks mitral cell input. Population rate for the full network is shown in grey for comparison. (C) Network activity after eliminating excitatory input to FFIs. Note that sustained FBI spiking increases without FFI, resulting in roughly constant levels of sustained spiking in pyramidal cells. (D) Network activity with no recurrent excitation onto either other pyramidal cells or FBIs. The transient activity peak seen in A-D is absent, and sustained activity is slightly higher. (E) Population rate plots for 3 different odors with the full network (black trace), no FFI (red trace) or no recurrent excitation/FBI (green trace). Insets expand the period around inhalation. Note the population spiking is higher and peaks slightly earlier without FFI, and that responses are slow without recurrent excitation. All population spike rate plots are averaged over 6 trials.

of the population response is largely unchanged (Figure 3C,E) and the fraction of cells activated

over the entire sniff only increases modestly (from 13% to 16%). When we selectively eliminated

feedback inhibition, odor induces unchecked runaway recurrent excitation, and all the cells end up

spiking vigorously (not shown). Finally, we simulated the circuit with pyramidal cell output

blocked, which eliminates both recurrent excitation and the recruitment of feedback inhibition.

Under these conditions pyramidal cell spiking initially ramps slowly, similar to the feedforward-

only condition, but then plateaus within ~50 ms, as feedforward inhibition is recruited (Figure

3D,E). Although the rapid peak in instantaneous spiking activity is substantially decreased under

these conditions, activity over the whole sniff increases, with 20% of the pyramidal cells respond-

ing. These results illustrate the istinct roles that different elements of the model PCx play in shap-

ing odor responses.

Piriform responses are shaped by early-responding glomeruli

The previous analysis showed that population spiking peaks early when recurrent excitation is

present (Figures 3A,C) but ramps up more slowly when it is eliminated (Figures 3B,D), indicating

a key role for intracortical circuitry in amplifying the initial response. We examined this directly

by comparing population spiking to the sequential activation of individual glomeruli (Figure 4A).

In the full network, population spiking peaks 34 ± 8.3 ms after inhalation onset (mean ± st. dev.

for 6 odors with ensemble averages of 6 trials per odor, at the reference concentration; Figure

4B,C). At this time, only 15 ± 1.4 glomeruli have been activated out of the 95 ± 6.0 glomeruli that

will eventually be activated across the full sniff. In other words, at its peak, PCx activity is driven

by the earliest ~15% of activated glomeruli. Mean responses peak slightly earlier when feedfor-

ward inhibition is eliminated (28 ± 4.5 ms; Figure 4B), but the peak activity is still driven by a

similar number of glomeruli (12 ± 0.80 glomeruli; Figure 4C); on individual trials, the peak re-

sponse is larger without feedforward inhibition (Figure 3C,D), but this difference is not captured

Figure 4. Earliest-active glomeruli define the cortical response.
(A) Normalized population spike rates (black) in response to an odor during the sniff cycle shown in grey. The red curve shows the cumulative number of glomeruli activated across the sniff. Note that population spiking peaks after only a small subset of glomeruli have been activated. (B) Normalized population spike rates for one odor for the full network (black trace), without FFI (red trace) and without recurrent excitation (green trace). Grey trace shows the cumulative number of activated glomeruli. (C) Fraction of peak population spike rate as a function of the cumulative number of activated glomeruli for 6 different odors. These curves indicate the central role recurrent excitation plays in amplifying the impact of early-responsive glomeruli. (D) Average correlation coefficients for repeated same-odor trials and pairs of different-odor trials measured over the full 200 ms inhalation. (E) As in D but measured over the first 50 ms after inhalation onset. (F) Ratios of correlations for same- vs. different-odor trials measured over the full sniff (grey bar on left) and over the first 50 ms (black bar on right).

in the average responses). In contrast, population spiking peaks much later without recurrent exci-

tation (139 ± 29 ms) at a time when most of the responsive glomeruli have been activated (66 ±

0.44; Figure 4C). Recurrent excitation therefore plays an important role in shaping the dynamics

of the cortical odor response by amplifying the impact of early-responsive glomeruli and discount-

ing the impact of later-responding glomeruli through the recruitment of strong feedback inhibition

(Franks et al. 2011).

We wondered whether the earliest part of the cortical response provides a more distinctive repre-

sentation of odor identity than the later response, so we quantified responses as vectors of spike

counts, one component for each pyramidal cell, either over either the full 200 ms inhalation or

only the first 50 ms after inhalation onset. We then compared pair-wise correlations between re-

sponse vectors on either same-odor trials or trials involving different odors. Even though glomer-

ulus onset latencies are identical in all same-odor trials, stochastic mitral cell firing results in con-

siderable trial-to-trial variability (Figure 1E). Consequently, correlations coefficients are low for

same-odor trial pairs over the full sniff (0.35 ± 0.010, mean ± st. dev. for multiple same-odor trial

pairs using 6 different odors; Figure 4D). Nevertheless, same-odor responses are considerably

more correlated than pairs of responses to different odors (0.11 ± 0.016; mean ± st. dev. for pairs

of the same 6 odors). Response correlations to both same-odor and different-odor responses are

lower when using only first 50 ms (same-odor, 0.24 ± 0.019; different-odor pairs, 0.044 ± 0.014;

Figure 4E). However, the ratio of correlations for same- vs. different-odor responses, which can

be thought of as a signal-to-noise ratio, is almost double in the first 50 ms versus the full 200 ms

inhalation (Figure 4F). Taken together, our model predicts that a cascade of cortical activity is

initiated by the earliest-responsive inputs and amplified by recurrent excitation, providing a dis-

tinctive odor representation, and then truncated by feedback inhibition. The cortical odor response

is therefore largely shaped by the glomeruli that respond earliest in the sniff.

<u>Distinct roles for feedforward and feedback inhibition in normalizing PCx output</u>

We next asked how cortical odor representations depend on odor concentration. Onset latencies

decrease with increasing concentrations of odorant (Cang and Isaacson 2003, Junek et al. 2010,

Sirotin et al. 2015), which we simulate in our olfactory bulb model (Figure 5A) by shrinking the

onset latencies from those at the reference concentration. In other words, to decrease odor concen-

tration, we uniformly stretch latencies causing fewer glomeruli to be activated within 200 ms, and

making those that do activate respond later. Conversely, we shrink the set of latencies to simulated

higher concentrations so that glomeruli are activated earlier. Importantly, stretching or shrinking

latencies does not change the sequence in which glomeruli become activated (Figure 5A). We

quantify odor concentration using the fraction of activated glomeruli. Note that given the non-

linear concentration-dependence of receptor activation and extensive normalization at each multi-

ple stages of the system upstream of the cortex (Cleland et al. 2011), a 10-fold increase in mitral

cell output is expected to correspond to a much, much greater range of concentration.

The number of responsive pyramidal cells could scale with the total amount of glomerular input,

which increases at higher concentrations. However, PCx odor ensembles are buffered against

changes in odor concentration (Figure 5B). Across the population, we found that the number of

responsive pyramidal cells only increases by 50% upon a 10-fold increase in input (Figure 5C).

This indicates that the size of cortical odor ensembles is only weakly concentration-dependent,

consistent with in vivo imaging studies (Stettler and Axel 2009). In addition, both the total number

of spikes across the population (Figure 5D) and the number of spikes evoked per responsive cell

(not shown**)** are only modestly concentration-dependent. Our model therefore shows that piriform

circuitry can normalize intensity-scaled input from the olfactory bulb.

To gain insight into how this form of response normalization is implemented, we again simulated

responses either without feedforward inhibition or without recurrent excitation and feedback inhi-

bition. Eliminating feedforward inhibition increases both the number of responsive cells (Figure

Figure 5. Cortical output is normalized across concentrations.
(A) Mitral cell rasters for 2 odors at 3 different concentrations defined by the fraction of co-active glomeruli. Glomerular onset latencies decrease with concentration allowing more glomeruli to become active over the sniff period. Odors are different from the odors in Figure 1. (B) Single-trial piriform response vectors for single trials over a concentration range corresponding to 3-30% active glomeruli. Note that activity does not dramatically increase despite the 10-fold increase in input. (C) Fraction of activated pyramidal cells at different odor concentrations for the full network (black trace), without FFI (red trace) and without recurrent excitation (green trace) for 4 different odors (open circles, thin lines) and averaged across odors (filled circles, thicker lines). Note that eliminating FFI primarily shifts the number of responsive cells, indicating that FFI is largely subtractive, whereas eliminating recurrent excitation alters the gain of the response. Note also that recurrent excitation amplifies the number of activated cells at low odor concentrations. (D) As in C but for the total number of spikes across the population. (E) Perceptron classifications of an odor at different concentrations on the basis of pyramidal cell activity. Either the transient cortical activity (first 50 ms of the inhalation; black curve) or the activity across the full inhalation (gray curve) was used for both training and testing. Training was solely on the basis of the concentration indicated by the black arrow. The dashed line shows the chance level of classification (50%).

5C) and total population spiking (Figure 5D). However, this increase is fairly uniform across con-

centrations and removing feedforward inhibition does not substantially change the gain of the re-

sponse (i.e. how rapidly these responses vary with input strength; Figure 5C&D), indicating that

the effect of feedforward inhibition is largely subtractive. In contrast, eliminating recurrent exci-

tation and feedback inhibition destroys concentration invariance by dramatically increasing re-

sponse gain, indicating that they implement divisive normalization (Carandini and Heeger 2012).

Interestingly, cortical output is decreased at low odor concentrations when recurrent excitatory and

feedback inhibition are removed, indicating that recurrent collateral excitation amplifies cortical

output in response to weak input (Figure 5C&D).

<u>Early-activated PCx cells support concentration-invariant odor recognition</u>

Because the sequence of glomerular activation latencies is preserved across concentrations, odor

representations defined largely by the earliest activated glomeruli could support concentration-

invariant odor recognition. We therefore asked whether the cortical odor representation in our

model was well-suited to this purpose. Specifically, we next asked if a downstream observer could

reliably identify an odor using population spiking, and whether the odor can be recognized when

presented at different concentrations. To do this we trained a perceptron to identify a specific odor

at one concentration (10% active glomeruli) and then asked how well it could identify that odor

presented at different concentrations. We used spike counts over either the full 200 ms inhalation

or the first 50 ms as input. Perceptron performance was excellent when trained and tested at a

single concentration, indicating that despite considerable trial-to-trial variability, responses to dif-

ferent odors can be distinguished reliably (Figure 5E). We then examined classifier performance

when tested on different concentrations without retraining. Performance using the full 200 ms re-

sponse was generally excellent but fell off at both the lowest and highest concentrations. However,

classification using only the early transient response as input was essentially perfect across all

tested concentrations. This decoding analysis supports the idea that the earliest cortical response

provides an especially good substrate for concentration-invariant odor identification.

<u>Strategies for encoding odor intensity</u>

How is odor intensity represented in PCx? Because both the number of responsive neurons and

total spiking are relatively constant across concentrations (Figure 5C-E), spike rate seems unlikely

to be used to represent intensity. We therefore examined the dynamics of population spiking in

response to odors at different concentrations (Figure 6A). As with the mitral cell activity, cortical

response latencies decrease at higher concentrations by a factor of about 2 over a 10-fold increase

in input from bulb (Figure 6B), suggesting that a latency code could be used. We also found that

the peak of the population response increases by more than a factor of 5 over this range of odor

concentration (Figure 6C). Given that total spike output is largely constant, this result indicates

that the synchrony of the population piriform response is particularly concentration-dependent.

Furthermore, unlike latency, synchrony thus provides an representation of odor intensity that does

not require an external reference. The increase in synchrony occurs primarily because activation

latencies of the earliest responding glomeruli, which cannot be activated before inhalation onset,

compress as odor concentrations increase. causing responses to become more coincident. Thus,

the early peak in the PCx response can be used to rapidly decode both odor identity and concentra-

tion.

Figure 6. Effect of odor concentration on response timing
(A). Normalized population spike rates for 3 odors at 3 concentrations denoted by percentage of glomeruli responding. Peak responses are higher and occur earlier at higher concentrations. Dashed lines indicate inhalation onset. Average latencies to peak (B) and peak rate (C) vs. number of activated glomeruli.

DISCUSSION

The model we have developed and analyzed provides mechanistic insight into the transformation from an odor code using latency in olfactory bulb to a spatial, ensemble code in the PCx (for an alternative model, see (Sanders et al. 2014)). Individual glomeruli are activated at different latencies after inhalation onset, with glomeruli corresponding to the highest affinity receptors for a given odorant activated earliest. Only a small subset of responsive piriform neurons receives input from many of these early-activated inputs. However, because pyramidal cells are connected via long-range recurrent collateral inputs, these few activated cells provide diffuse excitatory synaptic input across PCx that can bring a larger subset of pyramidal cells that received subthreshold bulb input to threshold. This cascade of cortical activity continues until feedback inhibitory neurons are activated, which then strongly suppresses subsequent cortical spiking. Thus, the earliest activated glomeruli play a dominant role generating PCx odor ensembles. Increasing odor concentration systematically decreases glomerular onset latencies but does not change the sequence in which they are activated. Consequently, a given odor will activate similar ensembles of PCx neurons over a large range of concentrations. In this way, odor identity can be decoded across a wide range of concentrations. Odor concentration, on the other hand, is primarily encoded by the degree of synchrony of the cortical response.

Different odors activate distinct subsets of glomeruli and the number of activated glomeruli increases with odorant concentration (Rubin and Katz 1999, Meister and Bonhoeffer 2001, Spors and Grinvald 2002). However, considerable processing of this input occurs within olfactory bulb so that mitral cells can either be activated or suppressed by odor (Shusterman et al. 2011, Fukunaga et al. 2012, Economo et al. 2016), and normalization within the bulb (Cleland et al. 2011) entails

that MTC output scales much less steeply with concentration than the number of activated glomer-

uli (Banerjee et al. 2015, Sirotin et al. 2015, Roland et al. 2016). Nevertheless, here we have used

a highly simplified olfactory bulb model in which mitral cells are only activated by odor and the

number of activated mitral cells increases steeply with concentration. This has allowed us to ex-

amine the sufficiency of the cortical circuitry in transforming the temporal bulb representation and

normalizing output. Adding suppressed responses would slightly decrease the peak and deepen the

subsequent trough of the cortical response (Figure 2), and some normalization across concentra-

tions within the bulb would result in even less concentration-dependence in PCx (Figure 5). We

have also not attempted to separately model mitral and tufted cell responses. In fact, odor-evoked

responses in tufted cells occur reliably with short latencies whereas mitral cells, whose activity we

simulate, fire slightly later and with latencies that decrease with concentration (Fukunaga et al.

2012). This would be expected to produce a somewhat biphasic bulb and cortical response, with a

more rapid and concentration-invariant component and a later, concentration-dependent compo-

nent. Recent simultaneous bulb-PCx recordings are consistent with these predictions (Bolding and

Franks, *submitted*). We have also not attempted to distinguish here between different subclasses

of principal neurons (e.g. semilunar cells versus superficial pyramidal cells) or inhibitory inter-

neurons (e.g. somatostatin versus parvalbumiun-expressing GABAergic interneurons) in this

model. No doubt incorporating these distinctions will alter network behavior somewhat but the

overall dynamics of this network and general conclusions that we draw here are expected to be

robust, and therefore these refinements are appropriate for future, updated versions of the model

but beyond the scope of this project.

Our model predicts considerable trial-to-trial variability for same-odor responses and very low

correlations between responses to different odors. Both these values are lower than recent experi-

mental measures obtained by in vivo imaging (Roland et al *submitted*) or extracellular spiking

(Bolding & Franks, *submitted*). Responses correlations to different odors in the model PCx are

close to zero, which is expected because bulb inputs are random and cells receive no other sources

of excitation. However, what underlies the high trial-to-trial variability? In our model, mitral cell

spiking is governed by a Poisson process. Because inputs from OB to PCx are randomly assigned,

this stochasticity is reflected in the input to individual pyramidal cells from mitral cells that are

innervated by the same glomerulus (i.e. sister mitral cells). However, sister mitral cells are electri-

cally coupled by gap junctions in their apical dendrites (Christie et al. 2005) and spiking in pairs

of sister mitral cells can be highly correlated (Schoppa 2006). Correlated spiking in sister mitral

cells would more synchronously innervate the set of piriform neurons receiving input from an

activated glomerulus. The effect of this electrical coupling could be determined directly by exam-

ining the reliability of cortical odor responses in mice lacking connexin-36. In addition, recurrent

connections between pyramidal cells in the model are also random. However, these synapses are

plastic and are thought to provide a substrate for forming assemblies of similarly responsive neu-

rons (Haberly 2001, Wilson and Sullivan 2011). Such assemblies could help complete ensembles

of cells that are regularly co-activated, thereby decreasing trial-to-trial variability. Thus, correlated

spiking in sister mitral cells and specific recurrent connectivity between cortical pyramidal cells

may increase the fidelity of cortical odor responses.

Piriform cortex neurons are interconnected via a diffuse, long-range recurrent collateral excitatory

plexus that also recruits strong and scaled feedback inhibition (Johnson et al. 2000, Franks et al.

2011, Suzuki and Bekkers 2012). Our model shows how this circuitry amplifies the impact of the

earliest responsive glomeruli and discounts the impact of glomeruli that respond later. Why would

a sensory system discard so much information about the stimulus? To represent a large extent of

potential odorant space, the olfactory system employs a huge number of distinct odorant receptors

that are each highly selective for a small fraction of potential odorants. Thus, any given odorant

will strongly activate only a few, optimally-tuned receptors, while many other receptors will be

moderately or weakly activated if they receive input from sensory neurons expressing receptors

with lower affinity for the odor, especially at high concentrations. However, the few odorant re-

ceptors with the highest affinities to a given odorant will always be more strongly activated, and

therefore their associated glomeruli and MTCs will be activated earlier than those receiving input

from lower affinity receptors, regardless of odorant concentration. By defining odors according to

the earliest responding glomeruli, the olfactory system retains the specificity of the odor represen-

tation and discards spurious information provided by non-specific receptor activations.

MATERIALS AND METHODS

Modeling

Model olfactory bulb

The model bulb includes 900 glomeruli with 25 model mitral cells assigned to each glumerulus.

Odor concentrations are defined by specifying a fraction f of glomeruli that respond to an odor at

that concentration. Then, for every odor, each glomerulus is assigned a reference onset latency

between 0 to 200 ms. At our reference concentration 10% of the glomeruli have onset latencies

<200 ms. The actual glomerular onset latencies for a given concentration are then obtained by

dividing the set of reference latencies by f. Glomeruli with latencies longer than the duration of

the inhalation are not activated. Mitral cell spiking is modeled as a Poisson process that generates

action potentials at specified rates; baseline spike rate is 1-2 Hz, which steps to 100 Hz when a

glomerulus is activated and then decays back to baseline with a time constant of 50 ms. This Pois-

son-generated mitral cell spiking introduces stochasticity into our olfactory bulb model.

Model piriform architecture and connectivity

The piriform model includes three types of model cells: 10,000 excitatory pyramidal cells, 1,225

feedforward inhibitory cells (FFI), and 1,225 feedback inhibitory cells (FBI). The model pyramidal

cells and FBIs are assigned to locations on a two-layer grid. Pyramidal cells and FBIs are uni-

formly spread over the grid on their respective layers. Each pyramidal cell receives an input from

1000 random other pyramidal cells and from 50 FFIs, both independent of location. Each pyram-

idal cell receives local input from the closest (on average) 12 FBIs. Each FBI receives input from

1,000 randomly chosen pyramidal cells and 8 (on average) closest FBIs. Each FFI receives input

from 50 other randomly chosen FFIs. Each mitral cell sends input to 25 random cells (either py-

ramidal cells or FFIs) in the pirifom. As a result, each pyramidal and FFI cell receives input from

approximately 50 random mitral cells.

<u>Piriform Dynamics</u>

The piriform model cells are modeled as leaky integrate-and-fire neurons with the membrane po-

tential V_i of model piriform cell following the dynamical equation

$$\tau_m \frac{dV_i}{dt} = (V_r - V_i) + I_{i,ex} - I_{i,in} + I_{i,ext}$$

Here $\tau_m = 15\text{ms}$ is the membrane time constant, $V_r = -65\text{mV}$ is the resting potential and $I_{i,ex}$, $I_{i,in}$

and $I_{i,ext}$ are the excitatory, inhibitory and external synaptic currents (of cell) respectively. We

have absorbed a factor of the membrane resistance into the definition of the input currents so they

are measured in the same units as the membrane potential (mV). When the membrane potential

V reaches the firing threshold $V_{th} = -50\text{mV}$ the neuron fires an action potential and the membrane

potential is reset to the resting value V_r, where it remains for a refractory period $\tau_{ref} = 1\text{ms}$.

The minimum allowed value for the membrane potential is $V_{min} = -75\text{mV}$. From this value the

membrane potential can only rise, following the dynamics, it cannot be decreased. Each action

potential fired by a neuron induces an instantaneous jump in the current of all its postsynaptic

targets by an amount proportional to the appropriate synaptic weight. Action potentials in FFIs

and FBIs affect the inhibitory current of their postsynaptic target neurons while action potentials

in the pyrmidal cells affect the excitatory current of their postsynaptic targets. Mitral cell spikes

also contribute to the excitatory current of the appropriate postsynaptic cells.

The inhibitory and excitatory currents dynamics are described by

$$\tau_{ex}\frac{dI_{i,ex}}{dt} = -I_{i,ex} + \tau_{ex}\sum_{j_{Ex}\to i} S_{ij,ex}\sum_{k}^{t_k^j<t}\delta(t_k^j - t) + \tau_{ex}\sum_{j_{Mitral}\to i} S_{ij,Mitral}\sum_{k}^{t_k^j<t}\delta(t_k^j - t)$$

$$\tau_{in}\frac{dI_{i,in}}{dt} = -I_{i,in} + \tau_{in}\sum_{j_{Ff;Fb}\to i} S_{ij,in}\sum_{k}^{t_k^j<t}\delta(t_k^j - t)$$

The synaptic time constants are given by $\tau_{ex} = 20\text{ms}$ and $\tau_{in} = 10\text{ms}$. The excitatory current com-

bines two components, AMPA and NMDA, into a single current. Because the NMDA synapses

are relatively slow and AMPA relatively fast, we choose the time constant of this composite cur-

rent in an intermediate range between these two extremes. In each term that includes summation

over contributions to the current, the first sum runs over all neurons that are presynaptic to neuron

and that contributes to the specified current. The second sum runs over all the times t_k^j of spikes

produced by presynaptic neuron prior to time , indexed by . The synaptic weight between presyn-

aptic Ex cell and postsynaptic Ex or Fb cell is given by $S_{ij}^{Ex\to Ex} = 0.25\text{mV}$ and $S_{ij}^{Ex\to Fb} = 1\text{mV}$ re-

spectively. The synaptic weight between presynaptic Mitral cell and postsynaptic pyramidal or

FFI is given by $S_{ij}^{Mitral\to Ex,Ff} = 10\text{mV}$. The synaptic weight between presynaptic FFI and postsynaptic

pyramidal or FFI cell is given by $S_{ij}^{Ff\to Ex,Ff} = 10\text{mV}$. The synaptic weight between presynaptic FBI

and postsynaptic pyramidal or FBI has the same strength, $S_{ij}^{Fb\to Ex,Fb} = 10\text{mV}$.

It is useful to describe the strengths of the different synapses we have discussed in terms of unitary

PSP sizes rather than using the S parameters given above. Using the constants we have given and

the synaptic and membrane dynamics, we calculated peak EPSPs and IPSPs. All inhibitory inputs

create IPSPs at the postsynaptic cell with a peak deflection of . EPSPs from mitral cell produce .

The EPSP for input from pyramidal to pyramidal or to FBI cell has or respectively.

The external input is constant in time but depend on cell identity. For all pyramidal cells it is given

by

$I_{i,ext} = 0.5 + I_{rand}(i)$

where $I_{rand}(i)$ is a random number generated from a Gaussian distribution with zero mean and

standard deviation . The external input for all other cells is zero.

Pyramidal cell population activity vectors

To analyze cortical responses, we define an activity vector . Each component of this vector, , is

the number of spikes generated by pyramidal neuron . Because we are interested in odor responses,

we start to count spikes at the beginning of the inhalation. The spikes count continues across the

full inhale, or stops after 50 ms in cases when we are interested in the initial response only. The

activity maps in the figures are a visual representation of the activity vectors created by reshaping

the vectors and assigning a color on the basis of their component values.

Activity correlation coefficients

To compute the correlation between two cortical responses on two different trials, we calculate the

correlation coefficient between the corresponding activity vectors, $\bar{r}_1$, $\bar{r}_2$,

$$c = \frac{\sum_{i=1}^{N}(r_{1,i}-<\bar{r}_1>)(r_{2,i}-<\bar{r}_2>)}{\sqrt{\sum_{i=1}^{N}(r_{1,i}-<\bar{r}_1>)^2}\sqrt{\sum_{i=1}^{N}(r_{2,i}-<\bar{r}_2>)^2}}$$

with $N = 10,000$ the number of pyramidal cells and $<\bar{r}>$ the average value of the entries of $\bar{r}$.

The perceptron

We use a perceptron, defined by a weights vector $\bar{w}$, to classify odor responses to odors on the

basis of the activity vectors explained above. Our goal is to train the perceptron such that trials

involving a chosen target odor are distinguished from trials using other odors. Because we gener-

ate odors randomly and all model mitral cells behave similarly, the results are independent of the

choice of target odor. Distinguishing the activity for a target odor from all other activity patterns

means that we wish to find $\bar{w}$ such that trials with a target odor have $\bar{w} \cdot \bar{r} > 0$ and trials with other

odors have $\bar{w} \cdot \bar{r} < 0$. Such a $\bar{w}$ only exists if trials using the target odor are linearly separable

from trials using other odors. If such a perceptron weight vector exists, we know that pyramidal

cell activity in response to a specific odor are inherently distinguishable from activity for other

odors. Furthermore, because the activity vectors we use do not contain temporal information, we

know that the discrimination relies solely on the identity of the active pyramidal cells, these are

the pyramidal cell ensembles, and their firing rates.

During training, 100 odors were presented at a specific concentration (10% activated glomeruli)

over a total of 600 trials. Odor 1 was chosen as the target, and the trials alternated between this

target odor and all odors. Thus, odor 1 was presented 303 times and every other odor 3 time. On

every trial, the quantity $\bar{w} \cdot \bar{r}$ was calculated, with $\bar{r}$ the activity vector for that trial and $\bar{w}$ the

current perceptron weight vector. Initially, $\bar{w}$ was zero. If classification was correct, meaning $\bar{w} \cdot$

$\bar{r} > 0$ for the target odor trial or $\bar{w} \cdot \bar{r} < 0$ for other odors, $\bar{w}$ was left unchanged. Otherwise $\bar{w}$

was updated to $\bar{w} + \bar{r}$ or $\bar{w} - \bar{r}$ for trials of odor 1 or for other odors, respectively. The entire

training procedure was repeated twice, once with activity vectors that included spikes counts

around the peak of the piriform activity (the first $50ms$ of inhale) and once using spikes counts

from the entire inhalation.

To test the perceptron, each odor was presented at many concentrations (even though training was

done only for a single fixed concentration). For the target odor 100 trials were tested at each

concentration (30 different concentrations ranging between 3% activated glomeruli and 30% acti-

vated glomeruli). Each trial that gave $\bar{w} \cdot \bar{r} > 0$ was considered a correct classification. For each

concentration, the percentage of trials that were correctly classified was calculated. Trials with

non-target odors were tested as well, one trial for each odor at each concentration. All the non-

target odors were correctly classified as not target ($\bar{w} \cdot \bar{r} < 0$) across all concentrations. The testing

procedure was done using both the peak and full activity vectors, with the corresponding percep-

tron weight vectors.

REFERENCES

Abraham, N. M., H. Spors, A. Carleton, T. W. Margrie, T. Kuner and A. T. Schaefer (2004).

"Maintaining accuracy at the expense of speed: stimulus similarity defines odor discrimination

time in mice." Neuron **44**(5): 865-876.

Apicella, A., Q. Yuan, M. Scanziani and J. S. Isaacson (2010). "Pyramidal Cells in Piriform Cortex

Receive Convergent Input from Distinct Olfactory Bulb Glomeruli." Journal of Neuroscience

30(42): 14255-14260.

Banerjee, A., F. Marbach, F. Anselmi, M. S. Koh, M. B. Davis, P. Garcia da Silva, K. Delevich,

H. K. Oyibo, P. Gupta, B. Li and D. F. Albeanu (2015). "An Interglomerular Circuit Gates

Glomerular Output and Implements Gain Control in the Mouse Olfactory Bulb." Neuron **87**(1):

193-207.

Bathellier, B., D. L. Buhl, R. Accolla and A. Carleton (2008). "Dynamic ensemble odor coding in

the mammalian olfactory bulb: sensory information at different timescales." Neuron **57**(4): 586-

598.

Bozza, T., J. P. McGann, P. Mombaerts and M. Wachowiak (2004). "In vivo imaging of neuronal

activity by targeted expression of a genetically encoded probe in the mouse." Neuron **42**(1): 9-21.

Cang, J. and J. S. Isaacson (2003). "In vivo whole-cell recording of odor-evoked synaptic

transmission in the rat olfactory bulb." J Neurosci **23**(10): 4108-4116.

Carandini, M. and D. J. Heeger (2012). "Normalization as a canonical neural computation." Nat

Rev Neurosci **13**(1): 51-62.

Carey, R. M., J. V. Verhagen, D. W. Wesson, N. Pirez and M. Wachowiak (2009). "Temporal

structure of receptor neuron input to the olfactory bulb imaged in behaving rats." J Neurophysiol

101(2): 1073-1088.

Christie, J. M., C. Bark, S. G. Hormuzdi, I. Helbig, H. Monyer and G. L. Westbrook (2005).

"Connexin36 mediates spike synchrony in olfactory bulb glomeruli." Neuron **46**(5): 761-772.

Cleland, T. A., S. Y. Chen, K. W. Hozer, H. N. Ukatu, K. J. Wong and F. Zheng (2011).

"Sequential mechanisms underlying concentration invariance in biological olfaction." Front

Neuroeng **4**: 21.

Cury, K. M. and N. Uchida (2010). "Robust odor coding via inhalation-coupled transient activity

in the mammalian olfactory bulb." Neuron **68**(3): 570-585.

Davison, I. G. and M. D. Ehlers (2011). "Neural Circuit Mechanisms for Pattern Detection and

Feature Combination in Olfactory Cortex." Neuron **70**(1): 82-94.

Economo, M. N., K. R. Hansen and M. Wachowiak (2016). "Control of Mitral/Tufted Cell Output

by Selective Inhibition among Olfactory Bulb Glomeruli." Neuron **91**(2): 397-411.

Franks, K. M. and J. S. Isaacson (2006). "Strong single-fiber sensory inputs to olfactory cortex:

implications for olfactory coding." Neuron **49**(3): 357-363.

Franks, K. M., M. J. Russo, D. L. Sosulski, A. A. Mulligan, S. A. Siegelbaum and R. Axel (2011).

"Recurrent circuitry dynamically shapes the activation of piriform cortex." Neuron **72**(1): 49-56.

Fukunaga, I., M. Berning, M. Kollo, A. Schmaltz and A. T. Schaefer (2012). "Two distinct

channels of olfactory bulb output." Neuron **75**(2): 320-329.

Ghosh, S., S. D. Larson, H. Hefzi, Z. Marnoy, T. Cutforth, K. Dokka and K. K. Baldwin (2011).

"Sensory maps in the olfactory cortex defined by long-range viral tracing of single neurons."

Nature **472**(7342): 217-220.

Gschwend, O., J. Beroud and A. Carleton (2012). "Encoding odorant identity by spiking packets

of rate-invariant neurons in awake mice." PLoS One **7**(1): e30155.

Haberly, L. B. (2001). "Parallel-distributed processing in olfactory cortex: new insights from

morphological and physiological analysis of neuronal circuitry." Chem Senses **26**(5): 551-576.

Haddad, R., A. Lanjuin, L. Madisen, H. Zeng, V. N. Murthy and N. Uchida (2013). "Olfactory

cortical neurons read out a relative time code in the olfactory bulb." Nat Neurosci **16**(7): 949-957.

Homma, R., L. B. Cohen, E. K. Kosmidis and S. L. Youngentob (2009). "Perceptual stability

during dramatic changes in olfactory bulb activation maps and dramatic declines in activation

amplitudes." Eur J Neurosci **29**(5): 1027-1034.

Hopfield, J. J. (1995). "Pattern recognition computation using action potential timing for stimulus

representation." Nature **376**(6535): 33-36.

Illig, K. R. and L. B. Haberly (2003). "Odor-evoked activity is spatially distributed in piriform

cortex." J Comp Neurol **457**(4): 361-373.

Johnson, D. M. G., K. R. Illig, M. Behan and L. B. Haberly (2000). "New features of connectivity

in piriform cortex visualized by intracellular injection of pyramidal cells suggest that "primary"

olfactory cortex functions like "association" cortex in other sensory systems." Journal of

Neuroscience **20**(18): 6974-6982.

Junek, S., E. Kludt, F. Wolf and D. Schild (2010). "Olfactory coding with patterns of response

latencies." Neuron **67**(5): 872-884.

Krone, D., M. Mannel, E. Pauli and T. Hummel (2001). "Qualitative and quantitative olfactometric

evaluation of different concentrations of ethanol peppermint oil solutions." Phytother Res **15**(2):

135-138.

Laing, D. G., P. K. Legha, A. L. Jinks and I. Hutchinson (2003). "Relationship between molecular

structure, concentration and odor qualities of oxygenated aliphatic molecules." Chem Senses

28(1): 57-69.

Large, A. M., N. W. Vogler, S. Mielo and A. M. Oswald (2016). "Balanced feedforward inhibition

and dominant recurrent inhibition in olfactory cortex." Proc Natl Acad Sci U S A **113**(8): 2276-

2281.

Luna, V. M. and N. E. Schoppa (2008). "GABAergic circuits control input-spike coupling in the

piriform cortex." Journal of Neuroscience **28**(35): 8851-8859.

Margrie, T. W. and A. T. Schaefer (2003). "Theta oscillation coupled spike latencies yield

computational vigour in a mammalian sensory system." J Physiol **546**(Pt 2): 363-374.

Meister, M. and T. Bonhoeffer (2001). "Tuning and topography in an odor map on the rat olfactory

bulb." J Neurosci **21**(4): 1351-1360.

Miura, K., Z. F. Mainen and N. Uchida (2012). "Odor representations in olfactory cortex:

distributed rate coding and decorrelated population activity." Neuron **74**(6): 1087-1098.

Miyamichi, K., F. Amat, F. Moussavi, C. Wang, I. Wickersham, N. R. Wall, H. Taniguchi, B.

Tasic, Z. J. Huang, Z. G. He, E. M. Callaway, M. A. Horowitz and L. Q. Luo (2011). "Cortical

representations of olfactory input by trans-synaptic tracing." Nature **472**(7342): 191-196.

Poo, C. and J. S. Isaacson (2009). "Odor Representations in Olfactory Cortex: "Sparse" Coding,

Global Inhibition, and Oscillations." Neuron **62**(6): 850-861.

Poo, C. and J. S. Isaacson (2011). "A major role ofor intracortical circuits in the strength and tuning

of odor-evoked excitation in olfactory cortex." Neuron **72**(1): 41-48.

Rennaker, R. L., C. F. Chen, A. M. Ruyle, A. M. Sloan and D. A. Wilson (2007). "Spatial and

temporal distribution of odorant-evoked activity in the piriform cortex." J Neurosci **27**(7): 1534-

1542.

Rinberg, D., A. Koulakov and A. Gelperin (2006). "Speed-accuracy tradeoff in olfaction." Neuron

51(3): 351-358.

Roland, B., R. Jordan, D. L. Sosulski, A. Diodato, I. Fukunaga, I. Wickersham, K. M. Franks, A.

T. Schaefer and A. Fleischmann (2016). "Massive normalization of olfactory bulb output in mice

with a 'monoclonal nose'." Elife **5**.

Rubin, B. D. and L. C. Katz (1999). "Optical imaging of odorant representations in the mammalian

olfactory bulb." Neuron **23**(3): 499-511.

Sanders, H., B. E. Kolterman, R. Shusterman, D. Rinberg, A. Koulakov and J. Lisman (2014). "A

network that performs brute-force conversion of a temporal sequence to a spatial pattern: relevance

to odor recognition." Front Comput Neurosci **8**: 108.

Schaefer, A. T. and T. W. Margrie (2007). "Spatiotemporal representations in the olfactory

system." Trends Neurosci **30**(3): 92-100.

Schaefer, A. T. and T. W. Margrie (2012). "Psychophysical properties of odor processing can be

quantitatively described by relative action potential latency patterns in mitral and tufted cells."

Front Syst Neurosci **6**: 30.

Schoppa, N. E. (2006). "Synchronization of olfactory bulb mitral cells by precisely timed

inhibitory inputs." Neuron **49**(2): 271-283.

Shusterman, R., M. C. Smear, A. A. Koulakov and D. Rinberg (2011). "Precise olfactory responses

tile the sniff cycle." Nat Neurosci **14**(8): 1039-1044.

Sirotin, Y. B., R. Shusterman and D. Rinberg (2015). "Neural Coding of Perceived Odor Intensity."

eNeuro **2**(6).

Sosulski, D. L., M. L. Bloom, T. Cutforth, R. Axel and S. R. Datta (2011). "Distinct representations

of olfactory information in different cortical centres." Nature **472**(7342): 213-216.

Spors, H. and A. Grinvald (2002). "Spatio-temporal dynamics of odor representations in the

mammalian olfactory bulb." Neuron **34**(2): 301-315.

Spors, H., M. Wachowiak, L. B. Cohen and R. W. Friedrich (2006). "Temporal dynamics and

latency patterns of receptor neuron input to the olfactory bulb." J Neurosci **26**(4): 1247-1259.

Stettler, D. D. and R. Axel (2009). "Representations of Odor in the Piriform Cortex." Neuron

63(6): 854-864.

Stokes, C. C. A. and J. S. Isaacson (2010). "From Dendrite to Soma: Dynamic Routing of

Inhibition by Complementary Interneuron Microcircuits in Olfactory Cortex." Neuron **67**(3): 452-

465.

Suzuki, N. and J. M. Bekkers (2006). "Neural coding by two classes of principal cells in the mouse

piriform cortex." J Neurosci **26**(46): 11938-11947.

Suzuki, N. and J. M. Bekkers (2012). "Microcircuits mediating feedforward and feedback synaptic

inhibition in the piriform cortex." J Neurosci **32**(3): 919-931.

Uchida, N. and Z. F. Mainen (2003). "Speed and accuracy of olfactory discrimination in the rat."

Nat Neurosci **6**(11): 1224-1229.

Uchida, N., C. Poo and R. Haddad (2014). "Coding and transformations in the olfactory system."

Annu Rev Neurosci **37**: 363-385.

Wesson, D. W., R. M. Carey, J. V. Verhagen and M. Wachowiak (2008). "Rapid encoding and

perception of novel odors in the rat." PLoS Biol **6**(4): e82.

Wilson, C., G. Serrano, A. Koulakov and D. Rinberg (2015). Primacy coding of odor identity

across concentrations. Society for Neuroscience. **229.01**.

Wilson, D. A. and R. M. Sullivan (2011). "Cortical processing of odor objects." Neuron **72**(4):

506-519.

Wilson, R. I. and Z. F. Mainen (2006). "Early events in olfactory processing." Annu Rev Neurosci

29: 163-201.

Zhan, C. and M. Luo (2010). "Diverse patterns of odor representation by neurons in the anterior

piriform cortex of awake mice." J Neurosci **30**(49): 16662-16672.

FIGURE LEGENDS

Figure 1. Odors activate distributed ensembles of PCx neurons.

(A) Raster plots showing spiking of 1,000 mitral cells (out of 22,500 total; 25 mitral cells/glomer-

ulus, 40 glomeruli shown) in response to 3 different odors. Each row represents a single mitral cell

and all mitral cells belonging to each glomerulus are clustered. Tick marks indicate spike times.

Inhalation begins at 0 ms and is indicated by the grey shaded region. The red curve shows the

cumulative number of glomeruli activated across the sniff, and the blue curve is the firing rate

averaged across all mitral cells.

(B) Schematic of the piriform cortex model.

(C) Voltage traces for three sequential sniffs in 4 model pyramidal cells. Time of inhalation is

indicated by the dashed line.

(D) Single-trial population activity map for all 10,000 pyramidal cells. Each pixel represents a

single cell, and pixel color indicates the number of spikes fired during the 200 ms inhalation. Ap-

proximately 13% of cells fired at least 1 action potential, with activated cells randomly distributed

across the cortex.

(E) Response vectors shown for 20 cells in response to different odors presented on 4 sequential

trials. Spiking levels are low for no-odor controls. Note the trial-to-trial variability and that indi-

vidual cells can be activated by different odors.

Supplementary Figure 1. Modeling odor responses in olfactory bulb

(A) Raster for all 22,500 mitral cells for one odor trial. Grey shading indicates inhalation phase.

Inset shows expanded view of 1,000 mitral cells belonging to 40 different glomeruli.

(B) Raster showing trial-to-trial variability for a single activated glomerulus. Each box represents

a different sister mitral cell, with trials 1-4 represented by the lines within each box.

Figure 2. Evolution of a cortical odor response.

(A) Raster for a single sniff showing spiking activity of a subset of mitral cells (2,250 out of

22,500), all 1,225 feedforward interneurons (FFIs), all 10,000 pyramidal cells, and all 1,225 feed-

forward interneurons (FBIs). Spiking rate for the population of pyramidal cells is shown at the

bottom (average of 6 trials). Note that the earliest activated glomeruli initiate a cascade of pyram-

idal cell spiking that peaks after ~50 ms and is abruptly truncated by synchronous spiking of FBIs.

(B) Single-trial voltage traces (black) for 3 pyramidal cells in response to the same odor. Inhalation

onset is indicated by the dashed line. The red traces show the bulb input and the green traces the

recurrent input received by each cell. Cell 1 receives strong bulb input and spikes soon after odor

presentation. Cell 2 receives subthreshold input from the bulb and only spikes after receiving

addition recurrent input from other pyramidal cells. Cell 3 receives no early odor-evoked input

from the bulb, and its recurrent input is subthreshold, so it does not spike over the time period

shown.

Figure 3. Distinct roles for different inputs in shaping ensemble response.

(A-D) Raster plots for presentations of the same odor using different circuit configurations. The

percentages of active pyramidal cells are shown in the insets.

(A) As in Figure 2A.

(B) Network activity when pyramidal cells get excitatory input from mitral cells but without FFI,

recurrent excitation or FBI. Pyramidal cell spiking tracks mitral cell input. Population rate for the

full network is shown in grey for comparison.

(C) Network activity after eliminating excitatory input to FFIs. Note that sustained FBI spiking

increases without FFI, resulting in roughly constant levels of sustained spiking in pyramidal cells.

(D) Network activity with no recurrent excitation onto either other pyramidal cells or FBIs. The

transient activity peak seen in A-D is absent, and sustained activity is slightly higher.

(E) Population rate plots for 3 different odors with the full network (black trace), no FFI (red trace)

or no recurrent excitation/FBI (green trace). Insets expand the period around inhalation. Note the

population spiking is higher and peaks slightly earlier without FFI, and that responses are slow

without recurrent excitation. All population spike rate plots are averaged over 6 trials.

Figure 4. Earliest-active glomeruli define the cortical response.

(A) Normalized population spike rates (black) in response to an odor during the sniff cycle shown

in grey. The red curve shows the cumulative number of glomeruli activated across the sniff. Note

that population spiking peaks after only a small subset of glomeruli have been activated.

(B) Normalized population spike rates for one odor for the full network (black trace), without FFI

(red trace) and without recurrent excitation (green trace). Grey trace shows the cumulative number

of activated glomeruli.

(C) Fraction of peak population spike rate as a function of the cumulative number of activated

glomeruli for 6 different odors. These curves indicate the central role recurrent excitation plays in

amplifying the impact of early-responsive glomeruli.

(D) Average correlation coefficients for repeated same-odor trials and pairs of different-odor trials

measured over the full 200 ms inhalation.

(E) As in D but measured over the first 50 ms after inhalation onset.

(F) Ratios of correlations for same- vs. different-odor trials measured over the full sniff (grey bar

on left) and over the first 50 ms (black bar on right).

Figure 5. Cortical output is normalized across concentrations.

(A) Mitral cell rasters for 2 odors at 3 different concentrations defined by the fraction of co-active

glomeruli. Glomerular onset latencies decrease with concentration allowing more glomeruli to be-

come active over the sniff period. Odors are different from the odors in Figure 1.

(B) Single-trial piriform response vectors for single trials over a concentration range corresponding

to 3-30% active glomeruli. Note that activity does not dramatically increase despite the 10-fold

increase in input.

(C) Fraction of activated pyramidal cells at different odor concentrations for the full network (black

trace), without FFI (red trace) and without recurrent excitation (green trace) for 4 different odors

(open circles, thin lines) and averaged across odors (filled circles, thicker lines). Note that elimi-

nating FFI primarily shifts the number of responsive cells, indicating that FFI is largely subtractive,

whereas eliminating recurrent excitation alters the gain of the response. Note also that recurrent

excitation amplifies the number of activated cells at low odor concentrations.

(D) As in C but for the total number of spikes across the population.

(E) Perceptron classifications of an odor at different concentrations on the basis of pyramidal cell

activity. Either the transient cortical activity (first 50 ms of the inhalation; black curve) or the

activity across the full inhalation (gray curve) was used for both training and testing. Training was

solely on the basis of the concentration indicated by the black arrow. The dashed line shows the

chance level of classification (50%).

Figure 6. Effect of odor concentration on response timing

(A). Normalized population spike rates for 3 odors at 3 concentrations denoted by percentage of

glomeruli responding. Peak responses are higher and occur earlier at higher concentrations.

Dashed lines indicate inhalation onset.

Average latencies to peak (B) and peak rate (C) vs. number of activated glomeruli.