Bailey SF, Guo Q and Bataillon T. 2018. **Identifying drivers of parallel evolution: A regression model approach**. *bioRxiv*, 118695, https://doi.org/10.1101/118695

Title: Identifying drivers of parallel evolution: A regression model approach

Authors: Susan F. Bailey[1,2,*], Qianyun Guo[1], and Thomas Bataillon[1]

Author affiliations:

[1] Bioinformatics Research Centre, Aarhus University, C.F. Møllers Allé 8, DK-8000 Aarhus C,

Denmark.

[2] Current affiliation: Department of Biology, Clarkson University, PO Box 5805, Potsdam, NY 13699-

5805

*Author for Correspondence: Susan F. Bailey, Department of Biology, Clarkson University, PO Box

5805, Potsdam, NY 13699-5805, phone: 315-268-4263, email: sbailey@clarkson.edu

Running head: Identifying drivers of parallel evolution

Keywords: parallel evolution, experimental evolution, Poisson regression, negative binomial

regression

Data archival location: Dryad, doi to be included later

Abstract

This preprint has been reviewed and recommended by Peer Community In Evolutionary Biology

(http://dx.doi.org/10.24072/pci.evolbiol.100045). Parallel evolution, defined as identical changes

arising in independent populations, is often attributed to similar selective pressures favoring the

fixation of identical genetic changes. However, some level of parallel evolution is also expected if

mutation rates are heterogeneous across regions of the genome. Theory suggests that mutation and

selection can have equal impacts on patterns of parallel evolution, however empirical studies have yet

to jointly quantify the importance of these two processes. Here, we introduce several statistical models

to examine the contributions of mutation and selection heterogeneity to shaping parallel evolutionary

changes at the gene-level. Using this framework we analyze published data from forty experimentally

evolved *Saccharomyces cerevisiae* populations. We can partition the effects of a number of genomic

variables into those affecting patterns of parallel evolution via effects on the rate of arising mutations,

and those affecting the retention versus loss of the arising mutations (i.e. selection). Our results suggest

that gene-to-gene heterogeneity in both mutation and selection, associated with gene length,

recombination rate, and number of protein domains drive parallel evolution at both synonymous and

nonsynonymous sites. While there are still a number of parallel changes that are not well described, we

show that allowing for heterogeneous rates of mutation and selection can provide improved predictions

of the prevalence and degree of parallel evolution.

Introduction

Documenting patterns of parallel evolution during the adaptive divergence of populations or during repeated bouts of adaptation in populations maintained in the lab is becoming increasingly feasible. Beyond the fascination for the pattern of repeatable evolution, an outstanding open question is to understand which underlying processes are driving the pattern of molecular evolution during adaptation. Theory makes clear cut predictions: in the absence of selective interference between beneficial mutations (the so called strong selection weak mutation, or SSWM, domain), heterogeneity in mutation rates and selection coefficients between loci are expected to have equal influence on patterns of parallel evolution (Chevin et al., 2010; Lenormand et al., 2016). So far very few empirical studies have attempted to jointly quantify the relative importance of these two processes in shaping patterns of parallel evolution in genetic data. One study has explored this indirectly by quantifying the contribution of these two processes in shaping the parallel evolution of heritable traits that are assumed to be associated with parallel genetic changes (Streisfeld and Rausher, 2011). Recent work by Bailey et al., 2017 outlines an approach for quantifying the effects of mutation and selection heterogeneity in driving parallel evolution in experimental evolution data, but this alternate approach can not identify potential genomic drivers of that heterogeneity, as we do here. Other previous studies looking explicitly at parallel genetic changes have focused on the impacts of either selection or mutation separately.

Parallel evolution is an identical change in independently evolving lineages, and the similar process, convergent evolution, occurs when different ancestral states change to the same descendant state in independently evolving lineages (Zhang and Kumar, 1997). These kinds of evolutionary changes are studied across many different levels of biological organization from nucleotides to genes to pathways and more. In this study, we focus on parallel evolution at the level of the gene.

Parallel, along with convergent, evolution has previously been considered strong evidence of

selection. A number of studies have examined gene-level mutation counts, looking for levels of parallel evolution that exceed what one would expect in the absence of selection (Caballero et al., 2015; Marvig et al., 2015; Woods et al., 2006), according to some null model, with an aim to identify genes that are under selection. For example, (Caballero et al., 2015) calculated the probability of instances of gene-level parallel evolution in whole genome sequences of *Pseudomonas aeruginosa* repeatedly sampled over the course of a year from the sputum of a cystic fibrosis patient assuming uniform re-sampling of ~150 mutation events across the approximately 6000 genes in the genome. The authors were able to identify 19 different genes for which there was significant deviation from their null model, and that pattern was interpreted as evidence for selection acting on these genes. However this study and other similar approaches do not account for the possibility of heterogeneity in mutation rate from gene-to-gene, a process that can generate false positives when using "abnormal" levels of parallel evolution as a means to detect selected genes.

Others have compared instances of parallel and convergent evolution across species (see Christin et al., 2010 for a review and examples). These studies also aim to identify genes under selection by searching for genes that exhibit a higher than expected number of instances of parallel evolution according to a specified null model for evolution. Many cross-species comparative studies report instances of parallel molecular evolution and readily interpret these as being driven by positive selection (e.g. Castoe et al., 2009; Feldman et al., 2012; Jost et al., 2008; Liu et al., 2014). However Zou and Zhang, 2015 show that in this type of analysis the choice of null model is crucial and suggest that many previously reported instances of parallel evolution driven by selection could in fact have resulted simply from mutation biases and mutational heterogeneity in the absence of selection.

In contrast to studies aimed at identifying selection, other work has focused on examining how heterogeneity in mutation rate can effect the distribution of mutations across a genome, and so the probability of parallel evolution. These studies focus exclusively on either those mutations that are

assumed to be to a first approximation neutral (e.g. synonymous mutations, Maddamsetti et al., 2015)

or mutations arising in the course of experiments where selection is minimal (e.g. mutations arising in a

mutation accumulation experiment, Ness et al., 2015). On the whole, these studies suggest substantial

gene-to-gene heterogeneity in mutation rate and this can arguably also generate differences in the

distribution of mutations across the genome (although studies differ in the factors identified that drive

that heterogeneity). However, it is not clear what the relative contribution of mutation rate

heterogeneity is when the mutations of interest also have the potential to be under varying degrees of

selection.

In this study we aim to explore the effects of both mutation and selection in generating the

mutations that are observed across the genome. By identifying and quantifying the processes that give

rise to mutations and how those vary from gene-to-gene, we can then begin to understand and predict

patterns of parallel evolution at the gene-level. To do this we propose a framework that explicitly

considers drivers of both selection and mutational heterogeneity. Using both Poisson and negative

binomial regression models, we analyze gene-level mutation count data obtained from whole genome

sequencing of a large set of yeast (*Saccharomyces cerevisiae*) experimental populations that were

adapted in parallel to identical environmental conditions in the lab (Lang et al., 2013). We find that the

best predictor of parallel mutations at the gene-level is simply the length of the gene, and along with

this, a few other genomic covariates – namely the number of protein domains and the rate of

recombination, and so it is variation in these variables that drives patterns of parallel evolution in this

system.

Models for identifying processes underlying parallel evolution

We are interested in quantifying heterogeneity in mutation rate and selection, and how these in turn are

driving patterns of parallel evolution, and identifying genomic variables that predict how the processes

of mutation and selection vary from gene-to-gene. To accomplish this, we need a framework that can

explicitly separate the effects of variation in mutation rate and variation in selection. We do this by

examining separately the observed synonymous and nonsynonymous mutations, making the

assumption (which we then explore) that gene-to-gene variation in the rate at which synonymous

mutations rise to observable frequencies is driven solely by variation in the mutation rate per gene,

while gene-to-gene variation in the rate at which nonsynonymous mutations have arisen may be driven

by heterogeneity in both mutation and selection processes. We describe the number of mutations

observed in gene i during the course of an experiment, as $X_i = X_i^S + X_i^N$, where X_i^S and X_i^N denote the

synonymous and nonsynonymous mutation counts respectively. We assume these mutations are Poisson

distributed with rates λ_i^S and λ_i^N respectively. For synonymous mutations, this Poisson rate can be

modeled as

$$\lambda_i^S = M_0 \, \mu_0 \, L_i \, \pi_0$$ Eqn (1)

Here, M_0 is a parameter that absorbs both time and population size at which the evolution occurred and

that is constant across the genome, μ_0 is the per-nucleotide mutation rate that we assume (and check) is

constant across the genome, L_i is the length of gene i in nucleotides, and π_0 is the probability of a

synonymous mutation rising to an observable frequency in the population (we assume that synonymous

mutations are selectively neutral and so this probability is assumed to be constant across the genome).

For nonsynonymous mutations,

$$\lambda_i^N = \lambda_i^S \, \pi_i \, ,$$ Eqn (2)

where π_i is a scalar that incorporates the effects of selection on the rate of fixation of non-synonymous

mutations arising in gene i. Specifically, π_i , is a function of the mean selection coefficient of gene i, s_i ,

and under strong-selection-weak-mutation (SSWM) conditions, $\pi_i \propto s_i$ (Gillespie, 1984). We assume

that the mean selection coefficient for non-synonymous mutations in a given gene can range from

deleterious, to neutral, to beneficial. The type of data used and the underlying assumptions are

summarized in Fig. 1.

Given these underlying assumptions about the processes giving rise to observable mutations in

the experimental sequence data, we can then use Poisson and negative binomial (NB) regression to

identify potential genomic variables that significantly explain variation in λ_i^N and λ_i^S, and thus

ultimately in the mutation and selection processes from gene-to-gene. The Poisson regression is used to

explore counts of rare events (i.e. the observed mutations) that have a fixed probability of being

observed, while for a NB regression, the rate of those rare events is itself a random variable that is

gamma-distributed. A NB regression incorporates an extra parameter beyond a Poisson rate, known as

the dispersion parameter (here denoted by θ), reflecting the amount of underlying variation in the rate

of observed mutations from gene-to-gene and governs the "extra" variance of the NB distribution

relative to a Poisson distribution with identical mean. If there is no heterogeneity among the rate of

observed mutations from gene-to-gene, the dispersion parameter θ goes to zero and we recover a

Poisson regression model. Therefore, the Poisson regression model is a special case of the NB

regression model, as $NB(\lambda_i, \theta)$ reduces to $Poisson(\lambda_i)$ at the limit of $\theta \rightarrow 0$ (see for instance Zuur,

2009). As a consequence, the Poisson and NB models are "nested" and their relative fit can be

compared using a likelihood ratio test when exploring the fit of both types of regression models in this

study.

More precisely, we use the models $X_i \sim Poisson(\lambda_i)$ or $X_i \sim NB(\lambda_i, \theta)$, fitting the following

regression:

$$\log(\boldsymbol{\lambda}) = constant + \alpha_1 \mathbf{A_1} + \alpha_2 \mathbf{A_2} + \ldots + \alpha_j \mathbf{A_j}, \quad\quad\quad\quad \text{Eqn (3)}$$

where $\boldsymbol{\lambda} = (\lambda_1, \ldots, \lambda_i, \ldots, \lambda_n)$ are the Poisson rates for all n genes, $\mathbf{A_1} \ldots \mathbf{A_j}$ are the j potential genomic

explanatory variables, and $\alpha_1 \ldots \alpha_j$, the estimated regression coefficients for those j variables. Thus, in

the case of the synonymous mutations, $constant = \log(M_0 \pi_0 \mu_0)$, $\mathbf{A_1} = \log(L_i)$ setting $\alpha_1 = 1$. For

nonsynonymous mutations, $\alpha_2 \mathbf{A_2} + \ldots + \alpha_j \mathbf{A_j} = \log(\pi_i)$. Details of the implementation of these models

is provided below.

Methods

The data

Evolution experiment data – We analyzed data obtained from whole genome re-sequencing of forty

populations of *S. cerevisiae* adapted in parallel in the lab (Lang et al., 2013). In this experiment, clonal

haploid yeast populations were grown in 128 µl of liquid YPD media and transferred every 12 to 24

hours to fresh media for approximately 1,000 generations (see Lang et al., 2011 for more details on the

experiment protocol). In our analysis we focus on all detected genic mutations (718 out of the total

1020 in the data set) from forty sequenced populations, i.e. all genic mutations that were able to escape

drift and so rise to frequencies of at least approximately 10% in the populations (mutations below this

frequency could not reliably be detected, see Lang et al., 2013). The mutations included in the data set

consisted of SNPs and small indels. Mutations were grouped by gene across all forty populations, and

categorized as synonymous (SYN) or nonsynonymous (NS), i.e. those that do not confer amino acid

changes, and those that do, respectively.

Comparative genomics data – We used a set of orthologuous gene alignments spanning four distinct

yeast species (*S. cerevisiae*, *S. paradoxus*, *S. bayanus*, and *S. mikatae*; available from

www.yeastgenome.org/download-data/genomics; Kellis et al., 2003; Cliften et al., 2003) to infer the

gene-to-gene heterogeneity of the substitution rates at synonymous sites and nonsynonymous sites,

hereafter *dS* and *dN* respectively. To do so, we first realigned the gene sequences using ClustalW

(Larkin et al., 2007) on the translated protein sequence data and then applied a number of filters to the

data with an aim at removing those gene alignments that might result in inaccurate codon substitution

model predictions. We removed alignments for those genes where sequences were not available from

all four species, alignments for which at least one sequence had <30% overlap with the one of the other

3 sequences, and alignments for which at least one sequence was <300 bps in length. We then used a

maximum likelihood codon based method (CodeML in the PAML software package; Yang, 2007) to

infer dS and dN, for each gene in our data set. We used a codon table model (i.e. seqtype = 1;

CodonFreq = 3) with a fixed tree topology (i.e. runmode = 0). A comparison of AICs among alternative

codon based models indicated this was the most appropriate model for the data set.

Additional genomic data – We included eight additional genomic variables in our analysis that we

expected could have the potential to effect the probability of a gene to harbor mutations. Our collection

of variables is not meant to be exhaustive, but simply meant to illustrate the potential for additional

genomic information to improve our predictions of which genes bear mutations across the genome. For

each gene we consider: gene length, % GC content, multi-functionality, degree of protein-protein

interaction (PPI), codon adaptation index (CAI), number of domains, level of expression (in the same

environment at the evolution experiment), local recombination rate, and essentiality of the gene. We

expect some of these variables may capture heterogeneity in the per-gene mutation rates, for example:

gene length, which likely captures variation in a gene's mutational target size, and local recombination

rate, which has been shown to be associated with mutability in yeast (Holbeck and Strathern, 1997;

Strathern et al., 1995). We expect other variables may capture heterogeneity in selection from gene-to-

gene, for example: multi-functionality and PPI, which may characterize aspects of how pleiotropic a

given gene is and so the level of evolutionary constraint it is under. We expect still other genomic

variables may capture heterogeneity in both mutation and selection. For example, level of expression of

a gene may be correlated with gene-to-gene variation in selection as highly expressed genes have been

shown to be more highly conserved, both specifically in yeast (Drummond et al., 2005; Pál et al., 2001)

and as a more general phenomenon across species (Drummond and Wilke, 2008). On the other hand,

level of expression of a gene has also been shown to be positively correlated with mutability (Ness et

al., 2015). Descriptions of the variables used in this study and sources from which the data were

obtained are provided in Table 1.

For the purposes of our analysis, we only used mutations in genes for which we were able to

obtain a full set of complementary genomic variables (393 of the total 718 genic mutations in Lang et

al., 2013). This final data set does not contain any examples of multiple mutations within the same gene

on the same genome and so we consider each mutation to be an independent mutational event. A data

set integrating the mutation counts originally made available by Lang et al., 2013 (from their

Supplementary Table 1) and all the genomic covariates that we aggregated for this study, as well as the

gene alignments used for estimating dN and dS are available on Dryad (doi will be inserted here).

Regression models

Regression models and explanatory variables tested – We used the Poisson and negative binomial

regression models described in the "Models" section above to examine how much of the variation in

our explanatory variables could account for patterns of variation in mutation counts per gene. We used

the 'glm' and 'glm.nb' functions in R (R Development Core Team, 2014) to implement these models. We

fit a series of models to synonymous and nonsynonymous mutation count data separately. To start, we

fit the synonymous mutations (model M_S), testing our assumptions that rate of observed mutations per

gene (totaled over all 40 populations in the data set) is proportional to number of nucleotide sites in the

gene (L_i), and the per nucleotide mutation rate does not vary significantly across the genome – i.e. a

model assuming μ_0 is a fixed parameter (Poisson regression) fits the data better than a model where μ_0

for each gene is drawn from a gamma distribution (NB regression). We also tested for significance of

each of the genomic variables included in this study by adding each of them to the Poisson model and

checking if the model fit is significantly improved.

After these assumptions were checked, we moved on to fit the nonsynonymous mutation data

(M_N), testing the 11 genomic variables listed in Table 1. We then examined an alternate model (M_N^{PC}),

fitting the nonsynonymous mutations using the principal components of the 11 genomic variables in

place of the raw variables. The reason we explore this model is that many genomic variables tend to be

correlated (for correlations between the particular variables used in this study, see supplementary Table

S1), and one approach to reducing potential problems with co-linearity is to transform the raw variables

into their principal components and use the resulting uncorrelated composite variables for the

regression analysis. We performed a principal component analysis on 11 genomic variables using the

'prcomp' function in R to obtain 11 principal components (PCs).

Model selection and significance of variables – For each variable and parameter of interest we tested

significance by comparing versions of the models with and without that variable or parameter of

interest through a likelihood-ratio test (LRT). Significance testing for LRTs was done using

permutation tests instead of relying on asymptotic distribution of the LRTs. Permutation was performed

by re-sampling the mutation data, re-fitting the models with and without the genomic variable or

parameter of interest and then calculating the LRT of those re-fitted models. This procedure allowed us

to approximate null distributions and obtaining P-values by calculating the frequency of permutations

that resulted in a likelihood-ratio greater than or equal to the true observed value. Variables found to

significantly improve model fit were retained in the final "best" model. We choose to test significance

using permutations given that asymptotic results on the distribution of the likelihood ratio test may

break down as the reduced model – the Poisson regression – lies at the boundaries of the parameter

space for θ, included in the NB regression (see for instance Self and Liang, 1987). In practice, 1000

permutations were used to approximate the null and obtain p-values on each variable (more

permutations might be required if needed to approximate p-values that are much smaller than 10^{-3}).

The two nonsynonymous mutation models M_N and M_N^{PC} were compared with each other using

Akaike information criterion (AIC; Akaike, 1973), and the proportion of variation explained (pseudo-

R^2) was estimated as the R^2 obtained from a linear regression (using 'lm' in R) between the observed

and predicted mutation counts for a given model. Note that this statistic is not used for any formal

goodness-of-fit but as an illustrative way to report how much of the whole variation is accounted for by

any model we fit to the mutation count data.

Predicting the degree of parallelism – We used the resulting best-fit parameters and distributions from

the regression models to simulate mutations for 40 populations and calculated the mean proportion of

shared genes bearing mutations for all pairwise combinations of those 40 populations using the Jaccard

Index, $\left(J_{G1,G2}=\dfrac{|G1\cap G2|}{|G1\cup G2|}\right)$. This measure of parallelism has been used in a number of previous

empirical comparisons (Bailey et al., 2017; Wong et al., 2012). We compared the simulated distribution

of J values to the distribution calculated from the real data set in order to assess the accuracy of the

regression models in predicting the degree of parallel evolution in this system. This can be used as a

predictive check for our model.

All statistical analyses were implemented in R (R Development Core Team, 2014) and an

example script for implementing our model framework and hypothesis testing is available on Dryad

(doi will be inserted here).

Results

The data

Mutation counts data – We used experimental data comprising all mutations detected at a frequency

over 10% in the forty evolved *S. cerevisiae* populations described in Lang et al., 2013. After removing

those genes for which we had incomplete or unreliable data (see Methods), we were left with 2891

genes out of a total of 6603. The filtered data set contained 357 nonsynonymous mutations distributed

across 267 genes, and 58 synonymous mutations distributed across 57 genes. The genes removed by

our filtering rules had disproportionately more mutations compared to those genes that were retained in

the data set (χ^2 = 50.57, df = 1, P < 0.001). This is not unexpected as highly divergent genes are more

likely to be filtered out due to alignment issues, and it is not surprising that highly divergent genes

would tend see more mutations than average, whether it be as a result of mutation and / or selection

mechanisms. This bias in the filtering means that our results are likely conservative in terms of

detecting significant relationships between long-term (from comparative genomics data) and short-term

(from experimental evolution data) measures of divergence.

Genomic variables – We used codon substitution models comparing four yeast species to estimate *dS*

and *dN/dS* for each gene. Estimates for *dS* ranged widely, from 0.21 to 68.7, however the vast majority

of *dS* estimates (~95%) were less than 4. Estimates for *dN/dS* ranged from 0.00010 to 0.43, and these

values are weakly negatively correlated with *dS* (r = -0.043, P = 0.021). We collated and/ or calculated

nine other genomic variables with the potential to effect the mutation and selection processes in this

system and estimated correlation coefficients between all pairs of explanatory variables used in this

study (Table S1). While the correlations between these variables tend to be quite weak, many are, in

fact, significant due to the large number of observations in the data set.

Mutation counts analysis

Synonymous mutations – We used regression models to test our assumption that gene-level mutation

rate can be adequately described as simply being directly proportional to gene length. Restricting the

data to the synonymous mutations, we compared Poisson regression models with and without gene

length included as an explanatory variable ($M_S0.P$: λ_S = *constant* and $M_S1.P$: λ_S = *constant**$(L_i)^{\alpha 1}$,

respectively), and a Poisson regression model where rate is restricted to be directly proportional to gene

length (i.e. $M_S2.P$: $\lambda_S = constant*L_i$). We also compared with negative binomial versions of these

model to look for the possibility of additional unexplained variation in the rate λ. A summary of the

results of these comparisons is shown in Table 2. Model $M_S2.P$ was the best model according to a

comparison of AICs. The fits of these models to the distribution of synonymous mutation counts per

gene are visualized in Fig. 2A. We also compared a series of Poisson models, each containing one of

the genomic variables included in this study (see supplementary information Table S2). Two variables

do significantly improve model fit – dN/dS and CAI, but very modestly, helping to explain only 0.24 %

and 0.11 % of the total variance respectively. Taken together, these tests suggest that the assumptions of

neutral selection at synonymous sites and a constant mutation rate across nucleotides are reasonable

simplifications for these data.

Nonsynonymous mutations – We fit regression models to the nonsynonymous mutation data, including

eleven genomic variables, trying to identify which of those variables could significantly explain

variation in the number of observed mutations per gene (totaled over all 40 populations in the data set).

We found that gene length (L), number of domains in the encoded protein (*num.dom*), and

recombination rate (r) were significant in our model (see model $M_N.NB$ in Table 3).

When we fit regression models using the principal components of the genomic variables in

place of the raw variables, we found that only a single principal component, PC10, was significant in

the model (see model $M_N.NB_{PC}$ in Table 3). PC10 is fairly evenly loaded with a number of genomic

variables (see Fig. 3), however the three significant genomic variables from $M_N.NB$ (L, *num.dom*, and

r) are among the variables more heavily loaded on PC10, so the two models seem to be roughly in

agreement. Further, these models both explain about 16% of the variance (as calculated from pseudo-r^2

estimates, see methods). A comparison of Poisson and negative binomial regression models, as well as

models including the raw genomic variables versus the transformed principal component variables,

suggests that the best model for these nonsynonymous mutation count data is a negative binomial

regression using the raw genomic variables (see AIC values in Table 4). The fits of these models to the

distribution of nonsynonymous mutation counts per gene are visualized in Fig. 2B.

Predicted parallelism – Using the predicted rates and distributions from the best-fit regression models

for both the synonymous and nonsynonymous mutations (M_S2 and $M_N.NB$, respectively), we simulated

mutations for a set of 40 populations and calculated the Jaccard index (J) as a measure of gene-level

parallelism between all pairs of those simulated populations. We performed the same calculations with

the 40 populations from the real data. Fig. 4 shows a comparison of those J values from the simulated

and real data. While the simulated data from our model does quite well at capturing the range of J

values, it does not do a great job of capturing the shape of the distribution.

<u>Discussion</u>

Here we present a modeling framework to infer what genomic variables may underlie gene to gene

variation in mutation rate and intensity of selection. We use these models to provide evidence that

parallel evolution at both nonsynonymous and synonymous sites is driven by non trivial amounts of

gene-to-gene heterogeneity in the mutation and selection processes. Using our modeling approach, we

identified a number of genomic variables that can significantly improve models predicting the

distribution of mutations observed across genes in experimentally evolved populations of *S. cerevisiae*

(Lang et al., 2013). We are also able to classify genomic variables into those that have affected

mutation counts 1) through their effect on the mutation rate (variables that significantly predict

synonymous mutations), and/ or 2) through their effect on the probability of a mutation being either

observed/ lost due to selection (variables that significantly predict nonsynonymous mutations). Out of

all the variables tested, we found that *gene length* explained the most variation in both synonymous and

nonsynonymous mutation counts per gene – plainly speaking, longer genes accumulate more

mutations. However, *number of domains* and *recombination* also had significant effects. Below we

discuss in detail these genomic variables and their potential contributions to the probability of parallel

evolution via the processes of mutation and selection.

Longer genes harbor more mutations – By far, the variable having the largest effect on variation in the

number of synonymous and nonsynonymous mutations observed was *gene length*. More specifically,

gene length positively affected the rate of mutation at the gene-level, meaning genes comprising more

nucleotides were more likely to harbor mutations. This result is not surprising and is in agreement with

recent analysis of synonymous mutation counts from Lenski's long term evolution experiment with *E.*

coli (Maddamsetti et al., 2015).

Long-term divergence does not predict short-term mutation counts – Our model for synonymous

mutation counts suggests that divergence estimates from long-term evolutionary comparisons at the

species level do not provide insight into expected mutation counts on the shorter time scale of evolution

in the lab, also in agreement with recent analysis of *E. coli* data (Maddamsetti et al., 2015).

Maddamsetti et al found that their proxy for long-term per gene mutation rate, θ_s (a measure of within-

species nucleotide diversity), did not explain gene-to-gene variation in synonymous mutation counts in

their data. The authors argued that horizontal gene transfer (HGT) is therefore likely a more important

process driving gene-to-gene variation in long-term divergence between naturally occurring *E. coli*

strains, and since HGT did not occur in their evolution experiment, it is not surprising that the

experiment's synonymous mutation counts did not correlate with θ_s . However, rates of HGT tend to be

higher in bacteria, and in particular *E. coli*, as compared to yeast and other eukaryotes (e.g. Boto,

2010). Furthermore, a recent mutation accumulation experiment with the eukaryote *Chlamydomonas*

reinhardtii showed a positive correlation between a proxy for long-term mutation rate (θ_s) and per site

mutability (Ness et al., 2015). Thus, it is somewhat surprising that we do not see a significant

relationship between *dS* and *dN/dS* and counts of synonymous and nonsynonymous mutations

respectively in our examination of the *S. cerevisiae* data used in this study. One possibility might also

be that d*S* and d*N*/d*S* are noisy to estimate at the gene level and that tends to downplay their predictive

power in our analysis of counts in an evolve and re-sequence experiment.

Nonsynonymous mutation counts show evidence of selection heterogeneity – As expected (Lenormand

et al., 2016), we see strong evidence that the distribution of nonsynonymous mutations across the

genome is driven in part by gene-to-gene heterogeneity in selection. Of those genomic variables tested,

we found three that were significant predictors of nonsynonymous mutation counts, suggesting that

those variables may drive or are correlated with processes that modulate the intensity of selection

across genes. The significant variables were *gene length*, *recombination rate*, and *number of protein*

domains.

We found that *gene length* predicts nonsynonymous mutation count via selection, over and

above its effects on per gene mutation rate – as estimated from models aimed at explaining the

synonymous mutation count only. While one might not expect *gene length* to have direct effects on

selection, we suggest that *gene length* may show a significant effect here because it is correlated with

other attributes of the genome that could have important effects on selection, for example *gene*

expression levels and *multifunctionality*. Because of these correlations, it could be that *gene length* acts

as a kind of summary variable for these covariates and other unidentified factors we have not captured

in these models. In fact, it is almost certainly the case that to some degree all the significant variables in

our model summarize variation from additional unknown factors we have not included in our data set.

In contrast to the positive relationship between *gene length* and number of nonsynonymous

mutations, we also found that the *number of protein domains* that a gene codes for (a variable that is

positively correlated with gene length; Table S1) actually negatively predicts the number of

nonsynonymous mutations. In other words, the more domains in the encoded protein of a gene, the

fewer mutations that gene is expected to incur in the course of the yeast evolution experiment analyzed

here. The mechanism behind this effect is not clear, but certainly protein structure has previously been

reported to have significant impacts on evolutionary rates in yeast (Bloom et al., 2006) and one can

also posit that genes encoding proteins with multiple domains and thereby involved in more numerous

interactions are – all else being equal – more severely constrained by purifying selection. It is

interesting that this effect can be observed in the course of relatively short time span (relative to

between species divergence times) through the relative paucity of nonsynonymous mutations in these

genes.

Our analysis also showed that *recombination rate* is a significant predictor of the number of

nonsynonymous mutations observed in a given gene in these data. Genes with higher recombination

rates are more likely to bear nonsynonymous mutations. We expect recombination rate to be correlated

with mutation, as previous studies in yeast have shown that recombinational repair of double strand

breaks substantially increases the frequency of point mutations in nearby intervals (e.g. Holbeck and

Strathern, 1997; Strathern et al., 1995). However, it is not clear how high recombination rates might

drive, or be correlated with other processes that drive, selection – as our models suggest is the case for

this data set. Another non-exclusive possibility might be the fact that biased gene conversion might

vary from gene to gene and also – like selection – affect the probability of detecting variants in evolve

and re-sequence experiments. However, if this was the case, we might expect GC content to be a

significant predictor of both synonymous and nonsynonymous mutations in our models, and it is not

(S: $P = 0.125$, NS: $P = 0.221$). We might also expect to see a bias towards GC in the observed

mutations, however this was not the case – only 33% of the SNPs in this data set were a change to G or

C.

Factors driving mutation and selection are complex – It is difficult to obtain any additional insights

from our models that include principal components of the genomic covariate data, however there is at

least some level of agreement between those variables that are significant (i.e. length, recombination,

and number of domains) and ones that are heavily weighted in PC10 – the principal component that

was found to be significant (see Fig. 3). The local properties of the genome do appear to drive some

heterogeneity in the selection processes, and in turn, shape the patterns of parallel evolution, however

individual effects that can be ascribed to individual variables are not easy to parse out.

We want to stress that while we were able to identify a number of factors affecting the count of

mutations observed in this evolution experiment data set, the total explained variance is still low: 1 %

and 16 % in the synonymous and nonsynonymous models respectively (calculated from pseudo-r^2

estimates of the "best" models, see methods). While the models do capture the general distribution of

mutation counts (Fig. 2) and so the degree of parallel evolution, accurately predicting on which genes

those mutations will fall is still very difficult. This is not surprising given the amount of stochasticity

involved in both the origin of new mutations and their evolutionary fate through drift and selection. A

clearer picture might emerge when using our modeling approach in a meta-analysis approach where

several evolve and re-sequence experiments are considered together (see Bailey et al., 2017 for a

similar approach on summary statistics of the amount of parallel evolution at the gene level across a

wide range of experimental studies in yeast and bacteria).

While we do find a number of genomic variables that significantly affect the distribution of

mutations across the genome, it is noteworthy that these models are still unable to capture the more

extreme patterns of parallel evolution observed in this data set. For example, one gene (IRA1) saw

mutations in over 50% of the populations sequenced in this experimental data set (discussed in more

detail in Lang et al., 2013). Such a mutation count is completely out of the range of likely outcomes

predicted by our models. Some of this discrepancy may be because of the simplifying assumptions

made about the process of selection. Our framework models the process of mutation and its

heterogeneity but while we account for the fact that newly arising mutations may have different

probabilities of reaching an observable frequency, the modeling of that process could be made more

precise by incorporating an explicit underlying distribution of fitness effects of new mutations at each

gene. Incorporating a selection process that allows for different amounts of both positive and negative

selection, as well as further details about the selection pressures in the particular environment of

interest – something we do not consider at all in this study – would likely improve prediction for some

of these more extreme events.

It is also important to note that the methods used in this study are focused on parallelism in

SNPs and small indels. While this focus is appropriate for the data set used here, it may not be

appropriate for other systems. For example, in an experimental evolution study with E. coli, (Tenaillon

et al., 2012) saw that much of the parallelism seen between populations was the result of IS elements

and large scale duplication and deletions. It may be important to try to account for this more diverse

range of mutational event types when trying to identify the drivers of parallel evolution other systems.

Can we use this modeling framework to predict parallel evolution? To some degree, yes – the

measures of parallel evolution between populations simulated using our model predictions span a

similar range to those calculated from the real population data (see Fig. 4). However, while this

congruence suggests we are on the right track, the shapes of the real and simulated distributions are still

quite different. For example, there is, on average, more parallelism between the real populations

compared to populations simulated from our models, and in particular, there seems to be a substantial

discrepancy between the number of real population-pairs and simulated population-pairs that have a

low level of parallelism (i.e. note the difference between real and simulated when J ranges from 0.02 to

0.04 in Fig. 4). This is further evidence suggesting that although our current models may be useful to

some extent, we are still missing some important factors driving heterogeneity in mutation and

selection across these genomes.

Advantages of this regression framework – Relying on the assumption that synonymous mutations are

selectively neutral (which does appear to be appropriate for these data), the regression models we use

in this study allow us to distinguish between genomic variables influencing the observed distribution of

mutations across a genome through their potential effects on both gene-to-gene heterogeneity in

mutation rate and gene-to-gene heterogeneity in selection. The great advantage of this is that it allows

us to begin to break down the importance of these two processes in shaping patterns of parallel

evolution we see, and move closer to the goal of predicting which genes will be involved in evolution

when organisms adapt to new environments. It will be interesting to apply this model framework to

other data sets of this type, as they become available, to see how general these patterns are across

different organisms and selection environments (Bailey and Bataillon, 2016).

Acknowledgments

We thank three PCI Evolutionary Biology recommenders for their constructive comments. This work

was supported by the European Research Council under the European Union's Seventh Framework

Program [FP7/20072013, ERC grant number 311341 to T.B.].

References

Akaike, H. (1973). Information theory and an extension of the maximum likelihood principle. In Proceeding of the Second International Symposium on Information Theory, (Budapest: Akademiai Kiado), pp. 267–281.

Bailey, S.F., and Bataillon, T. (2016). Can the experimental evolution programme help us elucidate the genetic basis of adaptation in nature? Mol. Ecol. *25*, 203–218.

Bailey, S.F., Blanquart, F., Bataillon, T., and Kassen, R. (2017). What drives parallel evolution? BioEssays *39*, 1–9.

Bloom, J.D., Drummond, D.A., Arnold, F.H., and Wilke, C.O. (2006). Structural Determinants of the Rate of Protein Evolution in Yeast. Mol. Biol. Evol. *23*, 1751–1761.

Boto, L. (2010). Horizontal gene transfer in evolution: facts and challenges. Proc. R. Soc. Lond. B Biol. Sci. *277*, 819–827.

Caballero, J.D., Clark, S.T., Coburn, B., Zhang, Y., Wang, P.W., Donaldson, S.L., Tullis, D.E., Yau, Y.C.W., Waters, V.J., Hwang, D.M., et al. (2015). Selective sweeps and parallel pathoadaptation drive Pseudomonas aeruginosa evolution in the cystic fibrosis lung. MBio *6*, e00981-15.

Castoe, T.A., de Koning, A.J., Kim, H.-M., Gu, W., Noonan, B.P., Naylor, G., Jiang, Z.J., Parkinson, C.L., and Pollock, D.D. (2009). Evidence for an ancient adaptive episode of convergent molecular evolution. Proc. Natl. Acad. Sci. *106*, 8986–8991.

Cherry, J.M., Hong, E.L., Amundsen, C., Balakrishnan, R., Binkley, G., Chan, E.T., Christie, K.R., Costanzo, M.C., Dwight, S.S., Engel, S.R., et al. (2012). Saccharomyces Genome Database: the genomics resource of budding yeast. Nucleic Acids Res. *40*, D700–D705.

Chevin, L.-M., Martin, G., and Lenormand, T. (2010). Fisher's model and the genomics of adaptation: restricted pleiotropy, heterogenous mutation, and parallel evolution. Evolution *64*, 3213–3231.

Christin, P.-A., Weinreich, D.M., and Besnard, G. (2010). Causes and evolutionary significance of genetic convergence. Trends Genet. *26*, 400–405.

Cliften, P., Sudarsanam, P., Desikan, A., Fulton, L., Fulton, B., Majors, J., Waterston, R., Cohen, B.A., and Johnston, M. (2003). Finding functional features in Saccharomyces genomes by phylogenetic footprinting. Science *301*, 71–76.

Drummond, D.A., and Wilke, C.O. (2008). Mistranslation-induced protein misfolding as a dominant constraint on coding-sequence evolution. Cell *134*, 341–352.

Drummond, D.A., Bloom, J.D., Adami, C., Wilke, C.O., and Arnold, F.H. (2005). Why highly expressed proteins evolve slowly. Proc. Natl. Acad. Sci. U. S. A. *102*, 14338–14343.

Feldman, C.R., Brodie, E.D., Brodie, E.D., and Pfrender, M.E. (2012). Constraint shapes convergence in tetrodotoxin-resistant sodium channels of snakes. Proc. Natl. Acad. Sci. *109,* 4556–4561.

Gillespie, J.H. (1984). Molecular evolution over the mutational landscape. Evolution *38*, 1116–1129.

Holbeck, S.L., and Strathern, J.N. (1997). A role for REV3 in mutagenesis during double-strand break repair in Saccharomyces cerevisiae. Genetics *147*, 1017–1024.

Holstege, F.C., Jennings, E.G., Wyrick, J.J., Lee, T.I., Hengartner, C.J., Green, M.R., Golub, T.R., Lander, E.S., and Young, R.A. (1998). Dissecting the regulatory circuitry of a eukaryotic genome. Cell *95*, 717–728.

Illingworth, C.J.R., Parts, L., Bergström, A., Liti, G., and Mustonen, V. (2013). Inferring Genome-Wide Recombination Landscapes from Advanced Intercross Lines: Application to Yeast Crosses. PLoS ONE *8*, e62266.

Jost, M.C., Hillis, D.M., Lu, Y., Kyle, J.W., Fozzard, H.A., and Zakon, H.H. (2008). Toxin-resistant sodium channels: Parallel adaptive evolution across a complete gene family. Mol. Biol. Evol. *25*, 1016–1024.

Kellis, M., Patterson, N., Endrizzi, M., Birren, B., and Lander, E.S. (2003). Sequencing and comparison of yeast species to identify genes and regulatory elements. Nature *423*, 241–254.

Koch, E.N., Costanzo, M., Bellay, J., Deshpande, R., Chatfield-Reed, K., Chua, G., D'Urso, G., Andrews, B.J., Boone, C., Myers, C.L., et al. (2012). Conserved rules govern genetic interaction degree across species. Genome Biol *13*, R57.

Lang, G.I., Botstein, D., and Desai, M.M. (2011). Genetic variation and the fate of beneficial mutations in asexual populations. Genetics *188*, 647–661.

Lang, G.I., Rice, D.P., Hickman, M.J., Sodergren, E., Weinstock, G.M., Botstein, D., and Desai, M.M. (2013). Pervasive genetic hitchhiking and clonal interference in forty evolving yeast populations. Nature *500*, 571–574.

Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H., Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., et al. (2007). Clustal W and Clustal X version 2.0. Bioinformatics *23*, 2947–2948.

Lenormand, T., Chevin, L.M., and Bataillon, T. (2016). Parallel evolution: what does it (not) tell us and why is it (still) interesting. In Chance in Evolution, (Chicago, Illinois: Chicago University Press), p.

Liu, Z., Qi, F.-Y., Zhou, X., Ren, H.-Q., and Shi, P. (2014). Parallel sites implicate functional convergence of the hearing gene prestin among echolocating mammals. Mol. Biol. Evol. *31*, 2415–2424.

Maddamsetti, R., Hatcher, P.J., Cruveiller, S., Médigue, C., Barrick, J.E., and Lenski, R.E. (2015). Synonymous genetic variation in natural isolates of Escherichia coli does not predict where synonymous substitutions occur in a long-term experiment. Mol. Biol. Evol. msv161.

Marvig, R.L., Sommer, L.M., Molin, S., and Johansen, H.K. (2015). Convergent evolution and adaptation of Pseudomonas aeruginosa within patients with cystic fibrosis. Nat. Genet. *47*, 57–64.

McVean, G.A.T., Myers, S.R., Hunt, S., Deloukas, P., Bentley, D.R., and Donnelly, P. (2004). The Fine-Scale Structure of Recombination Rate Variation in the Human Genome. Science *304*, 581–584.

Ness, R.W., Morgan, A.D., Vasanthakrishnan, R.B., Colegrave, N., and Keightley, P.D. (2015). Extensive de novo mutation rate variation between individuals and across the genome of Chlamydomonas reinhardtii. Genome Res. *25*, 1739–1749.

Pál, C., Papp, B., and Hurst, L.D. (2001). Highly expressed genes in yeast evolve slowly. Genetics *158*, 927–931.

Punta, M., Coggill, P.C., Eberhardt, R.Y., Mistry, J., Tate, J., Boursnell, C., Pang, N., Forslund, K., Ceric, G., Clements, J., et al. (2012). The Pfam protein families database. Nucleic Acids Res. *40*, D290–D301.

R Development Core Team (2014). R: a language and environment for statistical computing (R Foundation for Statistical Computing, Vienna, Austria).

Self, S.G., and Liang, K.-Y. (1987). Asymptotic properties of maximum likelihood estimators and likelihood ratio tests under nonstandard conditions. J. Am. Stat. Assoc. *82*, 605–610.

Sharp, P.M., and Li, W.-H. (1987). The Codon Adaptation Index - A measure of directional synonymous codon usage bias, and its potential applications. Nucleic Acids Res. *15*, 1281–1295.

Stark, C., Breitkreutz, B.-J., Reguly, T., Boucher, L., Breitkreutz, A., and Tyers, M. (2006). BioGRID: a general repository for interaction datasets. Nucleic Acids Res. *34*, D535-539.

Strathern, J.N., Shafer, B.K., and McGill, C.B. (1995). DNA Synthesis Errors Associated with Double-Strand-Break Repair. Genetics *140*, 965–972.

Streisfeld, M.A., and Rausher, M.D. (2011). Population genetics, pleiotropy, and the preferential fixation of mutations during adaptive evolution. Evolution *65*, 629–642.

Tenaillon, O., Rodríguez-Verdugo, A., Gaut, R.L., McDonald, P., Bennett, A.F., Long, A.D., and Gaut, B.S. (2012). The Molecular Diversity of Adaptive Convergence. Science *335*, 457–461.

Winzeler, E.A., Shoemaker, D.D., Astromoff, A., Liang, H., Anderson, K., Andre, B., Bangham, R., Benito, R., Boeke, J.D., Bussey, H., et al. (1999). Functional Characterization of the S. cerevisiae Genome by Gene Deletion and Parallel Analysis. Science *285*, 901–906.

Wong, A., Rodrigue, N., and Kassen, R. (2012). Genomics of adaptation during experimental evolution of the opportunistic pathogen Pseudomonas aeruginosa. PLoS Genet *8*, e1002928.

Woods, R., Schneider, D., Winkworth, C.L., Riley, M.A., and Lenski, R.E. (2006). Tests of parallel molecular evolution in a long-term experiment with Escherichia coli. Proc. Natl. Acad. Sci. *103*, 9107–9112.

Yang, Z. (2007). PAML 4: Phylogenetic Analysis by Maximum Likelihood. Mol. Biol. Evol. *24*, 1586–1591.

Zhang, J., and Kumar, S. (1997). Detection of convergent and parallel evolution at the amino acid sequence level. Mol. Biol. Evol. *14*, 527–536.

Zou, Z., and Zhang, J. (2015). Are convergent and parallel amino acid substitutions in protein evolution more prevalent than neutral expectations? Mol. Biol. Evol. *32*, 2085–2096.

Zuur, A.F. (2009). Mixed effects models and extensions in ecology with R (Springer).

FIGURES

Figure 1: Schematic showing how the mutation counts data are generated and general assumptions underlying these data.

487
488
489 **Figure 2**: Distribution of A) synonymous and B) nonsynonymous mutations per gene (totaled over all
490 40 populations in the data set) and predicted model distributions from M0.P (grey circles), M1.P (black
491 points), M2.P (green triangles), and $M_N.NB$ (blue squares), and $M_N.NB_{PC}$ (orange diamonds).

Figure 3: Loadings of the 11 genomic variables on PC10 – the only principal component that significantly explains variation in nonsynonymous mutation counts. Genomic variables are ordered from largest to smallest in terms of the absolute value of their loading.

498   **Figure 4**: Distribution of the degree of parallelism (estimated as the pairwise Jaccard Index, J) from the
499   real data and simulated data from the best-fit models.

TABLES

Table 1: Genomic variables used in this study.

Variable name	Description	Reference
dS	Number of synonymous substitutions per synonymous site, estimated from gene alignments of *S. cerevisiae*, *S. paradoxus*, *S. bayanus*, and *S. mikatae* (Cliften et al., 2003; Kellis et al., 2003).	Estimated for this study.
dN/dS	Number of nonsynonymous substitutions per nonsynonymous site, estimated from *S. cerevisiae*, *S. paradoxus*, *S. bayanus*, and *S. mikatae (Cliften et al., 2003; Kellis et al., 2003)*.	Estimated for this study.
Gene length *(L)*	The number of nucleotides.	(Cherry et al., 2012)
% GC content *(GC)*	Percentage of nucleotides in the gene sequence that are either guanine or cytosine.	(Cherry et al., 2012)
Multi-functionality *(multifunc)*	Number of different GO slim categories assigned to a gene.	(Cherry et al., 2012)
Degree of protein-protein interaction *(PPI)*	The number of physical interactions reported by BioGRID (Stark et al., 2006).	(Koch et al., 2012)
Codon adaptation index *(CAI)*	A measure of bias in the usage of synonymous codons, based on a comparison between codon frequencies in the gene and frequencies observed in a set of highly expressed genes (Sharp and Li, 1987).	(Koch et al., 2012)
Number of domains *(num.dom)*	The number of regions that Pfam (Punta et al., 2012) has identified as domains in the protein sequence of each gene.	(Koch et al., 2012)
Level of expression *(expr)*	A measure of mRNA level for each gene when grown in standard lab conditions.	(Holstege et al., 1998)
Local recombination rate *(r)*	Mean recombination rate for a given gene calculated from recombination rate estimate at 0.5 kb intervals using *LDhat* (McVean et al., 2004).	(Illingworth et al., 2013)
Essential genes *(essential)*	A true/ false indicator variable denoting whether or not a gene is essential, based on growth assays of deletion strains.	(Winzeler et al., 1999)

Table 2: 'M$_S$' models testing assumptions with the synonymous mutation data. Log-likelihoods, and AIC values are provided. The best model as determined by the lowest AIC with the fewest parameters is highlighted in grey.

Model		log-lik.	No. param.	AIC
M$_S$0.P:	Pois(λ_S = *constant*)	-283.0	1	568.2
M$_S$0.NB:	NB(λ_S = *constant*, θ_S)	-283.0	2	569.9
M$_S$1.P:	Pois(λ_S = *constant* * $L_i{}^{\alpha1}$)	-273.9	2	551.8
M$_S$1.NB:	NB(λ_S = *constant* * $L_i{}^{\alpha1}$, θ_S)	-273.9	3	553.8
M$_S$2.P:	Pois(λ_S = *constant* * L_i)	-274.0	1	549.9
M$_S$2.NB:	NB(λ_S = *constant* * L_i, θ_S)	-274.0	2	551.9

Table 3: 'M$_N$' models parameter estimates (*constant, α1, α2*, etc) and P-values for those estimates. Only those variables that significantly improved model fit are included.

M$_N$.NB: NB(λ_N = *constant* * L_i * $L_i{}^{\alpha1}$ * *num.dom$_i{}^{\alpha2}$* * $r_i{}^{\alpha3}$, θ_N)

		Estimate	P-value
λ_N ~	L	$\alpha1$ = 0.4405	0.001
	num.dom	$\alpha2$ = -0.4511	0.004
	r	$\alpha3$ = 0.1029	0.041
	constant	$8.084 * 10^{-6}$	<0.001
θ_N		0.3806	<0.001

M$_N$.NB$_{PC}$: NB(λ_N = *constant* * L_i * exp(*PC10$_i$*) $^{\alpha1}$, θ_N)

		Estimate	P-value
λ_N ~	exp(*PC10*)	$\alpha1$ = 0.2984	<0.001
	constant	$8.846 * 10^{-5}$	<0.001
θ_N		0.3988	<0.001

Table 4: Log-likelihoods, and AIC values for the 'M$_N$' models. The best model as determined by the lowest AIC with the fewest parameters is highlighted in grey.

Model	log-lik.	No. param.	AIC
M$_N$0.P:	-1159.9	1	2321.7
M$_N$0.NB:	-1022.5	2	2048.9
M$_N$2.P:	-1050.3	1	2102.7
M$_N$2.NB:	-956.6	2	1917.3
M$_N$.P	-1021.4	4	2050.8
M$_N$.P$_{PC}$	-1013.0	2	2030.1
M$_N$.NB	-944.8	5	1899.7
M$_N$.NB$_{PC}$	-947.9	3	1901.8