Title: Discovering *in vivo* cytokine eQTL interactions from a lupus
clinical trial

Authors: Emma E. Davenport,[1,2,3,4] Tiffany Amariuta,[1,2,3,4,5] Maria Gutierrez-
Arcelus,[1,2,3,4] Kamil Slowikowski,[1,2,3,4,5] Harm-Jan Westra,[1,2,3,4] Yang Luo,[1,2,3,4] Ciyue
Shen,[6] Deepak A. Rao,[7] Ying Zhang,[8] Stephen Pearson,[9] David von Schack,[8] Jean S.
Beebe,[8] Nan Bing,[8] Sally John,[10] Michael S. Vincent,[8] Baohong Zhang[8] and Soumya
Raychaudhuri[1,2,3,4,5,11*]

Affiliations:
[1]Center for Data Sciences, Brigham and Women's Hospital, Boston, MA, 02115, USA.
[2]Divisions of Genetics and Rheumatology, Department of Medicine, Brigham and
Women's Hospital, Harvard Medical School, Boston, MA, 02115, USA.
[3]Partners Center for Personalized Genetic Medicine, Boston, MA, 02115, USA.
[4]Program in Medical and Population Genetics, Broad Institute of MIT and Harvard,
Cambridge, MA, 02142, USA.
[5]Department of Biomedical Informatics, Harvard Medical School, Boston, MA, 02115,
USA.
[6]Department of Cell Biology, Harvard Medical School, Boston, MA, 02115, USA
[7]Division of Rheumatology, Allergy, Immunology, Brigham and Women's Hospital,
Harvard Medical School, MA 02115, USA.
[8]Pfizer Inc., Cambridge, MA, 02139, USA.
[9]Pfizer New Haven Clinical Research Unit, New Haven, CT, 06511, USA.
[10]Biogen, Cambridge, MA, 02142, USA.
[11]Faculty of Medical and Human Sciences, University of Manchester, Manchester, M13
9PL, UK.

*Correspondence to:
Soumya Raychaudhuri
77 Avenue Louis Pasteur
Harvard New Research Building, Suite 250D
Boston, MA 02446, USA.
soumya@broadinstitute.org; 617-525-4484 (tel); 617-525-4488 (fax)

Abstract:

Background: Cytokines are critical to human disease and are attractive therapeutic

targets given their widespread influence on gene regulation and transcription. Defining

the downstream regulatory mechanisms influenced by cytokines is central to defining

drug and disease mechanisms. One promising strategy is to use interactions between

expression quantitative trait loci (eQTLs) and cytokine levels to define target genes and

mechanisms.

Results: In a clinical trial for anti-IL-6 in patients with systemic lupus erythematosus we

measured interferon (IFN) status, anti-IL-6 drug exposure and genome-wide gene

expression at three time points (379 samples from 157 individuals). First, we show that

repeat transcriptomic measurements increases the number of *cis* eQTLs identified

compared to using a single time point by 64%. Then, after identifying 4,818 cis-eQTLs,

we observed a statistically significant enrichment of *in vivo* eQTL interactions with IFN

status (p<0.001 by permutation) and anti-IL-6 drug exposure (p<0.001). We observed

210 and 72 interactions for IFN and anti-IL-6 respectively (FDR<20%). Anti-IL-6

interactions have not yet been described while 99 of the IFN interactions are novel.

Finally, we found transcription factor binding motifs interrupted by eQTL interaction

SNPs, pointing to key regulatory mediators of these environmental stimuli and therefore

potential therapeutic targets for autoimmune diseases. In particular, genes with IFN

interactions are enriched for ISRE binding site motifs, while those with anti-IL-6

interactions are enriched for IRF4 motifs.

Conclusion: This study highlights the potential to exploit clinical trial data to discover *in*

vivo eQTL interactions with therapeutically relevant environmental variables.

Keywords: eQTL, interactions, clinical trials, cytokines

Background

Cytokines are critical signals used by the immune system to coordinate inflammatory

responses. These factors bind to specific receptors to induce widespread transcriptional

effects. Cytokines and their receptors are not only genetically associated with

susceptibility to a range of human diseases, they have also emerged as effective

therapeutic targets[1]. Blockade of tumour necrosis factor (TNF) was the first cytokine-

directed therapy to achieve widespread use and is now used broadly to treat multiple

inflammatory diseases including rheumatoid arthritis (RA), psoriasis, and inflammatory

bowel disease[2]. More recently, IL-6 has emerged as a compelling therapeutic target.

IL-6 levels are elevated in autoimmune diseases such as systemic lupus erythematosus

(SLE) and RA. The IL-6 receptor has been successfully targeted with tocilizumab in

RA[3] and giant cell arteritis[4], while IL-6 has been targeted directly with siltuximab for

successful treatment of Castleman's disease[5]. In SLE, IL-6 is thought to play a role in

the observed B cell hyperactivity and autoantibody production[6]. Targeting IL-6-R in

SLE has shown promise in phase I trials[7] and this has led to the development of other

biologics targeting IL-6 such as PF-04236921[8]. Interferon (IFN)-α, produced primarily

by plasmacytoid dendritic cells, has pleiotropic effects on the immune system. It has

been implicated as a key mechanism in SLE development and pathogenesis, and is

being investigated as a therapeutic target[9]. Agents targeting other inflammatory

cytokines, including Interleukin-1 (IL-1), IL-12, and IL-17A and IL-23 are also in clinical

use to treat autoimmune conditions. Interestingly, IL-1 blockade with canakinumab has

also been recently reported to reduce risk of heart attacks, stroke and cardiovascular

disease[10]. Therefore, defining the regulatory consequences of physiologic

perturbations of cytokine levels will inform our understanding of both disease and drug

mechanisms.

A *cis* expression quantitative trait locus (eQTL) contains a genetic variant that alters

expression of a nearby gene. *Cis* eQTLs are ubiquitous across the genome[11] and

while most are stable across tissues and conditions, environmental variables can alter

the effects of some of them[12–18]. If an environmental change leads to disruption of

regulators upstream of a gene, then it could magnify or dampen an eQTL effect,

resulting in a genotype-by-environment interaction (**Figure S1**). Therefore, observing a

set of eQTL interactions due to a perturbagen, such as a cytokine, can identify shared

upstream regulatory mechanisms, such as transcription factors and key pathways. Even

a single eQTL interaction where we can define mechanism can lead to insights about

the action of the perturbagen.

However, *cis* eQTL interactions with physiologic environmental factors in humans have

been challenging to discover *in vivo*[19–23] even with large cohorts[11, 17]. Success at

finding *cis* eQTL interactions has largely been found in studies using model

organisms[24, 25] or treating cells *in vitro* with non-physiologic conditions[26]. Thus far,

these studies might be limited in power since they often map eQTLs separately across

conditions and fail to exploit the power of repeat measurements[27]. In other instances,

they test for genetic variants associated with differential expression and miss

information about the magnitude of the eQTL effect in a specific condition[28].

We predicted that if the transcriptome is assayed at multiple time points under different

exposure states, then the repeat measurements could lead to an increase in power to

detect eQTLs and their interactions with environmental perturbations. If the same

individual is assessed at multiple times, then the noise in transcriptomic measurements

is reduced. Furthermore, repeat measurements from the same individuals when they

are both unexposed and exposed to an environmental perturbagen allow for more

accurate modelling of the effect of the perturbagen within those subjects.

Clinical trials, with their structured study design, may be the ideal setting to detect eQTL

interactions with therapeutically important variables. In clinical trials, it is becoming

increasingly common to collect transcriptional and genetic data alongside clinical and

physiological data[29]. This extensive phenotyping of therapeutically important variables

and biomarkers within the same individual at multiple time points provides a unique

opportunity to identify *in vivo* eQTL interactions.

Here, we examined the modulation of eQTL effects by environmental factors that alter

cytokine levels using data from a phase II clinical trial to evaluate the safety and efficacy

of a neutralizing IL-6 monoclonal antibody (PF-04236921) in 157 SLE patients[8]

(**Methods**). Many patients with SLE exhibit high levels of genes induced by type I IFN;

these genes, known as the IFN signature, are a marker of disease severity[30, 31] and

a pathogenic feature of SLE. This feature of the disease, together with exposure to anti-

IL-6 leads to cytokine fluctuations in this cohort yielding opportunities to assess the

impact of cytokine levels on eQTL effects. While this drug was not significantly different

from placebo for the primary efficacy endpoint (proportion of patients achieving the SLE

Responder Index (SRI-4) at week 24), biologically it effectively reduced free IL-6 protein

levels (**Figure S2**). Given the key role of IL-6 and IFN in a range of diseases, the

downstream regulatory effects of these cytokines are of great interest to study.

In this study, we leverage the power of repeat transcriptional and environmental

measurements from a lupus clinical trial to identify *in vivo* eQTL interactions with IFN

status and anti-IL-6 exposure. In the process, we define novel eQTL interactions for

both IFN and IL-6.

Results

We conducted whole blood high-depth RNA-seq profiling at 0, 12, and 24 weeks in anti-

IL-6 exposed and unexposed individuals with the Illumina TruSeq protocol. We

quantified 20,253 gene features and examined 1,595,793 genotyped and imputed

common variants genome-wide (**Methods**). Along with each RNA-seq assay, we

documented anti-IL-6 exposure and quantified IFN signature status with real-time PCR.

Mapping eQTL in SLE patients

We first mapped *cis* eQTLs and then tested them for interactions with IFN status and

anti-IL-6 exposure. eQTL interactions can be explored using our interactive visualisation

tool (http://baohongz.github.io/Lupus_eQTL, **Figure S3**).

To identify *cis* eQTLs, we examined the association between gene expression and

SNPs within 250kb upstream of the transcription start site and 250kb downstream of the

transcription end site. In order to account for repeat measurements, with up to three

RNA-seq assays per patient (**Figure 1A,** 379 samples from 157 patients, **Methods**), we

used a linear mixed model. We included 25 gene expression principal components to

maximise the number of eQTL detected and 5 genotyping principal components to

account for the heterogeneity in ethnicity in our cohort (**Methods**). We observed that the

multi-ethnic nature of our study did not confound our results, consistent with Stranger et

al.[32] (**Figure S4**).

To ensure we only tested for interactions in a set of highly confident eQTLs, we applied

a stringent correction for the total number of hypotheses tested. We recognized that this

approach might arguably be overly stringent for eQTL discovery, but we wanted to be

certain that we were only testing eQTLs for interactions that had a convincing main

effect. Since we tested a total of 5,872,001 SNP-gene pairs genomewide, we set a

significance threshold of $p_{eqtl} < 8.5 \times 10^{-9}$ (0.05/5,872,001 tests). We identified 4,818 *cis*

eQTL genes (**Figure 1B,1C, Table S1**). The summary statistics for all the gene SNP

pairs tested are available in **Table S2**.

To confirm the validity of our eQTLs, we compared them to a larger dataset. In the BIOS

cohort, consisting of 2,166 healthy individuals[11], we observed that 85.4% of our SLE

eQTL SNP-gene pairs are reported as eQTLs (FDR<0.05). Of these, 98.9% showed

consistent direction of effect ($p<5\times10^{-16}$, binomial test, **Figure 1D**), suggesting that our

results were highly concordant with those in this substantially larger study.

Repeat measurements increase power to detect eQTL

Under reasonable assumptions, we would expect repeat samples to increase our

power. Supporting that expectation, we detected 64% more *cis* eQTLs compared to the

2,934 genes from using a single sample (first available time point) per individual (**Figure**

1B). An alternative might have been to identify eQTLs separately from each of the three

time points; however, this approach identified only a total of 3,050 eQTL genes (**Figure**

S5). Modelling all three time points together results in 58% more *cis* eQTLs than

modelling each time point separately.

We speculated that while repeat measures did increase power over single measures,

that given a fixed number of samples, independent samples would lead to more power.

To this end, we conducted an analysis fixing the number of samples at 157 and using

53 individuals with repeat measures (with two missing samples). Unsurprisingly, we

found fewer eQTLs (2,215 genes) with the repeat measures alone compared to an

analysis with the same number of independent samples (2,934 genes).

IFN status eQTL interactions

For each of the 4,818 *cis* eQTL genes, we tested the most significantly associated SNP

for environmental interactions with our linear mixed model framework. We first explored

the influence of type I IFN on gene regulation after determining the IFN status of every

patient at each time point. We classified each sample as either IFN high or IFN low

using real-time PCR of 11 IFN-inducible genes[33] (**Methods, Figure 2A**).

We first wanted to assess whether our results were indeed enriched for interactions. To

do this, we identified those eQTLs with nominally significant interaction effects at

$p_{interact}<0.01$. We would expect ~48 out of 4,818 from chance alone. Surprisingly we

observed 182 IFN-eQTL interactions (**Table S1**) that were nominally significant at

$p_{interact}<0.01$ suggesting that there was evidence of enrichment for eQTL interactions.

We conducted permutations to ensure that these results were not the consequence of

potentially inflated statistics, which might be the result for example of low frequency

alleles, genes violating normality assumptions, or other technical artefacts. In each of

1,000 stringent permutations, we simply reassigned IFN status across samples and

retested for eQTL interactions. This permutation preserves the main eQTL effect, since

it maintains genotypes of the individuals with the associated expression data, but

disrupts any real interactions that might be present in the data. In 0 out of 1,000

instance did we observe 182 or more interactions at $p_{interact}<0.01$ suggesting that the

number of observed interactions is enriched and highly unlikely to have happened by

chance (**Figure S6**, $p_{permute}$~0/1,000 = <0.001).

We then went on to identify those specific IFN-eQTL interactions of greatest interest by

calculating a false discovery rate or q value for each interaction using the q value

package[34] (**Methods**). We observed a total of 210 interactions with an FDR<0.2

threshold (**Table S1**). We note that 11 of these genes have already been described as

having an interaction with a proxy gene for type I IFN signalling in the much larger BIOS

study[11]. For example, *SLFN5* expression is influenced by the rs12602407 SNP

($p_{interact}$ =1.3x10^{-10}, FDR <9.9x10^{-8}, **Figure 2B**) and this effect is magnified in IFN high

samples. Of these 210 IFN-eQTL interactions, 99 were not reported in the BIOS

study[11]. Indeed, applying a more stringent cut off of FDR<0.01, 27/34 of our

interactions are not previously reported and therefore are almost certainly novel IFN-

eQTL interactions with high confidence (**Figure S7**).

We speculated that groups of eQTL interactions might be driven by the same common

regulatory factor. We divided interactions into magnifiers, where the environmental

exposure increases the size of the eQTL effect, and dampeners where the

environmental exposure decreases the eQTL effect (**Figure S8**). We hypothesized that

the transcription factors driving the response to type I IFN may be different for the eQTL

interactions defined as magnifiers (n=127, FDR<0.2) and dampeners (n=83, FDR<0.2).

We applied HOMER[35] to assess overlap between transcription factor binding motifs

and the eQTL interaction SNPs (and SNPs in high linkage disequilibrium (LD, r^2>0.8) in

the *cis* window, **Methods**). We conducted two separate analyses: magnifying eQTL

interactions in the foreground with dampening interactions in the background and vice

versa. We found enrichment of motifs for key transcription factors involved in IFN

signalling including a statistically significant enrichment for the ISRE motif (HOMER

p=1x10^{-4}, **Table S3**). The ISRE motif disruption occurred for 11 genes with an eQTL

magnified in IFN high samples but for only 1 gene with an eQTL dampened

(permutation p<0.019, **Methods, Figure 2C**). An example is the *GTF2A2* rs2306355

eQTL ($p_{interact}$=8.7x10^{-3}, FDR<0.15, **Figure 2D**); rs2306355 is in tight LD (r^2=0.83 in

Europeans) with rs6494127, which interrupts the TTCNNTTT core of the ISRE motif

(**Figure 3C**). This SNP likely disrupts IRF9 and STAT2 binding in the ISGF3

complex[36], which binds to the ISRE motif. We observe greater expression of *GTF2A2*

in individuals with the rs2306355 A allele compared to G; this difference is magnified in

IFN high individuals (**Figure 2D**).

We considered that the principal components included as covariates in our model might

be mitigating power. For example, the 4th principal component of gene expression is

correlated with the IFN signature status of the sample (r_s= -0.7, **Table S4**) so we

repeated the IFN interaction analysis without correcting for principal component 4. For

all the eQTLs tested for an IFN interaction, we observed very similar results with highly

correlated z-scores (r_s=0.94, **Figure S9**).

Discovery of eQTL interactions with anti-IL-6 drug exposure

We then examined whether IL-6 blockade alters the relationship between genomic

variation and gene expression and induces drug-eQTL interactions. We wanted to first

test if there was evidence of such interactions in our data set. Again, using a threshold

of $p_{interact}$<0.01 for nominal significance for interactions, we observed 121 drug-eQTL

interactions with anti-IL-6 out of 4,818 eQTLs tested (**Table S1**); similar to IFN

interactions, this is far in excess of the ~48 we would expect by chance. As above, to

ensure that these results were not the consequence of statistical artefact, we applied

the same stringent permutation strategy, reassigning which samples were exposed or

not to anti-IL-6. After 1,000 permutations, we never observed as many as 121 drug-

eQTL interactions with $p_{interact}<0.01$ (**Figure S10**), suggesting that our eQTLs were

indeed highly enriched for those interacting with anti-IL-6 ($p_{permute}\sim0/1000<0.001$).

To identify specific eQTL events that interact with anti-IL-6, we again calculated a false

discovery rate. We observed that 72 of these interactions have an FDR<0.2. Only 8 of

these drug-eQTL interactions overlap with the interactions observed for IFN status

(**Table S1**). We note biologically relevant drug-eQTL interactions for *IL10*

($p_{interact}=2.6 \times 10^{-3}$, FDR<0.19, **Figure S11**), an anti-inflammatory cytokine, *CLEC4C*

($p_{interact}=2.9 \times 10^{-3}$, FDR<0.19) which has previously been associated in *trans* with an SLE

risk allele[37] and *CLEC18A* ($p_{interact}=5.1 \times 10^{-4}$, FDR<0.14, **Figure 3A**) another member

of the C-type lectin domain family.

Similar to the IFN-eQTL interactions, we divided the drug-eQTL interactions into

magnifiers (n=33, FDR<0.2) and dampeners (n=39, FDR<0.2) (**Figure S12**) and used

the approach as described above to define transcription factors potentially driving the

response to IL-6 blockade (**Table S5**). One of the motifs enriched for eQTLs magnified

after drug treatment (with dampeners in the background) was IRF4 (HOMER $p=1 \times 10^{-3}$).

The IRF4 motif disruption occurred for 9 genes, including *CLEC18A*, with an eQTL

magnified after drug treatment compared to 4 genes with an eQTL dampened (**Figure**

3B, **Methods**). We permuted the magnifying and dampening genes and found this ratio

for enrichment is suggestive at the gene level (p=0.058) but additional eQTL

interactions will be necessary to confirm.

Comparing differential expression to eQTL interactions

A more common strategy to determine the effect of an environmental variable is to use

differential gene expression. For differential expression following anti-IL-6 treatment, we

identified 415 genes with FDR<0.05 but modest effects (max fold change=1.3, **Figure**

S13). Intriguingly, only 1/72 drug-eQTL interaction genes also show evidence of

differential gene expression. This suggests that eQTL interactions offer independent

information from differential expression, which might contribute to defining mechanisms.

Concordance of drug-eQTL interactions with protein level interactions

We hypothesized that interactions due to drug exposure are likely driven by free IL-6

cytokine levels (our key clinical biomarker of interest). If this is the case, for eQTLs

dampened by drug exposure, an increase in free IL-6 should elicit an opposite

interaction effect and result in eQTL magnification. We assessed whether eQTL

interactions with free IL-6 protein levels measured in the patient serum samples were

consistent with those following IL-6 blockade. We observed enrichment in the overlap

between cytokine interactions and drug interactions (53/72 interactions in expected

opposite direction, **Figure 3C,** $p=3.8 \times 10^{-5}$, binomial test).

Discussion

In this study we mapped eQTLs in a clinical trial of SLE patients and discovered

interactions with IFN and IL-6, two clinically important cytokines. Our study had

dramatic variation in IL-6 that was therapeutically induced, and variation in IFN due to

the disease status of the SLE patients. This, together with the structured study design

with repeat measurements of gene expression across different conditions in the same

individual, allowed us to identify *in vivo* eQTL interactions.

eQTL interactions with drug interventions or other therapeutically relevant physiologic

variables are important to identify. We note that these results are independent of

differentially expressed genes, which are more likely to represent second order effects,

rather than primary genetic effects. Therefore, these eQTL interactions can point to

regulatory mechanisms, such as transcription factors or subclasses of enhancers,

acting downstream of the environmental condition of interest and driving groups of

eQTL interactions. The IFN status eQTL interactions we identified provide support for

this approach. By making use of the direction of effect for the eQTL interaction, we were

able to identify an enrichment of magnifying eQTL interaction SNPs interrupting the

binding sites of transcription factors known to be important in the response to IFN, such

as ISGF3 (the STAT1, STAT2 and IRF9 complex), which binds ISRE. Once we are able

to recognize the downstream drivers of therapeutically relevant clinical variables, then it

may become possible to define more mechanisms of action for drugs and more precise

drug targets.

As a powerful example, we note enrichment of magnifying anti-IL-6-eQTL interaction

SNPs interrupting the binding site of IRF4. It has been suggested that IRF4 works

downstream of IL-6 by binding BATF and co-ordinately regulating the production of IL10

and other genes[38]. Consistent with this, we observed that the *IL10* eQTL does indeed

interact with presence of anti-IL-6 (**Figure S11**). Previous studies have highlighted a

role for IRF4 in the pathogenesis of autoimmune diseases in mouse and humans. For

example in a murine model of SLE, *IRF4* knockout mice did not develop lupus

nephritis[39]. In humans, IRF4 is associated with RA[40], a disease in which anti-IL-6

treatment has been successful[3]. Our findings provide further support that IRF4 could

be a potential therapeutic target for autoimmune diseases such as RA where anti-IL-6 is

effective[41].

The ability to focus on interactions with specific patient phenotypes might point to key

targets for disease intervention. For example, IFN is a key immunophenotype in SLE

patients, and elevated in SLE compared to healthy controls[30, 31]. The IFN status

immunophenotype is already itself driving interest in therapeutic targets. A recent phase

II clinical trial has shown that an antagonist to the type I IFN receptor, acting upstream

of ISRE, reduced severity of symptoms in SLE. Interestingly, the antagonist was more

effective in the patients with a high baseline IFN status[42]. This example provides a

compelling case study for how understanding master regulators of key disease

phenotypes might lead to promising new therapeutic strategies. We speculate that this

provides a mechanism for stratified medicine for future studies, which may be applicable

to other diseases.

We recognized that computing eQTL interactions requires a robust statistical model that

accounts for genotype, environmental factor, RNA expression levels, repeat

measurements, and technical covariates. We were sensitive to the possibility that pre-

processing and normalisation of these factors could potentially have an impact on our

results. For this reason, we used stringent filtering and examined only variants that were

common and where the minor allele was present for each of the exposure groups. Next,

to confirm enrichment of eQTL interactions, we used a stringent permutation-based

strategy that preserved the distribution of genotypes and corresponding expression

values. Finally, we also utilized a standard normal transformation[43] (**Methods**) and

observed that this had little effect on the primary eQTL analysis (r_s=0.99 for z scores,

Figure S14) and interaction analyses (IFN r_s=0.84, drug r_s=0.76 for z scores, **Figure**

S15), or the observed enrichment over the null in our stringent permutation analysis

(**Figure S16**).

We speculate that drug-eQTL interactions might offer an alternative pharmacogenetic

strategy to assess drug response. For many biologic medications, predictive

pharmacogenetics through typical association studies has been challenging; for

example, studies trying to define genetic or transcriptomic biomarkers of anti-TNF

response have not been successful[44, 45]. An eQTL interaction approach can be used

to define a genotype-aware score reflecting the biological activity that a medication is

having upon an individual, given their allelic combination of multiple genetic markers.

For example, we can define a simple anti-IL-6 exposure score based on 7 anti-IL-6

eQTL interactions with a more stringent FDR (FDR<0.1). This score is based on

assessing whether the expression of the eQTL target gene was more consistent with

the drug exposed or the unexposed state for the corresponding interaction SNP

genotype. Unsurprisingly, we found a difference in drug exposure score between the

unexposed and exposed samples (**Figure S17**) (r_s=0.40, p=2.1x10^{-16}); these differences

reflect the fact that the eQTLs were themselves identified by examining samples with

and without drug exposure. However, while we did not utilize the administered drug

dose to identify drug-eQTL interactions; we observed a significant correlation between

drug dose (10, 50 or 200mg) and drug exposure score (r_s=0.16, p=0.02) in the drug-

exposed samples (**Figure S18**). A simple eQTL interaction score may therefore have

the potential to stratify individuals when assessing response to a medication for

example those with a higher drug exposure score may have a better response to

treatment. Similarly, this score could be correlated with adverse effects to capture

informative gene expression signatures.

We note a limitation of this study is that the drug itself did not achieve its primary

efficacy endpoint of improving SLE outcomes. Hence, while the drug exposure score for

this study tracked with the biological effect of the drug (reducing free IL-6 protein levels),

it might not be useful for SLE specifically. However, such a scoring system could be

implemented easily in most phase III trials for a broad range of therapeutics, where the

numbers of samples are far in excess of this phase II trial, ensuring better powered and

more accurate eQTL-interaction mapping.

Conclusion

We devised a framework for identifying *in vivo* eQTL interactions with therapeutically

relevant variables, exploiting repeat measurements from a clinical trial. We have applied

this approach to demonstrate how downstream regulatory effects of cytokine biology

can be elucidated. This same approach can be applied to a wide range of other

clinically important cytokines, their antagonists, or indeed other targeted biologic

therapies. We speculate that this approach might even be applied to the presence or

absence of disease, or disease activity. However, given the multifaceted nature of

disease effects, interpreting an eQTL interaction in that context might be more

challenging. Modern clinical cohorts and clinical trial data sets with RNA-seq data that

has been collected will make this approach easily applicable on a wide scale.

Methods

Study design

The objectives of this study were to map eQTLs in a cohort of lupus patients and

identify eQTL interactions with environmental perturbations such as drug treatment to

shed light on drug and disease mechanisms. SLE patients were recruited to a phase II

clinical trial to test the efficacy and safety of an IL-6 monoclonal antibody (PF-

04236921). The patient population recruited to this trial have been detailed extensively

by Wallace et al.[8] 183 patients (forming a multi-ethnic cohort) were randomized to

receive three doses of drug (10, 50 or 200mg) or placebo at three time points during the

trial (weeks 0, 8 and 16).

RNA-sequencing

We collected peripheral venous blood samples in PAXgene Blood RNA tubes

(PreAnalytiX GmbH, BD Biosciences) for high-depth RNA-seq profiling at 0, 12, and 24

weeks. We extracted total RNA from blood samples using the PAXgene Blood RNA kit

(Qiagen) at a contract lab using a customized automation method. We assessed the

yield and quality of the isolated RNA using Quant-iT™ RiboGreen® RNA Assay Kit

(ThermoFisher Scientific) and Agilent 2100 Bioanalyzer (Agilent Technologies),

respectively. Following quality assessment, we processed an aliquot of 500-1000 ng of

each RNA with a GlobinClear-Human kit, (ThermoFisher Scientific) to remove globin

mRNA. We then converted RNA samples to cDNA libraries using TruSeq RNA Sample

Prep Kit v2 (Illumina) and sequenced using Illumina HiSeq 2000 sequencers. We

generated an average of 40M 100bp pair-end reads per sample for downstream

analysis.

We successfully obtained 468 RNA-seq profiles from 180 patients. We aligned reads to

the reference genome and quantified gene expression using Subread[46] and

featureCounts[47] respectively. We included genes with at least 10 reads (CPM>0.38)

in at least 32 samples (minimum number of patients with both unexposed and exposed

RNA-seq assays in a drug group) prior to normalization. Following quality control (QC),

we removed 4 samples as outliers. We then normalized 20,253 transcripts using the

trimmed mean of M-values method and the edgeR R package[48]. Expression levels

are presented as $\log_2$(cpm +1) (**Table S6**).

Genotyping

We genotyped 160 individuals across 964,193 variants genome-wide with the Illumina

HumanOmniExpressExome-8v1.2 beadchip. We removed SNPs if they deviated from

Hardy-Weinberg Equilibrium (HWE) ($p < 1 \times 10^{-7}$), had a minor allele frequency <5%,

missingness >2% or a heterozygosity rate greater than 3 standard deviations from the

mean (PLINK[49, 50]). For mapping eQTLs, we removed SNPs on the Y chromosome.

Following QC, we used 608,017 variants for further analysis. We removed one sample

with high missingness and outlying heterozygosity rate from further analysis.

Imputation

We pre-phased the genotypes with SHAPEIT v2[51]. We imputed missing genotypes

and untyped SNPs using Impute2[52] in 5Mb chunks against the 1000 Genomes Phase

3[53] reference panel. To ensure only highly quality genotypes, and to avoid artefacts

that can be induced by imputation uncertainty, we removed SNPs with an info score <1,

MAF<0.05 or HWE $p<1 \times 10^{-7}$ leaving 1,595,793 SNPs for further analysis.

Interferon status

We classified the interferon (IFN) status of each sample at each time point from the

expression of 11 IFN response genes (*HERC5*, *IFI27*, *IRF7*, *ISG15*, *LY6E*, *MX1*, *OAS2*,

OAS3, *RSAD2*, *USP18*, *GBP5*) using TaqMan Low Density Arrays. These 11 genes

were selected by identifying transcripts for which there was both a measureable

response to IFN treatment *in vitro*, as well as differential expression (reduction in

expression level) between baseline and visits with clinical improvement in the BOLD

study[33]. There is no consensus set of genes to determine the IFN status of SLE

patients but these 11 genes do overlap with other published gene sets. For example

4/11 genes are also used in the 7-gene set defined by McBride et al[54] and 9/11 genes

overlap with the 21-gene set defined by Yao et al[55].

The first principal component of the expression of the 11-gene set captured 91.7% of

the variation (**Figure S19**). The distribution of this first principal component is nearly

bimodal with good separation (**Figure 2A**) and we classified samples as high or low IFN

based on this first principal component score. In our dataset, we see excellent

correlations (r_s=0.86-0.98) between the real-time PCR expression and the RNA-seq

expression for these 11 genes (**Figure S20**). The first PC of the IFN signature of RNA-

seq data is also strongly correlated with the first PC of the IFN signature of real-time

PCR (r_s=0.96, **Figure S21**). IFN status was available for 376 samples from 157

subjects.

Drug exposure

Samples were assigned as unexposed (placebo or week 0 samples) or drug exposed

(week 12 and week 24 samples in the drug groups).

Free IL-6 protein levels

We determined free IL-6 protein levels from serum using a commercial sandwich ELISA

selected for binding only free IL-6. The assay was validated according to FDA

biomarker and fit-for purpose guidelines. Free IL-6 protein levels were available for 311

samples from 145 subjects. Since the distribution of IL-6 levels was highly skewed, we

ranked samples in order of IL-6 protein levels and included in the model to identify drug-

eQTL interactions.

Statistical analysis

eQTL and interaction analysis

In total, 157 patients (with 379 RNA-seq samples) had good quality gene expression

and genotyping data for eQTL analysis. All statistical analyses were carried out in R[56].

We defined a *cis* eQTL as the SNP within 250kb upstream of the GENCODE[57]

transcription start site of the gene or 250kb downstream of the transcription end site.

We first applied a linear model for the first available time point (week 0 sample for

n=152, week 12 sample for n=5) to identify each eQTL using the first 25 principal

components of gene expression and the first 5 principal components of genotyping as

covariates.

To select the number of gene expression principal components to include, we counted

the number of eQTL genes identified after incrementally increasing the number of

principal components accounted for in the model from 0 to 50 by increments of five

(**Figure S22**). We selected 25 principal components of gene expression to maximise the

number of eQTL genes detected while minimising the number of principal components

we corrected for. We included 5 principal components of genotyping to account for the

heterogeneity in ethnicity in our cohort (**Figure S23**).

SNPs were encoded as 0, 1 and 2 with respect to the number of copies of the minor

allele. To adjust for multiple testing during eQTL discovery we used a stringent

Bonferroni corrected *p*-value threshold of 8.5×10^{-9} (0.05/ 5,872,001 tests). The

Bonferroni adjustment assumes independence among the tests and we therefore note

that it is a conservative multiple comparisons adjustment.

To map eQTLs using multiple samples for each individual, we applied a random

intercept linear mixed model using the first 25 principal components of gene expression

and the first 5 principal components of genotyping as covariates and patient as a

random effect:

$$E_{i,j} = \theta + \beta_{geno} \cdot g_j + (\kappa_j | i) + \sum_{l=1}^{25} \phi_l \cdot pc_{i,l} + \sum_{m=1}^{5} \gamma_m \cdot pc_{j,m}$$

Where $E_{i,j}$ is gene expression for the i^{th} sample from the j^{th} subject, θ is the intercept,

β_{geno} is the genotype effect (eQTL), $(\kappa_j | i)$ is the random effect for the i^{th} sample from the

j^{th} subject, $pc_{i,l}$ is principal component l of gene expression for sample i, $pc_{j,m}$ is principal

component m of genotyping for subject j.

We fitted the linear mixed models using the lme4 R package[58]. We assumed

covariance between samples from the same individual, but did not assume any

structure in this covariance.

We used the most significant SNP (with $p < 8.5 \times 10^{-9}$) from the 4,818 identified eQTL

genes to explore eQTL interactions. For each environmental interaction analysis, we

further filtered these eQTLs to include only those with at least two individuals

homozygous for the minor allele of the SNP being tested in each of the environmental

factor groups. For example, we required two of these individuals in each of the drug

exposed and drug unexposed groups. To identify eQTL interactions, we added an

additional covariate to the model for example drug exposure, and an interaction term

between this covariate and the genotype of the SNP:

$$E_{i,j} = \theta + \beta_{geno} \cdot g_j + (\kappa_j | i) + \sum_{l=1}^{25} \phi_l \cdot pc_{i,l} + \sum_{m=1}^{5} \gamma_m \cdot pc_{j,m} +$$
       $$\beta_{drug} \cdot d_i + \beta_x \cdot d_i \cdot g_j$$

Where $E_{i,j}$ is gene expression for the i^{th} sample from the j^{th} subject, θ is the intercept,

β_{geno} is the genotype effect (eQTL), $(\kappa_j | i)$ is the random effect for the i^{th} sample from the

j^{th} subject, $pc_{i,l}$ is principal component l of gene expression for sample i, $pc_{j,m}$ is principal

component m of genotyping for subject j, β_{drug} is the drug effect (differential gene

expression) and β_x is the interaction effect.

We determined the significance of the interaction term with a likelihood ratio test.

To rigorously confirm the relative enrichment of eQTL interactions, we shuffled the

interaction covariate (for example drug exposure) 1,000 times and calculated the

number of significant interactions observed in each permutation. Our primary goal for

the permutation analysis was to retain the main eQTL effect while examining only the

effect of the environmental factor on the interaction. In this study, the main purpose of

the covariates included in the model is to ensure the main eQTL effect is found. For IFN

high/low status, we shuffled across all samples. For drug interaction permutation

analysis, we maintained the number of individuals in the drug group and the number of

samples with exposure to drug. We calculated a qvalue for each interaction using the q

value package[34]. **Figure S24** shows the observed versus the expected p values for

the interaction analyses.

The expression of the majority of genes followed a normal distribution (**Figure S25**) but

to assess whether non-normality could be causing an inflation of our test statistic, we

repeated the identification of eQTLs and eQTL interactions following the standard

normal transformation. We transformed the expression values of each gene to their

respective quantiles of a normal distribution using the qqnorm function in R, breaking

any ties (for example expression levels of zero in some individuals) randomly.

Concordance with an eQTL study in healthy individuals

In the SLE cohort, we classified 4,818 *cis* eQTL genes (p<8.5×10^{-9}). The *z*-score for the

most associated SNP for each of these genes was compared to the *z*- score from a

previously published eQTL dataset from whole blood from 2,166 healthy individuals[11].

4,113/4818 SNP-gene pairs (85.4%) were also reported in the BIOS dataset

(FDR<0.05). After removing 301 SNPs, which could not be mapped to a strand

3,770/3,812 (98.9%) had a *z*-score (eQTL effect) in a consistent direction.

Magnifiers and Dampeners

An eQTL interaction can either magnify or dampen the original eQTL effect. We

multiplied the interaction *z*-score by the sign of the original eQTL effect (genotype beta)

and defined magnifiers as interactions with an adjusted *z*-score > 0 and dampeners as

interactions with an adjusted *z*-score < 0.

Differential gene expression analysis

To identify differentially expressed genes following drug exposure (unexposed or

exposed), we applied a random intercept linear mixed model using the first 25 principal

components of gene expression and the first 5 principal components of genotyping as

covariates and patient as a random effect. We calculated a q value using the q value

package[34].

Drug exposure score

We assigned a drug exposure score to each sample. We calculated a score for each

gene (see equation below) and then averaged across the 7 drug-eQTL genes

(FDR<0.1) to give the final drug exposure score.

$$Drug\ exposure\ score\ for\ gene\ = \frac{1}{2}(\frac{G - G_{Unexp}}{SE})^2 - \frac{1}{2}(\frac{G - G_{Exp}}{SE})^2$$

Where G is gene expression for a given sample, G_{Unexp} is predicted mean gene

expression for unexposed samples of the relevant SNP genotype, G_{Exp} is predicted

mean gene expression for exposed samples of the relevant SNP genotype and SE is

standard error for the intercept term of the model (unexposed expression for genotype

0).

HOMER analysis for transcription factor binding motif enrichment

We used the HOMER software suite[35] to look for enrichment of transcription factor

binding motifs in the 210 IFN-eQTL interactions (FDR<0.2) and the 72 drug-eQTL

interactions (FDR<0.2). Each eQTL interaction was identified using the most highly

associated SNP for that eQTL. However, as this SNP is not necessarily the functional

SNP, we additionally considered all those with an $r^2 \geq 0.8$ in the 1000 Genomes

European population[53] within the *cis* eQTL window. We defined our motif search

window as 20 bp on either side of each SNP (i.e. 41 bp wide).

For each environmental factor, we divided the eQTL interactions into magnifiers or

dampeners and conducted two separate HOMER analyses: one with magnifiers in the

foreground and dampeners in the background; the other with dampeners in the

foreground and magnifiers in the background. HOMER reported the transcription factor

motifs that were significantly enriched in the foreground relative to background. Motifs

were plotted using the SeqLogo R library[59].

We determined permutation p values for enrichment of the ISRE and IRF4 transcription

factor binding sites as follows. For ISRE, the motif is interrupted by interaction SNPs (or

SNPs in LD) corresponding to 11 magnifying genes and 1 dampening genes. We

permuted which genes were labelled as magnifiers or dampeners 100,000 times and

counted the number of genes in each category with an ISRE motif interrupted. We

found 1,855 occurrences from 100,000 trials with at least 11 magnifying genes

(p<0.019). For IRF4 the motif is interrupted by SNPs corresponding to 9 magnifying

genes and 4 dampening genes. Using the same permutation approach, we found 5,801

occurrences from 100,000 trials with at least 9 magnifying genes (p<0.058).

Declarations

Ethics approval and consent to participate

The protocol was approved by each institutional review board subject to applicable laws

and regulations and ethical principles consistent with the Declaration of Helsinki.

Availability of data and material

Data are available in the supplementary tables and at

http://baohongz.github.io/Lupus_eQTL.

Acknowledgements

This work is supported in part by funding from the National Institutes of Health

(U01GM092691, UH2AR067677, U19AI111224 (SR)), the Doris Duke Charitable

Foundation Grant #2013097, the Ruth L. Kirschstein National Research Service Award

(F31AR070582) from the National Institute of Arthritis and Musculoskeletal and Skin

Diseases (KS) and the Rheumatology Research Foundation Tobé and Stephen E.

Malawista, MD Endowment in Academic Rheumatology (D.A.R). This work is also

supported by unrestricted funding from Pfizer, Inc.

Author Contributions

The project was conceived and designed by EED, MSV, BZ and SR. Statistical analysis

was conducted by EED, TA, MG-A, KS, H-JW, YL and CS. Molecular data was

obtained, organized and analysed by YZ, SP, DvS, JSB, NB, MSV, BZ and DAR. The

initial manuscript was written by EED and SR. All authors edited and approved the

manuscript.

Competing Interests

The authors declare that they have no competing interests.

References

1. Rider P, Carmi Y, Cohen I. Biologics for Targeting Inflammatory Cytokines, Clinical
Uses, and Limitations. Int J Cell Biol. 2016;2016 iv.
2. Cessak G, Kuzawińska O, Burda A, Lis K, Wojnar M, Mirowska-Guzel D, et al. TNF
inhibitors - Mechanisms of action, approved and off-label indications. Pharmacol
Reports. 2014;66:836–44.
3. Tanaka Y, Mola EM. IL-6 targeting compared to TNF targeting in rheumatoid arthritis:
studies of olokizumab, sarilumab and sirukumab. Ann Rheum Dis. 2014;73:1595–7.
4. Stone JH, Tuckwell K, Dimonaco S, Klearman M, Aringer M, Blockmans D, et al. Trial
of Tocilizumab in Giant-Cell Arteritis. N Engl J Med. 2017;377:317–28.
doi:10.1056/NEJMoa1613849.
5. Van Rhee F, Wong RS, Munshi N, Rossi JF, Ke XY, Fosså A, et al. Siltuximab for
multicentric Castleman's disease: A randomised, double-blind, placebo-controlled trial.
Lancet Oncol. 2014;15:966–74.
6. Tackey E, Lipsky PE, Illei GG. Rationale for interleukin-6 blockade in systemic lupus
erythematosus. Lupus. 2004;13:339–43.
7. Illei GG, Shirota Y, Yarboro CH, Daruwalla J, Tackey E, Takada K, et al. Tocilizumab
in systemic lupus erythematosus: Data on safety, preliminary efficacy, and impact on
circulating plasma cells from an open-label phase I dosage-escalation study. Arthritis
Rheum. 2010;62:542–52.
8. Wallace DJ, Strand V, Merrill JT, Popa S, Spindler AJ, Eimon A, et al. Efficacy and
safety of an interleukin 6 monoclonal antibody for the treatment of systemic lupus
erythematosus: a phase II dose-ranging randomised controlled trial. Ann Rheum Dis.
2017;76:534–42. doi:10.1136/annrheumdis-2016-209668.
9. Bronson PG, Chaivorapol C, Ortmann W, Behrens TW, Graham RR. The genetics of
type I interferon in systemic lupus erythematosus. Curr Opin Immunol. 2012;24:530–7.
doi:10.1016/j.coi.2012.07.008.

10. Ridker PM, Everett BM, Thuren T, MacFadyen JG, Chang WH, Ballantyne C, et al.
Antiinflammatory Therapy with Canakinumab for Atherosclerotic Disease. N Engl J Med.
2017;377:119–1131. doi:10.1056/NEJMoa1707914.
11. Zhernakova D V, Deelen P, Vermaat M, van Iterson M, van Galen M, Arindrarto W,
et al. Identification of context-dependent expression quantitative trait loci in whole blood.
Nat Genet. 2017;49 November:139–45. doi:10.1038/ng.3737.
12. Fairfax BP, Humburg P, Makino S, Naranbhai V, Wong D, Lau E, et al. Innate
Immune Activity Conditions the Effect of Regulatory Variants upon Monocyte Gene
Expression. Science. 2014;343:1246949. doi:10.1126/science.1246949.
13. Fairfax BP, Makino S, Radhakrishnan J, Plant K, Leslie S, Dilthey A, et al. Genetics
of gene expression in primary immune cells identifies cell type-specific master
regulators and roles of HLA alleles. Nat Genet. 2012;44:502–10. doi:10.1038/ng.2205.
14. Raj T, Rothamel K, Mostafavi S, Ye C, Lee MN, Replogle JM, et al. Polarization of
the Effects of Autoimmune and Neurodegenerative Risk Alleles in Leukocytes. Science.
2014;344:519–23.
15. Ardlie KG, Deluca DS, Segre A V., Sullivan TJ, Young TR, Gelfand ET, et al. The
Genotype-Tissue Expression (GTEx) pilot analysis: Multitissue gene regulation in
humans. Science. 2015;348:648–60. doi:10.1126/science.1262110.
16. Nica AC, Parts L, Glass D, Nisbet J, Barrett A, Sekowska M, et al. The architecture
of gene regulatory variation across multiple human tissues: the MuTHER study. PLoS
Genet. 2011;7:e1002003. doi:10.1371/journal.pgen.1002003.
17. Kukurba KR, Parsana P, Balliu B, Smith KS, Zappala Z, Knowles DA, et al. Impact
of the X chromosome and sex on regulatory variation. Genome Res. 2016;26:768–77.
18. Buil A, Brown AA, Lappalainen T, Viñuela A, Davies MN, Zheng H-F, et al. Gene-
gene and gene-environment interactions detected by transcriptome sequence analysis
in twins. Nat Genet. 2015;47:88–91. doi:10.1038/ng.3162.
19. Maranville JC, Luca F, Stephens M, Di Rienzo A. Mapping gene-environment
interactions at regulatory polymorphisms: insights into mechanisms of phenotypic
variation. Transcription. 2012;3:56–62. doi:10.4161/trns.19497.
20. Idaghdour Y, Awadalla P. Exploiting gene expression variation to capture gene-
environment interactions for disease. Front Genet. 2013;4 MAY:1–7.
21. Idaghdour Y, Quinlan J, Goulet J-P, Berghout J, Gbeha E, Bruat V, et al. Evidence
for additive and interaction effects of host genotype and infection in malaria. Proc Natl
Acad Sci U S A. 2012;109:16786–93. doi:10.1073/pnas.1204945109.
22. Idaghdour Y, Czika W, Shianna K V, Lee SH, Visscher PM, Martin HC, et al.
Geographical genomics of human leukocyte gene expression variation in southern
Morocco. Nat Genet. 2010;42:62–7. doi:10.1038/ng.495.
23. Peters JE, Lyons PA, Lee JC, Richard AC, Fortune MD, Newcombe PJ, et al.
Insight into Genotype-Phenotype Associations through eQTL Mapping in Multiple Cell
Types in Health and Immune-Mediated Disease. PLoS Genet. 2016;12:e1005908.
doi:10.1371/journal.pgen.1005908.
24. Li Y, Álvarez OA, Gutteling EW, Tijsterman M, Fu J, Riksen JAG, et al. Mapping
determinants of gene expression plasticity by genetical genomics in C. elegans. PLoS
Genet. 2006;2:2155–61.
25. Smith EN, Kruglyak L. Gene-environment interaction in yeast gene expression.
PLoS Biol. 2008;6:810–24.

26. Maranville JC, Luca F, Richards AL, Wen X, Witonsky DB, Baxter S, et al.
Interactions between glucocorticoid treatment and Cis-regulatory polymorphisms
contribute to cellular response phenotypes. PLoS Genet. 2011;7:e1002162.
27. Barreiro LB, Tailleux L, Pai AA, Gicquel B, Marioni JC, Gilad Y. Deciphering the
genetic architecture of variation in the immune response to Mycobacterium tuberculosis
infection. Proc Natl Acad Sci U S A. 2012;109:1204–9.
28. Smirnov D, Morley M, Shin E. Genetic analysis of radiation-induced changes in
human gene expression. Nature. 2009;459:587–91. doi:10.1038/nature07940.Genetic.
29. Walsh AM, Whitaker JW, Huang CC, Cherkas Y, Lamberth SL, Brodmerkel C, et al.
Integrative genomic deconvolution of rheumatoid arthritis GWAS loci into gene and cell
type associations. Genome Biol. 2016;17:79. doi:10.1186/s13059-016-0948-6.
30. Baechler EC, Batliwalla FM, Karypis G, Gaffney PM, Ortmann W a, Espe KJ, et al.
Interferon-inducible gene expression signature in peripheral blood cells of patients with
severe lupus. Proc Natl Acad Sci U S A. 2003;100:2610–5.
31. Bennett L, Palucka a K, Arce E, Cantrell V, Borvak J, Banchereau J, et al.
Interferon and granulopoiesis signatures in systemic lupus erythematosus blood. J Exp
Med. 2003;197:711–23.
32. Stranger BE, Montgomery SB, Dimas AS, Parts L, Stegle O, Ingle CE, et al.
Patterns of Cis regulatory variation in diverse human populations. PLoS Genet. 2012;8.
33. Hill AA, Immermann FW, Zhang Y, Reddy PS, Zhou T, O'Toole M, et al. FRI0003
Determination of interferon (IFN) signatures for sle patients may be critical for optimal
treatment selection but depends on the genes chosen: report from the bold (biomarkers
of lupus disease) study. Ann Rheum Dis. 2013;72:A369–70. doi:10.1136/annrheumdis-
2013-eular.1131.
34. Storey JD, Tibshirani R. Statistical significance for genomewide studies. Proc Natl
Acad Sci. 2003;100:9440–5. doi:10.1073/pnas.1530509100.
35. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple
Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory
Elements Required for Macrophage and B Cell Identities. Mol Cell. 2010;38:576–89.
doi:10.1016/j.molcel.2010.05.004.
36. Au-Yeung N, Mandhana R, Horvath CM. Transcriptional regulation by STAT1 and
STAT2 in the interferon JAK-STAT pathway. Jak-Stat. 2013;2:e23931.
doi:10.4161/jkst.23931.
37. Westra H-J, Peters MJ, Esko T, Yaghootkar H, Schurmann C, Kettunen J, et al.
Systematic identification of trans eQTLs as putative drivers of known disease
associations. Nat Genet. 2013;45:1238–43. doi:10.1038/ng.2756.
38. Koch S, Mousset S, Graser A, Reppert S, Übel C, Reinhardt C, et al. IL-6 activated
integrated BATF/IRF4 functions in lymphocytes are T-bet-independent and reversed by
subcutaneous immunotherapy. Sci Rep. 2013;3:1754. doi:10.1038/srep01754.
39. Lech M, Weidenbusch M, Kulkarni OP, Ryu M, Darisipudi MN, Susanti HE, et al.
IRF4 Deficiency Abrogates Lupus Nephritis Despite Enhancing Systemic Cytokine
Production. J Am Soc Nephrol. 2011;22:1443–52. doi:10.1681/ASN.2010121260.
40. Okada Y, Wu D, Trynka G, Raj T, Terao C, Ikari K, et al. Genetics of rheumatoid
arthritis contributes to biology and drug discovery. Nature. 2014;506:376–81.
doi:10.1038/nature12873.
41. Xu WD, Pan HF, Ye DQ, Xu Y. Targeting IRF4 in autoimmune diseases. Autoimmun

Rev. 2012;11:918–24. doi:10.1016/j.autrev.2012.08.011.

42. Furie R, Merrill J, Werth V, Khamashta M, Kalunian K, Brohawn P, et al.
Anifrolumab, an Anti-Interferon Alpha Receptor Monoclonal Antibody, in Moderate to
Severe Systemic Lupus Erythematosus (SLE). Arthritis Rheumatol. 2017;69:376–86.

43. Lappalainen T, Sammeth M, Friedländer MR, 't Hoen PAC, Monlong J, Rivas MA, et
al. Transcriptome and genome sequencing uncovers functional variation in humans.
Nature. 2013;501:506–11. doi:10.1038/nature12531.

44. Cui J, Stahl EA, Saevarsdottir S, Miceli C, Diogo D, Trynka G, et al. Genome-wide
association study and gene expression analysis identifies CD84 as a predictor of
response to etanercept therapy in rheumatoid arthritis. PLoS Genet. 2013;9:e1003394.
doi:10.1371/journal.pgen.1003394.

45. Cui J, Diogo D, Stahl EA, Canhao H, Mariette X, Greenberg MPH JD, et al. The role
of rare protein-coding variants to anti-TNF treatment response in rheumatoid arthritis.
Arthritis Rheumatol. 2017;69:735–41.

46. Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable read
mapping by seed-and-vote. Nucleic Acids Res. 2013;41:e108–e108.
doi:10.1093/nar/gkt214.

47. Liao Y, Smyth GK, Shi W. FeatureCounts: An efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 2014;30:923–30.

48. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics.
2010;26:139–40. doi:10.1093/bioinformatics/btp616.

49. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation
PLINK: rising to the challenge of larger and richer datasets. Gigascience. 2015;4:7.
doi:10.1186/s13742-015-0047-8.

50. Purcell S, Chang C. PLINK 1.9. https://www.cog-genomics.org/plink2.

51. Delaneau O, Marchini J, Zagury J-F. A linear complexity phasing method for
thousands of genomes. Nat Methods. 2011;9:179–81. doi:10.1038/nmeth.1785.

52. Howie BN, Donnelly P, Marchini J. A Flexible and Accurate Genotype Imputation
Method for the Next Generation of Genome-Wide Association Studies. PLoS Genet.
2009;5:e1000529. doi:10.1371/journal.pgen.1000529.

53. Auton A, Abecasis GR, Altshuler DM, Durbin RM, Abecasis GR, Bentley DR, et al. A
global reference for human genetic variation. Nature. 2015;526:68–74.
doi:10.1038/nature15393.

54. McBride JM, Jiang J, Abbas AR, Morimoto A, Li J, Macluca R, et al. Safety and
pharmacodynamics of rontalizumab in patients with systemic lupus erythematosus:
Results of a phase I, placebo-controlled, double-blind, dose-escalation study. Arthritis
Rheum. 2012;64:3666–76.

55. Yao Y, Higgs BW, Morehouse C, de Los Reyes M, Trigona W, Brohawn P, et al.
Development of Potential Pharmacodynamic and Diagnostic Markers for Anti-IFN-α
Monoclonal Antibody Trials in Systemic Lupus Erythematosus. Hum Genomics
Proteomics. 2009;2009:374312-. doi:10.4061/2009/374312.

56. R Core Team. R: A Language and Environment for Statistical Computing. 2015.
https://www.r-project.org.

57. Harrow J, Frankish A, Gonzalez JM, Tapanari E, Diekhans M, Kokocinski F, et al.
GENCODE: the reference human genome annotation for The ENCODE Project.

Genome Res. 2012;22:1760–74. doi:10.1101/gr.135350.111.

58. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using

lme4. J Stat Softw. 2015;67:1–48. doi:10.18637/jss.v067.i01.

59. Bembom O. seqLogo: Sequence logos for DNA sequence alignments. 2016.

Figure 1. Identifying eQTLs in SLE patients (A) Clinical trial structure and sampling strategy. (B) Number of eQTL genes identified using a linear model (left) and a linear mixed model (right). For the linear model, we used the first available time point for each individual (week 0 sample for n=152, week 12 sample for n=5). (C) Volcano plot of eQTL effects for the most significantly associated SNP for each gene (red color indicates $p<8.5\times10^{-9}$). (D) Concordance of SLE eQTL effects ($p<8.5\times10^{-9}$) with eQTLs observed in the BIOS cohort[11] of healthy individuals (FDR <0.05). Each point represents the most significant SNP-gene pair for the SLE eQTL.

Figure 2

Figure 2. eQTL interactions with IFN status (**A**) Designation of IFN status for each sample from the real-time PCR expression of 11 genes (first principal component). (**B**) IFN status interaction with the *SLFN5* eQTL plotted with respect to rs12602407 genotype (left) and IFN status of the sample (right). (**C**) The ISRE motif enriched among eQTLs magnified in IFN high samples. Arrows indicate positions of the motif interrupted by interaction SNPs (or SNPs in strong LD). Red indicates these SNPs correspond to magnified eQTLs. (**D**) IFN status interaction with the *GTF2A2* eQTL plotted with respect to rs2306355 genotype (left) and IFN status of the sample (right).

Figure 3. eQTL interactions with drug exposure (A) Drug exposure interaction with the *CLEC18A* eQTL plotted with respect to rs3192882 genotype (left) and drug exposure (right). (B) The IRF4 motif enriched among eQTLs magnified following drug treatment. Arrows indicate positions of the motif interrupted by interaction SNPs (or SNPs in strong LD). Red and blue indicate SNPs corresponding to magnified and dampened eQTLs respectively. (C) Concordance of free IL-6 protein interaction effects with drug exposure interaction effects (grey indicates expected opposite direction).