1

Conserved microRNA targeting reveals preexisting gene dosage sensitivities that shaped amniote sex chromosome evolution

Sahin Naqvi[1, 2], Daniel W. Bellott[1], Kathy S. Lin[1, 3], & David C. Page[1, 2, 4]

[1] Whitehead Institute, Cambridge MA 02142;
[2] Department of Biology, Massachusetts Institute of Technology, Cambridge MA 02139
[3] Program in Computational and Systems Biology, Massachusetts Institute of Technology, Cambridge MA 02139
[4] Howard Hughes Medical Institute, Whitehead Institute, Cambridge MA 02142

Mammalian X and Y chromosomes evolved from an ordinary autosomal pair. Genetic

decay of the Y led to X chromosome inactivation (XCI) in females, but some Y-linked genes

were retained during the course of sex chromosome evolution, and many X-linked genes

did not become subject to XCI. We reconstructed gene-by-gene dosage sensitivities on the

ancestral autosomes through phylogenetic analysis of microRNA (miRNA) target sites and

compared these preexisting characteristics to the current status of Y-linked and X-linked

genes in mammals. Preexisting heterogeneities in dosage sensitivity, manifesting as

differences in the extent of miRNA-mediated repression, predicted either the retention of a

Y homolog or the acquisition of XCI following Y gene decay. Analogous heterogeneities

among avian Z-linked genes predicted the retention of a W homolog. Genome-wide

analyses of human copy number variation indicate that these heterogeneities consisted of

sensitivity to both increases and decreases in dosage. We propose a model of XY/ZW

evolution incorporating such preexisting dosage sensitivities in determining the

evolutionary fates of individual genes. Our findings thus provide a more complete view of

the role of dosage sensitivity in shaping the mammalian and avian sex chromosomes, and

reveal an important role for post-transcriptional regulatory sequences (miRNA target sites)

in sex chromosome evolution.

INTRODUCTION

The mammalian X and Y chromosomes evolved from a pair of ordinary autosomes over the past

300 million years (Lahn & Page, 1999). Only 3% of genes on the ancestral pair of autosomes

survive on the human Y chromosome (Bellott et al., 2010; Skaletsky et al., 2003), compared to

98% on the X chromosome (Mueller et al., 2013). In females, one copy of the X chromosome is

silenced by X inactivation (XCI); this silencing evolved on a gene-by-gene basis following Y

gene loss and compensatory X upregulation (Jegalian & Page, 1998; Ross et al., 2005), and some

genes escape XCI in humans (Carrel & Willard, 2005) and other mammals (Yang, Babak,

Shendure, & Disteche, 2010). In parallel, the avian Z and W sex chromosomes evolved from a

different pair of autosomes than the mammalian X and Y chromosomes (Bellott et al., 2010;

Nanda et al., 1999; Ross et al., 2005). Decay of the female-specific W chromosome was

similarly extensive, but birds did not evolve a large-scale inactivation of Z-linked genes

analogous to XCI in mammals (Itoh et al., 2007; Mank & Ellegren, 2009; Uebbing et al., 2015;

Wright, Zimmer, Harrison, & Mank, 2015). Thus, genes previously found on the ancestral

autosomes that gave rise to the mammalian or avian sex chromosomes have undergone

significant changes in gene dosage. In modern mammals, these molecular events have resulted in

three classes of ancestral X-linked genes representing distinct evolutionary fates: those with a

surviving Y homolog, those with no Y homolog and subject to XCI, and those with no Y

homolog but escaping XCI. In birds, two classes of ancestral Z-linked genes have arisen: those

with or without a W homolog. Identifying gene-by-gene properties that distinguish classes of X-

and Z-linked genes is thus crucial to understanding the selective pressures underlying the

molecular events of mammalian and avian sex chromosome evolution.

Emerging evidence suggests a role for gene dosage sensitivity in mammalian and avian

sex chromosome evolution. X- and Z-linked genes with surviving homologs on the mammalian

Y or avian W chromosomes are enriched for important regulatory functions and predictors of

haploinsufficiency compared to those lacking Y or W homologs (Bellott et al., 2014, 2017);

similar observations have been made in fish (White, Kitano, & Peichel, 2015) and Drosophila

(Kaiser, Zhou, & Bachtrog, 2011). Human X- and chicken Z-linked genes that show the

strongest signatures of dosage compensation in either lineage also show signs of dosage

sensitivity as measured by membership in large protein complexes (Pessia, Makino, Bailly-

Bechet, McLysaght, & Marais, 2012) or evolutionary patterns of gene duplication and retention

(Zimmer, Harrison, Dessimoz, & Mank, 2016). Despite these advances, little is known regarding

selective pressures resulting from sensitivity to dosage increases, as these studies either focused

on haploinsufficiency or employed less direct predictors of dosage sensitivity. Furthermore, it is

not known whether heterogeneities in dosage sensitivity among classes of sex-linked genes were

acquired during sex chromosome evolution, or predated the emergence of sex chromosomes, as

there has been no explicit, systematic reconstruction of dosage sensitivity on the ancestral

autosomes that gave rise to the mammalian and avian sex chromosomes.

To assess the role of preexisting dosage sensitivities in XY and ZW evolution, we sought

to employ a measure of dosage sensitivity that could be 1) demonstrably informative with respect

to sensitivity to dosage increases, and 2) explicitly reconstructed on the ancestral autosomes. We

focused on regulation by microRNAs (miRNAs), small noncoding RNAs that function as tuners

of gene dosage by lowering target mRNA levels through pairing to the 3` untranslated region

(UTR) (Bartel, 2009). The repressive nature of miRNA targeting is informative with respect to

sensitivity to dosage increases, allowing for a more complete understanding of the role of dosage

sensitivity in sex chromosome evolution. Both miRNAs themselves and their complementary

target sites can be preserved over millions of years of vertebrate evolution, facilitating the

reconstruction of miRNA targeting on the ancestral autosomes through cross-species sequence

alignments. As miRNA targeting occurs post-transcriptionally, reconstruction of its ancestral

state is decoupled from transcriptional regulatory mechanisms such as XCI that evolved

following X-Y differentiation.

First, through genome-wide analysis of human copy number variation, we established

conserved miRNA targeting as an indicator of sensitivity to dosage increases, which we used to

infer heterogeneities in dosage sensitivity between the classes of present-day X- and Z-linked

genes. We then reconstructed miRNA target sites on the ancestral autosomes, finding significant

heterogeneity in the extent to which the autosomal precursors of X- or Z-linked genes were

targeted by miRNAs. Finally, we reanalyzed publically available experimental datasets to

validate that this conserved miRNA targeting is operative in living cells. Combined with results

from previous studies, these findings provide the foundation for a model of X-Y and Z-W

evolution in which both survival on the sex-specific Y or W chromosome and the subsequent

evolution of XCI in mammals were determined by preexisting sensitivities to both decreases and

increases in gene dosage. Our findings also demonstrate distinct selective pressures acting on

post-transcriptional regulatory sequences that functioned to tune gene dosage both prior to and

during amniote sex chromosome evolution.

RESULTS

Analysis of human copy number variation indicates conserved microRNA targeting of

genes sensitive to dosage increases

We first sought to determine whether conserved targeting by microRNAs (miRNAs) correlates

with sensitivity to dosage increases across the human genome. To estimate pressure to maintain

miRNA targeting, we used published probabilities of conserved targeting (P_{CT} scores) for each

gene-miRNA interaction in the human genome. The P_{CT} score reflects an estimate of the

probability that a given gene-miRNA interaction is conserved due to miRNA targeting, obtained

by calculating the conservation of the relevant miRNA target sites relative to the conservation of

the entire 3` UTR (Friedman, Farh, Burge, & Bartel, 2009). In this manner, the P_{CT} score

intrinsically controls for differences in 3` UTR background conservation and sequence

composition due to differing rates of expression divergence and/or sequence evolution, making it

comparable across both genes and miRNA families. We refer to these P_{CT} scores as "miRNA

conservation scores" in the remainder of the text.

A recent study reported a correlation between these miRNA conservation scores and

predicted haploinsufficiency (Pinzón et al., 2016), indicating that conserved miRNA targeting

broadly corresponds to dosage sensitivity. However, such a correlation does not isolate the

effects of sensitivity to dosage increases, which we expect to be particularly important in the

context of miRNA targeting. We reasoned that genes for which increases in dosage are

deleterious should be depleted from the set of observed gene duplications in healthy human

individuals. We used a catalogue of rare genic copy number variation among 59,898 control

human exomes (Exome Aggregation Consortium, ExAC)(Ruderfer et al., 2016) to classify

autosomal protein-coding genes as exhibiting or lacking duplication or deletion in healthy

individuals (see Methods). We compared duplicated and non-duplicated genes with the same

deletion status in order to control for differences in sensitivity to underexpression. We found that

non-duplicated genes have significantly higher miRNA conservation scores than duplicated

genes, irrespective of deletion status (Figure 1A,B). Non-deleted genes also have significantly

higher scores than deleted genes irrespective of duplication status (Supplemental Figure S1), but

duplication status has a greater effect on miRNA conservation scores than does deletion status

(blue vs. orange boxes, Figure 1C). Thus, conserved miRNA targeting is a feature of genes

sensitive to changes in gene dosage in humans and is especially informative with regards to

sensitivity to dosage increases.

Figure 1: Conserved miRNA targeting of autosomal genes stratified by copy number variation in 59,898 human exomes. Probabilities of conserved targeting (P_{CT}) of all gene-miRNA interactions involving non-duplicated and duplicated genes, further stratified as (A) deleted (grey, n = 69,339 interactions from 4,118 genes; blue, n = 80,290 interactions from 3,976 genes) or (B) not deleted (orange, n = 51,514 interactions from 2,916 genes; purple, n = 72,826 interactions from 3,510 genes). *** p < 0.001, two-sided Kolmogorov-Smirnov test. (C) Mean gene-level P_{CT} scores. ** p < 0.01, *** p < 0.001, two-sided Wilcoxon rank-sum test.

X-Y pairs and X-inactivated genes have higher miRNA conservation scores than X escape

genes

We next assessed whether the three classes of X-linked genes differ with respect to dosage

sensitivity as inferred by conserved miRNA targeting. To delineate these classes, we began with

the set of ancestral genes reconstructed through cross-species comparisons of the human X

chromosome and orthologous chicken autosomes (Bellott et al., 2014, 2017, 2010; Hughes et al.,

2012; Mueller et al., 2013). We designated ancestral X-linked genes with a surviving human Y

homolog (Skaletsky et al., 2003) as X-Y pairs and also considered the set of X-linked genes with

a surviving Y homolog in any of eight mammals (Bellott et al., 2014) to increase the

phylogenetic breadth of findings regarding X-Y pairs. A number of studies have catalogued the

inactivation status of X-linked genes in various human tissues and cell-types. We used a meta-

analysis that combined results from three studies by assigning a "consensus" X-inactivation

status to each gene (Balaton, Cotton, & Brown, 2015) to designate the remainder of ancestral

genes lacking a Y homolog as subject to or escaping XCI. In summary, we classified genes as

either: 1) X-Y pairs, 2) lacking a Y homolog and subject to XCI (X-inactivated), or 3) lacking a

Y homolog but escaping XCI (X escape).

We found that human X-Y pairs have the highest miRNA conservation scores, followed

by X-inactivated and finally X escape genes (Figure 2A,B). The expanded set of X-Y pairs

across eight mammals also has significantly higher miRNA conservation scores than ancestral X-

linked genes with no Y homolog (Supplemental Figure S2). Observed differences between

miRNA conservation scores are not driven by distinct subsets of genes in each class, as indicated

by gene resampling with replacement (Supplemental Figure S3), and the decrease in miRNA

conservation scores of X escape genes relative to X-inactivated genes and X-Y pairs is not

driven by genes that escape XCI variably across individuals (Supplemental Figure S4). We also

verified that these differences were not driven by artificially inflated or deflated conservation

scores of certain target sites due to non-uniformity in 3` UTR conservation (Methods,

Supplemental Figure S5).

Finally, we assessed whether miRNA conservation scores distinguish the three classes by

providing additional information not accounted for by known factors (Bellott et al., 2014)

influencing evolutionary outcomes. We used logistic regression to model, for each gene, the

probability of falling into each of the three classes (X-Y pair, X-inactivated, or X escape) as a

linear combination of haploinsufficiency probability (pHI) (Huang, Lee, Marcotte, & Hurles,

2010), human expression breadth (GTEx Consortium, 2015), and purifying selection, measured

by the ratio of non-synonymous to synonymous substitution rates (dN/dS) between human and

mouse orthologs (Yates et al., 2016). Adding miRNA conservation as an additional predictor

resulted in a better model fit as measured by Aikake's information criterion (AIC), which

quantifies goodness of fit while also penalizing increases in model complexity (known factors

only, AIC 327.7; known factors plus miRNA conservation, AIC 321.5; lower AIC indicates

better model). Based on our analyses of autosomal copy number variation (Figure 1), we

attribute this improvement in model fit to the fact that miRNA conservation scores are most

informative with regards to sensitivity to dosage increases. Thus, there is significant

heterogeneity in dosage sensitivity, both upwards and downwards, among the three classes of

ancestral X-linked genes: X-Y pairs are the most dosage-sensitive, while X-inactivated genes are

of intermediate dosage sensitivity, and X escape genes are the least dosage-sensitive.

196
Figure 2. X-Y pairs and X-inactivated genes have higher miRNA conservation scores than
X escape genes. P_{CT} score distributions of all gene-miRNA interactions involving (A) human X-
Y pairs (n = 371 interactions from 15 genes), X-inactivated genes (n = 6,743 interactions from
329 genes), and X escape genes (n = 1,037 interactions from 56 genes). ** $p < 0.01$, two-sided
Kolmogorov-Smirnov test. (B) Mean gene-level P_{CT} scores. * $p < 0.05$, ** $p < 0.01$, two-sided
Wilcoxon rank-sum test.

203
204
205
Heterogeneities in X-linked miRNA targeting were present on the ancestral autosomes

We next asked whether differences in miRNA targeting were present on the ancestral autosomes

that gave rise to the mammalian X and Y chromosomes. To reconstruct the ancestral state of

miRNA targeting, we first focused on miRNA target sites in the 3` UTR of human orthologs that

align with perfect identity to a site in the corresponding chicken ortholog; these sites were likely

present in the common ancestor of mammals and birds (Figure 3A,B). We found that X-Y pairs

have the most human-chicken conserved target sites, followed by X-inactivated genes, and then

X escape genes (Figure 3C, top). Unlike the miRNA conservation scores used earlier, this metric

does not account for background conservation; we therefore estimated the background

conservation of each 3` UTR using shuffled miRNA family seed sequences (see Methods). X-Y

pairs, X-inactivated genes, and X escape genes do differ significantly with respect to background

conservation (data not shown), but these differences cannot account for the observed differences

in true human-chicken conserved sites (Figure 3C, bottom). We observed similar results for the

expanded set of X-Y pairs across 8 mammals (Supplemental Figure S6A).

Differences in the number of human-chicken conserved sites among the three classes of

X-linked genes could be explained by heterogeneity in miRNA targeting present on the ancestral

autosomes, or by ancestral homogeneity followed by different rates of target site loss during or

following X-Y differentiation. To distinguish between these two possibilities, we took advantage

of previous reconstructions of human sex chromosome evolution (Figure 3A) (Bellott et al.,

2014), which confirmed that, following the divergence of placental mammals from marsupials,

an X-autosome chromosomal fusion generated the X-added region (XAR) (Watson, Spencer,

Riggs, & Graves, 1990). Genes on the XAR are therefore X-linked in placental mammals, but

autosomal in marsupials such as the opossum. We limited our analysis to genes in the XAR and

target sites conserved between orthologous chicken and opossum 3` UTRs, ignoring site

conservation in humans; these sites were likely present in the common ancestor of mammals and

birds, and an absence of such sites cannot be explained by site loss following X-Y differentiation.

We observed the same pattern as with the human-chicken conserved sites, both before and after

accounting for background 3` UTR conservation (Figure 3D, three gene classes; Supplemental

Figure S6B, X-Y pairs across 8 mammals). These results demonstrate that the autosomal

precursors of X-Y pairs and X-inactivated genes were subject to more miRNA-mediated

regulation than X escape genes. Combined with our earlier results, we conclude that present-day

heterogeneities in dosage sensitivity on the mammalian X chromosome existed on the ancestral

autosomes from which it derived.

240
Figure 3. Heterogeneities in X-linked miRNA targeting were present on the ancestral
autosomes. (A) Example reconstruction of an ancestral miR-96 target site in the 3` UTR of
KDM6A, an X-linked gene in the X-added region (XAR) with a surviving Y homolog. Dots in
non-human species indicate identity with the human sequence, dashes gaps indicate gaps in the
multiple sequence alignment. (B) Distributions of sites conserved between 3` UTRs of human
and chicken orthologs (top) or comparisons to background expectation (bottom, see Methods) for
human X-Y pairs (n = 16), X-inactivated genes (n = 251), and X escape genes (n = 42). (C)

Statistics as in (B), but using sites conserved between chicken and opossum 3` UTRs only for
genes in the XAR; X-Y pairs (n = 11), X-inactivated genes (n = 58), and X escape genes (n = 27).
250
251
252
Z-W pairs have higher miRNA conservation scores than other ancestral Z-linked genes

We next assessed whether classes of avian Z-linked genes, those with and without a W homolog,

show analogous heterogeneities in sensitivity to dosage increases. We used the set of ancestral

genes reconstructed through cross-species comparisons of the avian Z chromosome and

orthologous human autosomes and focused on the set of Z-W pairs identified by sequencing of

the chicken W chromosome (Bellott et al., 2017, 2010). To increase the phylogenetic breadth of

our comparisons, we also included candidate Z-W pairs obtained through comparisons of male

and female genome assemblies (4 species set) or inferred by read-depth changes in female

genome assemblies (14 species set, see Methods for details) (Zhou et al., 2014). The more

complete 3` UTR annotations in the human genome relative to chicken allow for a more accurate

assessment of conserved miRNA targeting. Accordingly, we analyzed the 3` UTRs of the human

orthologs of chicken Z-linked genes, reasoning that important properties of chicken Z-linked

genes should be retained by their human autosomal orthologs.

We found that the human orthologs of Z-W pairs have higher miRNA conservation

scores than the human orthologs of other ancestral Z genes (Figure 4A, B). Differences in

miRNA conservation scores between Z-W pairs and other ancestral Z genes remained significant

when considering the expanded sets of Z-W pairs across four and 14 avian species

(Supplemental Figure S7). These differences are not driven by distinct subsets of genes, as

indicated by gene resampling with replacement (Supplemental Figure S8), and cannot be

accounted for by within-UTR variation in regional conservation (Supplemental Figure S9).

Logistic regression models indicate that miRNA conservation scores provide additional

information not captured by known factors (Bellott et al., 2017) influencing survival of W-linked

genes (known factors only model, AIC 137.8; known factors plus miRNA conservation model,

AIC 127.1; lower AIC indicates better model). Together, these results indicate that the autosomal

orthologs of Z-linked genes with a surviving W homolog are more sensitive to changes in dosage

-- both increases and decreases -- than are genes without a surviving W homolog.

280
281　**Figure 4. Z-W pairs have higher miRNA conservation scores than other ancestral Z-linked**
282　**genes.** P_{CT} score distributions of all gene-miRNA interactions involving the human orthologs of
283　(A) chicken Z-W pairs (n = 832 interactions from 28 genes) and other ancestral Z genes (n =
284　16,692 interactions from 657 genes). *** p < 0.001, two-sided Kolmogorov-Smirnov test. (B)
285　Mean gene-level P_{CT} scores. *** p < 0.001, two-sided Wilcoxon rank-sum test.
286
287
288
Heterogeneities in Z-linked miRNA targeting were present on the ancestral autosomes

We next asked whether differences in miRNA targeting between Z-W pairs and other ancestral

Z-linked genes were present on the ancestral autosomes that gave rise to the avian Z and W

chromosomes. We found that chicken Z-W pairs have more human-chicken-conserved miRNA

target sites than their Z-linked counterparts without surviving W homologs, both before (Figure

5C, top) and after (Figure 5C, bottom) accounting for the background conservation of each

individual 3` UTR. To confirm that these differences represent ancestral heterogeneity rather

than differential site loss during or following Z-W differentiation, we instead considered the

number of sites conserved between human and anolis lizard, which diverged from birds prior to

Z-W differentiation (Figure 5A). Chicken Z-W pairs contain an excess of human-anolis

conserved miRNA target sites, both before (Figure 5D, top) and after (Figure 5D, bottom)

accounting for the background conservation of each individual 3` UTR. We observed similar

results with the predicted four-species (Supplemental Figure S10) and 14-species (Supplemental

Figure S11) sets of Z-W pairs. Thus, the autosomal precursors of avian Z-W pairs were subject

to more miRNA-mediated regulation than the autosomal precursors of Z-linked genes that lack a

W homolog, indicating heterogeneity in dosage sensitivity among genes on the ancestral

autosomes that gave rise to the avian Z chromosome.

Figure 5. Heterogeneities in Z-linked miRNA targeting were present on the ancestral autosomes. (A) Example reconstruction of an ancestral miR-145 target site in the 3` UTR of RASA1, a Z-linked gene with a surviving W homolog.  Example of 3` UTR sequence alignment for RASA1, a Z-linked gene with a surviving W homolog, with a target site for miR-145 highlighted in gray. (B) Numbers of sites conserved between 3` UTRs of human and chicken orthologs (top) or comparisons to background expectation (bottom) for chicken Z-W pairs (n = 27) and other ancestral Z genes (n = 578). (C) Statistics as in (B), but using sites conserved between human and anolis 3` UTRs.

Analyses of experimental datasets validate miRNA target site function

Our results to this point, which indicate preexisting heterogeneities in dosage constraints among

X- or Z-linked genes as inferred by predicted miRNA target sites, lead to predictions regarding

the function of these sites in vivo. To test these predictions, we turned to publically available

experimental datasets consisting both of gene expression profiling following transfection or

knockout of individual miRNAs, and of high-throughput crosslinking-immunoprecipitation

(CLIP) to identify sites that bind Argonaute in vivo (see Methods). If the above-studied sites are

effective in mediating target repression, targets of an individual miRNA should show increased

expression levels or Argonaute binding following miRNA transfection, and decreased expression

levels following miRNA knockout. Together, our analyses of publically available datasets

fulfilled these predictions, validating the function of sites targeting ancestral X-linked genes and

the autosomal orthologs of ancestral Z-linked genes in multiple cellular contexts and species

(Figure 6). From the gene expression profiling data, we observed results consistent with effective

targeting by a) eleven different miRNA families in human HeLa cells (Supplemental Figure S12),

b) four different miRNAs in human HCT116 and HEK293 cells (Supplemental Figure S13), and

c) miR-155 in mouse B and Th1 cells (Supplemental Figure S14). In the CLIP data, the human

orthologs of X- or Z-linked targets of miR-124 are enriched for Argonaute-bound clusters that

appear following miR-124 transfection, while a similar but non-significant enrichment is

observed for miR-7 (Supplemental Figure S15). Thus, conserved miRNA target sites used to

infer dosage constraints on X-linked genes and the autosomal orthologs of Z-linked genes can

effectively mediate target repression in living cells.

Figure 6. Analyses of experimental datasets validate miRNA target site function. Responses to transfection (A,B,C) or knockout (D) of indicated miRNAs in human (A,B,C) or mouse (D) cell-types. Each panel depicts corresponding changes in mRNA levels (A,B), in fraction of Argonaute-bound genes (C), and in mRNA stability and translational efficiency as measured by ribosome protected fragments (RPF, D). In each case, X-linked genes and the human orthologs of Z-linked genes containing target sites with an assigned P_{CT} score (red) for the indicated miRNA were compared to all expressed genes lacking target sites (black); gene numbers are indicated in parentheses. (A,B,D) *** $p < 0.001$, two-sided Kolmogorov-Smirnov test. (C) * $p < 0.05$, two-sided Fisher's exact test.

DISCUSSION

Here, through the evolutionary reconstruction of microRNA (miRNA) target sites, we

demonstrate preexisting heterogeneities in dosage sensitivity among genes on the mammalian X

and avian Z chromosomes. We first showed that, across all human autosomal genes, dosage

sensitivity -- as indicated by patterns of genic copy number variation -- correlates with the degree

of conserved miRNA targeting. We found that conserved targeting correlates especially strongly

with sensitivity to dosage increases, consistent with miRNA targeting serving to reduce gene

expression. Turning to the sex chromosomes of mammals and birds, genes that retained a

homolog on the sex-specific Y or W chromosome (X-Y and Z-W pairs) are more dosage-

sensitive than genes with no Y or W homolog. In mammals, genes with no Y homolog that

became subject to XCI are more dosage-sensitive than those that continued to escape XCI

following Y gene decay. We then reconstructed the ancestral state of miRNA targeting,

observing significant heterogeneities in the extent of miRNA targeting, and thus dosage

sensitivity, on the ancestral autosomes that gave rise to the mammalian and avian sex

chromosomes. Finally, through analysis of publically available experimental datasets, we

validated the function, in living cells, of the miRNA target sites used to infer dosage sensitivity.

We thus conclude that differences in dosage sensitivity – both to increases and to decreases in

gene dosage -- among genes on the ancestral autosomes influenced their evolutionary trajectory

during sex chromosome evolution, not only on the sex-specific Y and W chromosomes, but also

on the sex-shared X chromosome.

Our findings build upon previous work in three important ways. First, our analysis of

miRNA-mediated repression indicates that these heterogeneities consist of sensitivities to dosage

increases and decreases, whereas previous studies had either focused on sensitivity to

underexpression or could not differentiate the two. Second, our reconstruction of miRNA

targeting on the ancestral autosomes provides direct evidence that heterogeneities in dosage

sensitivity among classes of X- and Z-linked were preexisting rather than acquired during sex

chromosome evolution. Finally, by pointing to specific regulatory sequences (miRNA target

sites) functioning to tune gene dosage both prior to and during sex chromosome evolution, our

study provides a view of dosage compensation encompassing post-transcriptional regulation.

Human disease studies support the claim that increased dosage of X-Y pairs and X-

inactivated genes is deleterious to fitness. Copy number gains of the X-linked gene *KDM6A*,

which has a surviving human Y homolog, are found in patients with developmental

abnormalities and intellectual disability (Lindgren et al., 2013). *HDAC6*, *CACNA1F*, *GDI1*, and

IRS4 all lack Y homologs and are subject to XCI in humans. A mutation in the 3` UTR of

HDAC6 abolishing targeting by miR-433 has been linked to familial chondrodysplasia in both

sexes (Simon et al., 2010). Likely gain-of-function mutations in *CACNA1F* cause congenital

stationary night blindness in both sexes (Hemara-Wahanui et al., 2005). Copy number changes of

GDI1 correlate with the severity of X-linked mental retardation in males, with female carriers

preferentially inactivating the mutant allele (Vandewalle et al., 2009). Somatic genomic deletions

downstream of *IRS4* lead to its overexpression in lung squamous carcinoma (Weischenfeldt et al.,

2017). Males with partial X disomy due to translocation of the distal long arm of the X

chromosome (Xq28) to the long arm of the Y chromosome show severe mental retardation and

developmental defects (Lahn et al., 1994). Most genes in Xq28 are inactivated in 46,XX females

but escape inactivation in such X;Y translocations, suggesting that increased dosage of Xq28

genes caused the cognitive and developmental defects. We anticipate that further studies will

reveal additional examples of the deleterious effects of increases in gene dosage of X-Y pairs

and X-inactivated genes.

We and others previously proposed that Y gene decay drove upregulation of homologous

X-linked genes in both males and females, and that XCI subsequently evolved at genes sensitive

to increased expression from two active X-linked copies in females (Jegalian & Page, 1998;

Ohno, 1967). Our finding that X-inactivated genes have higher miRNA conservation scores than

X escape genes is consistent with this aspect of the model. However, recent studies indicating

heterogeneity in dosage sensitivity between classes of mammalian X- or avian Z-linked genes

(Bellott et al., 2014, 2017; Pessia et al., 2012; Zimmer et al., 2016), combined with the present

finding that these dosage sensitivities existed on the ancestral autosomes, challenge the previous

assumption of a single evolutionary pathway for all sex-linked genes.

We therefore propose a revised model of X-Y and Z-W evolution in which the ancestral

autosomes that gave rise to the mammalian and avian sex chromosomes contained three (or two,

in the case of birds) classes of genes with differing dosage sensitivities (Figure 7A,B). For

ancestral genes with high dosage sensitivity, Y or W gene decay would have been highly

deleterious, and thus the Y- or W-linked genes were retained. According to our model, these

genes' high dosage sensitivity also precluded upregulation of the X- or Z-linked homolog, and,

in mammals, subsequent X-inactivation; indeed, their X-linked homologs continue to escape

XCI (Bellott et al., 2014). For ancestral mammalian genes of intermediate dosage sensitivity, Y

gene decay did occur, and was accompanied or followed by compensatory upregulation of the X-

linked homolog in both sexes; the resultant increased expression in females was deleterious and

led to the acquisition of XCI. Ancestral mammalian genes of low dosage sensitivity continued to

escape XCI following Y decay, and perhaps did not undergo compensatory X upregulation

(Figure 6A). These genes' dosage insensitivity set them apart biologically, and evolutionarily,

from the other class of X-linked genes escaping XCI -- those with a surviving Y homolog.

423
Figure 7. An evidence-based model of mammalian and avian sex chromosome evolution
from ancestral autosomes. In this model, preexisting heterogeneities in dosage sensitivity
determined the trajectory of Y/W gene loss in both mammals and birds, and of subsequent X-
inactivation in mammals. Colored arrow widths are scaled approximately to the number of
ancestral genes in each class. The dashed orange line represents the possibility that a subset of X-
linked genes may have not undergone compensatory X upregulation following Y gene decay.
Small black arrows adjacent to molecular evolutionary transitions indicate increases or decreases
in sex-linked gene dosage
432
433
Our revised model relates preexisting, gene-by-gene heterogeneities in dosage sensitivity

to the outcomes of sex chromosome evolution. However, the suppression of X-Y recombination

did not occur on a gene-by-gene basis, instead initiating Y gene decay and subsequent dosage

compensation through a series of large-scale inversions encompassing many genes (Lahn & Page,

1999). The timings and boundaries of these evolutionary strata varied among mammalian

lineages, thus leading to unique chromosome-scale evolutionary dynamics across mammals.

These large-scale changes would have then allowed for genic selection to take place according to

the preexisting dosage sensitivities outlined above. In this way, the course of sex chromosome

evolution in mammals is a composite of 1) preexisting, gene-by-gene dosage sensitivities and 2)

the manner in which the history of the X and Y unfolded in particular lineages via discrete, large-

scale inversions.

In this study, we have focused on classes of ancestral X-linked genes delineated by the

survival of a human Y homolog or by the acquisition of XCI in humans, but such evolutionary

states can differ among mammalian lineages and species. In mouse, for instance, both Y gene

decay (Bellott et al., 2014) and the acquisition of X-inactivation (Yang et al., 2010) are more

complete than in humans or other mammals. These observations could be explained by shortened

generation times in the rodent lineage, resulting in longer evolutionary times, during which the

forces leading to Y gene decay and the acquisition of X-inactivation could act (Charlesworth &

Crow, 1978; Jegalian & Page, 1998; Ohno, 1967). In the future, more complete catalogues of X-

inactivation and escape in additional mammalian lineages would make it possible to examine

whether analogous, preexisting dosage sensitivities differentiate the three classes of X-linked

genes (X-Y pairs, X-inactivated genes, and X escape genes) in other species.

Previous studies have sought evidence of X-linked upregulation during mammalian sex

chromosome evolution using comparisons of gene expression levels between the whole of the X

chromosome and all of the autosomes, with equal numbers of studies rejecting or finding

evidence consistent with upregulation (Deng et al., 2011; Julien et al., 2012; Kharchenko, Xi, &

Park, 2011; Lin, Xing, Zhang, & He, 2012; Xiong et al., 2010). This is likely due to gene-by-

gene heterogeneity in dosage sensitivities that resulted in a stronger signature of upregulation at

more dosage sensitive genes (Pessia et al., 2012). Similarly, studies of Z-linked gene expression

in birds provide evidence for the gene-by-gene nature of Z dosage compensation, as measured by

comparisons of gene expression levels between ZZ males and ZW females (Itoh et al., 2007;

Mank & Ellegren, 2009; Uebbing et al., 2015), and indicate a stronger signature of dosage

compensation at predicted dosage-sensitive genes (Zimmer et al., 2016). By showing that such

dosage sensitivities existed on the ancestral autosomes and consist of sensitivity to both increases

and decreases, our findings highlight an additional aspect of dosage compensation that affects

both birds and mammals.

In addition to revealing similarities between mammals and birds, our study provides a

view of dosage compensation that highlights post-transcriptional regulatory mechanisms. The

majority of previous studies, by relying upon measurements of mRNA abundance, could not

differentiate transcriptional and post-transcriptional effects (Deng et al., 2011; Itoh et al., 2007;

Julien et al., 2012; Kharchenko et al., 2011; Lin et al., 2012; Xiong et al., 2010; Zimmer et al.,

2016). Other studies have drawn inferences regarding post-transcriptional dosage compensation

from measurements of extant protein abundance (Chen & Zhang, 2015; Pessia et al., 2012;

Uebbing et al., 2015). Our study points to specific non-coding sequences with known

mechanisms (microRNA target sites) functioning across evolutionary time. Furthermore, our

finding of greater conserved miRNA targeting of X-inactivated genes relative to X escape genes

shows that it is possible to predict the acquisition of a transcriptional regulatory state (XCI)

during sex chromosome evolution on the basis of a preexisting, post-transcriptional regulatory

state. Perhaps additional post-transcriptional regulatory mechanisms and their associated

regulatory elements will be shown to play roles in mammalian and avian dosage compensation.

Recent work has revealed that the sex-specific chromosome -- the Y in mammals and the

W in birds -- convergently retained dosage-sensitive genes with important regulatory functions

(Bellott et al., 2014, 2017). Our study, by reconstructing the ancestral state of post-transcriptional

regulation, provides direct evidence that such heterogeneity in dosage sensitivity existed on the

ancestral autosomes that gave rise to the mammalian and avian sex chromosomes. This

heterogeneity influenced both survival on the sex-specific chromosomes in mammals and birds

and the evolution of XCI in mammals. Thus, two independent experiments of nature offer

empirical evidence that modern-day amniote sex chromosomes were shaped, during evolution,

by the properties of the ancestral autosomes from which they derive.

METHODS

Statistics

Details of all statistical tests (type of test, test statistic, and p-value) used in this manuscript are

provided in Supplemental Table S1.

Human genic copy number variation

To annotate gene deletions and duplications, we used data from the Exome Aggregation

Consortium (ExAC, RRID:SCR_004068)

(ftp://ftp.broadinstitute.org/pub/ExAC_release/release0.3.1/cnv/), which consists of autosomal

genic duplications and deletions (both full and partial) called in 59,898 exomes (Ruderfer et al.,

2016). We used the publicly available genic deletion counts but re-computed genic duplication

counts using only full duplications, reasoning that partial duplications are unlikely to result in

increased dosage of the full gene product. We thus required that an individual duplication fully

overlapped the longest protein-coding transcript (GENCODE v19) of a gene using BEDtools

(RRID:SCR_006646) (Quinlan & Hall, 2010). We removed genes flagged by ExAC as lying in

known regions of recurrent CNVs. This yielded 4,118 genes within duplications and deletions,

3,976 genes within deletions but not duplications, 2,916 genes within duplications but not

deletions, and 3,510 genes not subject to duplication or deletion. These gene assignments are

provided in Supplemental Table S2.

X-linked gene sets

Analyses of conserved miRNA targeting based on multiple species alignments are unreliable for

multicopy or ampliconic genes due to ambiguous sequence alignment between species. To avoid

such issues, we first removed multicopy and ampliconic genes (Mueller et al., 2013) from a

previously published set of human X genes present in the amniote ancestor (Bellott et al., 2014).

We then excluded genes in the human pseudoautosomal (PAR) regions since these genes have

not been exposed to the same evolutionary forces as genes in regions where X-Y recombination

has been suppressed. Of the remaining ancestral X genes, we classified the 15 genes with human

Y-linked homologs as X-Y pairs. We also analyzed the larger set of 32 X-Y pairs across eight

mammals (human, chimpanzee, rhesus macaque, marmoset, mouse, rat, bull, and opossum) with

sequenced Y chromosomes (Bellott et al., 2014).

To classify ancestral X-linked genes without Y homologs as subject to or escaping XCI

in humans, we used a collection of consensus XCI calls which aggregate the results of three

studies (Carrel & Willard, 2005; A. M. Cotton et al., 2013; a. M. Cotton et al., 2014) assaying

XCI escape (Balaton et al., 2015). Out of 472 ancestral X genes without a human Y homolog

assigned an XCI status by Balaton et al. (Balaton et al., 2015), 329 were subject to XCI ("Subject"

or "Mostly subject" in Balaton et al.), 26 displayed variable escape ("Variable escape" or

"Mostly variable escape") from XCI, and 30 showed consistent escape ("Escape" or "Mostly

escape"). We excluded 40 ancestral X genes with a "Discordant" XCI status as assigned by

Balaton et al. In the main text, we present results obtained after combining both variable and

consistent escape calls from Balaton et al. into one class, yielding the following counts: 15 X-Y

pairs, 329 ancestral X genes subject to XCI, and 56 ancestral X genes with evidence of escape

from XCI. We also performed analyses considering escape and variable escape genes separately.

Information on X-linked genes is provided in Supplemental Table S3.

Z-linked gene sets

We previously refined the ancestral gene content of the avian sex chromosomes to 685 Z-linked

genes with human orthologs by sequencing of the chicken Z chromosome and analysis of 13

other avian species with published female genomes (Bellott et al., 2017). Of these 685 ancestral

Z genes, 28 retained a homolog on the fully sequenced chicken W chromosome. Including three

additional avian species in which candidate W-linked genes were ascertained by directly

comparing male and female genome assemblies results in a total of 78 W-linked genes. Including

another 10 avian species in which W-linkage was inferred by read depth changes in a female

genome results in a total of 157 W-linked genes. Information on Z-linked genes is provided in

Supplemental Table S4.

microRNA target site P_{CT} scores

Pre-calculated P_{CT} scores for all gene-miRNA family interactions were obtained from

TargetScanHuman v7.1 (RRID:SCR_010845)

([http://www.targetscan.org/vert_71/vert_71_data_download/Summary_Counts.all_predictions.tx](http://www.targetscan.org/vert_71/vert_71_data_download/Summary_Counts.all_predictions.txt.zip)

[t.zip](http://www.targetscan.org/vert_71/vert_71_data_download/Summary_Counts.all_predictions.txt.zip)), (Friedman et al., 2009). We excluded mammalian-specific miRNA families based on

classifications by Friedman et al (Friedman et al., 2009) and updated in TargetScanHuman

v7.1(Agarwal, Bell, Nam, & Bartel, 2015). To account for gene-specific variability in the

number and P_{CT} score of gene-miRNA interactions within a group of genes, we sampled 1000x

with replacement from the same group of genes and computed the mean gene-miRNA P_{CT} score

for all associated gene miRNA interactions from each sampling. These 1000 samplings were

then used to estimate the median resampled gene-miRNA P_{CT} and 95% confidence intervals.

Variation in within- conservation bias

The P_{CT} of a given miRNA target site depends on the conservation of the site, as measured by the

total branch length of the phylogenetic tree containing the target sites (branch length score, BLS)

relative to the mean BLS of the whole 3` UTR. To address the possibility that non-uniformity in

the regional BLS could artificially inflate or deflate conservation scores of certain target sites, we

implemented a step-detection algorithm to segment 3` UTRs into regions of homogeneous BLS

values and calculated miRNA site conservation relative to these smaller regions. In order to call

steps within a 3` UTR, we computed the t-test p-value between the BLS values of the 50-nt

window upstream and downstream of each nucleotide position in the 3` UTR. Transitions were

called at a log p-value cutoff of -15. Because of noise in the BLS signal, the log p-value often

dips below -15 several times around each transition. If more than 1 position met the cutoff within

100 nucleotides of each other, we took only the one with the smaller p-value. We then computed,

for each miRNA site, the ratio of the mean BLS of its section to that of the entire 3` UTR; we

term this statistic the "within-UTR conservation bias". Values of this statistic greater than 1

indicate that the P_{CT} overestimates the relative conservation of a given target site, while values

less than 1 indicate that the P_{CT} underestimates conservation. For gene-miRNA interactions with

multiple sites, we used the mean within-UTR conservation bias for all sites. We also repeated

P_{CT} score comparisons between classes of X- and Z-linked genes with P_{CT} scores normalized by

the corresponding gene-miRNA within-UTR conservation bias (Supplemental Figure S5C,

Supplemental Figure S9B). These scores, corresponding to all gene-miRNA interactions, are

provided in Supplemental Table S5.

Logistic regression

Logistic regression models were constructed using the function 'multinom' in the R package

'nnet.' For Z-linked genes, this reduces to binomial logistic regression, with probabilities of a

binary outcome (surviving W homolog or not) modeled as a log-linear combination of predictor

variables. Since there are three classes of X-linked genes, class probabilities are modeled by a

multinomial distribution. To compare models including only known factors to ones with mean

P_{CT} scores as an addition predictor, we used Aikake's Information Criterion (AIC), since it

penalizes more complex models (such as those with P_{CT} as an additional predictor). We used

previously published values for known factors in the survival of Y-linked (Bellott et al., 2014)

and W-linked (Bellott et al., 2017) genes except for human expression breadth, which we

recalculated using data from the GTEx Consortium v6 data release (Consortium, 2015). Briefly,

kallisto was used to estimate transcript per million (TPM) values in the 10 male samples with the

highest RNA integrity numbers (RINs) from each of 37 tissues, and expression breadth across

tissues was calculated as described in (Bellott et al., 2014), using median TPM values for each

tissue.

Human-chicken conserved microRNA target sites

Site-wise alignment information was obtained from TargetScanHuman v7.1

(http://www.targetscan.org/vert_71/vert_71_data_download/Conserved_Family_Info.txt.zip). To

determine which target sites are present in the 3` UTRs of both human and chicken orthologs, we

counted, for genes with both a human and chicken ortholog, the number of miRNA interactions

that had at least one target site in both human and chicken. To control for gene-specific

background 3` UTR conservation, we generated six control k-mers for each miRNA family seed

sequence that were matched exactly for nucleotide and CpG content. Six was the maximum

number of unique control k-mers that could be generated for all sequences. We repeated the

above counting analysis with each of the control k-mers using scripts from TargetScan, and

compared, for each gene, the observed number of human-chicken-conserved miRNA interactions

(the observed conservation signal) to the average number from controls (the background

conservation). This same procedure was repeated for alternative pairs of species considered

(opossum-chicken and human-anolis lizard).

Gene expression profiling and crosslinking datasets

Fold-changes in mRNA expression from a compendium of small RNA (sRNA) transfections

(corresponding to twelve different miRNAs) in HeLa cells were obtained from Agarwal and

colleagues (Agarwal et al., 2015) (GSM210904, GSM37601, GSM210913, GSM210903,

GSM210911, GSM210898, GSM210897, GSM210897, GSM210901, GSM210909,

GSM119747; E-MEXP-1402(1595297513)). Further datasets describing the effects of

transfecting miR-103 in HCT116 cells (Linsley et al., 2007) (GSM156580), knocking down

miR-92a in HEK293 cells (Hafner et al., 2010) (GSM538818), transfecting miR-7 or miR-124 in

HEK293 cells (Hausser, Landthaler, Jaskiewicz, Gaidatzis, & Zavolan, 2009) (GSM363763,

GSM363766, GSM363769, GSM363772, GSM363775, GSM363778), or of knocking out miR-

155 in mouse B cells (Eichhorn et al., 2014)

(GSM1479572, GSM1479576, GSM1479580, GSM1479584), T cells (Loeb et al., 2012)

(GSM1012118, GSM1012119, GSM1012120, GSM1012121, GSM1012122, GSM1012123), or

Th1 and Th2 cells (Rodriguez et al., 2007) (E-TABM-232), processed as described in (Agarwal

et al., 2015), were provided by V. Agarwal. Targets for the PAR-CLIP study (Hafner et al.,

2010) were inferred from an online resource of HEK293 clusters observed after transfection of

either miR-124

(http://www.mirz.unibas.ch/restricted/clipdata/RESULTS/miR124_TRANSFECTION/miR124_

TRANSFECTION.html) or miR-7

(http://www.mirz.unibas.ch/restricted/clipdata/RESULTS/miR7_TRANSFECTION/miR7_TRA

NSFECTION.html). All fold-changes and CLIP targets are provided in Supplemental Table S6.

DATA ACCESS

Data supporting the findings of this study are available within the paper and its Supplemental

information files.

Code availability

A custom Python (RRID:SCR_008394) script utilizing Biopython (RRID:SCR_007173) was

used to generate shuffled miRNA family seed sequences. Identification of miRNA target site

matches using shuffled seed sequences was performed using the 'targetscan_70.pl' perl script

(http://www.targetscan.org/vert_71/vert_71_data_download/targetscan_70.zip). 3` UTR

segmentation was performed with the 'plot_transitions.py' python script. Code is available at:

https://github.com/snaqvi1990/Naqvi17-code

ACKNOWLEDGEMENTS

We thank V. Agarwal, S. Eichorn, S. McGeary, and D. Bartel for assistance with the TargetScan

database and helpful discussions; A. Godfrey for updated human-chicken orthology information;

and A. Godfrey, J. Hughes and H. Skaletsky for critical reading of the manuscript. This work

was supported by the National Institutes of Health and the Howard Hughes Medical Institute.

S.N. was supported under a research grant by Biogen.

Author contributions

S.N., D.W.B. and D.C.P designed the study. S.N. performed analyses with assistance from

D.W.B. K.S.L developed and implemented the step-detection algorithm. S.N. and D.C.P wrote

the paper.

DISCLOSURE DECLARATION

The authors declare no competing financial interests.

Agarwal, V., Bell, G. W., Nam, J., and Bartel, D. P. (2015). Predicting effective microRNA
target sites in mammalian mRNAs, 1–38. https://doi.org/10.7554/eLife.05005
Balaton, B. P., Cotton, A. M., and Brown, C. J. (2015). Derivation of consensus inactivation
status for X-linked genes from genome-wide studies. *Biology of Sex Differences*, *6*(1), 35.
https://doi.org/10.1186/s13293-015-0053-7
Bartel, D. P. (2009). MicroRNAs: target recognition and regulatory functions. *Cell*, *136*(2), 215–
33. https://doi.org/10.1016/j.cell.2009.01.002
Bellott, D. W., Hughes, J. F., Skaletsky, H., Brown, L. G., Pyntikova, T., Cho, T.-J., Koutseva,
N., Zaghlul, S., Graves, T., Rock, S., Kremitzki, C., Fulton, R. S., Dugan, S., Ding, Y.,
Morton, D., Khan, Z., Lewis, L., … Page, D. C. (2014). Mammalian Y chromosomes retain
widely expressed dosage-sensitive regulators. *Nature*, *508*(7497), 494–9.
https://doi.org/10.1038/nature13206
Bellott, D. W., Skaletsky, H., Cho, T.-J., Brown, L., Locke, D., Chen, N., Galkina, S., Pyntikova,
T., Koutseva, N., Graves, T., Kremitzki, C., Warren, W. C., Clark, A. G., Gaginskaya, E.,
Wilson, R. K., and Page, D. C. (2017). Avian W and mammalian Y chromosomes
convergently retained dosage-sensitive regulators. *Nature Genetics*, *in press*.
Bellott, D. W., Skaletsky, H., Pyntikova, T., Mardis, E. R., Graves, T., Kremitzki, C., Brown, L.
G., Rozen, S., Warren, W. C., Wilson, R. K., and Page, D. C. (2010). Convergent evolution
of chicken Z and human X chromosomes by expansion and gene acquisition. *Nature*,
466(7306), 612–616. https://doi.org/10.1038/nature09172
Carrel, L., and Willard, H. F. (2005). X-inactivation profile reveals extensive variability in X-
linked gene expression in females. *Nature*, *434*(March), 400–404.
https://doi.org/10.1038/nature03479
Charlesworth, B., and Crow, J. F. (1978). Model for evolution of Y chromosomes and dosage
compensation, *75*(11), 5618–5622.
Chen, X., and Zhang, J. (2015). No X-chromosome dosage compensation in human proteomes.
Molecular Biology and Evolution, *32*(6), 1456–1460.
https://doi.org/10.1093/molbev/msv036
Consortium, G. (2015). The Genotype-Tissue Expression (GTEx) pilot analysis: Multitissue
gene regulation in humans. *Science (New York, N.Y.)*, *348*(6235), 648–660.
Cotton, A. M., Ge, B., Light, N., Adoue, V., Pastinen, T., and Brown, C. J. (2013). Analysis of
expressed SNPs identifies variable extents of expression from the human inactive X
chromosome. *Genome Biology*, *14*(11), R122. https://doi.org/10.1186/gb-2013-14-11-r122
Cotton, a. M., Price, E. M., Jones, M. J., Balaton, B. P., Kobor, M. S., and Brown, C. J. (2014).
Landscape of DNA methylation on the X chromosome reflects CpG density, functional
chromatin state and X-chromosome inactivation. *Human Molecular Genetics*, *24*(6), 1528–
1539. https://doi.org/10.1093/hmg/ddu564
Deng, X., Hiatt, J. B., Nguyen, D. K., Ercan, S., Sturgill, D., Hillier, L. W., Schlesinger, F.,
Davis, C. a, Reinke, V. J., Gingeras, T. R., Shendure, J., Waterston, R. H., Oliver, B., Lieb,
J. D., and Disteche, C. M. (2011). Evidence for compensatory upregulation of expressed X-
linked genes in mammals, Caenorhabditis elegans and Drosophila melanogaster. *Nature*
Genetics, *43*(12), 1179–85. https://doi.org/10.1038/ng.948
Eichhorn, S. W., Guo, H., McGeary, S. E., Rodriguez-Mias, R. a, Shin, C., Baek, D., Hsu, S.-H.,
Ghoshal, K., Villén, J., and Bartel, D. P. (2014). mRNA Destabilization Is the Dominant
Effect of Mammalian MicroRNAs by the Time Substantial Repression Ensues. *Molecular*
Cell. https://doi.org/10.1016/j.molcel.2014.08.028

Friedman, R. C., Farh, K. K.-H., Burge, C. B., and Bartel, D. P. (2009). Most mammalian mRNAs are conserved targets of microRNAs. *Genome Research*, *19*(1), 92–105. https://doi.org/10.1101/gr.082701.108

Hafner, M., Landthaler, M., Burger, L., Khorshid, M., Berninger, P., Rothballer, A., Jr, M. A., Munschauer, M., Ulrich, A., Wardle, G. S., Dewell, S., Zavolan, M., and Tuschl, T. (2010). Transcriptome-wide identification of RNA-binding protein and microRNA target sites by PAR-CLIP. *Cell*, *141*(1), 129–141. https://doi.org/10.1016/j.cell.2010.03.009.Transcriptome-wide

Hausser, J., Landthaler, M., Jaskiewicz, L., Gaidatzis, D., and Zavolan, M. (2009). mRNA binding of Argonaute / EIF2C − miRNA complexes and the degradation of miRNA targets Relative contribution of sequence and structure features to the mRNA binding of Argonaute / EIF2C – miRNA complexes and the degradation of miRNA targets. *Genome Research*, 2009–2020. https://doi.org/10.1101/gr.091181.109

Hemara-Wahanui, A., Berjukow, S., Hope, C. I., Dearden, P. K., Wu, S.-B., Wilson-Wheeler, J., Sharp, D. M., Lundon-Treweek, P., Clover, G. M., Hoda, J.-C., Striessnig, J., Marksteiner, R., Hering, S., and Maw, M. a. (2005). A CACNA1F mutation identified in an X-linked retinal disorder shifts the voltage dependence of Cav1.4 channel activation. *Proceedings of the National Academy of Sciences of the United States of America*, *102*(21), 7553–7558. https://doi.org/10.1073/pnas.0501907102

Huang, N., Lee, I., Marcotte, E. M., and Hurles, M. E. (2010). Characterising and predicting haploinsufficiency in the human genome. *PLoS Genetics*, *6*(10), e1001154. https://doi.org/10.1371/journal.pgen.1001154

Hughes, J. F., Skaletsky, H., Brown, L. G., Pyntikova, T., Graves, T., Fulton, R. S., Dugan, S., Ding, Y., Buhay, C. J., Kremitzki, C., Wang, Q., Shen, H., Holder, M., Villasana, D., Nazareth, L. V, Cree, A., Courtney, L., … Page, D. C. (2012). Strict evolutionary conservation followed rapid gene loss on human and rhesus Y chromosomes. *Nature*, *483*(7387), 82–6. https://doi.org/10.1038/nature10843

Itoh, Y., Melamed, E., Yang, X., Kampf, K., Wang, S., Yehya, N., Van Nas, A., Replogle, K., Band, M. R., Clayton, D. F., Schadt, E. E., Lusis, A. J., and Arnold, A. P. (2007). Dosage compensation is less effective in birds than in mammals. *Journal of Biology*, *6*(1), 2. https://doi.org/10.1186/jbiol53

Jegalian, K., and Page, D. (1998). A proposed path by which genes common to mammalian X and Y chromosomes evolve to become X inactivated. *Nature*, *394*(August), 776–780. Retrieved from http://www.nature.com/nature/journal/v394/n6695/abs/394776a0.html

Julien, P., Brawand, D., Soumillon, M., Necsulea, A., Liechti, A., Schütz, F., Daish, T., Grützner, F., and Kaessmann, H. (2012, January). Mechanisms and evolutionary patterns of mammalian and avian dosage compensation. *PLoS Biology*. https://doi.org/10.1371/journal.pbio.1001328

Kaiser, V. B., Zhou, Q., and Bachtrog, D. (2011). Nonrandom gene loss from the drosophila miranda neo-Y chromosome. *Genome Biology and Evolution*, *3*, 1329–1337. https://doi.org/10.1093/gbe/evr103

Kharchenko, P. V, Xi, R., and Park, P. J. (2011). Evidence for dosage compensation between the X chromosome and autosomes in mammals. *Nature Genetics*, *43*(12), 1167-9–2. https://doi.org/10.1038/ng.991

Lahn, B. T., Ma, N., Breg, R. W., Stratton, R., Surti, U., and Page, D. C. (1994). Xq-Yq interchange resulting in supernormal X-linked gene expression in severely retarded males

with 46,XYq- karyotype. *Nature Genetics*, *8*, 362–369. https://doi.org/10.1038/ng1294-340

Lahn, B. T., and Page, D. (1999). Four Evolutionary Strata on the Human X Chromosome. *Science*, *286*(5441), 964–967. https://doi.org/10.1126/science.286.5441.964

Lin, F., Xing, K., Zhang, J., and He, X. (2012). Expression reduction in mammalian X chromosome evolution refutes Ohno's hypothesis of dosage compensation. *Proceedings of the National Academy of Sciences*, *109*(29), 11752–11757. https://doi.org/10.1073/pnas.1201816109

Lindgren, A. M., Hoyos, T., Talkowski, M. E., Hanscom, C., Blumenthal, I., Chiang, C., Ernst, C., Pereira, S., Ordulu, Z., Clericuzio, C., Drautz, J. M., Rosenfeld, J. a, Shaffer, L. G., Velsher, L., Pynn, T., Vermeesch, J., Harris, D. J., … Morton, C. C. (2013). Haploinsufficiency of KDM6A is associated with severe psychomotor retardation, global growth restriction, seizures and cleft palate. *Human Genetics*, *132*(5), 537–52. https://doi.org/10.1007/s00439-013-1263-x

Linsley, P. S., Schelter, J., Burchard, J., Kibukawa, M., Martin, M. M., Bartz, S. R., Johnson, J. M., Cummins, J. M., Raymond, C. K., Dai, H., Chau, N., Cleary, M., Jackson, A. L., Carleton, M., and Lim, L. (2007). Transcripts targeted by the microRNA-16 family cooperatively regulate cell cycle progression. *Mol Cell Biol*, *27*(6), 2240–2252. https://doi.org/10.1128/MCB.02005-06

Loeb, G. B., Khan, A. A., Canner, D., Hiatt, J. B., Shendure, J., Darnell, R. B., Leslie, C. S., and Rudensky, A. Y. (2012). Transcriptome-wide miR-155 Binding Map Reveals Widespread Noncanonical MicroRNA Targeting. *Molecular Cell*, *48*(5), 760–770. https://doi.org/http://dx.doi.org/10.1016/j.molcel.2012.10.002

Mank, J. E., and Ellegren, H. (2009). All dosage compensation is local: gene-by-gene regulation of sex-biased expression on the chicken Z chromosome. *Heredity*, *102*(3), 312–320. https://doi.org/10.1038/hdy.2008.116

Mueller, J. L., Skaletsky, H., Brown, L. G., Zaghlul, S., Rock, S., Graves, T., Auger, K., Warren, W. C., Wilson, R. K., and Page, D. C. (2013). Independent specialization of the human and mouse X chromosomes for the male germ line. *Nature Genetics*, *45*(9), 1083–1087. https://doi.org/10.1038/ng.2705

Nanda, I., Shan, Z., Schartl, M., Burt, D. W., Koehler, M., Nothwang, H., Grützner, F., Paton, I. R., Windsor, D., Dunn, I., Engel, W., Staeheli, P., Mizuno, S., Haaf, T., and Schmid, M. (1999). 300 million years of conserved synteny between chicken Z and human chromosome 9. *Nature Genetics*, *21*(march), 258–259. https://doi.org/10.1038/6769

Ohno, S. (1967). *Sex chromosomes and sex-linked genes*.

Pessia, E., Makino, T., Bailly-Bechet, M., McLysaght, A., and Marais, G. a B. (2012). Mammalian X chromosome inactivation evolved as a dosage-compensation mechanism for dosage-sensitive genes on the X chromosome. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(14), 5346–51. https://doi.org/10.1073/pnas.1116763109

Pinzón, N., Li, B., Martinez, L., Sergeeva, A., Presumey, J., Apparailly, F., and Seitz, H. (2016). The number of biologically relevant microRNA targets has been largely over-estimated The number of biologically relevant microRNA targets has been largely overestimated, (November), 1–11. https://doi.org/10.1101/gr.205146.116

Quinlan, A. R., and Hall, I. M. (2010). BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics*, *26*(6), 841–842. https://doi.org/10.1093/bioinformatics/btq033

Rodriguez, A., Vigorito, E., Clare, S., Warren, M. V, Couttet, P., Soond, D. R., Dongen, S. Van,
Grocock, R. J., Das, P. P., Miska, E. A., Vetrie, D., Okkenhaug, K., Enright, A. J., Dougan,
G., Turner, M., and Bradley, A. (2007). Requirement of bic/microRNA-155 for normal
immune function. *Science*, *981*(April), 608–611. https://doi.org/10.1126/science.1139253
Ross, M. T., Grafham, D. V, Coffey, A. J., Scherer, S., McLay, K., Muzny, D., and Platzer, M.
(2005). The DNA sequence of the human X chromosome. *Nature*, *434*(March), 325–337.
Ruderfer, D. M., Hamamsy, T., Lek, M., Karczewski, K. J., Kavanagh, D., Samocha, K. E.,
Exome Aggregation Consortium, Daly, M. J., MacArthur, D. G., Fromer, M., and Purcell, S.
M. (2016). Patterns of genic intolerance of rare copy number variation in 59,898 human
exomes. *Nature Genetics*, *48*(10). https://doi.org/10.1038/ng.3638
Simon, D., Laloo, B., Barillot, M., Barnetche, T., Blanchard, C., Rooryck, C., Marche, M.,
Burgelin, I., Coupry, I., Chassaing, N., Gilbert-Dussardier, B., Lacombe, D., Grosset, C.,
and Arveiler, B. (2010). A mutation in the 3′-UTR of the HDAC6 gene abolishing the post-
transcriptional regulation mediated by hsa-miR-433 is linked to a new form of dominant X-
linked chondrodysplasia. *Human Molecular Genetics*, *19*(10), 2015–2027.
https://doi.org/10.1093/hmg/ddq083
Skaletsky, H., Kuroda-kawaguchi, T., Minx, P. J., Cordum, H. S., Hillier, L., Brown, L. G.,
Repping, S., Pyntikova, T., Ali, J., Bieri, T., Chinwalla, A., Delehaunty, A., Delehaunty, K.,
Du, H., Fewell, G., Fulton, L., Fulton, R., … Page, D. C. (2003). The male-specific region
of the human Y chromosome is a mosaic of discrete sequence classes. *Nature*, *423*, 825–
838.
Uebbing, S., Konzer, A., Xu, L., Backström, N., Brunström, B., Bergquist, J., and Ellegren, H.
(2015). Quantitative Mass Spectrometry Reveals Partial Translational Regulation for
Dosage Compensation in Chicken. *Molecular Biology and Evolution*, *32*(10), 2716–2725.
https://doi.org/10.1093/molbev/msv147
Vandewalle, J., Van Esch, H., Govaerts, K., Verbeeck, J., Zweier, C., Madrigal, I., Mila, M.,
Pijkels, E., Fernandez, I., Kohlhase, J., Spaich, C., Rauch, A., Fryns, J. P., Marynen, P., and
Froyen, G. (2009). Dosage-Dependent Severity of the Phenotype in Patients with Mental
Retardation Due to a Recurrent Copy-Number Gain at Xq28 Mediated by an Unusual
Recombination. *American Journal of Human Genetics*, *85*(6), 809–822.
https://doi.org/10.1016/j.ajhg.2009.10.019
Watson, J. M., Spencer, J. A., Riggs, A. D., and Graves, J. A. (1990). The X chromosome of
monotremes shares a highly conserved region with the eutherian and marsupial X
chromosomes despite the absence of X chromosome inactivation. *Proceedings of the*
National Academy of Sciences, *87*(18), 7125–7129. https://doi.org/10.1073/pnas.87.18.7125
Weischenfeldt, J., Dubash, T., Drainas, A. P., Mardin, B. R., Chen, Y., Stütz, A. M., Waszak, S.
M., Bosco, G., Halvorsen, A. R., Raeder, B., Efthymiopoulos, T., Erkek, S., Siegl, C.,
Brenner, H., Brustugun, O. T., Dieter, S. M., Northcott, P. A., … Korbel, J. O. (2017). Pan-
cancer analysis of somatic copy-number alterations implicates IRS4 and IGF2 in enhancer
hijacking. *Nature Genetics*, (49), 65–74. https://doi.org/10.1038/ng.3722
White, M. a, Kitano, J., and Peichel, C. L. (2015). Purifying Selection Maintains Dosage-
Sensitive Genes during Degeneration of the Threespine Stickleback Y Chromosome.
Molecular Biology and Evolution, *32*(8), 1981–1995.
https://doi.org/10.1093/molbev/msv078
Wright, A. E., Zimmer, F., Harrison, P. W., and Mank, J. E. (2015). Conservation of Regional
Variation in Sex-Specific Sex Chromosome Regulation. *Genetics*, *201*(2), 587–598.

https://doi.org/10.1534/genetics.115.179234

Xiong, Y., Chen, X., Chen, Z., Wang, X., Shi, S., Wang, X., Zhang, J., and He, X. (2010). RNA
sequencing shows no dosage compensation of the active X-chromosome. *Nature Genetics*,
42(12), 1043–7. https://doi.org/10.1038/ng.711

Yang, F., Babak, T., Shendure, J., and Disteche, C. M. (2010). Global survey of escape from X
inactivation by RNA-sequencing in mouse. *Genome Research*, *20*(5), 614–622.
https://doi.org/10.1101/gr.103200.109

Yates, A., Akanni, W., Amode, M. R., Barrell, D., Billis, K., Carvalho-Silva, D., Cummins, C.,
Clapham, P., Fitzgerald, S., Gil, L., Girón, C. G., Gordon, L., Hourlier, T., Hunt, S. E.,
Janacek, S. H., Johnson, N., Juettemann, T., … Flicek, P. (2016). Ensembl 2016. *Nucleic*
Acids Research, *44*(D1), D710–D716. https://doi.org/10.1093/nar/gkv1157

Zhou, Q., Zhang, J., Bachtrog, D., An, N., Huang, Q., Jarvis, E. D., Gilbert, M. T. P., and Zhang,
G. (2014). Complex evolutionary trajectories of sex chromosomes across bird taxa. *Science*,
346(6215), 1246338–1246338. https://doi.org/10.1126/science.1246338

Zimmer, F., Harrison, P. W., Dessimoz, C., and Mank, J. E. (2016). Compensation of Dosage-
Sensitive Genes on the Chicken Z Chromosome. *Genome Biology and Evolution*, *8*(4),
1233–1242. https://doi.org/10.1093/gbe/evw075