1
8 **Conserved microRNA targeting reveals preexisting gene dosage**
9 **sensitivities that shaped amniote sex chromosome evolution**

Sahin Naqvi[1,2], Daniel W. Bellott[1], Kathy S. Lin[1,3], & David C. Page[1,2,4]

16
17
18
19

[1] Whitehead Institute, Cambridge MA 02142;
[2] Department of Biology, Massachusetts Institute of Technology, Cambridge MA 02139
[3] Program in Computational and Systems Biology, Massachusetts Institute of Technology, Cambridge MA 02139
[4] Howard Hughes Medical Institute, Whitehead Institute, Cambridge MA 02142

Mammalian X and Y chromosomes evolved from an ordinary autosomal pair. Genetic

decay of the Y led to X chromosome inactivation (XCI) in females, but some Y-linked genes

were retained during the course of sex chromosome evolution, and many X-linked genes

did not become subject to XCI. We reconstructed gene-by-gene dosage sensitivities on the

ancestral autosomes through phylogenetic analysis of microRNA (miRNA) target sites and

compared these preexisting characteristics to the current status of Y-linked and X-linked

genes in mammals. Preexisting heterogeneities in dosage sensitivity, manifesting as

differences in the extent of miRNA-mediated repression, predicted either the retention of a

Y homolog or the acquisition of XCI following Y gene decay. Analogous heterogeneities

among avian Z-linked genes predicted either the retention of a W homolog or gene-specific

dosage compensation following W gene decay. Genome-wide analyses of human copy

number variation indicate that these heterogeneities consisted of sensitivity to both

increases and decreases in dosage. We propose a model of XY/ZW evolution incorporating

such preexisting dosage sensitivities in determining the evolutionary fates of individual

genes. Our findings thus provide a more complete view of the role of dosage sensitivity in

shaping the mammalian and avian sex chromosomes, and reveal an important role for

post-transcriptional regulatory sequences (miRNA target sites) in sex chromosome

evolution.

INTRODUCTION

The mammalian X and Y chromosomes evolved from a pair of ordinary autosomes over the past

300 million years (Lahn & Page, 1999). Only 3% of genes on the ancestral pair of autosomes

survive on the human Y chromosome (Bellott et al., 2010; Skaletsky et al., 2003), compared to

98% on the X chromosome (Mueller et al., 2013). In females, one copy of the X chromosome is

silenced by X inactivation (XCI); this silencing evolved on a gene-by-gene basis following Y

gene loss and compensatory X upregulation (Berletch et al., 2015; Jegalian & Page, 1998; Ross

et al., 2005; Tukiainen et al., 2017), and some genes escape XCI in humans (Carrel & Willard,

2005) and other mammals (Yang, Babak, Shendure, & Disteche, 2010). In mammals, dosage

compensation, which refers to any mechanism restoring ancestral dosage following gene loss

from the sex-specific chromosome, thus consists of both X upregulation following Y gene loss

and the subsequent acquisition of XCI.

In parallel, the avian Z and W sex chromosomes evolved from a different pair of

autosomes than the mammalian X and Y chromosomes (Bellott et al., 2010; Nanda et al., 1999;

Ross et al., 2005). Decay of the female-specific W chromosome was similarly extensive, but

birds did not evolve a large-scale inactivation of Z-linked genes analogous to XCI in mammals

(Itoh et al., 2007). Dosage compensation, as measured by a male/female expression ratio close to

1, has been observed for some Z-linked genes in some tissues. (Mank & Ellegren, 2009; Uebbing

et al., 2015; Zimmer, Harrison, Dessimoz, & Mank, 2016). Thus, genes previously found on the

ancestral autosomes that gave rise to the mammalian or avian sex chromosomes have undergone

significant changes in gene dosage. In modern mammals, these molecular events have resulted in

three classes of ancestral X-linked genes representing distinct evolutionary fates: those with a

surviving Y homolog, those with no Y homolog and subject to XCI, and those with no Y

homolog but escaping XCI. In birds, two clear classes of ancestral Z-linked genes have arisen:

those with or without a W homolog, with additional heterogeneity among Z-linked genes without

a W homolog as a result of gene-specific dosage compensation. Identifying gene-by-gene

properties that distinguish classes of X- and Z-linked genes is thus crucial to understanding the

selective pressures underlying the molecular events of mammalian and avian sex chromosome

evolution.

Emerging evidence suggests a role for gene dosage sensitivity in mammalian and avian

sex chromosome evolution. X- and Z-linked genes with surviving homologs on the mammalian

Y or avian W chromosomes are enriched for important regulatory functions and predictors of

haploinsufficiency compared to those lacking Y or W homologs (Bellott et al., 2014, 2017);

similar observations have been made in fish (White, Kitano, & Peichel, 2015) and Drosophila

(Kaiser, Zhou, & Bachtrog, 2011). Human X- and chicken Z-linked genes that show the

strongest signatures of dosage compensation in either lineage also show signs of dosage

sensitivity as measured by membership in large protein complexes (Pessia, Makino, Bailly-

Bechet, McLysaght, & Marais, 2012) or evolutionary patterns of gene duplication and retention

(Zimmer et al., 2016). Despite these advances, little is known regarding selective pressures

resulting from sensitivity to dosage increases, as these studies either focused on

haploinsufficiency or employed less direct predictors of dosage sensitivity. Furthermore, it is not

known whether heterogeneities in dosage sensitivity among classes of sex-linked genes were

acquired during sex chromosome evolution, or predated the emergence of sex chromosomes, as

there has been no explicit, systematic reconstruction of dosage sensitivity on the ancestral

autosomes that gave rise to the mammalian and avian sex chromosomes.

To assess the role of preexisting dosage sensitivities in XY and ZW evolution, we sought

to employ a measure of dosage sensitivity that could be 1) demonstrably informative with respect

to sensitivity to dosage increases, and 2) explicitly reconstructed on the ancestral autosomes. We

focused on regulation by microRNAs (miRNAs), small noncoding RNAs that function as tuners

of gene dosage by lowering target mRNA levels through pairing to the 3` untranslated region

(UTR) (Bartel, 2009). The repressive nature of miRNA targeting is informative with respect to

sensitivity to dosage increases, allowing for a more complete understanding of the role of dosage

sensitivity in sex chromosome evolution. Both miRNAs themselves and their complementary

target sites can be preserved over millions of years of vertebrate evolution, facilitating the

reconstruction of miRNA targeting on the ancestral autosomes through cross-species sequence

alignments. As miRNA targeting occurs post-transcriptionally, reconstruction of its ancestral

state is decoupled from transcriptional regulatory mechanisms such as XCI that evolved

following X-Y differentiation.

RESULTS

Analysis of human copy number variation indicates conserved microRNA targeting of

genes sensitive to dosage increases

We first sought to determine whether conserved targeting by microRNAs (miRNAs) correlates

with sensitivity to dosage increases across the human genome. To estimate pressure to maintain

miRNA targeting, we used published probabilities of conserved targeting (P_{CT} scores) for each

gene-miRNA interaction in the human genome. The P_{CT} score reflects an estimate of the

probability that a given gene-miRNA interaction is conserved due to miRNA targeting, obtained

by calculating the conservation of the relevant miRNA target sites relative to the conservation of

the entire 3` UTR (Friedman, Farh, Burge, & Bartel, 2009). In this manner, the P_{CT} score

intrinsically controls for differences in background conservation and sequence composition, both

of which vary widely among 3`UTRs due to differing rates of expression divergence and/or

sequence evolution. We refer to these P_{CT} scores as "miRNA conservation scores" in the

remainder of the text.

A recent study reported a correlation between these miRNA conservation scores and

predicted haploinsufficiency (Pinzón et al., 2016), indicating that conserved miRNA targeting

broadly corresponds to dosage sensitivity. However, such a correlation does not isolate the

effects of sensitivity to dosage increases, which we expect to be particularly important in the

context of miRNA targeting. We reasoned that genes for which increases in dosage are

deleterious should be depleted from the set of observed gene duplications in healthy human

individuals. We used a catalogue of rare genic copy number variation among 59,898 control

human exomes (Exome Aggregation Consortium, ExAC)(Ruderfer et al., 2016) to classify

autosomal protein-coding genes as exhibiting or lacking duplication or deletion in healthy

individuals (see Methods). We compared duplicated and non-duplicated genes with the same

deletion status in order to control for differences in sensitivity to underexpression. We found that

non-duplicated genes have significantly higher miRNA conservation scores than duplicated

genes, irrespective of deletion status (Figure 1A,B). Non-deleted genes also have significantly

higher scores than deleted genes irrespective of duplication status (Supplemental Figure S1), but

duplication status has a greater effect on miRNA conservation scores than does deletion status

(blue vs. orange boxes, Figure 1C). Thus, conserved miRNA targeting is a feature of genes

sensitive to changes in gene dosage in humans and is especially informative with regards to

sensitivity to dosage increases, consistent with the known role of miRNAs in tuning gene dosage

by lowering target mRNA levels.

X-Y pairs and X-inactivated genes have higher miRNA conservation scores than X escape genes

We next assessed whether the three classes of X-linked genes differ with respect to dosage sensitivity as inferred by conserved miRNA targeting. To delineate these classes, we began with the set of ancestral genes reconstructed through cross-species comparisons of the human X chromosome and orthologous chicken autosomes (Bellott et al., 2014, 2017, 2010; Hughes et al., 2012; Mueller et al., 2013). We designated ancestral X-linked genes with a surviving human Y homolog (Skaletsky et al., 2003) as X-Y pairs and also considered the set of X-linked genes with a surviving Y homolog in any of eight mammals (Bellott et al., 2014) to increase the phylogenetic breadth of findings regarding X-Y pairs. A number of studies have catalogued the inactivation status of X-linked genes in various human tissues and cell-types. We used a meta-analysis that combined results from three studies by assigning a "consensus" X-inactivation status to each gene (Balaton, Cotton, & Brown, 2015) to designate the remainder of ancestral genes lacking a Y homolog as subject to or escaping XCI. In summary, we classified genes as either: 1) X-Y pairs, 2) lacking a Y homolog and subject to XCI (X-inactivated), or 3) lacking a Y homolog but escaping XCI (X escape).

We found that human X-Y pairs have the highest miRNA conservation scores, followed by X-inactivated and finally X escape genes (Figure 2A,B). The expanded set of X-Y pairs across eight mammals also has significantly higher miRNA conservation scores than ancestral X-linked genes with no Y homolog (Supplemental Figure S2). Observed differences between miRNA conservation scores are not driven by distinct subsets of genes in each class, as indicated by gene resampling with replacement (Supplemental Figure S3). The decrease in miRNA conservation scores of X escape genes relative to X-inactivated genes and X-Y pairs is not

driven by genes that escape XCI variably across individuals (Supplemental Figure S4), and was

consistent even when including ambiguous genes as either X-inactivated or X escape genes

(Supplemental Figure S5). We also verified that these differences were not driven by artificially

inflated or deflated conservation scores of certain target sites due to non-uniformity in 3` UTR

conservation (Methods, Supplemental Figure S6).

Finally, we assessed whether miRNA conservation scores distinguish the three classes by

providing additional information not accounted for by known factors (Bellott et al., 2014)

influencing evolutionary outcomes. We used logistic regression to model, for each gene, the

probability of falling into each of the three classes (X-Y pair, X-inactivated, or X escape) as a

linear combination of haploinsufficiency probability (pHI) (Huang, Lee, Marcotte, & Hurles,

2010), human expression breadth (GTEx Consortium, 2015), purifying selection, measured by

the ratio of non-synonymous to synonymous substitution rates (dN/dS) between human and

mouse orthologs (Yates et al., 2016), and mean gene-level miRNA conservation scores. We note

that pHI is a score composed of several genic features, one of which is the number of protein-

protein interactions, consistent with the idea that members of large protein complexes tend to be

dosage-sensitive (Papp, Pal, & Hurst, 2003; Pessia et al., 2012). Removing either miRNA

conservation or pHI as predictors from the full model resulted in inferior model fits as measured

by Aikake's information criterion (AIC) (full model, AIC 321.5; full model minus miRNA

conservation, AIC 327.7; full model minus pHI, AIC 327.3; higher AIC indicates inferior model).

Therefore, miRNA conservation and pHI contribute independent information that distinguishes

the 3 classes of X-linked genes. Based on our analyses of autosomal copy number variation

(Figure 1), we attribute this independence to the fact that miRNA conservation scores are most

informative with regards to sensitivity to dosage increases. Taken together, these results indicate

significant heterogeneity in dosage sensitivity, as inferred by miRNA target site conservation,

among the three classes of ancestral X-linked genes: X-Y pairs are the most dosage-sensitive,

while X-inactivated genes are of intermediate dosage sensitivity, and X escape genes are the

least dosage-sensitive.

Heterogeneities in X-linked miRNA targeting were present on the ancestral autosomes

We next asked whether differences in miRNA targeting were present on the ancestral autosomes

that gave rise to the mammalian X and Y chromosomes. To reconstruct the ancestral state of

miRNA targeting, we first focused on miRNA target sites in the 3` UTR of human orthologs that

align with perfect identity to a site in the corresponding chicken ortholog; these sites were likely

present in the common ancestor of mammals and birds (Figure 3A,B). We found that X-Y pairs

have the most human-chicken conserved target sites, followed by X-inactivated genes, and then

X escape genes (Figure 3C, top). Unlike the miRNA conservation scores used earlier, this metric

does not account for background conservation; we therefore estimated the background

conservation of each 3` UTR using shuffled miRNA family seed sequences (see Methods). X-Y

pairs, X-inactivated genes, and X escape genes do differ significantly with respect to background

conservation (data not shown), but these differences cannot account for the observed differences

in true human-chicken conserved sites (Figure 3C, bottom). We observed similar results for the

expanded set of X-Y pairs across 8 mammals (Supplemental Figure S7A).

Differences in the number of human-chicken conserved sites among the three classes of

X-linked genes could be explained by heterogeneity in miRNA targeting present on the ancestral

autosomes, or by ancestral homogeneity followed by different rates of target site loss during or

following X-Y differentiation. To distinguish between these two possibilities, we took advantage

of previous reconstructions of human sex chromosome evolution (Figure 3A) (Bellott et al.,

2014), which confirmed that, following the divergence of placental mammals from marsupials,

an X-autosome chromosomal fusion generated the X-added region (XAR) (Watson, Spencer,

Riggs, & Graves, 1990). Genes on the XAR are therefore X-linked in placental mammals, but

autosomal in marsupials such as the opossum. We limited our analysis to genes in the XAR and

target sites conserved between orthologous chicken and opossum 3` UTRs, ignoring site

conservation in humans; these sites were likely present in the common ancestor of mammals and

birds, and an absence of such sites cannot be explained by site loss following X-Y differentiation.

We observed the same pattern as with the human-chicken conserved sites, both before and after

accounting for background 3` UTR conservation (Figure 3D, three gene classes; Supplemental

Figure S7B, X-Y pairs across 8 mammals). These results demonstrate that the autosomal

precursors of X-Y pairs and X-inactivated genes were subject to more miRNA-mediated

regulation than X escape genes. Combined with our earlier results, we conclude that present-day

heterogeneities in dosage sensitivity on the mammalian X chromosome existed on the ancestral

autosomes from which it derived.

Z-W pairs have higher miRNA conservation scores than other ancestral Z-linked genes

We next assessed whether classes of avian Z-linked genes, those with and without a W homolog,

show analogous heterogeneities in sensitivity to dosage increases. We used the set of ancestral

genes reconstructed through cross-species comparisons of the avian Z chromosome and

orthologous human autosomes and focused on the set of Z-W pairs identified by sequencing of

the chicken W chromosome (Bellott et al., 2017, 2010). To increase the phylogenetic breadth of

our comparisons, we also included candidate Z-W pairs obtained through comparisons of male

and female genome assemblies (4 species set) or inferred by read-depth changes in female

genome assemblies (14 species set, see Methods for details) (Zhou et al., 2014). The more

complete 3` UTR annotations in the human genome relative to chicken allow for a more accurate

assessment of conserved miRNA targeting. Accordingly, we analyzed the 3` UTRs of the human

orthologs of chicken Z-linked genes.

We found that the human orthologs of Z-W pairs have higher miRNA conservation

scores than the human orthologs of other ancestral Z genes (Figure 4A, B). Differences in

miRNA conservation scores between Z-W pairs and other ancestral Z genes remained significant

when considering the expanded sets of Z-W pairs across four and 14 avian species

(Supplemental Figure S8). These differences are not driven by distinct subsets of genes, as

indicated by gene resampling with replacement (Supplemental Figure S9), and cannot be

accounted for by within-UTR variation in regional conservation (Supplemental Figure S10).

Logistic regression models indicate that miRNA conservation scores provide additional

information not captured by known factors (Bellott et al., 2017) influencing survival of W-linked

genes (full model model, AIC 127.1; full model minus miRNA conservation, AIC 137.8; full

model minus pHI 132.7; higher AIC indicates inferior model). Together, these results indicate

that Z-linked genes with a surviving W homolog are more sensitive to changes in dosage -- both

increases and decreases -- than are genes without a surviving W homolog.

While there are two clear classes of Z-linked genes -- those with or without a W homolog

-- studies of Z-linked gene expression have suggested additional heterogeneity among Z-linked

genes without a W homolog due to gene-specific dosage compensation (Mank & Ellegren, 2009;

Uebbing et al., 2015; Zimmer et al., 2016). If Z-linked genes with no W homolog exist upon a

continuum from non-compensated to dosage-compensated, those that are more compensated

should have more conserved miRNA target sites, reflective of greater dosage sensitivity. We

quantified the dosage compensation by using RNA sequencing data (Marin et al., 2017) to

compare, in 4 somatic tissues, the chicken male/female expression ratio to the analogous ratio in

human and anolis (see Methods). In the brain, kidney, and liver, Z-linked genes with no W

homolog and higher mean miRNA conservation scores had male/female expression ratios closer

to 1 (Supplemental Figure S11). Thus, in addition to the above-described differences between Z-

linked genes with or without a W homolog, Z-linked genes with no W homolog but with more

effective dosage compensation have more conserved miRNA target sites than non-compensated

genes.

Heterogeneities in Z-linked miRNA targeting were present on the ancestral autosomes

We next asked whether differences in miRNA targeting between Z-W pairs and other ancestral

Z-linked genes were present on the ancestral autosomes that gave rise to the avian Z and W

chromosomes. We found that chicken Z-W pairs have more human-chicken-conserved miRNA

target sites than their Z-linked counterparts without surviving W homologs, both before (Figure

5C, top) and after (Figure 5C, bottom) accounting for the background conservation of each

individual 3` UTR. To confirm that these differences represent ancestral heterogeneity rather

than differential site loss during or following Z-W differentiation, we instead considered the

number of sites conserved between human and anolis lizard, which diverged from birds prior to

Z-W differentiation (Figure 5A). Chicken Z-W pairs contain an excess of human-anolis

conserved miRNA target sites, both before (Figure 5D, top) and after (Figure 5D, bottom)

accounting for the background conservation of each individual 3` UTR. We observed similar

results with the predicted four-species (Supplemental Figure S12) and 14-species (Supplemental

Figure S13) sets of Z-W pairs. Thus, the autosomal precursors of avian Z-W pairs were subject

to more miRNA-mediated regulation than the autosomal precursors of Z-linked genes that lack a

W homolog. Furthermore, in the liver and brain, Z-linked genes with no W homolog with an

excess of human-chicken-conserved miRNA sites had male/female expression ratios closer to 1,

implying more effective dosage compensation (Supplementary Figure S14). Together, these

results indicate heterogeneity in dosage sensitivity among genes on the ancestral autosomes that

gave rise to the avian Z chromosome.

Analyses of experimental datasets validate miRNA target site function

Our results to this point, which indicate preexisting heterogeneities in dosage constraints among

X- or Z-linked genes as inferred by predicted miRNA target sites, lead to predictions regarding

the function of these sites in vivo. To test these predictions, we turned to publically available

experimental datasets consisting both of gene expression profiling following transfection or

knockout of individual miRNAs, and of high-throughput crosslinking-immunoprecipitation

(CLIP) to identify sites that bind Argonaute in vivo (see Methods). If the above-studied sites are

effective in mediating target repression, targets of an individual miRNA should show increased

expression levels or Argonaute binding following miRNA transfection, and decreased expression

levels following miRNA knockout. Together, our analyses of publically available datasets

fulfilled these predictions, validating the function of these sites in multiple cellular contexts and

species (Figure 6). From the gene expression profiling data, we observed results consistent with

effective targeting by a) eleven different miRNA families in human HeLa cells (Supplemental

Figure S15), b) four different miRNAs in human HCT116 and HEK293 cells (Supplemental

Figure S16), and c) miR-155 in mouse B and Th1 cells (Supplemental Figure S17). In the CLIP

data, the human orthologs of X- or Z-linked targets of miR-124 are enriched for Argonaute-

bound clusters that appear following miR-124 transfection, while a similar but non-significant

enrichment is observed for miR-7 (Supplemental Figure S18). Thus, conserved miRNA target

sites used to infer dosage constraints on X-linked genes and the autosomal orthologs of Z-linked

genes can effectively mediate target repression in living cells.

DISCUSSION

Here, through the evolutionary reconstruction of microRNA (miRNA) target sites, we provide

evidence for preexisting heterogeneities in dosage sensitivity among genes on the mammalian X

and avian Z chromosomes. We first showed that, across all human autosomal genes, dosage

sensitivity -- as indicated by patterns of genic copy number variation -- correlates with the degree

of conserved miRNA targeting. We found that conserved targeting correlates especially strongly

with sensitivity to dosage increases, consistent with miRNA targeting serving to reduce gene

expression. Turning to the sex chromosomes of mammals and birds, genes that retained a

homolog on the sex-specific Y or W chromosome (X-Y and Z-W pairs) have more conserved

miRNA target sites than genes with no Y or W homolog. In mammals, genes with no Y homolog

that became subject to XCI have more conserved sites than those that continued to escape XCI

following Y gene decay. In birds, across Z-linked genes with no W homolog, the degree of

conserved miRNA targeting correlates with the degree of gene-specific dosage compensation.

We then reconstructed the ancestral state of miRNA targeting, observing significant

heterogeneities in the extent of miRNA targeting, and thus dosage sensitivity, on the ancestral

autosomes that gave rise to the mammalian and avian sex chromosomes. Finally, through

analysis of publically available experimental datasets, we validated the function, in living cells,

of the miRNA target sites used to infer dosage sensitivity. We thus conclude that differences in

dosage sensitivity – both to increases and to decreases in gene dosage -- among genes on the

ancestral autosomes influenced their evolutionary trajectory during sex chromosome evolution,

not only on the sex-specific Y and W chromosomes, but also on the sex-shared X and Z

chromosomes.

Our findings build upon previous work in three important ways. First, our analysis of

miRNA-mediated repression indicates that these heterogeneities consist of sensitivities to dosage

increases and decreases, whereas previous studies had either focused on sensitivity to

underexpression or could not differentiate the two. Second, our reconstruction of miRNA

targeting on the ancestral autosomes provides direct evidence that heterogeneities in dosage

sensitivity among classes of X- and Z-linked were preexisting rather than acquired during sex

chromosome evolution. Finally, by pointing to specific regulatory sequences (miRNA target

sites) functioning to tune gene dosage both prior to and during sex chromosome evolution, our

study provides a view of dosage compensation encompassing post-transcriptional regulation.

Human disease studies support the claim that increased dosage of X-Y pairs and X-

inactivated genes is deleterious to fitness. Copy number gains of the X-linked gene *KDM6A*,

which has a surviving human Y homolog, are found in patients with developmental

abnormalities and intellectual disability (Lindgren et al., 2013). *HDAC6*, *CACNA1F*, *GDI1*, and

IRS4 all lack Y homologs and are subject to XCI in humans. A mutation in the 3` UTR of

HDAC6 abolishing targeting by miR-433 has been linked to familial chondrodysplasia in both

sexes (Simon et al., 2010). Likely gain-of-function mutations in *CACNA1F* cause congenital

stationary night blindness in both sexes (Hemara-Wahanui et al., 2005). Copy number changes of

GDI1 correlate with the severity of X-linked mental retardation in males, with female carriers

preferentially inactivating the mutant allele (Vandewalle et al., 2009). Somatic genomic deletions

downstream of *IRS4* lead to its overexpression in lung squamous carcinoma (Weischenfeldt et al.,

2017). Males with partial X disomy due to translocation of the distal long arm of the X

chromosome (Xq28) to the long arm of the Y chromosome show severe mental retardation and

developmental defects (Lahn et al., 1994). Most genes in Xq28 are inactivated in 46,XX females

but escape inactivation in such X;Y translocations, suggesting that increased dosage of Xq28

genes caused the cognitive and developmental defects. We anticipate that further studies will

reveal additional examples of the deleterious effects of increases in gene dosage of X-Y pairs

and X-inactivated genes.

We and others previously proposed that Y gene decay drove upregulation of homologous

X-linked genes in both males and females, and that XCI subsequently evolved at genes sensitive

to increased expression from two active X-linked copies in females (Jegalian & Page, 1998;

Ohno, 1967). Our finding that X-inactivated genes have higher miRNA conservation scores than

X escape genes is consistent with this aspect of the model. However, recent studies indicating

heterogeneity in dosage sensitivity between classes of mammalian X- or avian Z-linked genes

(Bellott et al., 2014, 2017; Pessia et al., 2012; Zimmer et al., 2016), combined with the present

finding that these dosage sensitivities existed on the ancestral autosomes, challenge the previous

assumption of a single evolutionary pathway for all sex-linked genes.

We therefore propose a revised model of X-Y and Z-W evolution in which the ancestral

autosomes that gave rise to the mammalian and avian sex chromosomes contained three (or two,

in the case of birds) classes of genes with differing dosage sensitivities (Figure 7A,B). For

ancestral genes with high dosage sensitivity, Y or W gene decay would have been highly

deleterious, and thus the Y- or W-linked genes were retained. According to our model, these

genes' high dosage sensitivity also precluded upregulation of the X- or Z-linked homolog, and,

in mammals, subsequent X-inactivation; indeed, their X-linked homologs continue to escape

XCI (Bellott et al., 2014). For ancestral mammalian genes of intermediate dosage sensitivity, Y

gene decay did occur, and was accompanied or followed by compensatory upregulation of the X-

linked homolog in both sexes; the resultant increased expression in females was deleterious and

led to the acquisition of XCI. Ancestral mammalian genes of low dosage sensitivity continued to

escape XCI following Y decay; heterogeneity in X upregulation may further subdivide such

genes (Figure 6A). These genes' dosage insensitivity set them apart biologically, and

evolutionarily, from the other class of X-linked genes escaping XCI -- those with a surviving Y

homolog.

Our revised model relates preexisting, gene-by-gene heterogeneities in dosage sensitivity

to the outcomes of sex chromosome evolution. However, the suppression of X-Y recombination

did not occur on a gene-by-gene basis, instead initiating Y gene decay and subsequent dosage

compensation through a series of large-scale inversions encompassing many genes (Lahn & Page,

1999). The timings and boundaries of these evolutionary strata varied among mammalian

lineages, thus leading to unique chromosome-scale evolutionary dynamics across mammals.

These large-scale changes would have then allowed for genic selection to take place according to

the preexisting dosage sensitivities outlined above. In this way, the course of sex chromosome

evolution in mammals is a composite of 1) preexisting, gene-by-gene dosage sensitivities and 2)

the manner in which the history of the X and Y unfolded in particular lineages via discrete, large-

scale inversions.

In this study, we have focused on classes of ancestral X-linked genes delineated by the

survival of a human Y homolog or by the acquisition of XCI in humans, but such evolutionary

states can differ among mammalian lineages and species. In mouse, for instance, both Y gene

decay (Bellott et al., 2014) and the acquisition of X-inactivation (Yang et al., 2010) are more

complete than in humans or other mammals, as exemplified by *RPS4X*, which retains a Y

homolog and continues to escape XCI in primates, but has lost its Y homolog and is subject to

XCI in rodents. These observations could be explained by shortened generation times in the

rodent lineage, resulting in longer evolutionary times, during which the forces leading to Y gene

decay and the acquisition of X-inactivation could act (Charlesworth & Crow, 1978; Jegalian &

Page, 1998; Ohno, 1967). Another case of lineage differences involves *HUWE1*, which lacks a Y

homolog and is subject to XCI in both human and mouse, but retains a functional Y homolog in

marsupials, where it continues to escape XCI. In the future, more complete catalogues of X-

inactivation and escape in additional mammalian lineages would make it possible to examine

whether analogous, preexisting dosage sensitivities differentiate the three classes of X-linked

genes (X-Y pairs, X-inactivated genes, and X escape genes) in other species.

Previous studies have sought evidence of X-linked upregulation during mammalian sex

chromosome evolution using comparisons of gene expression levels between the whole of the X

chromosome and all of the autosomes, with equal numbers of studies rejecting or finding

evidence consistent with upregulation (Deng et al., 2011; Julien et al., 2012; Kharchenko, Xi, &

Park, 2011; Lin, Xing, Zhang, & He, 2012; Xiong et al., 2010). This is likely due to gene-by-

gene heterogeneity in dosage sensitivities that resulted in a stronger signature of upregulation at

more dosage sensitive genes (Pessia et al., 2012). Similarly, studies of Z-linked gene expression

in birds provide evidence for the gene-by-gene nature of Z dosage compensation, as measured by

comparisons of gene expression levels between ZZ males and ZW females (Itoh et al., 2007;

Mank & Ellegren, 2009; Uebbing et al., 2015), and indicate a stronger signature of dosage

compensation at predicted dosage-sensitive genes (Zimmer et al., 2016). By showing that such

dosage sensitivities existed on the ancestral autosomes and consist of sensitivity to both increases

and decreases, our findings highlight an additional aspect of dosage compensation that affects

both birds and mammals.

In addition to revealing similarities between mammals and birds, our study provides a

view of dosage compensation that highlights post-transcriptional regulatory mechanisms,

pointing to specific non-coding sequences with known mechanisms (microRNA target sites)

functioning across evolutionary time. A recent study in birds showed a role for a Z-linked

miRNA, miR-2954-3p, in dosage compensation of some Z-linked genes (Warnefors et al., 2017).

Our study suggests an additional, broader role for miRNA targeting, with hundreds of different

miRNAs acting to tune gene dosage both before and during sex chromosome evolution.

Furthermore, our finding of greater conserved miRNA targeting of X-inactivated genes relative

to X escape genes shows that it is possible to predict the acquisition of a transcriptional

regulatory state (XCI) during sex chromosome evolution on the basis of a preexisting, post-

transcriptional regulatory state. Perhaps additional post-transcriptional regulatory mechanisms

and their associated regulatory elements will be shown to play roles in mammalian and avian

dosage compensation.

Recent work has revealed that the sex-specific chromosome -- the Y in mammals and the

W in birds -- convergently retained dosage-sensitive genes with important regulatory functions

(Bellott et al., 2014, 2017). Our study, by reconstructing the ancestral state of post-transcriptional

regulation, provides direct evidence that such heterogeneity in dosage sensitivity existed on the

ancestral autosomes that gave rise to the mammalian and avian sex chromosomes. This

heterogeneity influenced both survival on the sex-specific chromosomes in mammals and birds

and the evolution of XCI in mammals. Thus, two independent experiments of nature offer

empirical evidence that modern-day amniote sex chromosomes were shaped, during evolution,

by the properties of the ancestral autosomes from which they derive.

METHODS

Statistics

Details of all statistical tests (type of test, test statistic, and p-value) used in this manuscript are

provided in Supplemental Table S1.

Human genic copy number variation

To annotate gene deletions and duplications, we used data from the Exome Aggregation

Consortium (ExAC) (ftp://ftp.broadinstitute.org/pub/ExAC_release/release0.3.1/cnv/), which

consists of autosomal genic duplications and deletions (both full and partial) called in 59,898

exomes (Ruderfer et al., 2016). Further details are provided in Supplemental Methods in the

section titled 'Human genic copy number variation'. These gene assignments are provided in

Supplemental Table S2.

X- and Z-linked gene sets

We utilized our previous reconstructions of the ancestral mammalian X (Bellott et al., 2014) and

avian Z (Bellott et al., 2017) chromosomes, as well as information on multicopy and ampliconic

X-linked genes (Mueller et al., 2013) and XCI status in humans (Balaton et al., 2015) to

delineate classes of X- and Z-linked genes. Further details are provided in Supplemental

Methods under the sections titled 'X-linked gene sets' and 'Z-linked gene sets'. Information on

X-linked genes is provided in Supplemental Table S3. Information on Z-linked genes is provided

in Supplemental Table S4.

microRNA target sites

Pre-calculated P_{CT} scores for all gene-miRNA family interactions

(http://www.targetscan.org/vert_71/vert_71_data_download/Summary_Counts.all_predictions.tx

t.zip) and site-wise alignment information

(http://www.targetscan.org/vert_71/vert_71_data_download/Conserved_Family_Info.txt.zip)

were obtained from TargetScan Human v7. Details on the filtering of miRNAs and resampling-

based assessment of P_{CT} scores are provided in Supplemental Methods in the section titled

'microRNA target site P_{CT} scores'. Details regarding analysis of human-chicken or human-anolis

conserved sites, as well as approaches to control for background conservation, are provided in

Supplemental Methods in the section titled 'Human-chicken conserved microRNA target sites.'

Variation in within-UTR conservation bias

To address the possibility that non-uniformity in regional 3` UTR conservation could artificially

inflate or deflate conservation scores of certain target sites, we implemented a step-detection

algorithm to segment 3` UTRs into regions of homogeneous background conservation and

calculated miRNA site conservation relative to these smaller regions. These regionally

normalized scores, corresponding to all gene-miRNA interactions, are provided in Supplemental

Table S5. Details of the step-detection algorithm are provided in Supplemental Methods in the

section titled 'Variation in within-UTR conservation bias'.

Logistic regression

Logistic regression models were constructed using the function 'multinom' in the R package

'nnet.' We used previously published values for known factors in the survival of Y-linked

(Bellott et al., 2014) and W-linked (Bellott et al., 2017) genes except for human expression

breadth, which we recalculated using data from the GTEx Consortium v6 data release

(Consortium, 2015). Briefly, kallisto was used to estimate transcript per million (TPM) values in

the 10 male samples with the highest RNA integrity numbers (RINs) from each of 37 tissues, and

expression breadth across tissues was calculated as described in (Bellott et al., 2014), using

median TPM values for each tissue.

Assessing Z-linked dosage compensation using cross-species RNA-sequencing data

Raw data from Marin et al 2017 (add citation) was obtained, and kallisto and limma/voom were

used for abundance quantification and differential expression, respectively. Further details are

provided in Supplemental Methods in the section titled 'Assessing Z-linked dosage

compensation using cross-species RNA-sequencing data'.

Gene expression profiling and crosslinking datasets

Fold-changes in mRNA expression and targets of Argonaute as determined by high-throughput

crosslinking-immunoprecipitation (CLIP) were obtained from a variety of publically available

datasets. Further details are provided in Supplemental Methods in the section titled 'Gene

expression profiling and crosslinking datasets.' All fold-changes and CLIP targets are provided

in Supplemental Table S6.

Code availability

A custom Python (RRID:SCR_008394) script utilizing Biopython (RRID:SCR_007173) was

used to generate shuffled miRNA family seed sequences. Identification of miRNA target site

matches using shuffled seed sequences was performed using the 'targetscan_70.pl' perl script

(http://www.targetscan.org/vert_71/vert_71_data_download/targetscan_70.zip). 3` UTR

segmentation was performed with the 'plot_transitions.py' python script. Code is available at:

https://github.com/snaqvi1990/Naqvi17-code and as Supplementary Code.

Data access

Data supporting the findings of this study are available within the paper and its Supplemental

information files. Accession numbers for publically available datasets are provided when

appropriate in Methods sections.

ACKNOWLEDGEMENTS

We thank V. Agarwal, S. Eichorn, S. McGeary, and D. Bartel for assistance with the TargetScan

database and helpful discussions; A. Godfrey for updated human-chicken orthology information;

and A. Godfrey, J. Hughes and H. Skaletsky for critical reading of the manuscript. This work

was supported by the National Institutes of Health and the Howard Hughes Medical Institute.

S.N. was supported under a research grant by Biogen.

Author contributions

S.N., D.W.B. and D.C.P designed the study. S.N. performed analyses with assistance from

D.W.B. K.S.L developed and implemented the step-detection algorithm. S.N. and D.C.P wrote

the paper.

DISCLOSURE DECLARATION

The authors declare no competing financial interests.

Balaton, B. P., Cotton, A. M., and Brown, C. J. (2015). Derivation of consensus inactivation status for X-linked genes from genome-wide studies. *Biology of Sex Differences*, *6*, 35. https://doi.org/10.1186/s13293-015-0053-7

Bartel, D. P. (2009). MicroRNAs: target recognition and regulatory functions. *Cell*, *136*(2), 215–233. https://doi.org/10.1016/j.cell.2009.01.002

Bellott, D. W., Hughes, J. F., Skaletsky, H., Brown, L. G., Pyntikova, T., Cho, T.-J., Koutseva, N., Zaghlul, S., Graves, T., Rock, S., Kremitzki, C., Fulton, R. S., Dugan, S., Ding, Y., Morton, D., Khan, Z., Lewis, L., … Page, D. C. (2014). Mammalian Y chromosomes retain widely expressed dosage-sensitive regulators. *Nature*, *508*(7497), 494–499. https://doi.org/10.1038/nature13206

Bellott, D. W., Skaletsky, H., Cho, T.-J., Brown, L., Locke, D., Chen, N., Galkina, S., Pyntikova, T., Koutseva, N., Graves, T., Kremitzki, C., Warren, W. C., Clark, A. G., Gaginskaya, E., Wilson, R. K., and Page, D. C. (2017). Avian W and mammalian Y chromosomes convergently retained dosage-sensitive regulators. *Nature Genetics*, *in press*.

Bellott, D. W., Skaletsky, H., Pyntikova, T., Mardis, E. R., Graves, T., Kremitzki, C., Brown, L. G., Rozen, S., Warren, W. C., Wilson, R. K., and Page, D. C. (2010). Convergent evolution of chicken Z and human X chromosomes by expansion and gene acquisition. *Nature*, *466*(7306), 612–616. https://doi.org/10.1038/nature09172

Berletch, J. B., Ma, W., Yang, F., Shendure, J., Noble, W. S., Disteche, C. M., and Deng, X. (2015). Escape from X Inactivation Varies in Mouse Tissues. *PLOS Genetics*, *11*(3), e1005079. https://doi.org/10.1371/journal.pgen.1005079

Carrel, L., and Willard, H. F. (2005). X-inactivation profile reveals extensive variability in X-linked gene expression in females. *Nature*, *434*(March), 400–404. https://doi.org/10.1038/nature03479

Charlesworth, B., and Crow, J. F. (1978). Model for evolution of Y chromosomes and dosage compensation, *75*(11), 5618–5622.

Consortium, Gte. (2015). The Genotype-Tissue Expression (GTEx) pilot analysis: Multitissue gene regulation in humans. *Science (New York, N.Y.)*, *348*(6235), 648–660.

Deng, X., Hiatt, J. B., Nguyen, D. K., Ercan, S., Sturgill, D., Hillier, L. W., Schlesinger, F., Davis, C. a, Reinke, V. J., Gingeras, T. R., Shendure, J., Waterston, R. H., Oliver, B., Lieb, J. D., and Disteche, C. M. (2011). Evidence for compensatory upregulation of expressed X-linked genes in mammals, Caenorhabditis elegans and Drosophila melanogaster. *Nature Genetics*, *43*(12), 1179–1185. https://doi.org/10.1038/ng.948

Friedman, R. C., Farh, K. K.-H., Burge, C. B., and Bartel, D. P. (2009). Most mammalian mRNAs are conserved targets of microRNAs. *Genome Research*, *19*(1), 92–105. https://doi.org/10.1101/gr.082701.108

Hemara-Wahanui, A., Berjukow, S., Hope, C. I., Dearden, P. K., Wu, S.-B., Wilson-Wheeler, J., Sharp, D. M., Lundon-Treweek, P., Clover, G. M., Hoda, J.-C., Striessnig, J., Marksteiner, R., Hering, S., and Maw, M. a. (2005). A CACNA1F mutation identified in an X-linked retinal disorder shifts the voltage dependence of Cav1.4 channel activation. *Proceedings of the National Academy of Sciences of the United States of America*, *102*(21), 7553–7558. https://doi.org/10.1073/pnas.0501907102

Huang, N., Lee, I., Marcotte, E. M., and Hurles, M. E. (2010). Characterising and predicting haploinsufficiency in the human genome. *PLoS Genetics*, *6*(10), e1001154. https://doi.org/10.1371/journal.pgen.1001154

Hughes, J. F., Skaletsky, H., Brown, L. G., Pyntikova, T., Graves, T., Fulton, R. S., Dugan, S.,

Ding, Y., Buhay, C. J., Kremitzki, C., Wang, Q., Shen, H., Holder, M., Villasana, D.,
Nazareth, L. V, Cree, A., Courtney, L., … Page, D. C. (2012). Strict evolutionary
conservation followed rapid gene loss on human and rhesus Y chromosomes. *Nature*,
483(7387), 82–86. https://doi.org/10.1038/nature10843

Itoh, Y., Melamed, E., Yang, X., Kampf, K., Wang, S., Yehya, N., Van Nas, A., Replogle, K.,
Band, M. R., Clayton, D. F., Schadt, E. E., Lusis, A. J., and Arnold, A. P. (2007). Dosage
compensation is less effective in birds than in mammals. *Journal of Biology*, *6*(1), 2.
https://doi.org/10.1186/jbiol53

Jegalian, K., and Page, D. C. (1998). A proposed path by which genes common to mammalian X
and Y chromosomes evolve to become X inactivated. *Nature*, *394*(August), 776–780.
Retrieved from http://www.nature.com/nature/journal/v394/n6695/abs/394776a0.html

Julien, P., Brawand, D., Soumillon, M., Necsulea, A., Liechti, A., Schütz, F., Daish, T., Grützner,
F., and Kaessmann, H. (2012, January). Mechanisms and evolutionary patterns of
mammalian and avian dosage compensation. *PLoS Biology*.
https://doi.org/10.1371/journal.pbio.1001328

Kaiser, V. B., Zhou, Q., and Bachtrog, D. (2011). Nonrandom gene loss from the drosophila
miranda neo-Y chromosome. *Genome Biology and Evolution*, *3*, 1329–1337.
https://doi.org/10.1093/gbe/evr103

Kharchenko, P. V, Xi, R., and Park, P. J. (2011). Evidence for dosage compensation between the
X chromosome and autosomes in mammals. *Nature Genetics*, *43*(12), 1167–1169.
https://doi.org/10.1038/ng.991

Lahn, B. T., Ma, N., Breg, R. W., Stratton, R., Surti, U., and Page, D. C. (1994). Xq-Yq
interchange resulting in supernormal X-linked gene expression in severely retarded males
with 46,XYq- karyotype. *Nature Genetics*, *8*, 362–369. https://doi.org/10.1038/ng1294-340

Lahn, B. T., and Page, D. C. (1999). Four evolutionary strata on the human X chromosome.
Science, *286*(5441), 964–967. https://doi.org/10.1126/science.286.5441.964

Lin, F., Xing, K., Zhang, J., and He, X. (2012). Expression reduction in mammalian X
chromosome evolution refutes Ohno's hypothesis of dosage compensation. *Proceedings of
the National Academy of Sciences*, *109*(29), 11752–11757.
https://doi.org/10.1073/pnas.1201816109

Lindgren, A. M., Hoyos, T., Talkowski, M. E., Hanscom, C., Blumenthal, I., Chiang, C., Ernst,
C., Pereira, S., Ordulu, Z., Clericuzio, C., Drautz, J. M., Rosenfeld, J. a, Shaffer, L. G.,
Velsher, L., Pynn, T., Vermeesch, J., Harris, D. J., … Morton, C. C. (2013).
Haploinsufficiency of KDM6A is associated with severe psychomotor retardation, global
growth restriction, seizures and cleft palate. *Human Genetics*, *132*(5), 537–52.
https://doi.org/10.1007/s00439-013-1263-x

Mank, J. E., and Ellegren, H. (2009). All dosage compensation is local: gene-by-gene regulation
of sex-biased expression on the chicken Z chromosome. *Heredity*, *102*(3), 312–320.
https://doi.org/10.1038/hdy.2008.116

Marin, R., Cortez, D., Lamanna, F., Pradeepa, M. M., Leushkin, E., Julien, P., Liechti, A.,
Halbert, J., Brüning, T., Mössinger, K., Trefzer, T., Conrad, C., Kerver, H. N., Wade, J.,
Tschopp, P., and Kaessmann, H. (2017). Convergent origination of a Drosophila -like
dosage compensation mechanism in a reptile lineage. *Genome Research*, 1–14.
https://doi.org/10.1101/gr.223727.117

Mueller, J. L., Skaletsky, H., Brown, L. G., Zaghlul, S., Rock, S., Graves, T., Auger, K., Warren,
W. C., Wilson, R. K., and Page, D. C. (2013). Independent specialization of the human and

mouse X chromosomes for the male germ line. *Nature Genetics*, *45*(9), 1083–1087. https://doi.org/10.1038/ng.2705

Nanda, I., Shan, Z., Schartl, M., Burt, D. W., Koehler, M., Nothwang, H., Grützner, F., Paton, I. R., Windsor, D., Dunn, I., Engel, W., Staeheli, P., Mizuno, S., Haaf, T., and Schmid, M. (1999). 300 million years of conserved synteny between chicken Z and human chromosome 9. *Nature Genetics*, *21*(march), 258–259. https://doi.org/10.1038/6769

Ohno, S. (1967). *Sex chromosomes and sex-linked genes*. Springer-Verlag.

Papp, B., Pal, C., and Hurst, L. D. (2003). Dosage sensitivity and the evolution of gene families in yeast. *Nature*, *424*, 194–197. https://doi.org/10.1038/nature01713.1.

Pessia, E., Makino, T., Bailly-Bechet, M., McLysaght, A., and Marais, G. a B. (2012). Mammalian X chromosome inactivation evolved as a dosage-compensation mechanism for dosage-sensitive genes on the X chromosome. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(14), 5346–51. https://doi.org/10.1073/pnas.1116763109

Pinzón, N., Li, B., Martinez, L., Sergeeva, A., Presumey, J., Apparailly, F., and Seitz, H. (2016). The number of biologically relevant microRNA targets has been largely over-estimated The number of biologically relevant microRNA targets has been largely overestimated, (November), 1–11. https://doi.org/10.1101/gr.205146.116

Ross, M. T., Grafham, D. V, Coffey, A. J., Scherer, S., McLay, K., Muzny, D., and Platzer, M. (2005). The DNA sequence of the human X chromosome. *Nature*, *434*(March), 325–337.

Ruderfer, D. M., Hamamsy, T., Lek, M., Karczewski, K. J., Kavanagh, D., Samocha, K. E., Exome Aggregation Consortium, Daly, M. J., MacArthur, D. G., Fromer, M., and Purcell, S. M. (2016). Patterns of genic intolerance of rare copy number variation in 59,898 human exomes. *Nature Genetics*, *48*(10), 1107–1111. https://doi.org/10.1038/ng.3638

Simon, D., Laloo, B., Barillot, M., Barnetche, T., Blanchard, C., Rooryck, C., Marche, M., Burgelin, I., Coupry, I., Chassaing, N., Gilbert-Dussardier, B., Lacombe, D., Grosset, C., and Arveiler, B. (2010). A mutation in the 3′-UTR of the HDAC6 gene abolishing the post-transcriptional regulation mediated by hsa-miR-433 is linked to a new form of dominant X-linked chondrodysplasia. *Human Molecular Genetics*, *19*(10), 2015–2027. https://doi.org/10.1093/hmg/ddq083

Skaletsky, H., Kuroda-kawaguchi, T., Minx, P. J., Cordum, H. S., Hillier, L., Brown, L. G., Repping, S., Pyntikova, T., Ali, J., Bieri, T., Chinwalla, A., Delehaunty, A., Delehaunty, K., Du, H., Fewell, G., Fulton, L., Fulton, R., … Page, D. C. (2003). The male-specific region of the human Y chromosome is a mosaic of discrete sequence classes. *Nature*, *423*, 825–838.

Tukiainen, T., Villani, A.-C., Yen, A., Rivas, M. A., Marshall, J. L., Satija, R., Aguirre, M., Gauthier, L., Fleharty, M., Kirby, A., Cummings, B. B., Castel, S. E., Karczewski, K. J., Aguet, F., Byrnes, A., Aguet, F., Ardlie, K. G., … MacArthur, D. G. (2017). Landscape of X chromosome inactivation across human tissues. *Nature*, *550*(7675), 244–248. https://doi.org/10.1038/nature24265

Uebbing, S., Konzer, A., Xu, L., Backström, N., Brunström, B., Bergquist, J., and Ellegren, H. (2015). Quantitative mass spectrometry reveals partial translational regulation for dosage compensation in chicken. *Molecular Biology and Evolution*, *32*(10), 2716–2725. https://doi.org/10.1093/molbev/msv147

Vandewalle, J., Van Esch, H., Govaerts, K., Verbeeck, J., Zweier, C., Madrigal, I., Mila, M., Pijkels, E., Fernandez, I., Kohlhase, J., Spaich, C., Rauch, A., Fryns, J. P., Marynen, P., and

Froyen, G. (2009). Dosage-dependent severity of the phenotype in patients with mental retardation due to a recurrent copy-number gain at Xq28 mediated by an unusual recombination. *American Journal of Human Genetics*, *85*(6), 809–822. https://doi.org/10.1016/j.ajhg.2009.10.019

Warnefors, M., Mossinger, K., Halbert, J., Studer, T., VandeBerg, J. L., Lindgren, I., Fallahshahroudi, A., Jensen, P., and Kaessmann, H. (2017). Sex-biased microRNA expression in mammals and birds reveals underlying regulatory mechanisms and a role in dosage compensation. *Genome Research*, 1–13. https://doi.org/10.1101/gr.225391.117

Watson, J. M., Spencer, J. A., Riggs, A. D., and Graves, J. A. (1990). The X chromosome of monotremes shares a highly conserved region with the eutherian and marsupial X chromosomes despite the absence of X chromosome inactivation. *Proceedings of the National Academy of Sciences*, *87*(18), 7125–7129. https://doi.org/10.1073/pnas.87.18.7125

Weischenfeldt, J., Dubash, T., Drainas, A. P., Mardin, B. R., Chen, Y., Stütz, A. M., Waszak, S. M., Bosco, G., Halvorsen, A. R., Raeder, B., Efthymiopoulos, T., Erkek, S., Siegl, C., Brenner, H., Brustugun, O. T., Dieter, S. M., Northcott, P. A., … Korbel, J. O. (2017). Pan-cancer analysis of somatic copy-number alterations implicates IRS4 and IGF2 in enhancer hijacking. *Nature Genetics*, *49*, 65–74. https://doi.org/10.1038/ng.3722

White, M. a, Kitano, J., and Peichel, C. L. (2015). Purifying Selection Maintains Dosage-Sensitive Genes during Degeneration of the Threespine Stickleback Y Chromosome. *Molecular Biology and Evolution*, *32*(8), 1981–1995. https://doi.org/10.1093/molbev/msv078

Xiong, Y., Chen, X., Chen, Z., Wang, X., Shi, S., Wang, X., Zhang, J., and He, X. (2010). RNA sequencing shows no dosage compensation of the active X-chromosome. *Nature Genetics*, *42*(12), 1043–1047. https://doi.org/10.1038/ng.711

Yang, F., Babak, T., Shendure, J., and Disteche, C. M. (2010). Global survey of escape from X inactivation by RNA-sequencing in mouse. *Genome Research*, *20*(5), 614–622. https://doi.org/10.1101/gr.103200.109

Yates, A., Akanni, W., Amode, M. R., Barrell, D., Billis, K., Carvalho-Silva, D., Cummins, C., Clapham, P., Fitzgerald, S., Gil, L., Girón, C. G., Gordon, L., Hourlier, T., Hunt, S. E., Janacek, S. H., Johnson, N., Juettemann, T., … Flicek, P. (2016). Ensembl 2016. *Nucleic Acids Research*, *44*(D1), D710–D716. https://doi.org/10.1093/nar/gkv1157

Zhou, Q., Zhang, J., Bachtrog, D., An, N., Huang, Q., Jarvis, E. D., Gilbert, M. T. P., and Zhang, G. (2014). Complex evolutionary trajectories of sex chromosomes across bird taxa. *Science*, *346*(6215), 1246338. https://doi.org/10.1126/science.1246338

Zimmer, F., Harrison, P. W., Dessimoz, C., and Mank, J. E. (2016). Compensation of Dosage-Sensitive Genes on the Chicken Z Chromosome. *Genome Biology and Evolution*, *8*(4), 1233–1242. https://doi.org/10.1093/gbe/evw075

Figure 1: Conserved miRNA targeting of autosomal genes stratified by copy number

variation in 59,898 human exomes. Probabilities of conserved targeting (P_{CT}) of all gene-

miRNA interactions involving non-duplicated and duplicated genes, further stratified as (A)

deleted (grey, n = 69,339 interactions from 4,118 genes; blue, n = 80,290 interactions from 3,976

genes) or (B) not deleted (orange, n = 51,514 interactions from 2,916 genes; purple, n = 72,826

interactions from 3,510 genes). *** $p < 0.001$, two-sided Kolmogorov-Smirnov test. (C) Mean

gene-level P_{CT} scores. ** $p < 0.01$, *** $p < 0.001$, two-sided Wilcoxon rank-sum test.

Figure 2. X-Y pairs and X-inactivated genes have higher miRNA conservation scores than

X escape genes. P_{CT} score distributions of all gene-miRNA interactions involving (A) human X-

Y pairs (n = 371 interactions from 15 genes), X-inactivated genes (n = 6,743 interactions from

329 genes), and X escape genes (n = 1,037 interactions from 56 genes). ** $p < 0.01$, two-sided

Kolmogorov-Smirnov test. (B) Mean gene-level P_{CT} scores. * $p < 0.05$, ** $p < 0.01$, two-sided

Wilcoxon rank-sum test.

Figure 3. Heterogeneities in X-linked miRNA targeting were present on the ancestral

autosomes. (A) Example reconstruction of an ancestral miR-96 target site in the 3` UTR of

KDM6A, an X-linked gene in the X-added region (XAR) with a surviving Y homolog. Dots in

non-human species indicate identity with the human sequence, dashes gaps indicate gaps in the

multiple sequence alignment. (B) Distributions of sites conserved between 3` UTRs of human

and chicken orthologs (top) or comparisons to background expectation (bottom, see Methods) for

human X-Y pairs (n = 16), X-inactivated genes (n = 251), and X escape genes (n = 42). (C)

Statistics as in (B), but using sites conserved between chicken and opossum 3` UTRs only for

genes in the XAR; X-Y pairs (n = 11), X-inactivated genes (n = 58), and X escape genes (n = 27).

Figure 4. Z-W pairs have higher miRNA conservation scores than other ancestral Z-linked

genes. P_{CT} score distributions of all gene-miRNA interactions involving the human orthologs of

(A) chicken Z-W pairs (n = 832 interactions from 28 genes) and other ancestral Z genes (n =

16,692 interactions from 657 genes). *** $p < 0.001$, two-sided Kolmogorov-Smirnov test. (B)

Mean gene-level P_{CT} scores. *** $p < 0.001$, two-sided Wilcoxon rank-sum test.

Figure 5. Heterogeneities in Z-linked miRNA targeting were present on the ancestral

autosomes. (A) Example reconstruction of an ancestral miR-145 target site in the 3` UTR of

RASA1, a Z-linked gene with a surviving W homolog. Example of 3` UTR sequence alignment

for RASA1, a Z-linked gene with a surviving W homolog, with a target site for miR-145

highlighted in gray. (B) Numbers of sites conserved between 3` UTRs of human and chicken

orthologs (top) or comparisons to background expectation (bottom) for chicken Z-W pairs (n =

27) and other ancestral Z genes (n = 578). (C) Statistics as in (B), but using sites conserved

between human and anolis 3` UTRs.

Figure 6. Analyses of experimental datasets validate miRNA target site function. Responses

to transfection (A,B,C) or knockout (D) of indicated miRNAs in human (A,B,C) or mouse (D)

cell-types. Each panel depicts corresponding changes in mRNA levels (A,B), in fraction of

Argonaute-bound genes (C), and in mRNA stability and translational efficiency as measured by

ribosome protected fragments (RPF, D). In each case, X-linked genes and the human orthologs

of Z-linked genes containing target sites with an assigned P_{CT} score (red) for the indicated

miRNA were compared to all expressed genes lacking target sites (black); gene numbers are

indicated in parentheses. (A,B,D) *** p < 0.001, two-sided Kolmogorov-Smirnov test. (C) * p <

0.05, two-sided Fisher's exact test.

Figure 7. An evidence-based model of preexisting heterogeneities in dosage sensitivity

shaping mammalian and avian sex chromosome evolution. In this model, preexisting

heterogeneities in dosage sensitivity determined the trajectory of Y/W gene loss in both

mammals and birds, and of subsequent X-inactivation in mammals and dosage compensation in

birds. Colored arrow widths are scaled approximately to the number of ancestral genes in each

class. (A) The dashed orange line represents the possibility that a subset of X-linked genes may

have not undergone compensatory X upregulation following Y gene decay. (B) Ancestral Z

genes with no W homolog follow a gradient of preexisting dosage sensitivity (top, grey to white),

which determined the degree of dosage compensation following W gene loss (bottom).

Supplemental Figure S1: Effect of deletion status on autosomal P_{CT} scores. Probabilities of

conserved targeting (P_{CT}) of all gene-miRNA interactions involving non-deleted and deleted

genes, further stratified as (A) duplicated (grey, n = 69,339 interactions from 4,118 genes; orange,

n = 51,514 interactions from 2,916 genes) or (B) not duplicated (purple, n = 72,826 interactions

from 3,510 genes; blue, n = 80,290 interactions from 3,976 genes). *** p < 0.001, two-sided

Kolmogorov-Smirnov test. (C) P_{CT} scores for all gene sets in (A) and (B) superimposed on one

plot. (D) Mean gene-level P_{CT} scores when aggregating sets of duplicated/not duplicated (left) or

deleted/not deleted (right) genes. *** p < 0.0001, two-sided Wilcoxon rank-sum test.

Supplemental Figure S2: P_{CT} scores of X-Y pairs across 8 mammals. (A) P_{CT} score distributions of all gene-miRNA interactions involving X-Y pairs across eight sequenced mammalian Y chromosomes (n = 647 interactions from 32 genes) and other ancestral X genes (n = 8,831 interactions from 457 genes). ** p < 0.01, two-sided Kolmogorov-Smirnov test. (B) Gene-level mean P_{CT} scores. * p < 0.05, two-sided Wilcoxon rank-sum test.

Supplemental Figure S3: Resampled mean P_{CT} scores of X-linked genes. (A) Resampled gene-miRNA P_{CT} scores for human X-Y pairs (n = 15 genes), X-inactivated genes (n = 329 genes) and X escape genes (n = 56 genes). (B) Resampled gene-miRNA P_{CT} scores for X-Y pairs across eight mammals (n = 32 genes) and genes with no Y homolog in any of eight mammals (n = 457 genes). Points and error bars represent the median and 95% confidence intervals from 1,000 gene samplings with replacement. * p < 0.05, ** p < 0.01, empirical p-value computed as the fraction of random non-overlapping gene sets with a median difference in P_{CT} score at least as large as the true difference.

Supplemental Figure S4: P_{CT} score comparisons with consistent and variable escape genes separated. (A) P_{CT} score distributions of all gene-miRNA interactions involving X-Y pairs (n = 371 interactions from 16 genes), X-inactivated genes (n = 6743 interactions from 329 genes), consistent escape genes (n = 567 interactions from 30 genes), or variable escape genes (n = 470 interactions from 26 genes) as defined by Balaton et al (Balaton et al., 2015). * p < 0.05, ** p < 0.01, two-sided Kolmogorov-Smirnov test. (B) Resampled gene-miRNA P_{CT} scores of gene classes from (A). Points and error bars represent the median and 95% confidence intervals from

1,000 gene samplings with replacement. * $p < 0.05$, empirical p-value computed as the fraction

of random non-overlapping gene sets with a median difference in P_{CT} score at least as large as

the true difference.

Supplemental Figure S5: P_{CT} score comparisons with discordant genes included as X-

inactivated or escape. P_{CT} score distributions of all gene-miRNA interactions (A,C) or mean

gene-level P_{CT} score (B,D) of classes of X-linked genes with genes with a discordant XCI call (n

= 721 interactions from 40 genes) included as X-inactivated (A,B) or X escape (C,D). Numbers

of gene-miRNA interactions and genes as in Figure 1, but with the addition of discordant gene

numbers/interactions to X-inactivated genes (A,B) or X escape genes (C,D). * $p < 0.05$, ** $p <$

0.01, two-sided Kolmogorov-Smirnov (A,C) or Wilcoxon rank-sum (B,D) test.

Supplemental Figure S6: Variation in within-UTR conservation does not account for

observed differences in P_{CT} score among classes of X-linked genes. (A) Example of step-

detection to segment 3` UTRs. Top, base-wise branch length scores; bottom, probabilities of

transition to a new section. Dashed line indicates p-value cutoff used to delineate a new section

(plotted as alternating magenta/yellow points). (B) Boxplots of within-UTR conservation bias

(see Methods) for all gene-miRNA interactions involving classes of X-linked genes. (C)

Comparisons of P_{CT} scores normalized by within-UTR bias. **, $p < 0.01$, *** $p < 0.001$, two-

sided Kolmogorov-Smirnov test.

Supplemental Figure S7: Ancestral miRNA targeting of X-Y pairs across 8 mammals. (A)

Distributions of sites conserved between 3` UTRs of human and chicken orthologs (top) or

comparisons to background expectation (bottom, see Methods) for X-Y pairs across 8 mammals

(n = 25) and other ancestral X genes (n = 351). (D) Statistics as in (C), but using sites conserved

between chicken and opossum 3` UTRs only for genes in the XAR; X-Y pairs across 8 mammals

(n = 15), other ancestral X genes (n = 102).

Supplemental Figure S8: P_{CT} scores of Z-W pairs across 4 and 14 birds. (A,C) P_{CT} score

distributions of all gene-miRNA interactions (A) Z-W pairs including predictions from three

additional birds with male and female genome sequence (n = 2,187 interactions from 78 genes)

and other ancestral Z genes (n = 15,357 interactions from 607 genes), or (C) Z-W pairs including

read depth-based predictions from 10 additional birds with only female genome sequence (n =

4,458 interactions from 157 genes) and other ancestral Z genes (n = 13,086 interactions from 528

genes) *** $p < 0.001$, two-sided Kolmogorov-Smirnov test. (B,D) Gene-level mean P_{CT} scores.

*** $p < 0.01$, two-sided Wilcoxon rank-sum test.

Supplemental Figure S9: Resampled mean P_{CT} scores of Z-linked genes. Gene sets: (A)

chicken Z-W pairs (n = 28 genes) and other ancestral Z genes (n = 657 genes), (B) Z-W pairs

across four birds (n = 78 genes) compared to the remainder of ancestral Z genes (n = 607 genes),

and (C) Z-W pairs across 14 birds (n = 157 genes) compared to the remainder of ancestral Z

genes (n = 528 genes). Points and error bars represent the median and 95% confidence intervals

from 1,000 gene samplings with replacement. *** $p < 0.001$, empirical p-value computed as the

fraction of random non-overlapping gene sets with a median difference in P_{CT} score at least as

large as the true difference.

Supplemental Figure S10: Variation in within-UTR conservation cannot account for

observed differences in P_{CT} score among classes of Z-linked genes. (A) Boxplots of within-

UTR conservation bias (see Methods) for all gene-miRNA interactions involving chicken Z-W

pairs or other ancestral X genes. Numbers of interactions and genes as in Figure 4A. ** $p < 0.01$,

two-side Wilcoxon rank-sum test. (B) Comparisons of P_{CT} scores normalized by within-UTR

bias. *** $p < 0.001$, two-sided Kolmogorov-Smirnov test.

Supplemental Figure S11: Correlation of Z-linked gene-specific dosage compensation with

gene-level P_{CT} score. Distributions of chicken male/female expression ratio, normalized to that

of human and anolis (y-axis) as a function of mean gene-level P_{CT} quartile (x-axis) for all

expressed Z-linked gene with no W homolog. Expression ratios are plotted on a $\log_2$ scale;

values closer to 0 imply more effective dosage compensation following W gene loss.

Supplemental Figure S12: Ancestral miRNA targeting of Z-W pairs across 4 birds. (A)

Distributions of sites conserved between 3` UTRs of human and chicken orthologs (top) or

comparisons to background expectation (bottom, see Methods) for Z-W pairs across chicken and

three additional birds with male and female genome sequence (4 birds, n = 73) and other

ancestral Z genes (n = 532). (D) Statistics as in (C), but using sites conserved between human

and anolis 3` UTRs; Z-W pairs across 4 birds (n = 73), other ancestral Z genes (n = 527). *** $p <$

0.001, two-sided Fisher's exact test.

Supplemental Figure S13: Ancestral miRNA targeting of predicted Z-W pairs across 14

birds. (A) Distributions of sites conserved between 3` UTRs of human and chicken orthologs

(top) or comparisons to background expectation (bottom, see Methods) for Z-W pairs in chicken,

predicted in three additional birds with male and female genome sequence, and predicted based

on read depth from 10 additional birds with only female genome sequence (14 birds, n = 147)

and other ancestral Z genes (n = 458). (D) Statistics as in (C), but using sites conserved between

human and anolis 3` UTRs; Z-W pairs across 14 birds (n = 147), other ancestral Z genes (n =

453).

Supplemental Figure S14: Correlation of Z-linked gene-specific dosage compensation with

human-chicken-conserved site excess. Distributions of chicken male/female expression ratios,

normalized to those of human and anolis (y-axis) for expressed Z-linked genes with no W

homolog with (left) or without (right) an excess of human-chicken-conserved miRNA sites. ** p

< 0.01, *** $p < 0.0001$, Wilcoxon rank-sum test.

Supplemental Figure S15: Gene expression changes following small RNA transfections in

human HeLa cells. * $p < 0.05$, *** $p < 0.001$, two-sided K-S test.

Supplemental Figure S16: Gene expression changes following transfection or knockdown

of additional miRNAs in human HCT116 or HEK293 cells. *** $p < 0.001$, two-sided

Kolmogorov-Smirnov test.

Supplemental Figure S17: Changes in mRNA stability and translational efficiency and gene

expression following miR-155 knockout in mouse immune cells. In each case, mouse

orthologs of X- or Z-linked genes containing a human-mouse-conserved (hsa-mmu) miR-155

site were compared to mouse genes containing only nonconserved miR-155 sites. * $p < 0.05$, ***

$p < 0.001$, two-sided Kolmogorov-Smirnov test.

Supplemental Figure S18: Argonaute binding measured by high-throughput crosslinking-

immunoprecipitation (CLIP) following miRNA transfection in HEK293 cells. * $p < 0.05$,

two-sided Fisher's exact test.

A Human-chicken conserved sites

p = 1.10e-10,
Fisher's exact test

Fraction of genes

sites: 0, 1–10, >10

B Human-anolis conserved sites

p = 9.068e-09,
Fisher's exact test

Fraction of genes

sites: 0, 1–10, >10

Fraction with # true sites > bckgd.

***

Ancestral Z

Z-W pairs
(4 birds)

Fraction with # true sites > bckgd.

***

A Human-chicken conserved sites

p = 1.013e-12,
Fisher's exact test

B Human-anolis conserved sites

p = 8.557e-09,
Fisher's exact test

Ancestral Z

Z-W pairs
(14 birds)

A

post–miR–124 transfection

B

post–miR–7 transfection