1 **Estimation of genomic prediction accuracy from reference populations with**
2 **varying degrees of relationship**

4 S. Hong Lee[*], Sam Clark, and Julius H.J. van der Werf[*]

5 School of Environmental and Rural Science, University of New England, NSW 2351,

6 Australia

*Correspondence:

S. Hong Lee, School of Environmental and Rural Science, University of New

England, NSW 2351, Australia

Tel: +61 2 6773 3665

Email: hong.lee@une.edu.au

Or

Julius H.J. van der Werf, School of Environmental and Rural Science, University of

New England, NSW 2351, Australia

Tel: +61 2 6773 2092

Email: Julius.vanderwerf@une.edu.au

ABSTRACT

Genomic prediction is emerging in a wide range of fields including animal and plant

breeding, risk prediction in human precision medicine and forensic. It is desirable to

establish a theoretical framework for genomic prediction accuracy when the reference

data consists of information sources with varying degrees of relationship to the target

individuals. A reference set can contain both close and distant relatives as well as

'unrelated' individuals from the wider population in the genomic prediction. The

various sources of information were modeled as different populations with different

effective population sizes (N_e). Both the effective number of chromosome segments

(M_e) and N_e are considered to be a function of the data used for prediction. We

validate our theory with analyses of simulated as well as real data, and illustrate that

the variation in genomic relationships with the target is a predictor of the information

content of the reference set. With a similar amount of data available for each source,

we show that close relatives can have a substantially larger effect on genomic

prediction accuracy than lesser related individuals. We also illustrate that when

prediction relies on closer relatives, there is less improvement in prediction accuracy

with an increase in training data or marker panel density. We release software that can

estimate the expected prediction accuracy and power when combining different

reference sources with various degrees of relationship to the target, which is useful

when planning genomic prediction (before or after collecting data) in animal, plant

and human genetics.

INTRODUCTION

Genomic prediction of (additive) genetic effects and phenotypes is emerging in a wide range of fields including animal and plant breeding, risk prediction in human medicine and forensics[1-4]. Genomic prediction requires modeling of the association between genome-wide single nucleotide polymorphisms (SNPs) and phenotypes. The success of genomic prediction is measured by its accuracy, i.e. how reliable a future phenotype of target individuals can be predicted.

Genomic prediction requires a reference population of individuals having information on both genotype and phenotype. The accuracy of genomic prediction depends on various parameters, including sample size of the reference and its genetic structure. An important parameter in relation to the latter is the effective size of the population. The effective population size is a predictor of the effective number of chromosome segments that are represented in the population[5-7]. Theoretical predictions have usually considered a homogeneous population of individuals that are essentially unrelated. However, in most practical applications, the reference population used for genomic predictions possibly consists of many sub-groups with individuals having a variety of relatedness to the target individual, e.g. direct relatives, more distant relatives, and individuals that are considered unrelated. It is relevant to assess the contribution of these various sources to prediction accuracy before actually conducting an experiment.

A number of studies have shown that genomic predictions are more accurate if the genomic relationship between the proband and the reference population is higher, both in humans[8-11] and in other species[12-14]. Habier et al (2013)[15]

distinguished between three types of information in genomic prediction; linkage

disequilibrium, additive-genetic relationships and co-segregation of QTL predicted

from markers genotypes with a pedigree. They argued that it would be useful to

understand how these sources contribute to the accuracy of genomic predictions,

especially when designing reference populations for breeding programs. They show

these contributions via simulated examples but did not provide methods that allow

simple predictions for their contribution to accuracy. Pszczola et al. (2012)[16]

showed that the relationship between the reference population and the proband should

be maximized to achieve an optimal design using a simulation study. However, they

also did not attempt to derive the expected prediction accuracy from an optimal

design in advance. Hayes et al. (2009)[17] considered the influence of direct relatives

on genomic prediction. They followed the same approach as the general theory, i.e. by

considering the number of independently segregating chromosome segments within

families. They showed the accuracy of genomic prediction from varying sizes of the

first and second degree of relatives, but did not consider the information from

combined sources[18]. It should also be noted that those studies that derived genomic

prediction accuracy from theory using effective number of chromosome segments

(M_e)[5, 6, 19-21], did not consider the correlation between relatedness at different

chromosomes, therefore overestimating M_e and underestimating whole-genome

prediction accuracy[7].

Wientjes et al (2016)[22] proposed a simple selection index approach to combine

information from different populations. They considered a genetic correlation

between genetic effects expressed in different populations. We propose to use the

same approach to combine different sources of information from different subsets

within a population, where the different subsets have a different degree of relationship

with the target individual. To predict the accuracy, we derive the number of effective

chromosome segments from a hypothetical N_e associated with each subset, and we

show that that combining such subsets using selection index theory gives the same

result as using a prediction from an M_e derived from the variation in genomic

relationships between the overall reference data and the target. Prediction accuracy is

derived from variation in genomic relationship rather than the variation in genomic

relationship as a deviation for the expected relationship among members of the

reference set, as was proposed by Goddard et al (2011) and also applied by Wientjes

et al (2016). This approach leads to a theoretical concept useful for assessing the

accuracy of genomic predictions in advance, and we illustrate this with examples

based on real data.

MATERIALS AND METHODS

Predicting genomic selection accuracy

The accuracy of genomic breeding values (GBV) or (genomic profile score in the context of human risk prediction[23]) based on genome-wide SNP genotypes can be predicted from theory[5-7, 24], assuming that prediction is based on a reference population with phenotypes and genotypes for the same genome-wide SNPs that are linked to quantitative trait loci (QTL). The accuracy depends on i) the proportion of genetic variance at QTL captured by markers and ii) the accuracy of estimating marker effects. The proportion of genetic variance at QTL captured by markers (b) depends on linkage disequilibrium (LD) between markers and QTL, which in turn depends on the number of markers (M) and the number of 'effective chromosome segments' (M_e)[5], that is

$b = M / (M_e + M)$[5].

Various forms of prediction of M_e have been presented[5, 6, 21] that were however inconsistent to each other, and without considering the correlation between chromosomes. Recently, we presented a prediction formula with the form[7]

$$M_e = \frac{N_{chr}}{[\ln(2N_eL+1)+2N_eL(\ln(2N_eL+1)-1)]/(4N_e^2L^2)+(1/3N_e)\cdot(N_{chr}-1)}$$

(1)

where N_e = effective population size; L = average chromosome length; N_{chr} = number of chromosomes. This formula accounts for mutation, and that without considering mutation should be referred to equation (10) in Lee et al. (2017)[7]. If $N_{chr} = 30$ and $L = 1$, Eq. (1) can be approximately simplified as

$M_e = 2.938\, N_e^{0.965}$, and $N_e = 0.327\, M_e^{1.036}$.

(2)

The accuracy of the genomic prediction of a phenotype can be written as[5]

$$r_{y,\hat{g}} = h \cdot r_{g,\hat{g}} = h \cdot \sqrt{\frac{bh^2}{bh^2 + M_e/N}} = \sqrt{\frac{bh^4}{bh^2 + M_e/N}}$$

(3)

where $r_{y,\hat{g}}$ is the correlation coefficient between the true phenotypes (y) and estimated

GBV, h^2 is the heritability of the trait, and N is the number of phenotypic

observations. Other measures for genomic prediction accuracy, particularly for human

risk prediction, such as the area under the receiver operating characteristic curve

(AUC) or odds ratio of case-control status contrasting the higher or lower risk group

are described elsewhere[7].

M_e and genomic relationship

After collecting genotypic information of the reference data and the target individual,

it is possible to obtain an empirical M_e from a genomic relationship matrix (GRM).

The elements in the GRM are $G_{Tj} = \sum_{m=1}^{M} x_{Tm} x_{jm} / M$ where x_{Tm} and x_{jm} are the

standardised genotype coefficients (mean 0 and variance 1) for the target individual

(T) and jth individual in the reference data at the mth locus. It is possible to construct

a GRM for each locus, and the elements in the GRM at the mth locus are

$G_{Tj(m)} = x_{Tm} x_{jm}$. Then, the variance of the mean of $G_{Tj(m)}$ across all M_i SNPs in a single

chromosome is

149
$$
\begin{aligned}
\operatorname{var}\left(\sum_{m=1}^{M_i} G_{Tj(m)} / M_i\right) &= \frac{1}{M_i^2}\left[\sum_{m=1}^{M_i}\sum_{l=1}^{M_i} \operatorname{cov}(G_{Tj(l)}, G_{Tj(m)})\right] \\
&= \frac{1}{M_i^2}\left[\sum_{m=1}^{M_i}\sum_{l=1}^{M_i} \operatorname{cov}(x_{T(l)}x_{j(l)}, x_{T(m)}x_{j(m)})\right] \\
&= \frac{1}{M_i^2}\left[\sum_{m=1}^{M_i}\sum_{l=1}^{M_i} \operatorname{cov}(x_{j(l)}, x_{j(m)})\operatorname{cov}(x_{T(l)}, x_{T(m)})\right] \\
&= \frac{1}{M_i^2}\left[\sum_{m=1}^{M_i}\sum_{l=1}^{M_i} r_{m,l}^2\right] \\
&= \frac{1}{M_{e(i)}}
\end{aligned}
\tag{4}
$$

where $\text{cov}(x_{j(l)}, x_{j(m)}) = \text{cov}(x_{T(l)}, x_{T(m)}) = r_{m,l}$, which is a correlation between the m^{th}

and l^{th} SNP-genotype, because of $\text{var}(x) = 1$ and $\text{mean}(x) = 0$, i.e. the genotype

coefficients are standardized in the population. The term $M_{e(i)}$ is the effective number

of chromosome segments for the ith chromosome, and it is calculated from the

inverse of the average of the squared correlation between the M_i SNPs[6, 7]. When

considering multiple chromosomes, the covariance of the pairwise relationship

between two chromosomes is not negligible[7]. Assuming equal length and number of

SNPs for N_{chr} chromosomes, M_e for the whole genome can be derived from the

variance in the relationship across the whole genome as

$$\frac{1}{M_e} = \left[\text{var}\left(\sum_{m=1}^{M_i} G_{Tj(m)} / M_i \right) + \left(\frac{N_{chr} - 1}{3N_e} \right) \right] \cdot \frac{1}{N_{chr}} = \text{var}\left(\sum_{m=1}^{M} G_{Tj(m)} / M \right) = \text{var}(G_{Tj}) \qquad (5)$$

where N_{chr} is the number of chromosomes. This expression is equivalent to (1) when

using Sved (1971)[25] for deriving the expected squared correlation between

genotypes, which is dependent on N_e. An empirical M_e can be derived from the GRM

as the variance in relationships between a target individual T and N individuals in the

reference population. Goddard et al. (2011)[5] suggested their theoretical derivation had

to assume a homogeneous population of individuals that are essentially unrelated.

However, Eq. (4) show that the assumption about unrelated individuals is not

necessary so that any random samples from the population can be used, irrespective

of whether they are highly related or not (see Results).

Effective population size in a reference data set

One of critical parameters to determine the accuracy of genomic prediction is the

effective population size (N_e). It is not very common to represent a reference

population by a single value of N_e when it consists of several cohorts of individuals

with different relationships to the target individual. Wientjes et al. (2016)[22] used a

single value for M_e representing a reference set consisting of two populations. Here,

we generalized that concept for any number of subsets of the reference population

based on the relationship between N_e, M_e and var(G_{T*}), leading to a certain value for

M_e and N_e for a reference population consisting of several cohorts. For any subset of

the reference data set, there are realized relationships with the target sample. From

Eq. (5), a value of M_e, which is the inverse of the variance of the genomic

relationships between the target and the reference sample, can be assigned to the

reference data. Then, a single value of N_e, which is a function of M_e from Eq. (1) or

(2), can be obtained for the reference data. The effective population size of the

reference set is therefore a parameter specific to the data used and it describes

genomic diversity of the reference individuals used for prediction of genomic

breeding value relative to the target individual which breeding value is being

predicted. This N_e value can be smaller than the effective size of the population from

which the reference individuals were sampled, but it can also be larger, depending on

whether the reference individuals chosen were closer or more distantly related to the

reference set.

Based on this concept of N_e, reflecting information content of the reference sample in

relation to the target sample, we consider three information sources consisting of i)

close relatives of the proband, e.g. $N_e = 10$, ii) distant relatives or individuals from the

local area of the proband, e.g. $N_e = 100$ and iii) a wider population sample of

individuals that are not related to the proband, e.g. $N_e = 1,000$.

The GBV can be estimated based on each of these information sources, and the

accuracy of the estimation can be calculated as above, e.g. $r_{g,\hat{g}(i)}$ from Eq. (3) where i

represents the ith information source. It is also possible to estimate GBV based on

combined data of all three information sources. Assuming a random sample from the

same population for each source, the accuracy of the GBV based on the combined

data set can then be calculated using standard selection index theory as

$$
204 \quad r_{g,\hat{g}} = \sqrt{\mathbf{g}'\mathbf{P}^{-1}\mathbf{g}} = \sqrt{ \begin{bmatrix} r_{g,\hat{g}(1)}^2 \\ r_{g,\hat{g}(2)}^2 \\ r_{g,\hat{g}(3)}^2 \end{bmatrix}' \begin{bmatrix} r_{g,\hat{g}(1)}^2 & r_{g,\hat{g}(1)}^2 \cdot r_{g,\hat{g}(2)}^2 & r_{g,\hat{g}(1)}^2 \cdot r_{g,\hat{g}(3)}^2 \\ r_{g,\hat{g}(2)}^2 \cdot r_{g,\hat{g}(1)}^2 & r_{g,\hat{g}(2)}^2 & r_{g,\hat{g}(2)}^2 \cdot r_{g,\hat{g}(3)}^2 \\ r_{g,\hat{g}(3)}^2 \cdot r_{g,\hat{g}(1)}^2 & r_{g,\hat{g}(3)}^2 \cdot r_{g,\hat{g}(2)}^2 & r_{g,\hat{g}(3)}^2 \end{bmatrix}^{-1} \begin{bmatrix} r_{g,\hat{g}(1)}^2 \\ r_{g,\hat{g}(2)}^2 \\ r_{g,\hat{g}(3)}^2 \end{bmatrix} } \quad (6)
$$

205 where $\mathbf{g}$ is the vector with covariances between each of the GBV and the true

206 breeding value, and $\mathbf{P}$ is the variance-covariance matrix of the set of GBV. The

207 accuracy of the GBV based on the combined data set can also be estimated based on

208 the weighted M_e from the three information sources. Assuming a random sample from

209 the same population for each source, the weighted M_e can be obtained as

$$
210 \quad M_{e(weighted)} = \frac{1}{\sum_{k=1}^{N_{sub-sample}} \mathrm{var}(G_{Tj})_k\, p_k} = \frac{1}{\sum_{k=1}^{N_{sub-sample}} \frac{p_k}{M_{e(k)}}} \quad (7)
$$

where p_k is the proportion of the sample size over the total sample for each

information source. The accuracy of the GBV based on the weighted M_e is identical

with that using standard selection index theory above (Eq. (6)).

Following Wientjes et al. (2016)[22] we can further generalize for a case where

genetic correlations among multiple reference populations and those between

reference populations and the target are not one. Equation (6) can be generalized as

$$
217 \quad r_{g,\hat{g}} = \sqrt{\mathbf{g}'\mathbf{P}^{-1}\mathbf{g}} = \sqrt{ \begin{bmatrix} r_{G_{1,T}}^2 r_{g,\hat{g}(1)}^2 \\ \vdots \\ r_{G_{k,T}}^2 r_{g,\hat{g}(k)}^2 \end{bmatrix}' \begin{bmatrix} r_{G_{1,T}}^2 r_{g,\hat{g}(1)}^2 & \cdots & r_{g,\hat{g}(1)}^2 r_{g,\hat{g}(k)}^2 r_{G_{1,k}} r_{G_{1,T}} r_{G_{k,T}} \\ \vdots & \ddots & \vdots \\ r_{g,\hat{g}(k)}^2 r_{g,\hat{g}(1)}^2 r_{G_{k,1}} r_{G_{k,T}} r_{G_{1,T}} & \cdots & r_{G_{k,T}}^2 r_{g,\hat{g}(k)}^2 \end{bmatrix}^{-1} \begin{bmatrix} r_{G_{1,T}}^2 r_{g,\hat{g}(1)}^2 \\ \vdots \\ r_{G_{k,T}}^2 r_{g,\hat{g}(k)}^2 \end{bmatrix} } \quad (8)
$$

where $r_{G_{k,T}}$ is the genetic correlation between the k^{th} reference population and the

target set, and similarly, $r_{G_{i,j}}$ is the genetic correlation between the i^{th} and j^{th} reference

population ($i = j = 1 \sim k$).

Equation (8) can be used when multiple reference populations have both quantitative

traits and diseases. If the k^{th} reference population is measured for binary response

(disease or disorder), the reliability term, $r_{g,\hat{g}}^2$, in Eq. (8) can be replaced with

$$r_{u,\hat{u}}^2 = \frac{h^2 z^2}{h^2 z^2 + [K(1-K)]^2 \cdot M_e / [P(1-P) \cdot N]}$$

where u is a genetic profile score on the 0,1 disease scale[26, 27], K is the population

prevalence for the disease, P is the proportion of cases in the total sample N of cases

and controls, and z is the density at the threshold on the normal distribution in the

liability threshold model.

When the target data set is measured for a binary response (e.g. diseases), the AUC or

odds ratio of case-control status contrasting the higher or lower risk group can be also

estimated[7]. These measures for genomic prediction accuracy with multiple

heterogeneous reference populations can be obtained using MTG2, publicly available

software (https://sites.google.com/site/honglee0707/mtg2).

Power of genomic prediction

The power depends on the sample size in the target data set and the overall reliability

of the genomic prediction (Eq. (8)).

$NCP = N_T \cdot r_{y,\hat{g}}^2 / (1 - r_{y,\hat{g}}^2)$

where NCP is the non-centrality parameters, N_T is the sample size in the target data

and $r_{y,\hat{g}}^2$ is the coefficient of determination of the predictor. Then, the power can be

written as

$$power = 1 - \Phi\left[\Phi^{-1}\left(1 - \alpha/2\right) - \sqrt{NCP}\right] + \Phi\left[\Phi^{-1}\left(\alpha/2\right) - \sqrt{NCP}\right]$$

where Φ is the cumulative standard normal function and α is the significance level.

When using the odds ratio of case-control status contrasting the higher or lower risk

group, the non-centrality parameters can be derived as[28]

$$NCP = \ln(OR)^2\left[N_{top}P_{top}\left(1 - P_{top}\right)P_{bottom}\left(1 - P_{bottom}\right)\right]/\left[P_{bottom}\left(1 - P_{bottom}\right) + \left(N_{top}/N_{bottom}\right)P_{top}\left(1 - P_{top}\right)\right]$$

where N_{top} and N_{bottom} is the number of individuals in the top and bottom percentile,

and P_{top} and P_{bottom} is the proportion of cases in each group. The power can be

estimated as above.

Simulation

In a simulation, a stochastic gene-dropping method [29, 30] was used to simulate

4,000 SNPs for each of 30 chromosomes, each of length L=1 Morgan with $N_e = 50$,

500 and 1000 for 50, 500 and 1000 generations, respectively. Recombination and

mutations were modelled according to the genetic distance between SNPs and the

mutation rate of 1e-08 per site per generation[31]. In the final generation, we

constructed a genomic relationship matrix for a random set of 3000 individuals.

Among the 3000 individuals, we randomly selected 1000 individuals as target data

and 2000 individual as reference data, and estimated variance of the genomic

relationships between the target and reference data to validate Eq. (1), (2), (4) and (5).

Evaluation of the formulas

For each of the three information sources contributing to genomic prediction we

varied values for N_e, sample size in reference data and marker density. We compared

the expected accuracy of GBV from the sample of N_e =1000 with predictions that

additionally included information from the sample of N_e =100 and N_e =10. The total

number in the reference population was kept equal between the comparisons.

Real data analysis

We used publicly available data from the Framingham heart study

(phs000007.v26.p10.c1)[32]. There were 6950 individuals genotyped for 500,568

SNPs. Stringent quality control for genotype data and phenotype adjustment for

confounders were applied to the data (the details can be found in Lee et al. (2016)[7]).

The quality control included SNP call rate > 0.95, individual call rate > 0.95, HWE p-

value > 0.0001, MAF > 0.01 and individual population outliers < 6 SD from the first

and second principal components (PC). After QC, 6920 individuals and 389,265

SNPs remained. Among them, 4243 individuals were phenotyped for height and body

mass index (BMI).

We made three different information sources to form the reference data that were

tested in 100 replicated analyses (Table 1). Initially, we randomly selected 800

individuals out of 4243 phenotyped individuals as a target data set. For reference data

set #1, we selected 50% of individuals that were highly related (> relatedness of 0.3)

to the 800 target individuals (N_1 = 617 ± 19). For reference data set #2, we selected

80% of moderately related individuals (> relatedness of 0.1) of the 800 target

individuals (N_2 = 1254 ± 30). For reference data set #3, we took the rest of the

individuals that were not selected for reference data set #1 and #2 (N_3 = 1572 ± 33).

There was no overlap sample between target data set and reference data sets #1, #2

and #3.

Using the real genotype data, the genomic relationships between the reference and

target sample were constructed. Empirical M_e was estimated from equation (5) for

reference #1, 2 and 3, and that for combined data. We took a median rather than mean

because the distribution of variance of the genomic relationship between target and

reference sample was skewed. The correlation between the true phenotypes (that were

not used in the analyses) and estimated GBV in the target data set was estimated for

the combined data set, which was used as the genomic prediction accuracy ($r_{y,\hat{g}}$).

Phenotypes were adjusted for birth year, sex, and the first 10 PCs were used to control

non-genetic confounding effects, e.g. population stratification.

RESULTS

In the simulation study, as shown in Figure 1A, 1B and 1C, the expected (from Eq.

(1)) and empirically observed M_e from the simulated genotyped data (using Eq. (5))

are in good agreement, however, they are considerably lower than the expectation

from the previous formulas[5, 6, 21], which confirms the result from Lee et al.

(2017)[7]. It is noted that Eq. (5) is still valid in the subset with a smaller $N_e = 50$ that

has a significant proportion of high related individuals, indicating that the assumption

about unrelated individuals (made in Godard et al. (2011)[5]) can be relaxed. It is

shown that whether using a high or low effective population size, the mean and

variance of genomic relationships is generally agreed with the expectation from the

theory (Supplementary Figures S1 and S2).

In the evaluation of the formulas, we first tested how the prediction accuracy was

changed with varying marker density, using formula (1) and (3) and $b = M / (M_e + M)$

(Figure 2). For N_e=10,000, the accuracy gradually increased with marker density, but

the slope became flat when using the number of SNPs exceeded 100,000 (Figure 2A).

For N_e=1,000, the accuracy did not increase with marker density as long as the

number of SNPs was higher than 50,000 (Figure 2B). For N_e=100, there was no

improvement of the accuracy if the number of SNPs was more than 10,000 (Figure

2C). This would be expected because the proportion of genetic variance at QTL

captured by markers (b = $M / (M_e + M)$) approached one when the number of SNPs

(M) was more than 100,000, 50,000 and 10,000 for N_e = 10,000, 1000 and 100,

respectively (Figure 3), as M_e was equal to 21,248, 2,313 and 254 for these three

cases.

Next, we quantified the contribution of each information source when varying sample

size in the reference data using formula (1), (3) and (5) (Figure 4). It was assumed

that the number of SNPs was sufficient to capture most of causal variants (e.g. >

50,000). When adding 100 individuals of N_e=100 or N_e=10 to the reference sample

with N_e=1000, the accuracy was slightly or substantially improved (Figure 4A). The

improvement was larger when adding more individuals (500) (Figure 4B). Results

showed that an information source of a smaller N_e was more important when the

samples sizes of each information source were the same. When the total number of

reference data was increased, the importance of adding an information source of a

smaller N_e was relatively decreased (Figure 4). When heritability was higher, overall

accuracy was increased, and the relative contribution from an information source of a

smaller N_e, i.e. the close relatives, was reduced (Figure 5).

Figure 6 confirms again that the smaller N_e, the better the prediction accuracy when

using each information source separately. However, the sample sizes can be also

varied across the information sources, as there are generally a lot fewer close relatives

than individuals from the wider population. In Figure 6A, the accuracy at a sample

size of 100 for N_e=10 was 0.73, which was lower than that of a sample size of 1,000

for N_e=100 (0.81) or that of a sample size of 20,000 for N_e=1,000 (0.83). With a

higher heritability, the result is similar in that the 20,000 records in the information

source of N_e=1000 gave a better accuracy than the 100 records of close relatives

(N_e=10).

In real situations, the most common and desirable design may combine all of the

information sources to maximize the prediction accuracy. We plotted the accuracy

using a composite design consisting of N_e=1000 + N_e=100 (N=500) + N_e=10 (N=50),

compared to that using N_e=1000 (Figure 7). The accuracy for a composite design was

substantially increased especially when the total number of reference sample is low

(Figure 7).

Figure 8 illustrates the real data analyses. The median of empirically estimated M_e

from the inverse of the variance of the genomic relationship (Eq. 5) over 100

replicates was 2254 (SD=50), 3989 (SD=104) and 28848 (SD=920) for reference #1,

#2 and #3, respectively (Table 1). Empirically estimated M_e based on the combined

data was 4836 (SD=106) while expected M_e was 5309 (SD=88), approximately

confirming Eq. (7). The (small) difference between empirical observation and

expectation was probably due to skewed distribution of the variance of the genomic

relationships.

Given M_e, N and h^2, the expected accuracy agreed well with the observed accuracy

when using Framingham data (Figure 9). The reported heritabilities, h^2=0.8 [33-35]

for height and h^2=0.46[36, 37] for BMI, were used. We also quantified the importance

of marker density using the real data. In agreement with Figure 2, the prediction

accuracy is not much decreased even with 50,000 SNPs that were randomly selected

from 389,265 SNPs (Figure 10).

DISCUSSION

This work shows a simple approach for modeling genomic prediction in a reference

data set that contained several subpopulations that differ in relatedness to the target

set, and by modeling these subpopulations as having different effective population

size. The model allows assessing the prediction accuracy before actually conducting

an experiment so that designing genomic prediction can be precise and effective in

animal, plant and human genetics. For example, it can address a question how much

the prediction accuracy can be increased by adding 10,000 (conventionally) unrelated

individuals into the current experiment consisting of 100 relatives in the reference

data. The value for N_e in equation (1) can be approximated based on prior knowledge

of a population, and the relatedness of the sample with the target, possibly supported

by some genotype information that maybe available on cohorts, or samples thereof.

As a reference table, we added approximated value of N_e for structured population of

full and half sibs (Supplementary Tables 1 and 2). Prediction in advance indeed relies

on arbitrary modelling a number of cohorts, but it would be a useful exercise, as

illustrated in the results when considering marker density and various sizes of the

subsets of the training data. The theory is also useful for an animal breeder to predict

the value of genotyped animals in an own herd versus those in a wider references

population consisting of a larger number of more distantly related individuals.

The genotypic and phenotypic information of close and distant relatives of the

proband can be effectively used as a part of the unified reference panel that also

include a large number of individuals that are not related to the predicted subject to

improve the accuracy further as illustrated in Figure 7. For a random sample from the

same homogenous population, e.g. within the same breed or ethnicity, an optimal

design should consist of both close and distant relatives and unrelated individuals, e.g.

a composite design, to maximise the prediction accuracy (Figure 7). That is, the

composite design takes advantage of effective information from smaller number of

relatives while it also use information from a greater number of unrelated individual.

We showed that the prediction accuracy derived for a population with unrelated

individuals turns out to be higher, compared to previous quantifications that

overestimated M_e for a larger number of chromosomes[5, 6, 21, 38]. Using the same

theory, we also showed that the information from close relatives could increase the

accuracy even further, especially for smaller reference populations (Figures 3 − 6). It

is important to note that the assumption about using unrelated individuals in

estimating empirical M_e from genomic relationship[5] is not strictly necessary and can

be relaxed (Eq. (4) and (5)). The theory and empirical observation from simulation

study agreed well (Figure 1) even when using a population with a smaller effective

population size (N_e=50) that consisted of a significant proportion of high relatedness.

Previous studies related to genomic prediction accuracy have suggested that M_e can

be derived from the variation in the differences between realized and expected

relationships[6, 22], i.e. $\mathbf{D} = \mathbf{G} - \mathbf{A}$ where $\mathbf{G}$ is a genomic relationship matrix and $\mathbf{A}$ is

a numerator relationship matrix based on pedigree. Those studies validated their

results also with simulation. If the individuals used in the training set have a low

expected relationship to the target individuals, then there is not much difference

between the variations in **D** versus **G**. However, when some closer relatives are used,

var(**G**) is larger than var(**D**) and M_e is therefore smaller. Note that non-random

sampling of individuals used for the training set can cause a difference between the N_e

of the population that was simulated, and the N_e of the data set that was used for

prediction.

We have not tested the theory for multi-breed reference populations, i.e. those that are

heterogeneous in the sense of consisting of populations from different genetic

background, i.e. different breeds or ethnicities, each with different minor allele

frequencies, different LD structure and different effects for causal variants. Wientjes

et al. (2016)[22] explicitly addressed the problem of different effects for causal

variants (i.e. genetic correlation less than one) when combining data from two

populations. Individuals from different populations share genomic relationships that

are lower than those among members within each population. Evidence in literature

suggests low prediction accuracies when using information from different breeds or

populations, which could be viewed as predicting from populations with very large

N_e. Moreover, we have not considered historical population dynamics such as

bottleneck and admixture, but assumed a constant N_e over the historical generations,

which leads to simplifications that make the formulae tractable and easy to derive.

Further work is required to extend the theory accounting for admixture populations

and historical population dynamics.

We have shown an improved theory for the prediction of the effective number of

chromosome segments, which is a key parameter in genomic prediction accuracy[7].

The theory accounts for the correlation between relationships at different

chromosomes and as a result the effective number of chromosome segments is

smaller than predicted from previous theory[5, 6, 21]. As a result, the increase of the

genomic prediction accuracy appears to be less reliant on higher marker density

unless N_e is very large (e.g. > 10,000) (Figure 2), compared to what have been

quantified by previous theory[5, 6, 21]. The previous theory overestimates M_e (mostly

due to neglecting correlation between chromosomes), therefore underestimates the

proportion of genetic variance at QTL captured by markers. Little improvement of

prediction accuracy with increasing SNP marker density has been empirically

observed in a number of studies[39-41]. This may also have important implication in

genomic prediction as to designing marker density in animal, plant and human

genetics.

The ability to quantify the accuracy in relation to various degrees of relationships (e.g.

close relatives, distant relatives, local or extensive population sample) is important for

predicting outcomes of genomic prediction for specific designs. This study has

addressed this question, and the theory has been implemented in MTG2 software

(https://sites.google.com/site/honglee0707/mtg2). Therefore, a user can know the

expected prediction accuracy and the power[26] before designing an experiment of

genomic prediction. Our approach can be applied both before and after collecting the

data.

ACKNOWLEDGEMENTS

This research is supported by the Australian National Health and Medical Research Council (APP1080157), the Australian Research Council (DP160102126, FT160100229) and the Australian Sheep Industry Cooperative Research Centre. The Framingham Heart Study is conducted and supported by the National Heart, Lung, and Blood Institute (NHLBI) in collaboration with Boston University (Contract No. N01-HC-25195). This manuscript was not prepared in collaboration with investigators of the Framingham Heart Study and does not necessarily reflect the opinions or views of the Framingham Heart Study, Boston University, or NHLBI. Funding for SHARe Affymetrix genotyping was provided by NHLBI Contract N02-HL-64278. SHARe Illumina genotyping was provided under an agreement between Illumina and Boston University. The authors acknowledge useful discussion with Han Mulder that contributed to a clearer paper.

DISCLOSURE DECLARATION
The authors declare no competing financial interests.

486   Table 1. The sample size (N) and empirically observed M_e in each of three different

487   reference data sets and combined data set in the Framingham data analysis.

	N_i (standard deviation)	M_e (standard deviation)
Reference #1	N_1 = 617 (19)	2254 (50)
Reference #2	N_2 = 1254 (30)	3989 (104)
Reference #3	N_3 = 1572 (33)	28848 (920)
Combined all	N_{all} = 3443 (0)	4836 (106)

489   The values were averaged over 100 replicates.

References

1.   Meuwissen T, Hayes B, Goddard M. Prediction of total genetic value using genome-wide dense marker maps. Genetics. 2001;157(4):1819 - 29.
2.   Wray NR, Goddard ME, Visscher PM. Prediction of individual genetic risk to disease from genome-wide association studies. Genome Res. 2007;17:1520-8.
3.   Collins FS, Varmus H. A New Initiative on Precision Medicine. New England Journal of Medicine. 2015;372(9):793-5. doi: doi:10.1056/NEJMp1500523. PubMed PMID: 25635347.
4.   Jannink J-L, Lorenz AJ, Iwata H. Genomic selection in plant breeding: from theory to practice. Brief Funct Genomics. 2010;9(2):166-77. doi: 10.1093/bfgp/elq001.
5.   Goddard ME, Hayes BJ, Meuwissen THE. Using the genomic relationship matrix to predict the accuracy of genomic selection. Journal of Animal Breeding and Genetics. 2011;128(6):409-21. doi: 10.1111/j.1439-0388.2011.00964.x.
6.   Goddard ME. Genomic selection: prediction of accuracy and maximisation of long term response. Genetica. 2009;136:245-57. doi: 10.1007/s10709-008-9308-0.
7.   Lee SH, Weerasinghe WMSP, Wray N, Goddard M, Van der Werf J. Using information of relatives in genomic prediction to apply effective stratified medicine. Scientific Reports. 2017;7:42091. doi: DOI: 10.1038/srep42091.
8.   Tucker G, Loh P-R, MacLeod Iona M, Hayes Ben J, Goddard Michael E, Berger B, et al. Two-Variance-Component Model Improves Genetic Prediction in Family Datasets. The American Journal of Human Genetics. 2015;97(5):677-90. doi: 10.1016/j.ajhg.2015.10.002.
9.   de los Campos G, Vazquez AI, Fernando R, Klimentidis YC, Sorensen D. Prediction of Complex Human Traits Using the Genomic Best Linear Unbiased Predictor. PLoS Genet. 2013;9(7):e1003608. doi: 10.1371/journal.pgen.1003608.
10.   Makowsky R, Pajewski NM, Klimentidis YC, Vazquez AI, Duarte CW, Allison DB, et al. Beyond Missing Heritability: Prediction of Complex Traits. PLoS Genet. 2011;7(4):e1002051. doi: 10.1371/journal.pgen.1002051.
11.   Aulchenko YS, Struchalin MV, Belonogova NM, Axenovich TI, Weedon MN, Hofman A, et al. Predicting human height by Victorian and genomic methods. Eur J Hum Genet. 2009;17(8):1070-5. Epub 2009/02/19. doi: 10.1038/ejhg.2009.5. PubMed PMID: 19223933; PubMed Central PMCID: PMCPmc2986552.
12.   Lee S, van der Werf J, Hayes B, Goddard M, Visscher P. Predicting unobserved phenotypes for complex traits from whole-genome SNP data. PLOS Genet. 2008;4(10):e1000231. doi: 10.1371/journal.pgen.1000231. PubMed PMID: WOS:000261480900026.
13.   Clark SA, Hickey JM, Daetwyler HD, van der Werf JH. The importance of information on relatives for the prediction of genomic breeding values and the implications for the makeup of reference data sets in livestock breeding schemes. Genet Sel Evol. 2012;44:4. Epub 2012/02/11. doi: 10.1186/1297-9686-44-4. PubMed PMID: 22321529; PubMed Central PMCID: PMCPmc3299588.
14.   Legarra A, Robert-Granie C, Manfredi E, Elsen JM. Performance of genomic selection in mice. Genetics. 2008;180(1):611-8. Epub 2008/09/02. doi: 10.1534/genetics.108.088575. PubMed PMID: 18757934; PubMed Central PMCID: PMCPmc2535710.

15.   Habier D, Fernando RL, Garrick DJ. Genomic BLUP Decoded: A Look into
the Black Box of Genomic Prediction. Genetics. 2013;194(3):597-607. doi:
10.1534/genetics.113.152207.
16.   Pszczola M, Strabel T, Mulder HA, Calus MP. Reliability of direct genomic
values for animals with different relationships within and to the reference population.
J Dairy Sci. 2012;95(1):389-400. Epub 2011/12/24. doi: 10.3168/jds.2011-4338.
PubMed PMID: 22192218.
17.   Hayes B, Visscher P, Goddard M. Increased accuracy of artificial selection by
using the realized relationship matrix. Genet Res. 2009;91:47 - 60. PubMed PMID:
doi:10.1017/S0016672308009981.
18.   van der Werf JHJ, Clark SA, Lee SH, editors. Predicting genomic selection
accuracy from heterogeneous sources. Association for the Advancement of Animal
Breeding and Genetics Conference; 2015; Lorne, Australia: AAABG.
19.   Wientjes YC, Veerkamp RF, Calus MP. The effect of linkage disequilibrium
and family relationships on the reliability of genomic prediction. Genetics.
2013;193(2):621-31. Epub 2012/12/26. doi: 10.1534/genetics.112.146290. PubMed
PMID: 23267052; PubMed Central PMCID: PMCPMC3567749.
20.   Rabier C-E, Barre P, Asp T, Charmet G, Mangin B. On the Accuracy of
Genomic Selection. PLoS ONE. 2016;11(6):e0156086. doi:
10.1371/journal.pone.0156086.
21.   Meuwissen T, Hayes B, Goddard M. Accelerating Improvement of Livestock
with Genomic Selection. Annual Review of Animal Biosciences. 2013;1(1):221-37.
doi: doi:10.1146/annurev-animal-031412-103705. PubMed PMID: 25387018.
22.   Wientjes YCJ, Bijma P, Veerkamp RF, Calus MPL. An Equation to Predict
the Accuracy of Genomic Values by Combining Data from Multiple Traits,
Populations, or Environments. Genetics. 2016;202:799-823. doi:
10.1534/genetics.115.183269.
23.   Wray NR, Lee SH, Mehta D, Vinkhuyzen AAE, Dudbridge F, Middeldorp
CM. Research Review: Polygenic methods and their application to psychiatric traits.
Journal of Child Psychology and Psychiatry. 2014;55(10):1068-87. doi:
10.1111/jcpp.12295.
24.   Daetwyler HD, Villanueva B, Woolliams JA. Accuracy of predicting the
genetic risk of disease using a genome-wide approach. PLOS ONE.
2008;3(10):e3395.
25.   Sved JA. Linkage disequilibrium and homozygosity of chromosome segments
in finite populations. Theor Popul Biol. 1971;2(2):125-41. PubMed PMID: 5170716.
26.   Lee SH, Wray NR. Novel genetic analysis for case-control genome-wide
association studies: quantification of power and genomic prediction accuracy. PLOS
ONE. 2013;8(8):e71494. doi: 10.1371/journal.pone.0071494. PubMed PMID:
WOS:000323425700059.
27.   Lee SH, Wray NR, Goddard ME, Visscher PM. Estimating missing
heritability for disease from genome-wide association studies. Am J Hum Genet.
2011;88:294-305. PubMed PMID: S0002-9297(11)00020-6 DOI -
10.1016/j.ajhg.2011.02.002.
28.   Chow S, Sahao J, Wang H. Sample Size Calculations in Clinical Research.
2nd Ed ed2008.
29.   MacCluer JW, VandeBerg JL, Read B, Ryder OA. Pedigree analysis by
computer simulation. Zoo Biology. 1986;5:147-60.
30.   Lee S, van der Werf J. The efficiency of designs for fine-mapping of
quantitative trait loci using combined linkage disequilibrium and linkage. Genetics

Selection Evolution. 2004;36(2):145-61. doi: 10.1051/gse:2003056. PubMed PMID: WOS:000188936300001.

31.   Roach JC, Glusman G, Smit AFA, Huff CD, Hubley R, Shannon PT, et al. Analysis of Genetic Inheritance in a Family Quartet by Whole-Genome Sequencing. Science. 2010;328(5978):636-9. doi: 10.1126/science.1186802.

32.   Splansky GL, Corey D, Yang Q, Atwood LD, Cupples LA, Benjamin EJ, et al. The Third Generation Cohort of the National Heart, Lung, and Blood Institute's Framingham Heart Study: design, recruitment, and initial examination. Am J Epidemiol. 2007;165(11):1328-35. Epub 2007/03/21. doi: 10.1093/aje/kwm021. PubMed PMID: 17372189.

33.   Silventoinen K, Sammalisto S, Perola M, Boomsma DI, Cornes BK, Davis C, et al. Heritability of adult body height: a comparative study of twin cohorts in eight countries. Twin research : the official journal of the International Society for Twin Studies. 2003;6(5):399-408. Epub 2003/11/20. doi: 10.1375/136905203770326402. PubMed PMID: 14624724.

34.   Macgregor S, Cornes BK, Martin NG, Visscher PM. Bias, precision and heritability of self-reported and clinically measured height in Australian twins. Hum Genet. 2006;120(4):571-80. Epub 2006/08/26. doi: 10.1007/s00439-006-0240-z. PubMed PMID: 16933140.

35.   Visscher PM, Medland SE, Ferreira MAR, Morley KI, Zhu G, Cornes BK, et al. Assumption-free estimation of heritability from genome-wide identity-by-descent sharing between full siblings. PLoS Genet. 2006;2(3):e41.

36.   Elks CE, den Hoed M, Zhao JH, Sharp SJ, Wareham NJ, Loos RJ, et al. Variability in the heritability of body mass index: a systematic review and meta-regression. Frontiers in endocrinology. 2012;3:29. Epub 2012/05/31. doi: 10.3389/fendo.2012.00029. PubMed PMID: 22645519; PubMed Central PMCID: PMCPMC3355836.

37.   Wilson JG, Rotimi CN, Ekunwe L, Royal CD, Crump ME, Wyatt SB, et al. Study design for genetic analysis in the Jackson Heart Study. Ethnicity & disease. 2005;15(4 Suppl 6):S6-30-7. Epub 2005/12/02. PubMed PMID: 16317983.

38.   Goddard ME, Hayes BJ. Mapping genes for complex traits in domestic animals and their use in breeding programmes. Nat Rev Genet. 2009;10(6):381-91.

39.   Su G, Brøndum RF, Ma P, Guldbrandtsen B, Aamand GP, Lund MS. Comparison of genomic predictions using medium-density (~54,000) and high-density (~777,000) single nucleotide polymorphism marker panels in Nordic Holstein and Red Dairy Cattle populations. Journal of Dairy Science. 2012;95(8):4657-65. doi: http://dx.doi.org/10.3168/jds.2012-5379.

40.   VanRaden, Paul M., O'Connell, Jeffrey R., Wiggans GR, Weigel KA. Genomic evaluations with many more genotypes. Genetics Selection Evolution. 2011;43(1):1-11. doi: 10.1186/1297-9686-43-10.

41.   Moghaddar N, Swan AA, van der Werf JHJ. Accuracy of genomic prediction for Merino wool traits using high-density marker genotypes. Proc Assoc Advmt Breed Genet 2015;21:165-8.

634   A.

637   B.

640   C.

Figure 1. Expected effective number of chromosome segments (M_e) from previous
studies in 2009[6], 2011[5] and 2013[21] and from Eq. (1) in this study, compared to
empirically observed from M_e simulation when varying the number of chromosomes
each with 1 Morgan long. Effective population size was used as N_e =50 **(A)**, 500 **(B)**
and 1000 **(C)**. This confirms the result from Lee et al. (2017)[7].

661 Figure 2. Accuracy of GBV when varying the number of SNPs for $N_e = 10,000$ **(A)**,
662 1000 **(B)** and 100 **(C)**. The sample size in the reference data was N=12,000, 6000 or
663 3000. The heritability was 0.25.
664

Figure 3. The proportion of genetic variance at QTL captured by markers (b = M / (M_e + M)) when varying the number of SNPs for N_e = 10,000, 1000 and 100.

678 A.

679
680

681 B.

682
683

684 Figure 4. Accuracy of GBV when adding 100 individuals (*N*=100) **(A)** or 500
685 individuals (*N*=500) **(B)** of N_e=100 or N_e=10 to the reference population of N_e=1000.
686 The heritability was 0.25.

687
688

A.

Figure 5. Accuracy of GBV when adding 100 individuals (*N*=100) **(A)** or 500
individuals (*N*=500) **(B)** of N_e=100 or N_e=10 to the reference population of N_e=1000.
The heritability was 0.25. The heritability was 0.75.

701
702 A.

703
704
705 B.

706
707
708 Figure 6. Accuracy of GBV when using N_e=1000 only, N_e=100 only and N_e=10 only
709 with a heritability of 0.25 **(A),** and with a heritability of 0.75 **(B).**
710 For N_e=10 only, the accuracy at a sample size of 100 was 0.73 **(A)** and 0.88 **(B).** For
711 N_e=100 only, the accuracy at a sample size of 1000 was 0.81 **(A)** and 0.92 **(B).** For
712 N_e=1000 only, the accuracy at a sample size of 20,000 was 0.83 **(A)** and 0.93 **(B).**
713
714

715
716   A.

717
718
719   B.

720
721
722   Figure 7. Accuracy of GBV when using a composite design, e.g. N_e=1000 + N_e=100
723   (N=500) + N_e=10 (N=50), compared to Ne=1000 only with a heritability of 0.25 **(A)**
724   and with a heritability of 0.75 **(B).**
725
726

727
728
729 Figure 8. When using Framingham data, empirically estimated M_e based on each of
730 the reference data sets and combined data. Empirically estimated M_e based on
731 combined data is approximately agreed with that from theory (Eq. 5).
732

733
734

735
736
737 Figure 9. When using Framingham data, observed prediction accuracy and expected
738 prediction accuracy with given M_e and N (from Eq. (3)) are agreed well. The reported
739 heritability, h^2=0.8 [33-35] for height and h^2=0.46[36, 37] for BMI, were used.

Figure 10. When using Framingham data, the prediction accuracy is not much decreased even with 50,000 SNPs that were randomly selected from 389,265 SNPs.