Identification and characterization of domesticated bacterial transposases

Frederic Bertels[1,2,*], Jenna Gallie[2] and Paul B Rainey[1,3,4]

[1]New Zealand Institute for Advanced Study, Massey University at Albany, Auckland, New Zealand

[2]Max Planck Institute for Evolutionary Biology, Department of Evolutionary Theory, Plön, Germany

[3]Max Planck Institute for Evolutionary Biology, Department of Microbial Population Biology, Plön, Germany

[4]Ecole Supérieure de Physique et de Chimie Industrielles de la Ville de Paris (ESPCI ParisTech), CNRS UMR

8231, PSL Research University, 75231 Paris Cedex 05, France

* To whom correspondence should be addressed: bertels@evolbio.mpg.de

ABSTRACT

Selfish genetic elements (SGEs), such as insertion sequences (ISs) and transposons are found in most

genomes. Transposons are identifiable by their high copy number and capacity for horizontal transfer. In

contrast REP associated tyrosine transposases (RAYTs), a recently described class of bacterial

transposases, are typically present at just one copy *per* genome. This suggests that RAYTs no longer copy

themselves. Motivated by this possibility we interrogated thousands of fully sequenced bacterial genomes

in order to determine patterns of RAYT diversity, their distribution across chromosomes and accessory

elements, and rate of duplication. RAYTs encompass exceptional diversity and are divisible into at least

five distinct groups. They possess features more similar to housekeeping genes than insertion sequences,

are vertically transmitted and have persisted through evolutionary time to the point where they are now

found in ~19% of all fully sequenced bacterial genomes. Overall, the evolutionary dynamics of RAYTs

suggest that they have been co-opted by host genomes to perform an important but currently unknown

function.

Introduction
Mechanisms for the maintenance of selfish genetic elements (SGEs) have been discussed for over 40 years

(Nevers and Saedler 1977; Doolittle and Sapienza 1980; Charlesworth et al. 1994). To persist in the long term,

the subsequent host generation must contain more SGE copies on average than the previous host generation

(Burt and Trivers 2006). This can be achieved even when the element confers a fitness cost, either through

mechanisms that disproportionately increase copy number through self-replication and horizontal gene transfer

(Doolittle and Sapienza 1980), or by mechanisms that kill offspring devoid of the SGE (*e.g.*, meiotic drive or

toxin-antitoxin (TA) systems (Naito et al. 1995; Hurst et al. 1996; Gerdes et al. 2005)).

While SGEs need not benefit the host (Hickey 1982; Bichsel et al. 2013), many SGEs carry genes that do

contribute toward enhanced host fitness. In some cases SGEs have even been domesticated. That is, they have

evolved to perform functions that provide direct benefit to the host organism. Examples of such fitness

enhancing functions include adaptive immunity, where genes derived from transposases mediate rearrangement

of gene cassettes involved in antigen recognition (Jones and Gellert 2004), stress response systems derived from

bacterial TA systems (Van Melderen and De Bast 2009; Van Melderen 2010), and defence systems against

foreign DNA, such as caspases from CRISPRs (Krupovic et al. 2014).

Recently a new class of bacterial transposase has been described: REP associated tyrosine transposases

(RAYTs) are found in a wide range of bacteria (Nunvar et al. 2010; Bertels and Rainey 2011a; Ton-Hoang et al.

2012). Unlike typical transposases, which are part of insertion sequences (ISs; (Mahillon and Chandler 1998),

RAYTs occur as single copy elements. They are characteristically associated with short repetitive extragenic

palindromic sequences (REPs), and are typically arranged as pairs of REP sequences termed REPINs (REP

doublets forming hairpINs; (Bertels and Rainey 2011b; Bertels and Rainey 2011a). REPINs are non-

autonomous mobile elements that are significantly over-represented in many bacterial genomes, and are likely

dependent upon RAYTs for their dissemination (Nunvar et al. 2010; Bertels and Rainey 2011a; Ton-Hoang et

al. 2012).

Recently it has been argued that because REPs perform distinct functions within bacterial cells, ranging from

transcriptional termination (Espéli et al. 2001) to regulation of translation (Liang et al. 2015), RAYTs may be

domesticated transposable elements (Ton-Hoang et al. 2012; Siguier et al. 2014). While evidence supports the

notion that REP sequences have been co-opted to perform diverse functions, to our knowledge there is little

evidence to suggest this is also true for RAYTs; to date the only function assigned to RAYTs is movement of

REPINs (Nunvar et al. 2010; Bertels and Rainey 2011a; Ton-Hoang et al. 2012). Nonetheless, the idea that

RAYTs are domesticated transposons remains an intriguing hypothesis (Bertels and Rainey 2011a; Bertels and

Rainey 2011b).

Here we report an evolutionary analysis of RAYT diversity across thousands of bacterial genomes. Central to

our analysis is a comparison of the RAYT family with two families of well-characterized housekeeping genes

(*def* and *tpiA*) and two known IS families (IS*200* and IS*110*). We show that RAYTs tend to share characteristics

with housekeeping genes rather than with ISs. We also reveal five divergent types of RAYTs. Our data indicate

that RAYTs have no capacity for selfish replication and so, in addition to within-genome dissemination of

REPINs, RAYTs likely perform some currently unrecognized function that is central to their maintenance

within a broad range of bacterial genomes.

MATERIAL AND METHODS

Acquisition of genome sequences

Bacterial genome sequences were downloaded from the NCBI ftp site on the 11[th] of June 2013

(ftp://ftp.ncbi.nih.gov/genomes/Bacteria/). On that day 2,950 bacterial chromosomes and 2,106 plasmids were

fully sequenced and available for analysis.

Identification of family members by BLAST

For each gene family (RAYTs, IS*200*, IS*100*, *def*, *tpiA*) one arbitrarily chosen protein was used as a query for a

BLAST search. The query sequence for the RAYT family was YafM from *Pseudomonas fluorescens* SBW25

(PFLU_RS20900). Query sequences for housekeeping genes were *P. fluorescens* SBW25 peptide deformylase,

Def (PFLU_RS00090), and *P. fluorescens* SBW25 triosephosphate isomerase TpiA (PFLU_RS25840). Query

sequences for the ISs were ECIAI1_4438 from *Escherichia coli* IAI1 (IS*200* family) and, ISEc*32* from *E. coli*

S88 plasmid pECOS88 (IS*110* family, ECSMS35_RS25240). The sequence of each protein was used to

interrogate 2,950 chromosomes and 2,106 plasmids using TBLASTN (protein search against nucleotide

database).

Search results were analysed as follows. First, all database matches with an e-value of less than 1e-2, 1e-5, 1e-

15, and 1e-20 were recorded. Second, for each of the recorded database matches all coding sequences (CDS)

from Genbank annotations were determined. Third, for overlaps with a single gene, the DNA sequence of the

gene and its encoded amino acid sequence were extracted; for multiple overlaps the longest overlapping open

reading frame was used for sequence extraction; matches without overlapping genes were ignored. Fourth, in

addition to recording the DNA and amino acid sequence of the overlapping gene, the DNA sequence of the 5'

and 3' extragenic space immediately flanking the identified gene was stored.

Identification of duplication events (analysis 2)

For all homologs that occur in the same genome, nucleotide sequences were aligned using the Needleman-

Wunsch algorithm (Needleman and Wunsch 1970) and pairwise identities (the number of sites identical between

two sequences divided by the total number of sites) calculated. All pairs with an identity greater than 95% were

deemed duplicates.

Determination of genome-wide frequencies of flanking 16-mers (analysis 4)

Frequencies for all oligonucleotides of length 16 (16mers) from all replicons (chromosomes and plasmids) were

determined, according to Bertels and Rainey (2011b). In brief, extragenic 5' and 3' 16mer frequencies were

determined for each family member using a sliding window with a step size of one for both leading and lagging

strands. With knowledge of this frequency, the most abundant 16mer from each flanking non-coding DNA

sequence could be determined.

Calculating pairwise identity of amino acid sequences

Pairwise alignments between protein sequences were computed by applying the Needleman-Wunsch algorithm

(Needleman and Wunsch 1970). Pairwise identity is the number of identical sites within the alignment divided

by the total number of sites.

Definition of RAYT protein sequence groups

To determine sequence groups a Markov clustering algorithm (MCL) was applied to a matrix of pairwise

protein identities greater than 26% (Van Dongen 2000). For proteins longer than 80 amino acids a identity

threshold of 24.8% has previously been used for identifying homologs (Sander and Schneider 1991). The

resulting sequence clusters were visualized with Cytoscape (www.cytoscape.org) (Cline, et al. 2007).

Phylogenetic analyses

To build a combined RAYT/IS*200* phylogeny, three protein sequences from each RAYT and IS*200* cluster

containing more than 30 sequences (six RAYT and six IS*200* groups) were randomly selected. These 36

sequences were then aligned with MUSCLE (Edgar 2004) and a phylogeny was generated using PhyML

(Guindon and Gascuel 2003). The IS*200* Group 6 was excluded because sequences from this group could not be

aligned with the rest of the sequences.

RESULTS AND DISCUSSION

Overview of a comparison of RAYTs to IS and housekeeping gene families

ISs persist by replicating within and between genomes (Doolittle and Sapienza 1980; Bichsel et al. 2013), with

dissemination between genomes being mediated by elements capable of horizontal transmission (*e.g.*, plasmids).

Hence, ISs are expected to be present in multiple copies within the genomes of individual bacteria, and to be

over-represented on transmissible elements such as plasmids. In contrast, genes beneficial for bacteria (*e.g.*

housekeeping genes) typically occur in a single copy *per* genome, and are not dependent on horizontal transfer

to persist.

In order to assess whether RAYTs show patterns of diversity typical of ISs or housekeeping genes, we first

identified members of five gene families and then compared familial characteristics. The five gene families are

RAYTs, two IS families (IS*200* and IS*110*), an essential housekeeping gene family (*def*; peptide deformylase),

and a non-essential but highly conserved gene family (*tpiA*; triosephosphate isomerase). For each family, we

investigated the following four characteristics: (1) the average number of family members *per* replicon

(chromosome or plasmid), (2) the proportion of family members for which identical (or nearly identical) copies

are present on the same replicon, (3) the proportion of family members present on plasmids, and (4) the average

replicon-wide frequency of the most common 16bp sequence from the DNA sequences flanking each family

member. The results for the RAYT family were compared to those obtained for the other families.

Identification of members of the RAYT, IS*200*, IS*110*, *def* and *tpiA* gene families

We defined gene families as groups of genes that share a similar sequence with a single representative of that

gene family (the reference gene, arbitrarily chosen). To identify all members of a family, the reference protein

sequence was used in a tBLASTN (basic local alignment search tool; (Altschul et al. 1990)) search against all

publicly available, fully-sequenced bacterial genomes. Each BLAST search was performed at four different e-

value thresholds (1e-20, 1e-15, 1e-5 and 1e-2). The e-value is a measure for sequence similarity; low e-values

(1e-20) indicate high similarity and high e-values (1e-2) indicate low similarity. Hence gene families defined by

less restrictive e-values are expected to be larger as they encompass more divergent genes.

IS and housekeeping gene families are of comparable size (**Figure 1**). At the broadest definition of gene family

(*i.e.* 1e-2), the *def* gene family is the largest (3,841 members; members found in 92% of all bacterial genomes

searched) followed by IS*200* (3,232; members present in 26% of all bacterial genomes searched), IS*110* (2,806;

members found in 26% of all bacterial genomes searched) and *tpiA* (2,782; members present in 93% of all

bacterial genomes searched). The RAYT family is considerably smaller, containing 1,045 members with

members occurring in 19% of all bacterial genomes analysed.

Figure 1. Family size of ISs, RAYTs and housekeeping genes under four decreasingly stringent definitions of sequence relatedness.

As is expected of functionally constrained housekeeping genes, the diversity of these gene families is low, which is indicated by the small number of additional members added to each family with the addition of increasingly divergent sequences. Conversely, the size of the IS*110*, IS*200* and RAYT families increased considerably upon addition of more divergent members, indicating less sequence conservation. This effect was most notable for the RAYT family, for which the family size increased more than ten-fold from the most to the least stringent family definition.

Figure 2. Distribution characteristics of five gene families. (*A*) Average copy number of each gene family *per*

replicon (excluding replicons in which no family members occur). (*B*) Proportion of genes for which at least one

duplicate (pairwise nucleotide identity of ≥95%) was found on the same replicon. (*C*) Proportion of family

members found on plasmids. (*D*) Average frequency of the most abundant 16mer identified in the flanking 5′

and 3′ extragenic regions of each family member. All error bars show standard errors.

Analysis 1: Number of family members *per* replicon

Having determined the number of genes in each family, we next wanted to know how gene family members are

distributed across replicons. That is, for replicons that contain at least one member of a particular gene family,

how many members are present on average (**Figure 2A**)?

Under the most stringent family definition, replicons that harbour IS*200* or IS*110* contain on average between

two and five copies *per* replicon, respectively. Interestingly, the two IS families show opposite trends with

respect to e-value classes. The 531 members comprising the most stringently defined IS*200* gene family (**Figure**

178   **1**) occur on 250 different replicons, hence they are represented on average 2.1 times per replicon. This number
179   increases at lower stringency to approximately four occurrences *per* replicon. Contrastingly, the average number
180   of copies *per* replicon for the IS*110* family decreases from 4.8 to 3.5 as familial stringency is lowered. This
181   means that although IS*100* family size increases with decreasing stringency (**Figure 1**), the number of replicons
182   harbouring IS*100* elements increases at a greater rate.
183
For the two housekeeping gene families, the number of homologs *per* replicon increases with increasing family
size. The extent of the increase, however, is different for the two families. For the *tpiA* family the number of
copies *per* replicon increases only slightly (1.06 to 1.08). For the *def* family the increase in the number of family
members *per* replicon is higher (1.2 to 1.5). This larger increase may have been caused by an ancient
duplication event that led to the addition of a family of *def* paralogs (genes derived from ancient duplication
events) for less stringent family definitions (Bergthorsson et al. 2007; Näsvall et al. 2012).
190
191   RAYTs are present a little more than once *per* replicon when family size is defined by the most stringent e-
192   value criterion, and almost two *per* replicon under the least stringent criterion. This hints at diversification of the
193   RAYT family, which we analyse in detail in later sections.
194

Analysis 2: Proportion of family members that are duplicated on a replicon
Analysis 1 encompasses **all** members of a family present on a replicon. This includes recently duplicated genes,
genes derived from ancient duplications (paralogs) and laterally acquired genes. Of these, recently duplicated
genes are of greatest interest to our study, because they are likely to have arisen from recent self-duplication
events (*i.e.* SGE activity). Accordingly, one would expect the relative size of this gene subset to be considerably
larger for SGEs than for genes with a beneficial function. Thus, taking advantage of the fact that recent
duplication events will have resulted in the presence of gene copies with identical (or nearly so) DNA
sequences, we next identified recently duplicated genes for the five gene families at each e-value. Specifically,
we determined the proportion of each gene family for which at least one nearly identical copy (≥95%) exists on
the same replicon (**Figure 2B**).
205

For the least stringent gene family definition (1e-2), about 66% of all IS*200* (2127/3232) and IS*110* (1874/2806)
genes appear to have been recently duplicated (**Figure 2B**). For these families, the proportion of duplicated
genes correlates with the number of copies *per* replicon (**Figure 2A**; F-statistic for a linear fit of data for both IS
families P=6.5 x 10^{-5}). We observe that, with increasing proportion of duplications, there is an increase in the
number of gene copies *per* replicon in both IS families. No such correlation exists for either RAYTs or
housekeeping genes, as expected from the low proportion of gene duplications (see below). This means that for
ISs the change in the average number of family members *per* replicon (**Figure 1**) is not due to ancient
duplication events leading to diversification of the IS family; rather, it is due to differences in duplication rates
for IS family members more distantly related to the reference gene.
215

In contrast to ISs, the two housekeeping gene families show almost no evidence of recent duplication. At 1e-2,
there were no duplications among *def* gene family members (0/3841), and only 0.2% (3/2782) for the *tpiA*

family for the least stringent family definition. Of all the RAYT genes at 1e-2, 7.7% (80/1045) were found to

have recently duplicated with the additional copy being present on the same replicon. The proportion of RAYT

duplication events is less than that observed for IS*200* and IS*110*, but nevertheless higher than that of

housekeeping gene families. As we show in later sections, this pattern results from the addition of IS*200*

sequences to the RAYT family at less stringent e-values.

Analysis 3: Presence of gene family members on plasmids

In the above analyses, both chromosomes and plasmids were interrogated for the presence of homologous

genes. We next determined the proportion of family members that are found solely on plasmids (**Figure 2C**).

Under the assumption that plasmids facilitate the horizontal transfer of genes, we expect SGEs such as ISs to be

present more frequently than housekeeping genes on plasmids (Bichsel et al. 2013).

ISs are commonly found on plasmids; across all stringency classes, between 13% and 19.6% of IS*200* genes and

between 15% and 17.3% of IS*110* genes were found on plasmids. Contrastingly, members of the two

housekeeping gene families were almost never found on plasmids - only 0.2% to 0.4% of all *def* genes and 0.4%

to 0.7% of all *tpiA* genes occurred on plasmids. For RAYTs, 3.6% of family members occur on plasmids at the

least stringent gene family definition (1e-2). This value is between that obtained for ISs and housekeeping

genes. However at the slightly more stringent e-value of 1e-5, only 0.3% of all RAYTs occur on plasmids, a

value very similar to that found for the housekeeping gene families. Again this observation can be explained by

the addition of IS*200* members at higher e-values.

Analysis 4: Maximum 16mer frequencies in the flanking extragenic space

Many bacterial genomes contain short, palindromic sequences that are overrepresented in extragenic space.

These sequences have been named repetitive extragenic palindromic sequences (REPs) (Higgins et al. 1982;

Stern et al. 1984), and have been shown to replicate as part of a doublet termed a REPIN (Bertels and Rainey

2011a). REPINs are almost always found in the extragenic space flanking RAYT genes. These flanking REPINs

usually contain highly abundant 16bp long sequences (16mers) (Bertels and Rainey 2011a).We generalized this

finding and use the abundance of the most common 16mer of the 5′ and 3′ extragenic flanking regions as a

feature to distinguish RAYTs from other genes.

For the IS*200* and IS*110* families, we found higher frequencies of flanking 16mers than for housekeeping genes:

the average frequency of flanking 16mers ranged between 10 and 48. The most common flanking 16mer for ISs

are typically functionally linked to the IS (Mahillon and Chandler 1998) as they are part of the flanking terminal

repeats. Hence, a significant correlation is expected between the proportion of gene family members that are

duplicates and 16mer frequencies. The higher the proportion of duplicates the higher the flanking 16mer

frequency. Such a correlation is evident from **Figures 2D** and **2B** (F-statistic for a linear fit P=1.7 x 10^{-5})

supporting the notion that the most common flanking 16mer is part of the IS.

We found that 16mers in the adjacent extragenic space of housekeeping genes are repeated infrequently in the

remainder of the replicon: average frequencies range from 5.2 to 5.6 (for different e-values). For 100 randomly

assembled genomes of ~6.7Mb in length (the length of the *P. fluorescens* SBW25 genome), the most abundant

16mer occurs on average 5.08 times in the genome (Bertels and Rainey 2011a). The frequencies for 16mers

flanking housekeeping genes are only slightly higher than these values, indicating that duplications do not affect

flanking 16mer frequencies for housekeeping genes.

The RAYT gene family was found to have the highest frequencies of flanking 16mers across replicons, from an

average of 170 for the most stringent family definition to 41 for the least stringent. As expected there is no

correlation between 16mer frequencies and the proportion of duplicated genes (F-statistic for a linear fit P=0.24,

ranging from 2% for the most stringent family definition to 7.7%). Thus gene duplication events are not the

cause for the observed high frequencies of 16mers flanking RAYT genes. Instead high flanking 16mer

frequencies are caused by RAYTs disseminating neighbouring REPINs throughout the genome.

RAYT characteristics show large variations across sequence groups

In the above analyses, some characteristics of RAYT genes change dramatically with the inclusion of more

diverse family members. For example, the results of analysis 3 indicate that a subset of RAYTs identified only

at the most inclusive e-value (1e-2) is found on plasmids (**Figure 2C**). Further, as family members are added

from a stringent to very loose gene family definition, the size of the RAYT family increases ten-fold, more than

any other sequence family. Together these results raise the possibility that the increase in RAYT gene family

size is attributable to the existence of RAYT subgroups, with more distantly related RAYTs being incorporated

with each relaxation of the family definition. In the following section we investigate this hypothesis.

Figure 3. Protein similarity network and genomic distribution of RAYT groups. (A) RAYT protein

relationship map constructed by Cytoscape using the organic layout (1e-2; (Cline et al. 2007). RAYT groups

identified by MCL are highlighted according to the legend. Nodes represent RAYT proteins. Nodes are
connected if the pairwise amino acid identity is ≥26%. Differences between the Cytoscape protein clustering
(proximity of nodes) and the MCL algorithm (colour of nodes) are due to the difference in the information
provided to MCL and Cytoscape: Cytoscape only receives information on whether or not nodes are connected,
whereas MCL performs coloration based on the exact pairwise similarity between proteins. Graphs (B) to (F)
contain the same information as **Figures 1** and **2**, but for the six largest RAYT groups in (A) as opposed to the
five gene families.

RAYT protein sequences cluster into distinct subgroups
To determine whether the RAYT family consists of distinct subgroups we applied the MCL clustering algorithm
(Van Dongen 2000) to a protein similarity matrix of RAYTs at the least stringent family definition. In the
matrix, each node is a protein, and nodes with ≥26% amino acid identity are connected. This similarity matrix
contains the amino acid identity for each pair of proteins. The clustering algorithm uses this information to
divide the protein network into subgroups of strongly connected nodes. RAYT protein sequences cluster into at
least six subgroups, each containing 30 or more members (**Figure 3A**). For each of these subgroups we
determined the same genomic distribution characteristics as for the five gene families above, at the same e-
values (**Figure 3B-F**).

Characteristics of RAYT protein subgroups
As the protein similarity network in **Figure 3A** shows, the RAYT sequence family is divisible into at least six
groups. The different groups show significantly different responses to the inclusion of increasingly diverse
RAYT family members.
299
300 The effect of broadening family definition on group size is dramatically different between RAYT groups
301 (**Figure 3B**). Group 3 is essentially invariant to the effects of adding family members of increasing diversity,
302 which is a consequence of the fact that the RAYT query sequence stems from this group. Members of Groups 1,
303 2, 4, and 5 appear in the family at 1e-5, and are maximal at 1e-2. Members of Group 6, the group most distantly
304 related to the query sequence, are only included at the broadest family definition.
305
There is little variation in the number of group members *per* replicon at the broadest family definition; the
numbers range from 1.3 (Group 2) to 1.8 (Group 3) copies *per* replicon (**Figure 3C**). Despite this, the
proportion of group members with a nearly identical copy on the same replicon differs dramatically between
groups (**Figure 3D**). Group 6 members (30%) and Group 1 members (6.3%) have the highest proportion of
duplication events, whereas Group 3 has the lowest (0.8% at 1e-2). Surprisingly, although Group 3 has the
lowest proportion of duplicates, it is the group with the highest number of gene copies *per* replicon, possibly
indicating that this group is more functionally diverse than the other RAYT groups.
313
The proportion of group members found on plasmids shows a similar pattern to the proportion of duplicates
observed for each group (**Figure 3E**). Again Group 6 shows by far the highest proportion on plasmids (35% at
1e-2). The only other RAYT group for which any members were found on plasmids was Group 1 (1.2% at 1e-

2), which also had the second highest proportion of duplicates (**Figure 3D)**. Hence, of all RAYT subgroups,

Group 1 and 6 are the two that most closely resemble ISs in terms of their distribution characteristics.

The maximum frequency of 16mers found in extragenic space indicates which of the RAYT groups are likely to

be associated with REP sequences (**Figure 3F**). Only Group 2 (57 replicon-wide occurrences at 1e-2) and

Group 3 (140 occurrences) members are associated with 16mers that are more frequently found across replicons

than those associated with ISs. However, Group 1 (12 occurrences) and Group 4 (14 occurrences) are associated

with 16mers that are more common than 16mers from housekeeping genes. Group 5 (31 occurrences), despite

not showing signs of high duplication rates, is associated with 16mers that are as frequent as those associated

with insertion sequences. Group 6 (5), despite relatively high duplication rates is associated with 16mers that

show low abundance in the genome.

Broadly speaking, the results in this section show that RAYT protein relationships (**Figure 3A**) are correlated

with the similarities and differences observed in their characteristics. For example, in accordance with the

frequency of flanking 16mers, Group 2 and 3 are closely linked in the protein network. Group 1, 4 and 5 are

more loosely linked and Group 6 is the most distantly related group.

The difference in the characteristics of Group 6 to the RAYT family as well as the very loose connection to the

RAYT family in the protein relationship network raises the possibility that Group 6 RAYTs also belong to a

family of ISs. IS*200* is the most likely close relative (Nunvar et al. 2010; Bertels and Rainey 2011a; Ton-Hoang

et al. 2012). To investigate this possibility, the IS*200* family was subjected to identical analyses as the RAYT

family.

Figure 4. Protein similarity network and genomic distribution of IS*200* subgroups. (A) shows a protein

similarity network of the IS*200* family. The map was produced as described in the legend for **Figure 3A**.

Graphs (B) to (F) show the same genomic characteristics as in **Figures 2 and 3**.

Characteristics of IS*200* subgroups

The structure of the IS*200* family is much more homogenous than that of the RAYT family (**Figures 4A** and

3A, respectively). All subgroups are arranged around the largest groups, Group 1 and Group 2. The only

subgroup that is more separate is Group 3, a distinction that is also evident in the very different distribution

characteristics of Group 3.

Group sizes in IS*200* are less evenly distributed than RAYT groups (**Figure 4B**). Groups 1 and 2 (the central

subgroups in **Figure 4A**) have by far the most members (1946 and 894), whereas the remaining groups all have

less than 100 members.

As expected for ISs, the number of gene copies *per* replicon correlates (**Figure 4C**) well with the observed

proportion of duplicates for each group (**Figure 4D**). The proportion of duplicates is high for all IS*200* groups

except for Group 3, for which we did not find a single duplicate. Groups with a high proportion of duplicates

have a high number of gene copies *per* replicon (F-statistic for a linear fit for all groups $P=10^{-12}$). Again the

only outlier is Group 3, which has more gene copies *per* replicon than Group 4 and Group 5, but no duplicates.

The proportion of genes on plasmids is high for four out of the six IS subgroups (**Figure 4E**). As expected from

the previous results, we did not observe any Group 3 members on plasmids. More surprisingly, we also did not

observe Group 4 members on plasmids. This is rather unexpected for an IS with an extremely high replication

rate (proportion of duplicates is 61%).

Typical for ISs, the frequency of flanking 16mers again correlates with the proportion of duplicates (**Figure 4F;**

F-statistic for a linear fit for all groups $P=10^{-6}$), except for Group 3, which is associated with highly abundant

16mers (21 occurrences across replicons at 1e-2) but shows no signs of duplication.

Our analyses show that RAYTs and IS*200* sequences have very different genomic distribution characteristics.

RAYTs show features more typical of housekeeping genes and IS*200* shows features typical of ISs. However, in

each family we identified one subgroup that does not fit into the overall pattern. It is possible that the IS-like

RAYT group belongs to the IS*200* family, considering its distant relationship to the RAYT family. Similarly,

although not as distantly related to the IS*200* query sequence, it is possible that IS*200* Group 3 belongs to the

RAYT family. To investigate these possibilities we performed a combined phylogenetic and network analysis

for the RAYT and IS*200* Groups.

Figure 5. Phylogeny of RAYT and IS*200* family. (A) Phylogenetic tree. From each IS*200* and RAYT group,

three random proteins were selected and aligned with MUSCLE (Edgar 2004). We excluded IS*200* Group 6,

because the alignments could not be calculated correctly when members of this subgroup were included. The

phylogeny was built using PHYML (Guindon and Gascuel 2003). All RAYTs are found in the same clade

except for RAYT Group 6, which clusters within the IS*200* family. This is plausible considering the IS-like

genome distribution characteristics. Green bars indicate low duplication rates and a low presence on plasmids;

red bars indicate high duplication rate and high presence on plasmids. RAYT Group 1 is marked with a

red/green bar as within the RAYT family it has the highest duplication rate and the highest presence on

plasmids. (B) Protein similarity network of the RAYT (left) and IS*200* (right) gene family. We chose 30 random

proteins from the six largest RAYT and IS200 groups and displayed them with Cytoscape identically to **Figures**

3F and **4F**. RAYT Group 6 clusters within IS*200* Group 1 in line with the phylogeny in (A) and the genomic

distribution characteristics in **Figure 3**.

Combined RAYT and IS*200* analysis

The combined RAYT and IS*200* phylogeny shows that the IS*200* and RAYT families are distinct with the

exception of RAYT Group 6, which clusters within the IS*200* phylogeny (**Figure 5A**), In contrast, the RAYT-

like IS*200* Group 3 clusters as a distinct group with the IS*200* family. However, according to the phylogenetic

tree, IS*200* Group 3 is the most closely related IS*200* group to the RAYT family. This raises the possibility that

RAYTs have been domesticated only once and that the IS*200* Group 3 has maintained sequence features similar

to IS*200* sequences. This hypothesis, however, needs to be treated with caution as the high divergence between

the different genes, potential convergent evolution and horizontal gene transfer hinders ability to infer the

phylogenetic history with confidence.

The separation between RAYTs and IS*200* in the phylogenetic tree is not as clear in the protein network (**Figure**

5B). Although RAYTs seem to form relatively coherent clusters, these clusters do not include the RAYT-like

Group 3 from the IS*200* family.

Conclusion

RAYTs are a recently described class of bacterial transposases that share a distant evolutionary relationship with

the IS*200* gene family (Lam and Roth 1983; Lam and Roth 1986; Barabas et al. 2008). The suggestion of a

distant evolutionary relationship comes from the fact that both RAYTs and IS*200* share an identical catalytic

centre (Nunvar et al. 2010; Bertels and Rainey 2011a; Ton-Hoang et al. 2012). Despite this similarity, the data

presented here show that the RAYT and IS*200* families display different characteristics. Members of the IS*200*

gene family show characteristics typical of ISs, whereas members of the RAYT family are, in many respects,

almost indistinguishable from housekeeping genes. The housekeeping-like nature of RAYTs provides a firm

basis for the earlier suggestion that RAYTs are domesticated transposons (Bertels and Rainey 2011a; Bertels

and Rainey 2011b; Ton-Hoang et al. 2012; Siguier et al. 2014).

The housekeeping role of RAYTs is unclear, but the existence of five distinct subfamilies (**Figures 3A & 5B**)

raises the possibility of five different functions. Biochemical studies on RAYTs from *E. coli* provide evidence

of *in vitro* endonuclease and DNA binding activity (Ton-Hoang et al. 2012). Such activity is thought to be

integral to the capacity of RAYTs to catalyse transposition of non-autonomous REP / REPIN elements *via* an

IS*200*-like mechanism (Barabas et al. 2008; Ton-Hoang et al. 2010). But this role alone is insufficient to explain

the maintenance of RAYTs within genomes. In the absence of RAYTs catalysing their own transposition (for

which there is no evidence), there is no mechanism for RAYTs to escape mutational decay, leading to the

prediction that RAYTs must perform some function that benefits the host cell (Bertels and Rainey 2011a;

Bertels and Rainey 2011b).

Although there are various plausible scenarios, it is challenging to conceive of a single RAYT function that

defines both its interaction with REPINs and its beneficial effect on the host bacterium. One possibility is that

the various previously described (and likely co-opted) functions of REPs and REPINs, including effects on

translation, transcription and DNA organization (Newbury et al. 1987; Yang and Ames 1988; Espéli et al. 2001;

Voineagu et al. 2008; Liang et al. 2015), depend on some functional interaction with RAYTs. For example,

binding of the RAYT protein to a particular REP or REPIN sequence may modulate its effects, with the

different classes of RAYT possibly reflecting different specialisations in different bacterial groups. An alternate

idea that we find intriguing — and which draws upon the endonuclease function of RAYTs — is that RAYTs

function in concert with REPINs in a CRISPR-like way (Horvath and Barrangou 2010), with the palindromic

REPIN structure acting as bait for incoming foreign DNA (Tobes and Pareja 2006; Treangen et al. 2009) that is

then inactivated or degraded *via* the endonuclease activity of the RAYT. It is not inconceivable that such a

scenario might also generate antagonistic co-evolution between bait and RAYT promoting the evolution of

divergent RAYT subfamilies.

REFERENCES

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. J Mol Biol
215:403–410.

Barabas O, Ronning DR, Guynet C, Hickman AB, Ton-Hoang B, Chandler M, Dyda F. 2008. Mechanism of
IS*200*/IS*605* family DNA transposases: activation and transposon-directed target site selection. Cell
132:208–220.

Bergthorsson U, Andersson DI, Roth JR. 2007. Ohno's dilemma: evolution of new genes under continuous
selection. Proc Natl Acad Sci U S A 104:17004–17009.

Bertels F, Rainey PB. 2011a. Within-Genome Evolution of REPINs: a New Family of Miniature Mobile DNA
in Bacteria. PLoS Genet. 7:e1002132.

Bertels F, Rainey PB. 2011b. Curiosities of REPINs and RAYTs. Mob Genet Elements 1:262–268.

Bichsel M, Barbour AD, Wagner A. 2013. Estimating the fitness effect of an insertion sequence. J Math Biol
66:95–114.

Burt A, Trivers R. 2006. Genes in Conflict: The Biology of Selfish Genetic Elements. Cambridge,
Massachusetts: Belknap Press of Harvard University Press

Charlesworth B, Sniegowski P, Stephan W. 1994. The evolutionary dynamics of repetitive DNA in eukaryotes.
Nature 371:215–220.

Cline MS, Smoot M, Cerami E, Kuchinsky A, Landys N, Workman C, Christmas R, Avila-Campilo I, Creech
M, Gross B, et al. 2007. Integration of biological networks and gene expression data using Cytoscape. Nat
Protoc 2:2366–2382.

Doolittle WF, Sapienza C. 1980. Selfish genes, the phenotype paradigm and genome evolution. Nature
284:601–603.

Espéli O, Moulin L, Boccard F. 2001. Transcription attenuation associated with bacterial repetitive extragenic
BIME elements. J Mol Biol 314:375–386.

Gerdes K, Christensen SK, Løbner-Olesen A. 2005. Prokaryotic toxin-antitoxin stress response loci. Nat Rev
Micro 3:371–382.

Hickey DA. 1982. Selfish DNA: a sexually-transmitted nuclear parasite. Genetics 101:519–531.

Higgins CF, Ames GF, Barnes WM, Clement JM, Hofnung M. 1982. A novel intercistronic regulatory element
of prokaryotic operons. Nature 298:760–762.

Horvath P, Barrangou R. 2010. CRISPR/Cas, the immune system of bacteria and archaea. Science 327:167–
170.

Hurst LD, Atlan A, Bengtsson BO. 1996. Genetic conflicts. Q Rev Biol 71:317–364.

Jones JM, Gellert M. 2004. The taming of a transposon: V(D)J recombination and the immune system. Immunol
Rev 200:233–248.

Krupovic M, Makarova KS, Forterre P, Prangishvili D, Koonin EV. 2014. Casposons: a new superfamily of
self-synthesizing DNA transposons at the origin of prokaryotic CRISPR-Cas immunity. BMC Biol 12:36.

Lam S, Roth JR. 1983. IS*200*: a *Salmonella*-specific insertion sequence. Cell 34:951–960.

Lam S, Roth JR. 1986. Structural and functional studies of insertion element IS*200*. J Mol Biol 187:157–167.

Liang W, Rudd KE, Deutscher MP. 2015. A Role for REP Sequences in Regulating Translation. Mol Cell
58:431–439.

Mahillon J, Chandler M. 1998. Insertion sequences. Microbiol Mol Biol Rev 62:725–774.

Naito T, Kusano K, Kobayashi I. 1995. Selfish behavior of restriction-modification systems. Science 267:897–
899.

Näsvall J, Sun L, Roth JR, Andersson DI. 2012. Real-time evolution of new genes by innovation, amplification,
and divergence. Science 338:384–387.

Nevers P, Saedler H. 1977. Transposable genetic elements as agents of gene instability and chromosomal
rearrangements. Nature 268:109–115.

Newbury SF, Smith NH, Robinson EC, Hiles ID, Higgins CF. 1987. Stabilization of translationally active
mRNA by prokaryotic REP sequences. Cell 48:297–310.

Nunvar J, Huckova T, Licha I. 2010. Identification and characterization of repetitive extragenic palindromes
(REP)-associated tyrosine transposases: implications for REP evolution and dynamics in bacterial
genomes. BMC Genomics 11:44.

Sander C, Schneider R. 1991. Database of homology-derived protein structures and the structural meaning of
sequence alignment. Proteins 9:56–68.

Siguier P, Gourbeyre E, Chandler M. 2014. Bacterial insertion sequences: their genomic impact and diversity.
FEMS Microbiol Rev 38:865–891.

Stern MJ, Ames GF, Smith NH, Robinson EC, Higgins CF. 1984. Repetitive extragenic palindromic sequences:
a major component of the bacterial genome. Cell 37:1015–1026.

Tobes R, Pareja E. 2006. Bacterial repetitive extragenic palindromic sequences are DNA targets for Insertion
Sequence elements. BMC Genomics 7:62.

Ton-Hoang B, Pasternak C, Siguier P, Guynet C, Hickman AB, Dyda F, Sommer S, Chandler M. 2010. Single-
stranded DNA transposition is coupled to host replication. Cell 142:398–408.

Ton-Hoang B, Siguier P, Quentin Y, Onillon S, Marty B, Fichant G, Chandler M. 2012. Structuring the bacterial
genome: Y1-transposases associated with REP-BIME sequences. Nucleic Acids Res 40:3596–3609.

Treangen TJ, Abraham A-L, Touchon M, Rocha EPC. 2009. Genesis, effects and fates of repeats in prokaryotic
genomes. FEMS Microbiol Rev 33:539–571.

Van Dongen S. 2000. A cluster algorithm for graphs. Report-Information systems:1–40.

Van Melderen L, De Bast MS. 2009. Bacterial toxin-antitoxin systems: more than selfish entities? PLoS Genet.
5:e1000437.

Van Melderen L. 2010. Toxin-antitoxin systems: why so many, what for? Curr Opin Microbiol 13:781–785.

Voineagu I, Narayanan V, Lobachev KS, Mirkin SM. 2008. Replication stalling at unstable inverted repeats:
interplay between DNA hairpins and fork stabilizing proteins. Proc Natl Acad Sci U S A 105:9936–9941.

Yang Y, Ames GF. 1988. DNA gyrase binds to the family of prokaryotic repetitive extragenic palindromic
sequences. Proc Natl Acad Sci U S A 85:8850–8854.