Escape from bacterial diversity: potential enemy release in invading yellow

starthistle (*Centaurea solstitialis*) microbiomes

Patricia Lu-Irving[1], Julia Harenčár[1,2], Hailey Sounart[1,3], Shana R Welles[1], Sarah

M Swope[3], David A Baltrus[4,5], and Katrina M Dlugosch[1]

[1] Department of Ecology and Evolutionary Biology, University of Arizona, Tucson,

Arizona, USA; [2] Current address: Biological Sciences Department, California

Polytechnic State University, San Luis Obispo, California, USA; [3] Department of

Biology, Mills College, Oakland, California, USA; [4] School of Plant Sciences,

University of Arizona, Tucson, Arizona, USA; [5] School of Animal and

Comparative Biomedical Sciences, University of Arizona, Tucson, Arizona, USA.

Author for correspondence:

Patricia Lu-Irving

Email: luirving@email.arizona.edu

Total word count (excluding summary, references and legends):	5753	No. of figures:	7
Summary:	195	No. of Tables:	0
Introduction:	1147	No of Supporting Information files:	1
Materials and Methods:	1908		
Results:	803		
Discussion:	1762		
Acknowledgements:	133		

SUMMARY

- Invasive species may benefit from introduction to new regions where they can escape their natural enemies. Here we examined whether geographic patterns of microbial community composition support a role for enemy escape in the invasion of California, USA by yellow starthistle, a highly invasive plant in western North America.

- We used high-throughput sequencing of the 16S V4 region to characterize bacterial community composition in the phyllosphere, rhizosphere, leaves, and roots of plants from seven populations in California and eight populations in the native European range. We compared bacterial diversity between the native and invaded ranges, and with previously published estimates of plant genetic diversity within each population.

- Bacterial communities differed significantly among plant compartments, and between native and invaded ranges within compartments, with consistently lower diversity in the invaded range. Plant genetic diversity did not explain this pattern in bacterial diversity, but a positive relationship was found within ranges between bacterial diversity in roots and plant genetic diversity within populations.

- Our observation of lower bacterial diversity in the invaded relative to the native range of yellow starthistle is consistent with potential enemy escape, providing some of the first evidence for this scenario in plant microbiomes.

KEYWORDS

bacteria, *Centaurea solstitialis*, endophyte, genetic diversity, invasive species, microbiome, phyllosphere, rhizosphere

INTRODUCTION

Humans continue to transport plant species around the globe, and increasing numbers of these translocations result in the invasive expansion of non-native species into recipient communities (Lonsdale, 1999; Butchart *et al.*, 2010; Essl *et*

al., 2011; Ellis *et al.*, 2012). There is a longstanding hypothesis that many species become invasive after escaping from enemies that reduce invader fitness and limit their populations in their native ranges (Darwin, 1859; Williamson, 1996). Known as the 'Enemy Release' hypothesis, this idea is highly intuitive and forms a basis for the biological control of invasive species (Keane & Crawley, 2002). Initial tests of enemy release focused on quantifying visible changes in above-ground herbivore damage (Keane & Crawley, 2002), but there has been increasing recognition that microbial enemies above- and below-ground can have large effects on plant fitness, and could thus determine whether invasive plants benefit from escaping negative species interactions (Callaway *et al.*, 2004; Colautti *et al.*, 2004; Torchin & Mitchell, 2004; Agrawal *et al.*, 2005; Mitchell *et al.*, 2006; Kulmatiski *et al.*, 2008; van der Putten *et al.*, 2013; Dawson & Schrama, 2016; Faillace *et al.*, 2017).

In recent years, microbial communities have emerged as particularly likely candidates for generating enemy release. Although many interactions between plants and microbes can be beneficial, microbial communities often appear to have negative net effects on plant fitness which may become more negative over time, e.g., via plant-soil feedbacks (Bever, 2003; Reinhart & Callaway, 2006; Kulmatiski *et al.*, 2008; Petermann *et al.*, 2008). It is now apparent that interactions between plants and their microbiomes can vary over space and environment (Nemergut *et al.*, 2013; van der Putten *et al.*, 2013; terHorst & Zee, 2016), creating opportunities for introduced plants to escape the microbial communities that characterize their native ranges. Moreover, evidence is building that reductions in microbial diversity are occurring in response to environmental change and human disturbances, and these reductions in diversity may reduce the resistance of ecosystems to invasion (Schnitzer *et al.*, 2011; Wagg *et al.*, 2014; Dawson & Schrama, 2016; van der Putten *et al.*, 2016).

Invasive plant species have provided some of the best evidence to date that microbial interactions can be locally evolved, and can vary considerably over

geographic regions (Rout & Callaway, 2012). Invaders have been shown to vary

in their response to soil communities from their native and invaded ranges, and

there are now many examples of more favorable interactions between plants and

soil from the invaded range, consistent with escape from enemies (Reinhart *et*

al., 2003; Callaway *et al.*, 2004; Mitchell *et al.*, 2006; Engelkes *et al.*, 2008;

Kulmatiski *et al.*, 2008; Maron *et al.*, 2014; van der Putten *et al.*, 2016). Plant-

microbe interactions which provide net benefits to invasive species can be

explained by reduced negative effects of key microbial pathogens, increased

direct beneficial effects of mutualistic taxa, or increased indirect benefits from

taxa that affect competitors more negatively than they do the invader (Dawson &

Schrama, 2016). These mechanisms should manifest as differences in the

microbial communities associated with invading vs. native plants, specifically as

divergence in taxonomic composition, reduction in diversity, and/or the loss or

gain of groups known to have pathogenic or mutualistic effects, where taxonomic

resolution permits inference of function (Herrera Paredes & Lebeis, 2016).

Release from enemies is expected to be beneficial in and of itself, but it may

further promote invasion by changing the pattern of natural selection on resource

allocation by the invader (Sakai *et al.*, 2001). Plants that require reduced

defenses against negative enemy interactions have the potential to adapt to

invest a larger proportion of resources in traits that increase competitiveness,

reproduction, and/or spread. This idea, known as the Evolution of Increased

Competitive Ability (EICA) hypothesis (Blossey & Notzold, 1995) has received a

great deal of attention but mixed empirical support (Maron *et al.*, 2004; Bossdorf

et al., 2005; Mitchell *et al.*, 2006; Felker-Quinn *et al.*, 2013). Potential contributors

to evolutionary responses to enemy release are likely to become better resolved

as our understanding of microbial community interactions increases, particularly

since adaptive responses to microbial disease are known to be among the most

rapid evolutionary changes that occur in any organism (Tiffin & Moeller, 2006;

Bomblies *et al.*, 2007; Salvaudon *et al.*, 2008).

Here we conduct one of the first comparisons of plant microbiomes between

invading populations and their native source region, explicitly testing for patterns

consistent with enemy release (see also Gundale *et al.*, 2016). We ask whether

changes in plant-associated microbial communities have the potential to

generate enemy escape in the highly invasive plant yellow starthistle (*Centaurea*

solstitialis). Yellow starthistle is native to a wide region of Eurasia and was

introduced to South America in the 1600's and North America in the 1800's as a

contaminant of alfalfa seed (Gerlach, 1997). This herbaceous annual is a

colonizer of grassland ecosystems, and is often called one of the '10 Worst

Weeds of the West' in North America (DiTomaso & Healy, 2007). Its extensive

invasion of California in the USA (>14 million acres; Pitcairn *et al.*, 2006) is well-

studied, and invading genotypes in this region have been shown to grow larger

and produce more flowers than plants in the native range, suggesting an

adaptive shift in resource allocation and an increase in invasiveness (Widmer *et*

al., 2007; Eriksen *et al.*, 2012; Dlugosch *et al.*, 2015). Previous research has

demonstrated that yellow starthistle throughout all of its native and invaded

ranges experiences net fitness reductions when grown with its local soil

communities (Andonian *et al.*, 2011, 2012; Andonian & Hierro, 2011). However,

these studies have also indicated that this negative interaction is weaker (more

favorable) in California, raising the possibility that escape from microbial enemies

has promoted this aggressive invasion.

We sample microbial communities associated with leaves (phyllosphere and

endosphere) and roots (rhizosphere and endosphere) of yellow starthistle plants

in both the California invasion and its source region in Europe. Previous

experiments with fungicide treatments have shown that plant-soil interactions

between yellow starthistle and fungi in California are *more* negative (less

favorable) than those in the native range, inconsistent with a role for fungi in

escape from microbial enemies (Hierro *et al.*, 2016). Here, we focus on bacterial

communities as candidates for a potential role in enemy escape in this system.

We use high-throughput sequencing of prokaryotic ribosomal 16S sequences to

quantify diversity and relative abundance of taxa in yellow starthistle

microbiomes, designing a novel modified peptide nucleic acid clamp (Lundberg

et al., 2013) to reduce non-target sequencing of host plastids. We ask whether

there are patterns of reduced taxonomic diversity and/or potential loss of

pathogens in the invaded range, and whether patterns of diversity in plant-

associated bacteria can be explained by geographic patterns of plant genetic

diversity. Loss of potential pathogens would be consistent with opportunities for

enemy escape that could contribute to the evolution of increased invasiveness in

yellow starthistle's highly successful invasion of California.

MATERIALS AND METHODS

Study species

Yellow starthistle (*Centaurea solstitialis* L., Asteraceae) is an obligately

outcrossing annual plant, diploid throughout its range (Heiser & Whitaker, 1948;

Widmer *et al.*, 2007; Öztürk *et al.*, 2009). Plants form a taproot and grow as a

rosette through mild winter and/or spring conditions, bolting and producing

flowering heads (capitula) throughout the summer. The species is native to

Eurasia, where distinct genetic subpopulations have been identified in

Mediterranean western Europe, central-eastern Europe, Asia (including the

Middle East), and the Balkan-Apennine peninsulas (Barker *et al.*, 2017). The

invasion of California as well as those in South America appear to be derived

almost entirely from western European genotypes (Fig. 1; Barker *et al.*, 2017).

Sample collection

Fifteen populations of yellow starthistle were sampled for microbial communities:

seven populations across the invasion of California, six in western Europe, and

two in eastern Europe (Fig. 1; Supporting Information Table S1). At each

population, plants were sampled every meter (or to the nearest meter mark)

along a 25 meter transect, to yield 25 individuals per population. Individuals in

rosette or early bolting stages were preferentially selected. In one population

(HU29), low plant density yielded 20 individuals along the 25 meter transect.

Using sterile technique, plants were manually pulled and each individual sampled

for phyllosphere, rhizosphere, leaves, and roots using modified versions of

protocols by Lundberg *et al.* (2013) and Lebeis *et al.* (2015) as described below.

A control (blank) sample was collected for each population. Plants were pressed

and dried after sampling, and submitted to the University of Arizona Herbarium

(ARIZ; Supporting Information Table S1).

Phyllosphere and rhizosphere — one to three basal, non-senescent leaves were

collected from each plant, as well as the upper 2-5 cm of the taproot, together

with accompanying lateral roots (excess soil was brushed or shaken off). Leaf

and root samples were placed in individual 50 ml tubes containing 25 ml of sterile

wash solution (45.9 mM NaH_2PO_4, 61.6 mM Na_2HPO_4, 0.1% Tween 20). Tubes

were shaken by hand for one minute (timed). Leaf and root samples were then

removed and stored on ice in separate tubes (leaves in empty tubes, roots in

tubes containing 10 ml of wash solution) until further processing. Wash samples

were stored on ice during transport, then refrigerated at $4^{o}C$. Phyllosphere and

rhizosphere washes were pooled per population, then centrifuged at 2,200 g at

$4^{o}C$ for 15 minutes. Supernatants were discarded, and pellets were air-dried and

stored at $-20^{o}C$ until DNA extraction.

Leaf endosphere — leaves were surface sterilized by submerging in bleach

solution (10% commercial bleach, 0.1% tween 20) for two minutes. Leaves were

then rinsed in distilled water, patted dry using clean kimwipe, and sealed in

individual sterile surgical envelopes (Fisherbrand #01-812-50). Envelopes were

kept in silica gel desiccant until leaf tissue was completely dry, then stored at

room temperature until DNA extraction.

Root surface and endosphere (hereafter 'whole root') — roots were further

washed by shaking in 10 ml of wash solution until visible residual soil was

removed. Washed roots were stored and dried as described above for leaves.

Controls — at each collection site, a tube of sterile wash solution was left

uncapped while sampling plants. Disinfected tools were periodically swished in

the blank wash tube before sterilization and use for the next sample collection.

For each population, rinse water and wipes used to process tissue samples were

represented in controls by rinsing and wiping flame-sterilized forceps, then

swishing the forceps in the blank wash tube. Controls were stored and processed

in the same manner as phyllosphere and rhizosphere samples.

DNA extraction

Extractions were carried out using sterile technique in a laminar flow hood. Leaf

and root DNA was extracted as bulk samples from tissue pooled by population

(15 total populations), and as individual samples from 8 plants from each of 10

populations (80 total individuals). For pooled tissue extractions, equal sections of

leaf tissue (50 mm^2) and root tissue (12.5 mm^3 plus 10 mm of lateral roots) were

collected from each individual sample per population and pooled prior to

extraction. Control (blank) samples were collected for each batch of extractions

by swabbing tools and surfaces, then extracting DNA from the swab head.

All DNA samples were extracted using the MO BIO PowerSoil kit (MO BIO

Laboratories, Inc.). Phyllosphere and rhizosphere DNA was extracted from up to

0.25 g of wash pellets following the standard kit protocol. Leaf and root tissues

were ground to powder or sawdust consistency in liquid nitrogen using sterile

mortars and pestles. Leaf and root DNA was extracted from 20 mg (leaf) or 100

mg (root) of ground tissue with the following modification to the standard

protocol: tissue was incubated at 65^{o}C for 10 minutes in extraction buffer, then

vortexed for 1 minute, followed by a second 10 minute incubation (as described

under "alternative lysis methods" in the kit protocol). Control DNA was extracted

by placing whole swab heads directly into extraction tubes. Extracted DNA was

eluted in PCR-grade water and stored at -20^{o}C pending library preparation.

Library preparation and sequencing

To remove secondary compounds inhibiting PCR, DNA extracted from root and

leaf tissue (together with corresponding blanks) was purified using a ZR-96

genomic DNA clean-up kit (Zymo Research). All DNA concentrations were

quantified using a Qubit fluorometer high-sensitivity assay for double-stranded

DNA (Invitrogen), and standardized to equimolar amounts.

Library preparation followed a dual barcoded two-step PCR protocol. In the first

step (target-specific PCR), the V4 region of the 16S rRNA gene was amplified

using target specific primers (515F and 806R) appended with common sequence

(CS) tags through a linker sequence which varied from two to five nucleotides in

length. Target-specific PCR was carried out using Phusion Flash master mix

(Thermo Scientific) in 25 µl reaction volume in a Mastercycler pro thermocycler

(Eppendorf) under the following conditions: 25 cycles of 1 s at 98°C, 5 s at 78°C,

5 s at 57°C, 15 s at 72°C. Products were visualized on an agarose gel and

diluted by up to 1:15 (depending on yield); 1 µl of diluted product was then used

as template in the second step (barcode-adapter attachment PCR). Using

reagents and equipment as described above, barcoded primer pairs

incorporating Illumina P5 and P7 adapters were used to amplify products from

target-specific PCR in 25 µl reaction volumes under the following conditions: 10

cycles of 1 s at 98°C, 5 s at 78°C, 5 s at 51°C, 15 s at 72°C. Barcoded amplicons

were quantified by fluorometry, pooled in equimolar amounts, cleaned, and

submitted to the University of Idaho's IBEST Genomic Resources Core for QC

and sequencing. Amplicons were multiplexed to use half the capacity of one 2 ×

300 bp run on an Illumina MiSeq platform. Raw sequence data are deposited in

the NCBI Short Read Archive under accession number XXXXXX [pending

submission].

Peptide nucleic acid clamps (PNAs) were included in both PCR steps of library

preparation to block amplification of plant chloroplast and mitochondrial 16S as

recommended by Lundberg *et al.* (2013). Clamp sequences published by

Lundberg *et al.* (2013) were compared with chloroplast and mitochondrial 16S

sequences from yellow starthistle and three other species of Asteraceae with

published organellar genomes (*Centaurea diffusa*, *Helianthus annuus*, *Lactuca*

sativa). We found a single nucleotide mismatch between the Asteraceae

chloroplast 16S and the plastid PNA sequences, and designed an alternative

plastid PNA specific to the Asteraceae sequence (5'—

GGCTCAACTCTGGACAG—3'). All samples for this study were amplified using

the plastid PNA of our design, together with the mitochondrial PNA published by

Lundberg *et al.* (2013). To gauge the effectiveness of our alternative PNA, two

duplicate samples were processed using both PNAs published by Lundberg *et al.*

(2013).

Identification of operational taxa

Demultiplexed paired reads were merged and quality filtered using tools from the

USEARCH package version 9.0.2132 (Edgar & Flyvbjerg, 2015). Merged reads

were truncated to uniform length and primer sequences were removed using a

combination of the seqtk toolkit version 1.2 (github.com/lh3/seqtk) and a custom

script. The UPARSE pipeline (Edgar, 2013) implemented in the USEARCH

package was used for further data processing and analysis: unique sequences

were identified, and those represented only once or twice in the processed read

set were discarded as likely PCR or sequencing errors. Remaining sequences

were clustered into operational taxonomic units (OTUs) at a 97% threshold,

chimeras were filtered out, and per-sample OTU read counts were tabulated

using the UPARSE-OTU algorithm. Assignment of OTUs to nearest taxonomic

match in the Greengenes database (McDonald *et al.*, 2012) was carried out

using the UCLUST algorithm implemented in QIIME version 1.9.1 (Caporaso *et*

al., 2010; Edgar, 2010). Data were further processed using tools from the QIIME

package: reads mapping to chloroplast and mitochondrial OTUs were removed,

and samples were rarefied by plant compartment. Rarefaction levels were

chosen to reflect the distribution of read counts per sample within plant

compartments, subsampling to the minimum number of reads necessary to

include all samples except those that were outliers for low read count.

Microbial community analyses

All statistical analyses were performed in R (R Core Team, 2015). We evaluated
overall differences in bacterial community composition between plant
compartments, and between native and invaded ranges within plant
compartments, by performing non-metric multidimensional scaling (NMDS) using
the R packages vegan (Oksanen *et al.*, 2016) and MASS (Venables & Ripley,
2002). Ordinations were based on Bray-Curtis distances, and were performed
using a two-dimensional configuration to minimize stress, using Wisconsin
double standardized and square root transformed data, with expanded weighted
averages of species scores added to the final NMDS solution. Significant
differences among plant compartments and between native and invaded samples
were assessed using the envfit function in vegan. Ellipses were drawn on NMDS
plots using the vegan function ordiellipse, representing 95% confidence limits of
the standard error of the weighted average of scores.

We further explored the underlying correlates of bacterial community variation
using Principal Components Analysis (PCA; using R function prcomp) for
samples from native and invaded ranges within each plant compartment. We
identified the OTUs with the highest loading on the dominant PC axis of variation
by examining the matrix of variable loadings produced by prcomp. The OTU
composition of samples pooled by population (phyllosphere, rhizosphere, and
bulk root samples; hereafter 'bulk samples') was visualized using a heatmap
generated in ggplot2 (Wickham, 2009). Bulk samples were hierarchically
clustered by Bray-Curtis dissimilarity (hclust function in R) using McQuitty's
method (McQuitty, 1966).

We compared the diversity of OTUs between the native and invaded range for
each plant compartment using both richness (R) and the Shannon diversity index
(H'; Shannon, 1948), which reflects the contributions of both taxonomic richness
and evenness to diversity. These values were calculated using the vegan
package, and compared between native and invaded ranges using a

nonparametric Kruskal-Wallis rank sum test on rarefied read counts. For plant

tissue samples that included multiple individuals per site, we compared the

diversity among sites using a Kruskal-Wallis test within regions.

Finally, we asked whether the geographic distribution of plant-associated

bacterial diversity could be explained by the geographic distribution of genetic

diversity in the plants. Measurements of plant genetic diversity at each of our

sampling sites were obtained from previously published genome-wide marker

analyses by Barker and colleagues (Barker *et al.*, 2017), calculated as the

average proportion of pairwise nucleotide differences between alleles (π) at

variable sites across the yellow starthistle genome. Diversity estimates (H') for

each plant compartment were predicted using linear models that included fixed

effects of plant genetic diversity, region (native vs. invaded), and the interaction

between these two effects.

RESULTS

Sequencing and data processing

Sequencing yielded 9,672,898 read pairs, of which 6,217,852 remained after

merging and quality control; these were 253 bp in length after removing artificial

and primer sequences. The number of raw read counts per sample ranged from

16 to 306,200 with a median of 21,964.

Analysis of the merged and processed reads resulted in 4,014 OTUs, of which 60

were identified as plastid or mitochondrial. Sequences representing yellow

starthistle chloroplast and mitochondrial 16S accounted for 40% and 1% of all

reads, respectively. Amplification of host chloroplast in samples using the

Asteraceae-specific plastid PNA was reduced by up to 51% compared with the

Lundberg *et al.* (2013) PNA (Supporting Information Table S2). This is consistent

with results from a broader comparison of the two PNAs, using five Asteraceae

species, which also found that blocking of host chloroplast amplification was

improved by using the Asteraceae-specific PNA (FitzPatrick *et al.*, unpublished).

Despite PNA blocking activity, 83% of the total reads from leaf endosphere

samples were yellow starthistle chloroplast sequences. This might potentially be

attributable to high chloroplast DNA concentrations relative to endophyte DNA in

leaf tissue total DNA extracts. After removal of chloroplast and mitochondrial

reads, remaining read counts for most leaf endosphere samples were low (Fig.

2), so no further analysis of leaf endosphere bacterial communities was

performed.

Rarefaction levels (chosen to reflect the minimum number of reads per sample

by compartment, not including outliers) were 18,000 reads per sample for

phyllosphere, 17,000 for rhizosphere, and 5,000 for whole root samples. These

levels resulted in the exclusion of six samples which were outliers for low read

count: one phyllosphere (DIA), one rhizosphere (SAZ), one bulk root (CAN), and

three individual root samples (two from SAZ; one from SIE).

Microbial community analyses

Results from NMDS ordinations indicated that bacterial communities differed

overall among the phyllosphere, rhizosphere, and whole root compartments (Fig.

3a; P = 0.001). Within compartments, NMDS further revealed significant

differences between native and invaded range whole root samples (Fig. 3b; P =

0.001) and rhizosphere samples (P = 0.001). Native and invaded range

phyllosphere samples differed with marginal significance (P = 0.05). The

dominant phyla among bacterial communities were Proteobacteria,

Actinobacteria, Bacteroidetes, and Firmicutes (Fig. 4), which is consistent with

the findings of previous characterizations of plant-associated bacterial

communities (reviewed by Bulgarelli *et al.*, 2013). Principal component analyses

suggested that the strongest contributions to changes in bacterial community

composition between the native and invaded ranges were made by shifts in the

representation of *Pseudomonas*, *Erwinia*, *Chryseobacterium*,

Xanthomonadaceae, and *Bacillus* taxa (Supporting Information Fig. S1; Table

S3). Clustering analyses within the phyllosphere and rhizosphere compartments

consistently grouped invaded range samples together, as well as samples from

the source region in western Europe (Supporting Information Fig. S2). Native

range samples from eastern Europe (HU01 and HU29) clustered together in

these compartments but were variable in their relationship to the other regions.

Bulk root samples showed less consistent clustering by range.

Bacterial OTU diversity (H': Fig. 5) was significantly lower in the invaded range in

the phyllosphere (X^2_1 = 5.36, P = 0.02) and rhizosphere (X^2_1 = 6.21, P = 0.01),

and marginally lower in bulk whole root samples (X^2_1 = 3.01, P = 0.08). Diversity

in whole root individual samples (Fig. 5d) did not vary significantly among

populations within the native range (X^2_4 = 1.82, P = 0.77), but did vary

significantly within the invaded range (X^2_4 = 15.30, P = 0.004), such that the two

most extreme populations (TRI, SIE) were significantly different from one another

but not the remaining sites. Bacterial OTU richness (R) values showed patterns

similar to H' in general, but differences between native and invading regions were

much weaker overall (Supporting Information Fig. S3), indicating that both

richness and evenness of OTU representation contributed to differences in

diversity between the ranges. For whole roots, our most extensively sampled

plant compartment, an analysis of OTUs observed across all bulk and individual

samples combined indicated that the native and invaded range shared 51% of

observed OTUs, with 31% fewer unique OTUs observed in the invaded relative

to the native range (Fig. 6a). These patterns were reflected across both major

groups of Proteobacteria and Actinobacteria (Fig. 6b,c).

A linear model predicting microbial diversity (H') from plant diversity was

significant for bulk whole root samples (Fig. 7; $F_{(2,12)}$ = 4.99; P = 0.02; r^2_{adj} =

0.36), with significant main effects of both plant genetic diversity (P = 0.04) and

region (native vs. invaded; P = 0.009). The interaction between these two effects

was not significant (P = 0.69) and was removed from the final model. Similar

linear models did not identify significant effects of plant diversity when predicting

phyllosphere ($P = 0.42$) or rhizosphere ($P = 0.11$) diversity, nor the median

diversity of individual whole root samples at a site ($P = 0.95$).

DISCUSSION

Our study revealed strikingly lower diversity of bacterial communities in yellow

starthistle's invasion of California (USA), relative to native European populations

within and beyond the source region for the invasion. Reduced bacterial diversity

was apparent across phyllosphere, rhizosphere, and whole root communities,

and across dominant bacterial phyla. These patterns are consistent with

opportunities for enemy escape in this invasion and could explain why soils have

more favorable (less negative) effects on yellow starthistle fitness in California

relative to other parts of its range (Andonian *et al.*, 2011, 2012; Andonian &

Hierro, 2011).

In line with other surveys of plant microbiomes, we found that differences in

bacterial communities were greatest among plant compartments (i.e.,

phyllosphere, rhizosphere, and roots; (Bulgarelli *et al.*, 2013; Vandenkoornhuyse

et al., 2015). The numbers and diversity of taxa (OTUs) that we recovered in

samples from each compartment were generally similar in magnitude to those

reported in other studies of prokaryotic 16S sequences, from angiosperm groups

as diverse as e.g. Agavaceae (Coleman-Derr *et al.*, 2016), Brassicaceae

(Bodenhausen *et al.*, 2013), Cactaceae (Fonseca-García *et al.*, 2016), and other

Asteraceae (Leff *et al.*, 2016). Notably, we found that diversity was approximately

twice as high in the roots than the rhizosphere. Higher root endosphere diversity

relative to the rhizosphere is also reported in some other recent studies

(Fonseca-García *et al.*, 2016; Leff *et al.*, 2016), but previous reviews have

concluded that root endosphere communities are typically less diverse than

those in the rhizosphere (Bulgarelli *et al.*, 2013; Vandenkoornhuyse *et al.*, 2015).

Our root collections were not surface sterilized and may represent some of the

rhizoplane/rhizosphere in addition to the endosphere, elevating our estimates of

diversity, though it is also possible that yellow starthistle deviates from previous

trends. In contrast to phyllosphere, rhizosphere, and root compartments, we

recovered few microbial sequences for leaf bacterial endophytes, even when

compared to control (blank) samples, suggesting low sequence coverage due to

persistent chloroplast contamination, and potentially low overall bacterial loads

within yellow starthistle leaves.

Within compartments, differences in community composition between ranges

were substantial and dominated by differences in OTU diversity, which ranged

from 18-40% lower in the invaded range. This variation in diversity is similar in

scale to other studies that have sampled distant geographic locations (e.g.

across regions of North America: (Bodenhausen *et al.*, 2013; Peiffer *et al.*, 2013).

A variety of factors may explain this pattern, including environmental differences

across sites (Fierer & Jackson, 2006; Bulgarelli *et al.*, 2013; Nemergut *et al.*,

2013; Vandenkoornhuyse *et al.*, 2015). Soil type appears to have a particularly

strong influence on microbial communities (e.g. Bulgarelli *et al.*, 2012; Lundberg

et al., 2012), and is known to differ broadly across yellow starthistle's range

(Hierro *et al.*, 2016). In addition, populations in California include temperature

and precipitation environments that are on the warm and dry extreme of yellow

starthistle's climatic niche (Dlugosch *et al.*, 2015), and our sampling was

conducted at the end of a period of severe drought in California (Griffin &

Anchukaitis, 2014; Diffenbaugh *et al.*, 2015), which could have amplified

microbial differences related to climate (Schrama & Bardgett, 2016).

Interestingly, a recent study of grassland plants found that microbial diversity

increased under drought, whereas we found reduced diversity in our drought-

affected invaded range (terHorst *et al.*, 2014).

We also observed an effect of plant genotypic diversity on microbial diversity in

bulk samples of roots. We note that bacterial diversity at the level of the

individual plant did not covary with plant population genetic diversity, indicating

that it was only when samples from different plants were combined that an effect

of plant genotype variation was apparent. Such within-species plant genotype

effects have been observed in other studies and may interact with the effect of

environment to shape microbial communities (Peiffer *et al.*, 2013; terHorst & Zee,

2016). In many cases, it appears that genotype effects are minor relative to site

effects (e.g. (Lundberg *et al.*, 2012; Peiffer *et al.*, 2013; Bodenhausen *et al.*,

2014; Bulgarelli *et al.*, 2015), and we also found that genotypic effects were not

stronger than between-region variation in yellow starthistle. Moreover, plant

genotype effects were only significant in the roots, the only endophytic

compartment analyzed: consistent with plant genotype having the strongest

influence on microbial taxa colonizing within the plant itself (Bulgarelli *et al.*,

2012; Lundberg *et al.*, 2012; Lebeis *et al.*, 2015). The ability of plants to shape

the composition and quantity of endophytic microbial taxa in particular has been

suggested to contribute to a 'core' microbiome shared across environments

(Lundberg *et al.*, 2012; Shade & Handelsman, 2012; Reisberg *et al.*, 2013), and

indeed we found that the majority of OTUs observed in yellow starthistle roots

were shared across invading and native populations on different continents.

Importantly, yellow starthistle's invasion to high density could be a cause rather

than an effect of low microbial diversity. Species invasions and range expansions

have been shown to change microbial composition over short timescales (Collins

et al., 2016; Gibbons *et al.*, 2017). Yellow starthistle invasions are denser than

populations surveyed in the native range by an order of magnitude or more

(Uygur *et al.*, 2004; Andonian *et al.*, 2011). Invaded communities that are

dominated by yellow starthistle include lower diversity of plant species overall

(Seabloom *et al.*, 2003; Zavaleta & Hulvey, 2004; D'Antonio *et al.*, 2007), and low

plant diversity may depress the diversity of plant-associated microbes in the

environment (Garbeva *et al.*, 2004; Schnitzer *et al.*, 2011; Coleman-Derr *et al.*,

2016). Such an effect of plant density could provide an explanation for a general

pattern of weaker plant-soil interactions for invasive species in their introduced

ranges (Kulmatiski *et al.*, 2008; Dawson & Schrama, 2016). These results

reinforce a growing need for explicit observational and experimental tests of the

association between microbial diversity and the potential influences of plant

density, plant community diversity, and environmental gradients (Dawson &

Schrama, 2016).

This study is among the first to examine differences in microbial taxa between

the native and introduced ranges of an invasive species. Gundale and colleagues

(Gundale *et al.*, 2014) also identified more favorable soil interactions in invasions

of lodgepole pine (*Pinus contorta*), and explored potential enemy escape in its

fungal endophyte community (Gundale *et al.*, 2016). For lodgepole pine,

microbial communities differed among regions, but there was no consistent

pattern of loss of potential fungal pathogens or gain of mutualists in the invaded

range, and it remains unclear what part of the soil community is responsible for

observed differences in interactions across ranges (Gundale *et al.*, 2016). Finkel

and colleagues (Finkel *et al.*, 2011; 2016) similarly explored the phyllosphere

community of multiple species of *Tamarix* in native and introduced parts of their

range, finding that communities are most strongly structured by geographic

region. Our study reveals that this type of comparative microbiome approach can

be fruitful for identifying changes in species interactions that might be

contributing to invasion success.

One of the central challenges in testing the hypothesis that invaders are

benefitting from enemy release is quantifying the impact of all types of enemies,

with the microbial community being historically the hardest to observe (Keane &

Crawley, 2002; Beckstead & Parker, 2003; Dawson & Schrama, 2016; Müller *et*

al., 2016; van der Putten *et al.*, 2016; Crawford & Knight, 2017). For yellow

starthistle, a great deal of effort has gone into the identification of potential native

herbivores/seed predators that could be used as biocontrol in California. Six

specialist biocontrol insect species and one fungal foliar pathogen have been

released into this area without resulting in effective control (DiTomaso *et al.*,

2006; Swope & Parker, 2012), suggesting that escape from these species has

not facilitated the invasion. Our finding that invaded populations have not only

unique, but also less diverse bacterial communities, suggests particularly strong

opportunities for pathogen escape in this system.

We have previously argued that yellow starthistle has invaded into a low

competition environment in California, benefitting from the historical loss of plant

competitors for water in this system (Dlugosch *et al.*, 2015). Disturbance of the

native community is critical for yellow starthistle establishment, and functionally

similar native species compete well against it in experiments; however, key

competitors have been lost from the ecosystem due to a variety of perturbations

prior to the yellow starthistle invasion (Zavaleta & Hulvey, 2004; Hooper &

Dukes, 2010; Hierro *et al.*, 2011, 2016; Hulvey & Zavaleta, 2012). Any benefits to

yellow starthistle of reduced bacterial diversity could be independent of these

interactions with native plant species, but there are clear opportunities for these

factors to be related. If a lack of competition allowed yellow starthistle to increase

in density, then this could have reduced plant-associated microbial diversity in

the environment, as noted above. However, while this scenario could explain

lower diversity among bacteria, it appears that density is unlikely to explain

reduced negative interactions with the soil community in California. Yellow

starthistle experiences some of its strongest negative plant-soil feedbacks across

generations in California soils (Andonian *et al.*, 2011), suggesting that the build

up of high plant densities is unlikely to explain patterns of enemy release.

Alternatively, the historical loss of native species diversity in California (D'Antonio

et al., 2007) could have resulted in the loss of associated microbial diversity,

generating particularly strong opportunities for invasion into a system with both

reduced competition and reduced pathogen diversity. Microbial surveys of

remnant native communities, as well as across densities of yellow starthistle

would help to clarify alternative interacting effects of plant and microbial diversity,

and it may be particularly enlightening to explore microbial communities

preserved on native plant specimens pre-dating the extensive invasion of yellow

starthistle into this region.

Conclusions

574
To our knowledge, our study is the first to find evidence consistent with
opportunities for release from microbial enemies during invasion. We find lower
overall bacterial diversity in invading plant populations, similar in scale to
geographic variation in bacterial diversity that has been observed in other
studies. These patterns suggest that yellow starthistle may have benefitted from
introduction into disturbed plant communities with relatively low microbial
diversity. Microbial interactions appear to be important for plant fitness in this
system, but may interact with other factors shaping invasiveness, including
disturbance and lack of effective competition from native plant species. In
particular, escape from both microbial enemies and plant competitors might have
created an opportunity for adaptive allocation of resources away from defensive
functions and towards reproduction and the evolution of increased invasiveness
in yellow starthistle. Comparative surveys of the microbiome in invading and
native populations, as presented here, can reveal important variation in the
species interactions that are shaping patterns of invasion.
590

ACKNOWLEDGMENTS
We thank D. Lundberg and S. Lebeis for helpful discussions regarding sampling
design; G. Reardon and C. Chandeyson for assistance with field collections; E.
Arnold, J. Aspinwall, J. Braasch, E. Carlson, K. Hockett, T. O'Connor, M.
Schneider, J. U'Ren, and N. Zimmerman for 16S library preparation assistance
and discussion; A. Gerritsen, D. New and staff at iBEST for assistance with
sequencing; K. Andonian, J. Hierro, and # reviewers for helpful feedback on the
manuscript. We are particularly indebted to C.E. Morris for hosting the European
sample processing at INRA Station de Pathologie Végétale, Montfavet, France.
Data collection and analyses performed by the IBEST Genomics Resources
Core at the University of Idaho were supported in part by NIH COBRE grant
P30GM103324. This study was supported by USDA grant 2015-67013-23000 to
K.M.D., D.A.B., and S.M.S.
604

AUTHOR CONTRIBUTIONS

P.L-I., D.A.B., and K.M.D. designed the study. P.L-I. and J.H. collected the samples with assistance from H.S., S.M.S., and S.R.W. P.L-I conducted the microbial sequencing and bioinformatics. P.L-I, S.R.W., and K.M.D. analyzed the data. P.L-I and K.M.D. wrote the manuscript, which was edited by all authors.

REFERENCES

Agrawal AA, Kotanen PM, Mitchell CE, Power AG, Godsoe W, Klironomos J. **2005**. Enemy release? an experiment with congeneric plant pairs and diverse above- and belowground enemies. *Ecology* **86**: 2979–2989.

Andonian K, Hierro JL. **2011**. Species interactions contribute to the success of a global plant invader. *Biological Invasions* **13**: 2957–2965.

Andonian K, Hierro JL, Khetsuriani L, Becerra PI, Janoyan G, Villareal D, Cavieres LA, Fox LR, Callaway RM. **2012**. Geographic mosaics of plant-soil microbe interactions in a global plant invasion. *Journal of Biogeography* **39**: 600–608.

Andonian K, Hierro JL, Khetsuriani L, Becerra P, Janoyan G, Villarreal D, Cavieres L, Fox LR, Callaway RM. **2011**. Range-expanding populations of a globally introduced weed experience negative plant-soil feedbacks. *PLoS One* **6**: e20117.

Barker BS, Andonian K, Swope SM, Luster DG, Dlugosch KM. **2017**. Population genomic analyses reveal a history of range expansion and trait evolution across the native and invaded range of yellow starthistle (Centaurea solstitialis). *Molecular Ecology* **26**: 1131–1147.

Beckstead J, Parker IM. **2003**. Invasiveness of Ammophila arenaria: release from soil-borne pathogens? *Ecology* **84**: 2824–2831.

Bever JD. **2003**. Soil community feedback and the coexistence of competitors: conceptual frameworks and empirical tests. *The New Phytologist* **157**: 465–473.

Blossey B, Notzold R. **1995**. Evolution of increased competitive ability in
invasive nonindigenous plants: a hypothesis. *The Journal of Ecology* **83**:
887–889.

Bodenhausen N, Bortfeld-Miller M, Ackermann M, Vorholt JA. **2014**. A
synthetic community approach reveals plant genotypes affecting the
phyllosphere microbiota. *PLoS Genetics* **10**: e1004283.

Bodenhausen N, Horton MW, Bergelson J. **2013**. Bacterial communities
associated with the leaves and the roots of Arabidopsis thaliana (AM
Ibekwe, Ed.). *PloS one* **8**: e56329.

Bomblies K, Lempe J, Epple P, Warthmann N, Lanz C, Dangl JL, Weigel D.
2007. Autoimmune response as a mechanism for a Dobzhansky-Muller-
type incompatibility syndrome in plants. *PLoS Biology* **5**: e236.

Bossdorf O, Auge H, Lafuma L, Rogers WE, Siemann E, Prati D. **2005**.
Phenotypic and genetic differentiation between native and introduced
plant populations. *Oecologia* **144**: 1–11.

Bulgarelli D, Garrido-Oter R, Münch PC, Weiman A, Dröge J, Pan Y,
McHardy AC, Schulze-Lefert P. **2015**. Structure and function of the
bacterial root microbiota in wild and domesticated barley. *Cell Host &*
Microbe **17**: 392–403.

Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, Ahmadinejad
N, Assenza F, Rauf P, Huettel B, Reinhardt R, Schmelzer E, *et al.*
2012. Revealing structure and assembly cues for Arabidopsis root-
inhabiting bacterial microbiota. *Nature* **488**: 91–95.

Bulgarelli D, Schlaeppi K, Spaepen S, Ver Loren van Themaat E, Schulze-
Lefert P. **2013**. Structure and functions of the bacterial microbiota of
plants. *Annual Review of Plant Biology* **64**: 807–838.

Butchart SHM, Walpole M, Collen B, van Strien A, Scharlemann JPW,
Almond REA, Baillie JEM, Bomhard B, Brown C, Bruno J, *et al.* **2010**.
Global biodiversity: indicators of recent declines. *Science* **328**: 1164–
1168.

Callaway RM, Thelen GC, Rodriguez A, Holben WE. **2004**. Soil biota and
exotic plant invasion. *Nature* **427**: 731–733.

Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD,
Costello EK, Fierer N, Peña AG, Goodrich JK, Gordon JI, *et al.* **2010**.
QIIME allows analysis of high-throughput community sequencing data.
Nature Methods **7**: 335–336.

Colautti RI, Ricciardi A, Grigorovich IA, MacIsaac HJ. **2004**. Is invasion
success explained by the enemy release hypothesis? *Ecology Letters* **7**:
721–733.

Coleman-Derr D, Desgarennes D, Fonseca-Garcia C, Gross S, Clingenpeel
S, Woyke T, North G, Visel A, Partida-Martinez LP, Tringe SG. **2016**.
Plant compartment and biogeography affect microbiome composition in
cultivated and native Agave species. *The New Phytologist* **209**: 798–811.

Collins CG, Carey CJ, Aronson EL, Kopp CW, Diez JM. **2016**. Direct and
indirect effects of native range expansion on soil microbial community
structure and function. *The Journal of Ecology* **104**: 1271–1283.

Crawford KM, Knight TM. **2017**. Competition overwhelms the positive plant-soil
feedback generated by an invasive plant. *Oecologia* **183**: 211–220.

D'Antonio CM, Malmstrom C, Reynolds SA, Gerlach J. **2007**. Ecology of
invasive non-native species in California grassland. In: Stromberg MR,, In:
Corbin JD,, In: D'Antonio CM, eds. California grasslands: ecology and
management. Berkeley, California, USA: University of California Press,
67–83.

Darwin C. **1859**. *The Origin of Species*. New York: Random House.

Dawson W, Schrama M. **2016**. Identifying the role of soil microbes in plant
invasions. *The Journal of Ecology* **104**: 1211–1218.

Diffenbaugh NS, Swain DL, Touma D. **2015**. Anthropogenic warming has
increased drought risk in California. *Proceedings of the National Academy*
of Sciences **112**: 3931–3936.

DiTomaso JM, Healy EA. **2007**. *Weeds of California and other western states*.

Oakland, CA: University of California Department of Agriculture and

Natural Resources.

DiTomaso JM, Kyser GB, Pitcairn MJ. **2006**. *Yellow starthistle management*

guide. Berkeley, CA: California Invasive Plant Council.

Dlugosch KM, Alice Cang F, Barker BS, Andonian K, Swope SM, Rieseberg

LH. **2015**. Evolution of invasiveness through increased resource use in a

vacant niche. *Nature Plants* **1**: 15066.

Edgar RC. **2010**. Search and clustering orders of magnitude faster than BLAST.

Bioinformatics **26**: 2460–2461.

Edgar RC. **2013**. UPARSE: highly accurate OTU sequences from microbial

amplicon reads. *Nature Methods* **10**: 996–998.

Edgar RC, Flyvbjerg H. **2015**. Error filtering, pair assembly and error correction

for next-generation sequencing reads. *Bioinformatics* **31**: 3476–3482.

Ellis EC, Antill EC, Kreft H. **2012**. All is not loss: plant biodiversity in the

anthropocene. *PLoS One* **7**: e30535.

Engelkes T, Morriën E, Verhoeven KJF, Bezemer TM, Biere A, Harvey JA,

McIntyre LM, Tamis WLM, van der Putten WH. **2008**. Successful range-

expanding plants experience less above-ground and below-ground enemy

impact. *Nature* **456**: 946–948.

Eriksen RL, Desronvil T, Hierro JL, Kesseli R. **2012**. Morphological

differentiation in a common garden experiment among native and non-

native specimens of the invasive weed yellow starthistle (Centaurea

solstitialis). *Biological Invasions* **14**: 1459–1467.

Essl F, Dullinger S, Rabitsch W, Hulme PE, Hülber K, Jarošík V, Kleinbauer

I, Krausmann F, Kühn I, Nentwig W, *et al.* 2011. Socioeconomic legacy

yields an invasion debt. *Proceedings of the National Academy of Sciences*

108: 203–207.

Faillace CA, Lorusso NS, Duffy S. **2017**. Overlooking the smallest matter:

viruses impact biological invasions. *Ecology Letters* **EarlyView**.

Felker-Quinn E, Schweitzer JA, Bailey JK. **2013**. Meta-analysis reveals
evolution in invasive plant species but little support for Evolution of
Increased Competitive Ability (EICA). *Ecology and Evolution* **3**: 739–751.

Fierer N, Jackson RB. **2006**. The diversity and biogeography of soil bacterial
communities. *Proceedings of the National Academy of Sciences* **103**:
626–631.

Finkel OM, Burch AY, Lindow SE, Post AF, Belkin S. **2011**. Geographical
location determines the population structure in phyllosphere microbial
communities of a salt-excreting desert tree. *Applied and Environmental*
Microbiology **77**: 7647–7655.

Finkel OM, Delmont TO, Post AF, Belkin S. **2016**. Metagenomic signatures of
bacterial adaptation to life in the phyllosphere of a salt-secreting desert
tree. *Applied and Environmental Microbiology* **82**: 2854–2861.

Fonseca-García C, Coleman-Derr D, Garrido E, Visel A, Tringe SG, Partida-
Martínez LP. **2016**. The cacti microbiome: interplay between habitat-
filtering and host-specificity. *Frontiers in Microbiology* **7**: 150.

Garbeva P, van Veen JA, van Elsas JD. **2004**. Microbial diversity in soil:
selection microbial populations by plant and soil type and implications for
disease suppressiveness. *Annual Review of Phytopathology* **42**: 243–270.

Gerlach JD. **1997**. How the West was lost: reconstructing the invasion dynamics
of yellow starthistle and other plant invaders of western rangelands and
natural areas. *California Exotic Pest Plant Council Symposium*
Proceedings **3**: 67–72.

Gibbons SM, Lekberg Y, Mummey DL, Sangwan N, Ramsey PW, Gilbert JA,
Shade A. **2017**. Invasive plants rapidly reshape soil properties in a
grassland ecosystem. *mSystems* **2**: e00178–16.

Griffin D, Anchukaitis KJ. **2014**. How unusual is the 2012–2014 California
drought? *Geophysical Research Letters* **41**: 2014GL062433.

Gundale MJ, Almeida JP, Wallander H, Wardle DA, Kardol P, Nilsson M-C,
Fajardo A, Pauchard A, Peltzer DA, Ruotsalainen S, *et al.* 2016.
Differences in endophyte communities of introduced trees depend on the

phylogenetic relatedness of the receiving forest. *The Journal of Ecology*
104: 1219–1232.

Gundale MJ, Kardol P, Nilsson M-C, Nilsson U, Lucas RW, Wardle DA. **2014**.
Interactions with soil biota shift from negative to positive when a tree
species is moved outside its native range. *The New Phytologist* **202**: 415–
421.

Heiser CB Jr, Whitaker TW. **1948**. Chromosome number, polyploidy, and
growth habit in California weeds. *American Journal of Botany* **35**: 179–
186.

Herrera Paredes S, Lebeis SL. **2016**. Giving back to the community: microbial
mechanisms of plant–soil interactions. *Functional Ecology* **30**: 1043–1052.

Hierro JL, Khetsuriani L, Andonian K, Eren Ö, Villarreal D, Janoian G,
Reinhart KO, Callaway RM. **2016**. The importance of factors controlling
species abundance and distribution varies in native and non-native
ranges. *Ecography* **EarlyView**.

Hierro JL, Lortie CJ, Villarreal D, Estanga-Mollica ME, Callaway RM. **2011**.
Resistance to Centaurea solstitialis invasion from annual and perennial
grasses in California and Argentina. *Biological Invasions* **13**: 2249–2259.

Hooper DU, Dukes JS. **2010**. Functional composition controls invasion success
in a California serpentine grassland. *The Journal of Ecology* **98**: 764–777.

Hulvey KB, Zavaleta ES. **2012**. Abundance declines of a native forb have
nonlinear impacts on grassland invasion resistance. *Ecology* **93**: 378–388.

Keane R, Crawley MJ. **2002**. Exotic plant invasions and the enemy release
hypothesis. *Trends in Ecology & Evolution* **17**: 164–170.

Kulmatiski A, Beard KH, Stevens JR, Cobbold SM. **2008**. Plant–soil
feedbacks: a meta-analytical review. *Ecology Letters* **11**: 980–992.

Lebeis SL, Paredes SH, Lundberg DS, Breakfield N, Gehring J, McDonald
M, Malfatti S, Glavina del Rio T, Jones CD, Tringe SG, *et al.* 2015.
Salicylic acid modulates colonization of the root microbiome by specific
bacterial taxa. *Science* **349**: 860–864.

Leff JW, Lynch RC, Kane NC, Fierer N. **2016**. Plant domestication and the
assembly of bacterial and fungal communities associated with strains of
the common sunflower, Helianthus annuus. *The New Phytologist*.
Lonsdale WM. **1999**. Global patterns of plant invasions and the concept of
invasibility. *Ecology* **80**: 1522–1536.
Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S,
Tremblay J, Engelbrektson A, Kunin V, del Rio TG, *et al.* 2012.
Defining the core Arabidopsis thaliana root microbiome. *Nature* **488**: 86–
90.
Lundberg DS, Yourstone S, Mieczkowski P, Jones CD, Dangl JL. **2013**.
Practical innovations for high-throughput amplicon sequencing. *Nature*
Methods **10**: 999–1002.
Maron JL, Klironomos J, Waller L, Callaway RM. **2014**. Invasive plants escape
from suppressive soil biota at regional scales. *The Journal of Ecology* **102**:
19–27.
Maron JL, Vilà M, Arnason J. **2004**. Loss of enemy resistance among
introduced populations of St. John's Wort (Hypericum perforatum).
Ecology **85**: 3243–3253.
McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A,
Andersen GL, Knight R, Hugenholtz P. **2012**. An improved Greengenes
taxonomy with explicit ranks for ecological and evolutionary analyses of
bacteria and archaea. *The ISME Journal* **6**: 610–618.
McQuitty LL. **1966**. Similarity Analysis by Reciprocal Pairs for Discrete and
Continuous Data. *Educational and psychological measurement* **26**: 825–
831.
Mitchell CE, Agrawal AA, Bever JD, Gilbert GS, Hufbauer RA, Klironomos
JN, Maron JL, Morris WF, Parker IM, Power AG, *et al.* 2006. Biotic
interactions and plant invasions. *Ecology Letters* **9**: 726–740.
Müller G, Horstmeyer L, Rönneburg T, van Kleunen M, Dawson W. **2016**.
Alien and native plant establishment in grassland communities is more

strongly affected by disturbance than above- and below-ground enemies.
The Journal of Ecology **104**: 1233–1242.

Nemergut DR, Schmidt SK, Fukami T, O'Neill SP, Bilinski TM, Stanish LF,
Knelman JE, Darcy JL, Lynch RC, Wickey P, *et al.* **2013**. Patterns and
processes of microbial community assembly. *Microbiology and Molecular*
Biology Reviews **77**: 342–356.

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D,
Minchin PR, O'Hara RB, Simpson GL, Solymos P, *et al.* **2016**. vegan:
Community Ecology Package.

Öztürk M, Martin E, Dinç M, Duran A, Özdemir A, Çetin Ö. **2009**. A
cytogenetical study on some plants taxa in Nizip region (Aksaray, Turkey).
Turkish Journal of Biology **33**: 35–44.

Peiffer JA, Spor A, Koren O, Jin Z, Tringe SG, Dangl JL, Buckler ES, Ley
RE. **2013**. Diversity and heritability of the maize rhizosphere microbiome
under field conditions. *Proceedings of the National Academy of Sciences*
110: 6548–6553.

Petermann JS, Fergus AJF, Turnbull LA, Schmid B. **2008**. Janzen-Connell
effects are widespread and strong enough to maintain diversity in
grasslands. *Ecology* **89**: 2399–2406.

Pitcairn M, Schoenig S, Yacoub R, Gendron J. **2006**. Yellow starthistle
continues its spread in California. *California Agriculture* **60**: 83–90.

van der Putten WH, Bardgett RD, Bever JD, Bezemer TM, Casper BB,
Fukami T, Kardol P, Klironomos JN, Kulmatiski A, Schweitzer JA, *et*
al. **2013**. Plant–soil feedbacks: the past, the present and future
challenges. *The Journal of Ecology* **101**: 265–276.

van der Putten WH, Bradford MA, Pernilla Brinkman E, van de Voorde TFJ,
Veen GF. **2016**. Where, when and how plant–soil feedback matters in a
changing world. *Functional Ecology* **30**: 1109–1121.

R Core Team. **2015**. *R: a language and environment for statistical computing*.
Vienna, Austria: R Foundation for Statistical Computing.

Reinhart KO, Callaway RM. **2006**. Soil biota and invasive plants. *The New*
Phytologist **170**: 445–457.

Reinhart KO, Packer A, Van der Putten WH, Clay K. **2003**. Plant–soil biota
interactions and spatial distribution of black cherry in its native and
invasive ranges. *Ecology letters* **6**: 1046–1050.

Reisberg EE, Hildebrandt U, Riederer M, Hentschel U. **2013**. Distinct
phyllosphere bacterial communities on Arabidopsis wax mutant leaves.
PLoS One **8**: e78613.

Rout ME, Callaway RM. **2012**. Interactions between exotic invasive plants and
soil microbes in the rhizosphere suggest that 'everything is not
everywhere'. *Annals of Botany* **110**: 213–222.

Sakai AK, Allendorf FW, Holt JS, Lodge DM, Molofsky J, With KA,
Baughman S, Cabin RJ, Cohen JE, Ellstrand NC, *et al.* 2001. The
population biology of invasive species. *Annual Review of Ecology and*
Systematics **32**: 305–332.

Salvaudon L, Giraud T, Shykoff JA. **2008/4**. Genetic diversity in natural
populations: a fundamental component of plant–microbe interactions.
Current Opinion in Plant Biology **11**: 135–143.

Schnitzer SA, Klironomos JN, HilleRisLambers J, Kinkel LL, Reich PB, Xiao
K, Rillig MC, Sikes BA, Callaway RM, Mangan SA, *et al.* 2011. Soil
microbes drive the classic plant diversity–productivity pattern. *Ecology* **92**:
296–303.

Schrama M, Bardgett RD. **2016**. Grassland invasibility varies with drought
effects on soil functioning. *The Journal of Ecology* **104**: 1250–1258.

Seabloom EW, Harpole WS, Reichman OJ, Tilman D. **2003**. Invasion,
competitive dominance, and resource use by exotic and native California
grassland species. *Proceedings of the National Academy of Sciences* **100**:
13384–13389.

Shade A, Handelsman J. **2012**. Beyond the Venn diagram: the hunt for a core
microbiome. *Environmental Microbiology* **14**: 4–12.

Shannon CE. **1948**. A Mathematical Theory of Communication. *Bell System*
Technical Journal **27**: 379–423.
Swope SM, Parker IM. **2012**. Complex interactions among biocontrol agents,
pollinators, and an invasive weed: a structural equation modeling
approach. *Ecological Applications* **22**: 2122–2134.
terHorst CP, Lennon JT, Lau JA. **2014**. The relative importance of rapid
evolution for plant-microbe interactions depends on ecological context.
Proceedings of the Royal Society: Biological Sciences **281**: 20140028.
terHorst CP, Zee PC. **2016**. Eco-evolutionary dynamics in plant–soil feedbacks.
Functional Ecology **30**: 1062–1072.
Tiffin P, Moeller DA. **2006**. Molecular evolution of plant immune system genes.
Trends in Genetics **22**: 662–670.
Torchin ME, Mitchell CE. **2004**. Parasites, pathogens, and invasions by plants
and animals. *Frontiers in Ecology and the Environment* **2**: 183–190.
Uygur S, Smith L, Uygur FN, Cristofaro M, Balciunas J. **2004**. Population
densities of yellow starthistle (Centaurea solstitialis) in Turkey. *Weed*
Science **52**: 746–753.
Vandenkoornhuyse P, Quaiser A, Duhamel M, Le Van A, Dufresne A. **2015**.
The importance of the microbiome of the plant holobiont. *The New*
Phytologist **206**: 1196–1206.
Venables WN, Ripley BD. **2002**. Random and Mixed Effects. Statistics and
Computing. Modern Applied Statistics with S. New York: Springer, 271–
300.
Wagg C, Bender SF, Widmer F, van der Heijden MGA. **2014**. Soil biodiversity
and soil community composition determine ecosystem multifunctionality.
Proceedings of the National Academy of Sciences **111**: 5266–5270.
Wickham H. **2009**. *ggplot2: elegant graphics for data analysis*. New York:
Springer-Verlag.
Widmer TL, Guermache F, Dolgovskaia MY, Reznik SY. **2007**. Enhanced
growth and seed properties in introduced vs. native populations of yellow
starthistle (Centaurea solstitialis). *Weed Science* **55**: 465–473.

Williamson M. **1996**. *Biological Invasions*. London: Chapman & Hall.

Zavaleta ES, Hulvey KB. **2004**. Realistic species losses disproportionately

reduce grassland resistance to biological invaders. *Science* **306**: 1175–

1177.

Fig. 1. The distribution (gray) of yellow starthistle and sampling sites for this study. Maps detail the native range in Eurasia (a) and the invasion of western North America (b). Previous work has indicated that western Europe is the source for the severe invasion of California, USA (both in dark shading; Barker et al. 2017). Sampling included seven locations in California (b, filled circles), six locations in western Europe and an additional two locations in eastern Europe (a, open circles).

Fig. 2. Distribution of read counts for bulk samples from all four compartments (native and invading population samples combined), as well as control (blank) samples.

Fig. 3. NMDS plots of bacterial OTU composition in phyllosphere (green), rhizosphere (light blue), and whole root (dark blue) samples for native (open symbols) and invading (closed symbols) populations. Plotted are a) bulk samples for each population, showing overall separation by compartment and by range within compartment, and b) individual whole root samples within native and invading populations. Ellipses indicate 95% confidence intervals for samples grouped by range (native range: dashed line; invaded range: solid line).

(a)   **(b)**

Fig 4. Relative abundance of (proportion of reads mapping to) dominant phyla in (a) phyllosphere, (b) rhizosphere, and (c) whole root bulk samples from native and invaded ranges.

Fig. 5. Comparison of diversity (H') between samples from native and invaded ranges for (a) phyllosphere, (b) rhizosphere, (c) bulk whole roots by population, (d) individual whole roots. Significance levels from Kruskal-Wallis tests indicated with asterisks: *$P <$ 0.05, (*)$P <$ 0.1.

Fig 6. Venn diagrams indicating the number of OTUs shared between native and invaded ranges, and unique to each range, for whole root samples. Shown are OTUs from bulk and individual samples combined, for (a) all OTUs, and for the dominant phyla Proteobacteria (b) and Actinobacteria (c).

Fig 7. Bacterial diversity (H') in bulk whole root samples for each population as a function of the genetic diversity among plants in those populations (calculated as the average proportion of pairwise nucleotide differences between alleles (π) at variable sites across the genome; from Barker *et al.*, 2017). Lines show significant positive relationships (linear model: $P < 0.02$) between microbial and plant diversity in both the native range (open symbols, dashed line) and invaded range (closed symbols, solid line).