Native and invading yellow starthistle (*Centaurea solstitialis*) microbiomes differ in

composition and diversity of bacteria

Patricia Lu-Irving[1,2], Julia Harenčár[1,3], Hailey Sounart[1,4], Shana R Welles[1], Sarah M Swope[4],

David A Baltrus[5,6], and Katrina M Dlugosch[1]

[1]Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, Arizona,

USA; [2]Evolutionary Ecology, Royal Botanic Gardens Sydney, NSW, Australia; [3]Biological

Sciences Department, California Polytechnic State University, San Luis Obispo, California,

USA; [4]Department of Biology, Mills College, Oakland, California, USA; [5]School of Plant

Sciences, University of Arizona, Tucson, Arizona, USA; [6]School of Animal and Comparative

Biomedical Sciences, University of Arizona, Tucson, Arizona, USA.

Author for correspondence:

Patricia Lu-Irving

Email: patricia.luirving@bgcp.nsw.gov.au

Total word count (excluding summary, references and legends):	6323	No. of figures:	6
Summary:	200	No. of Tables:	0
Introduction:	1026	No of Supporting Information files:	1
Materials and Methods:	2122		
Results:	1139		
Discussion:	1899		
Acknowledgements:	136		

SUMMARY

- Invasive species could benefit from introduction to locations with favorable species interactions. Microbiomes are an important source of interactions that vary across regions. We examine whether bacterial communities could explain more favorable microbial interactions in highly invasive populations of yellow starthistle.

- We sequenced amplicons of prokaryotic 16S rRNA genes to characterize bacterial community composition in the phyllosphere, ectorhizosphere, and endorhizosphere of plants from seven invading populations in California, USA and eight native populations in Europe. We tested for differentiation of microbiomes by geography, plant compartment, and plant genotype.

- Bacterial communities differed significantly between native and invaded ranges within plant compartments, with consistently lower diversity in plants from the invaded range. Genera containing known plant pathogens also showed lower diversity in invaded range plants. The diversity of bacteria in roots was positively correlated with plant genotype diversity within both ranges, but this relationship did not explain microbial differences between ranges.

- Our findings reveal changes in the composition and diversity of bacterial interactions in invading plants, consistent with observations of altered soil interactions in this invasion. These results call for further study of the sources of variation in microbiomes and the potential for bacteria to facilitate invasion success.

KEYWORDS

species introductions, invasive species, microbial communities, phyllosphere, rhizosphere, endophytes, genetic diversity

INTRODUCTION

Humans continue to transport plant species around the globe, and increasing numbers of these translocations result in the invasive expansion of non-native species into recipient communities (Lonsdale, 1999; Butchart *et al.*, 2010; Essl *et al.*, 2011; Ellis *et al.*, 2012). While there are undoubtedly many reasons that species introductions lead to invasions, there is growing evidence that novel species interactions may facilitate the invasive spread of populations (Mitchell *et al.*, 2006; Pearson *et al.*, 2018). Initially, hypotheses about the contribution of species interactions to invasions focused on the potential for non-native species to escape from above-ground herbivores, which are easily observed (Keane & Crawley, 2002), though it is not clear that herbivore escape is a frequent mechanism of invasion (Maron & Vila, 2001; Levine *et al.*, 2004; Agrawal *et al.*, 2005; Felker-Quinn *et al.*, 2013). More recently, there has been increasing recognition that microbial taxa above- and below-ground can have large effects on plant fitness, both positive and negative, and could thus determine whether invasive plants benefit from novel species interactions (Callaway *et al.*, 2004; Colautti *et al.*, 2004; Torchin & Mitchell, 2004; Agrawal *et al.*, 2005; Mitchell *et al.*, 2006; Kulmatiski *et al.*, 2008; van der Putten *et al.*, 2016; Faillace *et al.*, 2017). Plant-associated microbial communities have been historically difficult to observe, however, and studies that leverage newly-available tools to quantify differences in these interactions across native and invading populations are needed to test hypotheses for invasion success (Dawson & Schrama, 2016).

Microbial communities have emerged as particularly likely candidates for facilitating invasions. Although many interactions between plants and microbes can be beneficial, soil microbial communities often appear to have negative net effects on plant fitness which may become more negative over time, e.g., via plant-soil feedbacks (Bever, 2003; Reinhart & Callaway, 2006; Kulmatiski *et al.*, 2008; Petermann *et al.*, 2008). These interactions between plants and their microbiomes can vary over space and environment (Nemergut *et al.*, 2013; van der Putten *et al.*, 2013; terHorst & Zee, 2016; Lankau & Keymer, 2018), creating opportunities for introduced plants to escape negative interactions that might characterize their native ranges. Moreover, reductions in microbial diversity are occurring in response to environmental change and human disturbances, and lowered microbial diversity may reduce the resistance of ecosystems to invasion (Schnitzer *et al.*, 2011; Wagg *et al.*, 2014; Dawson & Schrama, 2016; van der Putten *et*

al., 2016).

Invasive plant species have provided some of the best evidence to date that microbial interactions

can be locally evolved, and can vary considerably over geographic regions (Rout & Callaway,

2012). Introduced plants have been shown to vary in their response to soil communities from

their native and invaded ranges, and there are now many examples of more favorable interactions

between plants and soil from their invaded range, consistent with escape from enemies or gain of

mutualists during invasion (Reinhart *et al.*, 2003; Callaway *et al.*, 2004; Mitchell *et al.*, 2006;

Engelkes *et al.*, 2008; Kulmatiski *et al.*, 2008; Maron *et al.*, 2014; van der Putten *et al.*, 2016).

Plant-microbe interactions which provide relative benefits to invasive species can be explained

by reduced negative effects of key microbial pathogens, increased direct beneficial effects of

mutualistic taxa, or increased indirect benefits from taxa that affect competitors more negatively

than they do the invader (Dawson & Schrama, 2016). It is also possible that invaders could

benefit from a reduced diversity of enemy interactions, as a result of an associated reduction in

ecological costs that derive from simultaneously deploying different defense responses against

many different enemies (Agrawal, 2007; Colautti & Lau, 2015; Smakowska *et al.*, 2016). These

hypotheses all require that there are differences in the microbial communities associated with

invading vs. native plants, specifically changes in taxonomic composition, and the loss or gain of

groups with pathogenic or mutalistic effects (Dawson & Schrama, 2016).

Here we conduct one of the first comparisons of plant microbiomes between invading

populations and populations in their native source region. We survey plant-associated microbial

communities in the highly invasive forb yellow starthistle (*Centaurea solstitialis*). Yellow

starthistle is native to a wide region of Eurasia and was introduced from western Europe to South

America in the 1600's and North America in the 1800's as a contaminant of alfalfa seed

(Gerlach, 1997; Barker *et al.*, 2017). This herbaceous annual is a colonizer of grassland

ecosystems, and is often cited as one of the '10 Worst Weeds of the West' in North America

(DiTomaso & Healy, 2007). Its extensive invasion of California in the USA (>14 million acres;

Pitcairn *et al.*, 2006) is particularly well-studied, and invading genotypes in this region have

evolved to grow larger and produce more flowers than plants in the native range, suggesting a

shift in resource allocation that has favored invasiveness (Widmer *et al.*, 2007; Eriksen *et al.*,

2012; Dlugosch *et al.*, 2015). Previous research has demonstrated that yellow starthistle

throughout all of its native and invaded ranges experiences net fitness reductions when grown

with its local soil communities (Andonian *et al.*, 2011, 2012; Andonian & Hierro, 2011). These

studies have also indicated that this negative interaction is weaker (more favorable) in California,

raising the possibility that changes in the microbial community have promoted an aggressive

invasion.

We sample microbial communities associated with leaves (phyllosphere and endosphere) and

roots (ectorhizosphere and endorhizosphere) of yellow starthistle plants in both the California

invasion and native regions in Europe. Previous experiments with fungicide treatments have

shown that plant-soil interactions between yellow starthistle and fungi in California are *more*

negative (less favorable) than those in the native range, inconsistent with a role for fungi in

beneficial species interactions in this invasion (Hierro *et al.*, 2017). Here, we focus on bacterial

communities, using high-throughput sequencing of prokaryotic ribosomal 16S amplicon

sequences to quantify taxonomic composition and diversity of bacteria in yellow starthistle

microbiomes. Microbial communities are known to differ among plant compartments (Bulgarelli

et al., 2013) and to be influenced by individual plant genotype (Lundberg *et al.*, 2012; Peiffer *et*

al., 2013; Bodenhausen *et al.*, 2014), and so we explicitly test for differences in the microbiomes

of native and invaded range plants relative to the influence of both plant compartment and plant

genotype.

MATERIALS AND METHODS

Study species

Yellow starthistle (*Centaurea solstitialis* L., Asteraceae) is an obligately outcrossing annual

plant, diploid throughout its range (Irmina *et al.*, 2017). Plants form a taproot and grow as a

rosette through mild winter and/or spring conditions, bolting and producing flowering heads

(capitula) throughout the summer. The species is native to Eurasia, where distinct genetic

subpopulations have been identified in Mediterranean western Europe, central-eastern Europe,

Asia (including the Middle East), and the Balkan-Apennine peninsulas (Barker *et al.*, 2017). The

invasion of California as well as invasions in South America appear to be derived almost entirely

from western European genotypes (Fig. 1; Barker *et al.*, 2017).

Sample collection

Fifteen populations of yellow starthistle were sampled for microbial communities: seven
sampling sites across the invasion of California, six in western Europe, and two in eastern
Europe (Fig. 1; Supporting Information Table S1). At each location, plants were sampled every
meter (or to the nearest meter mark) along a 25 meter transect, to yield microbial samples from
25 individual plants per population. Individuals in rosette or early bolting stages were
preferentially selected. In one population (HU29), low plant density yielded 20 individuals along
the 25 meter transect. Using sterile technique, plants were manually pulled and each individual
sampled using modified versions of protocols by Lundberg *et al.* (2013) and Lebeis *et al.* (2015)
as described below. Plants were pressed and dried after sampling, and submitted to the
University of Arizona Herbarium (ARIZ; Supporting Information Table S1).

Phyllosphere and ectorhizosphere One to three basal, non-senescent leaves were collected from
each plant, as well as the upper 2-5 cm of the taproot, together with accompanying lateral roots
(excess soil was brushed or shaken off). Leaf and root samples were placed in individual 50 ml
tubes containing 25 ml of sterile wash solution (45.9 mM NaH_2PO_4, 61.6 mM Na_2HPO_4, 0.1%
Tween 20). Tubes were shaken by hand for one minute (timed). Leaf and root samples were then
removed and stored on ice in separate tubes (leaves in empty tubes, roots in tubes containing 10
ml of wash solution) until further processing. Wash samples were stored on ice during transport,
then refrigerated at 4^oC. Phyllosphere and ectorhizosphere washes were pooled across all (20 or
25) plants at a location, then centrifuged at 2,200 g at 4^oC for 15 minutes. Supernatants were
discarded, and pellets were air-dried and stored at -20^oC until DNA extraction.

Leaf endosphere Leaves were surface sterilized by submerging in bleach solution (10%
commercial bleach, 0.1% tween 20) for two minutes. Leaves were then rinsed in distilled water,
patted dry using a kimwipe, and sealed in individual sterile surgical envelopes (Fisherbrand #01-
812-50). Envelopes were kept in silica gel desiccant until leaf tissue was completely dry, then
stored at room temperature until DNA extraction.

Endorhizosphere Roots were further washed by shaking in 10 ml of wash solution until visible

residual soil was removed. Washed roots were stored and dried as described above for leaves.

Controls At each collection site, a tube of sterile wash solution was left uncapped while

sampling plants. Disinfected tools were periodically agitated in the blank wash tube before

sterilization and use for the next sample collection. For each population, rinse water and wipes

used to process tissue samples were represented in controls by rinsing and wiping flame-

sterilized forceps, then agitating the forceps in the blank wash tube. Controls were stored and

processed in the same manner as phyllosphere and ectorhizosphere samples.

DNA extraction

Extractions were carried out using sterile technique in a laminar flow hood. Leaf and root DNA

were extracted from tissue pooled by sampling site (15 total populations), and as individual plant

samples from 8 plants from each of 10 populations (80 total individuals). For pooled tissue

extractions, equal sections of leaf tissue (50 mm^2) and root tissue (12.5 mm^3 plus 10 mm of

lateral roots) were collected from each individual sample per location and pooled prior to

extraction. Control (blank) samples were collected for each batch of extractions by swabbing

tools and surfaces, then extracting DNA from the swab head.

All DNA samples were extracted using the MO BIO PowerSoil kit (MO BIO Laboratories, Inc.).

Phyllosphere and ectorhizosphere DNA was extracted from up to 0.25 g of wash pellets

following the standard kit protocol. Leaf and root tissues were ground to powder or sawdust

consistency in liquid nitrogen using sterile mortars and pestles. Leaf and root DNA was extracted

from 20 mg (leaf) or 100 mg (root) of ground tissue with the following modification to the

standard protocol: tissue was incubated at 65$^{\mathrm{o}}$C for 10 minutes in extraction buffer, then vortexed

for 1 minute, followed by a second 10 minute incubation (as described under "alternative lysis

methods" in the kit protocol). Control DNA was extracted by placing whole swab heads directly

into extraction tubes. Extracted DNA was eluted in PCR-grade water and stored at -20$^{\mathrm{o}}$C

pending library preparation.

Library preparation and sequencing

To remove secondary compounds inhibiting PCR, DNA extracted from root and leaf tissue

(together with corresponding blanks) was purified using a ZR-96 genomic DNA clean-up kit

(Zymo Research). All DNA concentrations were quantified using a Qubit fluorometer high-

sensitivity assay for double-stranded DNA (Invitrogen), and standardized to equimolar amounts.

Library preparation followed a dual-index approach (Kozich *et al.*, 2013) using a two-step PCR

protocol as follows. In the first step (target-specific PCR), the V4 region of the 16S rRNA gene

was amplified using target specific primers (515F and 806R; Caporaso *et al.*, 2011) appended

with common sequence (CS) tags through a linker sequence which varied from two to five

nucleotides in length. Target-specific PCR was carried out using Phusion Flash master mix

(Thermo Scientific) in 25 μl reaction volume in a Mastercycler pro thermocycler (Eppendorf)

under the following conditions: 25 cycles of 1 s at 98^{o}C, 5 s at 78^{o}C, 5 s at 57^{o}C, 15 s at 72^{o}C.

Products were visualized on an agarose gel and diluted by up to 1:15 (depending on yield); 1 μl

of diluted product was then used as template in the second step (barcode-adapter attachment

PCR). Using reagents and equipment as described above, barcoded primer pairs incorporating

Illumina P5 and P7 adapters were used to amplify products from target-specific PCR in 25 μl

reaction volumes under the following conditions: 10 cycles of 1 s at 98^{o}C, 5 s at 78^{o}C, 5 s at

51^{o}C, 15 s at 72^{o}C. Barcoded amplicons were quantified by fluorometry, pooled in equimolar

amounts, cleaned, and submitted to the University of Idaho's IBEST Genomic Resources Core

for QC and sequencing. Amplicons were multiplexed to use half the capacity of one 2 × 300 bp

run on an Illumina MiSeq platform. Raw sequence data are deposited in the NCBI Short Read

Archive under accession number XXXXXX [pending submission].

Peptide nucleic acid clamps (PNAs) were included in both PCR steps of library preparation to

block amplification of plant chloroplast and mitochondrial 16S as recommended by Lundberg *et

al.* (2013). Clamp sequences published by Lundberg *et al.* (2013) were compared with

chloroplast and mitochondrial 16S sequences from yellow starthistle and three other species of

Asteraceae with published organellar genomes (*Centaurea diffusa*, *Helianthus annuus*, *Lactuca

sativa*). We found a single nucleotide mismatch between the Asteraceae chloroplast 16S and the

plastid PNA sequences, and designed an alternative plastid PNA specific to the Asteraceae

sequence (5'—GGCTCAACTCTGGACAG—3') (Fitzpatrick *et al.*, in revision). All samples for

this study were amplified using the plastid PNA of our design, together with the mitochondrial

PNA published by Lundberg *et al.* (2013). To gauge the effectiveness of our alternative PNA,

two duplicate samples were processed using both PNAs published by Lundberg *et al.* (2013).

Identification of operational taxa and potential plant pathogens

Demultiplexed paired reads were merged and quality filtered using tools from the USEARCH

package version 9.0.2132 (Edgar & Flyvbjerg, 2015). Merged reads were truncated to uniform

length and primer sequences were removed using a combination of the seqtk toolkit version 1.2

(github.com/lh3/seqtk) and a custom script. The UPARSE pipeline (Edgar, 2013) implemented

in the USEARCH package was used for further data processing and analysis: unique sequences

were identified, and those represented only once or twice in the processed read set were

discarded as likely PCR or sequencing errors. Remaining sequences were clustered into

operational taxonomic units (OTUs) at a 97% threshold, chimeras were filtered out, and per-

sample OTU read counts were tabulated using the UPARSE-OTU algorithm. Assignment of

OTUs to nearest taxonomic match in the Greengenes database (McDonald *et al.*, 2012) was

carried out using the UCLUST algorithm implemented in QIIME version 1.9.1 (Caporaso *et al.*,

2010; Edgar, 2010). Data were further processed using tools from the QIIME package: reads

mapping to chloroplast and mitochondrial OTUs were removed, and samples were rarefied by

plant compartment. Rarefaction levels were chosen to reflect the distribution of read counts per

sample within plant compartments, subsampling to the minimum number of reads necessary to

include all samples except those that were outliers for low read count.

Taxa known to contain plant pathogens were identified using the FAPROTAX database (version

1.1, (Louca *et al.*, 2016). A list of all genera included under the "plant pathogen" functional

category was used to filter our OTU tables by taxonomic assignment. While pathogen-containing

genera are likely to include both pathogenic and non-pathogenic strains (Newton et al., 2010),

our pathogen-containing OTU dataset should be enriched for potential plant pathogens, relative

to the full dataset.

Microbial community analyses

All statistical analyses were performed in R (R Core Team, 2015). We evaluated differences in

bacterial community composition between plant compartments, and between native and invaded

ranges within plant compartments, by performing non-metric multidimensional scaling (NMDS)

using the R packages vegan (Oksanen *et al.*, 2016) and MASS (Venables & Ripley, 2002).

Individual plant samples or samples pooled within sampling site provided replicates in these

comparisons. Ordinations were based on Bray-Curtis distances, and were performed using a two-

dimensional configuration to minimize stress, using Wisconsin double standardized and square

root transformed data, with expanded weighted averages of species scores added to the final

NMDS solution. Significant differences among plant compartments and between native and

invaded samples were assessed using the envfit function in vegan. Ellipses were drawn on

NMDS plots using the vegan function ordiellipse, representing 95% confidence limits of the

standard error of the weighted average of scores.

We further explored the underlying correlates of bacterial community variation using Principal

Components Analysis (PCA; using R function prcomp) for samples from native and invaded

ranges within each plant compartment. Prior to performing the PCA, we performed Hellinger's

transformation to minimize the influence of OTUs with low counts or many zeros (Legendre &

Legendre, 1998; Legendre & Gallagher, 2001; Ramette, 2007). We then identified the OTUs

with the highest loading on the dominant PC axis of variation by examining the matrix of

variable loadings produced by prcomp. The OTU composition of samples pooled by sampling

site (phyllosphere, ectorhizosphere, and endorhizosphere samples; hereafter 'pooled samples')

was visualized using a heatmap generated in ggplot2 (Wickham, 2009), and samples were

hierarchically clustered by Bray-Curtis dissimilarity (hclust function in R) using McQuitty's

method (McQuitty, 1966).

We compared the diversity of OTUs between the native and invaded range for each plant

compartment using richness (R), evenness (Pielou's J; Pielou, 1966), and their combined effects

via the Hill series exponent $e^{H'}$ (Hill, 1973) of the Shannon diversity index (H'; Shannon, 1948).

Again, individual plant samples or samples pooled by sampling site provided replicates in these

comparisons. Diversity values were calculated using the packages vegan and iNEXT (Hsiegh *et*

al., 2016), and compared between native and invaded ranges using a nonparametric Kruskal-

Wallis rank sum test on rarefied read counts. For plant tissue samples that included multiple

individuals per site, we compared diversity between regions using a nested ANOVA with fixed

effects of region and population nested within region, and among sites using a post hoc Kruskal-

Wallis test within regions. We conducted these comparisons on a dataset including all OTUs, and

a reduced dataset including only OTUs assigned to genera with known plant pathogens

according to the FAPROTAX database, as described above.

Finally, we examined the influence of plant genotype on microbial composition. Our geographic

regions correspond to genetically differentiated subpopulations, and within these regions, our

study sites are also known to vary in plant genetic diversity (Barker *et al.*, 2017). Microbial

diversity estimates ($e^{H'}$) for each plant compartment were predicted using linear models that

included fixed effects of plant genetic diversity, region (native vs. invaded), and the interaction

between these two effects. Measurements of plant genetic diversity at each of our sampling sites

were obtained from previously published genome-wide marker analyses by Barker and

colleagues (Barker *et al.*, 2017), calculated as the average proportion of pairwise nucleotide

differences between alleles (π) at variable positions across the yellow starthistle genome.

RESULTS

Sequencing and data processing

Sequencing yielded 9,672,898 read pairs, of which 6,217,852 remained after merging and quality

control; these were 253 bp in length after removing artificial and primer sequences. The number

of raw read counts per sample ranged from 16 to 306,200 with a median of 21,964. Analysis of

the merged and processed reads resulted in 4,014 OTUs, of which 60 were identified as plastid or

mitochondrial, and 428 were unidentifiable (11%). Of the remaining 3,526, 206 were identified

to species (6%), 1,084 to genus (27%), and 2,229 to family (56%). A total of 103 OTUs (3%)

were identified as members of the 49 genera with known plant pathogens in the FAPROTAX

v.1.1 database.

Sequence reads representing yellow starthistle chloroplast and mitochondrial 16S accounted for

40% and 1% of all reads, respectively. Amplification of host chloroplast in samples using the

Asteraceae-specific plastid PNA was reduced by up to 51% compared with the Lundberg *et al.*

(2013) PNA (Supporting Information Table S2). Despite PNA blocking activity, 83% of the total

reads from leaf endosphere samples were yellow starthistle chloroplast sequences. After removal

of chloroplast and mitochondrial reads, remaining read counts for most leaf endosphere samples

were low relative to controls (Supporting Information Fig. S1), so no further analysis of leaf

endosphere bacterial communities was performed.

Rarefaction levels (chosen to reflect the minimum number of reads per sample by compartment,

not including outliers) were 18,000 reads per sample for phyllosphere, 17,000 for

ectorhizosphere, and 5,000 for endorhizosphere samples (Supporting Information Fig. S1). These

levels were also higher than nearly all control samples. Rarefaction cutoffs resulted in the

exclusion of five non-control samples which were outliers for low read count: one phyllosphere

(DIA), one ectorhizosphere (SAZ), and three individual endorhizosphere samples (two from

SAZ; one from SIE). An NMDS ordination of all un-rarefied samples showed that the controls

clustered together and were clearly differentiated from all samples in all plant compartments

other than the leaf endosphere (Supporting Information Fig. S2).

Microbial community analyses

Results from NMDS ordinations indicated that bacterial communities differed overall among the

phyllosphere, ectorhizosphere, and endorhizosphere compartments (Fig. 2a; stress = 0.14, P =

0.001). Within compartments, NMDS further revealed significant differences between native and

invaded range endorhizosphere samples (Fig. 2b; stress = 0.16, P = 0.001) and ectorhizosphere

samples (P = 0.001). Native and invaded range phyllosphere samples differed with marginal

significance (P = 0.05). Clustering analyses within the phyllosphere and ectorhizosphere

compartments consistently grouped invaded range samples together, as well as samples from the

source region in western Europe (Supporting Information Fig. S3). Native range samples from

eastern Europe (HU01 and HU29) clustered together in these compartments but were variable in

their relationship to the other regions. Endorhizosphere samples pooled by location showed less

consistent clustering by range.

The dominant phyla among all bacterial communities were Proteobacteria, Actinobacteria,

Bacteroidetes, and Firmicutes (Fig. 3). Principal component analyses suggested that the strongest

contributions to changes in bacterial community composition between the native and invaded

ranges were made by shifts in the representation of *Bacillus* (Firmicutes), *Chryseobacterium*

(Bacteroidetes), and the Proteobacteria taxa *Erwinia*, *Pseudomonas*, and Xanthomonadaceae

(Supporting Information Table S3; Fig. S4). All of these taxa other than *Chryseobacterium*

include known plant pathogens in the FAPROTAX v.1.1 database (Louca *et al.*, 2016).

In general, bacterial OTUs showed a pattern of lower median richness (R), evenness (J), and

diversity ($e^{H'}$) for plants from invaded range sites in all compartments, with the exception of

richness in the phyllosphere (Figs 4, 5). Within the phyllosphere, invaders were not significantly

different in richness ($\chi^2_1 = 1.67$, $P = 0.20$), were significantly lower in evenness ($\chi^2_1 = 8.07$, $P = $

0.005), and were marginally lower in Hill diversity ($\chi^2_1 = 3.75$, $P = 0.05$), relative to native range

plants. Similarly, the ectorhizosphere of invaders was not significantly different in richness ($\chi^2_1 = $

0.69, $P = 0.41$), and was significantly lower in both evenness ($\chi^2_1 = 5.0$, $P = 0.03$) and diversity

($\chi^2_1 = 6.21$, $P = 0.01$). In contrast, endorhizosphere samples pooled by site did not differ

significantly in evenness ($\chi^2_1 = 2.26$, $P = 0.13$), but were marginally significantly lower in

richness ($\chi^2_1 = 3.43$, $P = 0.06$) and diversity ($\chi^2_1 = 3.01$, $P = 0.08$). Nested ANOVA of individual

endorhizosphere samples indicated strongly significant reductions in richness, evenness, and

diversity in invading plants (Fig 5; fixed effect of region: all $P < 0.001$). For individual

endorhizosphere samples, populations did not differ significantly in any metrics within

native/invaded regions, with the exception of significantly higher evenness in plants at site SIE

relative to site TRI in the invaded range.

Filtering the rarefied datasets for genera containing known plant pathogens resulted in 49

phyllosphere OTUs, 69 ectorhizosphere OTUs, and 88 endorhizosphere OTUs enriched for

potential plant pathogens. *Pseudomonas* and *Erwinia* were among the most common pathogen-

containing genera encountered in each plant compartment in both ranges. The phyllosphere also

included a high frequency of *Janthinobacterium*, with a relative increase in *Serratia* in the

invaded range. In the ectorhizosphere, *Serratia* was common in both ranges, but invading plants

showed a large relative decline in *Erwinia* and increase in *Pseudomonas*. In the endorhizosphere,

invading plant harbored less *Pseudomonas* and more *Bacillus* and *Streptomyces* than native

plants (Supporting Information Fig. S5). Diversity of these OTUs showed similar trends to total

diversity, with lower median values in invaded range root compartments. No differences between

regions were statistically significant for the phyllosphere ($\chi^2_1 = 0.60$, $P = 0.44$) or

ectorhizosphere ($\chi^2_1 = 0.49$, $P = 0.48$). For the endorhizosphere samples pooled by site,

significantly lower diversity was indicated in the invaded range ($\chi^2_1 = 4.34$, $P = 0.04$). For

individual endorhizosphere samples, nested ANOVA also indicated significantly lower diversity

in the invaded range ($P < 0.0001$), and no significant differences among populations within

regions.

The diversity of bacteria in root compartments showed evidence of associations with plant

genetic diversity at sampling sites. A linear model predicting microbial diversity ($e^{H'}$) from plant

genetic diversity was significant for endorhizosphere samples pooled by site (Fig. 6; $F_{(2,12)} =$

5.89; $P = 0.02$; $r^2_{adj} = 0.41$), with significant main effects of both plant genetic diversity ($P =$

0.03) and region (native vs. invaded; $P = 0.006$). The interaction between these two effects was

not significant ($P = 0.71$) and was removed from the final model. This same pattern was

marginally significant when using only OTUs from pathogen-containing genera in the

endorhizosphere samples (effect of plant genetic diversity: $P = 0.08$). Similar linear models did

not identify significant effects of plant genetic diversity when predicting the median diversity of

individual plant endorhizosphere samples ($P = 0.74$), or diversity in the phyllosphere ($P = 0.35$).

There was a marginally significant positive effect of plant genetic diversity on diversity in the

ectorhizosphere ($P = 0.08$) again in addition to the effect of region ($P = 0.002$).

DISCUSSION

Introduced plants will encounter a variety of novel species interactions as they establish across

biogeographic regions. For many plants, severe invasions are associated with more favorable

interactions with soil microbial communities (Andonian & Hierro, 2011; Dawson & Schrama,

2016). We found that bacterial microbiomes of invading yellow starthistle were unique in

composition and lower in diversity relative to bacterial microbiomes of native plants, differences

that persisted within plant compartments and across variation in plant genetic diversity.

As observed in other species, bacterial communities differed most among plant compartments

(Bulgarelli *et al.*, 2013; Vandenkoornhuyse *et al.*, 2015). The numbers and diversity of taxa

within each compartment were similar in magnitude to those reported in other studies of

prokaryotic 16S sequences, e.g. from Agavaceae (Coleman-Derr *et al.*, 2016), Brassicaceae

(Bodenhausen *et al.*, 2013), Cactaceae (Fonseca-García *et al.*, 2016), and other Asteraceae (Leff

et al., 2016). The dominant phyla were Proteobacteria, Actinobacteria, Bacteroidetes, and

Firmicutes, which are also characteristic of plant-associated bacterial communities surveyed to

date (Bulgarelli *et al.*, 2013). The exception was the leaf endosphere, where a paucity of

sequences relative to controls suggests that persistent chloroplast contamination obscured low

frequency endophytes, despite our development of an Asteraceae-specific PNA (FitzPatrick *et*

al., submitted). A targeted survey is needed to better characterize this compartment (e.g. qPCR,

(Fierer et al., 2005).

Notably, diversity was approximately twice as high in the endorhizosphere as in the

ectorhizosphere. Current reviews have concluded that root endosphere communities are typically

less diverse than those in the ectorhizosphere (Bulgarelli *et al.*, 2013; Vandenkoornhuyse *et al.*,

2015). Our root collections were washed but not surface sterilized and may represent some of the

rhizoplane/rhizosphere in addition to the endosphere, elevating our estimates of diversity. It is

also possible that yellow starthistle deviates from initially reported patterns, which have also

been challenged by other recent studies (Fonseca-García *et al.*, 2016; Leff *et al.*, 2016).

Within compartments, community composition was consistently different between samples from

native and invaded ranges and included shifts in taxa across all major groups. Our native range

samples represented a larger geographic area and spanned distinct genetic subpopulations of

yellow starthistle, but native range sites clustered together in overall community composition and

there was little evidence of individual site differences within ranges. Between ranges, the

diversity of OTUs was lower in the invaded range, a pattern that was dominated by lower

evenness of OTUs in both the phyllosphere and ectorhizosphere, and by lower richness of OTUs

in the endorhizosphere. Thus invading plants were more strongly dominated by a few taxa at

high relative abundance on root and leaf surfaces, and harbored fewer bacterial taxa in their root

endophytic communities.

We observed a significant positive association between root microbial diversity and genotypic

diversity among plants at the population scale. This association was strongest in the

endorhizosphere, the only endophytic compartment in our analysis, consistent with plant

genotype having the largest influence on microbial taxa colonizing within the plant itself

(Bulgarelli *et al.*, 2012; Lundberg *et al.*, 2012; Lebeis *et al.*, 2015). Within-species plant

genotype effects have been observed previously and may interact with the effect of environment

to shape microbial communities (Peiffer *et al.*, 2013; terHorst & Zee, 2016). Interactions

between plant and microbial diversity could be particularly important for invasive species, where

genetic bottlenecks during establishment and range expansion can reduce genetic diversity

among plants (Dlugosch & Parker, 2008; Excoffier *et al.*, 2009; Uller & Leimu 2011).

Nevertheless, it appears that genotype effects are often minor relative to site effects (Lundberg *et*

al., 2012; Peiffer *et al.*, 2013; Bodenhausen *et al.*, 2014; Bulgarelli *et al.*, 2015), and we found

that genotypic effects were evident only within region and did not explain microbial diversity

differences between regions.

The microbial differences that we observed across ranges are similar to those observed at

continental scales (Bodenhausen *et al.*, 2013; Peiffer *et al.*, 2013). A variety of factors may

explain these patterns, particularly abiotic differences (Fierer & Jackson, 2006; Bulgarelli *et al.*,

2013; Nemergut *et al.*, 2013; Vandenkoornhuyse *et al.*, 2015). Soil type appears to have a strong

influence on microbial communities (e.g. Bulgarelli *et al.*, 2012; Lundberg *et al.*, 2012), and is

known to differ broadly across yellow starthistle's range (Hierro *et al.*, 2017). In addition,

populations in California are at the warm and dry extreme of yellow starthistle's climatic niche

(Dlugosch *et al.*, 2015), and our sampling was conducted at the end of a period of severe drought

(Griffin & Anchukaitis, 2014; Diffenbaugh *et al.*, 2015), which could have amplified microbial

differences related to climate (Schrama & Bardgett, 2016). Interestingly, a recent study of

grassland plants found that microbial diversity increased under drought, whereas we found

reduced diversity in the drought-affected range (terHorst *et al.*, 2014).

Importantly, yellow starthistle's invasion could also cause changes to microbial communities.

Species invasionss have been shown to alter microbial composition over short timescales

(Collins *et al.*, 2016; Gibbons *et al.*, 2017), though long term effects are less clear (Lankau,

2011; Day *et al.*, 2015). Yellow starthistle invasions are denser than populations in the native

range by an order of magnitude or more (Uygur *et al.*, 2004; Andonian *et al.*, 2011), and include

lower diversity of plant species overall (Seabloom *et al.*, 2003; Zavaleta & Hulvey, 2004;

D'Antonio *et al.*, 2007). Low plant diversity can depress the diversity of microbes in the

environment and in plants (Garbeva *et al.*, 2004; Schnitzer *et al.*, 2011; Coleman-Derr *et al.*,

2016). Invading plants in California also grow to larger size, and could allow particular strains to

proliferate if size increases were gained at a cost to defenses (Huot *et al.*, 2014; Dlugosch *et al.*,

2015). These possibilities mean that invasions could be a cause rather than an effect of weaker

plant-soil interactions for invasive species in their introduced ranges (Kulmatiski *et al.*, 2008;

Dawson & Schrama, 2016), reinforcing a growing need for explicit observational and

experimental tests that disentangle associations between microbial diversity and plant density,

plant community diversity, and environmental gradients (Dawson & Schrama, 2016).

To date, few studies have compared microbial community compositionl between the native and

introduced ranges of invasive plants. McGinn and colleagues (2016) reported no differences in

the diversity of mutualistic fungal taxa associated with the roots of multiple species of European

Trifolium introduced to New Zealand, despite more favorable soil interactions in the invasions

(McGinn et al., 2018; but see Parker & Gilbert, 2007. Johansen and colleagues (2017) found

increased diversity of fungal communities on the roots of European *Ammophila arenaria*

invading Australia and New Zealand, though there appear to be no differences in interactions

with soil microbial communities in its invasions (in North America; Beckstead & Parker, 2003).

Gundale and colleagues explored the potential contribution of fungal endophyte communities to

more favorable (negative) soil interactions observed in introduced plantations of lodgepole pine

(*Pinus contorta*) from North America (Gundale *et al.*, 2014; 2016). For lodgepole pine,

microbial communities differed among several global regions examined, but there was no

consistent pattern of loss of potential fungal pathogens or gain of mutualists in the introductions,

and it remains unclear what part of the soil community is responsible for observed differences in

interactions across ranges (Gundale *et al.*, 2016). Reinhart and colleagues (2010) focused

specifically on *Phythium* fungal pathogens and quantified their virulence on North American

Prunus serotina introduced to Europe. They found that the most virulent strains occurred only in

the native range, consistent with benefits to invading plants escaping components this specific

pathogenic group. In the only microbiome comparison that included bacterial taxa, Finkel and

colleagues (Finkel *et al.*, 2011; 2016) explored the phyllosphere community of multiple species

of *Tamarix* in native and introduced parts of their ranges, finding that microbial communities

were in general most strongly structured by geographic region.

Our study is the first to find consistent differences in the microbiomes of native and invading

plants which coincide with fitness differences in plant interactions with soil communities.

Among the few invader microbiome studies to date, ours is unusual in focusing on the bacterial

community. Fungi have historically received more attention for their fitness effects on plants, but

bacteria can also play a critical role both as pathogens and mutualists (Haney *et al.*, 2015;

Vandenkoornhuyse *et al.*, 2015; Herrera Paredes & Lebeis, 2016). For yellow starthistle,

previous experiments have demonstrated that fungal communities are not responsible for more

favorable conditions in the invaded range (Hierro *et al.*, 2017), and our findings indicate that

bacterial communities warrant further investigation as the potential source of these differences.

The fitness effects of our specific OTUs are unknown, however, and identifying the bacterial

OTUs that accumulate during interactions with plants would help to elucidate important

pathogenic or mutualistic taxa, and allow field surveys to explicitly test hypotheses that these

strains are lost or gained in the invaded environment.

We have previously argued that yellow starthistle has benefitted from the historical loss of plant

competitors in California (Dlugosch *et al.*, 2015). Disturbance is critical for yellow starthistle

establishment, and functionally similar native species compete well against it in experiments;

however, key competitors have been lost from the ecosystem due toperturbations prior to yellow

starthistle invasion (Zavaleta & Hulvey, 2004; Hooper & Dukes, 2010; Hierro *et al.*, 2011, 2017;

Hulvey & Zavaleta, 2012). Any benefits of altered bacterial communities could be independent

of competition with native plant species, but these factors might also interact. Increased density

due to a lack of competition could have reduced plant-associated microbial diversity, as

described above. Yellow starthistle experiences negative plant-soil feedbacks across generations

(Andonian *et al.*, 2011), however, and the build up of high plant densities is therefore unlikely to

generate more favorable soil interactions. Alternatively, the historical loss of native species

diversity in California (D'Antonio *et al.*, 2007) could have resulted in the loss of associated

microbial diversity (Garbeva *et al.*, 2004; Schnitzer *et al.*, 2011; Coleman-Derr *et al.*, 2016),

generating particularly strong opportunities for invasion into a system with both reduced plant

competition and reduced pathogen diversity. Microbial surveys of remnant native communities,

as well as across densities of yellow starthistle would facilitate tests of alternative hypotheses for

interacting effects of plant and microbial diversity, and it may be informative to explore

microbial communities preserved on native plant specimens pre-dating the extensive invasion of

yellow starthistle into this region.

Conclusions

We found consistent differences between native and invading yellow starthistle plants in their

bacterial microbiomes. These differences were robust to additional variation associated with

plant compartment and the diversity of plant genotypes. Invaded range microbiomes differed in

composition across major taxonomic groups, and harbored a lower diversity of bacteria,

including reduced evenness on the surface of leaves and roots and reduced richness of root

endophytes. We suggest that bacteria could be the source of more favorable microbial

interactions that have been observed in this invasion. Our findings also raise questions about 1)

whether lower bacterial diversity is a feature of the invaded environment or whether it is caused

by the invasion itself, and 2) how differences in the microbial community might have interacted

with other changes in plant competitive interactions and the abiotic environment to affect plant

fitness. These questions highlight the need for additional studies that compare microbial

communities (including bacteria) associated with native and invading populations, that couple

microbial community identification with plant-soil feedback and fitness experiments, and that

examine the interaction of environment, plant diversity, and plant density on microbial

communities and their fitness effects on plants.

ACKNOWLEDGMENTS

We thank D. Lundberg and S. Lebeis for helpful discussions regarding sampling design; G.

Reardon and C. Chandeyson for assistance with field collections; E. Arnold, J. Aspinwall, J.

Braasch, E. Carlson, K. Hockett, P. Humphries, T. O'Connor, M. Schneider, J. U'Ren, and N.

Zimmerman for 16S library preparation assistance and discussion; A. Gerritsen, D. New and

staff at iBEST for assistance with sequencing; K. Andonian, J. Hierro, and 3 reviewers for

helpful feedback on the manuscript. We are particularly indebted to C.E. Morris for hosting the

European sample processing at INRA Station de Pathologie Végétale, Montfavet, France. Data

collection and analyses performed by the IBEST Genomics Resources Core at the University of

Idaho were supported in part by NIH COBRE grant P30GM103324. This study was supported

by USDA grant 2015-67013-23000 to K.M.D., D.A.B., and S.M.S.

AUTHOR CONTRIBUTIONS

P.L-I., D.A.B., and K.M.D. designed the study. P.L-I. and J.H. collected the samples with

assistance from H.S., S.M.S., and S.R.W. P.L-I conducted the microbial sequencing and

bioinformatics. P.L-I, S.R.W., and K.M.D. analyzed the data. P.L-I and K.M.D. wrote the

manuscript, which was edited by all authors.

REFERENCES

Agrawal AA. **2007**. Macroevolution of plant defense strategies. *Trends in Ecology & Evolution*

22: 103–109.

Agrawal AA, Kotanen PM, Mitchell CE, Power AG, Godsoe W, Klironomos J. **2005**.

Enemy release? an experiment with congeneric plant pairs and diverse above- and

belowground enemies. *Ecology* **86**: 2979–2989.

Andonian K, Hierro JL. **2011**. Species interactions contribute to the success of a global plant

invader. *Biological Invasions* **13**: 2957–2965.

Andonian K, Hierro JL, Khetsuriani L, Becerra PI, Janoyan G, Villareal D, Cavieres LA,

Fox LR, Callaway RM. **2012**. Geographic mosaics of plant-soil microbe interactions in a

global plant invasion. *Journal of Biogeography* **39**: 600–608.

Andonian K, Hierro JL, Khetsuriani L, Becerra P, Janoyan G, Villarreal D, Cavieres L,

Fox LR, Callaway RM. **2011**. Range-expanding populations of a globally introduced weed

experience negative plant-soil feedbacks. *PLoS One* **6**: e20117.

Barker BS, Andonian K, Swope SM, Luster DG, Dlugosch KM. **2017**. Population genomic

analyses reveal a history of range expansion and trait evolution across the native and invaded

range of yellow starthistle (*Centaurea solstitialis*). *Molecular Ecology* **26**: 1131–1147.

Beckstead J, Parker IM. **2003**. Invasiveness of *Ammophila arenaria*: release from soil-borne

pathogens? *Ecology* **84**: 2824–2831.

Bever JD. **2003**. Soil community feedback and the coexistence of competitors: conceptual

frameworks and empirical tests. *The New Phytologist* **157**: 465–473.

Bodenhausen N, Bortfeld-Miller M, Ackermann M, Vorholt JA. **2014**. A synthetic

community approach reveals plant genotypes affecting the phyllosphere microbiota. *PLoS Genetics* **10**: e1004283.

Bodenhausen N, Horton MW, Bergelson J. **2013**. Bacterial communities associated with the leaves and the roots of *Arabidopsis thaliana*. *PLoS One* **8**: e56329.

Bulgarelli D, Garrido-Oter R, Münch PC, Weiman A, Dröge J, Pan Y, McHardy AC, Schulze-Lefert P. **2015**. Structure and function of the bacterial root microbiota in wild and domesticated barley. *Cell Host & Microbe* **17**: 392–403.

Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, Ahmadinejad N, Assenza F, Rauf P, Huettel B, Reinhardt R, Schmelzer E, *et al.* 2012. Revealing structure and assembly cues for *Arabidopsis* root-inhabiting bacterial microbiota. *Nature* **488**: 91–95.

Bulgarelli D, Schlaeppi K, Spaepen S, Ver Loren van Themaat E, Schulze-Lefert P. **2013**. Structure and functions of the bacterial microbiota of plants. *Annual Review of Plant Biology* **64**: 807–838.

Butchart SHM, Walpole M, Collen B, van Strien A, Scharlemann JPW, Almond REA, Baillie JEM, Bomhard B, Brown C, Bruno J, *et al.* 2010. Global biodiversity: indicators of recent declines. *Science* **328**: 1164–1168.

Callaway RM, Thelen GC, Rodriguez A, Holben WE. **2004**. Soil biota and exotic plant invasion. *Nature* **427**: 731–733.

Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña AG, Goodrich JK, Gordon JI, *et al.* 2010. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* **7**: 335–336.

Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, Fierer N, Knight R. **2011**. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proceedings of the National Academy of Sciences* **108 Suppl 1**: 4516–4522.

Colautti RI, Lau JA. **2015**. Contemporary evolution during invasion: evidence for differentiation, natural selection, and local adaptation. *Molecular Ecology* **24:** 1999-2017.

Colautti RI, Ricciardi A, Grigorovich IA, MacIsaac HJ. **2004**. Is invasion success explained by the enemy release hypothesis? *Ecology Letters* **7**: 721–733.

Coleman-Derr D, Desgarennes D, Fonseca-Garcia C, Gross S, Clingenpeel S, Woyke T, North G, Visel A, Partida-Martinez LP, Tringe SG. **2016**. Plant compartment and

biogeography affect microbiome composition in cultivated and native *Agave* species. *The*

New Phytologist **209**: 798–811.

Collins CG, Carey CJ, Aronson EL, Kopp CW, Diez JM. **2016**. Direct and indirect effects of

native range expansion on soil microbial community structure and function. *The Journal of*

Ecology **104**: 1271–1283.

D'Antonio CM, Malmstrom C, Reynolds SA, Gerlach J. **2007**. Ecology of invasive non-

native species in California grassland. In: Stromberg MR, Corbin JD, D'Antonio CM, eds.

California grassland: ecology and management. Berkeley, CA, USA: University of

California Press, 67–86.

Dawson W, Schrama M. **2016**. Identifying the role of soil microbes in plant invasions. *The*

Journal of Ecology **104**: 1211–1218.

Day NJ, Dunfield KE, Antunes PM. **2015**. Temporal dynamics of plant-soil feedback and root-

associated fungal communities over 100 years of invasion by a non-native plant. *The Journal*

of Ecology **103**: 1557–1569.

Diffenbaugh NS, Swain DL, Touma D. **2015**. Anthropogenic warming has increased drought

risk in California. *Proceedings of the National Academy of Sciences* **112**: 3931–3936.

DiTomaso JM, Healy EA. **2007**. *Weeds of California and other western states*. Oakland, CA,

USA: University of California Department of Agriculture and Natural Resources.

Dlugosch KM, Cang FA, Barker BS, Andonian K, Swope SM, Rieseberg LH. **2015**.

Evolution of invasiveness through increased resource use in a vacant niche. *Nature Plants* **1**:

15066.

Dlugosch KM, Parker IM. **2008**. Founding events in species invasions: genetic variation,

adaptive evolution, and the role of multiple introductions. *Molecular Ecology* **17**: 431–449.

Edgar RC. **2010**. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*

26: 2460–2461.

Edgar RC. **2013**. UPARSE: highly accurate OTU sequences from microbial amplicon reads.

Nature Methods **10**: 996–998.

Edgar RC, Flyvbjerg H. **2015**. Error filtering, pair assembly and error correction for next-

generation sequencing reads. *Bioinformatics* **31**: 3476–3482.

Ellis EC, Antill EC, Kreft H. **2012**. All is not loss: plant biodiversity in the anthropocene. *PLoS*

One **7**: e30535.

Engelkes T, Morriën E, Verhoeven KJF, Bezemer TM, Biere A, Harvey JA, McIntyre LM,
Tamis WLM, van der Putten WH. **2008**. Successful range-expanding plants experience
less above-ground and below-ground enemy impact. *Nature* **456**: 946–948.

Eriksen RL, Desronvil T, Hierro JL, Kesseli R. **2012**. Morphological differentiation in a
common garden experiment among native and non-native specimens of the invasive weed
yellow starthistle (*Centaurea solstitialis*). *Biological Invasions* **14**: 1459–1467.

Essl F, Dullinger S, Rabitsch W, Hulme PE, Hülber K, Jarošík V, Kleinbauer I,
Krausmann F, Kühn I, Nentwig W, *et al.* **2011**. Socioeconomic legacy yields an invasion
debt. *Proceedings of the National Academy of Sciences* **108**: 203–207.

Excoffier L, Foll M, Petit RJ. **2009**. Genetic consequences of range expansions. *Annual Review*
of Ecology, Evolution, and Systematics **40**: 481–501.

Faillace CA, Lorusso NS, Duffy S. **2017**. Overlooking the smallest matter: viruses impact
biological invasions. *Ecology Letters* **EarlyView**.

Felker-Quinn E, Schweitzer JA, Bailey JK. **2013**. Meta-analysis reveals evolution in invasive
plant species but little support for Evolution of Increased Competitive Ability (EICA).
Ecology and Evolution **3**: 739–751.

Fierer N, Jackson JA, Vilgalys R, Jackson RB. **2005**. Assessment of soil microbial community
structure by use of taxon-specific quantitative PCR assays. *Applied and Environmental*
Microbiology **71**: 4117–4120.

Fierer N, Jackson RB. **2006**. The diversity and biogeography of soil bacterial communities.
Proceedings of the National Academy of Sciences **103**: 626–631.

Finkel OM, Burch AY, Lindow SE, Post AF, Belkin S. **2011**. Geographical location
determines the population structure in phyllosphere microbial communities of a salt-
excreting desert tree. *Applied and Environmental Microbiology* **77**: 7647–7655.

Finkel OM, Delmont TO, Post AF, Belkin S. **2016**. Metagenomic signatures of bacterial
adaptation to life in the phyllosphere of a salt-secreting desert tree. *Applied and*
Environmental Microbiology **82**: 2854–2861.

Fitzpatrick C; Lu-Irving P; Copeland J; Guttman DS; Wang PW; Baltrus DA; Dlugosch
KM; Johnson MTJ. Chloroplast sequence variation and the efficacy of peptide nucleic acids
for blocking host amplification in plant microbiome studies. *Microbiome* **In minor revision**

Fonseca-García C, Coleman-Derr D, Garrido E, Visel A, Tringe SG, Partida-Martínez LP.

2016. The cacti microbiome: interplay between habitat-filtering and host-specificity. *Frontiers in Microbiology* **7**: 150.

Garbeva P, van Veen JA, van Elsas JD. **2004**. Microbial diversity in soil: selection microbial populations by plant and soil type and implications for disease suppressiveness. *Annual Review of Phytopathology* **42**: 243–270.

Gerlach JD. **1997**. How the West was lost: reconstructing the invasion dynamics of yellow starthistle and other plant invaders of western rangelands and natural areas. *California Exotic Pest Plant Council Symposium Proceedings* **3**: 67–72.

Gibbons SM, Lekberg Y, Mummey DL, Sangwan N, Ramsey PW, Gilbert JA, Shade A. **2017**. Invasive plants rapidly reshape soil properties in a grassland ecosystem. *mSystems* **2**: e00178–16.

Griffin D, Anchukaitis KJ. **2014**. How unusual is the 2012–2014 California drought? *Geophysical Research Letters* **41**: 2014GL062433.

Gundale MJ, Almeida JP, Wallander H, Wardle DA, Kardol P, Nilsson M-C, Fajardo A, Pauchard A, Peltzer DA, Ruotsalainen S, *et al.* 2016. Differences in endophyte communities of introduced trees depend on the phylogenetic relatedness of the receiving forest. *The Journal of Ecology* **104**: 1219–1232.

Gundale MJ, Kardol P, Nilsson M-C, Nilsson U, Lucas RW, Wardle DA. **2014**. Interactions with soil biota shift from negative to positive when a tree species is moved outside its native range. *The New Phytologist* **202**: 415–421.

Haney CH, Samuel BS, Bush J, Ausubel FM. **2015**. Associations with rhizosphere bacteria can confer an adaptive advantage to plants. *Nature Plants* **1**.

Herrera Paredes S, Lebeis SL. **2016**. Giving back to the community: microbial mechanisms of plant–soil interactions. *Functional Ecology* **30**: 1043–1052.

Hierro JL, Khetsuriani L, Andonian K, Eren Ö, Villarreal D, Janoian G, Reinhart KO, Callaway RM. **2017**. The importance of factors controlling species abundance and distribution varies in native and non-native ranges. *Ecography* **40:** 991-1002.

Hierro JL, Lortie CJ, Villarreal D, Estanga-Mollica ME, Callaway RM. **2011**. Resistance to *Centaurea solstitialis* invasion from annual and perennial grasses in California and Argentina. *Biological Invasions* **13**: 2249–2259.

Hill MO. **1973**. Diversity and evenness: a unifying notation and its consequences. *Ecology* **54**:

731   427–432.

Hooper DU, Dukes JS. **2010**. Functional composition controls invasion success in a California serpentine grassland. *The Journal of Ecology* **98**: 764–777.

Hulvey KB, Zavaleta ES. **2012**. Abundance declines of a native forb have nonlinear impacts on grassland invasion resistance. *Ecology* **93**: 378–388.

Huot B, Yao J, Montgomery BL, He SY. **2014**. Growth–defense tradeoffs in plants: a balancing act to optimize fitness. *Molecular Plant* **7**: 1267–1287.

Hsieh TC, Ma KH, Chao A. **2016**. iNEXT: an R package for rarefaction and extrapolation of species diversity (Hill numbers). *Methods in Ecology and Evolution* **7**: 1451–1456.

Johansen RB, Johnston P, Mieczkowski P, Perry GLW, Robeson MS, Vilgalys R, Burns BR. **2017**. Scattered far and wide: A broadly distributed temperate dune grass finds familiar fungal root associates in its invasive range. *Soil Biology & Biochemistry* **112**: 177–190.

Keane R, Crawley MJ. **2002**. Exotic plant invasions and the enemy release hypothesis. *Trends in Ecology & Evolution* **17**: 164–170.

Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. **2013**. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Applied and Environmental Microbiology* **79**: 5112–5120.

Kulmatiski A, Beard KH, Stevens JR, Cobbold SM. **2008**. Plant–soil feedbacks: a meta-analytical review. *Ecology Letters* **11**: 980–992.

Lankau RA. **2011**. Resistance and recovery of soil microbial communities in the face of *Alliaria petiolata* invasions. *The New Phytologist* **189**: 536–548.

Lankau RA, Keymer DP. **2018**. Simultaneous adaptation and maladaptation of tree populations to local rhizosphere microbial communities at different taxonomic scales. *The New Phytologist* **217**: 1267–1278.

Lebeis SL, Paredes SH, Lundberg DS, Breakfield N, Gehring J, McDonald M, Malfatti S, Glavina del Rio T, Jones CD, Tringe SG, *et al.* 2015. Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa. *Science* **349**: 860–864.

Leff JW, Lynch RC, Kane NC, Fierer N. **2016**. Plant domestication and the assembly of bacterial and fungal communities associated with strains of the common sunflower, *Helianthus annuus*. *The New Phytologist* **214:** 412–423.

Legendre P, Gallagher ED. **2001**. Ecologically meaningful transformations for ordination of
species data. *Oecologia* **129**: 271–280.

Legendre P, Legendre L. **1998**. *Numerical Ecology*. Amsterdam: Elsevier Science.

Levine JM, Adler PB, Yelenik SG. **2004**. A meta-analysis of biotic resistance to exotic plant
invasions. *Ecology Letters* **7**: 975–989.

Lonsdale WM. **1999**. Global patterns of plant invasions and the concept of invasibility. *Ecology*
80: 1522–1536.

Louca S, Jacques SMS, Pires APF, Leal JS, Srivastava DS, Parfrey LW, Farjalla VF,
Doebeli M. **2016**. High taxonomic variability despite stable functional structure across
microbial communities. *Nature Ecology & Evolution* **1**: s41559–016–0015.

Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, Tremblay J,
Engelbrektson A, Kunin V, del Rio TG, *et al.* **2012**. Defining the core *Arabidopsis*
thaliana root microbiome. *Nature* **488**: 86–90.

Lundberg DS, Yourstone S, Mieczkowski P, Jones CD, Dangl JL. **2013**. Practical innovations
for high-throughput amplicon sequencing. *Nature Methods* **10**: 999–1002.

Maron JL, Klironomos J, Waller L, Callaway RM. **2014**. Invasive plants escape from
suppressive soil biota at regional scales. *The Journal of Ecology* **102**: 19–27.

Maron JL, Vilà M. **2001**. When do herbivores affect plant invasion? Evidence for the natural
enemies and biotic resistance hypotheses. *Oikos* **95**: 361–373.

McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A, Andersen GL,
Knight R, Hugenholtz P. **2012**. An improved Greengenes taxonomy with explicit ranks for
ecological and evolutionary analyses of bacteria and archaea. *The ISME Journal* **6**: 610–618.

McGinn KJ, van der Putten WH, Duncan RP, Shelby N, Weser C, Hulme PE. **2016**.
Trifolium species associate with a similar richness of soil-borne mutualists in their introduced
and native ranges. *Journal of Biogeography* **43**: 944–954.

McGinn KJ, van der Putten WH, Hulme PE, Shelby N, Weser C, Duncan RP. **2018**. The
influence of residence time and geographic extent on the strength of plant-soil feedbacks for
naturalised *Trifolium*. *The Journal of Ecology* **106**: 207–217.

McQuitty LL. **1966**. Similarity analysis by reciprocal pairs for discrete and continuous data.
Educational and Psychological Measurement **26**: 825–831.

Mitchell CE, Agrawal AA, Bever JD, Gilbert GS, Hufbauer RA, Klironomos JN, Maron

JL, Morris WF, Parker IM, Power AG, *et al.* 2006. Biotic interactions and plant invasions. *Ecology Letters* **9**: 726–740.

Nemergut DR, Schmidt SK, Fukami T, O'Neill SP, Bilinski TM, Stanish LF, Knelman JE, Darcy JL, Lynch RC, Wickey P, *et al.* 2013. Patterns and processes of microbial community assembly. *Microbiology and Molecular Biology Reviews* **77**: 342–356.

Newton AC, Fitt BDL, Atkins SD, Walters DR, Daniell TJ. 2010. Pathogenesis, parasitism and mutualism in the trophic space of microbe–plant interactions. *Trends in Microbiology* **18**: 365–373.

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB, Simpson GL, Solymos P, *et al.* 2016. vegan: community ecology package. R package version 2.4-3.

Parker IM, Gilbert GS. 2007. When there is no escape: the effects of natural enemies on native, invasive, and noninvasive plants. *Ecology* **88**: 1210–1224.

Pearson DE, Ortega YK, Eren Ö, Hierro JL. 2018. Community assembly theory as a framework for biological invasions. *Trends in Ecology & Evolution* **33**: 313–325.

Peiffer JA, Spor A, Koren O, Jin Z, Tringe SG, Dangl JL, Buckler ES, Ley RE. 2013. Diversity and heritability of the maize rhizosphere microbiome under field conditions. *Proceedings of the National Academy of Sciences* **110**: 6548–6553.

Petermann JS, Fergus AJF, Turnbull LA, Schmid B. 2008. Janzen-Connell effects are widespread and strong enough to maintain diversity in grasslands. *Ecology* **89**: 2399–2406.

Pielou EC. 1966. The measurement of diversity in different types of biological collections. *Journal of Theoretical Biology* **13**: 131–144.

Pitcairn M, Schoenig S, Yacoub R, Gendron J. 2006. Yellow starthistle continues its spread in California. *California Agriculture* **60**: 83–90.

R Core Team. 2015. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.

Ramette A. 2007. Multivariate analyses in microbial ecology. *FEMS Microbiology Ecology* **62**: 142–160.

Reinhart KO, Callaway RM. 2006. Soil biota and invasive plants. *The New Phytologist* **170**: 445–457.

Reinhart KO, Packer A, Van der Putten WH, Clay K. 2003. Plant–soil biota interactions and

824 spatial distribution of black cherry in its native and invasive ranges. *Ecology Letters* **6**: 1046–
825 1050.

826 **Reinhart KO, Tytgat T, Van der Putten WH, Clay K**. **2010**. Virulence of soil-borne
827 pathogens and invasion by *Prunus serotina*. *The New Phytologist* **186**: 484–495.

828 **Rout ME, Callaway RM**. **2012**. Interactions between exotic invasive plants and soil microbes in
829 the rhizosphere suggest that 'everything is not everywhere'. *Annals of Botany* **110**: 213–222.

830 **Schnitzer SA, Klironomos JN, HilleRisLambers J, Kinkel LL, Reich PB, Xiao K, Rillig**
831 **MC, Sikes BA, Callaway RM, Mangan SA,** *et al.* **2011**. Soil microbes drive the classic
832 plant diversity–productivity pattern. *Ecology* **92**: 296–303.

833 **Schrama M, Bardgett RD**. **2016**. Grassland invasibility varies with drought effects on soil
834 functioning. *The Journal of Ecology* **104**: 1250–1258.

835 **Seabloom EW, Harpole WS, Reichman OJ, Tilman D**. **2003**. Invasion, competitive
836 dominance, and resource use by exotic and native California grassland species. *Proceedings*
837 *of the National Academy of Sciences* **100**: 13384–13389.

838 **Shannon CE**. **1948**. A mathematical theory of communication. *Bell System Technical Journal*
839 **27**: 379–423.

840 **Smakowska E, Kong J, Busch W, Belkhadir Y**. **2016**. Organ-specific regulation of growth-
841 defense tradeoffs by plants. *Current Opinion in Plant Biology* **29**: 129–137.

842 **terHorst CP, Lennon JT, Lau JA**. **2014**. The relative importance of rapid evolution for plant-
843 microbe interactions depends on ecological context. *Proceedings of the Royal Society:*
844 *Biological Sciences* **281**: 20140028.

845 **terHorst CP, Zee PC**. **2016**. Eco-evolutionary dynamics in plant–soil feedbacks. *Functional*
846 *Ecology* **30**: 1062–1072.

847 **Torchin ME, Mitchell CE**. **2004**. Parasites, pathogens, and invasions by plants and animals.
848 *Frontiers in Ecology and the Environment* **2**: 183–190.

849 **Uller T, Leimu R**. **2011**. Founder events predict changes in genetic diversity during human-
850 mediated range expansions. *Global Change Biology* **17**: 3478–3485.

851 **Uygur S, Smith L, Uygur FN, Cristofaro M, Balciunas J**. **2004**. Population densities of yellow
852 starthistle (*Centaurea solstitialis*) in Turkey. *Weed Science* **52**: 746–753.

853 **Vandenkoornhuyse P, Quaiser A, Duhamel M, Le Van A, Dufresne A**. **2015**. The importance
854 of the microbiome of the plant holobiont. *The New Phytologist* **206**: 1196–1206.

van der Putten WH, Bardgett RD, Bever JD, Bezemer TM, Casper BB, Fukami T, Kardol

P, Klironomos JN, Kulmatiski A, Schweitzer JA, *et al.* **2013**. Plant–soil feedbacks: the

past, the present and future challenges. *The Journal of Ecology* **101**: 265–276.

van der Putten WH, Bradford MA, Pernilla Brinkman E, van de Voorde TFJ, Veen GF.

2016. Where, when and how plant–soil feedback matters in a changing world. *Functional*

Ecology **30**: 1109–1121.

Venables WN, Ripley BD. **2002**. *Statistics and computing: modern applied statistics with S*.

New York, NY, USA: Springer.

Wagg C, Bender SF, Widmer F, van der Heijden MGA. **2014**. Soil biodiversity and soil

community composition determine ecosystem multifunctionality. *Proceedings of the*

National Academy of Sciences **111**: 5266–5270.

Wickham H. **2009**. *ggplot2: elegant graphics for data analysis*. New York: Springer-Verlag.

Widmer TL, Guermache F, Dolgovskaia MY, Reznik SY. **2007**. Enhanced growth and seed

properties in introduced vs. native populations of yellow starthistle (*Centaurea solstitialis*).

Weed Science **55**: 465–473.

Zavaleta ES, Hulvey KB. **2004**. Realistic species losses disproportionately reduce grassland

resistance to biological invaders. *Science* **306**: 1175–1177.

SUPPORTING INFORMATION

Fig. S1. Frequency distributions of readcounts per sample.

Fig. S2. NMDS plot of OTU composition across all sample types.

Fig. S3. Clustering dendrograms for bulk samples from three plant compartments.

Fig. S4. Heat maps showing relative abundance of top 25 OTUs per bulk sample from three plant

compartments.

Table S1. Collecting information for populations sampled in this study.

Table S2. Extent of host chloroplast amplification with different PNA clamps.

Table S3. Axis loadings from Principal Component Analysis of three plant compartments.

FIGURE LEGENDS

Fig. 1. The distribution (gray) of yellow starthistle and sampling sites (circles) for this study.

Maps detail the native range in Eurasia (a) and the invasion of western North America (b).

Previous work has indicated that western Europe is the source for the severe invasion of

California, USA (both in dark shading; Barker et al. 2017). Sampling included seven locations in

California (b, filled circles), six locations in western Europe and an additional two locations in

eastern Europe (a, open circles).

Fig 2. NMDS plots of bacterial OTU composition in phyllosphere (green), endorhizosphere

(light blue), and ectorhizosphere (dark blue) samples from native (open symbols) and invaded

(closed symbols) ranges. Plotted are a) pooled samples for each sampling location, showing

overall separation by range within compartment (stress = 0.14), and b) whole root samples for

individual plants within native and invading populations (stress = 0.16). Ellipses indicate 95%

confidence intervals for samples grouped by range (native range: dashed lines; invaded range:

solid lines).

Fig 3. Relative abundance of (proportion of reads mapping to) phyla in yellow starthistle

phyllosphere, ectorhizosphere, and endorhizosphere samples from native and invaded ranges.

Fig 4. Distributions of OTU (a) richness, (b) evenness, and (c,d) diversity ($e^{H'}$) among samples

(pooled plants) from each location in the native and invaded ranges for phyllosphere,

ectorhizosphere, and endorhizosphere compartments. (a-c) show values for all OTUs and (d)

shows values based on OTUs from known pathogen-containing genera. Significant differences

from Kruskal-Wallis tests are indicated with asterisks: $*P < 0.05,$ $^{(*)}P < 0.1.$

Fig 5. Distributions of endorhizosphere OTU (a) richness, (b) evenness, and (c,d) diversity ($e^{H'}$)

among individual plants at each location in the native and invaded ranges. (a-c) show values for

all OTUs and (d) shows values based on OTUs from known pathogen-containing genera.

Significant differences from Kruskal-Wallis tests are indicated with asterisks: $*P < 0.05,$ $^{(*)}P <$

0.1.

Fig 6. Bacterial diversity ($e^{H'}$) in endorhizosphere samples pooled by sampling location, as a

function of the genetic diversity among plants at the same sites (calculated as the average

proportion of pairwise nucleotide differences between alleles (π) at variable positions in the

genome; from Barker *et al.*, 2017). Lines show significant positive relationships (linear model: P

= 0.02) between microbial and plant diversity at sampling locations in both the native range

(open symbols, dashed line) and the invaded range (closed symbols, solid line).

Fig. 1 The distribution (gray) of yellow starthistle and sampling sites (circles) for this study. Maps detail the native range in Eurasia (a) and the invasion of western North America (b). Previous work has indicated that western Europe is the source for the severe invasion of California, USA (both in dark shading; Barker et al. 2017). Sampling included seven locations in California (b, filled circles), six locations in western Europe and an additional two locations in eastern Europe (a, open circles).

Fig 2. NMDS plots of bacterial OTU composition in phyllosphere (green), endorhizosphere (light blue), and ectorhizosphere (dark blue) samples from native (open symbols) and invaded (closed symbols) ranges. Plotted are a) pooled samples for each sampling location, showing overall separation by range within compartment (stress = 0.14), and b) whole root samples for individual plants within native and invading populations (stress = 0.16). Ellipses indicate 95% confidence intervals for samples grouped by range (native range: dashed lines; invaded range: solid lines).

Fig 3. Relative abundance of (proportion of reads mapping to) phyla in yellow starthistle phyllosphere, ectorhizosphere, and endorhizosphere samples from native and invaded ranges.

Fig 4. Distributions of OTU (a) richness, (b) evenness, and (c,d) diversity ($e^{H'}$) among samples (pooled plants) from each location in the native and invaded ranges for phyllosphere, ectorhizosphere, and endorhizosphere compartments. (a-c) show values for all OTUs and (d) shows values based on OTUs from known pathogen-containing genera. Significant differences from Kruskal-Wallis tests are indicated with asterisks: *$P < 0.05$, (*)$P < 0.1$.

Fig 5: Distributions of endorhizosphere OTU (a) richness, (b) evenness, and (c,d) diversity ($e^{H'}$) among individual plants at each location in the native and invaded ranges. (a-c) show values for all OTUs and (d) shows values based on OTUs from known pathogen-containing genera. Significant differences from Kruskal-Wallis tests are indicated with asterisks: *P < 0.05, (*)P < 0.1.

Fig 6. Bacterial diversity ($e^{H'}$) in endorhizosphere samples pooled by sampling location, as a function of the genetic diversity among plants at the same sites (calculated as the average proportion of pairwise nucleotide differences between alleles ( ) at variable positions in the genome; from Barker *et al.*, 2017). Lines show significant positive relationships (linear model: *P* = 0.02) between microbial and plant diversity at sampling locations in both the native range (open symbols, dashed line) and the invaded range (closed symbols, solid line).

Supporting Information

Fig. S1 Frequency distributions of un-rarefied read counts for samples from all four plant compartments (native and invading population samples combined), as well as control (blank) samples. Vertical red lines indicate thresholds for rarefaction where relevant.

Fig. S2 NMDS plot of bacterial OTU composition in unrarefied control, phyllosphere, leaf endosphere, ectorhizosphere, and endorhizosphere samples (stress = 0.10).

Fig. S3 Yellow starthistle populations clustered according to Bray-Curtis dissimilarity between rarefied, square-root transformed pooled samples from three plant compartments.

Fig. S4 Heat maps showing the fraction of reads mapping to each of the top 25 OTUs in rarefied bulk samples from three plant compartments. The top five OTUs contributing to the first principal component separating native and invaded samples in each compartment are indicated with an asterisk. Taxonomic assignments for each OTU are given to the lowest taxonomic rank identified.

Fig. S5 Proportion of rarefied read counts assigned to pathogen-containing genera in all samples according to plant compartment and range.

Table S1 Collecting information for populations sampled in this study, including designation codes, localities, coordinates, sampling dates, and specimen accession numbers of plant collections at ARIZ.

Code	Lat.	Long.	Locality	Date	ARIZ specimen numbers
SAL	N40.99112	W005.65831	Avenida de la Merced, ~1.5km N of Salamanca. Dirt path between road and field, on E side of road.	22-Jun-15	426137, 426032, 425636, 426125, 426034, 426033, 426035, 426126, 426140, 426068, 426127, 426128, 426129, 426130, 426131, 426132, 426133, 426134, 426135, 426136, 426138, 425635, 426139
CAN	N41.00085	W004.89715	Junction AV800 & road to Canales. On E side of Canales road, in the narrow strip between road and field edge.	22-Jun-15	425643, 426065, 426064, 426063, 426055, 426054, 425646, 426067, 426053, 426052, 425642, 426066, 426051, 425641, 426062, 426061, 426059, 426058, 426060, 426057, 426056
GRA	N37.26843	W003.66488	GR300 ~100m from junction w 213, ~20m up dirt path to the E side of the road. Weedy waste area near edge of olive orchard.	23-Jun-15	425716, 425715, 425714, 425713, 425739, 425738, 425737, 425736, 425735, 425734, 425733, 425732, 425728, 425740, 425729, 425711, 425702, 425703, 425704, 425705, 425706, 425707, 425708, 425709, 425710
SAZ	N39.83529	W002.50997	Gravel road by W side of CM3118 (road to Villares del Saz), just next to underpass under A3. Dirt farming path (for tractors) on S side of road, between sunflower fields.	24-Jun-15	425678, 425741, 425679, 425731, 425680, 425681, 425682, 425683, 425684, 425685, 425757, 425755, 425753, 425751, 425749, 425746, 425744, 425743, 425758, 425756, 425752, 425750, 425748, 425747, 425745, 425742
CUE	N40.12939	W002.13880	CM2105 between Cuenca & Tragacete ~8km W of Cuenca. Gravel road to the N of 2105, between field and river.	24-Jun-15	425424, 425421, 425420, 425423, 425425, 425426
CAZ	N43.74958	E003.77119	D113, ~1km from junction w 986, near Cazevielle. Flat, gravelly turnout & intersection w unpaved road, & gravelly roadsides.	20-Jun-15	425691, 425690, 425971, 425688, 425689, 425972, 425692, 425970, 425693
HU01	N47.18240	E18.08928	Unnamed dirt roads about 200m NW of E66/8, running parallel. Access by perpendicular unnamed paved road.	7-Jul-15	425655, 425660, 425661, 425653, 425652, 425658, 425657, 425656, 425654, 425631, 425687, 425634, 425633, 425686, 425630, 425628, 425629, 425627, 425662

HU29	N47.31785	E21.03385	Margin of irrigation ditch bordered by hay and sunflower fields on one side and service road parallel to main road (E60/4) on the other side. Site to the south of E60/4.	8-Jul-15	425730, 425677, 425676, 425640, 425675, 425674, 425673, 425726, 425725, 425724, 425723, 425722, 425721, 425720, 425719, 425718, 425717, 425727
DIA	N37.86275	W121.98003	Hillside to the West of parking lot at the end of Green Valley Road.	8-Jun-15	425394, 425400, 425401, 425402, 425403, 425404, 425405, 425406, 425407, 425408, 425409, 425115, 425381, 425410, 425382, 425383, 425384, 425385, 425386, 425387, 425388, 425389, 425390, 425391, 425392
GIL	N37.03389	W121.53611	Disturbed grassy area adjacent to turn out on South side of Leavesley Rd. near intersection with New Ave. Grassy patch between road and tilled field.	7-Jun-15	425374, 425373, 425380, 425379, 425378, 425377, 425376, 425375, 425124, 425123, 425122, 425121, 425119, 425120, 425118, 425114, 425113, 425399, 425112, 425395, 425111, 425398, 425397, 425396
RB	N40.27085	W122.27103	Lightly grassy patch to north-west of rest stop (Herbert S. Miles rest area) between Cottonwood and Red Bluff on southbound I-5.	9-Jun-15	425372, 425371, 425370, 425369, 425368, 425418, 425365, 425415, 425367, 425366, 425416, 425417, 425419
CLV	N36.91603	W119.79341	Gully on eastern side of the Yosemite fwy (41) adjacent to olive orchards.	6-Jun-15	426046, 426045, 426044, 426043, 426042, 426036, 426037, 425637, 426038, 426039, 426040, 426041, 425639, 426026, 426027, 426028, 426029, 426030, 426031, 425638
LEB	N34.82844	W118.87369	Cal trans maintained, weedy area adjacent to SW of rest stop off of south bound I-5.	5-Jun-15	425968, 425695, 425969, 425966, 425967, 426047, 425694, 426050, 426049, 426048
TRI	N37.46159	W119.79399	Strip of grassland along south side of Hwy 49 at intersection with Triangle road. Strip between road and fenced off rangeland.	11-Jun-15	425116, 425117, 425414, 425412, 425413, 425411, 425393
SIE	N38.78162	W120.41639	Site upslope of Weber Mill rd. about ¼ mile from the turnoff from Ice House road.	10-Jun-15	425648, 425647, 425649, 425644, 425645

Table S2 Comparison of host chloroplast blocking levels achieved by different PNA clamps in pooled endorhizosphere samples from two populations. Total read counts, number of reads mapping to host chloroplast, and host chloroplast reads as percentage of total reads are reported for each sample.

Sample	plastid PNA	total	chloroplast	% chloroplast
CAN pooled endorhizosphere	Lundberg *et al.* 2013 (GGCTCAACCCTGGACAG)	14,763	2,873	19
CAN pooled endorhizosphere	Asteraceae-specific (GGCTCAACTCTGGACAG)	12,579	91	1
CUE pooled endorhizosphere	Lundberg *et al.* 2013 (GGCTCAACCCTGGACAG)	11,962	6,318	53
CUE pooled endorhizosphere	Asteraceae-specific (GGCTCAACTCTGGACAG)	14,608	314	2

Table S3 Loading by top 10 individual OTUs along first principal component axis from analysis of Hellinger transformed data matrices for three plant compartments. Column A: individual OTU unique identification number; column B: individual OTU lowest taxonomic assignment; column C: axis loading per individual OTU.

Phyllosphere				Ectorhizosphere				Endorhizosphere		
A	B	C		A	B	C		A	B	C
15	*Chryseobacterium*	0.51		5	*Pseudomonas*	0.62		2	*Erwinia*	0.56
4	*Pseudomonas viridiflava*	0.41		17	Xanthomonadaceae	0.44		3	*Bacillus flexus*	0.41
17	Xanthomonadaceae	0.32		31	*Achromobacter*	0.24		5	*Pseudomonas*	0.41
221	*Pseudomonas*	0.27		26	*Acinetobacter*	0.24		6	*Streptomyces*	0.17
2	*Erwinia*	0.26		15	*Chryseobacterium*	0.20		10	Kineosporiaceae	0.15
20	*Chryseobacterium*	0.22		48	*Sphingobacterium faecium*	0.15		11	Micromonosporaceae	0.13
136	*Janthinobacterium*	0.20		75	Comamonadaceae	0.14		4	*Pseudomonas viridiflava*	0.13
7	Aeromonadaceae	0.18		4	*Pseudomonas viridiflava*	0.13		27	*Streptomyces*	0.11
49	*Wautersiella*	0.16		221	*Pseudomonas*	0.13		13	*Caulobacter henricii*	0.10
31	*Achromobacter*	0.16		2329	Enterobacteriaceae	0.12		23	Actinomycetales	0.10