Precursor processes of human self-initiated action

Short title: Precursor processes of self-initiated action

Nima Khalighinejad[a1], Aaron Schurger[b], Andrea Desantis[a2], Leor Zmigrod[c],

Patrick Haggard[a*]

[a] Institute of Cognitive Neuroscience, University College London, London WC1N 3AR, UK

[b] Cognitive Neuroimaging Unit, CEA DRF/I2BM, INSERM, Université Paris-Sud, Université Paris-Saclay, NeuroSpin center, 91191 Gif/Yvette, France

[c] Department of Psychology, University of Cambridge, Cambridge CB2 3EB, UK

* Correspondence should be addressed to Patrick Haggard, Institute of Cognitive Neuroscience, University College London, Alexandra House, 17 Queen Square, London WC1N 3AR, UK. Tel: +44 (0)20 7679 1149, E-mail: p.haggard@ucl.ac.uk

[1] NK is currently at Wellcome Centre for Integrative Neuroimaging, Department of Experimental Psychology, University of Oxford, Oxford, UK

[2] AD is currently at Laboratoire Psychologie de la Perception, Universite Paris Descartes & CNRS (UMR 8242), Paris, France

Abstract

A gradual buildup of electrical potential over motor areas precedes self-initiated movements. Recently, such "readiness potentials" (RPs) were attributed to stochastic fluctuations in neural activity. We developed a new experimental paradigm that operationalised self-initiated actions as endogenous 'skip' responses while waiting for target stimuli in a perceptual decision task. We compared these to a block of trials where participants could not choose when to skip, but were instead instructed to skip. Frequency and timing of motor action were therefore balanced across blocks, so that conditions differed only in how the timing of skip decisions was generated. We reasoned that across-trial variability of EEG could carry as much information about the source of skip decisions as the mean RP. EEG variability decreased more markedly prior to self-initiated compared to externally-triggered skip actions. This convergence suggests a consistent preparatory process prior to self-initiated action. A leaky stochastic accumulator model could reproduce this convergence given the additional assumption of a systematic decrease in input noise prior to self-initiated actions. Our results may provide a novel neurophysiological perspective on the topical debate regarding whether self-initiated actions arise from a deterministic neurocognitive process, or from neural stochasticity. We suggest that the key precursor of self-initiated action may manifest as a reduction in neural noise.

Keywords: self-initiated action; externally-triggered action; readiness potential; stochastic

fluctuations; human

1. Introduction

Functional and neuroanatomical evidence has been used to distinguish between two broad classes of human actions: self-initiated actions that happen endogenously, in the absence of any specific stimulus (Haggard, 2008; Passingham et al., 2010), and reactions to external cues. Endogenous actions are distinctive in several ways. First, they depend on an internal decision to act and are not triggered by external stimuli. In other words, the agent decides internally what to do, or when to do it, without any external cue specifying the action (Passingham et al., 2010). Second, we often deliberate and consider reasons before choosing and performing one course of action rather than an alternative. Thus, endogenous actions should be responsive to reasons (Anscombe, 2000).

Many neuroscientific studies of self-initiated action lack this reasons-responsive quality. They often involve the paradoxical instruction to 'act *freely*' e.g., "press a key when you feel the urge to do so" (Cunnington et al., 2002; Jahanshahi et al., 1995; Libet et al., 1983; Wiese et al., 2004). However, this instruction has been justifiably criticised (Nachev and Hacker, 2014). Here, we adapted for humans a paradigm previously used in animal research (Murakami et al., 2014), which embeds endogenous actions within the broader framework of decision-making. Participants responded to the direction of unpredictably-occurring dot motion stimuli by pressing left or right arrow keys (Gold and Shadlen, 2007). Importantly, they could also choose to skip waiting for the stimuli to appear, by pressing both keys simultaneously whenever they wished. The skip response thus reflects a purely endogenous decision to act, without any direct external stimulus, and provides an operational definition of a self-initiated action. Self-initiated 'skip' responses were compared to a block where participants made the same bilateral 'skip' actions in response to an unpredictable change in the fixation point (Figure. 1).

Controversies regarding precursor processes have been central to neuroscientific debates about volition (Dennett, 2015; Libet et al., 1983). The classical neural precursor is the

readiness potential (RP: (Kornhuber and Deecke, 1965). The RP is taken to be "the electro-

physiological sign of planning, preparation, and initiation of volitional acts" (Kornhuber and

Deecke, 1990) and was considered a pre-requisite of the conscious intention to act (Libet et

al., 1983; Sinnott-Armstrong and Nadel, 2010).

Classical studies explicitly or implicitly assume that the RP reflects a putative 'internal

volitional signal', with a constant, characteristic ramp-like form, necessarily preceding action

initiation - although this signal is heavily masked by noise on any individual trial (Dirnberger

et al., 2008). However, the idea that the RP reflects a specific precursor process has been

recently challenged. Instead, the time of crossing a threshold for movement could depend in

part on stochastic fluctuations in neural activity (Murakami et al., 2014; Schurger et al.,

2012). Crucially, averaging such fluctuations time-locked to action initiation reproduced the

"build-up" pattern of the mean RP, suggesting that the classical interpretation of RP as a

stable precursor of voluntary action could be deceptive. On this account, RP is not a specific,

goal-directed process that triggers action, but is rather an artefact of biased sampling and

averaging of neural noise (Murakami et al., 2014; Schurger et al., 2012).

However, classical and stochastic models offer different explanations for the variability of

EEG signals prior to self-initiated action. On the stochastic model, neural activity eventually

and necessarily converges because stochastic fluctuations must approach the motor

threshold from below. The degree to which the EEG signal converges prior to action and the

timing of that convergence should depend only on the parameters of the accumulator, and

the temporal structure of the noise input to the accumulator. In contrast, classical models

would attribute the convergence of single trial RPs to consistent precursor processes of

action preparation that reliably precede self-initiated action. While variability of RP activity

has rarely been studied previously (but see (Dirnberger et al., 2008), several studies of

externally-triggered processing have used variability of neural responses to identify neural

codes. For example, variability goes down in the interval between a go-cue and movement

onset (Churchland et al., 2006), and during perceptual processing (He, 2013; Schurger et

al., 2015). We thus compared EEG variability prior to self-initiated skip actions with variability

prior to externally-triggered actions occurring at a similar time. We used a systematic

modelling approach to show that a stochastic accumulator framework could indeed explain

the pattern of EEG variability, but only by assuming an additional process modulating the

level of neural noise.

2. Materials and Methods

2.1. Participants.

24 healthy volunteers, aged 18-35 years of age (9 male, mean age = 23 years), were

recruited from the Institute of Cognitive Neuroscience subject data pool. Two participants

were excluded before data analysis (they provided insufficient EEG data because of

excessive blinking). All participants were right handed, had normal or corrected to normal

vision, had no history or family history of seizure, epilepsy or any neurologic or psychiatric

disorder. Participants affirmed that they had not participated in any brain stimulation

experiment in the last 48 h, nor had consumed alcohol in the last 24 h. Participants were

paid an institution-approved amount for participating in the experiment. Experimental design

and procedure were approved by the UCL research ethics committee, and followed the

principles of the Declaration of Helsinki.

2.2. Behavioural task and procedure.

Participants were placed in an electrically shielded chamber, 55 cm in front of a computer

screen (60 Hz refresh rate). After signing the consent form, the experimental procedure was

explained and the EEG cap was set up. The behavioural task was as follows: participants

were instructed to look at a fixation cross in the middle of the screen. The colour of the

fixation cross changed slowly and continuously throughout the trial. This colour always

started from 'black' and then gradually changed to other colours in a randomised order. The

fixation cross changed colour gradually (e.g., from green to pink), taking 2.57 s. The fixation

cross was initially black, but the sequence of colours thereafter was random. At the same

time, participants waited for a display of randomly moving dots (displayed within a circular

aperture of 7º of diameter with a density of 14.28 dots/degree), initially moving with 0%

coherence with a speed of 2º/s (Desantis et al., 2016a, 2016b), to move coherently (step

change to 100% coherence) towards the left or right. They responded with the left or right

hand by pressing a left or right arrow key on a keyboard, accordingly. The change in dot

motion coherence happened abruptly. Correct responses were rewarded (2p). Conversely,

participants lost money (-1p) for giving a wrong answer (responding with the left hand when

dots were moving to right or vice versa), for responding before dots start moving, or not

responding within 2 s after dot motion. The trial was interrupted while such error feedback

was given. Importantly, the time of coherent movement onset was drawn unpredictably from

an exponential distribution (min = 2 s, max = 60 s, mean = 12 s), so waiting was sometimes

extremely long. However, this wait could be avoided by a 'skip' response (see later).

Participants could lose time by waiting, but receive a big reward (2p) if they responded

correctly, or could save time by 'skipping' but collect a smaller reward (1p) (Fig. 1A). The

experiment was limited to one hour, so using the skip response required a general

understanding of the trade-off between time and money. Participants were carefully informed

in advance of the rewards for responses to dot motion, and for skip responses, and were

clearly informed that the experiment had a fixed duration of one hour.

There were two blocked conditions, which differed only in the origin of the skip response. In

the 'self-initiated' condition blocks, participants could skip waiting if they chose to, by

pressing the left and right response keys simultaneously. The skip response saved time, but

produced a smaller reward (1p) than a response to dot motion. Each block consisted of 10

trials. To ensure consistent visual attention, participants were required to monitor the colour

of the fixation cross, which cycled through an unpredictable sequence of colours. At the end

of each block they were asked to classify the number of times the fixation cross turned

'yellow', according to the following categories: never, less than 50%, 50%, more than 50%.

They lost money (-1p) for giving a wrong answer. At the end of each block, participants

received feedback of total reward values, total elapsed time, and number of skips. They

could use this feedback to adjust their behaviour and maximise earnings, by regulating the

number of endogenous 'skip' responses.

In the *'externally-triggered'* condition blocks, participants could not choose for themselves

when to skip. Instead, they were instructed to skip only in response to an external signal.

The external signal was an unpredictable change in the colour of the fixation cross to 'red'

(Fig. 1B). Participants were instructed to make the skip response *as soon as* they detected

the change. The time of the red colour appearance was yoked to the time of the participant's

own previous skip responses in the immediately preceding self-initiated block, in a

randomised order. For participants who started with the externally-triggered block, the timing

of the red colour appearance in the first block only was yoked to the time of the previous

participant's last self-initiated block. The colour cycle of the fixation cross had a random

sequence, so that the onset of a red fixation could not be predicted. The fixation cross

ramped to 'red' from its previous colour in 300 ms. Again, a small reward (1p) was given for

skipping. The trial finished and the participant lost money (-1p) if s/he did not skip within 2.5

s from beginning of the ramping colour of the fixation cross. The 'red' colour was left out of

the colour cycle in the self-initiated blocks. To control for any confounding effect of attending

to the fixation cross, participants were also required to attend to the fixation cross in the self-

initiated blocks and to roughly estimate the number of times the fixation cross turned 'yellow'

(see previous). Each externally-triggered block had 10 trials, and after each block feedback

was displayed. Each self-initiated block was interleaved with an externally-triggered block,

and the order of the blocks was counterbalanced between the participants. The behavioural

task was designed in Psychophysics Toolbox Version 3 (Brainard, 1997).

2.3. EEG recording.

While participants were performing the behavioural task in a shielded chamber, EEG signals

were recorded and amplified using an ActiveTwo Biosemi system (BioSemi, Amsterdam,

The Netherlands). Participants wore a 64-channel EEG cap. To shorten the preparation

time, we recorded from a subset of electrodes that mainly covers central and visual areas:

F3, Fz, F4, FC1, FCz, FC2, C3, C1, Cz, C2, C4, CP1, CPz, CP2, P3, Pz, P4, O1, Oz, O2.

Bipolar channels placed on the outer canthi of each eye and below and above the right eye

were used to record horizontal and vertical electro-oculogram (EOG), respectively. The

Biosemi Active electrode has an output impedance of less than 1 Ohm. EEG signals were

recorded at a sampling rate of 2048 Hz.

2.4. EEG preprocessing.

EEG data preprocessing was performed in Matlab (MathWorks, MA, USA) with the help of

EEGLAB toolbox (Delorme and Makeig, 2004). Data were downsampled to 250 Hz and low-

pass filtered at 30 Hz. No high-pass filtering and no detrending were applied, to preserve

slow fluctuations. All electrodes were referenced to the average of both mastoid electrodes.

Separate data epochs of 4 s duration were extracted for self-initiated and externally-

triggered skip actions. Data epochs started from 3 s before to 1 s after the action. To avoid

EEG epochs overlapping each other any trial in which participants skipped earlier than 3 s

from trial initiation was removed. On average, 5% and 4% of trials were removed from the

self-initiated and externally-triggered conditions, respectively.

RP recordings are conventionally baseline-corrected by subtracting the average signal value

during a window from, for example, 2.5 until 2 s before action. This involves the implicit

assumption that RPs begin only in the 2 s before action onset (Shibasaki and Hallett, 2006),

but this assumption is rarely articulated explicitly, and is in fact questionable (Verbaarschot

et al., 2015). We instead took a baseline from -5 ms to +5 ms with respect to action onset.

This choice avoids making any assumption about how or when the RP starts. To ensure this

choice of baseline did not capitalize on chance, we performed parallel analyses on

demeaned data (effectively taking the entire epoch as baseline), with consistent results (see

Figure. S3). Finally, to reject non-ocular artefacts, data epochs from EEG channels (not

including EOG) with values exceeding a threshold of ±150 μv were removed. On average

7% and 8% of trials were rejected from self-initiated and externally-triggered conditions,

respectively. In the next step, Independent Component Analysis (ICA) was used to remove

ocular artefacts from the data. Ocular ICA components were identified by visual inspection.

Trials with artefacts remaining after this procedure were excluded by visual inspection.

2.5. EEG analysis.

Preliminary inspection showed a typical RP-shaped negative-going slow component that

was generally maximal at FCz. Therefore, data from FCz was chosen for subsequent

analysis. Time series analysis was performed in Matlab (MathWorks) with the help of the

FieldTrip toolbox (Oostenveld et al., 2010). We measured two dependent variables as

precursors of both *self-initiated* and *externally-triggered* skip actions: (1) mean RP amplitude

across trials and (2) variability of RP amplitudes across and within trials, measured by

standard deviation (SD). To compare across-trials SD between the two conditions, data

epochs were divided into four 500 ms windows, starting 2 s before action onset: [-2, -1.5 s],

[-1.5, -1 s], [-1, -0.5 s], [-0.5, 0 s]. All p-values were Bonferroni corrected for four

comparisons. To get a precise estimate of the standard error of the difference between

conditions, paired-samples t-tests were performed on jack-knifed data (Efron and Stein,

1981; Kiesel et al., 2008). Unlike the traditional methods, this technique compares variation

of interest across subsets of the total sample rather than across individuals, by temporarily

leaving each subject out of the calculation. In addition, we also performed cluster-based

permutation tests on SD (Maris and Oostenveld, 2007). These involve a priori identification

of a set of electrodes and a time-window of interest, and incorporate appropriate corrections

for multiple comparisons. Importantly, they avoid further arbitrary assumptions associated

with selecting specific sub-elements of the data of interest, such as individual electrodes,

time-bins or ERP components. The cluster-based tests were performed using the following

parameters: time interval = [-2 - 0 s relative to action], minimum number of neighbouring

electrodes required = 2, number of draws from the permutation distribution = 1000.

To measure variability of RP amplitudes within each individual trial, the SD of the EEG signal

from FCz was measured across time in a 100 ms window. This window was applied

successively in 30 time bins from the beginning of the epoch (3 s prior to action) to the time

of action onset. We used linear regression to calculate the slope of the within-trial SD as a

function of time (Figure. 6A). This was performed separately for each trial and each

participant. Slopes greater than 0 indicate that EEG within the 100 ms window becomes

more variable with the approach to action onset. Finally, we compared slopes of this within-

trial SD measure between self-initiated and externally-triggered conditions in a multilevel

model with single trials as level 1 and participants as level 2 variables. Multilevel analysis

was performed in R (R Core Team, Vienna, Austria).

Time-frequency analysis was performed with custom written Matlab scripts. The

preprocessed EEG time series were decomposed into their time-frequency representation by

using Complex Morlet wavelets with 20 frequencies, ranging linearly from 5 to 30 Hz. The

number of wavelet cycles increased from 3 to 7 in the same number of steps used to

increase the frequency of the wavelets from 5 to 30 Hz. Power at each trial, each frequency

and each time point was measured by convolving the raw time series with the wavelet and

squaring the resulting complex number. The power at each frequency and each time point

was then averaged across trials for each participant. Edge artefacts were removed by

discarding the first and last 500 ms of the epoch. Baseline time window was defined as the

first 500 ms of the epoch (after removal of edge artefacts: 2.5 - 2 s prior to skip action).

Changes in power during action preparation were subsequently expressed as the

percentage of change relative to the average power during the baseline time window, across

time at a specific frequency. Baseline normalisation was performed by using the following

equation:

$$percentage_{tf} = 100 \; \frac{power_{tf} - \overline{baseline_{f}}}{baseline_{f}}$$

Values > 0 indicates that power at a specific frequency (f) and a specific time (t) is higher

relative to the average power at the same frequency during the first 500 ms of the epoch.

Finally, we asked whether percentage change in power relative to baseline differs between

self-initiated and externally-triggered skip conditions in the beta band (15 – 30 Hz). Beta

band Event-related Desynchronization (ERD) during action preparation is a well-established

phenomenon (Bai et al., 2005; Doyle et al., 2005; Pfurtscheller and Lopes da Silva, 1999).

Beta power was calculated in a 500 ms window starting from 1 s and ending 0.5 s prior to

skip action. We avoided analysing later windows (e.g., 0.5 - 0 s prior to action) to avoid

possible contamination from action execution following presentation of the red fixation cross

that cued externally-triggered responses. The average normalised power across all pixels

within the selected window was then calculated for each participant and compared across

conditions using paired-samples t-tests.

2.6. Modelling and simulations.

All simulations were done in Matlab (MathWorks). We used a modified version of the Leaky

Stochastic Accumulator Model (Usher and McClelland, 2001), in which the activity of

accumulators increases stochastically over time but is limited by leakage.

$$\delta x \ = \ (I - kx)\Delta t + c\xi\sqrt{\Delta t}$$

Where I is drift rate, k is leak (exponential decay in x), ξ is Gaussian noise, c is noise scaling

factor, and Δt is the discrete time step (we used Δt = 0.001). This leaky stochastic

accumulator has been used previously to model the neural decision of 'when' to move in a

self-initiated movement task (Schurger et al., 2012). In that experiment, I was defined as the

general imperative to respond (with a constant rate). This imperative, if appropriately small in

magnitude, moves the baseline level of activity closer to the threshold, but not over it. Thus,

imperative alone does not trigger action, but does increase the likelihood of a random

threshold-crossing event triggering action. In the original model, c was assumed to be

constant and was fixed at 0.1. In a departure from the original model, we assumed that the

noise scaling factor could change linearly from an initial value of c_1 to a final value of c_2,

during action preparation. Consequently, Δc was defined as the magnitude of change in the

noise scaling factor during the trial.

240
$$\Delta c = c_2 - c_1$$

241   A negative Δc means that the signal becomes less noisy as it approaches the threshold for

242   action. Therefore, the modified model in our experiment had five free parameters: *I, k, c_1, c_2*

243   and threshold.

244
$$\delta x = (I - kx)\Delta t + c_t \xi \sqrt{\Delta t}$$

Where c_t is noise scaling factor at time *t*. The threshold was expressed as a percentile of the

output amplitude over a set of 1000 simulated trials (each of 50,000 time steps each).

Epochs of simulated data were matched to epochs of actual EEG data by identifying the

point of first threshold crossing event within each simulated trial and then extracting an

epoch from 3000 time steps before to 1000 time steps after the threshold crossing.

Parameter estimation for self-initiated skip actions was performed by fitting the model

against the *real mean* RP amplitude of each participant in the self-initiated condition. First,

1000 unique trials of Gaussian noise, each 50,000 time steps, were generated for each

participant and were fed into the model. The initial values of the model's parameters were

derived from a previous study (Schurger et al., 2012). The output of the model was then

averaged across trials and was down-sampled to 250 Hz to match the sampling rate of the

real EEG data. A *least squares* approach was used to minimise root mean squared deviation

(RMSD) between the *simulated* and *real* mean RP, by adjusting the free parameters of the

model for each participant (by using the MATLAB 'fminsearch' function). Note that this

procedure optimised the model parameters to reproduce the mean RP, rather than individual

trials.

To fit the model to our externally-triggered skip condition, we fixed the threshold of each

participant at their best fitting threshold from the self-initiated condition. We wanted to keep

the threshold the same in both conditions so that we could test the effect of changing noise

levels for a given threshold. Importantly, we also fixed the value of c_1 at its optimal value

form the self-initiated condition. By using this strategy, we can ask how noisiness of the

signal *changes*, from its initial value, and we can compare this change in noise between

conditions. We additionally performed parallel simulations without the assumption of a

common initial noise level, and obtained essentially similar results. Specifically, Δc in the all-

parameter-free model (mean= 0.02, SD = 0.06) was similar to the Δc in the model with c_1

and threshold fixed (mean= 0.02, SD = 0.05). The remaining parameters (l, k, c_2) were

optimised by minimising the deviation between the *simulated mean* RP and the *real mean*

RP in the externally-triggered condition.

Finally, we tested the model on the across-trial variability of RP epochs, having *fitted* the

model parameters to the mean RP. All parameters of the model were fixed at each

participant's optimised values for the self-initiated condition, and for the externally-triggered

condition respectively. The model was run 44 times (22 participants, x 2 conditions) with the

appropriate parameters, and 1000 separate trials were generated, each corresponding to a

putative RP exemplar. The Gaussian noise element of the model ensured that these 1000

exemplars were non-identical. The standard deviation across trials was calculated from

these 1000 simulated RP exemplars, for each participant and each condition. Importantly,

this procedure fits the model to each participant's *mean* RP amplitude, but then tests the fit

on the *standard deviation* across the 1000 simulated trials. Finally, to assess similarity

between the real and predicted SD reduction, the predicted SD in self-initiated and

externally-triggered conditions was plotted as a function of time and the area between the

two curves was computed. We then compared the area between the SD curves in a 2 s

interval prior to self-initiated and externally-triggered conditions for all participants' simulated

data, and actual data (Figure 5), using Pearson's correlation.

3. Results

3.1. Behavioural data.

Participants (n=22) waited for a display of random dots to change from 0% to 100% coherent

motion to the left or right. They responded by pressing a left or right arrow key on a

keyboard, accordingly, receiving a reward for correct responses. However, the time of

movement onset was drawn unpredictably from an exponential distribution, so waiting times

could be sometimes extremely long. In the 'self-initiated' condition blocks (Figure 1A),

participants could choose to skip waiting, by pressing both left and right response keys

simultaneously. This produced a smaller reward than a response to coherent dot motion.

Participants were informed that the experiment was limited to one hour, so that appropriate

use of the skip response implied a general understanding of the trade-off between time and

money.

Crucially, this design meant that the skip response reflected a purely endogenous decision

to act, without any direct external instruction or imperative stimulus, but rather reflecting the

general trade-off between smaller, earlier vs later, larger rewards (Green and Myerson,

2004). This operational definition of volition captures some important features of voluntary

control, such as the link between internally-generated action and a general understanding of

the distributional landscape for reasons-based decision-making (Schüür and Haggard,

2011).

We compared self-initiated skip decisions to skips in 'externally-triggered' blocks, where

participants could not choose for themselves when to skip. Instead, they were instructed to

make skip responses by a change in the fixation cross colour (Figure 1B) (see materials and

methods), yoked to the time of their own volitional skip decisions in previous blocks. Thus,

self-initiated and instructed blocks were behaviourally identical, but differed in that

participants had internal control over the hazard function in the former, but not the latter

condition.

Figure 1. Timeline of an experimental trial. Participants responded to the direction of dot-motion with

left and right keypresses. Dot-motion could begin unpredictably, after a delay drawn from an

exponential distribution. A. In the 'self-initiated' blocks participants waited for an unpredictably

occurring dot-motion stimulus, and were rewarded for correct left-right responses to motion

direction. They could decide to skip long waits for the motion stimulus, by making a bilateral

keypress. They thus decided between waiting, which lost time but brought a large reward, and

'skipping', which saved time but brought smaller rewards. The colour of the fixation cross changed

continuously during the trial, but was irrelevant to the decision task. B. In the 'externally-triggered'

blocks, participants were instructed to make bilateral skip keypresses when the fixation cross became

red, and not otherwise.

On average participants skipped 108 (SD = 16) and 106 (SD = 17) times in the self-initiated

and externally-triggered conditions, respectively. They responded to coherent dot motion in

the remaining trials (N = 177, SD = 61), with a reaction time of 767 ms (SD = 111 ms).

Those responses were correct on 86% (SD = 4%) of trials. The average waiting time before

skipping in the self-initiated condition (7.3 s, SD = 1.6) was similar to that in the externally-

triggered condition (7.6 s, SD = 1.6), confirming the success of our yoking procedure (see

materials and methods). The SD across trials had a mean of 3.17 s (SD across participants

= 1.42 s) for self-initiated skips. Our yoking procedure ensured similar values for externally-

triggered skips (mean of SD across trials 3.15 s, SD = 1.43 s). In the externally-triggered

condition, the average reaction time to the fixation cross change was 699 ms (SD = 67 ms).

On average participants earned £2.14 (SD = £0.33) from skipping and £2.78 (SD = £0.99)

from correctly responding to dot motion stimuli. This reward supplemented a fixed fee for

participation. The mean and distribution of waiting time before skip actions of each

participant are presented in Table S1 and Figure S1.

3.2. EEG variability decreases disproportionately prior to action in self-initiated and

externally-triggered conditions.

EEG data were pre-processed and averaged separately for self-initiated and externally-

triggered conditions (see materials and methods for full details). Figure 2A shows the grand

average RP amplitude in both conditions (see Figure S2A for the relation between RP-peak

amplitude and waiting time before skipping). The mean RP for self-initiated actions showed

the familiar negative-going ramp. Note that our choice to baseline-correct at the time of the

action itself (see materials and methods and Figure S3) means that the RP never in fact

reaches negative voltage values. This negative-going potential is absent from externally-

triggered skip actions (Jahanshahi et al., 1995; Papa et al., 1991). The morphology of the

mean RP might simply reflect the average of stochastic fluctuations, rather than a goal-

directed build-up. However, these theories offer differing interpretations of the variability of

individual EEG trajectories across trials (see intro).

Figure 2. EEG activity prior to skip actions. The red and blue lines represent self-initiated and externally-triggered skip conditions, respectively. Data is time-locked to the skip action (black vertical line), baseline-corrected in a 10 ms window around the skip, and recorded from FCz electrode. The average time of the skip instruction (fixation cross changing to red) in the externally-triggered condition is shown as a grey vertical line. A. Grand average RP amplitude ± standard error of the mean across participants (SEM). B. Standard deviation across trials averaged across participants ± SEM. Shaded grey area shows a significant difference between standard deviation traces across central electrodes, detected by cluster-based permutation test.

To investigate this distribution we computed standard deviation of individual trial EEG, and found a marked decrease prior to self-initiated skip action. This decrease is partly an artefact of the analysis technique: individual EEG epochs were time-locked and baseline-corrected at action onset, making the across-trial standard deviation at the time of action necessarily zero (but see Figure S3). However, this premovement drop in EEG standard deviation was more marked for self-initiated than for externally-triggered skip actions, although the analysis techniques were identical. Paired-samples t-test on jack-knifed data showed that this difference in SD was significant in the last three of the four pre-movement time bins before

skip actions (see materials and methods): that is from -1.5 to -1 s (t(21) = 4.32, p < 0.01 , d_z

= 0.92, p values are Bonferroni corrected for four comparisons), -1 to -0.5 s (t(21) = 5.97, p <

0.01 , d_z = 1.27), and -0.5 to 0 s (t(21) = 5.39, p < 0.01 , d_z = 1.15).

To mitigate any effects of arbitrary selection of electrodes or time-bins, we also performed

cluster-based permutation tests (see materials and methods). For the comparison between

SDs prior to self-initiated vs externally-triggered skip actions, a significant cluster (p < 0.01)

was identified extending from 1488 to 80 ms premovement (Figure 2B, see also Figure S3

for a different baseline and Figure S2B for the relation between EEG convergence and

waiting time before skipping). This suggests that neural activity gradually converges towards

an increasingly reliable pattern prior to self-initiated actions. Importantly, this effect is not

specific to FCz but could be observed over a wide cluster above central electrodes (Figure

S4). However, the bilateral skip response used here makes the dataset suboptimal for

thoroughly exploring the fine spatial topography of these potentials, which we hope to

address in future research.

We also analysed mean and SD EEG amplitude prior to stimulus-triggered responses to

coherent dot motion (as opposed to skip responses). Importantly, because coherent dot-

motion onset is highly unpredictable, any *general* difference in brain state between the self-

initiated skip blocks and the externally-triggered skip blocks should also be apparent prior to

coherent dot-motion onset. We did not observe any negative-going potential prior to

coherent dot motion (Figure 3A). More importantly, the SD of EEG prior to coherent dot

motion onset did not differ between conditions in any time window (p > 0.5, Bonferroni

corrected for four comparisons) (Figure 3B). This suggests that the disproportionate drop in

SD prior to skip actions (Figure 2) cannot be explained by some general contextual

difference between the two conditions, such as differences in expectation of dot stimuli, task

difficulty, temporal monitoring related to discounting and hazard function. If the decreased

variability prior to self-initiated skips had merely reflected a background, contextual process

of this kind, low variability should also be present when this process was unpredictably

interrupted by coherent dot motion. However, this was not found. Rather, reduced variability

was associated only with the period prior to self-initiated action (Figure 2), and not with any

difference in background cognitive processing between the conditions (Figure 3).

Figure 3. EEG activity prior to response to coherent dot motion direction. The red and blue lines

represent activity in self-initiated and externally-triggered blocks, respectively. Data is time-locked to

the response to coherent dot motion direction (black vertical line), baseline-corrected in a 10 ms

window around the response, and recorded from FCz electrode. The average time of the coherent dot

motion onset is shown as a grey vertical line. A. Grand average ERPs ± SEM across participants. B.

Standard deviation across trials, averaged across participants ± SEM across participants.

Finally, variability in the reaction time to respond to externally-triggered skip cues could

potentially smear out stimulus-driven preparation of skip actions. Such jitter in RT would

have the artefactual effect of increasing EEG variability across trials. To rule out this

possibility we checked whether across-trial EEG convergence was correlated across

participants with variability in behavioural reaction time to the skip response cue, but found

no significant correlation between the two variables. This suggests that the difference in

EEG convergence between self-initiated and externally-triggered skip conditions could not

be explained by mere variability in RT to skip cues (Figure S5).

3.3. Modelling the converging EEG distribution of self-initiated actions.

Leaky stochastic accumulator models have been used previously to explain the neural

decision of 'when' to move in a self-initiated task (Schurger et al., 2012). A general

imperative to perform the task shifts premotor activity up closer to threshold and then a

random threshold-crossing event provides the proximate cause of action. Hence, the precise

time of action is driven by accumulated internal physiological noise, and could therefore be

viewed as random, rather than decided (Schurger et al., 2012). However, the across-trial

variability of cortical potentials in our dataset suggests that neural activity converges on a

fixed pattern prior to self-initiated actions, to a greater extent than for externally-triggered

actions. This differential convergence could reflect a between-condition difference in the

autocorrelation function of the EEG. The early and sustained additional reduction in SD

before self-initiated actions motivated us to hypothesise an additional process of noise

control associated with self-initiated actions.

3.3.1. Sensitivity analysis

To investigate this hypothesis we first performed a sensitivity analysis by investigating how

changing key parameters of the model could influence across-trial variability of the output

(for details see materials and methods). We modelled the hypothesised process of noise

control by allowing a gradual *change in noise (Δc)* prior to action. We also explored how

changes in the key *drift (I)* and *leak (k)* parameters would influence the trial-to-trial variability

of RP. We gradually changed each parameter while holding the others fixed, and simulated

RP amplitude in 1000 trials time locked to a threshold-crossing event. SD was then

measured across these simulated trials. Simulated across-trial SDs showed that lower drift

rates and shorter leak constants were associated with a higher across-trial SD. Conversely,

reductions in noise were associated with a lower across-trial SD (Figure 4A-C). Thus, for the

model to reproduce the differential EEG convergence found in our EEG data, either the *drift*

or the *leak* should be higher, or the *change in noise* parameter should be lower, in self-

initiated compared to externally-triggered skip action conditions.

3.3.2. Model fitting and optimal parameters

We next fitted the model on the mean RP amplitude of each participant, separately for the

self-initiated and externally triggered conditions (Table S2, S3). The best fitting parameters

were then compared between the two conditions. The *drift* was significantly lower (t(21) = -

4.47, p < 0.001, after Bonferroni correction for the three parameters tested) in the self-

initiated (mean across participants = 0.09, SD = 0.03) compared to the externally-triggered

condition (mean across participants = 0.13 , SD = 0.04) (Figure 4D). The *leak* was also

significantly lower (t(21) = -4.20, p < 0.001, Bonferroni corrected) in the self-initiated (mean

across participants = 0.40, SD = 0.21) compared to the externally-triggered condition (mean

across participants = 0.58, SD = 0.20) (Figure 4E). The *change in noise* was negative in the

self-initiated (mean across participants = -0.03, SD = 0.05) but positive in the externally-

triggered condition (mean across participants = 0.02, SD = 0.05). This difference was

significant between the conditions (t(21) = -5.38, p < 0.001, Bonferroni corrected) (Figure

4F). Finally, to investigate which parameters were most sensitive to the difference between

self-initiated and externally-triggered conditions, we expressed the effect of condition on

each parameter as an effect size (standardized mean difference, Cohen's dz). Importantly,

the effect size for the between-condition difference in the *change in noise* parameter (dz =

1.15, 95%CI = [0.60 1.68]) was larger than that for the *drift* (dz = 0.95, 95%CI = [0.44 1.45])

or the *leak* (dz = 0.89, 95%CI = [0.39 1.38]) parameters (Figure 4G).

462
Figure 4 A-C. Results of sensitivity analysis. Effects of changing parameters of a stochastic

accumulator model on SD across 1000 model runs. (A) *Drift* gradually changed from 0.05 (cyan) to

0.15 (blue) in 0.02 steps, while other parameters were kept fixed. (B) *Leak* gradually changed from

0.3 (magenta) to 0.7 (blue) in 0.1 steps, while other parameters were kept fixed. (C) *Change in noise*

gradually changed from -0.05 (yellow) to 0.05 (green) in 0.02 steps, while other parameters were kept

fixed. D-F. The best fitting parameters to real mean RP amplitude in self-initiated (red) and externally-

triggered (blue) conditions. Asterisks show significant difference ($p < 0.001$). Error bars show SD

across participants. G. Effect sizes (d_z) for the between-condition difference in fitted *drift*, the *leak* and

the *change in noise* parameters. Error bars show 95% confidence interval.

So far, we fitted model parameters to the mean RP amplitude, and noted through separate

sensitivity analysis their implications for across-trial SD. Next, we directly predicted the drop

in across-trial SD of simulated RP data in self-initiated compared to externally-triggered

conditions, using the optimal model parameters for each participant in each condition. We

therefore simulated 22 RP data sets, using each participant's best fitting parameters in each

condition (see materials and methods), and computed the SD across the simulated trials. We

observed a marked additional drop in simulated across-trial SD in the self-initiated compared

to externally-triggered condition (Figure 5A, B). The differential convergence between

conditions in the simulated data closely tracked the differential convergence in our EEG data

(Correlation across participants, Pearson's r = 0.90, p < 0.001) (Figure 5C).

Figure 5. (A) Observed SD across trials averaged across participants ± SEM. Data are baselined to a

10 ms window around the skip and are recorded from FCz electrode. (B) Simulated SD across trials

averaged across participants ± SEM. The red and blue lines represent activity in self-initiated and

externally-triggered blocks, respectively. The black vertical line is the moment of skip action. (C)

Correlation between observed and simulated EEG convergence. EEG convergence was measured by

subtracting the area under the SD curve in self-initiated from the externally-triggered condition.

3.4. Within-trial reduction in EEG variability

Optimum parameter values from the model suggest that a consistent process of noise

reduction reliably occurs prior to self-initiated actions. This theory predicts that, compared to

externally-triggered actions, EEG variability should reduce more strongly not only across

trials but also within each single self-initiated action trial. To test this prediction we measured

SD within a 100 ms sliding window for each trial, and each condition (see materials and

methods) (Figure. 6A). We then used linear regression to calculate the slope of the within-

trial SD change for each trial, and compared slopes between the self-initiated and externally-

triggered conditions using a multilevel model with single trials as level 1 and participants as

level 2 variables. While EEG variability decreased within self-initiated skip trials (mean slope

= -0.01 μV/sample, SD across participants = 0.02 μV), it increased within externally-triggered

trials (mean slope = 0.01 μV/sample, SD across participants = 0.02 μV). The between-

condition difference in slopes was highly significant (t(4102) = 3.39, p < 0.001; Figure. 6B),

consistent with a progressive reduction of EEG variability prior to self-initiated actions.

Figure 6. Within-trial EEG variability. (A) SD was measured within 100ms windows for each trial and

each condition. Red and blue bars show within-trial SD in each time bin in self-initiated and externally-

triggered conditions, respectively. The solid red and blue lines show the linear fit to the time bins in

self-initiated and externally-triggered conditions, respectively. (B) The slope of the change in within-

trial variability was then compared between the self-initiated (red) and externally-triggered (blue) skip

conditions. Error bars show SEM across participants.

Figure 7. Percentage change in total EEG power compared to baseline (2.5 – 2 s prior to action) in
self-initiated (A), and externally-triggered skip conditions (B). In each condition, the percentage
change in power was computed 1 – 0.5 s prior to skip action, and from 15 – 30 Hz based on previous
literature (region of interest shown by black box). (C) The percentage change from baseline was
compared between the self-initiated (red bar) and externally-triggered (blue bar) conditions. Error bars
show SEM across participants.

Previous discussions of amplitude variation in EEG focussed on synchronised activity within

specific spectral bands (Pfurtscheller and Neuper, 1994). Preparatory decrease in beta-band

power has been used as a reliable biomarker of voluntary action (Kristeva et al., 2007).

While time-series methods identify activity that is phase-locked, spectral methods identify

EEG power that is both phase-locked and non-phase-locked, within each specific frequency

band (Cohen, 2014; Pfurtscheller and Lopes da Silva, 1999). Since motor threshold models

simply accumulate all neural activity, whether stochastic or synchronised, we reasoned that

reduction in the noise scaling factor within an accumulator model might be associated with

reduction in the synchronised activity. We therefore also investigated the decreasing

variability of neural activity prior to self-initiated action using spectral methods (see materials

and methods). Specifically, we focused on the event-related desynchronization (ERD) of

beta band activity (Bai et al., 2011; Calmels et al., 2006; Stancák and Pfurtscheller, 1996).

We compared ERD between the self-initiated and externally-triggered conditions in a 500 ms

window (1 – 0.5 s prior to action, based on previous reports (Tzagarakis et al., 2010)). Beta

power in this period decreased prior to self-initiated skip (mean percentage change = -%9.3,

SD = %7.4) (Figure. 7A), but not before externally-triggered skip actions (mean percentage

change = -%0.6, SD = %6.9) (Figure. 7B). Importantly, percentage change in beta power

was significantly different between the two conditions ($t(21) = -4.16$, $p < 0.001$) (Figure 7C).

4. Discussion

The capacity for endogenous voluntary action lies at the heart of human nature, but the brain

mechanisms that enable this capacity remain unclear. A key research bottleneck has been

the lack of convincing experimental paradigms for studying volition. Many existing paradigms

rely on paradoxical instructions equivalent to "be voluntary" or "act freely" (Haggard, 2005;

Libet et al., 1983). In a novel paradigm, we operationalized self-initiated actions as

endogenous 'skip' responses embedded in a perceptual decision task, with a long, random

foreperiod. Participants could decide to skip waiting for an imperative stimulus, by

endogenously initiating a bilateral keypress. Although previous studies in animals also used

'Giving up waiting' to study spontaneous action decisions (Murakami et al., 2014), we

believe this is the first use of this approach to study self-initiated actions in humans.

The skip action in our task has many features traditionally associated with volition including

internal-generation (Passingham et al., 2010), reasons-responsiveness (Anscombe, 2000),

freedom from immediacy (Shadlen and Gold, 2004), and a clear counterfactual alternative

(Pereboom, 2011). Crucially, operationalising self-initiated voluntary action in this way avoids

explicit instructions to "act freely", and avoids subjective reports about "volition". We

compared such actions to an exogenous skip response to a visual cue in control blocks.

The expectation of visual stimulation (coherent dot motion), and the occurrence and timing of

skip responses were all balanced across the two blocks, so the key difference is that

participants had voluntary control over skips in the self-initiated, but not the externally-

triggered blocks. We noted above that voluntary control in turn involves a number of

components, including decision, initiation, and expectation of action. We cannot be certain

how much each of these components contributes to our electrophysiological results.

However, these different components are all considered important markers of self-initiated

voluntary action.

The neural activity that generates self-initiated voluntary actions remains controversial.

Several theories attribute a key role to medial frontal regions (Krieghoff et al., 2011; Nachev

et al., 2008; Passingham, 1995). Averaged scalp EEG in humans revealed a rising negativity

beginning 1 s or more before the onset of endogenous actions (Kornhuber and Deecke,

1965), and appearing to originate in medial frontal cortex (Boschert et al., 1983; Deecke and

Kornhuber, 1978). Since this 'readiness potential' does not occur before involuntary or

externally-triggered movements, it has been interpreted as the electro-physiological sign of

planning, preparation, and initiation of self-initiated actions (Keller and Heckhausen, 1990;

Kornhuber and Deecke, 1990). RP-like brain activities preceding self-initiated actions were

also reported at the single-neuron level (Fried et al., 2011). However, the view of the RP as

a causal signal for voluntary action has been challenged, because simply averaging random

neural fluctuations that could trigger movement also produces RP-like patterns (Schurger et

al., 2012). Such stochastic accumulator models were subsequently used to predict humans'

(Schurger et al., 2012) and rats' self-initiated actions in a task similar to ours (Murakami et

al., 2014). Thus, it remains highly controversial whether the RP results from a fixed

precursor process that prepares self-initiated actions, or from random intrinsic fluctuations.

We combined an experimental design that provides a clear operational definition of volition,

and an analysis of distribution of pre-movement EEG *across and within individual trials*. We

report the novel finding that self-initiated movements are reliably preceded by a process of

variability reduction, measured as a decreasing variability in individual trial RPs, over the 1.5

s prior to movement.

Importantly, this variability reduction was specifically associated with the premovement

period before a self-initiated action: First, variability reduction was stronger prior to self-

initiated skip actions than prior to externally-triggered skip actions. Second, and crucially, the

variability reduction did not reflect any general contextual factor that might differ between

these two conditions. In our task, the onset of coherent dot motion provides an unexpected

snapshot of the brain state in the specific context provided by each condition, but not at the

time of the skip event. Figure 3 showed that any such *contextual* differences between

conditions did not affect EEG variability, and thus could not explain the reduced variability

prior to self-initiated skip actions. Thus, the reduced variability in our self-initiated skip

condition is linked to the impending action itself, and not to any general difference in the

cognitive processing or task demands between the two conditions. This pattern of results

suggests a neural precursor of self-initiated action, rather than other background contextual

factors unrelated to action preparation.

Measurement of inter-trial variability has been extensively used in the analysis of neural data

(Averbeck and Lee, 2003; Churchland et al., 2011, 2010, 2006; He, 2013; Saberi-

Moghadam et al., 2016; Schurger et al., 2015). For example, presenting a target stimulus

decreases inter-trial variability of neural firing rate in premotor cortex (Churchland et al.,

2006). Interestingly, RTs to external stimuli are shortest when variability is lowest,

suggesting that a decrease in neural variability is a marker of motor preparation (Churchland

et al., 2006). Moreover, reducing neural variability is characteristic of cortical responses to

any external stimulus (Churchland et al., 2010), and could be a reliable signature of

conscious perception (Schurger et al., 2015). Importantly, in previous studies, the decline in

neural variability was *triggered* by a target stimulus, i.e. decreasing neural variability was

triggered exogenously (Churchland et al., 2010). Our results show that inter-trial variability

also decreases prior to a self-initiated action, in the absence of any external target.

Classical models might attribute variability reduction prior to self-initiated action to a

consistent process of preparation. In contrast, stochastic fluctuation models have been

recently used to account for the neural activity preceding self-initiated actions in humans

(Schurger et al., 2012) and rodents (Murakami et al., 2014). We did not aim in this

experiment to compare these models directly, but to investigate their predictions regarding

the shape and variability of the RP. Our modelling showed variability reduction could be

explained within a stochastic fluctuation model, *with the additional assumption* of progressive

decrease in the input noise level. In the absence of external evidence, stochastic models

depend only on internal physiological noise to determine the time of action. Thus, Schurger

et al.'s model first shifts premotor activity closer to a motor threshold, while the actual

threshold-crossing event is triggered by accumulating stochastic fluctuations (Schurger et

al., 2012). By fitting a modified version of the leaky stochastic accumulator model on each

participant's mean RP amplitude, we observed that integration of internal noise evolves

differently prior to self-initiated and externally-triggered skip actions. The rate of the *drift* and

the *leak* was lower and the *change in noise* was negative prior to self-initiated actions,

compared to externally-triggered actions. Importantly, by fitting model parameters to each

participant's mean RP, and testing the variability of EEG data generated with those

parameters, we found that variability reduction before self-initiated action was mainly driven

by a gradually-reducing noise level.

Previous studies show that changes in noise level influence choice, RT and confidence in

accumulation-to-bound models of perceptual decision making (Fetsch et al., 2014;

Furstenberg et al., 2015; Kiani et al., 2008; Zylberberg et al., 2016). Interestingly, the

motivating effects of reward on speed and accuracy of behaviour were recently shown to be

attributable to active control of internal noise (Manohar et al., 2015). In general, previous

studies show an important role of active noise control in tasks requiring responses to

external stimuli (Kool and Botvinick, 2013; Manohar et al., 2015). We have shown that

similar processes may underlie self-initiated action, and that a consistent process of noise

reduction may be a key precursor of self-initiated voluntary action. This additional process of

noise control may make a stochastic approach more similar to classical models of voluntary

action.

Finally, we showed that a decrease in premotor neural variability prior to self-initiated action

is not only observed across-trials, but is also realised within-trial and as a reduction in EEG

power in the beta frequency band. The observed reduction in beta-band power is entirely

consistent with the proposed reduction in neural noise preceding self-initiated action that

was suggested by our modelling. Clearly, any natural muscular action must have *some*

precursors. Sherrington's final common path concept proposed that descending neural

commands from primary motor cortex necessarily preceded voluntary action (Sherrington,

1906). However, it remains unclear *how long* before action such precursor processes can be

identified. Our result provides a new method for addressing this question. The question is

theoretically important, because cognitive accounts of self-initiated action control divide into

two broad classes. In classical accounts, a fixed, and relatively long-lasting precursor

process is caused by a prior decision to act (Anscombe, 2000; Kornhuber and Deecke,

1990). In other recent accounts, stochastic fluctuations would dominate until a relatively late

stage, and fixed precursor processes would be confined to brief, motoric execution

processes (Schurger et al., 2012).

The precursor processes that our method identifies may be necessary for self-initiated

action, but may not be sufficient: identifying a precursor process prior to self-initiated

movement says nothing about whether and how often such a process might also be present

in the absence of movement. On one view, the precursor process might occur quite

frequently, but a last-minute decision might influence whether a given precursor event

completes with a movement, or not. Our movement-locked analyses cannot identify any

putative precursor processes or precursor-like processes that failed to result in a movement.

However, our spectral analyses (Figure 7) make this possibility unlikely. They show a

gradual decline in total beta-band power beginning around 1 s prior to self-initiated action.

Any putative unfulfilled precursor processes would presumably produce partial versions of

this effect throughout the epoch, but these are not readily apparent. Lastly, there might be a

nonlinear relation between the recorded signals and the decision process. Our analyses

assumed a simple, linear relation between the decision process and the measured variables.

This assumption may be simplistic, but almost all analyses of neural data make similar

assumptions at some level.

Interestingly, our endogenous skip response resembles the decision to explore during

foraging behaviour (Constantino and Daw, 2015; Kolling et al., 2012). That is, endogenous

skip responses amounted to deciding to look out for dot-motion stimuli in forthcoming time-

periods, rather than the present one. This prompts the speculation that spontaneous

transition from rest to foraging or vice-versa could be an early evolutionary antecedent of

human volition.

In conclusion, we show that self-initiated actions have a reliable precursor, namely a

consistent process of neural variability reduction prior to movement. We showed that this

variability reduction was not due to a background contextual process that differed between

self-initiated and externally-triggered conditions, but was related to self-initiated action. We

began this paper by distinguishing between a classical model, in which a fixed preparation

process preceded self-initiated action, and a fully stochastic model, in which the triggering of

self-initiated action is essentially random – although the artefact of working with movement-

locked epochs might give the appearance of a specific causal signal such as the RP. We

found that the precursor process prior to self-initiated action could be modelled within a

stochastic framework, given the additional assumption of a progressive reduction in input

noise. Future research might usefully investigate whether the precursor process we have

identified is the cause or the consequence of the subjective 'decision to act'.

5. Author Contributions

Conceptualization, N.K., P.H., A.S., and A.D.; Methodology, N.K., A.S., and A.D.; Formal

Analysis, N.K., L.Z., and P.H.; Investigation, N.K., L.Z.; Writing-original draft, N.K., Writing-

review & editing, P.H., A.S., Supervision, P.H. and A.S.

6. Acknowledgement

This work was supported by the European Research Council Advanced Grant HUMVOL

(Grant number: 323943). This work was additionally supported by a grant from

The Leverhulme Trust (Ref. RPG-2016-378) to PH. AS was supported by a starting grant

from the European Research Council (Grant number 640626). The authors declare no

competing financial interest.

References

Anscombe, G.E.M., 2000. Intention. Harvard University Press.

Averbeck, B.B., Lee, D., 2003. Neural noise and movement-related codes in the macaque supplementary motor area. J. Neurosci. Off. J. Soc. Neurosci. 23, 7630–7641.

Bai, O., Mari, Z., Vorbach, S., Hallett, M., 2005. Asymmetric spatiotemporal patterns of event-related desynchronization preceding voluntary sequential finger movements: a high-resolution EEG study. Clin. Neurophysiol. Off. J. Int. Fed. Clin. Neurophysiol. 116, 1213–1221. doi:10.1016/j.clinph.2005.01.006

Bai, O., Rathi, V., Lin, P., Huang, D., Battapady, H., Fei, D.-Y., Schneider, L., Houdayer, E., Chen, X., Hallett, M., 2011. Prediction of human voluntary movement before it occurs. Clin. Neurophysiol. Off. J. Int. Fed. Clin. Neurophysiol. 122, 364–372. doi:10.1016/j.clinph.2010.07.010

Boschert, J., Hink, R.F., Deecke, L., 1983. Finger movement versus toe movement-related potentials: further evidence for supplementary motor area (SMA) participation prior to voluntary action. Exp. Brain Res. 52, 73–80.

Brainard, D.H., 1997. The Psychophysics Toolbox. Spat. Vis. 10, 433–436.

Calmels, C., Holmes, P., Jarry, G., Lévèque, J.-M., Hars, M., Stam, C.J., 2006. Cortical activity prior to, and during, observation and execution of sequential finger movements. Brain Topogr. 19, 77–88. doi:10.1007/s10548-006-0014-x

Churchland, A.K., Kiani, R., Chaudhuri, R., Wang, X.-J., Pouget, A., Shadlen, M.N., 2011. Variance as a signature of neural computations during decision making. Neuron 69, 818–831. doi:10.1016/j.neuron.2010.12.037

Churchland, M.M., Yu, B.M., Cunningham, J.P., Sugrue, L.P., Cohen, M.R., Corrado, G.S., Newsome, W.T., Clark, A.M., Hosseini, P., Scott, B.B., Bradley, D.C., Smith, M.A., Kohn, A., Movshon, J.A., Armstrong, K.M., Moore, T., Chang, S.W., Snyder, L.H., Lisberger, S.G., Priebe, N.J., Finn, I.M., Ferster, D., Ryu, S.I., Santhanam, G., Sahani, M., Shenoy, K.V., 2010. Stimulus onset quenches neural variability: a widespread cortical phenomenon. Nat. Neurosci. 13, 369–378. doi:10.1038/nn.2501

Churchland, M.M., Yu, B.M., Ryu, S.I., Santhanam, G., Shenoy, K.V., 2006. Neural Variability in Premotor Cortex Provides a Signature of Motor Preparation. J. Neurosci. 26, 3697–3712. doi:10.1523/JNEUROSCI.3762-05.2006

Cohen, M.X., 2014. Analyzing Neural Time Series Data: Theory and Practice, 1 edition. ed. MIT Press, Cambridge, Massachusetts.

Constantino, S.M., Daw, N.D., 2015. Learning the opportunity cost of time in a patch-foraging task. Cogn. Affect. Behav. Neurosci. 15, 837–853. doi:10.3758/s13415-015-0350-y

Cunnington, R., Windischberger, C., Deecke, L., Moser, E., 2002. The preparation and execution of self-initiated and externally-triggered movement: a study of event-related fMRI. NeuroImage 15, 373–385. doi:10.1006/nimg.2001.0976

Deecke, L., Kornhuber, H.H., 1978. An electrical sign of participation of the mesial "supplementary" motor cortex in human voluntary finger movement. Brain Res. 159, 473–476.

Delorme, A., Makeig, S., 2004. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. J. Neurosci. Methods 134, 9–21. doi:10.1016/j.jneumeth.2003.10.009

Dennett, D.C., 2015. Elbow Room: The Varieties of Free Will Worth Wanting, New edition edition. ed. MIT Press, Cambridge, Massachusetts ; London, England.

Desantis, A., Waszak, F., Gorea, A., 2016a. Agency alters perceptual decisions about action-outcomes. Exp. Brain Res. doi:10.1007/s00221-016-4684-7

Desantis, A., Waszak, F., Moutsopoulou, K., Haggard, P., 2016b. How action structures time: About the perceived temporal order of action and predicted outcomes. Cognition 146, 100–109. doi:10.1016/j.cognition.2015.08.011

Dirnberger, G., Lang, W., Lindinger, G., 2008. A new method to determine temporal variability in the period of pre-movement electroencephalographic activity. Int. J. Psychophysiol. Off. J. Int. Organ. Psychophysiol. 70, 165–170. doi:10.1016/j.ijpsycho.2008.08.006

Doyle, L.M.F., Yarrow, K., Brown, P., 2005. Lateralization of event-related beta desynchronization in the EEG during pre-cued reaction time tasks. Clin. Neurophysiol. Off. J. Int. Fed. Clin. Neurophysiol. 116, 1879–1888. doi:10.1016/j.clinph.2005.03.017

Efron, B., Stein, C., 1981. The Jackknife Estimate of Variance. Ann. Stat. 9, 586–596.

Fetsch, C.R., Kiani, R., Newsome, W.T., Shadlen, M.N., 2014. Effects of Cortical Microstimulation on Confidence in a Perceptual Decision. Neuron 83, 797–804. doi:10.1016/j.neuron.2014.07.011

Fried, I., Mukamel, R., Kreiman, G., 2011. Internally generated preactivation of single neurons in human medial frontal cortex predicts volition. Neuron 69, 548–562. doi:10.1016/j.neuron.2010.11.045

Furstenberg, A., Breska, A., Sompolinsky, H., Deouell, L.Y., 2015. Evidence of Change of Intention in Picking Situations. J. Cogn. Neurosci. 1–14. doi:10.1162/jocn_a_00842

Gold, J.I., Shadlen, M.N., 2007. The Neural Basis of Decision Making. Annu. Rev. Neurosci. 30, 535–574. doi:10.1146/annurev.neuro.29.051605.113038

Green, L., Myerson, J., 2004. A Discounting Framework for Choice With Delayed and Probabilistic Rewards. Psychol. Bull. 130, 769–792. doi:10.1037/0033-2909.130.5.769

Haggard, P., 2008. Human volition: towards a neuroscience of will. Nat. Rev. Neurosci. 9, 934–946. doi:10.1038/nrn2497

Haggard, P., 2005. Conscious intention and motor cognition. Trends Cogn. Sci. 9, 290–295. doi:10.1016/j.tics.2005.04.012

He, B.J., 2013. Spontaneous and task-evoked brain activity negatively interact. J. Neurosci. Off. J. Soc. Neurosci. 33, 4672–4682. doi:10.1523/JNEUROSCI.2922-12.2013

Jahanshahi, M., Jenkins, I.H., Brown, R.G., Marsden, C.D., Passingham, R.E., Brooks, D.J., 1995. Self-initiated versus externally triggered movements. I. An investigation using measurement of regional cerebral blood flow with PET and movement-related potentials in normal and Parkinson's disease subjects. Brain J. Neurol. 118 (Pt 4), 913–933.

Keller, I., Heckhausen, H., 1990. Readiness potentials preceding spontaneous motor acts: voluntary vs. involuntary control. Electroencephalogr. Clin. Neurophysiol. 76, 351–361.

Kiani, R., Hanks, T.D., Shadlen, M.N., 2008. Bounded Integration in Parietal Cortex Underlies Decisions Even When Viewing Duration Is Dictated by the Environment. J. Neurosci. 28, 3017–3029. doi:10.1523/JNEUROSCI.4761-07.2008

Kiesel, A., Miller, J., Jolicoeur, P., Brisson, B., 2008. Measurement of ERP latency differences: a comparison of single-participant and jackknife-based scoring methods. Psychophysiology 45, 250–274. doi:10.1111/j.1469-8986.2007.00618.x

Kolling, N., Behrens, T.E.J., Mars, R.B., Rushworth, M.F.S., 2012. Neural mechanisms of foraging. Science 336, 95–98. doi:10.1126/science.1216930

Kool, W., Botvinick, M., 2013. The intrinsic cost of cognitive control. Behav. Brain Sci. 36, 697-698; discussion 707-726. doi:10.1017/S0140525X1300109X

Kornhuber, H.H., Deecke, L., 1990. Readiness for movement - The Bereitschaftspotential story. Citation Classic Commentary. Curr. Contents Life Sci. 33.

Kornhuber, H.H., Deecke, L., 1965. Hirnpotentialänderungen bei Willkürbewegungen und passiven Bewegungen des Menschen: Bereitschaftspotential und reafferente Potentiale. Pflüg. Arch. Für Gesamte Physiol. Menschen Tiere 284, 1–17. doi:10.1007/BF00412364

Krieghoff, V., Waszak, F., Prinz, W., Brass, M., 2011. Neural and behavioral correlates of intentional actions. Neuropsychologia 49, 767–776. doi:10.1016/j.neuropsychologia.2011.01.025

Kristeva, R., Patino, L., Omlor, W., 2007. Beta-range cortical motor spectral power and corticomuscular coherence as a mechanism for effective corticospinal interaction during steady-state motor output. NeuroImage 36, 785–792. doi:10.1016/j.neuroimage.2007.03.025

Libet, B., Gleason, C.A., Wright, E.W., Pearl, D.K., 1983. Time of conscious intention to act in relation to onset of cerebral activity (readiness-potential). The unconscious initiation of a freely voluntary act. Brain J. Neurol. 106 (Pt 3), 623–642.

Manohar, S.G., Chong, T.T.-J., Apps, M.A.J., Batla, A., Stamelou, M., Jarman, P.R., Bhatia, K.P., Husain, M., 2015. Reward Pays the Cost of Noise Reduction in Motor and Cognitive Control. Curr. Biol. CB 25, 1707–1716. doi:10.1016/j.cub.2015.05.038

Maris, E., Oostenveld, R., 2007. Nonparametric statistical testing of EEG- and MEG-data. J. Neurosci. Methods 164, 177–190. doi:10.1016/j.jneumeth.2007.03.024

Murakami, M., Vicente, M.I., Costa, G.M., Mainen, Z.F., 2014. Neural antecedents of self-initiated actions in secondary motor cortex. Nat. Neurosci. doi:10.1038/nn.3826

Nachev, P., Hacker, P., 2014. The neural antecedents to voluntary action: a conceptual analysis. Cogn. Neurosci. 5, 193–208. doi:10.1080/17588928.2014.934215

Nachev, P., Kennard, C., Husain, M., 2008. Functional role of the supplementary and pre-supplementary motor areas. Nat. Rev. Neurosci. 9, 856–869. doi:10.1038/nrn2478

Oostenveld, R., Fries, P., Maris, E., Schoffelen, J.-M., Oostenveld, R., Fries, P., Maris, E., Schoffelen, J.-M., 2010. FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data, FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. Comput. Intell. Neurosci. Comput. Intell. Neurosci. 2011, 2011, e156869. doi:10.1155/2011/156869, 10.1155/2011/156869

Papa, S.M., Artieda, J., Obeso, J.A., 1991. Cortical activity preceding self-initiated and externally triggered voluntary movement. Mov. Disord. Off. J. Mov. Disord. Soc. 6, 217–224. doi:10.1002/mds.870060305

Passingham, R., 1995. The Frontal Lobes and Voluntary Action. Oxford University Press.

Passingham, R.E., Bengtsson, S.L., Lau, H.C., 2010. Medial frontal cortex: from self-generated action to reflection on one's own performance. Trends Cogn. Sci. 14, 16–21. doi:10.1016/j.tics.2009.11.001

Pereboom, D., 2011. Free Will Skepticism and Meaning in Life, in: Kane, R. (Ed.), The Oxford Handbook of Free Will. Oxford University Press.

Pfurtscheller, G., Lopes da Silva, F.H., 1999. Event-related EEG/MEG synchronization and desynchronization: basic principles. Clin. Neurophysiol. Off. J. Int. Fed. Clin. Neurophysiol. 110, 1842–1857.

Pfurtscheller, G., Neuper, C., 1994. Event-related synchronization of mu rhythm in the EEG over the cortical hand area in man. Neurosci. Lett. 174, 93–96.

Saberi-Moghadam, S., Ferrari-Toniolo, S., Ferraina, S., Caminiti, R., Battaglia-Mayer, A., 2016. Modulation of Neural Variability in Premotor, Motor, and Posterior Parietal Cortex during Change of Motor Intention. J. Neurosci. Off. J. Soc. Neurosci. 36, 4614–4623. doi:10.1523/JNEUROSCI.3300-15.2016

Schurger, A., Sarigiannidis, I., Naccache, L., Sitt, J.D., Dehaene, S., 2015. Cortical activity is more stable when sensory stimuli are consciously perceived. Proc. Natl. Acad. Sci. U. S. A. 112, E2083-2092. doi:10.1073/pnas.1418730112

Schurger, A., Sitt, J.D., Dehaene, S., 2012. An accumulator model for spontaneous neural activity prior to self-initiated movement. Proc. Natl. Acad. Sci. 109, E2904–E2913. doi:10.1073/pnas.1210467109

Schüür, F., Haggard, P., 2011. What are self-generated actions? Conscious. Cogn. 20, 1697–1704. doi:10.1016/j.concog.2011.09.006

Shadlen, M.N., Gold, J.I., 2004. , in: Gazzaniga, M.S. (Ed.), The Cognitive Neurosciences. MIT Press, pp. 1229–1241.

Sherrington, C.S., 1906. The integrative action of the nervous system. New York, C Scribner's sons.

Shibasaki, H., Hallett, M., 2006. What is the Bereitschaftspotential? Clin. Neurophysiol. Off. J. Int. Fed. Clin. Neurophysiol. 117, 2341–2356. doi:10.1016/j.clinph.2006.04.025

Sinnott-Armstrong, W., Nadel, L. (Eds.), 2010. Conscious Will and Responsibility: A Tribute to Benjamin Libet, 1 edition. ed. OUP USA, Oxford ; New York.

Stancák, A., Pfurtscheller, G., 1996. Event-related desynchronisation of central beta-rhythms during brisk and slow self-paced finger movements of dominant and nondominant hand. Brain Res. Cogn. Brain Res. 4, 171–183.

Tzagarakis, C., Ince, N.F., Leuthold, A.C., Pellizzer, G., 2010. Beta-band activity during motor planning reflects response uncertainty. J. Neurosci. Off. J. Soc. Neurosci. 30, 11270–11277. doi:10.1523/JNEUROSCI.6026-09.2010

Usher, M., McClelland, J.L., 2001. The time course of perceptual choice: the leaky, competing accumulator model. Psychol. Rev. 108, 550–592.

Verbaarschot, C., Farquhar, J., Haselager, P., 2015. Lost in time...: The search for intentions and Readiness Potentials. Conscious. Cogn. 33C, 300–315. doi:10.1016/j.concog.2015.01.011

Wiese, H., Stude, P., Nebel, K., de Greiff, A., Forsting, M., Diener, H.C., Keidel, M., 2004. Movement preparation in self-initiated versus externally triggered movements: an event-related fMRI-study. Neurosci. Lett. 371, 220–225. doi:10.1016/j.neulet.2004.08.078

Zylberberg, A., Fetsch, C.R., Shadlen, M.N., 2016. The influence of evidence volatility on choice, reaction time and confidence in a perceptual decision. eLife 5. doi:10.7554/eLife.17688

Supporting Information

Figure S1. Histogram of waiting times before skip actions in self-initiated condition, across all trials and all participants. The dashed red line shows the average waiting time.

A

B

Figure S2. Correlation between participants' mean waiting time (s) before skipping in the self-initiated condition and RP-peak amplitude (A), and EEG convergence (B). EEG convergence was measured by subtracting the area under the SD curve in self-initiated from the externally-triggered condition. There was a significant negative correlation between waiting time and RP-peak amplitude (Pearson's r = -0.69, p < 0.01), and maximum EEG convergence (Pearson's r = -0.66, p < 0.01).

Figure S3. Demeaned EEG activity prior to skip actions. The red and blue lines represent self-initiated and externally-triggered skips, respectively. Data is time-locked to the skip action (black vertical line), and is baselined to the mean of entire epoch (i.e., demeaned), and recorded from FCz electrode. A. Grand average RP amplitude. B. Standard deviation across trials averaged across participants. Shaded area show significant clusters across central electrodes, detected by cluster-based permutation test. Whereas baselining to a limited time window forces a low SD within the baseline time window, and a progressive rise in SD with temporal distance before or after the baseline, the use of a broad baseline time window, as here, reduces this artefactual effect of baseline-correction on variability of time-locked data. Nevertheless, the difference in SD between conditions remains significant.

Figure S4. Topography of the difference in SDs between self-initiated and externally-triggered conditions. Small circles represent EEG electrodes across which the permutation test was performed. Electrodes that showed significant difference between conditions have been marked *. The time interval (s) is indicated above each subplot.

Fig S5. No significant correlation (Pearson's r = 0.16, p = 0.46) across participants between standard deviation of each participant's RT to externally-triggered skip cues (ms), and EEG convergence. EEG convergence was measured by subtracting the area under the SD curve in self-initiated from the externally-triggered condition.

Subject	Mean wait (s) self-initiated	Mean wait (s) externally-triggered	SD (s) self-initiated	SD (s) externally-triggered
1	11.76	12.09	6.82	6.79
2	5.38	5.76	2.27	2.30
3	6.14	6.45	4.58	4.61
4	7.16	7.45	3.25	3.22
5	8.25	8.39	4.57	3.97
6	7.21	7.70	3.22	3.27
7	5.92	6.20	1.66	1.67
8	7.83	8.16	3.20	3.23
9	6.08	6.48	1.26	1.20
10	6.32	6.54	2.96	2.68
11	5.34	5.85	1.00	1.01
12	6.89	7.28	2.72	2.70
13	4.83	5.23	1.83	1.82
14	8.06	8.23	2.66	2.54
15	8.97	9.44	3.60	3.60
16	9.25	9.66	5.85	5.91
17	6.30	6.81	2.69	2.61
18	7.72	8.14	2.53	2.57
19	6.90	7.16	2.94	2.93
20	7.32	7.80	2.40	2.62
21	9.44	9.85	2.93	2.94
22	7.14	7.73	4.74	5.19

Table S1. Mean and standard deviation of waiting time before skipping in self-initiated and externally-triggered conditions.

Drift (I)	Leak (k)	Initial noise (c_1)	Final noise (c_2)	Threshold	Change in noise (Δc)
0.04	-0.18	0.08	0.01	0.10	-0.07
0.11	0.55	0.12	0.13	0.14	0.00
0.11	0.57	0.13	0.11	0.18	-0.02
0.11	0.52	0.12	0.12	0.17	0.00
0.09	0.23	0.19	0.14	0.41	-0.05
0.01	0.38	0.23	0.13	0.42	-0.10
0.10	0.50	0.14	0.13	0.20	-0.01
0.12	0.62	0.10	0.14	0.08	0.04
0.11	0.57	0.11	0.13	0.11	0.01
0.02	0.27	0.25	0.12	0.44	-0.13
0.08	0.14	0.21	0.13	0.38	-0.08
0.11	0.53	0.12	0.12	0.13	0.01
0.09	0.11	0.21	0.12	0.35	-0.09
0.06	0.17	0.21	0.12	0.39	-0.09
0.07	0.28	0.19	0.14	0.36	-0.05
0.11	0.53	0.12	0.11	0.12	-0.01
0.11	0.54	0.12	0.12	0.13	0.01
0.08	0.22	0.18	0.12	0.29	-0.06
0.11	0.60	0.12	0.13	0.09	0.01
0.11	0.53	0.12	0.12	0.13	0.01
0.17	0.59	0.18	0.24	0.74	0.06
0.11	0.55	0.13	0.12	0.15	0.00

Table S2. Optimum parameters for self-initiated skip action. The values were detected by fitting the model against the mean RP amplitude of each participant in self-initiated condition. Δc was measured by subtracting the initial noise level (c_1) from the final noise level (c_2).

Drift (l)	Leak (k)	Initial noise (c_1)	Final noise (c_2)	Threshold	Change in noise (Δc)
0.09	0.42	0.08	0.10	0.10	0.02
0.11	1.01	0.12	0.21	0.14	0.09
0.10	0.35	0.13	0.13	0.18	0.00
0.16	0.70	0.12	0.16	0.17	0.04
0.12	0.48	0.19	0.18	0.41	-0.01
0.10	0.48	0.23	0.17	0.42	-0.05
0.09	0.26	0.14	0.07	0.20	-0.06
0.17	0.93	0.10	0.24	0.08	0.13
0.18	0.44	0.11	0.19	0.11	0.08
0.05	0.44	0.25	0.22	0.44	-0.03
0.10	0.38	0.21	0.17	0.38	-0.04
0.19	0.72	0.12	0.15	0.13	0.03
0.10	0.41	0.21	0.18	0.35	-0.03
0.08	0.54	0.21	0.17	0.39	-0.04
0.12	0.51	0.19	0.23	0.36	0.05
0.16	0.87	0.12	0.15	0.12	0.04
0.18	0.69	0.12	0.16	0.13	0.04
0.10	0.45	0.18	0.15	0.29	-0.03
0.21	0.75	0.12	0.17	0.09	0.05
0.17	0.77	0.12	0.20	0.13	0.09
0.10	0.61	0.18	0.20	0.74	0.01
0.16	0.58	0.13	0.16	0.15	0.03

Table S3. Optimum parameters for externally-triggered skip action. The values were detected by fitting the model against the mean RP amplitude of each participant in externally-triggered condition. Δc was measured by subtracting the initial noise level (c_1) from the final noise level (c_2). c_1 and the threshold were fixed at their optimum values in self-initiated condition (see materials and methods for more details)