Human self-initiated action is reliably preceded by a process of noise reduction

Short title: Neural precursors of self-initiated action

Nima Khalighinejad[1], Aaron Schurger[2], Andrea Desantis[1], Leor Zmigrod[3], Patrick Haggard[1*]

[1] Institute of Cognitive Neuroscience, University College London, London WC1N 3AR, UK

[2] Cognitive Neuroimaging Unit, CEA DRF/I2BM, INSERM, Université Paris-Sud, Université Paris-Saclay, NeuroSpin center, 91191 Gif/Yvette, France

[3] Department of Psychology, University of Cambridge, Cambridge CB2 3EB, UK

* Correspondence should be addressed to Patrick Haggard**,** Institute of Cognitive Neuroscience, University College London, Alexandra House, 17 Queen Square, London WC1N 3AR, UK. Tel: +44 (0)20 7679 1149, E-mail: p.haggard@ucl.ac.uk

NK is currently at Department of Experimental Psychology, University of Oxford, Oxford, UK

AD is currently at Laboratoire Psychologie de la Perception, Universite Paris Descartes & CNRS (UMR 8242), Paris, France

Abstract

A gradual buildup of electrical potential over motor areas precedes self-initiated movements. These "readiness potentials" (RPs) could simply reflect stochastic fluctuations in neural activity. We operationalised self-initiated actions as endogenous 'skip' responses while waiting for target stimuli in a perceptual decision task. Across-trial variability of EEG decreased more markedly prior to self-initiated compared to externally-triggered skip actions. This convergence towards a fixed pattern suggests a consistent preparatory process prior to self-initiated action. A leaky stochastic accumulator model could reproduce these features of the data, given the additional assumption of a decrease in noise level at the input to the accumulator prior to self-initiated, but not externally-triggered actions. The assumed reduction in neural noise was supported by analyses of both within-trial EEG variability and of spectral power. We suggest that a process of noise reduction is consistently recruited prior to self-initiated action. This precursor event may underlie the emergence of RP.

Keywords: self-initiated action; externally-triggered action; readiness potential; stochastic fluctuations; human

1   Introduction

Functional and neuroanatomical evidence has been used to distinguish between two broad

classes of human actions: self-initiated actions that happen endogenously, in the absence of

any specific stimulus (Haggard, 2008; Passingham, Bengtsson, & Lau, 2010), and reactions

to external cues. Endogenous actions are distinctive in several ways. First, they depend on

an internal decision to act and are not triggered by external stimuli. In other words, the agent

decides internally what to do, or when to do it, without any external cue specifying the action

(Passingham et al., 2010). Second, we often deliberate and consider reasons before

choosing and performing one course of action rather than an alternative. Thus, endogenous

actions should be responsive to reasons (Anscombe, 2000). These features of endogenous

action capture many key attributes of human volition. While there has been extensive

research on action selection and initiation in response to external cues, the brain

mechanisms for endogenous, self-initiated actions have been less studied.

Many studies of human voluntary action involve the paradoxical instruction to 'act *freely*' e.g.,

"press a key when you feel the urge to do so" (Cunnington, Windischberger, Deecke, &

Moser, 2002; Jahanshahi et al., 1995; Libet, Gleason, Wright, & Pearl, 1983; Wiese et al.,

2004). However, the situation and task demands of such experiments are complex, and

have been justly criticised (Nachev & Hacker, 2014). We adapted for humans a paradigm

previously used in animal research (Murakami, Vicente, Costa, & Mainen, 2014), which

embeds endogenous actions within the broader framework of reward-guided perceptual

decision-making. Participants responded to the direction of unpredictably-occurring dot

motion stimuli by pressing left or right arrow keys (Gold & Shadlen, 2007). Importantly, they

could also choose to skip waiting for the stimuli to appear, by pressing both keys

simultaneously whenever they wished. The skip response thus reflects a purely endogenous

decision to act, without any direct external stimulus, and provides an operational definition of

a self-initiated action. Self-initiated 'skip' responses were compared to a block where

participants made the same bilateral 'skip' actions in response to an unpredictable change in

the fixation point (Figure. 1).

Controversies regarding precursor processes have been central to neuroscientific debates

about volition (Dennett, 2015; Libet et al., 1983). The classical neural marker of precursor

processes for endogenous action is the readiness potential (RP: Kornhuber & Deecke,

1965). The RP is taken to be "the electro-physiological sign of planning, preparation, and

initiation of volitional acts" (Kornhuber & Deecke, 1990) and was considered a pre-requisite

of the conscious intention to act (Libet et al., 1983; Sinnott-Armstrong & Nadel, 2010).

Classical studies explicitly or implicitly assume that the RP reflects a putative 'internal

volitional signal', with a constant, characteristic ramp-like form, necessarily preceding action

initiation - although this signal is heavily masked by noise in any individual trial (Dirnberger,

Lang, & Lindinger, 2008). However, the very idea that the RP reflects a specific precursor

process leading to endogenous action has been recently challenged. Alternative models

suggest that the rising ramp pattern of the mean RP does not reflect a goal-directed process

but rather reflects subthreshold stochastic fluctuations that influence the precise time of

crossing the threshold for movement (Murakami et al., 2014; Schurger, Sitt, & Dehaene,

2012). Crucially, averaging these random fluctuations time-locked to action initiation results

in the rising ramp pattern of the mean RP, with its appearance of a stable ERP component.

According to the stochastic account, the RP reflects cross-trial averaging of data epochs

time locked to crests in autocorrelated neural noise, rather than a specific, goal-directed

process that causes action. This view, which can be formally expressed in a quantitative

model, and tested against neural data (Murakami et al., 2014; Schurger et al., 2012), has

radically revised neuroscientific theories of voluntary action.

Both classical and stochastic models can reproduce the existence of mean RP, treating it as

signal or as averaged noise, respectively. However, the two models make different

predictions about EEG variability prior to action. On the stochastic model, neural activity

eventually and necessarily converges because stochastic fluctuations must always approach

the motor threshold from below. The time of convergence depends only on the temporal

autocorrelation of the EEG signal. On the classical model, the distribution of single trial RPs

additionally converges because the RP marks a consistent precursor process that reliably

precedes self-initiated action. While variability of RP activity has rarely been studied

previously (but see Dirnberger et al., 2008), several studies of externally-triggered

processing have used variability of neural responses to identify neural codes. For example,

variability goes down in the interval between a go cue and movement onset (Churchland,

Yu, Ryu, Santhanam, & Shenoy, 2006), and during perceptual processing (He, 2013;

Schurger, Sarigiannidis, Naccache, Sitt, & Dehaene, 2015). We thus hypothesised that

variability of neural activity should decrease more markedly prior to self-initiated skip actions.

The *additional* decrease in variability would be a marker of a consistent preparatory process,

or precursor for self-initiated action.

We found an additional drop in inter-trial variability beginning 1.5 s before self-initiated skips,

compared to externally-driven skip actions. This differential convergence suggests a

consistent precursor process that reliably precedes self-initiated action. The gradual onset of

differential convergence suggests that the precursor process has a variable duration, rather

than a fixed duration (Dirnberger et al., 2008), although we could not formally compare these

alternative models using our methods. We showed that a modified version of an established

computational model based on stochastic fluctuations on neural activity (Schurger et al.,

2012; Usher & McClelland, 2001) could capture the patterns of EEG variability found in our

data. Crucially, the model required a specific modification to explain the data, namely a

neurocognitive process of noise reduction, time-locked to the moment of action. Our model

further predicts that decreases in within-trial EEG fluctuation and spectral power should be

observed prior to self-initiated action. These predictions were confirmed. Importantly, these

differences between conditions began well before the stimulus onset in our externally-

triggered condition (Figures 2 & 7), so presumably reflects changes in ongoing EEG prior to

self-initiated action. Thus, impending self-initiated voluntary action appears to be associated

with a distinctive, time-specific stabilisation of premovement EEG patterns.

Results

Behavioural data.

Participants (n=22) waited for a display of random dots to move coherently (step change

from 0% to 100% coherence) towards the left or right. They responded with the left or right

hand by pressing a left or right arrow key on a keyboard, accordingly. They received a

reward for correct responses. However, the time of movement onset was drawn

unpredictably from an exponential distribution, so waiting was sometimes extremely long. In

the '*self-initiated*' condition blocks (Figure 1A), participants could skip waiting if they chose

to, by pressing the left and right response keys simultaneously. The skip response saved

time, but produced a smaller reward than a response to dot motion. The experiment was

limited to one hour, so using the skip response implied a general understanding of the trade-

off between time and money. A skip response thus reflects a purely endogenous decision to

act, in the absence of any external instruction to act, and based on the tradeoff between

later, larger, and smaller, earlier rewards. This provides an operational definition of volition

within our experimental design, which captures some of the important features of

endogenous voluntary control, as well as the linkage of self-initiated action to other aspects

of cognition, such as decision-making and judgement (Schüür & Haggard, 2011). In the

'*externally-triggered*' condition blocks, participants could not choose for themselves when to

skip. Instead, they were instructed to make skip responses by an external signal (Figure 1B)

(see materials and methods).

A

B

✖ Fixation cross changes colour during the trial

Figure 1. Timeline of an experimental trial. Participants responded to the direction of dot-motion with left and right keypresses. Dot-motion could begin unpredictably, after a delay drawn from an exponential distribution. A. In the 'self-initiated' blocks participants waited for an unpredictably occurring dot-motion stimulus, and were rewarded for correct left-right responses to motion direction. They could decide to skip long waits for the motion stimulus, by making a bilateral keypress. They thus decided between waiting, which lost time but brought a large reward, and 'skipping', which saved time but brought smaller rewards. The colour of the fixation cross changed continuously during the trial, but was irrelevant to the decision task. B. In the 'externally-triggered' blocks, participants were instructed to make bilateral skip keypresses when the fixation cross became red, and not otherwise.

On average participants skipped 108 (SD = 16) and 106 (SD = 17) times in the self-initiated and externally-triggered conditions, respectively. They responded to coherent dot motion in

the remaining trials (N = 177, SD = 61), with a reaction time of 767 ms (SD = 111 ms).

Those responses were correct on 86% (SD = 4%) of trials. The average waiting time before

skipping in the self-initiated condition (7.3 s, SD = 1.6) was similar to that in the externally-

triggered condition (7.6 s, SD = 1.6), confirming the success of our yoking procedure (see

materials and methods). The waiting time varied more across trials within each individual,

than across individuals, suggesting that self-initiated skip responses represented an on-line

decision to act, rather than a pre-decided stereotyped response. Thus, the SD across trials

had a mean of 3.17 s (SD across participants = 1.42 s) for self-initiated skips. Our yoking

procedure ensured similar values for externally-triggered skips (mean of SD across trials

3.15 s, SD = 1.43 s). In the externally-triggered condition, the average reaction time to the

fixation cross change was 699 ms (SD = 67 ms). On average participants earned £2.14 (SD

= £0.33) from skipping and £2.78 (SD = £0.99) from correctly responding to dot motion

stimuli. This reward supplemented a fixed fee for participation. The mean and distribution of

waiting time before skip action of each participant are presented in Table S1 and Figure S1.

EEG variability decreases disproportionately prior to action in self-initiated and externally-

triggered conditions.

EEG data were pre-processed and averaged separately for self-initiated and externally-

triggered conditions. Figure 2A shows the grand average RP amplitude in both conditions.

The mean RP for self-initiated actions showed the familiar negative-going ramp. Note that

our choice to baseline-correct at the time of the action itself (see materials and methods)

means that the RP never in fact reaches negative voltage values. This negative-going

potential is absent from externally-triggered skip actions (Jahanshahi et al., 1995; Papa,

Artieda, & Obeso, 1991). The morphology of the mean RP might simply reflect the average

of stochastic fluctuations, rather than a goal-directed build-up. However, these theories offer

differing interpretations of the variability of individual EEG trajectories across trials (see

intro).

Figure 2. EEG activity prior to skip actions. The red and blue lines represent self-initiated and externally-triggered skip conditions, respectively. Data is time-locked to the skip action (black vertical line), baseline-corrected in a 10 ms window around the skip, and recorded from FCz electrode. The average time of the skip instruction (fixation cross changing to red) in the externally-triggered condition is shown as a grey vertical line. A. Grand average RP amplitude ± standard error of the mean across participants (SEM). B. Standard deviation across trials averaged across participants ± SEM. Shaded grey area shows a significant difference between standard deviation traces across central electrodes, detected by cluster-based permutation test.

To investigate this distribution we computed standard deviation of individual trial EEG, and found a marked decrease prior to self-initiated skip action. This decrease is partly an artefact of the analysis technique: individual EEG epochs were time-locked and baseline-corrected at action onset, making the across-trial standard deviation at the time of action necessarily zero (but see Figure S2). However, this premovement drop in EEG standard deviation was more marked for self-initiated than for externally-triggered skip actions, although the analysis techniques were identical. Paired-samples t-test on jack-knifed data showed that this difference in SD was significant in the last three of the four pre-movement time bins before

skip actions (see materials and methods): that is from -1.5 to -1 s (t(21) = 4.32, p < 0.01 , d_z

= 0.92, p values are Bonferroni corrected for four comparisons), -1 to -0.5 s (t(21) = 5.97, p <

0.01 , d_z = 1.27), and -0.5 to 0 s (t(21) = 5.39, p < 0.01 ,d_z = 1.15).

To mitigate any effects of arbitrary selection of electrodes or time-bins, we also performed

cluster-based permutation tests (see materials and methods). For the comparison between

SDs prior to self-initiated vs externally-triggered skip actions, a significant cluster (p < 0.01)

was identified extending from 1488 to 80 ms premovement (Figure 2B, see also Figure S2

for a different baseline). This suggests that neural activity gradually converges towards an

increasingly reliable pattern prior to self-initiated actions. Importantly, this effect is not

specific to FCz but could be observed over a wide cluster above central electrodes (Figure

S3). However, the bilateral skip response used here makes the dataset suboptimal for

thoroughly exploring the fine spatial topography of these potentials, which we hope to

address in future research.

It has been shown that stimulus anticipation is preceded by a cortical negative wave, the

contingent negative variation (CNV) (Walter, Cooper, Aldridge, McCallum, & Winter, 1964).

The CNV has been associated with expectation and temporal processing (Casini & Vidal,

2011; Van Rijn, Kononowicz, Meck, Ng, & Penney, 2011). Hence, our measures in the self-

initiated condition could reflect both accumulating conditions that make a skip action

desirable (e.g., passage of time without dot motion onset), and the preparation of the skip

action itself. However, our externally-triggered skip condition controls for effects of mere

passage of time, and expectation of dot motion onset.

To ensure that the key cognitive factors in the task were balanced between self-initiated and

externally-triggered conditions, we also analysed mean and SD EEG amplitude prior to

stimulus-triggered responses to coherent dot motion (as opposed to skip responses). We did

not observe any negative-going potential prior to coherent dot motion (Figure 3A), again

suggesting that temporal expectation did not strongly contribute to our ERPs. More

importantly, the SD of EEG prior to coherent dot motion onset did not differ between

conditions in any time window (p > 0.5, Bonferroni corrected for four comparisons) (Figure

3B). This suggests that the disproportionate drop in SD prior to skip actions cannot be

explained merely by a difference in expectation of dot stimuli or temporal processing.

Moreover, an explanation based on temporal processing would presumably predict stronger

EEG convergence when participants wait longer before skip action. In fact, we found a

negative correlation between EEG convergence and waiting time (Figure S4).

Figure 3. EEG activity prior to response to coherent dot motion direction. The red and blue lines
represent activity in self-initiated and externally-triggered blocks, respectively. Data is time-locked to
the response to coherent dot motion direction (black vertical line), baseline-corrected in a 10 ms
window around the response, and recorded from FCz electrode. The average time of the coherent dot
motion onset is shown as a grey vertical line. A. Grand average ERPs ± SEM across participants. B.
Standard deviation across trials, averaged across participants ± SEM across participants.

Finally, variability in the reaction time to respond to externally-triggered skip cues could

potentially smear out stimulus-driven preparation of skip actions. Such jitter in RT would

have the artefactual effect of increasing EEG variability across trials. To rule out this

possibility we checked whether across-trial EEG convergence was correlated across

participants with variability in behavioural reaction time to the skip response cue, but found

no significant correlation between the two variables. This suggests that the difference in

EEG convergence between self-initiated and externally-triggered skip conditions could not

be explained by mere variability in RT to skip cues (Figure S5).

Modelling the converging EEG distribution of self-initiated actions.

Leaky stochastic accumulator models have been used previously to explain the neural

decision of 'when' to move in a self-initiated task (Schurger et al., 2012). A general

imperative to perform the task shifts the premotor activity up closer to threshold and then a

random threshold-crossing event provides the proximate cause of action. Hence, the precise

time of action is driven by accumulated internal physiological noise, and could therefore be

viewed as random, rather than decided (Schurger et al., 2012). However, the across-trial

variability of cortical potentials in our dataset suggests that neural activity converges on a

fixed pattern prior to self-initiated actions, to a greater extent than for externally-triggered

actions. This differential convergence could reflect a between-condition difference in the

autocorrelation function of the EEG. The early and sustained additional reduction in SD

before self-initiated actions motivated us to hypothesise an additional process of noise

control associated with self-initiated actions.

Sensitivity analysis

To investigate this hypothesis we first performed a sensitivity analysis by investigating how

changing key parameters of the model could influence across-trial variability of the output

(for details see materials and methods). We modelled the hypothesised process of noise

control by allowing a gradual *change in noise (Δc)* prior to action. We also explored how

changes in the key *drift (I)* and *leak (k)* parameters would influence the trial-to-trial variability

of RP. We gradually changed each parameter while holding the others fixed, and simulated

RP amplitude in 1000 trials time locked to a threshold-crossing event. SD was then

measured across these simulated trials. Simulated across-trial SDs showed that lower drift

rates and shorter leak constants were associated with a higher across-trial SD. Conversely,

reductions in noise were associated with a lower across-trial SD (Figure 4A-C). Thus, for the

model to reproduce the differential EEG convergence found in our EEG data, either the *drift*

or the *leak* should be higher, or the *change in noise* parameter should be lower, in self-

initiated compared to externally-triggered skip action conditions.

Model fitting and optimal parameters

We next fitted the model on the mean RP amplitude of each participant, separately for the

self-initiated and externally triggered conditions (Table S2, S3). The best fitting parameters

were then compared between the two conditions. The *drift* was significantly lower (t(21) = -

4.47, p < 0.001, after Bonferroni correction for the three parameters tested) in the self-

initiated (mean across participants = 0.09, SD = 0.03) compared to the externally-triggered

condition (mean across participants = 0.13 , SD = 0.04) (Figure 4D). The *leak* was also

significantly lower (t(21) = -4.20, p < 0.001, Bonferroni corrected) in the self-initiated (mean

across participants = 0.40, SD = 0.21) compared to the externally-triggered condition (mean

across participants = 0.58, SD = 0.20) (Figure 4E). The *change in noise* was negative in the

self-initiated (mean across participants = -0.03, SD = 0.05) but positive in the externally-

triggered condition (mean across participants = 0.02, SD = 0.05). This difference was

significant between the conditions (t(21) = -5.38, p < 0.001, Bonferroni corrected) (Figure

4F). Finally, to investigate which parameters were most sensitive to the difference between

self-initiated and externally-triggered conditions, we expressed the effect of condition on

each parameter as an effect size (standardized mean difference, Cohen's dz). Importantly,

the effect size for the between-condition difference in the *change in noise* parameter (dz =

1.15, 95%CI = [0.60 1.68]) was larger than that for the *drift* (dz = 0.95, 95%CI = [0.44 1.45])

or the *leak* (dz = 0.89, 95%CI = [0.39 1.38]) parameters (Figure 4G).

254
Figure 4 A-C. Results of sensitivity analysis. Effects of changing parameters of a stochastic

accumulator model on SD across 1000 model runs. (A) *Drift* gradually changed from 0.05 (cyan) to

0.15 (blue) in 0.02 steps, while other parameters were kept fixed. (B) *Leak* gradually changed from

0.3 (magenta) to 0.7 (blue) in 0.1 steps, while other parameters were kept fixed. (C) *Change in noise*

gradually changed from -0.05 (yellow) to 0.05 (green) in 0.02 steps, while other parameters were kept

fixed. D-F. The best fitting parameters to real mean RP amplitude in self-initiated (red) and externally-

triggered (blue) conditions. Asterisks show significant difference (p < 0.001). Error bars show SD

across participants. G. Effect sizes (d_z) for the between-condition difference in fitted *drift*, the *leak* and

the *change in noise* parameters. Error bars show 95% confidence interval.

So far, we fitted model parameters to the mean RP amplitude, and noted through separate

sensitivity analysis their implications for across-trial SD. Next, we directly predicted the drop

in across-trial SD of simulated RP data in self-initiated compared to externally-triggered

conditions, using the optimal model parameters for each participant in each condition. We

therefore simulated 22 RP data sets, using each participant's best fitting parameters in each

condition (see materials and methods), and computed the SD across the simulated trials. We

observed a marked additional drop in simulated across-trial SD in the self-initiated compared

to externally-triggered condition (Figure 5A, B). The differential convergence between

conditions in the simulated data closely tracked the differential convergence in our EEG data

(Correlation across participants, Pearson's r = 0.90, p < 0.001) (Figure 5C).

Figure 5. (A) Observed SD across trials averaged across participants ± SEM. Data are baselined to a

10 ms window around the skip and are recorded from FCz electrode. (B) Simulated SD across trials

averaged across participants ± SEM. The red and blue lines represent activity in self-initiated and

externally-triggered blocks, respectively. The black vertical line is the moment of skip action. (C)

Correlation between observed and simulated EEG convergence. EEG convergence was measured by

subtracting the area under the SD curve in self-initiated from the externally-triggered condition.

Within-trial reduction in EEG variability

Optimum parameter values from the model suggest that a consistent process of noise

reduction reliably occurs prior to self-initiated actions. This theory predicts that, compared to

externally-triggered actions, EEG variability should reduce more strongly not only across

trials but also within each single self-initiated action trial. To test this prediction we measured

SD within a 100 ms sliding window for each trial, and each condition (see materials and

methods) (Figure. 6A). We then used linear regression to calculate the slope of the within-

trial SD change for each trial, and compared slopes between the self-initiated and externally-

triggered conditions using a multilevel model with single trials as level 1 and participants as

level 2 variables. While EEG variability decreased within self-initiated skip trials (mean slope

= -0.01 µV/sample, SD across participants = 0.02 µV), it increased within externally-triggered

trials (mean slope = 0.01 µV/sample, SD across participants = 0.02 µV). The between-

condition difference in slopes was highly significant (t(4102) = 3.39, p < 0.001; Figure. 6B),

consistent with a progressive reduction of EEG variability prior to self-initiated actions.

Figure 6. Within-trial EEG variability. (A) SD was measured within 100ms windows for each trial and

each condition. Red and blue bars show within-trial SD in each time bin in self-initiated and externally-

triggered conditions, respectively. The solid red and blue lines show the linear fit to the time bins in

self-initiated and externally-triggered conditions, respectively. (B) The slope of the change in within-

trial variability was then compared between the self-initiated (red) and externally-triggered (blue) skip

conditions. Error bars show SEM across participants.

Figure 7. Percentage change in total EEG power compared to baseline (2.5 – 2 s prior to action) in

self-initiated (A), and externally-triggered skip conditions (B). In each condition, the percentage

change in power was computed 1 – 0.5 s prior to skip action, and from 15 – 30 Hz based on previous

literature (region of interest shown by black box). (C) The percentage change from baseline was

compared between the self-initiated (red bar) and externally-triggered (blue bar) conditions. Error bars

show SEM across participants.

Previous discussions of amplitude variation in EEG focussed on synchronised activity within

specific spectral bands (Pfurtscheller & Neuper, 1994). Preparatory decrease in beta-band

power has been used as a reliable biomarker of voluntary action (Kristeva, Patino, & Omlor,

2007). While time-series methods identify activity that is phase-locked, spectral methods

identify EEG power that is both phase-locked and non-phase-locked, within each specific

frequency band (Cohen, 2014; Pfurtscheller & Lopes da Silva, 1999). Since motor threshold

models simply accumulate all neural activity, whether stochastic or synchronised, we

reasoned that reduction in the noise scaling factor within an accumulator model might be

associated with reduction in the synchronised activity. We therefore also investigated the

decreasing variability of neural activity prior to self-initiated action using spectral methods

(see materials and methods). Specifically, we focused on the event-related

desynchronization (ERD) of beta band activity (Bai et al., 2011; Calmels et al., 2006;

Stancák & Pfurtscheller, 1996). We compared ERD between the self-initiated and externally-

triggered conditions in a 500 ms window (1 – 0.5 s prior to action, based on previous reports

(Tzagarakis, Ince, Leuthold, & Pellizzer, 2010)). Beta power in this period decreased prior to

self-initiated skip (mean percentage change = -%9.3, SD = %7.4) (Figure. 7A), but not

before externally-triggered skip actions (mean percentage change = -%0.6, SD = %6.9)

(Figure. 7B). Importantly, percentage change in beta power was significantly different

between the two conditions ($t(21) = -4.16$, $p < 0.001$) (Figure 7C).

Discussion

The capacity for endogenous voluntary action lies at the heart of human nature, but the brain

mechanisms that enable this capacity remain unclear. A key research bottleneck has been

the lack of convincing experimental paradigms for studying volition. Many existing paradigms

rely on paradoxical instructions equivalent to "be voluntary" or "act freely" (Haggard, 2005;

Libet et al., 1983). In a novel paradigm, we operationalized self-initiated actions as

endogenous 'skip' responses embedded in a perceptual decision task, with a long, random

foreperiod. Participants could decide to skip waiting for an imperative stimulus, by

endogenously initiating a bilateral keypress. Although previous studies in animals also used

'Giving up waiting' to study spontaneous action decisions (Murakami et al., 2014), we

believe this is the first use of this approach to study self-initiated actions in humans.

Our experimental task provided a purely operational definition of self-initiated, endogenous

skip responses. Thus, the skip action can be understood from a strict behaviourist

perspective. However, skip responses have many of the hallmarks of volition traditionally

used in philosophy, including internal-generation (Passingham et al., 2010), reasons-

responsiveness (Anscombe, 2000), freedom from immediacy (Shadlen & Gold, 2004), and a

clear counterfactual alternative (Pereboom, 2011). Crucially, operationalising self-initiated

voluntary action in this way avoids explicit instructions to "act freely", and avoids subjective

reports about "volition". We compared such actions to an exogenous skip response triggered

by a visual cue in control blocks.

The neural activity that generates self-initiated voluntary actions remains controversial.

Several theories attribute a key role to medial frontal regions (Krieghoff, Waszak, Prinz, &

Brass, 2011; Nachev, Kennard, & Husain, 2008; Passingham, 1995). Averaged scalp EEG

in humans revealed a rising negativity beginning 1 s or more before the onset of

endogenous actions (Kornhuber & Deecke, 1965), and appearing to originate in medial

frontal cortex (Boschert, Hink, & Deecke, 1983; Deecke & Kornhuber, 1978). Since this

'readiness potential' does not occur before involuntary or externally-triggered movements, it

has been interpreted as the electro-physiological sign of planning, preparation, and initiation

of self-initiated actions (Keller & Heckhausen, 1990; Kornhuber & Deecke, 1990). RP-like

brain activities preceding self-initiated actions were also reported at the single-neuron level

(Fried, Mukamel, & Kreiman, 2011). However, the view of the RP as a causal signal for

voluntary action has been challenged, because simply averaging random neural fluctuations

that trigger a motor action also produces RP-like patterns (Schurger et al., 2012). Such

stochastic accumulator models were subsequently used to predict humans' (Schurger et al.,

2012) and rats' self-initiated actions in a task similar to ours (Murakami et al., 2014). Thus, it

remains highly controversial whether the RP results from a specific precursor process that

prepares self-initiated actions, or from random intrinsic fluctuations. We combined an

experimental design that provides a clear operational definition of volition, and an analysis of

distribution across and within individual trials of pre-movement EEG. Our results support a

novel combination of both the classical and the stochastic views. We report the novel finding

that self-initiated movements are reliably preceded by a process of noise reduction. This

process alters the pattern of stochastic fluctuations that accumulate towards the motor

threshold.

EEG showed decreased trial-to-trial variability prior to skip actions. This partly reflects the

time-locking and baseline-correction at the time of action: ERP methods necessarily imply

zero variability at the baseline (Luck, 2005). However, around 1.5 s prior to skip actions, the

decrease in variability became more marked for self-initiated compared to control externally-

triggered skip actions. Since the skip action in the externally-triggered control condition has

no endogenous volitional component, the decrease in variability prior to skip actions in the

control condition presumably reflects only the effects of time-locking, and the temporal

autocorrelation of the background EEG. However, the *additional* decrease in variability prior

to self-initiated action may reflect convergence of neural activity towards a steady trajectory

that precedes self-initiated actions. We hypothesised that this could indicate a consistent

preparatory process leading to self-initiated voluntary action.

Measurement of inter-trial variability has been extensively used in the analysis of neural data

(Averbeck & Lee, 2003; Churchland et al., 2011; Churchland et al., 2010, 2006; He, 2013;

Saberi-Moghadam, Ferrari-Toniolo, Ferraina, Caminiti, & Battaglia-Mayer, 2016; Schurger et

al., 2015). For example, presenting a target stimulus decreases inter-trial variability of neural

firing rate in premotor cortex (Churchland et al., 2006). Interestingly, RTs to external stimuli

are shortest when variability is lowest, suggesting that a decrease in neural variability is a

marker of motor preparation (Churchland et al., 2006). Moreover, reducing neural variability

is characteristic of cortical responses to any external stimulus (Churchland et al., 2010), and

could be a reliable signature of conscious perception (Schurger et al., 2015). Importantly, in

previous studies, the decline in neural variability was *triggered* by a target stimulus, i.e.

decreasing neural variability was triggered exogenously (Churchland et al., 2010). Our

results show that inter-trial variability also decreases prior to a self-initiated action, in the

absence of any external target.

Integration to bound models have been recently used to account for the neural activity

preceding self-initiated actions in humans (Schurger et al., 2012) and rodents (Murakami et

al., 2014). In the absence of external evidence, these models are fed solely with internal

physiological noise. Importantly, when signal-to-noise ratio is low the timing of the decision

to move is mainly determined by random fluctuations. Schurger et al.'s model first shifts

premotor activity closer to a motor threshold. This is followed by a threshold-crossing event,

triggered by stochastic fluctuations (Schurger et al., 2012). By fitting a modified version of

the leaky stochastic accumulator model on each participant's mean RP amplitude, we

observed that integration of internal noise evolves differently prior to self-initiated and

externally-triggered skip actions. The rate of the *drift* and the *leak* was lower and the *change*

in noise was negative prior to self-initiated actions, compared to externally-triggered actions.

Importantly, analysis of across-trial variability of simulated data, using model parameters

optimised for each participant, implied that the marked drop in variability that we observed

prior to self-initiated action was mainly driven by a gradually reducing noise level. Previous

studies show that changes in noise level influences choice, RT and confidence in

accumulation-to-bound models of perceptual decision (Fetsch, Kiani, Newsome, & Shadlen,

2014; Furstenberg, Breska, Sompolinsky, & Deouell, 2015; Kiani, Hanks, & Shadlen, 2008;

Zylberberg, Fetsch, & Shadlen, 2016). Interestingly, the motivating effects of reward on

speed and accuracy of behaviour were recently shown to be attributable to active control of

internal noise (Manohar et al., 2015). In general, previous studies show an important role of

active noise control in tasks requiring responses to external stimuli (Kool & Botvinick, 2013;

Manohar et al., 2015). We have shown that similar processes may underlie self-initiated

action, and that a consistent process of noise reduction may be a key precursor of self-

initiated voluntary action.

Finally, we showed that a decrease in premotor neural variability prior to self-initiated action

is not only observed across-trials, but is also realised within-trial and as a reduction in power

of beta frequency band. Clearly, any natural muscular action must have *some* precursors.

Sherrington's final common path concept proposed that descending neural commands from

primary motor cortex necessarily preceded voluntary action (Sherrington, 1906). However, it

remains unclear *how long* before action such precursor processes can be identified. Our

result provides a new method for addressing this question. The question is theoretically

important, because cognitive accounts of self-initiated action control divide into two broad

classes. In classical accounts, a fixed, and relatively long-lasting precursor process is

caused by a prior decision to act (Anscombe, 2000; Kornhuber & Deecke, 1990). In other

recent accounts, stochastic fluctuations dominate until a relatively late stage, and fixed

precursor processes would be confined to brief, motoric execution processes (Schurger et

al., 2012).

Our study cannot show whether self-initiated voluntary actions are caused by prior decisions,

or by randomness. However, our results do suggest that the contribution of stochastic

fluctuations is supplemented by a precursor process of noise reduction starting from around

1.5 s prior to action. Importantly, and by the same token, our results cannot show whether

the precursor process of noise reduction is initiated by some top-down decision, or is itself

triggered by some ongoing spontaneous fluctuations. Further, the precursor processes that

our method identifies may be necessary for self-initiated action, but may not be sufficient:

identifying a precursor process prior to self-initiated movement says nothing about whether

and how often such a process might also be present in the absence of movement. On one

view, the precursor process might occur quite frequently, but a last-minute decision might

influence whether the precursor process completes with a movement, or is vetoed. Our

movement-locked analyses cannot identify any putative vetoed precursor processes, or

precursor-like processes that failed to result in a movement. However, our spectral analyses

(Figure 7) make this possibility unlikely. They show a gradual decline in total beta-band

power beginning around 1 s prior to self-initiated action. Any putative vetoed precursor

processes would produce partial versions of this effect at other time-points in the epoch, but

these are not readily apparent.

Inter-trial variability provides an additional dimension to information coding in the brain (He,

2013; Schurger et al., 2015; Stein, Gossen, & Jones, 2005). We showed that both inter- and

within-trial variability decreases prior to a self-initiated action, akin to a reliable preparatory

process. Our computational modelling further suggests that this preparatory stabilising

process may itself reflect some as yet unknown mechanism of noise reduction. Actively

regulating noise at optimal levels typically enhances system performance (Faisal, Selen, &

Wolpert, 2008; Fitts, 1954; Groen & Wenderoth, 2016; Shu, Hasenstaub, Badoual, Bal, &

McCormick, 2003). However, noise regulation can also arise incidentally, as a result of

attractor dynamics in the motor system, as in the "optimal subspace hypothesis" (Shenoy,

Sahani, & Churchland, 2013). Further, the noise reduction mechanism in our data could itself

be triggered either by a specific top-down signal or a stochastic event. Finally, we speculate

that the process of noise reduction not only explains the reduction in inter- and within-trial

EEG variability prior to self-initiated action, but also generates the slow rising negativity of

the RP, and the well-known beta ERD that precedes voluntary actions. ERD represents an

activation in cortical areas that produce motor behaviour (Pfurtscheller, 1992; Pfurtscheller &

Lopes da Silva, 1999). Factors such as effort and attention enhance the ERD (Defebvre,

Bourriez, Destée, & Guieu, 1996). However, the relation between ERD and RP remained

unclear. Our modelling results suggest that a reliable process of noise reduction could

explain both the ERD, and the RP.

Interestingly, our endogenous skip response resembles the decision to explore during

foraging behaviour (Constantino & Daw, 2015; Kolling, Behrens, Mars, & Rushworth, 2012).

That is, endogenous skip responses amounted to deciding to look out for dot-motion stimuli

in forthcoming time-periods, rather than the present one. This prompts the speculation that

spontaneous transition from rest to foraging or vice-versa could be an early evolutionary

antecedent of human volition.

In conclusion, we show that self-initiated actions have a reliable precursor, namely a

consistent process of neural noise reduction prior to movement. We began this paper by

distinguishing between a classical model, in which a fixed preparation process consistently

preceded self-initiated action, and a fully stochastic model, in which the triggering of self-

initiated action is essentially random – though the artefact of working with movement-locked

epochs might give the appearance of a consistent precursor event such as the RP. We have

identified a reliable precursor process, but this precursor process can be accommodated as

a parameter change within the stochastic model framework. Future research might usefully

investigate whether the precursor process is the cause or the consequence of the subjective

'decision to act'.

Materials and Methods

Participants.

24 healthy volunteers, aged 18-35 years of age (9 male, mean age = 23 years), were

recruited from the Institute of Cognitive Neuroscience subject data pool. Two participants

were excluded before data analysis (they provided insufficient EEG data because of

excessive blinking). All participants were right handed, had normal or corrected to normal

vision, had no history or family history of seizure, epilepsy or any neurologic or psychiatric

disorder. Participants affirmed that they had not participated in any brain stimulation

experiment in the last 48 h, nor had consumed alcohol in the last 24 h. Participants were

paid an institution-approved amount for participating in the experiment. Experimental design

and procedure were approved by the UCL research ethics committee, and followed the

principles of the Declaration of Helsinki.

 Behavioural task and procedure.

Participants were placed in an electrically shielded chamber, 55 cm in front of a computer

screen (60 Hz refresh rate). After signing the consent form, the experimental procedure was

explained and the EEG cap was set up. The behavioural task was as follows: participants

were instructed to look at a fixation cross in the middle of the screen. The colour of the

fixation cross changed slowly and continuously throughout the trial. This colour always

started from 'black' and then gradually changed to other colours in a randomised order. The

fixation cross changed colour gradually (e.g., from green to pink), taking 2.57 s. The fixation

cross was initially black, but the sequence of colours thereafter was random. At the same

time, participants waited for a display of randomly moving dots (displayed within a circular

aperture of 7° of diameter with a density of 14.28 dots/degree, initially moving with 0%

coherence with a speed of 2°/s (Desantis, Waszak, & Gorea, 2016; Desantis, Waszak,

Moutsopoulou, & Haggard, 2016), to move coherently (step change to 100% coherence)

towards the left or right. They responded with the left or right hand by pressing a left or right

arrow key on a keyboard, accordingly. The change in dot motion coherence happened

abruptly. Correct responses were rewarded (2p). Conversely, participants lost money (-1p)

for giving a wrong answer (responding with the left hand when dots were moving to right or

vice versa), for responding before dots start moving, or not responding within 2 s after dot

motion. The trial was interrupted while such error feedback was given. Importantly, the time

of coherent movement onset was drawn unpredictably from an exponential distribution (min

= 2 s, max = 60 s, mean = 12 s), so waiting was sometimes extremely long. However, this

wait could be avoided by a 'skip' response (see later). Participants could lose time by

waiting, but receive a big reward (2p) if they responded correctly, or could save time by

'skipping' but collect a smaller reward (1p) (Fig. 1A). The experiment was limited to one

hour, so using the skip response required a general understanding of the trade-off between

time and money. Participants were carefully informed in advance of the rewards for

responses to dot motion, and for skip response, and were clearly informed that the

experiment had a fixed duration of one hour.

There were two blocked conditions, which differed only in the origin of the skip response. In

the 'self-initiated' condition blocks, participants could skip waiting if they chose to, by

pressing the left and right response keys simultaneously. The skip response saved time, but

produced a smaller reward (1p) than a response to dot motion. Each block consisted of 10

trials. To ensure consistent visual attention, participants were required to monitor the colour

of the fixation cross, which cycled through an unpredictable sequence of colours. At the end

of each block they were asked to classify the number of times the fixation cross turned

'yellow', according to the following categories : never, less than 50%, 50%, more than 50%.

They lost money (-1p) for giving a wrong answer. At the end of each block, participants

received feedback of total reward values, total elapsed time, and number of skips. They

could use this feedback to adjust their behaviour and maximise earnings, by regulating the

number of endogenous 'skip' responses.

In the 'externally-triggered' condition blocks, participants could not choose for themselves

when to skip. Instead, they were instructed to make skip responses by an external signal.

The external signal was an unpredictable change in the colour of the fixation cross to 'red'

(Fig. 1B). Participants were instructed to make the skip response as soon as they detected

the change. The time of the red colour appearance was yoked to the time of participant's

own previous skip responses in the immediately preceding self-initiated block, in a

randomised order. For participants who started with the externally-triggered block, the timing

of the red colour appearance in the first block only was yoked to the time of previous

participant's last self-initiated block. The colour cycle of the fixation cross had a random

sequence, so that the onset of a red fixation could not be predicted. The fixation cross

ramped to 'red' from its previous colour in 300 ms. Again, a small reward (1p) was given for

skipping. The trial finished and the participant lost money (-1p) if s/he did not skip within 2.5

s from beginning of the ramping colour of the fixation cross. The 'red' colour was left out of

the colour cycle in the self-initiated blocks. To control for any confounding effect of attending

to the fixation cross, participants were also required to attend to the fixation cross in the self-

initiated blocks and to roughly estimate the number of times the fixation cross turned 'yellow'

(see previous). Each externally-triggered block had 10 trials, and after each block feedback

was displayed. Each self-initiated block was interleaved with an externally-triggered block,

and the order of the blocks was counterbalanced between the participants. The behavioural

task was designed in Psychophysics Toolbox Version 3 (Brainard, 1997).

EEG recording.

While participants were performing the behavioural task in a shielded chamber, EEG signals

were recorded and amplified using an ActiveTwo Biosemi system (BioSemi, Amsterdam,

The Netherlands). Participants wore a 64-channel EEG cap. To shorten the preparation time

we recorded from a subset of electrodes that mainly covers central and visual areas: F3, Fz,

F4, FC1, FCz, FC2, C3, C1, Cz, C2, C4, CP1, CPz, CP2, P3, Pz, P4, O1, Oz, O2. Bipolar

channels placed on the outer canthi of each eye and below and above the right eye were

used to record horizontal and vertical electro-oculogram (EOG), respectively. The Biosemi

Active electrode has an output impedance of less than 1 Ohm. EEG signals were recorded

at a sampling rate of 2048 Hz.

EEG preprocessing.

EEG data preprocessing was performed in Matlab (MathWorks, MA, USA) with the help of

EEGLAB toolbox (Delorme & Makeig, 2004). Data were downsampled to 250 Hz and low-

pass filtered at 30 Hz. No high-pass filtering and no detrending were applied, to preserve

slow fluctuations. All electrodes were referenced to the average of both mastoid electrodes.

Separate data epochs of 4 s duration were extracted for self-initiated and externally-

triggered skip actions. Data epochs started from 3 s before to 1 s after the action. To avoid

EEG epochs overlapping each other any trial in which participants skipped earlier than 3 s

from trial initiation was removed. On average, 5% and 4% of trials were removed from the

self-initiated and externally-triggered conditions, respectively.

RP recordings are conventionally baseline-corrected using a baseline 2.5 until 2 s before

action. This involves the implicit assumption that RPs begin only in the 2 s before action

onset (Shibasaki & Hallett, 2006), but this assumption is rarely articulated explicitly, and is in

fact questionable (Verbaarschot, Farquhar, & Haselager, 2015). We instead took a baseline

from -5 ms +5 ms with respect to action onset. This choice avoids making any assumption

about how or when the RP starts. To ensure this choice of baseline did not capitalize on

chance, we performed parallel analyses on demeaned data (effectively taking the entire

epoch as baseline), with consistent results (see Figure. S2). Finally, to reject non-ocular

artefacts, data epochs from EEG channels (not including EOG) with values exceeding a

threshold of ±150 μv were removed. On average 7% and 8% of trials were rejected from

self-initiated and externally-triggered conditions, respectively. In the next step, Independent

Component Analysis (ICA) was used to remove ocular artefacts from the data. Ocular ICA

components were identified by visual inspection. Trials with artefacts remaining after this

procedure were excluded by visual inspection.

EEG analysis.

Preliminary inspection showed a typical RP-shaped negative-going slow component that

was generally maximal at FCz. Therefore, data from FCz was chosen for subsequent

analysis. Time series analysis was performed in Matlab (MathWorks) with the help of the

FieldTrip toolbox (Oostenveld et al., 2010). We measured two dependent variables as

precursors of both *self-initiated* and *externally-triggered* skip actions: mean RP amplitude

across trials and variability of RP amplitudes across and within trials (measured by SD). To

compare across-trials SD between the two conditions, data epochs were divided into four

500 ms windows, starting 2 s before action onset: [-2, -1.5 s], [-1.5, -1 s], [-1, -0.5 s], [-0.5, 0

s]. All p-values were Bonferroni corrected for four comparisons. To get a precise estimate of

the standard error of the difference between conditions, paired-samples t-tests were

performed on jack-knifed data (Efron & Stein, 1981; Kiesel, Miller, Jolicoeur, & Brisson,

2008). Unlike the traditional methods, this technique compares variation of interest across

subsets of the total sample rather than across individuals, by temporarily leaving each

subject out of the calculation. In addition, we also performed cluster-based permutation tests

on SD (Maris & Oostenveld, 2007). These involve a priori identification of a set of electrodes

and a time-window of interest, and incorporate appropriate corrections for multiple

comparisons. Importantly, they avoid further arbitrary assumptions associated with selecting

specific sub-elements of the data of interest, such as individual electrodes, time-bins or ERP

components. The cluster-based tests were performed using the following parameters: time

interval = [-2 - 0 s relative to action], minimum number of neighbouring electrodes required =

2, number of draws from the permutation distribution = 1000.

To measure variability of RP amplitudes within each individual trial, the SD of the EEG signal

from FCz was measured across time in a 100 ms window. This window was applied

successively in 30 time bins from the beginning of the epoch (3 s prior to action) to the time

of action onset. We used linear regression to calculate the slope of the within-trial SD as a

function of time (Figure. 6A). This was performed separately for each trial and each

participant. Slopes greater than 0 indicate that EEG within the 100 ms window becomes

more variable with the approach to action onset. Finally, we compared slopes of this within-

trial SD measure between self-initiated and externally-triggered conditions in a multilevel

model with single trials as level 1 and participants as level 2 variables. Multilevel analysis

was performed in R (R Core Team, Vienna, Austria).

Time-frequency analysis was performed with custom written Matlab scripts. The

preprocessed EEG time series were decomposed into their time-frequency representation by

using Complex Morlet wavelet with 20 frequencies, ranging linearly from 5 to 30 Hz. The

number of wavelet cycles increased from 3 to 7 in the same number of steps used to

increase the frequency of the wavelets from 5 to 30 Hz. Power at each trial, each frequency

and each time point was measured by convolving the raw time series with the wavelet and

squaring the resulting complex number. The power at each frequency and each time point

was then averaged across trials for each participant. Edge artefacts were removed by

discarding the first and last 500 ms of the epoch. Baseline time window was defined as the

first 500 ms of the epoch (after removal of edge artefacts: 2.5 - 2 s prior to skip action).

Changes in power during action preparation was subsequently expressed as the percentage

of change relative to the average power during the baseline time window, across time at a

specific frequency. Baseline normalisation was performed by using the following equation:

$$percentage_{tf} = 100 \; \frac{power_{tf} - \overline{baseline_f}}{\overline{baseline_f}}$$

Values > 0 indicates that power at a specific frequency (f) and a specific time (t) is higher

relative to the average power at the same frequency during the first 500 ms of the epoch.

Finally, we asked whether percentage change in power relative to baseline differs between

self-initiated and externally-triggered skip conditions in the beta band (15 – 30 Hz). Beta

band Event-related Desynchronization (ERD) during action preparation is a well-established

phenomenon (Bai, Mari, Vorbach, & Hallett, 2005; Doyle, Yarrow, & Brown, 2005;

Pfurtscheller & Lopes da Silva, 1999). Beta power was calculated in a 500 ms window

starting from 1 s and ending 0.5 s prior to skip action. We avoided analysing later windows

(e.g., 0.5 - 0 s prior to action) to avoid possible contamination from action execution following

presentation of the red fixation cross that cued externally-triggered responses. The average

normalised power across all pixels within the selected window was then calculated for each

participant and compared across conditions using paired-samples t-tests.

Modelling and simulations.

All simulations were done in Matlab (MathWorks). We used a modified version of the Leaky

Stochastic Accumulator Model (Usher & McClelland, 2001), in which the activity of

accumulators increases stochastically over time but is limited by leakage.

$$\delta x = (I - kx)\Delta t + c\xi\sqrt{\Delta t}$$

Where I is drift rate, k is leak (exponential decay in x), ξ is Gaussian noise, c is noise scaling

factor, and Δt is the discrete time step (we used $\Delta t = 0.001$). This leaky stochastic

accumulator has been used previously to model the neural decision of 'when' to move in a

self-initiated task (Schurger et al., 2012). In that experiment, I was defined as the general

imperative to respond (with a constant rate). This imperative, if appropriately small in

magnitude, moves the baseline level of activity closer to the threshold, but not over it. Thus,

imperative alone does not trigger action, but does increase the likelihood of a random

threshold-crossing event triggering action. In the original model, c was assumed to be

constant and was fixed at 0.1. In a departure from the original model, we assumed that the

noise scaling factor could change linearly from an initial value of c_1 to a final value of c_2,

during action preparation. Consequently, Δc was defined as the magnitude of change in the

noise scaling factor during the trial.

$$\Delta c = c_2 - c_1$$

664   A negative Δc means that signal becomes less noisy as it approaches the threshold for

665   action. Therefore, the modified model in our experiment had five free parameters: I, k, c_1, c_2

666   and threshold.

$$\delta x = (I - kx)\Delta t + c_t\xi\sqrt{\Delta t}$$

Where c_t is noise scaling factor at time t. The threshold was expressed as a percentile of the

output amplitude over a set of 1000 simulated trials (each of 50,000 time steps each).

Epochs of simulated data were matched to epochs of actual EEG data by identifying the

point of first threshold crossing event within each simulated trial and then extracting an

epoch from 3000 time steps before to 1000 time steps after the threshold crossing.

Parameter estimation for self-initiated skip action was performed by fitting the model against

the *real mean* RP amplitude of each participant in self-initiated condition. First, 1000 unique

trials of Gaussian noise, each 50,000 time steps, were generated for each participant and

were fed into the model. The initial values of the model's parameters were derived from

previous studies (Schurger et al., 2012). The output of the model was then averaged across

trials and was down sampled to 250 Hz to match the sampling rate of the real EEG data. A

least squares approach was used to minimise root mean squared deviation (RMSD)

between the *simulated* and *real* mean RP, by adjusting the free parameters of the model for

each participant (by using the MATLAB 'fminsearch' function). Note that this procedure

optimised the model parameters to reproduce the mean RP, rather than individual trials.

To fit the model to our externally-triggered skip condition, we fixed the threshold of each

participant at their best fitting threshold from the self-initiated condition. We wanted to keep

the threshold the same in both conditions so that we could test the effect of changing noise

levels for a given threshold. Importantly, we also fixed the value of c_1 at its optimal value

form the self-initiated condition. By using this strategy, we can ask how noisiness of the

signal *changes*, from its initial value, and we can compare this change in noise between

conditions. We additionally performed parallel simulations without the assumption of a

common initial noise level, and obtained essentially similar results. Specifically Δc in the all-

parameter-free model (mean= 0.02, SD = 0.06) was similar to the Δc in the model with c_1

and threshold fixed (mean= 0.02, SD = 0.05). The remaining parameters (I, k, c_2) were

optimised by minimising the deviation between the *simulated mean* RP and the *real mean*

RP in externally-triggered condition.

Finally, we tested the model on the across-trial variability of RP epochs, having *fitted* the

model parameters to the mean RP. All parameters of the model were fixed at each

participant's optimised values for the self-initiated condition, and for the externally-triggered

condition respectively. The model was run 44 times (22 participants, x 2 conditions) with the

appropriate parameters, and 1000 separate trials were generated, each corresponding to a

putative RP exemplar. The Gaussian noise element of the model ensured that these 1000

exemplars were non-identical. The standard deviation across trials was calculated from

these 1000 simulated RP exemplars, for each participant and each condition. Importantly,

this procedure fits the model to each participant's *mean* RP amplitude, but then tests the fit

on the *standard deviation* across the 1000 simulated trials. Finally, to assess similarity

between the real and predicted SD reduction, the predicted SD in self-initiated and

externally-triggered conditions was plotted as a function of time and the area between the

two curves was computed. We then compared the area between the SD curves in a 2 s

interval prior to self-initiated and externally-triggered conditions for all participants' simulated

data, and actual data (Figure 5), using Pearson's correlation.

Author Contributions

710   Conceptualization, N.K., P.H., A.S., and A.D.; Methodology, N.K., A.S., and A.D.; Formal

711   Analysis, N.K., L.Z., and P.H.; Investigation, N.K., L.Z.; Writing-original draft, N.K., Writing-

712   review & editing, P.H., A.S., Supervision, P.H. and A.S.

Acknowledgement

714   This work was supported by the European Research Council Advanced Grant HUMVOL

715   (Grant number: 323943). This work was additionally supported by a grant from

716   The Leverhulme Trust (Ref. RPG-2016-378) to PH. AS was supported by a starting grant

717   from the European Research Council (Grant number 640626). The authors declare no

718   competing financial interest.

References

Anscombe, G. E. M. (2000). *Intention*. Harvard University Press.

Averbeck, B. B., & Lee, D. (2003). Neural noise and movement-related codes in the macaque supplementary motor area. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *23*(20), 7630–7641.

Bai, O., Mari, Z., Vorbach, S., & Hallett, M. (2005). Asymmetric spatiotemporal patterns of event-related desynchronization preceding voluntary sequential finger movements: a high-resolution EEG study. *Clinical Neurophysiology: Official Journal of the International Federation of Clinical Neurophysiology*, *116*(5), 1213–1221. https://doi.org/10.1016/j.clinph.2005.01.006

Bai, O., Rathi, V., Lin, P., Huang, D., Battapady, H., Fei, D.-Y., … Hallett, M. (2011). Prediction of human voluntary movement before it occurs. *Clinical Neurophysiology: Official Journal of the International Federation of Clinical Neurophysiology*, *122*(2), 364–372. https://doi.org/10.1016/j.clinph.2010.07.010

Boschert, J., Hink, R. F., & Deecke, L. (1983). Finger movement versus toe movement-related potentials: further evidence for supplementary motor area (SMA) participation prior to voluntary action. *Experimental Brain Research*, *52*(1), 73–80.

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

Calmels, C., Holmes, P., Jarry, G., Lévèque, J.-M., Hars, M., & Stam, C. J. (2006). Cortical activity prior to, and during, observation and execution of sequential finger movements. *Brain Topography*, *19*(1–2), 77–88. https://doi.org/10.1007/s10548-006-0014-x

Casini, L., & Vidal, F. (2011). The SMAs: neural substrate of the temporal accumulator? *Frontiers in Integrative Neuroscience*, 35. https://doi.org/10.3389/fnint.2011.00035

Churchland, A. K., Kiani, R., Chaudhuri, R., Wang, X.-J., Pouget, A., & Shadlen, M. N. (2011). Variance as a signature of neural computations during decision making. *Neuron*, *69*(4), 818–831. https://doi.org/10.1016/j.neuron.2010.12.037

Churchland, M. M., Yu, B. M., Cunningham, J. P., Sugrue, L. P., Cohen, M. R., Corrado, G. S., … Shenoy, K. V. (2010). Stimulus onset quenches neural variability: a widespread cortical phenomenon. *Nature Neuroscience*, *13*(3), 369–378. https://doi.org/10.1038/nn.2501

Churchland, M. M., Yu, B. M., Ryu, S. I., Santhanam, G., & Shenoy, K. V. (2006). Neural Variability in Premotor Cortex Provides a Signature of Motor Preparation. *The Journal of Neuroscience*, *26*(14), 3697–3712. https://doi.org/10.1523/JNEUROSCI.3762-05.2006

Cohen, M. X. (2014). *Analyzing Neural Time Series Data: Theory and Practice* (1 edition). Cambridge, Massachusetts: MIT Press.

Constantino, S. M., & Daw, N. D. (2015). Learning the opportunity cost of time in a patch-foraging task. *Cognitive, Affective & Behavioral Neuroscience*, *15*(4), 837–853. https://doi.org/10.3758/s13415-015-0350-y

Cunnington, R., Windischberger, C., Deecke, L., & Moser, E. (2002). The preparation and execution of self-initiated and externally-triggered movement: a study of event-related fMRI. *NeuroImage*, *15*(2), 373–385. https://doi.org/10.1006/nimg.2001.0976

Deecke, L., & Kornhuber, H. H. (1978). An electrical sign of participation of the mesial "supplementary" motor cortex in human voluntary finger movement. *Brain Research*, *159*(2), 473–476.

Defebvre, L., Bourriez, J. L., Destée, A., & Guieu, J. D. (1996). Movement related desynchronisation pattern preceding voluntary movement in untreated Parkinson's disease. *Journal of Neurology, Neurosurgery & Psychiatry*, *60*(3), 307–312. https://doi.org/10.1136/jnnp.60.3.307

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Dennett, D. C. (2015). *Elbow Room: The Varieties of Free Will Worth Wanting* (New edition). Cambridge, Massachusetts ; London, England: MIT Press.

Desantis, A., Waszak, F., & Gorea, A. (2016). Agency alters perceptual decisions about action-outcomes. *Experimental Brain Research*. https://doi.org/10.1007/s00221-016-4684-7

Desantis, A., Waszak, F., Moutsopoulou, K., & Haggard, P. (2016). How action structures time: About the perceived temporal order of action and predicted outcomes. *Cognition*, *146*, 100–109. https://doi.org/10.1016/j.cognition.2015.08.011

Dirnberger, G., Lang, W., & Lindinger, G. (2008). A new method to determine temporal variability in the period of pre-movement electroencephalographic activity. *International Journal of Psychophysiology: Official Journal of the International Organization of Psychophysiology*, *70*(3), 165–170. https://doi.org/10.1016/j.ijpsycho.2008.08.006

Doyle, L. M. F., Yarrow, K., & Brown, P. (2005). Lateralization of event-related beta desynchronization in the EEG during pre-cued reaction time tasks. *Clinical Neurophysiology: Official Journal of the International Federation of Clinical Neurophysiology*, *116*(8), 1879–1888. https://doi.org/10.1016/j.clinph.2005.03.017

Efron, B., & Stein, C. (1981). The Jackknife Estimate of Variance. *The Annals of Statistics*, *9*(3), 586–596.

Faisal, A. A., Selen, L. P. J., & Wolpert, D. M. (2008). Noise in the nervous system. *Nature Reviews Neuroscience*, *9*(4), 292–303. https://doi.org/10.1038/nrn2258

Fetsch, C. R., Kiani, R., Newsome, W. T., & Shadlen, M. N. (2014). Effects of Cortical Microstimulation on Confidence in a Perceptual Decision. *Neuron*, *83*(4), 797–804. https://doi.org/10.1016/j.neuron.2014.07.011

Fitts, P. M. (1954). The information capacity of the human motor system in controlling the amplitude of movement. *Journal of Experimental Psychology*, *47*(6), 381–391. https://doi.org/10.1037/h0055392

Fried, I., Mukamel, R., & Kreiman, G. (2011). Internally generated preactivation of single neurons in human medial frontal cortex predicts volition. *Neuron*, *69*(3), 548–562. https://doi.org/10.1016/j.neuron.2010.11.045

Furstenberg, A., Breska, A., Sompolinsky, H., & Deouell, L. Y. (2015). Evidence of Change of Intention in Picking Situations. *Journal of Cognitive Neuroscience*, 1–14. https://doi.org/10.1162/jocn_a_00842

Gold, J. I., & Shadlen, M. N. (2007). The Neural Basis of Decision Making. *Annual Review of Neuroscience*, *30*(1), 535–574. https://doi.org/10.1146/annurev.neuro.29.051605.113038

Groen, O. van der, & Wenderoth, N. (2016). Transcranial Random Noise Stimulation of Visual Cortex: Stochastic Resonance Enhances Central Mechanisms of Perception. *Journal of Neuroscience*, *36*(19), 5289–5298. https://doi.org/10.1523/JNEUROSCI.4519-15.2016

Haggard, P. (2005). Conscious intention and motor cognition. *Trends in Cognitive Sciences*, *9*(6), 290–295. https://doi.org/10.1016/j.tics.2005.04.012

Haggard, P. (2008). Human volition: towards a neuroscience of will. *Nature Reviews. Neuroscience*, *9*(12), 934–946. https://doi.org/10.1038/nrn2497

He, B. J. (2013). Spontaneous and task-evoked brain activity negatively interact. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *33*(11), 4672–4682. https://doi.org/10.1523/JNEUROSCI.2922-12.2013

Jahanshahi, M., Jenkins, I. H., Brown, R. G., Marsden, C. D., Passingham, R. E., & Brooks, D. J. (1995). Self-initiated versus externally triggered movements. I. An investigation using measurement of regional cerebral blood flow with PET and movement-related potentials in normal and Parkinson's disease subjects. *Brain: A Journal of Neurology*, *118 (Pt 4)*, 913–933.

Keller, I., & Heckhausen, H. (1990). Readiness potentials preceding spontaneous motor acts: voluntary vs. involuntary control. *Electroencephalography and Clinical Neurophysiology*, *76*(4), 351–361.

Kiani, R., Hanks, T. D., & Shadlen, M. N. (2008). Bounded Integration in Parietal Cortex Underlies Decisions Even When Viewing Duration Is Dictated by the Environment. *Journal of Neuroscience*, *28*(12), 3017–3029. https://doi.org/10.1523/JNEUROSCI.4761-07.2008

Kiesel, A., Miller, J., Jolicoeur, P., & Brisson, B. (2008). Measurement of ERP latency differences: a comparison of single-participant and jackknife-based scoring methods. *Psychophysiology*, *45*(2), 250–274. https://doi.org/10.1111/j.1469-8986.2007.00618.x

Kolling, N., Behrens, T. E. J., Mars, R. B., & Rushworth, M. F. S. (2012). Neural mechanisms of foraging. *Science (New York, N.Y.)*, *336*(6077), 95–98. https://doi.org/10.1126/science.1216930

Kool, W., & Botvinick, M. (2013). The intrinsic cost of cognitive control. *The Behavioral and Brain Sciences*, *36*(6), 697-698-726. https://doi.org/10.1017/S0140525X1300109X

Kornhuber, H. H., & Deecke, L. (1965). Hirnpotentialänderungen bei Willkürbewegungen und passiven Bewegungen des Menschen: Bereitschaftspotential und reafferente Potentiale. *Pflüger's Archiv für die gesamte Physiologie des Menschen und der Tiere*, *284*(1), 1–17. https://doi.org/10.1007/BF00412364

Kornhuber, H. H., & Deecke, L. (1990). Readiness for movement - The Bereitschaftspotential story. Citation Classic Commentary. *Current Contents Life Sciences*, *33*(4).

Krieghoff, V., Waszak, F., Prinz, W., & Brass, M. (2011). Neural and behavioral correlates of intentional actions. *Neuropsychologia*, *49*(5), 767–776. https://doi.org/10.1016/j.neuropsychologia.2011.01.025

Kristeva, R., Patino, L., & Omlor, W. (2007). Beta-range cortical motor spectral power and corticomuscular coherence as a mechanism for effective corticospinal interaction during steady-state motor output. *NeuroImage*, *36*(3), 785–792. https://doi.org/10.1016/j.neuroimage.2007.03.025

Libet, B., Gleason, C. A., Wright, E. W., & Pearl, D. K. (1983). Time of conscious intention to act in relation to onset of cerebral activity (readiness-potential). The unconscious initiation of a freely voluntary act. *Brain: A Journal of Neurology*, *106 (Pt 3)*, 623–642.

Luck, S. J. (2005). *An Introduction to the Event-Related Potential Technique*. Cambridge, Mass: MIT Press.

Manohar, S. G., Chong, T. T.-J., Apps, M. A. J., Batla, A., Stamelou, M., Jarman, P. R., … Husain, M. (2015). Reward Pays the Cost of Noise Reduction in Motor and Cognitive Control. *Current Biology: CB*, *25*(13), 1707–1716. https://doi.org/10.1016/j.cub.2015.05.038

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024

Murakami, M., Vicente, M. I., Costa, G. M., & Mainen, Z. F. (2014). Neural antecedents of self-initiated actions in secondary motor cortex. *Nature Neuroscience.* https://doi.org/10.1038/nn.3826

Nachev, P., & Hacker, P. (2014). The neural antecedents to voluntary action: a conceptual analysis. *Cognitive Neuroscience*, *5*(3–4), 193–208. https://doi.org/10.1080/17588928.2014.934215

Nachev, P., Kennard, C., & Husain, M. (2008). Functional role of the supplementary and pre-supplementary motor areas. *Nature Reviews Neuroscience*, *9*(11), 856–869. https://doi.org/10.1038/nrn2478

Oostenveld, R., Fries, P., Maris, E., Schoffelen, J.-M., Oostenveld, R., Fries, P., … Schoffelen, J.-M. (2010). FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Computational Intelligence and Neuroscience*, *2011*, e156869. https://doi.org/10.1155/2011/156869

Papa, S. M., Artieda, J., & Obeso, J. A. (1991). Cortical activity preceding self-initiated and externally triggered voluntary movement. *Movement Disorders: Official Journal of the Movement Disorder Society*, *6*(3), 217–224. https://doi.org/10.1002/mds.870060305

Passingham, R. (1995). *The Frontal Lobes and Voluntary Action*. Oxford University Press.

Passingham, R. E., Bengtsson, S. L., & Lau, H. C. (2010). Medial frontal cortex: from self-generated action to reflection on one's own performance. *Trends in Cognitive Sciences*, *14*(1), 16–21. https://doi.org/10.1016/j.tics.2009.11.001

Pereboom, D. (2011). Free Will Skepticism and Meaning in Life. In R. Kane (Ed.), *The Oxford Handbook of Free Will*. Oxford University Press.

Pfurtscheller, G. (1992). Event-related synchronization (ERS): an electrophysiological correlate of cortical areas at rest. *Electroencephalography and Clinical Neurophysiology*, *83*(1), 62–69.

Pfurtscheller, G., & Lopes da Silva, F. H. (1999). Event-related EEG/MEG synchronization and desynchronization: basic principles. *Clinical Neurophysiology: Official Journal of the International Federation of Clinical Neurophysiology*, *110*(11), 1842–1857.

Pfurtscheller, G., & Neuper, C. (1994). Event-related synchronization of mu rhythm in the EEG over the cortical hand area in man. *Neuroscience Letters*, *174*(1), 93–96.

Saberi-Moghadam, S., Ferrari-Toniolo, S., Ferraina, S., Caminiti, R., & Battaglia-Mayer, A. (2016). Modulation of Neural Variability in Premotor, Motor, and Posterior Parietal Cortex during Change of Motor Intention. *The Journal of Neuroscience: The Official*

Journal of the Society for Neuroscience, *36*(16), 4614–4623.
https://doi.org/10.1523/JNEUROSCI.3300-15.2016

Schurger, A., Sarigiannidis, I., Naccache, L., Sitt, J. D., & Dehaene, S. (2015). Cortical
activity is more stable when sensory stimuli are consciously perceived. *Proceedings
of the National Academy of Sciences of the United States of America*, *112*(16),
E2083-2092. https://doi.org/10.1073/pnas.1418730112

Schurger, A., Sitt, J. D., & Dehaene, S. (2012). An accumulator model for spontaneous
neural activity prior to self-initiated movement. *Proceedings of the National Academy
of Sciences*, *109*(42), E2904–E2913. https://doi.org/10.1073/pnas.1210467109

Schüür, F., & Haggard, P. (2011). What are self-generated actions? *Consciousness and
Cognition*, *20*(4), 1697–1704. https://doi.org/10.1016/j.concog.2011.09.006

Shadlen, M. N., & Gold, J. I. (2004). In M. S. Gazzaniga (Ed.), *The Cognitive Neurosciences*
(3rd edn, pp. 1229–1241). MIT Press.

Shenoy, K. V., Sahani, M., & Churchland, M. M. (2013). Cortical control of arm movements:
a dynamical systems perspective. *Annual Review of Neuroscience*, *36*, 337–359.
https://doi.org/10.1146/annurev-neuro-062111-150509

Sherrington, C. S. (1906). *The integrative action of the nervous system*. New York, C
Scribner's sons. Retrieved from http://archive.org/details/integrativeacti02shergoog

Shibasaki, H., & Hallett, M. (2006). What is the Bereitschaftspotential? *Clinical
Neurophysiology: Official Journal of the International Federation of Clinical
Neurophysiology*, *117*(11), 2341–2356. https://doi.org/10.1016/j.clinph.2006.04.025

Shu, Y., Hasenstaub, A., Badoual, M., Bal, T., & McCormick, D. A. (2003). Barrages of
synaptic activity control the gain and sensitivity of cortical neurons. *The Journal of
Neuroscience: The Official Journal of the Society for Neuroscience*, *23*(32), 10388–
10401.

Sinnott-Armstrong, W., & Nadel, L. (Eds.). (2010). *Conscious Will and Responsibility: A
Tribute to Benjamin Libet* (1 edition). Oxford ; New York: OUP USA.

Stancák, A., & Pfurtscheller, G. (1996). Event-related desynchronisation of central beta-
rhythms during brisk and slow self-paced finger movements of dominant and
nondominant hand. *Brain Research. Cognitive Brain Research*, *4*(3), 171–183.

Stein, R. B., Gossen, E. R., & Jones, K. E. (2005). Neuronal variability: noise or part of the
signal? *Nature Reviews Neuroscience*, *6*(5), 389–397.
https://doi.org/10.1038/nrn1668

Tzagarakis, C., Ince, N. F., Leuthold, A. C., & Pellizzer, G. (2010). Beta-band activity during
motor planning reflects response uncertainty. *The Journal of Neuroscience: The
Official Journal of the Society for Neuroscience*, *30*(34), 11270–11277.
https://doi.org/10.1523/JNEUROSCI.6026-09.2010

Usher, M., & McClelland, J. L. (2001). The time course of perceptual choice: the leaky, competing accumulator model. *Psychological Review*, *108*(3), 550–592.

Van Rijn, H., Kononowicz, T. W., Meck, W. H., Ng, K. K., & Penney, T. B. (2011). Contingent negative variation and its relation to time estimation: a theoretical evaluation. *Frontiers in Integrative Neuroscience*, *5*, 91. https://doi.org/10.3389/fnint.2011.00091

Verbaarschot, C., Farquhar, J., & Haselager, P. (2015). Lost in time...: The search for intentions and Readiness Potentials. *Consciousness and Cognition*, *33C*, 300–315. https://doi.org/10.1016/j.concog.2015.01.011

Walter, W. G., Cooper, R., Aldridge, V. J., McCALLUM, W. C., & Winter, A. L. (1964). Contingent Negative Variation : An Electric Sign of Sensori-Motor Association and Expectancy in the Human Brain. *Nature*, *203*(4943), 380–384. https://doi.org/10.1038/203380a0

Wiese, H., Stude, P., Nebel, K., de Greiff, A., Forsting, M., Diener, H. C., & Keidel, M. (2004). Movement preparation in self-initiated versus externally triggered movements: an event-related fMRI-study. *Neuroscience Letters*, *371*(2–3), 220–225. https://doi.org/10.1016/j.neulet.2004.08.078

Zylberberg, A., Fetsch, C. R., & Shadlen, M. N. (2016). The influence of evidence volatility on choice, reaction time and confidence in a perceptual decision. *eLife*, *5*. https://doi.org/10.7554/eLife.17688

Supplementary Figures & Tables

Figure S1. Histogram of waiting times before skip actions in self-initiated condition, across all trials and all participants. The dashed red line shows the average waiting time.

Figure S2. Demeaned EEG activity prior to skip actions. The red and blue lines represent self-initiated and externally-triggered skips, respectively. Data is time-locked to the skip action (black vertical line), and is baselined to the mean of entire epoch (i.e., demeaned), and recorded from FCz electrode. A. Grand average RP amplitude. B. Standard deviation across trials averaged across participants. Shaded area show significant clusters across central electrodes, detected by cluster-based permutation test. Whereas baselining to a limited time window forces a low SD within the baseline time window, and a progressive rise in SD with temporal distance before or after the baseline, the use of a broad baseline time window, as here, reduces this artefactual effect of baseline-correction on variability of time-locked data. Nevertheless, the difference in SD between conditions remains significant.

Figure S3. Topography of the difference in SDs between self-initiated and externally-triggered conditions. Small circles represent EEG electrodes across which the permutation test was performed. Electrodes that showed significant difference between conditions have been marked *. The time interval (s) is indicated above each subplot.

Figure S4. Correlation between participants' mean waiting time (s) before skipping in the self-initiated condition and EEG convergence. EEG convergence was measured by subtracting the area under the SD curve in self-initiated from the externally-triggered condition. There was a significant negative correlation between waiting time and EEG convergence (Pearson's r = -0.66, p < 0.01).

Fig S5. No significant correlation (Pearson's r = 0.16, p = 0.46) across participants between standard deviation of each participant's RT to externally-triggered skip cues (ms), and EEG convergence. EEG convergence was measured by subtracting the area under the SD curve in self-initiated from the externally-triggered condition.

Subject	Mean wait (s) self-initiated	Mean wait (s) externally-triggered	SD (s) self-initiated	SD (s) externally-triggered
1	11.76	12.09	6.82	6.79
2	5.38	5.76	2.27	2.30
3	6.14	6.45	4.58	4.61
4	7.16	7.45	3.25	3.22
5	8.25	8.39	4.57	3.97
6	7.21	7.70	3.22	3.27
7	5.92	6.20	1.66	1.67
8	7.83	8.16	3.20	3.23
9	6.08	6.48	1.26	1.20
10	6.32	6.54	2.96	2.68
11	5.34	5.85	1.00	1.01
12	6.89	7.28	2.72	2.70
13	4.83	5.23	1.83	1.82
14	8.06	8.23	2.66	2.54
15	8.97	9.44	3.60	3.60
16	9.25	9.66	5.85	5.91
17	6.30	6.81	2.69	2.61
18	7.72	8.14	2.53	2.57
19	6.90	7.16	2.94	2.93
20	7.32	7.80	2.40	2.62
21	9.44	9.85	2.93	2.94
22	7.14	7.73	4.74	5.19

Table S1. Mean and standard deviation of waiting time before skipping in self-initiated and externally-triggered conditions.

Drift (l)	Leak (k)	Initial noise (c_1)	Final noise (c_2)	Threshold	Change in noise (Δc)
0.04	-0.18	0.08	0.01	0.10	-0.07
0.11	0.55	0.12	0.13	0.14	0.00
0.11	0.57	0.13	0.11	0.18	-0.02
0.11	0.52	0.12	0.12	0.17	0.00
0.09	0.23	0.19	0.14	0.41	-0.05
0.01	0.38	0.23	0.13	0.42	-0.10
0.10	0.50	0.14	0.13	0.20	-0.01
0.12	0.62	0.10	0.14	0.08	0.04
0.11	0.57	0.11	0.13	0.11	0.01
0.02	0.27	0.25	0.12	0.44	-0.13
0.08	0.14	0.21	0.13	0.38	-0.08
0.11	0.53	0.12	0.12	0.13	0.01
0.09	0.11	0.21	0.12	0.35	-0.09
0.06	0.17	0.21	0.12	0.39	-0.09
0.07	0.28	0.19	0.14	0.36	-0.05
0.11	0.53	0.12	0.11	0.12	-0.01
0.11	0.54	0.12	0.12	0.13	0.01
0.08	0.22	0.18	0.12	0.29	-0.06
0.11	0.60	0.12	0.13	0.09	0.01
0.11	0.53	0.12	0.12	0.13	0.01
0.17	0.59	0.18	0.24	0.74	0.06
0.11	0.55	0.13	0.12	0.15	0.00

Table S2. Optimum parameters for self-initiated skip action. The values were detected by fitting the model against the mean RP amplitude of each participant in self-initiated condition. Δc was measured by subtracting the initial noise level (c_1) from the final noise level (c_2).

Drift (l)	Leak (k)	Initial noise (c_1)	Final noise (c_2)	Threshold	Change in noise (Δc)
0.09	0.42	0.08	0.10	0.10	0.02
0.11	1.01	0.12	0.21	0.14	0.09
0.10	0.35	0.13	0.13	0.18	0.00
0.16	0.70	0.12	0.16	0.17	0.04
0.12	0.48	0.19	0.18	0.41	-0.01
0.10	0.48	0.23	0.17	0.42	-0.05
0.09	0.26	0.14	0.07	0.20	-0.06
0.17	0.93	0.10	0.24	0.08	0.13
0.18	0.44	0.11	0.19	0.11	0.08
0.05	0.44	0.25	0.22	0.44	-0.03
0.10	0.38	0.21	0.17	0.38	-0.04
0.19	0.72	0.12	0.15	0.13	0.03
0.10	0.41	0.21	0.18	0.35	-0.03
0.08	0.54	0.21	0.17	0.39	-0.04
0.12	0.51	0.19	0.23	0.36	0.05
0.16	0.87	0.12	0.15	0.12	0.04
0.18	0.69	0.12	0.16	0.13	0.04
0.10	0.45	0.18	0.15	0.29	-0.03
0.21	0.75	0.12	0.17	0.09	0.05
0.17	0.77	0.12	0.20	0.13	0.09
0.10	0.61	0.18	0.20	0.74	0.01
0.16	0.58	0.13	0.16	0.15	0.03

Table S3. Optimum parameters for externally-triggered skip action. The values were detected by fitting the model against the mean RP amplitude of each participant in externally-triggered condition. Δc was measured by subtracting the initial noise level (c_1) from the final noise level (c_2). c_1 and the threshold were fixed at their optimum values in self-initiated condition (see materials and methods for more details)