Contribution of *trans* regulatory eQTL to cryptic genetic variation in *C. elegans*

L. Basten Snoek[1,3], Mark G. Sterken[1,3], Roel P. J. Bevers[1], Rita J. M. Volkers[1], Arjen van 't Hof[2], Rachel

Brenchley[2], Joost A. G. Riksen[1], Andrew Cossins[2], Jan E. Kammenga[1,4]

[1]Laboratory of Nematology, Wageningen University and Research, 6708 PB, The Netherlands.

[2]Centre for Genome research, Institute of Integrative Biology, Biosciences Building, University of Liverpool,

Liverpool L69 7ZB, United Kingdom.

[3]Authors contributed equally

[4]Corresponding author

Keywords:

Caenorhabditis elegans, genetical genomics, eQTL, heat stress, cryptic genetic variation, *trans*-band, eQTL

hotspot

Corresponding author:

Jan E. Kammenga

Wageningen University and Research, laboratory of Nematology

Droevendaalsesteeg 1, 6708PB, Wageningen, The Netherlands

0031317482998

Jan.kammenga@wur.nl

Abstract

Background

Cryptic genetic variation (CGV) is the hidden genetic variation that can be unlocked by perturbing normal

conditions. CGV can drive the emergence of novel complex phenotypes through changes in gene expression.

Although our theoretical understanding of CGV has thoroughly increased over the past decade, insight into

polymorphic gene expression regulation underlying CGV is scarce. Here we investigated the transcriptional

architecture of CGV in response to rapid temperature changes in the nematode *Caenorhabditis elegans*. We

analyzed regulatory variation in gene expression (and mapped eQTL) across the course of a heat stress and

recovery response in a recombinant inbred population.

Results

We measured gene expression over three temperature treatments: i) control, ii) heat stress, and iii) recovery from

heat stress. Compared to control, exposure to heat stress affected the transcription of 3305 genes, whereas 942

were affected in recovering animals. These affected genes were mainly involved in metabolism and

reproduction. The gene expression pattern in recovering animals resembled both the control and the heat-stress

treatment. We mapped eQTL using the genetic variation of the recombinant inbred population and detected 2626

genes with an eQTL in the heat-stress treatment, 1797 in the control, and 1880 in the recovery. The *cis*-eQTL

were highly shared across treatments. A considerable fraction of the *trans*-eQTL (40-57%) mapped to 19

treatment specific *trans*-bands. In contrast to *cis*-eQTL, *trans*-eQTL were highly environment specific and thus

cryptic. Approximately 67% of the *trans*-eQTL were only induced in a single treatment, with heat-stress

showing the most unique *trans*-eQTL.

Conclusions

These results illustrate the highly dynamic pattern of CGV across three different environmental conditions that

can be evoked by a stress response over a relatively short time-span (2 hours) and that CGV is mainly

determined by response related *trans* regulatory eQTL.

Background

Many organisms can respond to sudden changes in the ambient environmental conditions by adjusting their gene

expression levels [1]. In particular, invertebrates are prone to environment-induced rapid gene-expression

changes. For instance, gene expression in the nematode *Caenorhabditis elegans* can swiftly change due to

exposure to pathogens, temperature, and toxicants [2-6]. A common denominator of many of these studies is that

they provide snapshots in time of the responses elicited by these environments. As such, they provide static

profiles of gene expression at a given moment. Further insight into the dynamics of gene expression and gene-

expression regulation can be achieved by following responses, such as development or aging, over time [7-9].

Gene-expression regulation can be studied by investigating the transcriptional response in the context of

natural variation using genetical genomics [10]. In this approach, a genetically segregated population (*e.g.*

recombinant inbred lines, RILs) is used in a transcriptomics experiment to determine the genetic architecture of

gene expression. Genetical genomics has been used in many species, including *C. elegans*, and many

environments. One of the marked observations resulting from a comparison of different environments is the

change in genetic architectures, revealing additional variation (as conceptualized by [11]). For example, the

transcriptional architecture of *C. elegans* grown at 16^{o}C or 24^{o}C differs markedly [12]. Such studies show that

the genetic architecture of gene expression is a dynamic process affected by relatively long-term differences in

environment or age.

Gene expression is also highly dynamic during development and growth. For example, development in

C. elegans is a tightly regulated process that has strongly correlated patterns of gene expression [8, 9, 13]. These

developmental expression dynamics can be affected by natural genetic variation, for example between two

commonly used divergent strains N2 and CB4856 [14]. Depending on the developmental stage, up to 10% of the

genes in these strains show differential expression linked to genotype. This enabled the mapping of QTLs for

gene expression dynamics during development [15].

Environmental changes can unlock genetic variation that remains hidden when in one condition but

becomes apparent in another, a phenomenon called cryptic genetic variation (CGV) [16]. CGV has received

quite some renewed interest over the past decade and it is suggested that CGV provides the raw material of

evolution and adaptation under different environmental conditions [17]. Unlocking CGV can be achieved by

altered gene-expression regulation, such as the transcriptional response to changing ambient conditions. Gene

regulatory networks play an important role in understanding how environmental cues affect cryptic genetic

variation [18]. Here we aim to investigate the CGV of the genetic architecture over the course of a strong

environmental stimulus across a recombinant inbred line (RIL) population in the nematode *C. elegans*. As

stimulus we used a heat stress, since *C. elegans* strongly reacts to temperature differences [19]. Furthermore, the

two parental strains of the RIL population, Bristol (N2) and Hawaii (CB4856), display extensive variation in

response to heat [12, 19-21]. First, we obtained the transcriptomes of the RILs over three treatments: (i) control,

48 hours at 20°C; (ii) heat stress, 46 hours at 20$^°$C followed by 2 hours exposure to 35$^°$C; (iii) recovery, same as

the heat stress, followed by an additional 2 hours at 20$^°$C. The transcriptomes were used for comparing the

transcriptional differences as well as the identified eQTL between the three treatments. The treatments resulted

in strong differences in gene expression, whereby the recovery treatment showed characteristics of both the

control and the heat-stress treatment. Comparative analysis over the identified eQTL per treatment showed that

cis-eQTL were strongly conserved over treatments. *Trans*-eQTL were more dynamic and display little overlap

between treatments. We show that CGV is mainly manifested by *trans*-eQTL of specific sets of genes in specific

environments. This makes the genetic architecture of gene expression variation an even more complex and

cryptic phenomenon than previously thought.

Methods

Strains used

The wild-types N2 and CB4856 and 54 RILs derived from a CB4856 x N2 cross were used (strains generated in

[12]). For 49/54 of these strains low-coverage sequencing was applied to construct a more detailed genetic map

(see also [22]). A matrix with the strain names and the genetic map can be found in **Additional file 1**.

Nematode culturing

The strains were kept on 6-cm Nematode Growth Medium (NGM) dishes containing *Escherichia coli* strain

OP50 as food source [23]. Strains were kept in maintenance culture at 12^{o}C, the standard growing temperature

for experiments was 20^{o}C. Fungal and bacterial infections were cleared by bleaching [23]. The strains were

cleared of males prior to the experiments by selecting L2 larvae and placing them individually in a well in a 12-

wells plate at 20^{o}C. Thereafter, the populations were screened for male offspring after 3 days and only the 100%

hermaphrodite populations were transferred to fresh 9-cm NGM dishes containing *E. coli* OP50 and grown until

starved.

Control, heat stress, and recovery from heat stress experiments for transcriptomics

The experiments were started by transferring a starved population to a fresh 9-cm NGM dish. This population

was grown for 60 hours at 20^{o}C to obtain egg-laying adults, which were bleached in order to synchronize the

population. The eggs were transferred to a fresh 9-cm NGM dish. Three growing conditions were applied: (i) the

control treatment was grown for 48 hours at 20^{o}C, (ii) the heat-stress treatment was grown for 46 hours at 20^{o}C

followed by 2 hours at 35^{o}C, and (iii) the recovery treatment was grown for 46 hours at 20^{o}C, followed by 2

hours at 35^{o}C and thereafter 2 hours at 20^{o}C. Before the start of the treatment, the developmental stage of the

population was determined by observing the developmental stage of the vulva in multiple individuals.

Populations not consisting of L4 larvae were not isolated. Directly at the end of the treatment, the population was

washed off the plate with M9 buffer and collected in an Eppendorf tube, which was flash frozen in liquid

nitrogen. In this manner, 48 RILs per condition were assayed.

Genotypes and genetic map construction

Previously, 49 lines were sequenced and aligned. The single-nucleotide polymorphism (SNP) calls per strain

were taken for constructing the genetic map [22]. The SNP density was determined per 10 kb bins and

recombination events were recognized as transition of an area where there were no CB4856 SNPs in 10

consecutive bins into an area where there were CB4856 SNPs and the other way around. It was not allowed to

have two recombination events within 10 consecutive bins (100 kb). The 10 kb bin where the first SNPs were

detected was marked as the recombination event. Before use in mapping, the map was filtered for informative

markers – that is - markers indicating a recombination event in at least one of the lines. This resulted in a map of

729 informative markers, each indicating the location of the recombination events within 10 kb (see the figure in

Additional file 2).

The genetic map was investigated by correlation analysis to assess the linkage between markers.

Markers on the centers of the chromosomes showed strong linkage (see also [24]). No strong in between

chromosome correlations were found (see the figure in **Additional file 3**).

Transcript profiling

RNA isolation

The RNA of the RIL samples was isolated using the RNeasy Micro Kit from Qiagen (Hilden, Germany). The

'Purification of Total RNA from Animal and Human Tissues' protocol was followed, with a modified lysing

procedure; frozen pellets were lysed in 150 µl RLT buffer, 295 µl RNAse-free water, 800 µg/ml proteinase K

and 1% ß-mercaptoethanol. The suspension was incubated at 55°C at 1000 rpm in a Thermomixer (Eppendorf,

Hamburg, Germany) for 30 minutes or until the sample was clear. After this step the manufacturer's protocol

was followed.

cDNA synthesis, labelling and hybridization

The 'Two-Color Microarray-Based Gene Expression Analysis; Low Input Quick Amp Labeling' -protocol,

version 6.0 from Agilent (Agilent Technologies, Santa Clara, CA, USA) was followed, starting from step five.

The *C. elegans* (V2) Gene Expression Microarray 4X44K slides, manufactured by Agilent were used. Before

starting cDNA synthesis, quality and quantity of the RNA were measured using the NanoDrop-1000

spectrophotometer (Thermo Scientific, Wilmington DE, USA) and RNA integrity was determined by agarose gel

electrophoresis (3 μL of sample RNA on 1% agarose gel).

Data extraction and normalization

The microarrays were scanned by an Agilent High Resolution C Scanner with the recommended settings. The

data was extracted with Agilent Feature Extraction Software (version 10.5), following manufacturers' guidelines.

Normalization of the data was executed in two parts, first the RILs and the ILs, second the mutant strains. For

normalization, "R" (version 3.3.1 x 64) with the Limma package was used. The data was not background

corrected before normalization (as recommended by [25]). Within-array normalization was done with the Loess

method and between-array normalization was done with the Quantile method [26]. The obtained single channel

normalized intensities were log2 transformed and used for further analysis.

Environmental responses

The transcriptional response to heat stress was determined by explaining the gene expression over the treatment

with a linear model,

165
$$y_i \sim T + e_i$$

where y is the log2-normalized intensity as measured by microarray of spot i (i = 1, 2, ..., 45220), and T is the

treatment (either control, heat stress, or recovery from heat stress). This analysis ignored genotype.

The significances were corrected for multiple testing by applying the Benjamini Yekutieli method in

p.adjust (R, version 3.3.1 Windows x64) at FDR = 0.05 [27]. Thresholds of -log10(p) ≥ 2.87 for the control

versus heat-stress treatment, -log10(p) ≥ 3.09 for the control versus recovery treatment, and -log10(p) ≥ 3.02 for

the heat-stress versus recovery treatment were determined.

Developmental variation

Due to the setup of our experiment, potential variation in development could exist among the RILs and the

treatments. The recovery animals were sampled two hours later than the control and heat-stress animals,

furthermore, heat stress slows the developmental rate [19]. We estimated the relative age by using a set of ~100

genes that show a strong, positive, linear response during development [9]. By setting the average age of the

control RILs to 48 hours we could estimate and compare the RILs in all treatments (**Additional file 8**).

Principal component analysis

A principal component analysis was conducted on the gene-expression data of the RILs over the three

treatments. For this purpose, the data was transformed to a log2 ratio with the mean, using

183
$$R_{i,j} = \log 2(\frac{y_{i,j}}{\bar{y}_i})$$

where R is the log2 relative expression of spot i (i = 1, 2, ..., 45220) in strain j (RIL) over all three conditions (n

= 48 per condition), and *y* is the intensity (not the log2-transformed intensity) of spot i in strain j.

The transformed data was used in a principal component analysis, where the first six axes were further

examined.

Expression quantitative trait locus analysis

eQTL mapping and threshold determination

The eQTL mapping was done in "R" (version 3.3.1 Windows x64). The gene-expression data was fitted to the

linear model,

193
$$y_{i,j} \sim x_j + e_j$$

where y is the log2-normalized intensity as measured by microarray of spot i (i = 1, 2, ..., 45220) of RIL j. This

is explained over the genotype (either CB4856 or N2) on marker location x (x = 1, 2, ..., 729) of RIL j.

The genome-wide significance threshold was determined via permutation, where the log2-normalized

intensities were randomly distributed per gene over the genotypes. The randomized data was tested using the

same model as for the eQTL mapping. This was repeated for ten randomized datasets. A false discovery rate was

used to determine the threshold (as recommended for multiple

testing under dependency) [27],

201
$$\frac{FDS}{RDS} \leq \frac{m_0}{m} q \cdot \log(m)$$

where FDS is the outcome of the permutations and RDS is the outcome of the eQTL mapping at a specific

significance level. The value of m_0, the number of true null hypotheses tested, was 45220-RDS, and for the value

of m, the number of hypotheses tested, the number of spots (45220) was taken. The q-value was set at 0.05. This

yielded a threshold of –log10(p) > 3.9 for the control, –log10(p) > 3.5 for the heat stress, and –log10(p) > 3.9 for

the recovery treatment. For the analyses we used the most conservative thresholds measured, -log10(p) > 3.9, for

all the sets.

Statistical power calculations

In order to determine the statistical power at the set FDR threshold, QTL were simulated using the genetic map

of the strains used per condition (n=48 per condition). For each marker location, ten QTL were simulated that

explained 20-80% of the variation (in increments of 5%). Random variation was introduced based on a normal

distribution with sigma = 1 and mu = 0 and a peak of the corresponding size (e.g. a peak size of 1 corresponds to

20% explained variation) was simulated in this random variation. From the simulation, the number of correctly

detected QTL, the number of false positives and the number of undetected QTL were counted. This was based

on the thresholds determined in the permutations, –log10(p) > 3.9. Furthermore, the precision of the effect-size

estimation and the precision of the QTL location (based on a –log10(p) drop of 1.5 compared to the peak) were

determined. A table summarizing the results can be found in **Additional file 9**.

eQTL analysis

The distinction between *cis-* and *trans-*eQTL was made on the distance between the physical location of the gene

and the location of the eQTL-peak. For *cis-*eQTL the gene lies within 1 Mb of the peak or within the confidence

interval of the eQTL. The confidence interval was based on a –log10(p) drop of 1.5 compared to the peak.

The amount of variation explained per microarray spot with an eQTL was calculated by ANOVA, by

analysis of the gene expression explained over the peak-marker. For spots with multiple peaks, this analysis was

conducted per peak, not using a full model, since a single-marker model was used in the analysis.

In order to identify *trans-*bands (an enrichment of *trans-*eQTL), a Poisson distribution of the mapped

*trans-*eQTL was assumed (as in [28]). Therefore the number of *trans-*eQTL per 0.5 Mb bin were counted. Since

*trans-*eQTL peaks were mapped to 107, 106, and 103 bins (respectively in control, heat stress, and recovery), it

was expected that 9.16, 20.64, and 9.01 spots with a *trans-*eQTL were to be found at each of these markers.

Based on a Poisson distribution, it was calculated how many *trans-*eQTL needed to be found to represent an

overrepresentation. For example, for p < 0.001 there should be 20, 36, or 20 spots with a *trans-*eQTL at a

specific marker (respectively in control, heat stress, and recovery).

To test for polymorphisms in genes with eQTL, we used the data from the CB4856 reference genome

[22]. The genes with eQTL were matched to the polymorphisms. The frequencies of polymorphisms in each of

the groups (genes with *cis*-eQTL, genes with *trans*-eQTL, and genes without eQTL) were counted and compared

versus each other by a chi-squared test in "R" (version 3.3.1, x64).

Detection of eQTL across treatments

Two criteria were used to detect the occurrence of eQTL over multiple treatments.

In the first criterion, it was tested whether or not an eQTL was mapped in treatment one versus

treatment two, by simply comparing the tables listing the eQTL. This allowed for comparison of the actual

mapped peaks and for comparison of eQTL effects of *trans*-eQTL regulated from different loci. In order to

estimate the false-discovery rate associated with this comparison, the same analysis was applied to ten

permutated datasets per condition, using the $-\log10(p) > 3.9$ for eQTL discovery.

The second criterion compared the occurrence of eQTL at the exact same marker location. In this

comparison, the eQTL mapped in one treatment were taken as lead for the occurrence of the same eQTL in the

other two treatments. This comparison allowed for direct comparison of the eQTL effect at the locus. Based on

observations on the effect distribution, this approach was used to estimate the number of *trans*-eQTL not

detected due to statistical power or not detected due to absence of the eQTL in a treatment (see also text in

Additional file 15).

Functional enrichment analysis

Gene group enrichment analysis was done using a hypergeometric test and several databases with annotations.

The databases used were: the WS220 gene class annotations, the WS256 GO-annotation, anatomy terms,

phenotypes, RNAi phenotypes, developmental stage expression, and disease related genes (www.wormbase.org)

[29]; the MODENCODE release 32 transcription factor binding sites (www.modencode.org) [30, 31], which

were mapped to transcription start sites (according to [32]); and the KEGG pathway release 65.0 (Kyoto

Encyclopedia of Genes and Genomes, www.genome.jp/kegg/) [33].

Enrichments were selected based on the following criteria: size of the category n>3, size of the overlap

n>2. The overlap was tested using a hypergeometric test, of which the p-values were corrected for multiple

testing using Bonferroni correction (as provided by p.adjust in R, 3.3.1, x64). Enrichments were calculated based

on gene names, not on spots.

Results

Transcriptional response over the course of heat stress

To better understand the transcriptional response to heat stress, we obtained the transcriptomes of 48 recombinant inbred lines (RILs) at the L4 stage in each of three treatments: control, heat stress, and recovery from heat stress (**Figure 1A**). The effects of the treatments on gene-expression levels were analyzed using a linear model for pairwise comparisons between each of the conditions (see volcano plots in **Additional file 4** and a list of affected spots in **Additional file 5**). In this way, we identified 7720 differentially expressed genes over the course of the three treatments (FDR = 0.05; **Figure 1B**). We found that both control and heat stress had many unique differentially expressed genes: 2321 genes were only differently expressed in the comparisons of the control treatment to the other two treatments and 3305 genes were only differently expressed in the comparisons of the heat-stress treatment with the other two treatments. In the comparisons with the recovery treatment, only 942 genes were found unique for that treatment. Furthermore, many differentially expressed genes were shared in the comparison of the recovery treatment versus the control and heat-stress treatment (2251 genes). Again, the control and recovery (1152) and heat stress and recovery (1189) shared fewer genes. There were 1255 genes that were differentially expressed between all three conditions and were therefore highly treatment dependent. These results indicate that the control and heat-stress treatments are strongly contrasting in gene expression, whereas the recovery treatment shares characteristics with both other treatments.

To follow up on this interpretation, a principal component analysis (PCA) was conducted on the gene-expression data, transformed to the log2 ratio with the mean. The first axis (20.0% of the variation) captures variation mostly related to the heat-shock treatment, which is expected as this treatment specifically affect the largest number of genes (**Additional file 6**). The second axis (11.4% of the variation) captures variation mostly related to the control treatment, which also fits the analysis with the linear model as the control treatment was the second most distinct treatment. Together, these two axes also place the recovery treatment in between the control and heat-stress treatment, showing the contrast with the other two treatments was lower and possibly indicating transcript levels in the recovery treatment are returning to normal. The third principal component (10.1% of the variation) captures variation that sets the heat-stress treatment completely apart from the other two treatments, which is as expected since the heat-stress treatment has the most unique differentially expressed genes.

In order to gain further insight into the functional differences between the treatments, an enrichment analysis was conducted on genes belonging to the different overlap groups as shown in **Figure 1** (For example

genes differentially expressed in only the control treatment, see the list in **Additional file 7**). Each of the groups

was enriched for many processes, showing that the treatments had a profound impact on gene expression.

Interestingly, we found that genes specific for each of the three treatments were enriched for genes expressed in

the intestine. Furthermore, the control and heat-stress treatments were strongly enriched for genes expressed in

the germline. These enrichments indicated that the expression of genes involved in metabolism and reproduction

(or development of the reproductive organs) were strongly altered during the heat-stress response. As this may

be caused by a developmental difference in the sampled populations, we estimated the developmental age using

a transcriptional ruler (see **Materials and methods** for details) [9]. It was found that the control population was

transcriptionally slightly younger than the heat shock (estimated ~1 h older) and the recovery population

(estimated 1.3 h older; **Additional file 8**).

Gene expression linked to genetic variation

Linkage mapping was performed using 48 RILs for each of the three treatments. Statistical power analysis

showed that this population has the power to detect 80% of the eQTL that explain at least 35% of the variation

(see Materials and methods and the table in **Additional file 9**). Identified eQTL (FDR ≤ 0.05; **Figure 2;** a table

with all eQTL is given in **Additional file 10**) were compared between the treatments (**Table 1**). Most genes with

an eQTL were found in the heat-stress treatment (2626), whereas the control (1797) and recovery (1880) had

similar numbers. This increase in genes with eQTL was primarily caused by the larger number of genes with

trans-eQTL in the heat-stress treatment (1560; ~57% of total eQTL) compared to the control (751; ~40% of

total) and recovery (739; ~38% of total). The number of *cis*-eQTL was almost identical among conditions (**Table**

1).

Table 1: Number of genes with an eQTL			
	Control	**Heat stress**	**Recovery**
cis-eQTL[1]	1138	1186	1215
N2 higher	809 (71.1%)	802 (67.6%)	883 (72.7%)
CB4856 higher	329 (29.9%)	384 (32.4%)	332 (27.3%)
trans-eQTL	751	1560	739
N2 higher	300 (39.9%)	583 (37.4%)	393 (53.2%)

	CB4856 higher	451 (60.1%)	977 (62.6%)	346 (46.8%)
Total[2]		1797	2626	1880

[1]: *cis*-eQTL were called if the QTL peak lies within 1 Mb of the affected gene, or if the affected gene lies within the 1.5 LOD-drop confidence interval.

[2]: The differences between the total and the summation of the individual *cis*- and *trans*-eQTL is due to genes with both a *cis*- and *trans*-eQTL.

The *cis*-eQTL showed a bias for higher expression if the regulatory locus had the N2 allele (on average

70% of the genes with a *cis*-eQTL). This bias was absent in the *trans*-eQTL where, on average, 43% of the genes

with a *trans*-eQTL was more highly expressed if the locus had the N2 allele (illustrated by the figure in

Additional file 11). This discrepancy was to be expected, since the microarray platform used to measure the

transcripts was designed for the N2 genotype. Therefore, part of this variation is likely due to mis-hybridization.

However, previous studies have shown that this does not explain all the variation (see [28]). In congruency, we

found genes with a *cis*-eQTL to be more polymorphic than genes with a *trans*-eQTL and genes without an eQTL

(summarized in **Additional file 12**). For example, genes with a *cis*-eQTL were more likely to be fully deleted in

CB4856 (9.5% of the genes, compared to 0.9% in genes without an eQTL or 1.1% in genes with a *trans*-eQTL;

Chi-squared test, $P < 1*10^{-34}$). Furthermore, supporting that not all *cis*-eQTL stem from mis-hybridization,

polymorphisms in the flanking 3' and 5' regions were about two times more likely to occur near *cis*-eQTL

compared to genes without a *cis*-eQTL (Chi-squared test, $P < 1*10^{-6}$). When these enrichments for 3' and 5'

polymorphisms were compared between *cis*-eQTL with an N2 or CB4856 higher effect, we found no significant

difference.

The *trans*-eQTL were mainly found in 19 treatment-specific *trans*-bands, loci that regulate the

abundance of many transcripts. The 19 *trans*-bands were identified by analysis of the occurrence of *trans*-eQTL

across the genome (Poisson distribution, P<0.001; listed in **Additional file 13**). The seven *trans*-bands detected

in the heat-stress treatment affected most genes (1141; ~73% of all *trans*-eQTL). In the control treatment, five

trans-bands were found (325 genes; ~43% of all *trans*-eQTL) and in recovery treatment seven *trans*-bands were

identified (343 genes; ~46% of all *trans*-eQTL). Six out of the 19 *trans*-bands individually affected >100 genes,

one located at chromosome X: 0.5-2.0 Mb in control treatment; four in heat-stress treatment at chromosome I:

2.0-3.5 Mb, II: 12.0-13.5 Mb, IV: 1.0-2.5 Mb, and V:1.0-3.0 Mb; and one in recovery treatment at chromosome

I: 1.5-3.0 Mb. Importantly, the distribution across the genome of *trans*-bands and eQTL is treatment-specific. To

further investigate this, we determined the overlap in mapped eQTL over the course of the heat-stress response.

In contrast to cis-*eQTL,* trans-*eQTL were environment-specific*

Comparing the genes with a *cis*-eQTL among the three treatments, we found 1086 out of 1789 unique genes

(~61%) with a *cis*-eQTL in more than one treatment and 664 (~37%) in all three treatments (**Figure 3A**).

Because *cis*-eQTL can be caused by mis-hybridizations, we also calculated the overlap for *cis*-eQTL with an N2

higher effect. In that selection, 303 out of 615 unique genes (~49%) had a *cis*-eQTL in more than one treatment

and 157 (~26%) in all three treatments. For the whole set of genes with a *trans*-eQTL the overlap was much

smaller (**Figure 3B**); 360 out of 2610 genes (~14%) were found in more than one treatment and only 80 genes

(~3%) in all three treatments. By definition, the locus at which *cis*-eQTL were detected was the same between

treatments. Furthermore, the *cis*-eQTL effect sizes and directions were highly comparable (Pearson correlation

coefficients between 0.94-0.96; shown in a figure in **Additional file 14**). The *cis*-eQTL that were detected in

only one treatment were probably missed due to the small amount of variation explained by these eQTL (see the

text in **Additional file 15**). Interestingly, there were only three genes with *cis*-eQTL showing genotypic

plasticity: C52E2.4, C54D10.9, and *nhr-226*. This meant that we observed allelic variation acting in opposite

directions between environments (**Figure 3C**).

The effect sizes and direction of genes with a *trans*-eQTL found in more than one treatment were very

similar (Pearson correlation coefficients between 0.90-0.93; figure in **Additional file 14**). However, only a few

genes with a *trans*-eQTL were found over multiple treatments, far lower than the overlap in *cis*-eQTL between

conditions (**Figure 3A** and **B**). Since *trans*-eQTL are not by definition regulated from the same location, the

overlap between treatments declines even further if location is taken into account. Taking the *trans*-eQTL from

the three treatments together, only ~38% of the multi-treatment *trans*-eQTL were located at a different locus

from one treatment to another (~28% if only loci at different chromosomes were counted), see **Additional file**

16 (figure A). Interestingly, although the regulatory locus was located elsewhere, the genotypic effect of the

trans-eQTL was almost identical (**Additional file 16,** figure B), yet most genes with *trans*-eQTL only displayed

an eQTL in one treatment (text in **Additional file 15**). The likely reason that the majority of *trans*-eQTL was not

detected across treatments is that most *trans*-eQTL are environment-specific and therefore highly cryptic.

Trans-*eQTL display two types of cryptic variation*

For treatment-specific, cryptic *trans*-eQTL, we found that a regulator is active when the eQTL is detected and it

is not active when no eQTL is detected. With on and off switching regulators between treatments, genes can

have dynamic *trans*-eQTL, which appear as a treatment-specific switch of regulatory loci (**Figure 3 - 5**). Since

the majority of genes with a *trans*-eQTL have one unique *trans*-eQTL in only one treatment (**Figure 3B**), a

switch in regulatory loci seems to occur less frequently compared to the on/off switch.

As the majority of the *trans*-eQTL was treatment-specific, we investigated whether detection in only

one treatment was a result of the statistical power of our study or if it was a biological phenomenon. As *trans*-

eQTL explain 34.2% of variation on average, compared to 52.1% of variation for *cis*-eQTL (text in **Additional**

file 15), it is possible that the detection of *trans*-eQTL was more affected by lack of statistical power. By

simulations, we estimated that this only affected 20.6% of the *trans*-eQTL that were not detected in multiple

treatments (for the detailed analysis, see the text in **Additional file 15**), which argues for the cryptic nature of

trans-eQTL. Another line of evidence for this is the low overlap in affected genes in co-locating *trans*-bands

across treatments (13/19 *trans*-bands co-locate). We only found significant overlap in three pairs of *trans*-bands,

where 6.5-11.1% of the affected genes overlap (text in **Additional file 15;** hypergeometric test $p < 1*10^{-4}$). For

example, one of these, a major *trans*-band at chromosome IV:1-2.5 Mb in the heat-stress treatment, affected 244

genes of which 22 overlapped with the 31 genes in the recovery *trans*-band on chromosome IV:1-2 Mb.

Together, these results show that the majority (67.3%) of *trans*-eQTL indeed are cryptic. For example, the

transcript levels of *flp-22*, *pqm-1*, and *sod-5* were affected by treatment (**Figure 4A**) and showed a *trans*-eQTL

in only one treatment (**Figure 4B**).

As mentioned before, only a minority of the genes with a *trans*-eQTL have different eQTL across

treatments. Of those genes with different *trans*-eQTL over treatments, the genotypic effects of the eQTL were

similar across treatments, even if the loci were different (**Additional file 16,** figure B). Only between

chromosome IV and V were eQTL with changes in effect directions observed. One of the genes displaying this

pattern was *gei-7* (also known as *icl-1*). This gene was represented by three different micro-array probes, all

showing the same pattern: a primary eQTL on chromosome V at ~12.0Mb in all three conditions and a

secondary eQTL in the heat-stress treatment at chromosome IV at ~1.5Mb (**Figure 5**).

Functional enrichment of eQTL

To find which biological processes were affected by genetic variation on a gene-expression level we looked for

enrichment in gene classes, phenotypes, KEGG pathways, GO terms, and anatomy terms (see the list in

Additional file 17). For *cis*-eQTL enriched categories were similar in all three treatments, as expected by the

consistent nature of *cis*-eQTL. For *cis*-eQTL, we found enrichments for the gene classes *bath*, *math*, *btb*, and

fbxa, which were previously found to be highly polymorphic between CB4856 and N2 [22]. Moreover, we found

enrichment for genes involved in the innate immune response and protein homo-oligomerization. It should be

noted that these enrichments are likely due to hybridization differences for the polymorphic genes (as *cis*-eQTL

with a positive N2 effect are enriched for exactly these categories, **Additional file 17**).

The *trans*-eQTL were also enriched for genes functioning in the innate immune response, especially for

genes where the N2 allele leads to higher expression. Furthermore, genes expressed in the intestine were

enriched in the *trans*-eQTL found in control and heat-stress conditions. Contrasting to the genes with *cis*-eQTL,

the genes with *trans*-eQTL were enriched for many different transcription factor binding sites, indicating active

regulation of *trans*-eQTL.

Consistent *trans*-eQTL were found in all three treatments for the enriched NSPC (nematode specific

peptide family, group C) gene class. This was remarkable as only a very small part of the *trans*-eQTL were

shared over the three treatments. For the heat stress and recovery *trans*-eQTL, genes expressed in the

dopaminergic neuron were enriched, with the strongest enrichment in the heat-stress treatment. These genes

were also enriched in the control treatment, however, in a group of *trans*-eQTL mapping to a different trans-

band. In the heat stress and recovery treatment the dopaminergic neuron-specific genes showed *trans*-eQTL at

the IV:1-2MB locus, whereas in the control they showed *trans*-eQTL at the X:4-6 locus (**Figure 6**).

Discussion

Transcriptional response over the course of a heat stress

Here we present a comprehensive study of the effects of an induced heat-stress treatment on the genetic

architecture of gene expression. The obtained transcriptomes were analyzed in the context of the treatments and

in the context of the genetic variation present in the strains used in the experiments.

We found that many of the genes that are affected over the heat-stress course are associated with

expression in the germline and intestine. These findings are partially in line with findings from an investigation

on the heat-shock regulatory network using a genome-wide RNAi screen in *C. elegans* [5]. Their heat-stress

conditions were 31.5°C for two hours followed by 24 hours of recovery at 20°C. The authors found that genes

associated with the proteasome induced heat-stress-specific gene expression only in the intestine and

spermatheca, which corroborates with our results. Differences with Guisbert *et al.* (2013), could be explained by

the differences in larval stage used (L4 vs L2) as well as differences in temperature and duration of the heat

stress (and recovery). Another study exposed *C. elegans* at the L4 stage to a heat stress of 30 min at 33℃, and

measured transcriptome differences using RNA-seq [34]. In support of our study, they also detected genes

associated with metabolism and reproduction, whereas, in contrast to our findings, they found a strong link with

cuticle specific genes. This contrast could be due to the different experimental conditions of Bunquell *et al.*

(2016), as they used RNAi treatments (empty vector or against *hsf-1*), a different heat shock duration, method of

synchronization (additional L1 arrest by [34]), and rearing temperature before heat shock (23^{o}C versus 20^{o}C in

this study).

We hypothesize that ultimately these discrepancies are likely to result from developmental differences.

The transcriptional program in *C. elegans* differs strongly during development [8, 9]. Therefore, application of a

heat shock on L2 larvae has a very different developmental (and therefore transcriptional) starting point

compared to a heat shock applied on L4 larvae. Furthermore, within the L4 stage there is a strong difference in

gene expression in early-stage L4 and late-stage L4. We estimated that the heat-stress and recovery RILs were

one hour older than the control RILs suggesting that the former had an accelerated development as these

populations were physically the same age as the control population. The main processes that affect transcription

in the L4 stage are reproduction and development [9]. These are exactly the processes that are halted upon

induction of a heat shock [5, 34]. Therefore, it is likely that the state in which these processes are strongly affects

the possible routes for down regulation. It would therefore be very interesting to study the effect of heat shock in

relation to the developmental dynamics.

If the effect of a heat shock is indeed dependent on developmental status, then the transcriptomes of the

RILs presented in this study could be indicative of the phenotypic outcome (*e.g.* heat-stress survival). The reason

being the developmental gradient generated by RILs (as shown by [15] within a single experiment, or explicitly

by [35] between different stages), which mainly affects *trans*-eQTL. Alternatively, our experiment could also be

analysed by including the variation among the estimated age of the RILs in the mapping model together with

treatment effects. A thorough interpretation of these models requires a better understanding of the transcriptional

dynamics over the heat-shock and recovery response, a goal we are currently actively pursuing [36].

In contrast to trans-*eQTL,* cis-*eQTL are directly linked to polymorphisms*

The *cis*-eQTL over the three treatments, which strongly overlapped, are highly enriched for polymorphic genes.

This has been reported before in *C. elegans*, but also in *A. thaliana*, *Mus musculus*, and for human *cis*-eQTL [12,

28, 37-41]. This can result in the detection of transcriptional variation that is actually caused by hybridization

differences [28, 40]. Analysis of the bias in *cis*-eQTL with higher expression in N2 (the strain for which the

microarray was developed) versus CB4856 indeed shows that a proportion of the *cis*-eQTL is likely to stem from

hybridization differences. Also apparent from the gene-enrichment analysis, *cis*-eQTL were overrepresented for

polymorphic gene classes such as *bath*, *math*, *btb*, and *fbxa*, which are also divergent among other wild strains

[42, 43]. Other experimental methods could limit such 'false positives', for example, RNA sequencing is

expected not to suffer from such biases [44].

Interestingly, genes with a *cis*-eQTL were also strongly enriched for polymorphisms in regulatory

regions. For these *cis*-eQTL, it could be true that the expression is affected by transcription factor (TF) binding

sites [45], yet we did not detect any enrichment for such sites as mapped by ModEncode [30, 31]. An

explanation for this is that genes with a *cis*-eQTL are regulated by different TFs, therefore the affected TF

binding site is different per gene with a *cis*-eQTL making an overrepresentation among all *cis*-eQTL unlikely.

CGV of transcriptional architecture is determined by trans-*eQTL*

Although previous studies in *C. elegans* have focused on continuous (thermal or developmental) treatments or

gradual change over time [12, 15, 35], only few genetical genomics experiments have measured the effect of

acute perturbation, where an organism is suddenly exposed to a different environment [39, 46]. This acutely

affected the transcriptional architecture of gene expression, which consists of *cis-* and *trans*-acting eQTL. We

found that *cis*-eQTL were robust across all three treatments, including heat stress, which was found before in

studies on *C. elegans* and other species where eQTL patterns have been studied in different conditions. For

example, Li *et al.* (2006) found that more than 50% of all *trans*-eQTL were affected by temperature compared to

cis-eQTL [12], and Smith and Kruglyak (2008) also reported that *cis*-eQTLs were hardly affected by external

conditions as compared to *trans*-eQTL [47]. Also, in other species, like *Arabidopsis thaliana*, it was reported

that *cis*-eQTL were robust to light perturbation, whereas *trans*-eQTL were more affected by different light

regimes [39]. In humans, *cis*-eQTL showed a very high correlation between tissues compared to *trans*-eQTL,

and are replicated across populations in lymphoblastoid cell lines [48, 49].

By inducing a strong environmental perturbation in the form of a heat shock, a high amount of different

trans-eQTL were detected across the control, heat-stress, and recovery treatments. One of the things we noted is

the increase in the number of *trans*-eQTL and *trans*-bands in the heat-stress treatment compared to the other two

treatments. This strong transcriptional variation could underlie – or result from - the phenotypic trait differences

observed between N2 and CB4856 in temperature experiments. However, it should be noted that also

developmental differences due to timing between the three treatments can contribute [9, 15]. Temperature-

affected trait differences have been observed for age at maturity, fertility, body size, vulval induction, and

lifespan [19, 50-53]. Furthermore, these strains also display behavioural differences in heat avoidance and

thermal preference [20, 21]. Likely candidates for these trait differences could be found in loci affecting the

expression of many *trans*-eQTL. For example, the left arm of chromosome IV harbours a *trans*-band affecting

the expression of 244 genes and coincides with a QTL affecting lifespan after heat shock [19]. Analogously, the

laboratory allele of *npr-1* affects many trait differences between N2 and CB4856 (as reviewed in [54]). Most

importantly, *npr-1* affects the behaviour of the animal, which probably results in gene-expression differences, as

part of the expression differences can be mimicked by starving nematodes [55]. These gene-expression

differences can be picked up as a *trans*-band [28]. The latter example illustrates both a link between gene-

expression and classical traits and that caution is required for inferring the direction of causality.

Why is CGV mainly affecting *trans*-eQTL? We hypothesize that this is due to the versatile nature of

trans-eQTL. First, *trans*-eQTL are loci that are statistically associated with variation in transcript abundance

from genes elsewhere on the genome. The ultimate causes for this association can be manifold, from direct

interactions such as polymorphic transcription factors that affect gene expression to indirect interactions such as

receptor-kinase interactions [46], receptors [46, 55, 56], or effects at the behavioural level that result in

expression differences [55]. Therefore, an environment can require the organism to respond, thereby requiring

specific polymorphic genes to react, ultimately leading to the expression of cryptic variation.

The trans-*eQTL architecture is comprised of treatment-specific genes*

The *trans*-eQTL architecture is remarkably unique over the three treatments tested. We only observed 3%

overlap in *trans*-eQTL in the three treatments, for which the main cause was treatment specificity of *trans*-

eQTL. Surprisingly, genetic variation affects the expression of genes in only one direction; only in rare cases

does allelic variation change the sign of the effect and this was only observed for *cis*-eQTL (C52E2.4,

C54D10.9, and *nhr-226*). On the one hand, this is in congruency with other eQTL studies comparing different

environments; *trans*-eQTL are strongly affected by different environments (for example, see [12, 39, 47]). On

the other hand, it raises questions about the genetic architecture of *trans*-bands; co-localizing *trans*-bands are

generally not affecting the expression of the same genes (see text in **Additional file 15**), which can imply the

involvement of multiple regulators (causal genes).

However, it seems unlikely that an abundance of novel *trans*-eQTL also implies an abundance of novel

causal genes. First of all, over all three treatments, the majority of *trans*-eQTL are located in *trans*-bands, which

are mostly non-overlapping between treatments. This is an observation that extends to other studies and other

species in which eQTL have been mapped: the majority of *trans*-eQTL are found on a few regulatory hotspots

(for example, see [28, 38, 57]). Therefore, it is logical to assume that a small set of causal genes ultimately

explains the majority of *trans*-eQTL. Second, the allelic effect only has one direction, which is easily aligned

with the notion of a few regulators instead of many. Together, these observations can help in further dissecting

loci to identify causal genes. *Trans*-band regulators might play a role in the dynamic response, which could aid

in narrowing down candidate genes.

However, it should be reiterated that the route from genetic variation resulting in transcriptional effect

can be manifold, which can also obscure the ultimate cause of the observed trait variation. An organism is an

intricate web of interdependencies leading to the phenotype. Therefore, upon further dissection of the loci, a

single eQTL might prove to be many. Furthermore, it should be noted that *trans*-eQTL explain less variation

compared to *cis*-eQTL, which has been established across species [28, 58]. Therefore, it is more likely that *trans*-

eQTL are not detected. However, the treatment specificity and direction of allelic effects of *trans*-eQTL across

three treatments robustly show *trans*-eQTL architecture is comprised of treatment-specific genes.

Conclusion

Here we present the contribution of CGV on eQTL across three treatments in the nematode *C. elegans*. We find

that mainly *trans*-eQTL are affected by CGV, in contrast to *cis*-eQTL, which are highly similar across

treatments. Furthermore, we show that most CGV results in unique genes with a *trans*-eQTL, instead of different

allelic effects and/or different eQTL for the same genes. This shows the highly dynamic nature of CGV.

Declarations

Availability of data and materials

All strains used can be requested from the authors. The transcriptome datasets generated and the mapped eQTL

profiles can be interactively accessed via (http://www.bioinformatics.nl/EleQTL). Moreover, the transcriptome

datasets are also deposited at ArrayExpress (E-MTAB-5779).

Competing interests

The authors declare that they have no competing interests.

Funding

LBS was funded by ERASysbio-plus ZonMW project GRAPPLE - Iterative modelling of gene regulatory

interactions underlying stress, disease and ageing in *C. elegans* (project 90201066) and The Netherlands

Organization for Scientific Research (project no. 823.01.001). The funding bodies had no role in the design nor

the collection, analysis, and interpretation of the data, nor the writing of the manuscript.

Authors' contributions

LBS and JK conceived and designed the experiments. MGS, RPJB, RJMV, JAGR, and RB conducted the

experiments. AVH, RB conducted the sequencing of the strains. LBS, MGS, and RPJB conducted transcriptome

and main analyses. LBS, MGS, and JEK wrote the manuscript. RPJB, RJMV, and AC provided comments on the

manuscript.

Acknowledgements

The authors are grateful for the discussions and support from the GRAPPLE project partners: Olga Valba,

Sreenivas Chavali, Benjamin Lang, Mirko Francesconi, Sergei Nechaev, Olga Vasieva, M. Madan Babu, and

Ben Lehner. We are also grateful for Katharina Jovic, Lisa van Sluijs and two anonymous reviewers for

feedback on the manuscript. We also want to thank Harm Nijveen for making our data available in EleQTL.

Figure legends

Figure 1: Effect of treatment on gene expression. (**A**) The RIL populations were exposed to three treatments:

control (48h at 20^{o}C), heat stress (46h at 20^{o}C and 2h at 35^{o}C), and recovery (as heat stress, with an additional 2h

at 20^{o}C). At the end of these treatments, the nematodes were in the L4 stage, and were harvested. Thereafter

RNA was isolated and the transcriptome was measured by microarray. (**B**) The outcome of the treatment

analysis. On the left a legend is included to clarify which contrasts are compared. On the right, the overlap in

differentially expressed genes is shown per treatment comparison. For example, 942 genes are uniquely

differentially expressed in the recovery treatment, these genes are differently expressed between recovery and

heat stress and between recovery and control, but not between heat stress and control.

Figure 2: Identified eQTL in control (left), heat stress (middle), and recovery (right) treatments, with a

threshold of -log10(p) > 3.9 (FDR ≤ 0.05) in each treatment. The eQTL peak position is shown on the x-axis and

gene position is shown on the y-axis. The *cis*-eQTLs (within 1Mb of the gene) are shown in black and the *trans*-

eQTLs in blue (control), red (heat stress), or green (recovery). The horizontal bars indicate the confidence

interval of the eQTL. The chromosomes are indicated on the top and right of the plot. The histogram under the

plot shows the eQTL density per 0.5 Mb bin.

Figure 3: Overlap in *cis*- and *trans*-eQTL between conditions. (**A**) The overlap in genes with a *cis*-eQTL

between conditions. QTL were selected based on FDR ≤ 0.05 per condition, a QTL was scored as overlapping if

the same gene had a *cis*-eQTL in another condition. (**B**) The overlap in genes with a *trans*-eQTL between

conditions. QTL were selected based on FDR ≤ 0.05 per condition, a QTL was scored as overlapping if the same

gene had a *trans*-eQTL in another condition. The location was not yet considered in this analysis (see

Supplementary figure 7A). (**C**) The three genes with a *cis*-eQTL displaying genotypic plasticity, where the

direction of the effect switches between environments. For all three genes: C52E2.4, C54D10.9, and *nhr-226* the

plastic response is apparent between the heat stress and recovery treatment.

Figure 4: Genes with treatment specific *trans*-eQTL. (**A**) The expression patterns over the three treatments are

shown for *flp-22*, *pqm-1*, and *sod-5*, the location mentioned is the location of the gene. The x-axis is organized

per treatment (ct, control; hs, heat stress; rec, recovery). On the y-axis the log2 normalized expression is shown

(**B**) The eQTL patterns for the same three genes. The x-axis shows the position along the chromosomes and the

y-axis the significance of the association. The horizontal dashed line indicates the FDR ≤ 0.05 (-log10(p) > 3.9).

Colors indicate the three treatments.

Figure 5: The *trans*-eQTL of *gei-7*. (**A**) The eQTL patterns for *gei-7*, the x-axis shows the position along the

chromosomes and the y-axis the significance of the association. The horizontal dashed line indicates the FDR $\leq$

0.05 (-log10(p) > 3.9). Colors indicate the three treatments (control, blue; heat stress, red; recovery, green). (**B**)

The genotype effects split out at the minor heat stress QTL (chromosome IV). (**C**) The genotype effects split out

at the major QTL (chromosome V).

Figure 6: Genes belonging to the dopaminergic neuron anatomy term with a *trans*-eQTL. The *trans*-eQTL

position of the genes is shown (n = 133 in control, n = 281 in heat stress, and n = 91 in recovery).

Additional files

Additional file 1: Strains and genotypes. Matrix with the strain names and genotypes of the recombinant

inbred lines used in this study. The genotypes are based on genome sequencing (see Materials and Methods).

Additional file 2: A figure of the location of the 729 markers. The marker locations are plotted across the

genome. Locations are based on WS256.

Additional file 3: **A figure of the marker correlation analysis**. Correlations between the 729 markers in the

sequenced RIL population. The markers are plotted at their physical locations across the chromosomes.

Additional file 4: Figure of the treatment comparisons. Volcano plots of the expression comparisons per

treatment (n = 48 RILs per treatment). The horizontal line in the plots indicates the FDR = 0.05 threshold. The

colored dots indicate spots that are significantly different between treatments. Blue indicates spots more highly

expressed in the control treatment, red indicates spots more highly expressed in the heat-stress treatment, and

green indicates spots more highly expressed in the recovery treatment. (**A**). The comparison between control and

heat stress, threshold: -log10(p) $\geq$ 2.87. (**B**). The comparison between control and recovery, threshold: -log10(p)

$\geq$ 3.09. (**C**). The comparison between heat stress and recovery, threshold: -log10(p) $\geq$ 3.02.

Additional file 5: Table of treatment comparison results. A table with the spots that are significantly different

between treatments. The spot number, the comparison in which the spot was different, and the characteristics of

the difference (significance and effect) are given. Also information about the gene represented by the spot is

shown (WormBase identifier, sequence name, public name, and the location on the genome). The last column

indicates to which group the spot belongs (*e.g.* specific for heat-stress treatment, or significantly different

between all three treatments).

Additional file 6: Figure of the principal component analysis. The first six axes of the principal component

analysis are shown, plotted against each other. The axis capturing most variation (PC1), captures 20% and the

sixth axis captures 4.7%. The dots represent individual samples (n = 48 RILs per treatment) and are colored

according to treatment: blue for control, red for heat stress, and green for recovery. The first axis separates the

heat-stress treatment from the control and recovery treatment, but also captures some technical variation (19

samples that fall to the right of the plot). The second axis (11.4%) places the recovery treatment in between the

other two treatments, whereas the third axis (10.1%) separates the heat stress from the control and recovery

treatment.

Additional file 7: A list of the enrichment analysis treatment responsive genes. Enrichment analysis on the

genes with transcriptional responses by heat stress. The database used for enrichment (Annotation) and the

category (Group), and the number of genes on the array that are in the group (Genes_in_group) are also

indicated. Furthermore, the overlap with the cluster (Overlap) and the Bonferroni-corrected significance of that

overlap are shown.

Additional file 8: Developmental ruler applied to the RIL populations. The age of the samples was estimated

using the transcriptional ruler from [9], the reported ages are relative, where the average age of the control

samples were set to 48 hours. Each point represents one sample, whereof the relative age was determined by

assessing the expression of about 100 genes.

Additional file 9: Table summarizing the statistical power calculations. Outcome of the statistical power

calculations conducted for the RIL population of each treatment (n = 48 RILs per treatment). The outcomes are

ordered per treatment population, and per simulated QTL peak size. All peaks were simulated in random

variation generated by a standard normal distribution. The simulation reports on (i) QTL detection, *e.g.* how

many of the simulated QTL were detected, how many false QTL were reported; (ii) QTL effect size estimation,

mapped effect/simulated effect (reported in quantiles); (iii) QTL location estimation, |mapped location – simulated location|.

Additional file 10: Table of the mapped eQTL in the control, heat stress, and recovery treatment. The eQTL are given per trait (Spot) and treatment. The QTL type, location and confidence interval is listed, as is the significance and effect. The effect is higher in N2 (positive numbers) or higher in CB4856 (negative numbers) loci. Furthermore, information about the affected gene represented by the microarray spot is shown (name, and location).

Additional file 11: Figure of eQTL effect distribution. (**A**) Volcano plots of the eQTL mapped per type (*cis* or *trans*) per treatment. On the x-axis the effect is plotted and on the y-axis the significance of the association is plotted (-log10(p)). Each dot represents a microarray spot and only the significant associations are shown (FDR ≤ 0.05, -log10(p) > 3.9 in all three treatments). (**B**) A histogram of the eQTL effect sizes, per type (*cis* or *trans*) per treatment. Again, the number of significantly associated spots are counted.

Additional file 12: Table listing polymorphisms in eQTL. The polymorphism between and CB4856 and N2 were taken from Thompson *et al*., 2015 and counted in the genes with *cis*-, *trans*-, or no eQTL. Furthermore, the *cis*- and *trans*-eQTL have also been compared based on effect (higher expression in CB4856 or N2). The occurrences of polymorphic genes in the three sets were compared by a chi-squared test.

Additional file 13: A list of *trans*-bands. This table lists the identified *trans*-bands per treatment. The number of affected genes and the number of spots with a *trans*-eQTL these genes were represented by are shown.

Additional file 14: Figure comparing allelic effects of genes with an eQTL between treatments. A scatter plot of the effects of the eQTL of genes with an eQTL found in multiple treatments. Each dot represents a spot. The Pearson correlation values between the different comparisons are: R=0.94 and 0.91 for *cis*- and *trans*-eQTL in control versus heat stress, R=0.96 and 0.93 for *cis*- and *trans*-eQTL in control versus recovery, and R=0.94 and 0.89 for *cis*- and *trans*-eQTL in heat stress versus recovery. The striped diagonal lines are shown as an optical reference.

Additional file 15: Text detailing the calculations on overlap in *cis*- and *trans*-eQTL.

Additional file 16: Figure comparing genes with a *trans*-eQTL in different treatments. (**A**). A plot of the

eQTL location in treatment 1 versus the eQTL-location in treatment 2. Treatment 1 is the first treatment listed in

the legend, for example: the magenta dots represent eQTL where the first treatment is control and the second

treatment is heat stress. The grey lines represent the confidence interval of the eQTL based on a 1.5 drop in –

log10(p). The diagonal band in this plot represents *trans*-eQTL that are regulated from the same location across

treatments. (**B**) The eQTL effects of the *trans*-eQTL shown in (**A**), ordered per chromosome. The striped

diagonal lines are shown as an optical reference.

Additional file 17: A list of the enrichment analysis of genes with an eQTL. Enrichment analysis on the

genes with eQTL across different conditions. The sets were also analyzed for genes with an eQTL with a specific

allelic effect, belonging to a specific *trans*-band and found, and belonging to a specific treatment. The allelic

effects are indicated based on direction (*e.g. cis*-CB4856 means *cis*-eQTL higher expressed in CB4856). The

database used for enrichment (Annotation) and the category (Group), and the number of genes on the array that

are in the group (Genes_in_group) are indicated. Furthermore, the overlap with the cluster (Overlap) and the

Bonferroni-corrected significance of that overlap are shown.

References
712
713   1.   Grishkevich V, Yanai I: **The genomic determinants of genotype x environment interactions in gene**
714   **expression**. *Trends in genetics : TIG* 2013, **29**(8):479-487.
715   2.   GuhaThakurta D, Palomar L, Stormo GD, Tedesco P, Johnson TE, Walker DW, Lithgow G, Kim S,
716   Link CD: **Identification of a novel cis-regulatory element involved in the heat shock response in**
717   **Caenorhabditis elegans using microarray gene expression and computational methods**. *Genome*
718   *research* 2002, **12**(5):701-712.
719   3.   Bakowski MA, Desjardins CA, Smelkinson MG, Dunbar TL, Lopez-Moyado IF, Rifkin SA, Cuomo
720   CA, Troemel ER: **Ubiquitin-mediated response to microsporidia and virus infection in C. elegans**.
721   *PLoS pathogens* 2014, **10**(6):e1004200.
722   4.   Nakad R, Snoek LB, Yang W, Ellendt S, Schneider F, Mohr TG, Rosingh L, Masche AC, Rosenstiel
723   PC, Dierking K *et al*: **Contrasting invertebrate immune defense behaviors caused by a single gene,**
724   **the Caenorhabditis elegans neuropeptide receptor gene npr-1**. *BMC genomics* 2016, **17**:280.
725   5.   Guisbert E, Czyz DM, Richter K, McMullen PD, Morimoto RI: **Identification of a tissue-selective**
726   **heat shock response regulatory network**. *PLoS genetics* 2013, **9**(4):e1003466.
727   6.   Vinuela A, Snoek LB, Riksen JA, Kammenga JE: **Gene expression modifications by temperature-**
728   **toxicants interactions in Caenorhabditis elegans**. *PLoS one* 2011, **6**(9):e24676.
729   7.   Golden TR, Melov S: **Microarray analysis of gene expression with age in individual nematodes**.
730   *Aging cell* 2004, **3**(3):111-124.
731   8.   Kim D, Grun D, van Oudenaarden A: **Dampening of expression oscillations by synchronous**
732   **regulation of a microRNA and its target**. *Nature genetics* 2013, **45**(11):1337-1344.
733   9.   Snoek LB, Sterken MG, Volkers RJ, Klatter M, Bosman KJ, Bevers RP, Riksen JA, Smant G, Cossins
734   AR, Kammenga JE: **A rapid and massive gene expression shift marking adolescent transition in C.**
735   **elegans**. *Sci Rep* 2014, **4**:3912.

10. Jansen RC, Nap JP: **Genetical genomics: the added value from segregation**. *Trends Genet* 2001,
17(7):388-391.
11. Li Y, Breitling R, Jansen RC: **Generalizing genetical genomics: getting added value from**
environmental perturbation. *Trends Genet* 2008, **24**(10):518-524.
12. Li Y, Alvarez OA, Gutteling EW, Tijsterman M, Fu J, Riksen JA, Hazendonk E, Prins P, Plasterk RH,
Jansen RC *et al*: **Mapping determinants of gene expression plasticity by genetical genomics in C.**
elegans. *PLoS genetics* 2006, **2**(12):e222.
13. Hendriks GJ, Gaidatzis D, Aeschimann F, Grosshans H: **Extensive oscillatory gene expression during**
C. elegans larval development. *Molecular cell* 2014, **53**(3):380-392.
14. Capra EJ, Skrovanek SM, Kruglyak L: **Comparative developmental expression profiling of two C.**
elegans isolates. *PloS one* 2008, **3**(12):e4055.
15. Francesconi M, Lehner B: **The effects of genetic variation on gene expression dynamics during**
development. *Nature* 2014, **505**(7482):208-211.
16. Paaby AB, Rockman MV: **Cryptic genetic variation: evolution's hidden substrate**. *Nature reviews*
Genetics 2014, **15**(4):247-258.
17. Ledon-Rettig CC, Pfennig DW, Chunco AJ, Dworkin I: **Cryptic genetic variation in natural**
populations: a predictive framework. *Integrative and comparative biology* 2014, **54**(5):783-793.
18. Iwasaki WM, Tsuda ME, Kawata M: **Genetic and environmental factors affecting cryptic variations**
in gene regulatory networks. *BMC evolutionary biology* 2013, **13**:91.
19. Rodriguez M, Snoek LB, Riksen JA, Bevers RP, Kammenga JE: **Genetic variation for stress-response**
hormesis in C. elegans lifespan. *Exp Gerontol* 2012, **47**(8):581-587.
20. Gaertner BE, Parmenter MD, Rockman MV, Kruglyak L, Phillips PC: **More than the sum of its parts:**
a complex epistatic network underlies natural variation in thermal preference behavior in
Caenorhabditis elegans. *Genetics* 2012, **192**(4):1533-1542.
21. Glauser DA, Chen WC, Agin R, Macinnis BL, Hellman AB, Garrity PA, Tan MW, Goodman MB:
Heat avoidance is regulated by transient receptor potential (TRP) channels and a neuropeptide
signaling pathway in Caenorhabditis elegans. *Genetics* 2011, **188**(1):91-103.
22. Thompson OA, Snoek LB, Nijveen H, Sterken MG, Volkers RJ, Brenchley R, Van't Hof A, Bevers RP,
Cossins AR, Yanai I *et al*: **Remarkably Divergent Regions Punctuate the Genome Assembly of the**
Caenorhabditis elegans Hawaiian Strain CB4856. *Genetics* 2015, **200**(3):975-989.
23. Brenner S: **The genetics of Caenorhabditis elegans**. *Genetics* 1974, **77**(1):71-94.
24. Rockman MV, Kruglyak L: **Recombinational landscape and population genomics of**
Caenorhabditis elegans. *PLoS genetics* 2009, **5**(3):e1000419.
25. Zahurak M, Parmigiani G, Yu W, Scharpf RB, Berman D, Schaeffer E, Shabbeer S, Cope L: **Pre-**
processing Agilent microarray data. *BMC bioinformatics* 2007, **8**:142.
26. Smyth GK, Speed T: **Normalization of cDNA microarray data**. *Methods* 2003, **31**(4):265-273.
27. Benjamini Y, Yekutieli D: **The control of the false discovery rate in multiple testing under**
dependency. *Ann Stat* 2001, **29**(4):1165-1188.
28. Rockman MV, Skrovanek SS, Kruglyak L: **Selection at linked sites shapes heritable phenotypic**
variation in C. elegans. *Science* 2010, **330**(6002):372-376.
29. Harris TW, Baran J, Bieri T, Cabunoc A, Chan J, Chen WJ, Davis P, Done J, Grove C, Howe K *et al*:
WormBase 2014: new views of curated biology. *Nucleic acids research* 2014, **42**(Database
issue):D789-793.
30. Gerstein MB, Lu ZJ, Van Nostrand EL, Cheng C, Arshinoff BI, Liu T, Yip KY, Robilotto R,
Rechtsteiner A, Ikegami K *et al*: **Integrative analysis of the Caenorhabditis elegans genome by the**
modENCODE project. *Science* 2010, **330**(6012):1775-1787.
31. Niu W, Lu ZJ, Zhong M, Sarov M, Murray JI, Brdlik CM, Janette J, Chen C, Alves P, Preston E *et al*:
Diverse transcription factor binding features revealed by genome-wide ChIP-seq in C. elegans.
Genome research 2011, **21**(2):245-254.
32. Tepper RG, Ashraf J, Kaletsky R, Kleemann G, Murphy CT, Bussemaker HJ: **PQM-1 Complements**
DAF-16 as a Key Transcriptional Regulator of DAF-2-Mediated Development and Longevity.
Cell 2013, **154**(3):676-690.
33. Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M: **KEGG: Kyoto Encyclopedia of Genes**
and Genomes. *Nucleic acids research* 1999, **27**(1):29-34.
34. Brunquell J, Morris S, Lu Y, Cheng F, Westerheide SD: **The genome-wide role of HSF-1 in the**
regulation of gene expression in Caenorhabditis elegans. *BMC genomics* 2016, **17**:559.
35. Vinuela A, Snoek LB, Riksen JA, Kammenga JE: **Genome-wide gene expression regulation as a**
function of genotype and age in C. elegans. *Genome Res* 2010, **20**(7):929-937.

36. Jovic K, Sterken MG, Grilli J, Bevers RPJ, Rodriguez M, Riksen JAG, Allesina S, Kammenga JE,
Snoek LB: **A Point Of No Return Leading To Death During Heat-Shock In C. elegans**.
bioRxiv 2017.
37. Gan X, Stegle O, Behr J, Steffen JG, Drewe P, Hildebrand KL, Lyngsoe R, Schultheiss SJ, Osborne EJ,
Sreedharan VT *et al*: **Multiple reference genomes and transcriptomes for Arabidopsis thaliana**.
Nature 2011, **477**(7365):419-423.
38. Keurentjes JJ, Fu J, Terpstra IR, Garcia JM, van den Ackerveken G, Snoek LB, Peeters AJ,
Vreugdenhil D, Koornneef M, Jansen RC: **Regulatory network construction in Arabidopsis by using
genome-wide gene expression quantitative trait loci**. *Proceedings of the National Academy of
Sciences of the United States of America* 2007, **104**(5):1708-1713.
39. Snoek LB, Terpstra IR, Dekter R, Van den Ackerveken G, Peeters AJ: **Genetical Genomics Reveals
Large Scale Genotype-By-Environment Interactions in Arabidopsis thaliana**. *Frontiers in genetics*
2012, **3**:317.
40. Alberts R, Terpstra P, Li Y, Breitling R, Nap JP, Jansen RC: **Sequence polymorphisms cause many
false cis eQTLs**. *PloS one* 2007, **2**(7):e622.
41. Li Y, Breitling R, Snoek LB, van der Velde KJ, Swertz MA, Riksen J, Jansen RC, Kammenga JE:
Global genetic robustness of the alternative splicing machinery in Caenorhabditis elegans.
Genetics 2010, **186**(1):405-410.
42. Thompson O, Edgley M, Strasbourger P, Flibotte S, Ewing B, Adair R, Au V, Chaudhry I, Fernando L,
Hutter H *et al*: **The million mutation project: a new approach to genetics in Caenorhabditis
elegans**. *Genome research* 2013, **23**(10):1749-1762.
43. Volkers RJ, Snoek LB, Hubar CJ, Coopman R, Chen W, Yang W, Sterken MG, Schulenburg H,
Braeckman BP, Kammenga JE: **Gene-environment and protein-degradation signatures
characterize genomic and phenotypic diversity in wild Caenorhabditis elegans populations**. *BMC
biology* 2013, **11**:93.
44. Panousis NI, Gutierrez-Arcelus M, Dermitzakis ET, Lappalainen T: **Allelic mapping bias in RNA-
sequencing is not a major confounder in eQTL studies**. *Genome biology* 2014, **15**(9):467.
45. Chen K, van Nimwegen E, Rajewsky N, Siegal ML: **Correlating gene expression variation with cis-
regulatory polymorphism in Saccharomyces cerevisiae**. *Genome biology and evolution* 2010, **2**:697-
707.
46. Terpstra IR, Snoek LB, Keurentjes JJ, Peeters AJ, van den Ackerveken G: **Regulatory network
identification by genetical genomics: signaling downstream of the Arabidopsis receptor-like
kinase ERECTA**. *Plant physiology* 2010, **154**(3):1067-1078.
47. Smith EN, Kruglyak L: **Gene-environment interaction in yeast gene expression**. *PLoS biology* 2008,
6(4):e83.
48. Grundberg E, Small KS, Hedman AK, Nica AC, Buil A, Keildson S, Bell JT, Yang TP, Meduri E,
Barrett A *et al*: **Mapping cis- and trans-regulatory effects across multiple tissues in twins**. *Nature
genetics* 2012, **44**(10):1084-1089.
49. Stranger BE, Montgomery SB, Dimas AS, Parts L, Stegle O, Ingle CE, Sekowska M, Smith GD, Evans
D, Gutierrez-Arcelus M *et al*: **Patterns of cis regulatory variation in diverse human populations**.
PLoS genetics 2012, **8**(4):e1002639.
50. Gutteling EW, Doroszuk A, Riksen JA, Prokop Z, Reszka J, Kammenga JE: **Environmental influence
on the genetic correlations between life-history traits in Caenorhabditis elegans**. *Heredity* 2007,
98(4):206-213.
51. Gutteling EW, Riksen JA, Bakker J, Kammenga JE: **Mapping phenotypic plasticity and genotype-
environment interactions affecting life-history traits in Caenorhabditis elegans**. *Heredity* 2007,
98(1):28-37.
52. Kammenga JE, Doroszuk A, Riksen JA, Hazendonk E, Spiridon L, Petrescu AJ, Tijsterman M, Plasterk
RH, Bakker J: **A Caenorhabditis elegans wild type defies the temperature-size rule owing to a
single nucleotide polymorphism in tra-3**. *PLoS genetics* 2007, **3**(3):e34.
53. Duveau F, Felix MA: **Role of pleiotropy in the evolution of a cryptic developmental variation in
Caenorhabditis elegans**. *PLoS biology* 2012, **10**(1):e1001230.
54. Sterken MG, Snoek LB, Kammenga JE, Andersen EC: **The laboratory domestication of
Caenorhabditis elegans**. *Trends in genetics : TIG* 2015, **31**(5):224-231.
55. Andersen EC, Bloom JS, Gerke JP, Kruglyak L: **A variant in the neuropeptide receptor npr-1 is a
major determinant of Caenorhabditis elegans growth and physiology**. *PLoS genetics* 2014,
10(2):e1004156.
56. Yvert G, Brem RB, Whittle J, Akey JM, Foss E, Smith EN, Mackelprang R, Kruglyak L: **Trans-acting
regulatory variation in Saccharomyces cerevisiae and the role of transcription factors**. *Nature
genetics* 2003, **35**(1):57-64.

57.   Brem RB, Yvert G, Clinton R, Kruglyak L: **Genetic dissection of transcriptional regulation in**
855   **budding yeast**. *Science* 2002, **296**(5568):752-755.
856   58.   West MA, Kim K, Kliebenstein DJ, van Leeuwen H, Michelmore RW, Doerge RW, St Clair DA:
857   **Global eQTL mapping reveals the complex genetic architecture of transcript-level variation in**
858   **Arabidopsis**. *Genetics* 2007, **175**(3):1441-1450.

A

B

Genes differentially expressed in:

1. **control** versus **heat stress** & **recovery**

2. **Control** & **recovery** versus **heat stress**

3. All three conditions when compared