Disparate patterns of thermal adaptation between life stages in temperate vs. tropical

Drosophila melanogaster

Brent L. Lockwood*, Tarun Gupta, and Rosemary Scavotto

Department of Biology, The University of Vermont, Burlington, VT 05405

*Corresponding author: Brent.Lockwood@uvm.edu

Summary

Many terrestrial ectothermic species exhibit limited variation in upper thermal tolerance across

latitude. However, these trends may not signify limited adaptive capacity to increase thermal

tolerance in the face of climate change. Instead, thermal tolerance may be similar among

populations because behavioral thermoregulation by mobile organisms or life stages may buffer

natural selection for thermal tolerance. We compared thermal tolerance of adults and embryos

among natural populations of *Drosophila melanogaster* from a broad range of thermal habitats

around the globe to assess natural variation of thermal tolerance in mobile vs. immobile life

stages. We found no variation among populations in adult thermal tolerance, but embryonic

thermal tolerance was higher in tropical than in temperate regions. Average maximum

temperature of the warmest month of the year predicted embryonic thermal tolerance in tropical

but not temperate sites. We further report that embryos live closer to their upper thermal limits

than adults—i.e., thermal safety margins are smaller for embryos than adults. F1 hybrid embryos

from crosses between temperate and tropical populations had thermal tolerance that matched that

of tropical embryos, suggesting phenotypic dominance of heat-tolerant alleles. Together our

findings demonstrate that embryonic thermal tolerance readily evolves and suggest that selection

for thermal tolerance may be limited in adults. Further, our results suggest that thermal traits

should be measured across life stages in order to better predict adaptive limits.

Key words: *Drosophila*, embryo, heat tolerance, phenotypic dominance, thermal adaptation,

thermal safety margin

Introduction

Extreme temperatures, which may be encountered at the edge of a species' geographic range

(Hilbish *et al.* 2010) or episodically during the hottest or coldest days of the year (Hoffmann

2010; Kingsolver, Diamond & Buckley 2013; Dowd, King & Denny 2015; Buckley & Huey

2016), can cause populations to experience mortality (Helmuth *et al.* 2002; Denny, Miller &

Harley 2006) and ultimately lead to thermal adaptation (Lenski & Bennett 1993; Mongold,

Bennett & Lenski 1999; Hangartner & Hoffmann 2015). However, recent work suggests that

thermal adaptation of upper thermal limits might be evolutionarily constrained (Hoffmann,

Chown & Clusella-trullas 2013; Schou *et al.* 2014; Hangartner & Hoffmann 2015; Kristensen *et*

al. 2015; van Heerwaarden, Kellermann & Sgrò 2016), such that the evolution of increased heat

tolerance might be a relatively slow process that takes many millions of years to achieve

(Kellermann *et al.* 2012). If this is the case, global climate change, which has led to rapid

increases in mean temperatures and the frequency of extreme thermal events (Katz & Brown

1992; Meehl *et al.* 2000; Cai *et al.* 2014), may cause shifts in geographic distributions (Rank &

Dahlhoff 2002; Burrows *et al.* 2011; Thomas *et al.* 2012; Sunday, Bates & Dulvy 2012) as

populations may not be able to adapt fast enough to persist in hotter environments (Jezkova &

Wiens 2016).

But thermal adaptation depends on the strength of selection (Bennett, Lenski & Mittler

1992; Rudolph *et al.* 2010), and studies that focus on thermal tolerance of mobile organisms or

life stages may overestimate the degree to which these organisms encounter thermal selection in

nature. In other words, thermal safety margins—i.e., the difference between upper thermal limits

and maximum habitat temperature—may be larger than predicted because thermal environmental

heterogeneity allows mobile organisms to avoid thermal extremes via behavioral

thermoregulation (Dillon *et al.* 2009; Gunderson & Leal 2012; Buckley, Ehrenberger &

Angilletta 2015; Llewelyn *et al.* 2016; Munoz *et al.* 2016). To date, there have been relatively

few studies that examine thermal tolerance in immobile organisms or life stages, particularly in

the terrestrial realm (Angilletta *et al.* 2013; MacLean *et al.* 2016), and immobile organisms may

represent ideal study systems to investigate the evolutionary potential of thermal tolerance. In

support of this conjecture, broad scale patterns of thermal tolerance are more tightly correlated

with habitat temperatures in marine systems than in terrestrial systems (Sunday, Bates & Dulvy

2011), perhaps due to the more limited range of thermal microhabitats in the marine realm

(Denny *et al.* 2011) that makes behavioral thermoregulation a less effective buffering mechanism.

Here we sought to compare adult and embryonic heat tolerance among populations of

fruit flies, *Drosophila melanogaster*, from a broad range of thermal habitats across the world to

ascertain the degree to which thermal selection has shaped the evolution of thermal tolerance

across immobile vs. mobile life stages. Adult thermal tolerance has been extensively studied in

natural populations of *D. melanogaster* (Hoffmann, Anderson & Hallas 2002; Hoffmann &

Weeks 2007; Sgro *et al.* 2010; Adrion, Hahn & Cooper 2015; Buckley & Huey 2016), but to a

large extent the thermal physiology of the early embryonic life stage of *D. melanogaster* has not

been characterized in natural populations (Kristensen *et al.* 2015). Studies of lab-bred *D.*

melanogaster have shown that early embryos (0 – 2 hours post-fertilization) are more thermally

sensitive than later stages (Walter, Biessmann & Petersen 1990), perhaps due to the reduced

heat-shock response in early embryos (Graziosi *et al.* 1980; Welte *et al.* 1993). Thus, we

compared heat tolerance of adults and early stage embryos to determine whether or not

differences in thermal sensitivity, as well as mobility, lead to different patterns of thermal

adaptation across life stages. The thermal environment of *D. melanogaster* can change rapidly

(+18°C h^{-1}) and reach extreme values (> 40°C) (Feder, Blair & Figueras 1997; Terblanche *et al.*

2011). Therefore, we designed our thermal stress experiments to mimic sudden (acute) changes

in temperature that are characteristic of the variable thermal environments that flies experience in

nature (Terblanche *et al.* 2011). We report higher embryonic thermal tolerance in tropical

(hotter) vs. temperate (cooler) populations but no difference in adult thermal tolerance, and thus

we demonstrate that selection for thermal tolerance likely varies across life stages. Moreover, our

data suggest that there is significant adaptive variation for upper thermal tolerance in natural

populations in the earliest and most thermally sensitive life stage.

Materials and methods

Fly strains

We obtained 19 isofemale genetic lines that were collected from temperate locations in the USA

as a generous gift from B.S. Cooper and K.L. Montooth: 6 lines from Raleigh, NC (NC); 5 lines

from Beasley Orchard, IN (IN); and 8 lines from East Calais, VT (VT). The creation of these

lines has been previously described (Cooper, Hammad & Montooth 2014). We obtained 5

isofemale lines from the Drosophila Species Stock Center at the University of California, San

Diego that were collected from tropical locations around the world: 1 line each from Accra,

Ghana (GH); Mumbai, India (MU); Guam, USA (GU); Chiapas, Mexico (CH); and Monkey Hill,

St. Kitts (SK). Geographic coordinates of collection locations and stock numbers of isofemale

lines are provided in Supplementary Table S1. We maintained flies under common-garden

conditions on cornmeal-yeast-molasses medium at 25°C on a 12:12 light cycle for at least two

generations prior to measuring thermal tolerance.

Adult critical thermal maximum (CT$_{max}$)

We assayed thermal tolerance of adult male flies by measuring the temperature at which flies

incurred a loss of motor response along a heat ramp—i.e., the critical thermal maximum (CT_{max}).

3 to 5 day-old adult male flies were individually placed into glass vials with rubber stoppers and

submerged in a water bath that was programmed to increase $0.1°C$ min^{-1} upwards from $25°C$. We

chose this rate of temperature increase based on previously published studies (Chown *et al.*

2009; Sgro *et al.* 2010; Kellermann *et al.* 2012) and to mimic the variable thermal environments

that flies encounter in nature (Terblanche *et al.* 2011). Flies were regularly checked for

responsiveness along the heat ramp by gently tapping the vial, and the temperature at which a fly

lost the ability to move was recorded. We scored CT_{max} for each genotype via a least-squares

regression model of the logistic equation among 10 flies per genotype and extrapolated CT_{max}

from the inflection points of the logistic curves. We conducted these curve fitting analyses in

GraphPad Prism 7 for Mac OS X (GraphPad Software, La Jolla, CA).

Embryonic thermal tolerance (LT_{50})

We assayed embryonic thermal tolerance by measuring survival (hatching success) of early stage

embryos, 0 to 1 h post-fertilization, exposed to a 45-minute heat treatment across a range of

temperatures, from $25°C$ to $42°C$. We chose this assay method because embryos do not possess

behavioral characteristics that would permit the assessment of thermal tolerance via CT_{max}, and

measuring hatching success allowed us to overcome this obstacle. Previous studies conducted in

a wide range of species have shown that temperature adaptive variation in survival following

heat stress follows the same trends as CT_{max} (Huey, Patridge & Fowler 1991; Gilchrist, Huey &

Partridge 1997; Stillman & Somero 2000; Somero 2002; Stenseng *et al.* 2005; Braby & Somero

2006; Miller & Stillman 2012; Dowd & Somero 2013). We designed our heat treatments to

mimic sudden increases in temperature that frequently occur in nature where the temperature of

necrotic fruit can increase rapidly on hot days (Feder *et al.* 1997; Terblanche *et al.* 2011). 3 to 5

day-old adult flies were allowed to mate and lay eggs on grape juice agar plates for 1 h at 25°C.

Egg plates were then wrapped in Parafilm, submerged in a water bath, and heat shocked for 45

minutes. The rate of temperature change averaged +0.4°C min^{-1}. This rate of increase is within

the range of measured rates of change in nature (Feder *et al.* 1997). Following heat shock, 20

eggs were transferred on a piece of grape juice agar to fresh food vials and placed at 25°C.

Hatching success was scored as the proportion of larvae that successfully hatched by 48 h. We

conducted 4 to 6 replicate treatments at each of 9 temperatures (25°C, 28°C, 30°C, 32°C, 34°C,

36°C, 38°C, 40°C, and 42°C) for each genotype. We used these data to calculate the lethal

temperature at which 50% of the embryos failed to hatch (LT_{50}) via a least-squares regression

model of the logistic equation. In our logistic model, we allowed the y-intercept to vary between

0 and 1 and extrapolated the LT_{50} from the inflection point of the logistic curve fit. This

approach allowed us to infer thermal tolerance independently from other confounding factors

that may influence the measurement of hatching success, such as the presence of unfertilized

eggs. We conducted these curve fitting analyses in GraphPad Prism 7.

PCA of bioclimatic variables

We examined the correspondence of adult and embryonic thermal tolerance to local climate

conditions by examining the variation among our collection sites in 19 bioclimatic (Bioclim)

variables. Bioclim variables represent biologically meaningful climate predictors that are derived

from monthly temperature and rainfall data from the years 1960 to 1990 (Busby 1991). We

downloaded Bioclim variables from the WorldClim database (Hijmans *et al.* 2005)

(www.worldclim.org) that corresponded to the GPS coordinates of the collection sites of each

population (see Table S1). Because Bioclim variables tend to covary, we used a principal

components analysis (PCA) to characterize the major axes of variation in climate among our

sites. We then conducted a multiple least-squares regression analysis to assess the degree to

which variation in thermal tolerance could be explained by the primary principal components of

these climate data. We conducted these analyses in R version 3.3.2 (Team 2016).

Statistical comparisons of thermal tolerance and thermal safety margins

We compared adult (CT_{max}) and embryonic (LT_{50}) thermal tolerances among temperate sites (VT,

IN, and NC) and all tropical sites pooled together (CH, SK, GH, MU, and GU) with ANOVA.

Pairwise differences were assessed with Tukey's multiple comparison post-hoc test. We report

our data as mean values with 95% confidence intervals. We calculated thermal safety margins as

the difference between thermal tolerance (adult CT_{max} or embryo LT_{50}) and maximum

temperature of the warmest month (T_{max}) at each site (see above for description of Bioclim data).

Fine-scale spatial temperature data are not available for these collection sites, but while T_{max} may

not perfectly match the thermal environment experienced by flies, variation in T_{max} should reflect

relative differences in the thermal environments among locations. We compared thermal safety

margins across latitude, as in Kellerman et al. (Kellermann *et al.* 2012), between adults and

embryos using non-linear least-squares regression followed by the extra sum-of-squares F-test to

compare model fits. In a separate analysis, we assessed the main effects of region (temperate vs.

tropical), life stage (adult vs. embryo), and their interaction on thermal safety margins via a 2-

way ANOVA. Least-squares linear regression was used to assess the relationship between

embryonic thermal tolerance and significant climate variables. ANCOVA was used to assess the

difference in slopes of regression lines fit to data from temperate vs. tropical sites. We assessed

the genetic basis of differences in embryonic thermal tolerance by conducting reciprocal crosses

between the two parental strains that had the highest and lowest LT_{50}, Chiapas, MX (CH) and

Vermont, USA strain #12 (VT-12), respectively, and measured thermal tolerance of F1 progeny.

We used logistic models to fit the hatching success data, as described above, and compared

LT_{50}s of the parental strains and their F1 progeny by an extra sum-of-squares F-test of the

extrapolated LT_{50}s. We conducted these analyses in GraphPad Prism 7.

Results

Thermal tolerance and thermal safety margins across life stages

We found no difference in adult thermal tolerance among all sites (Figs. 1B and 1C; ANOVA,

$F_{3,9} = 2.378$, $P = 0.1375$), with an overall mean CT_{max} (± 95% C.I.) of 38.8 ± 0.45°C. Embryonic

thermal tolerance (LT_{50}) did not differ among the three temperate sites but was significantly

higher in tropical vs. temperate embryos (Figs. 1D and 1E; ANOVA, $F_{3,20} = 10.16$, $P = 0.0003$;

Tukey's test, VT vs. IN, $q = 2.428$, $P = 0.3416$, VT vs. NC, $q = 0.4268$, $P = 0.9902$, IN vs. NC, q

$= 2.666$, $P = 0.2656$, tropical vs. VT, $q = 6.909$, $P = 0.0005$, tropical vs. IN, $q = 4.04$, $P = 0.0444$,

tropical vs. NC, $q = 4.04$, $P = 0.0005$). Overall, tropical embryos were more heat tolerant; the

average LT_{50} was approximately 1°C higher in tropical embryos (35.8 ± 0.45°C) than in

temperate embryos (34.88 ± 0.18°C). There was no significant relationship between adult CT_{max}

and embryo LT_{50} (Fig. S1; Least-squares linear regression, $R^2 = 0.0005$).

Thermal safety margins—i.e., the difference between thermal tolerance (CT_{max} or LT_{50})

and maximum habitat temperature (T_{max})—were consistently smaller for embryos than adults and

smallest in the tropics. This pattern was consistent regardless of whether thermal safety margins

were compared across latitude (Fig. 2A; Extra sum-of-squares F-test on fitness of two curves,

$F_{3,32} = 15.95$, $P < 0.0001$; Non-linear least-squares regression, embryo, $R^2 = 0.8019$, $y =$

$0.01563x^2 - 0.7249x + 9.97$, adult, $R^2 = 0.6359$, $y = 0.01437x^2 - 0.6424x + 12.44$) or by region

(temperate vs. tropical) (Fig. 2B; ANOVA, main effect of region, $F_{1,34} = 6.482$, $P = 0.0156$, main

effect of life stage, $F_{1,34} = 11.15$, $P = 0.002$, region x life stage interaction, $F_{1,34} = 0.0006$, $P =$

0.98).

Climate variability and thermal tolerance

While there is a consistent difference between our temperate and tropical sites in mean annual

temperature, VT = 5.4°C, IN = 10.8°C, NC = 15.4°C, and all tropical = 26.1 ± 1.4°C (mean ±

95% C.I.), a principal components analysis of 19 Bioclim variables indicated that the greatest

difference in climate among the collection locations was in temperature seasonality—defined as

the yearly standard deviation in temperature. PC1 explained 95.6% of the variation in climate

data among the collection sites, and this pattern largely characterized the differences in climate

between temperate vs. tropical sites (Fig. S2). Temperature seasonality was the variable with the

top component loading of PC1, explaining 98% of the variation in PC1. There was a significant

negative relationship between PC1 and embryonic heat tolerance (Multiple least-squares linear

regression with PC1 and PC2 as covariates, $F_{1,4} = 8.1474$, $P = 0.0462$), but this pattern was

driven by the bimodal distribution of LT_{50}s between temperate vs. tropical sites and was not

consistent with temperature seasonality within each geographic region (Fig. S3). PC2 explained

4% of the variation in climate data among collection sites, mostly characterizing the differences

in climate among tropical sites (Fig. S2). Annual precipitation, precipitation of the wettest

quarter, and precipitation of the wettest month were the top three component loadings of PC2,

explaining 49%, 40%, and 8% of variation in PC2, respectively. But there was no significant

relationship between PC2 and embryonic heat tolerance (Least-squares linear regression, $F_{1,4} =$

0.367, $P = 0.577$).

Maximum temperature of the warmest month (i.e., maximum habitat temperature or T_{max})

spanned a range of 8.4°C among all sites, from 25.7°C in Vermont, USA (VT) to 34.1°C in

Chiapas, MX (CH). While this variation in T_{max} explained only a small fraction of the variation

in Bioclim variables among sites (0.0016% of the variation in PC1 and 0.00015% of the

variation in PC2), previous studies have shown T_{max} to be correlated with adult heat tolerance

(CT_{max}) among many species of *Drosophila* (Kellermann *et al.* 2012). In contrast to this previous

work, our populations of *D. melanogaster* showed no significant relationship between adult heat

tolerance and T_{max} (Fig. S4; Least-squares linear regression, $R^2 = 0.055$). However, the

embryonic life stage exhibited a different pattern than the adults, and the relationship between

embryonic heat tolerance and T_{max} was distinct between temperate and tropical regions (Fig. 3;

ANCOVA, $F_{1,4} = 10.26$, $P = 0.0328$). Among temperate populations there was a 6°C range in

T_{max}, but this produced no correlated response in the thermal tolerance of embryos (Fig. 3; Least-

squares regression, $R^2 = 0.0015$, $P = 0.9751$, $y = 0.00282x + 34.82$). But among tropical

populations, the approximate 4°C range in T_{max} corresponded to a positive relationship between

embryonic thermal tolerance and T_{max} (Fig. 3; Least-squares regression, $R^2 = 0.9478$, $P = 0.0051$,

$y = 0.2199x + 28.75$).

Embryonic thermal tolerance in F1 progeny from Chiapas x Vermont

Offspring from reciprocal genetic crosses between the most heat tolerant tropical genotype (CH)

and the least heat tolerant temperate genotype (VT-12) had thermal tolerances that closely

resembled that of the heat tolerant CH genotype, regardless of the direction of the cross (Fig. 4).

Embryonic LT_{50}s of F1 progeny of both crosses (CH♀ x VT♂ = 35.83°C and VT♀ x CH♂ =

35.80°C) were statistically indistinguishable from the LT_{50} of CH (36.24°C) but significantly

higher than the LT_{50} of VT-12 (34.23°C; Fig. 4; Logistic model, Extra sum-of-squares F-test on

lower LT_{50} of VT-12, $F_{3,166} = 6.695$).

Discussion

Despite the potential for thermal adaptation across the broad range of thermal habitats

represented in this study, our data suggest that natural selection on thermal tolerance does not act

equally across life stages in *D. melanogaster*. Rather, we provide evidence of adaptive variation

in upper thermal limits in the thermally sensitive and immobile embryonic life stage but not in

the more thermally tolerant and mobile adult stage. This is perhaps not surprising, given that

lower thermal tolerance in early embryos translates into smaller thermal safety margins. Thus,

we predict that embryos encounter lethal temperatures more frequently than adults, particularly

because embryos lack the ability to behaviorally avoid thermally stressful conditions, and this

likely drives divergence in embryonic thermal tolerance between temperate and tropical

populations.

Recent estimates of divergence in adult thermal tolerance among populations of *D.*

melanogaster have brought into question the degree of adaptive potential in upper thermal limits

in this species, as comparisons of populations across latitude have yielded mixed results

depending on assay methods (Sgro *et al.* 2010) and the laboratory in which thermal tolerance

was measured (Hoffmann *et al.* 2002; Hoffmann 2010; Buckley & Huey 2016). Our estimates of

D. melanogaster adult male CT_{max} are consistent with previous reports (Gilchrist *et al.* 1997;

Chown *et al.* 2009; Kellermann *et al.* 2012), and while we report novel findings on the

adaptation of embryonic thermal tolerance, our results are not unprecedented. Coyne et al.

(Coyne, Bundgaard & Prout 1983) reported a similar discrepancy in thermal adaptation between

mobile and immobile life stages among populations of *Drosophila pseudoobscura*—pupal

thermal tolerance, but not adult thermal tolerance, was higher in populations from warmer

locations. The interplay of population genetic factors in natural populations of *D. melanogaster*

suggest that this species harbors a high level of genetic diversity (Karasov, Messer & Petrov

2010) and that natural selection has led to allelic divergence among populations across the

genome (Hoffmann & Weeks 2007; Fabian *et al.* 2012; Adrion *et al.* 2015). In light of these

trends in population genomics, and the adaptive variation in embryonic thermal tolerance

presented in this study, it seems probable that there is significant natural variation of upper

thermal limits in *D. melanogaster* but that this variation may only be revealed in the embryonic

and other immobile life stages.

It is important to note that lab selection experiments in *D. melanogaster*, *Escherichia coli*,

and marine copepods (*Tigriopus californicus*) that imposed strong selection on thermal tolerance

reported significant potential for adaptation of upper thermal limits, but the response to selection

eventually plateaued after many generations, presumably when standing genetic diversity had

been exhausted (Huey *et al.* 1991; Gilchrist *et al.* 1997; Gilchrist & Huey 1999; Rudolph *et al.*

2010; Kelly, Sanford & Grosberg 2012; Hangartner & Hoffmann 2015). Thus, there may likely

be potential for adaptation of upper thermal limits, and in natural populations greater levels of

standing genetic variation may be able to sustain adaptive responses to thermal selection.

This study characterizes thermal tolerance among populations that span a large portion of

the *D. melanogaster* biogeographic range in the northern hemisphere, and while there is clear

evidence of adaptation of embryonic thermal tolerance between temperate and tropical regions,

the patterns of thermal adaptation are not consistent within each region. Tropical embryos

sampled from locations with higher T_{max} showed higher thermal tolerances, yet temperate

populations did not follow this trend. Why were there no observed differences in embryonic

thermal tolerance among temperate populations when temperate sites spanned a broader range of

thermal habitats than tropical populations? It is possible that gene flow between Vermont,

Indiana, and North Carolina overwhelms local adaptation, but recent studies show evidence of

adaptive divergence among *D. melanogaster* populations in eastern North America (Fabian *et al.*

2012; Bergland *et al.* 2016; Machado *et al.* 2016). Therefore, a more likely explanation is that

seasonal fluctuations in the activity of temperate populations (Cogni *et al.* 2014), may limit the

frequency at which temperate embryos encounter thermal selection. In addition, spatial and

temporal microclimatic variability in temperate sites may provide more choices for females to

lay their eggs at permissive temperatures (Allemand & David 1976; Dahlgaard, Hasson &

Loeschcke 2001; Huey & Pascual 2009; Dillon *et al.* 2009). We also note that our data constitute

thermal tolerances of multiple isofemale lines from each of the three temperate sites but just one

isofemale line from each of the five tropical sites. Thus, we have not captured the full range of

genetic variation at each tropical site. The fact that the tropical flies originated from

geographically distant locations suggests that our observed differences in embryonic thermal

tolerance between temperate and tropical populations represent consistent patterns of thermal

adaptation. However, our observation that embryonic thermal tolerance positively correlates with

maximum temperature at tropical sites is a result that warrants further investigation. It remains to

be determined whether or not this pattern will hold when a greater sample of genetic diversity is

surveyed at these topical locations.

Investigations into the genetic basis of variation in thermal tolerance in natural

populations have yielded somewhat conflicting results, depending on the species (Pereira,

Barreto & Burton 2014; Sato *et al.* 2015) and the way thermal tolerance was measured (Gilchrist

& Huey 1999; Mitchell & Hoffmann 2010; Hangartner & Hoffmann 2015). In this study,

embryonic LT_{50} spanned a range of approximately 2°C among all genotypes, and crosses

between genotypes at the high and low ends of this range produced F1 progeny that resembled

the phenotype of the heat-tolerant parent. This result indicates phenotypic dominance of heat-

tolerant alleles, at least for Chiapas vs. Vermont-12 embryos. This is a surprising result for two

reasons. First, this inheritance pattern is not expected because thermal physiological traits are

complex and depend on multiple interacting factors, such as lipid membrane composition

(Dahlhoff & Somero 1993; Cooper *et al.* 2012, 2014), protein stability (Lockwood & Somero

2012; Fields *et al.* 2015; Leuenberger *et al.* 2017), and transcriptional and translational responses

(Leemans *et al.* 2000; Lockwood, Sanders & Somero 2010; Tomanek & Zuzow 2010; Chen,

Nolte & Schlotterer 2015; Kristensen *et al.* 2016). Accordingly, previous work has shown that *D.*

melanogaster adult and larval thermal tolerance are quantitative traits influenced by at least 7

loci (Morgan & Mackay 2006; Sambucetti *et al.* 2013). Second, because we examined early *D.*

melanogaster embryos at 0 – 1 h post-fertilization (hpf), and zygotic genome activation has been

reported to not begin until approximately 2 hpf in this species (Tadros & Lipshitz 2009; Blythe

& Wieschaus 2015), we expected thermal tolerance to be determined by maternal effects. Instead,

we found embryonic thermal tolerance to match that of the Chiapas strain regardless of maternal

genotype. This suggests that either (1) the zygotic genome is being activated earlier than

expected in response to heat shock (Graziosi *et al.* 1980), which would reveal adaptive variation

in zygotic gene expression, or (2) that the effect is mediated at the level of the chromosomes,

perhaps due to thermally-induced DNA damage (Yao & Somero 2012) that differentially affects

different genotypes (Svetec *et al.* 2016). Either way, the unknown genetic basis of embryonic

thermal tolerance warrants future study.

Author's Contributions

BL conceived the ideas and designed the methodology; BL, TG, and RS collected the data; BL

analyzed the data; BL wrote the manuscript. All authors gave final approval for publication.

Acknowledgements

We thank Kristi Montooth, Brandon Cooper, Cole Julick and the UC San Diego Drosophila

Species Stock Center for providing fly stocks. We thank Andrew Nguyen for R code that was

helpful in generating the map shown in Figure 1. We thank Melissa Pespeni and Brandon Cooper

for helpful comments on this manuscript. This work was funded by the University of Vermont.

Data Accessibility

Fly stock information is included in Table S1, including geographical coordinates of sampling

locations, stock numbers, and thermal tolerance data.

References

Adrion, J.R., Hahn, M.W. & Cooper, B.S. (2015) Revisiting classic clines in *Drosophila*

melanogaster in the age of genomics. *Trends in Genetics*, **31**, 434–444.

Allemand, R. & David, J.R. (1976) The circadian rhythm of oviposition in *Drosophila*

melanogaster: a genetic latitudinal cline in wild populations. *Experientia*, **32**, 1403–1404.

Angilletta, M.J., Zelic, M.H., Adrian, G.J., Hurliman, a. M. & Smith, C.D. (2013) Heat

tolerance during embryonic development has not diverged among populations of a

widespread species (*Sceloporus undulatus*). *Conservation Physiology*, **1**, cot018-cot018.

Bennett, A.F., Lenski, R.E. & Mittler, J.E. (1992) Evolutionary Adaptation to Temperature . I .

Fitness Responses of *Escherichia coli* to Changes in its Thermal Environment. *Evolution*,

46, 16–30.

Bergland, A.O., Tobler, R., Gonzalez, J., Schmidt, P. & Petrov, D. (2016) Secondary contact and

local adaptation contribute to genome-wide patterns of clinal variation in *Drosophila*

melanogaster. *Molecular Ecology*, **25**, 1157–1174.

Blythe, S.A. & Wieschaus, E.F. (2015) Coordinating Cell Cycle Remodeling with

Transcriptional Activation at the *Drosophila* MBT. *Current Topics in Developmental*

Biology, **113**, 113–148.

Braby, C.E. & Somero, G.N. (2006) Following the heart: temperature and salinity effects on

heart rate in native and invasive species of blue mussels (genus *Mytilus*). *Journal of*

Experimental Biology, **209**, 2554–2566.

Buckley, L.B., Ehrenberger, J.C. & Angilletta, M.J. (2015) Thermoregulatory behaviour limits

local adaptation of thermal niches and confers sensitivity to climate change. *Functional*

Ecology, **29**, 1038–1047.

Buckley, L.B. & Huey, R.B. (2016) How Extreme Temperatures Impact Organisms and the

Evolution of their Thermal Tolerance. *Integrative and Comparative Biology* , 1–12.

Burrows, M.T., Schoeman, D.S., Buckley, L.B., Moore, P., Poloczanska, E.S., Brander, K.M.,

Brown, C., Bruno, J.F., Duarte, C.M., Halpern, B.S., Holding, J., Kappel, C. V, Kiessling,

W., O'Connor, M.I., Pandolfi, J.M., Parmesan, C., Schwing, F.B., Sydeman, W.J. &

Richardson, A.J. (2011) The pace of shifting climate in marine and terrestrial ecosystems.

Science, **334**, 652–655.

Busby, J.R. (1991) BIOCLIM-a bioclimate analysis and prediction system. *Plant Protection*

Quarterly (Australia).

Cai, W., Borlace, S., Lengaigne, M., van Rensch, P., Collins, M., Vecchi, G., Timmermann, A.,

Santoso, A., McPhaden, M.J., Wu, L., England, M.H., Wang, G., Guilyardi, E. & Jin, F.-F.

(2014) Increasing frequency of extreme El Niño events due to greenhouse warming. *Nature*

Climate Change, **5**, 1–6.

Chen, J., Nolte, V. & Schlotterer, C. (2015) Temperature-Related Reaction Norms of Gene

Expression: Regulatory Architecture and Functional Implications. *Molecular Biology and*

Evolution.

Chown, S.L., Jumbam, K.R., Sørensen, J.G. & Terblanche, J.S. (2009) Phenotypic variance,

plasticity and heritability estimates of critical thermal limits depend on methodological

context. *Functional Ecology*, **23**, 133–140.

Cogni, R., Kuczynski, C., Koury, S., Lavington, E., Behrman, E.L., O'Brien, K.R., Schmidt, P.S.

& Eanes, W.F. (2014) The Intensity of Selection Acting on the Couch Potato Gene -

Spatial-Temporal Variation in a Diapause Cline. *Evolution*, **68**, 538–548.

Cooper, B.S., Hammad, L. a., Fisher, N.P., Karty, J. a. & Montooth, K.L. (2012) In a variable

thermal environment selection favors greater plasticity of cell membranes in *Drosophila*

melanogaster. *Evolution*, **66**, 1976–1984.

Cooper, B.S., Hammad, L. a & Montooth, K.L. (2014) Thermal adaptation of cellular

membranes in natural populations of *Drosophila melanogaster*. *Functional Ecology*, **28**,

886–894.

Coyne, J. a., Bundgaard, J. & Prout, T. (1983) Geographic Variation of Tolerance to

Environmental Stress in *Drosophila pseudoobscura*. *The American Naturalist*, **122**, 474.

Dahlgaard, J., Hasson, E. & Loeschcke, V. (2001) Behavioral differentiation in oviposition

activity in *Drosophila buzzatii* from highland and lowland populations in Argentina:

plasticity or thermal adaptation? *Evolution*, **55**, 738–47.

Dahlhoff, E. & Somero, G.N. (1993) Effects of temperature on mitochondria from Abalone

(genus *Haliotis*) - adaptive plasticity and its limits. *Journal of Experimental Biology*, **185**,

151–168.

Denny, M.W., Dowd, W.W., Bilir, L. & Mach, K.J. (2011) Spreading the risk: Small-scale body

temperature variation among intertidal organisms and its implications for species

persistence. *Journal of Experimental Marine Biology and Ecology*, **400**, 175–190.

Denny, M.W., Miller, L.P. & Harley, C.D.G. (2006) Thermal stress on intertidal limpets: long-

term hindcasts and lethal limits. *The Journal of Experimental Biology*, **209**, 2420–2431.

Dillon, M.E., Wang, G., Garrity, P. a & Huey, R.B. (2009) Review: Thermal preference in

Drosophila. *Journal of thermal biology*, **34**, 109–119.

Dowd, W.W., King, F.A. & Denny, M.W. (2015) Thermal variation, thermal extremes and the

physiological performance of individuals. *Journal of Experimental Biology*, **218**, 1956–

1967.

Dowd, W.W. & Somero, G.N. (2013) Behavior and survival of *Mytilus* congeners following

episodes of elevated body temperature in air and seawater. *The Journal of Experimental*

Biology, **216**, 502–14.

Fabian, D.K., Kapun, M., Nolte, V., Kofler, R., Schmidt, P.S., Schlötterer, C. & Flatt, T. (2012)

Genome-wide patterns of latitudinal differentiation among populations of *Drosophila*

melanogaster from North America. *Molecular Ecology*, **21**, 4748–69.

Feder, M.E., Blair, N. & Figueras, H. (1997) Natural thermal stress and heat shock protein

expression in *Drosophila* larvae and pupae. *Functional Ecology*, **11**, 90–100.

Fields, P.A., Dong, Y., Meng, X. & Somero, G.N. (2015) Adaptations of protein structure and

function to temperature: there is more than one way to "skin a cat." *Journal of Experimental*

Biology, **218**, 1801–1811.

Gilchrist, G.W. & Huey, R.B. (1999) The direct response of *Drosophila melanogaster* to

selection on knockdown temperature. *Heredity*, **83 (Pt 1)**, 15–29.

Gilchrist, G.W., Huey, R. & Partridge, L. (1997) Thermal Sensitivity of *Drosophila*

melanogaster: Evolutionary Responses of Adults and Eggs to Laboratory Natural Selection

at Different Temperatures. *Physiological Zoology*, **70**, 403–414.

Graziosi, G., Micali, F., Marzari, R., de Cristini, F. & Savoini, A. (1980) Variability of response

of early *Drosophila* embryos to heat shock. *The Journal of Experimental Zoology*, **214**,

141–145.

Gunderson, A.R. & Leal, M. (2012) Geographic variation in vulnerability to climate warming in

a tropical Caribbean lizard. *Functional Ecology*, **26**, 783–793.

Hangartner, S. & Hoffmann, A. a. (2015) Evolutionary potential of multiple measures of upper

thermal tolerance in *Drosophila melanogaster*. *Functional Ecology*, **30**, n/a-n/a.

van Heerwaarden, B., Kellermann, V. & Sgrò, C.M. (2016) Limited scope for plasticity to

increase upper thermal limits. *Functional Ecology*.

Helmuth, B., Harley, C.D.G., Halpin, P.M., O'Donnell, M., Hofmann, G.E. & Blanchette, C.A.

(2002) Climate change and latitudinal patterns of intertidal thermal stress. *Science*, **298**,

1015–1017.

Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G. & Jarvis, A. (2005) Very high resolution

interpolated climate surfaces for global land areas. *International Journal of Climatology*, **25**,

1965–1978.

Hilbish, T.J., Brannock, P.M., Jones, K.R., Smith, A.B., Bullock, B.N., Wethey, D.S. & Smith,

AB. (2010) Historical changes in the distributions of invasive and endemic marine

invertebrates are contrary to global warming predictions: the effects of decadal climate

oscillations. *Journal of Biogeography*, **37**, 423–431.

Hoffmann, A.A. (2010) Physiological climatic limits in *Drosophila*: patterns and implications.

Journal of Experimental Biology, **213**, 870–880.

Hoffmann, A.A., Anderson, A. & Hallas, R. (2002) Opposing clines for high and low

temperature resistance in *Drosophila melanogaster*. *Ecology Letters*, **5**, 614–618.

Hoffmann, A.A., Chown, S.L. & Clusella-trullas, S. (2013) Upper thermal limits in terrestrial

ectotherms: how constrained are they? *Functional Ecology*, **27**, 934–949.

Hoffmann, A.A. & Weeks, A.R. (2007) Climatic selection on genes and traits after a 100 year-

old invasion: A critical look at the temperate-tropical clines in *Drosophila melanogaster*

from eastern Australia. *Genetica*, **129**, 133–147.

Huey, R.B. & Pascual, M. (2009) Partial thermoregulatory compensation by a rapidly evolving

invasive species along a latitudinal cline. *Ecology*, **90**, 1715–1720.

Huey, R.B., Patridge, L. & Fowler, K. (1991) Thermal sensitivity of *Drosophila melanogaster*

responds rapidly to laboratory natural selection. *Evolution*, **45**, 751–756.

Jezkova, T. & Wiens, J.J. (2016) Rates of change in climatic niches in plant and animal

populations are much slower than projected climate change. *Proceedings of the Royal*

Society B: Biological Sciences, **283**, 20162104.

Karasov, T., Messer, P.W. & Petrov, D. a. (2010) Evidence that adaptation in *Drosophila* is not

limited by mutation at single sites. *PLoS genetics*, **6**, e1000924.

Katz, R.W. & Brown, B.G. (1992) Extreme events in a changing climate: Variability is more

important than averages. *Climatic Change*, **21**, 289–302.

Kellermann, V., Overgaard, J., Hoffmann, A. a, Fløjgaard, C., Svenning, J.-C. & Loeschcke, V.

(2012) Upper thermal limits of *Drosophila* are linked to species distributions and strongly

constrained phylogenetically. *Proceedings of the National Academy of Sciences of the*

United States of America, **109**, 16228–33.

Kelly, M.W., Sanford, E. & Grosberg, R.K. (2012) Limited potential for adaptation to climate

change in a broadly distributed marine crustacean. *Proceedings of the Royal Society B-*

Biological Sciences, **279**, 349–356.

Kingsolver, J.G., Diamond, S.E. & Buckley, L.B. (2013) Heat stress and the fitness

consequences of climate change for terrestrial ectotherms. *Functional Ecology*, **27**, 1415–

1423.

Kristensen, T.N., Kjeldal, H., Schou, M.F. & Nielsen, J.L. (2016) Proteomic data reveals a

physiological basis for costs and benefits associated with thermal acclimation. *Journal of*

Experimental Biology, in press.

Kristensen, T.N., Overgaard, J., Lassen, J., Hoffmann, A. a. & Sgrò, C. (2015) Low evolutionary

potential for egg-to-adult viability in *Drosophila melanogaster* at high temperatures.

Evolution, **69**, 803–814.

Leemans, R., Egger, B., Loop, T., Kammermeier, L., He, H., Hartmann, B., Certa, U., Hirth, F.

& Reichert, H. (2000) Quantitative transcript imaging in normal and heat-shocked

Drosophila embryos by using high-density oligonucleotide arrays. *Proc Natl Acad Sci U S*

A, **97**, 12138–12143.

Lenski, R.E. & Bennett, A.F. (1993) Evolutionary Response of *Escherichia coli* to Thermal

Stress. *The American Naturalist*, **142**, S47–S64.

Leuenberger, P., Ganscha, S., Kahraman, A., Cappelletti, V., Boersema, P.J., von Mering, C.,

Claassen, M. & Picotti, P. (2017) Cell-wide analysis of protein thermal unfolding reveals

determinants of thermostability. *Science*, **355**, eaai7825.

Llewelyn, J., Macdonald, S., Hatcher, A., Moritz, C. & Phillips, B.L. (2016) Thermoregulatory

behaviour explains countergradient variation in the upper thermal limit of a rainforest skink.

Oikos, **doi:10.111**, 1–10.

Lockwood, B.L., Sanders, J.G. & Somero, G.N. (2010) Transcriptomic responses to heat-stress

in invasive and native blue mussels (genus *Mytilus*): molecular correlates of invasive

success. *The Journal of Experimental Biology*, **213**, 3548–3558.

Lockwood, B.L. & Somero, G.N. (2012) Functional determinants of temperature adaptation in

enzymes of cold- versus warm-adapted mussels (Genus *Mytilus*). *Molecular Biology and*

Evolution, **29**, 3061–70.

Machado, H.E., Bergland, A.O., O'Brien, K.R., Behrman, E.L., Schmidt, P.S. & Petrov, D.A.

(2016) Comparative population genomics of latitudinal variation in *D. simulans* and *D.*

melanogaster. *Molecular Ecology*.

MacLean, H.J., Higgins, J.K., Buckley, L.B. & Kingsolver, J.G. (2016) Geographic divergence

in upper thermal limits across insect life stages: does behavior matter? *Oecologia*.

Meehl, G. a., Zwiers, F., Evans, J., Knutson, T., Mearns, L. & Whetton, P. (2000) Trends in

extreme weather and climate events: Issues related to modeling extremes in projections of

future climate change. *Bulletin of the American Meteorological Society*, **81**, 427–436.

Miller, N. a. & Stillman, J.H. (2012) Neural Thermal Performance in Porcelain Crabs, Genus

Petrolisthes. *Physiological and Biochemical Zoology*, **85**, 29–39.

Mitchell, K. a. & Hoffmann, A. a. (2010) Thermal ramping rate influences evolutionary potential

and species differences for upper thermal limits in *Drosophila*. *Functional Ecology*, **24**,

694–700.

Mongold, J.A., Bennett, A.F. & Lenski, R.E. (1999) Evolutionary adaptation to temperature. VII.

Extension of the upper thermal limit of *Escherichia coli*. *Evolution*, **53**, 386–394.

Morgan, T. & Mackay, T. (2006) Quantitative trait loci for thermotolerance phenotypes in

Drosophila melanogaster. *Heredity*, **96**, 232–242.

Munoz, M.M., Langham, G.M., Brandley, M.C., Rosauer, D.F., Williams, S.E. & Moritz, C.

(2016) Basking behavior predicts the evolution of heat tolerance in Australian rainforest

lizards. *Evolution*, **70**, 2537–2549.

Pereira, R.J., Barreto, F.S. & Burton, R.S. (2014) Ecological novelty by hybridization:

Experimental evidence for increased thermal tolerance by transgressive segregation in

Tigriopus californicus. *Evolution*, **68**, 204–215.

Rank, N.E. & Dahlhoff, E.P. (2002) Allele frequency shifts in response to climate change and

physiological consequences of allozyme variation in a montane insect. *Evolution*, **56**, 2278–

2289.

Rudolph, B., Gebendorfer, K.M., Buchner, J. & Winter, J. (2010) Evolution of *Escherichia coli*

for Growth at High Temperatures. *Journal of Biological Chemistry*, **285**, 19029–19034.

Sambucetti, P., Scannapieco, a C., Loeschcke, V. & Norry, F.M. (2013) Heat-stress survival in

the pre-adult stage of the life cycle in an intercontinental set of recombinant inbred lines of

Drosophila melanogaster. *The Journal of Experimental Biology*, **216**, 2953–9.

Sato, A., Kawashima, T., Fujie, M., Hughes, S., Satoh, N. & Shimeld, S.M. (2015) Molecular

basis of canalization in an ascidian species complex adapted to different thermal conditions.

Scientific Reports, **5**, 1–9.

Schou, M.F., Kristensen, T.N., Kellermann, V., Schl??tterer, C. & Loeschcke, V. (2014) A

Drosophila laboratory evolution experiment points to low evolutionary potential under

increased temperatures likely to be experienced in the future. *Journal of Evolutionary*

Biology, **27**.

Sgro, C.M., Overgaard, J., Kristensen, T.N., Mitchell, K.A., Cockerell, F.E. & Hoffmann, A.A.

(2010) A comprehensive assessment of geographic variation in heat tolerance and

hardening capacity in populations of *Drosophila melanogaster* from Eastern Australia.

Journal of Evolutionary Biology, **23**, 2484–2493.

Somero, G.N. (2002) Thermal Physiology and Vertical Zonation of Intertidal Animals: Optima,

Limits, and Costs of Living. *Integrative and Comparative Biology*, **42**, 780–789.

Stenseng, E., Braby, C.E., Somero, G.N. & Anonymous. (2005) Evolutionary and acclimation-

induced variation in the thermal limits of heart function in congeneric marine snails (Genus

Tegula): Implications for vertical zonation. *Biological Bulletin (Woods Hole)*, **208**, 138–144.

Stillman, J.H. & Somero, G.N. (2000) A comparative analysis of the upper thermal tolerance

limits of eastern Pacific porcelain crabs, genus *Petrolisthes*: influences of latitude, vertical

zonation, acclimation, and phylogeny. *Physiological and Biochemical Zoology*, **73**, 200–

208.

Sunday, J.M., Bates, A.E. & Dulvy, N.K. (2011) Global analysis of thermal tolerance and

latitude in ectotherms. *Proceedings. Biological sciences / The Royal Society*, **278**, 1823–30.

Sunday, J.M., Bates, A.E. & Dulvy, N.K. (2012) Thermal tolerance and the global redistribution

of animals. *Nature Climate Change*, **2**, 686–690.

Svetec, N., Cridland, J.M., Zhao, L. & Begun, D.J. (2016) The Adaptive Significance of Natural

Genetic Variation in the DNA Damage Response of *Drosophila melanogaster*. *PLOS*

Genetics, **12**, e1005869.

Tadros, W. & Lipshitz, H.D. (2009) The maternal-to-zygotic transition: a play in two acts.

Development (Cambridge, England), **136**, 3033–42.

Team, R.C. (2016) *R: A Language and Environment for Statistical Computing*. R Foundation for

Statistical Computing, Vienna, Austria.

Terblanche, J.S., Hoffmann, A.A., Mitchell, K.A., Rako, L., le Roux, P.C. & Chown, S.L. (2011)

Ecologically relevant measures of tolerance to potentially lethal temperatures. *Journal of*

Experimental Biology, **214**, 3713–3725.

Thomas, M.K., Kremer, C.T., Klausmeier, C. a & Litchman, E. (2012) A global pattern of

thermal adaptation in marine phytoplankton. *Science*, **338**, 1085–1088.

Tomanek, L. & Zuzow, M.J. (2010) The proteomic response of the mussel congeners *Mytilus*

galloprovincialis and *M. trossulus* to acute heat stress: implications for thermal tolerance

limits and metabolic costs of thermal stress. *The Journal of experimental biology*, **213**,

3559–3574.

Walter, M.F., Biessmann, H. & Petersen, N.S. (1990) Heat shock causes the collapse of the

intermediate filament cytoskeleton in *Drosophila* embryos. *Developmental genetics*, **11**,

270–279.

Welte, M.A., Tetrault, J.M., Dellavalle, R.I.P. & Lindquist, S.L. (1993) A new method for

manipulating transgenes: engineering heat tolerance in a complex, multicellular organism.

Current Biology, **3**, 842–853.

Yao, C.-L. & Somero, G.N. (2012) The impact of acute temperature stress on hemocytes of

invasive and native mussels (*Mytilus galloprovincialis* and *Mytilus californianus*): DNA

damage, membrane integrity, apoptosis and signaling pathways. *The Journal of*

Experimental Biology, **215**, 4267–77.

Figures

Figure 1.

Flies from different populations around the world exhibited differences in embryonic

thermal tolerance but not adult thermal tolerance.

(**A**) Sampling locations of isofemale lines. Color indicates maximum temperature of warmest

month (T_{max}), extrapolated from the WorldClim database. Tropical populations lie between the

Equator (solid horizontal line) and the Tropic of Cancer (dotted horizontal line) and were

sampled in Chiapas, Mexico (CH), Saint Kitts (SK), Accra, Ghana (GH), Mumbai, India (MU),

and Guam (GU). Temperate populations lie north of the Tropic of Cancer and were sampled in

Vermont (VT), Indiana (IN), and North Carolina (NC). (**B**) Proportion of adult male flies

responsive along a heat ramp (+1°C min^{-1} from 25°C). Tropical lines are indicated in solid pink.

Temperate lines are indicated in dotted blue. (**C**) Adult male CT_{max} was consistent across all

populations (ANOVA, $F_{3,9} = 2.378$, $P = 0.1375$). CT_{max} extrapolated from the survival curves in

B. Solid horizontal lines show means. Error bars indicate 95% confidence intervals. (**D**)

Proportion of eggs successfully hatched following heat shock (45 min at indicated temperature).

Tropical lines are indicated in solid pink. Temperate lines are indicated in dotted blue. (**E**)

Embryonic thermal tolerance (LT_{50}) was higher in tropical lines than temperate lines (ANOVA,

$F_{3,20} = 10.16$, $P = 0.0003$; Tukey's test, VT vs. IN, $q = 2.428$, $P = 0.3416$, VT vs. NC, $q = 0.4268$,

$P = 0.9902$, IN vs. NC, $q = 2.666$, $P = 0.2656$, tropical vs. VT, $q = 6.909$, $P = 0.0005$, tropical vs.

IN, $q = 4.04$, $P = 0.0444$, tropical vs. NC, $q = 4.04$, $P = 0.0005$). LT_{50} extrapolated from the

survival curves in D. Solid horizontal lines show means. Error bars indicate 95% confidence

intervals. *$P < 0.05$.

Figure 2.

Thermal safety margin differs by life stage and geographic region.

(**A**) Thermal safety margins were smaller for embryos than adults across latitude (Extra sum-of-

squares F-test on fitness of two curves, $F_{3,32} = 15.95$, $P < 0.0001$; Non-linear least-squares

regression, embryo, $R^2 = 0.8019$, $y = 0.01563x^2 - 0.7249x + 9.97$, adult, $R^2 = 0.6359$, $y =$

$0.01437x^2 - 0.6424x + 12.44$). Adults are indicated in purple filled circles with dashed purple

regression line. Embryos are indicated in orange open circles with solid orange regression line.

(**B**) Thermal safety margins were smaller for embryos than adults and smaller in the tropics than

temperate sites (ANOVA, main effect of region, $F_{1,34} = 6.482$, $P = 0.0156$, main effect of life

stage, $F_{1,34} = 11.15$, $P = 0.002$, region x life stage interaction, $F_{1,34} = 0.0006$, $P = 0.98$). Adults

are indicated in purple filled circles, and embryos are indicated in orange open circles. Solid

horizontal lines show means. Error bars indicate 95% confidence intervals. *$P < 0.05$, **$P <$

0.01.

Figure 3.

Embryonic thermal tolerance and maximum habitat temperature (T_{max}) by region.

Embryonic thermal tolerance was positively correlated with T_{max} among tropical populations

(Least-squares regression, $R^2 = 0.9478$, $P = 0.0051$, $y = 0.2199x + 28.75$) but not temperate

populations (Least-squares regression, $R^2 = 0.0015$, $P = 0.9751$, $y = 0.00282x + 34.82$). Tropical

genotypes are indicated in pink circles, with a solid black regression line fit. Temperate

genotypes are indicated in blue triangles, with a dashed black regression line fit.

Figure 4.

F1 progeny from tropical x temperate parents have high embryonic heat tolerance.

Proportion of eggs successfully hatched following heat shock (45 min at indicated temperature) among two parental genotypes that had the highest and lowest LT_{50} of all strains in this study, CH (Chiapas, Mexico) and VT-12 (Vermont, USA), respectively, along with F1 progeny from reciprocal crosses of these two parental lines, CH♀ x VT♂ and VT♀ x CH♂ (♀ = dam; ♂ = sire). Note that VT-12 is labeled "VT" in the legend. LT_{50}: CH = 36.24°C, VT-12 = 34.23°C, CH♀ x VT♂ = 35.83°C, VT♀ x CH♂ = 35.80°C (Logistic model, Extra sum-of-squares F-test on lower LT_{50} of VT-12, $F_{3,166}$ = 6.695, ***P = 0.0003).

645   **Supplemental Figure S1.**

646   **No significant relationship between adult CT_{max} and embryo LT_{50}.**

647   Adult thermal tolerance and embryonic thermal tolerance were not correlated among all

648   isofemale lines (Least-squares linear regression, $R^2 = 0.0005$). Tropical isofemale lines are

649   shown in pink circles and temperate isofemale lines are shown in blue triangles.

Supplemental Figure S2.

Principal components analysis of 19 Bioclim variables at each of the 8 collection sites.

Plotted are scores of the first two principle components (PC1 vs. PC2), which describe the major

axes of variation in 19 bioclimatic variables among the collection sites. PC1 and PC2 explained

95.6% and 4%, respectively, of the variation in climate among sites. Temperature seasonality

was the top component loading on PC1, explaining 98% of the variation in PC1, and annual

precipitation was the top component loading on PC2, explaining 49% of the variation in PC2.

Temperate sites are shown in blue and tropical sites in pink.

A

B

Supplemental Figure S3.

Relationship between embryonic thermal tolerance and (A) temperature seasonality and

(B) mean annual temperature.

(**A**) Temperature seasonality is the standard deviation in temperature (°C) over the course of the

year, averaged across the years 1960 to 1990 for each of the sites. Solid black line represents the

regression line fit, which had a significantly negative slope (Least-squares linear regression, R^2 =

0.42, $P = 0.0006$, $y = -0.01x + 35.82$). (**B**) Mean annual temperature is the average yearly

temperature from 1960 to 1990 for each of the sites. Solid black line represents the regression

line fit, which had a significantly positive slope (Least-squares linear regression, $R^2 = 0.37$, $P =$

0.0016, $y = 0.041x + 34.52$). In both A and B, tropical isofemale lines are shown in pink circles

and temperate isofemale lines are shown in blue triangles.

Supplemental Figure S4.

No significant relationship between adult thermal tolerance (CT_{max}) and maximum habitat

temperature (T_{max}).

Variation in adult thermal tolerance showed no correspondence to variation in T_{max} among all

collection sites (Least-squares linear regression, $R^2 = 0.055$). Tropical isofemale lines are shown

in pink circles and temperate isofemale lines are shown in blue triangles.

682   **Tables**

683   **Supplemental Table S1.**

684   Stock information, collection locales, and thermal tolerance data for isofemale lines used in this

685   study.

Stock No.	Locale	State/Country	Lat. (°N)	Long. (°E)	Adult CT$_{max}$ (°C)	Embryo LT$_{50}$ (°C)
VTECK_2	East Calais	Vermont, USA	44.3664	-72.4300	39.69	35.18
VTECK_4	East Calais	Vermont, USA	44.3664	-72.4300	-	35.03
VTECK_5	East Calais	Vermont, USA	44.3664	-72.4300	38.27	34.75
VTECK_8	East Calais	Vermont, USA	44.3664	-72.4300	-	34.81
VTECK_9	East Calais	Vermont, USA	44.3664	-72.4300	-	35.27
VTECK_10	East Calais	Vermont, USA	44.3664	-72.4300	40.52	34.17
VTECK_12	East Calais	Vermont, USA	44.3664	-72.4300	-	34.23
VTECK_14	East Calais	Vermont, USA	44.3664	-72.4300	-	35.01
BEA_5	Beasley Orchard	Indiana, USA	39.7637	-86.4766	-	35.13
BEA_16	Beasley Orchard	Indiana, USA	39.7637	-86.4766	37.62	35.11
BEA_21	Beasley Orchard	Indiana, USA	39.7637	-86.4766	38.33	34.94
BEA_32	Beasley Orchard	Indiana, USA	39.7637	-86.4766	-	35.30
BEA_36	Beasley Orchard	Indiana, USA	39.7637	-86.4766	38.07	35.29
RFM_4	Raleigh	North Carolina, USA	35.7636	-78.6627	-	34.98
RFM_6	Raleigh	North Carolina, USA	35.7636	-78.6627	-	34.95
RFM_16	Raleigh	North Carolina, USA	35.7636	-78.6627	-	34.03
RFM_19	Raleigh	North Carolina, USA	35.7636	-78.6627	39.60	34.63
RFM_34	Raleigh	North Carolina, USA	35.7636	-78.6627	37.61	34.96
RFM_48	Raleigh	North Carolina, USA	35.7636	-78.6627	37.61	34.94
14021-0231.22	Chiapa de Corzo	Chiapas, Mexico	16.7022	-93.0081	39.13	36.24
14021-0231.34	Monkey Hill	St. Kitts	17.3240	-62.7250	39.05	35.44
14021-0231.182	Accra	Ghana	5.5557	-0.1963	38.74	35.65
14021-0231.45	Mumbai	Maharashtra, India	19.0760	72.8777	39.50	36.14
14021-0231.198	Guam	Guam, USA	13.4443	144.7937	39.01	35.51