1 ***Brucella abortus* induces a Warburg shift in host metabolism that enhances**

2 **intracellular replication of the pathogen**

Daniel M. Czyż[a,b], Jonathan Willett[a,b], Sean Crosson[a,b,c#]

Howard Taylor Ricketts Laboratory, University of Chicago, Argonne National
Laboratory, Lemont, Illinois, USA[a]; Department of Biochemistry and Molecular Biology,
University of Chicago, Chicago, Illinois, USA[b]; Department of Microbiology, University of
Chicago, Chicago, Illinois, USA[c]

Running title: *Brucella abortus* infection and host cell metabolism

[#]Address correspondence to Sean Crosson, scrosson@uchicago.edu

KEYWORDS: Metabolism, Warburg effect, lactate, Brucella, therapeutics

ABSTRACT

Intracellular bacterial pathogens exploit host cell resources to replicate and survive inside the host. Targeting these host systems is one promising approach to developing novel antimicrobials for intracellular infections. We show that human macrophage-like cells infected with *Brucella abortus* undergo a metabolic shift characterized by attenuated tricarboxylic acid cycle metabolism, altered mitochondrial morphology and increased lactate production. This shift to an aerobic glycolytic state resembles the Warburg effect, a change in energy production that is well-described in cancer cells, and also occurs in activated inflammatory cells. *B. abortus* efficiently uses lactic acid as a carbon and energy source and requires the ability to metabolize lactate for normal replication in human macrophage-like cells. We further demonstrate that chemical inhibitors of host glycolysis and lactate production do not affect *in vitro* growth of *B. abortus* in axenic culture, but decrease its replication in the intracellular niche. Our data support a model in which infection shifts host metabolism to a Warburg-like state, and *B. abortus* uses this change in metabolism to promote intracellular replication. Pharmacological perturbation of these features of host cell metabolism may be a useful strategy to inhibit infection by intracellular pathogens.

IMPORTANCE

Brucella spp. are intracellular pathogens that cause disease in a range of mammals, including livestock. Transmission from livestock to humans is common and can lead to chronic human disease. Human macrophage-like cells infected with *Brucella abortus* undergo a Warburg-like metabolic shift to an aerobic glycolytic state that produces lactic acid, which *B. abortus* can use as a carbon and energy source to promote its growth.

We provide evidence that the pathogen exploits this change in host metabolism to

replicate in the intracellular niche. Drugs that target this feature of host cell metabolism

inhibit intracellular replication of *B. abortus* and may be broadly useful therapeutics for

intracellular infections.

INTRODUCTION

Macrophages can be activated by pathogen-associated pro-inflammatory

molecules such as lipopolysaccharide (LPS). Classically activated (i.e. M1)

macrophages undergo a major metabolic shift in which the tricarboxylic acid (TCA)

cycle is downregulated, and energy production transitions from oxidative

phosphorylation to the less efficient process of aerobic glycolysis (1, 2). This change in

metabolism, known as the Warburg effect (3), is a well-described metabolic feature of

cancer cells (4). Increased glucose consumption through glycolysis is thought to

increase adenosine triphosphate (ATP) levels; a corresponding increase in pentose

phosphate pathway activity generates nicotinamide adenine dinucleotide phosphate

(NADPH), which is subsequently used by NADPH oxidases to generate reactive oxygen

species (ROS) (5). This metabolic shift is important in enabling cells to launch general

antimicrobial defense mechanisms by increasing concentrations of ROS and reactive

nitrogen intermediates (6).

The intracellular pathogen *Brucella abortus* invades host cells and is trafficked to

a compartment known as the Brucella containing vacuole, where it replicates. To

successfully find its way to this niche, *B. abortus* must avoid numerous chemical insults

by the host cell and adapt to the nutrient resources available within the host. The

mechanisms by which *B. abortus* evades host defenses and takes advantage of host

metabolic responses remain understudied. *B. abortus* induces a Warburg-like effect in

mammalian host cells. We demonstrate that this metabolic shift is advantageous for *B.*

abortus replication, and that chemical inhibition of Warburg metabolism in macrophage-

like cells attenuates intracellular replication of *B. abortus*. Thus, *B. abortus* exploits an

innate immune metabolic response to support infection.

RESULTS

***Brucella abortus* infection of human macrophage-like cells disrupts mitochondrial**

function and morphology

To characterize the effect of intracellular colonization by *B. abortus* on human

cell metabolism, we employed mammalian cell phenotype microarrays (PM) to measure

respiration in terminally differentiated, metabolically active THP-1 cells. These cells

were cultivated in minimal essential medium in the presence of central metabolic

substrates specifically selected to measure mitochondrial function. Three of the

substrates, α-ketoglutaric acid, β-hydroxybutyric acid, and pyruvic acid are metabolized

in mitochondria. We assessed the respiration of the human cells solely utilizing the

provided substrate as a primary energy source by colorimetrically measuring reduction

of a redox indicator dye. This assay allows us to assess the cells' ability to metabolize

substrates across different molecular pathways, and thus provides a measurement of

metabolic pathway activity upon infection (Figure 1A). We infected the cells with *B.*

abortus at increasing multiplicities of infection (MOI). To eliminate the possibility that

changes in the metabolic activity of human cells are caused by *Brucella*-induced cell death, we removed medium containing mitochondrial substrates and replaced it with a regular growth medium containing glucose, followed by assessment of cell viability using the MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay. The number of viable cells remained the same at different *Brucella* MOI (Figure 1B). Naïve cells utilized all of the mitochondrial substrates (Figure 1C), but their ability to utilize those substrates decreased in cells infected with *B. abortus* (Figure 1D). This decrease was dependent on MOI (Figures 1A). *Brucella*-infected cells, unlike with all other substrates, continued to utilize glucose (Figure 1C–D). In fact, infected cells increased glucose consumption with increasing MOI (Fig. S1).

The marked inhibition of mitochondrial substrate utilization suggests that intracellular Brucellae diminish mitochondrial function. Parasites (7), viruses (8), and intracellular bacterial pathogens including *B. melitensis* (9) and *Listeria monocytogenes* (10) perturb mitochondrial function in host cells. To obtain additional evidence that mitochondrial function is disrupted during *B. abortus* infection, we examined mitochondrial morphology in uninfected and infected HeLa cells (because THP-1 cells are not amenable to mitochondrial staining or imaging). Disruptions of mitochondrial structure were observed in HeLa cells with detectable *B. abortus*, whereas uninfected cells in the same field of view had normal cellular and mitochondrial morphology (Figure 2). Using an MTT cell viability assay we detected no loss in host viability with increasing MOI within the same timeframe (Figure 1B). Therefore, it is unlikely that the mitochondrial disruption observed by fluorescence microscopy results from host cell death.

Most amino acids can be metabolized to enter the TCA cycle through anaplerotic

reactions (11). To determine whether the observed *B. abortus*-mediated disruption of

host mitochondrial function directly affects amino acid metabolism, we assessed the

effect of infection on the host's ability to utilize amino acids as a carbon source. While

glucose metabolism was unaffected, infected cells could not metabolize any of the

amino acids known to enter the TCA cycle (Figure 3). Together, the results of these

experiments are consistent with a model in which infection of human macrophages with

B. abortus results in major metabolic changes that are linked to disruption of

mitochondrial function.

**Brucella*-dependent induction of lactic acid production in human cells*

Macrophages and dendritic cells are central to innate immunity. Upon activation, they

can undergo a shift in energy metabolism by shutting down oxidative phosphorylation

and increasing the rate of aerobic glycolysis in a process known as the Warburg effect

(3). A hallmark of this effect is production of lactic acid in the presence of oxygen. An

increase in lactate production in non-polarized macrophages infected with *Brucella* spp.

has been previously noted (12). To further validate that THP-1 infected with *B. abortus*-

are shifted toward a Warburg-like state, we measured the production of lactic acid in

host cells as a function of increasing MOI. THP-1 cells increase lactic acid production in

an MOI-dependent manner (Figure 4A). These results, together with the data presented

in Figures 1-3 support a model in which *Brucella* promotes a Warburg-like shift in host

cell metabolism by disrupting mitochondrial function.

129 *Brucella abortus* **lactate dehydrogenase is required for proper intracellular**

130 **replication**

131 *Brucella* and *Salmonella* both replicate intracellularly and can persist in the host for long

132 periods by shifting host metabolism to make glucose more available (12-14). We

133 measured the growth of *B. abortus* on two preferred substrates, glucose and erythritol,

134 and compared these data to growth on lactic acid. *B. abortus* grows on lactic acid as a

135 sole carbon and energy source as well as it does on glucose or erythritol in standing

136 culture *in vitro* (Figure 4B). We postulated that, during infection, *B. abortus* may exploit

137 the Warburg-type shift in host metabolism by utilizing lactic acid as a carbon and energy

138 source.

139 To define a genetic determinant of *B. abortus* lactic acid metabolism, we

140 engineered a strain harboring an in-frame deletion of the gene encoding L-lactate

141 dehydrogenase (LDH), *Bab2_0315* (Δ*lldD*). To determine whether *lldD* deletion affects

142 utilization of other carbon sources, we used PM to compare the metabolic activity of

143 Δ*lldD* to wild type (WT), and found that utilization of L-lactic acid was the most

144 statistically significant loss-of-function phenotype (Figure 5A). WT *B. abortus* exhibited

145 robust growth on lactic acid as a carbon and energy source, but Δ*lldD* did not grow in

146 this condition (Figure 5B). The Δ*lldD* mutant had no apparent growth defect in *Brucella*

147 broth supplemented with glucose (Figure 5C), but the numbers of colony forming units

148 (CFU) of Δ*lldD* were significantly reduced in THP-1 macrophages relative to WT at the

149 48-hour time point (Figure 5D). Complementation of the mutation by introduction of *lldD*

150 at its native locus rescued the THP-1 growth/replication defect. Together, our results

151 indicate that, upon infection, human cells increase lactic acid production and provide an

152 advantage to intracellular *B. abortus*, which can utilize host lactic acid as a carbon and

153 energy source.

155 Small-molecule glycolysis inhibitors inhibit intracellular replication of *Brucella*

156 *abortus* but do not inhibit axenic growth

Identification of pathogen-induced changes in host metabolism offers new avenues for

the development of anti-infectives. Since *Brucella*-infected cells exhibited attenuated

mitochondrial function and increased glycolysis, we used small-molecule inhibitors of

glucose metabolism (Figure 6A) to determine whether inhibition of the host glycolytic

pathway affected intracellular growth/replication of *B. abortus*. We tested: 3-

bromopuruvic acid (3-BPA), an inhibitor of energy metabolism that completely

eradicated cancer in animal models with no apparent cytotoxicity (15, 16); methyl 1-

hydroxy-6-phenyl-4-(trifluoromethyl)-1H-indole-2-carboxylate (NHI-2), a potent

anticancer inhibitor of lactate dehydrogenase A (LDH-A) (17, 18); and 2-deoxy-D-

glucose (2-dG), a non-metabolizable homolog of D-glucose that slows glycolysis (19).

We counted *B. abortus* cells located inside THP-1 cells treated with each of

these anti-metabolic compounds at various concentrations. Since some of these

compounds may be toxic to cells at the tested concentrations, results were expressed

as the number of *Brucella* per THP-1 nucleus; hence, the data represent bacterial

counts in live cells only. Each anti-metabolic drug significantly inhibited

growth/replication of intracellular *Brucella* in live cells (Figure 6B).

To determine whether the antimicrobial activity of these compounds coincides

with their cytotoxicity to human cells, we assessed their cytotoxicity using an XTT cell

viability assay. As expected, each metabolic inhibitor was cytotoxic at higher

concentrations. Treatment of THP-1 cells with 3-BPA, the most potent compound,

inhibited *Brucella* infection at non-cytotoxic concentrations. NHI-2 and 2-dG were less

potent, but effectively inhibited growth/replication of intracellular *Brucella* (Figure 6C).

Both 3-BPA and NHI-2 significantly inhibited intracellular *Brucella* over a range of

concentrations that did not affect host cell viability.

Finally, to determine whether the antimicrobial effect of these compounds occurs

by host targeting, or whether they also directly inhibit bacterial growth, we measured

growth and viability of *B. abortus* in axenic culture in the presence of 3-BPA, NHI-2, and

2-dG. Neither 3-BPA nor NHI-2 affected growth and viability of *Brucella* in axenic

medium, though 2-dG exhibited an inhibitory effect at higher concentrations ($\geq$ 50 mM).

These data support a model in which inhibition of host glycolysis and lactate

production with 3-BPA and NHI-2 inhibits intracellular *B. abortus* replication without

directly targeting *B. abortus* growth (Figure 6D). Together, these results demonstrate

that infection-induced changes in host central metabolic pathways are required for

proper growth/replication of intracellular *B. abortus*. Inhibition of the glycolytic pathway

with host-targeting anti-metabolic compounds provides an effective means to decrease

intracellular bacterial load without directly targeting the bacterium (Figure 6E).

DISCUSSION

We have identified metabolic changes in host cells upon exposure to an intracellular

bacterial pathogen, *B. abortus*. Intracellular *B. abortus* perturb the morphology and

function of host mitochondria, as characterized by loss of the ability to utilize

mitochondrial substrates (Figure 1 and 2). The host metabolic effects we observed upon

B. abortus infection resemble the effect of adding mitochondrial inhibitors (20), and are

consistent with the previously reported downregulation of mitochondrion-associated

gene expression in macrophages infected with *B. melitensis* (21). Many bacteria are

known to target host mitochondria (22), but only a few species have been reported to

disrupt mitochondrial morphology (9, 10, 23). The structures of mitochondria observed

in our study resemble those of cells treated with uncouplers in combination with a

respiration inhibitor; though this treatment resulted in energy deprivation and

aggregation of mitochondria, the cells remained viable (24).

Intracellular pathogens compete with the host cell for available nutrients,

including central metabolites and amino acids (25). As demonstrated using Biolog

mammalian PM, THP-1 cells utilized amino acids as an energy source, but this is

inhibited by *B. abortus* infection (Figure 3). The inability of infected cells to utilize amino

acids suggests that either intracellular Brucellae compete for the pool of available amino

acids, or the cells cease these catabolic reactions because of a change in mitochondrial

(or other) physiology. Inhibiting the TCA cycle and increasing glucose metabolism can

allow cells to rapidly generate the biomass required for cancer growth. Analogously,

induction of a Warburg-like state upon *B. abortus* infection may enable synthesis of

building blocks required for intracellular bacterial growth.

Intracellular pathogens rely on host resources for survival and replication. It is

therefore not surprising that other species of bacteria also affect host metabolism upon

infection. For example, *Mycobacterium tuberculosis* enhances glycolysis in THP-1 cells

(26). Early studies of the infection-induced changes in host metabolism showed that

cells infected with *Chlamydia* increase their utilization of glucose through glycolysis and

increase lactate production (27). Other pathogens, including the parasite *Plasmodium*

falciparum (28), and Mayaro virus (29), also induce similar changes in host metabolism.

Because the metabolic changes observed in these hosts are induced upon exposure to

various stimuli, it is unlikely that the changes observed in our study are specifically

induced by Brucellae.

Erythritol and glucose are preferred energy sources for *Brucella* (30), and our

results show that, in axenic culture, *Brucella* can utilize lactate as efficiently as erythritol

or glucose (Figure 4). Hence, we hypothesized that intracellular Brucellae can use

abundant host lactate as an energy source, which would enable successful intracellular

colonization and replication. We took two approaches to test whether host lactate

affects intracellular growth. First, we created a lactate dehydrogenase knockout strain

Δ*lldD*, which is unable to utilize lactate in axenic culture (Figure 5A-B). While Δ*lldD*

showed no growth defects across a range of axenic media, its intracellular

replication/survival was attenuated by ~1 log after 48 h in host THP-1 cells (Figure 5D).

This suggests that the ability to utilize lactate is necessary for proper replication in the

intracellular niche. We note there are parallels between *B. abortus* pathogenicity and

that of intracellular *Neisseria* spp., which is affected by host lactic acid (31). In the

second approach, we targeted host lactate synthesis using the small molecule inhibitors

NHI-2, 3-BPA, and 2-dG, which decrease lactate levels and inhibit aerobic glycolysis

(17, 18, 32, 33). Each of these metabolic inhibitors decreased intracellular growth

and/or replication of *Brucella* (Figure 6). Each of these compounds is known for its

anticancer properties, but their anti-infective properties were not previously reported.

Collectively, the data presented in this study support a model in which intracellular *B.*

abortus exploits host metabolic pathways (Figure 7). Hence, the host anti-metabolic

compounds identified here and in previous studies (34) may provide a novel therapeutic

approach to treat *B. abortus* and other bacterial infections that rely on host metabolic

intermediates.

MATERIALS AND METHODS

Bacterial strains

All studies on live *B. abortus* strain 2308 were conducted at Biosafety Level 3 (BSL3) at

the Howard Taylor Ricketts Regional Biocontainment Laboratory, University of Chicago,

according to US Federal Select Agent Program guidelines. *B. abortus* expressing

mCherry was previously generated from the WT *B. abortus* 2308 parent strain by

integrating mini-Tn7 expressing a single copy of *mCherry* at the *glmS* locus. Bacterial

strains used in this study are listed in Table S1.

Mammalian tissue culture

Prior to each experiment, THP-1 macrophage-like cells were grown to a maximum density of 1 x 10^6 mL^{-1} in complete RPMI-1640 medium (HyClone) supplemented with 2 mM glutamine (Gibco) and 10% heat-inactivated fetal bovine serum (FBS) (HyClone). HeLa cells were grown in Dulbecco Modified Eagles Medium (DMEM; HyClone) supplemented with 10% heat-inactivated FBS. All tissue culture cells were grown at 37°C in a humidified environment with 5% CO_2.

Mammalian Phenotype MicroArray: mitochondria toxicity

THP-1 cells were seeded at 50,000 cells/well in two half-area 96-well plates in the presence of 40 ng mL^{-1} phorbol myristate acetate (PMA), and allowed to differentiate for 72 h at 37°C. *B. abortus* 2308 strain was streaked onto Schaedler blood agar (SBA) plates before growing for 2 d, re-streaking, and growing for another 2 d. At the day of infection, *Brucella* cells were scraped off the SBA plate, and suspended and washed in RPMI-1640 growth medium. Cell concentration was measured by the optical density method at 600 nm (OD_{600}). Bacteria were added to the last column of a 96-well plate at 1000 MOI, and serially diluted by a factor of two. The first column was left uninfected. Plates were centrifuged at 2170 rpm for 10 min, followed by a 1-h incubation at 37°C and 5% CO_2. Gentamicin was added to each well at a final concentration of 100 µg mL^{-1} and the plates were further incubated for 1 h. Following incubation, the

medium was removed and replaced with fresh medium containing 25 µg mL^{-1}

gentamicin. Cells were incubated for 24 h. Twenty-four hours after infection, two PM-M

TOX1 MicroPlates (Biolog) were reconstituted with 50 µL (per well) of MC-0 medium

(Biolog IF-M1 medium, 0.3 mM L-Gln, 5% dialyzed FBS, 25 µg mL^{-1} gentamicin). The

culture medium was removed from infected cells and replaced with the contents of

reconstituted PM-M TOX1 plates. *Brucella*-infected cells were incubated in the presence

of mitochondrial substrates for another 24 h. Following this incubation, 10 µL of PM-A

dye (Biolog) was added to each well, and the absorbance measured every 15 min (up to

150 min) at 590 nm and 750 nm using a Tecan Infinite 200 PRO microplate reader.

Plates were incubated at 37°C and 5% CO_2 between readings. The collected

absorbance 590 nm data (minus reference absorbance at 750 nm) were analyzed as a

function of time.

Following the experiment, cell viability was assessed by removing all liquid from

the PM-M TOX1 plate, replacing it with glucose-supplemented growth medium

containing 10% FBS, and 0.5 mg mL^{-1} MTT reagent, then measuring the absorbance at

570 nm after 24 h.

Mammalian Phenotype MicroArray: utilization of amino acids

THP-1 and *B. abortus* 2308 cells were prepared as described above. The infection

protocol was as described above with small changes. Bacteria were added to

differentiated THP-1 cells in a 96-well plate at 100 MOI. Plates were centrifuged at

2170 rpm for 10 min followed by 1-h incubation at 37°C and 5% CO_2. Gentamicin was

added to each well at a final concentration of 100 µg mL^{-1} and plates were further

incubated for 1 h. Following incubation, the medium was removed and replaced with a

fresh medium containing 25 µg mL^{-1} gentamicin. Cells were incubated for 24 h. Twenty-

four hours after infection, the content of the mammalian phenotype microarray PM-M2

(Biolog) was reconstituted with 50 µL (per well) of MC-0 medium (Biolog IF-M1 medium,

0.3 mM L-Gln, 5% dialyzed FBS, 25 µg mL^{-1} gentamicin). The culture medium was

removed from infected cells and replaced with the contents of reconstituted PM-M2

plates. *Brucella*-infected cells were incubated in the presence of various amino acids for

another 24 h. Following this incubation, 10 µL of dye PM-A (Biolog) was added to each

well, and the absorbance was measured every 15 min (up to 120 min) at 590 nm and

750 nm using a Tecan Infinite 200 PRO microplate reader. Plates were incubated at

37°C and 5% CO_2 between readings. Absorbance at 590 nm data (minus reference

absorbance at 750 nm) was analyzed as a function of time. Results were analyzed by

hierarchical clustering using Gene Cluster 3.0 and visualized using Java TreeView

(v1.1.6r4).

Bacterial Phenotype MicroArray

Utilization of glucose, erythritol, and lactic acid was measured using PM as previously

described (35). Briefly, *B. abortus* strains (WT 2308 and Δ*bab2_0315*) were streaked

onto an SBA plate, cultivated at 37°C and 5% CO_2 for 48 h, re-streaked, and grown for

another 48 h. Cells were scraped off the plate and resuspended in 1 x IF-0a medium

(Biolog) at a final density equivalent to 5% transmittance (OD$_{600nm}$). PM inoculating

fluids (PM1 and PM2) were prepared from 0.2-µm-filtered and sterilized stock solutions

at the following final concentrations: 2 mM $MgCl_2$, 1 mM $CaCl_2$, 25 µM L-arginine,

50 µM L-glutamic acid, 5 µM β-NAD, 25 µM hypoxanthine, 25 µM 5′-UMP, 25 µM L-

cystine (pH 8.5), 0.005% yeast extract, and 0.005% Tween 40. PM1 and PM2

inoculating fluids were prepared in 1 x IF-0a GN/GP medium (Biolog). Dye mix G

(Biolog) was added to each solution to a final concentration of 1X. A 1:13.6 dilution of

5% transmittance *Brucella* suspension was added to each inoculating fluid. Per well,

100 µL of *Brucella*-containing inoculating fluid was dispensed into each PM plate. PM

plates were incubated at 37°C and 5% CO_2, and the absorbance at 630 nm was

measured every 24 h for up to 5 d, using a Tecan Infinite 200 PRO microplate reader.

Imaging of mitochondria

HeLa cells were seeded onto 22 x 22 mm cover slides at 50,000 cells/well in a 6-well

plate in 2 mL of DMEM medium per well. Cells were grown for 24 h prior to infection.

Prior to infection, a culture of *B. abortus* constitutively expressing mCherry was

inoculated in *Brucella* broth, and grown overnight at 37°C on orbital shaker. On the day

of infection, *Brucella* cells were collected, resuspended in DMEM, and opsonized by

incubating for 30 min at 37°C with 1:1000 dilution of serum obtained from *Brucella*-

infected mice, as previously described (36). Following opsonization, cells were washed

and added to HeLa cells at 10,000 MOI. The plate was centrifuged at 2170 rpm for

20 min, and placed for 24 h at 37°C and 5% CO_2. Following incubation, cells were

washed with DMEM, re-infected at 10,000 MOI, and incubated for another 24 h. After

this incubation, cells were washed with DMEM, the cover slide was transferred to a

50 mL tube containing 15 mL DMEM and 200 µg mL^{-1} gentamicin. The slide was

incubated at 37°C and 5% CO_2 for 1 h. Following the incubation, 15 µL of 10 mg mL^{-1}

Hoechst stain and 2 µL of 1 mM MitoTracker (Molecular Probes) dye were added and

incubated for another 20 min. The cover slide was washed and placed on a microscope

slide with cells facing up. A drop of DMEM was placed on top of the cover slide and

covered by another 22 x 22 mm cover slide. To prevent medium evaporation, the edges

were quickly sealed with nail polish, and live cells were immediately imaged using a

Leica-DMI6000B fluorescence microscope equipped with a HC PL APO 63X/1.4 NA Oil

PH3 CS2 objective. Images were captured using Leica Application Suite X and

Hamamatsu Orca-R2 camera.

Quantification of lactic acid

Intracellular lactic acid levels were quantified according to the manufacturer's protocol

using a lactate assay kit (Sigma-Aldrich). Briefly, THP-1 cells were seeded and infected

with *B. abortus* at increasing MOI, as described above. Infected cells were incubated for

48 h before quantifying lactate levels. After incubation, cells were washed with PBS,

homogenized in 60 µL of lactate assay buffer, and cell debris was removed by

centrifuging at 14,000 *g* for 10 min. Fifty microliters of supernatant was collected from

each sample, and endogenous lactate dehydrogenase was removed by centrifuging

supernatants at 14,000 *g* for 30 min using 10 kDa MWCO Ultra-0.5 mL spin filter

(Amicon). Twenty microliters of lactate assay buffer was added to each sample to bring

the volume up to 50 µL. Fifty microliters of master mix (lactate assay buffer, enzyme

mix, and probe) was added to each well, mixed, and incubated in the dark at room

temperature for 30 min before measuring absorbance at 570 nm. Lactate concentration

was calculated based on a standard lactate curve and normalized to the number of cells

per well.

371 Construction of *B. abortus lldD* chromosomal deletion

A chromosomal deletion of *B. abortus lldD* (*bab2_0315*) was obtained using a double

recombination strategy. Primers used in this study are given in Table S2. The 500 base

pairs upstream (using primers 1775 and 1779) and downstream (using primers 1776

and 1777) of *lldD* were Gibson-cloned into pNPTS138, which was digested with *Bam*HI

and *Hin*dIII (New England Biolabs). Constructs were validated by DNA sequencing. To

delete *lldD*, the deletion plasmid was transformed into *B. abortus*, and primary

integrants were selected by plating on SBA supplemented with 50 μg mL^{-1} kanamycin

(SBA-Kan). The resulting colonies were grown overnight in *Brucella* broth without

selection and plated on SBA supplemented with 5% sucrose to select for second

crossovers. The resulting colonies were screened for the *lldD* deletion by sequencing

using the sequencing primers 1788/1789.

To complement the *lldD* deletion strain, the *lldD* gene plus the 500 bp upstream

and downstream (using primers 1775/1777) were Gibson-cloned into pNPTS138, and

restored at the native site using a two-step recombination strategy as outlined above.

Brucella axenic growth

The concentration of *Brucella* cells (WT 2308 and Δ*bab2_0315*) was adjusted to 20% transmittance at OD_{600} and diluted into glucose-containing *Brucella* broth (Difco) at 1:20. MTT (3-(4.5-dimethylthiazol-2-yl)-2.5-diphenyltetrazolium bromide) reagent was added to the media to a final concentration of $0.5 \, mg \, mL^{-1}$, and cells were continuously incubated at 37°C in a Tecan plate reader, taking absorbance readings at 570 nm every 10 min for 72 h.

Brucella axenic growth: compound cytotoxicity

NHI-2 (Sigma-Aldrich), 3-BPA, and 2-dG (Santa Cruz Biotechnology) were freshly prepared and serially diluted to test concentrations in a half-area 96-well plate in a total volume of 50 µL of PM9 medium (2 mM $MgCl_2 \bullet 6H_2O$, 1 mM $CaCl_2 \bullet 2H_2O$, 0.005% yeast extract, 0.005% Tween 40, 2.5 mM D-glucose, 5 mM sodium pyruvate, and 1X Dye Mix G [Biolog]). WT 2308 *Brucella* was prepared from a plate culture in PM9 medium at 20% transmission at 600 nm optical density, and 50 µL was added to each well for a total volume of 100 µL per well. Cells were incubated at 37°C, and absorbance at 630 nm was read every 2 h for 72 h using a Tecan microplate reader.

XTT cell viability assay

The cytotoxicity of NHI-2, 3-BPA, and 2-dG to THP-1 cells was assessed using the XTT cell proliferation and viability assay kit (Roche), according to the supplier's protocol. Briefly, cells were seeded at 50,000 cells/well. Various concentrations of compounds

were added to each well, in triplicate, in the presence of 25 µg mL^{-1} of gentamicin. Cells

were incubated for 72 h before measuring viability using the XTT assay. XTT reagents

were prepared according to the manufacturer's instructions. Following incubation, 50 µL

of XTT labeling mixture was added per well, and cells were further incubated for 4 h at

37°C and 5% CO_2. Absorbance measurements were collected at 590 nm and 750 nm

(reference wavelength). Final data were expressed as absorbance readings at 590 nm

(minus 750 nm reference absorbance) at the respective compound concentrations.

Quantifying *Brucella* infection of THP-1 cells

THP-1 cells were seeded in a black-well, clear-bottom 96-well plate (Costar) at 50,000

cells/well, differentiated with 40 ng mL^{-1} PMA for 72 h. Two hours before infection, cells

were treated with compounds, and infected with *B. abortus* at 100 MOI, as described

above. Fresh compounds were added back into media containing 25 µg mL^{-1}

gentamicin, and cells were further incubated for 72 h. After incubation, cells were

washed three times with 1X PBS, and fixed for 10 min with 4% paraformaldehyde,

followed by three washes with 1X PBS. Cell nuclei were labeled with 1 µg mL^{-1} Hoechst

stain. Cells were imaged using a Leica-DMI6000B fluorescence microscope equipped

with a 20X/0.4 NA objective, Hamamatsu Orca-R2 camera, and Leica Application Suite

X. Forty images were captured in four replicate experiments: ten images per condition

per replicate. Cell Profiler (v2.1.1) was used to measure the number of intracellular

fluorescent Brucellae, and the number of THP-1 nuclei. The effect of the compounds on

B. abortus intracellular growth was assessed by calculating the number of bacteria per

cell, represented as a ratio of the total number of bacteria to the total number of nuclei

(*Brucella* per THP-1 nucleus) per image, which was calculated as an average of 40

images ± standard error of the mean. Statistical significance was determined using one-

way analysis of variance (ANOVA) followed by Dunnett's test.

To enumerate CFU at 1, 24, and 48 hours post-infection of THP-1, *B. abortus*

cells were resuspended in warmed RPMI and added at a multiplicity of infection (MOI)

of 100 CFU per macrophage cell. To synchronize infections after the addition of

Brucella cells, plates were centrifuged at 200 x *g* for 5 minutes. After a 30-minute

incubation, media was removed and replaced with RPMI supplemented with 100 µg mL$^-$

^{1} gentamycin to kill extracellular bacteria. Afterwards, samples were incubated for 1

hour, after which monolayers were washed with PBS and THP-1 cells were lysed by

addition of 0.01% Triton X-100. Serial dilutions were prepared and viable cells were

determined by plating on TSA plates. For longer time points, the RPMI was removed,

and replaced with RPMI supplemented with 50 µg mL^{-1} gentamycin. Additional samples

were lysed as above and serial dilutions plated. All THP-1 assays were performed in

triplicate utilizing independent *B. abortus* cultures.

ACKNOWLEDGEMENTS

We acknowledge the support staff of the Howard Taylor Ricketts Regional

Biocontainment Laboratory. We also acknowledge and thank Barry Bochner and Aretha

Fiebig for thoughtful feedback and critical evaluation of the manuscript. This study was

funded in part by National Institutes of Health grant R01AI107159 to SC, and the

AUTHOR CONTRIBUTIONS

DC and SC conceived and designed the experiments, analyzed the data, and wrote the

manuscript. DC and JW performed the experiments. All authors reviewed and approved

the final manuscript.

REFERENCES

1. **Galvan-Pena S, O'Neill LA.** 2014. Metabolic reprograming in macrophage

    polarization. Front Immunol **5:**420.

2. **Pearce EL, Pearce EJ.** 2013. Metabolic pathways in immune cell activation and

    quiescence. Immunity **38:**633-643.

3. **Kelly B, O'Neill LA.** 2015. Metabolic reprogramming in macrophages and

    dendritic cells in innate immunity. Cell Res **25:**771-784.

4. **Warburg O, Wind F, Negelein E.** 1927. The metabolism of tumors in the body. J

    Gen Physiol **8:**519-530.

5. **Bedard K, Krause KH.** 2007. The NOX Family of ROS-Generating NADPH

    Oxidases: Physiology and Pathophysiology. Physiol Rev **87:**245-313.

6. **Nathan CF, Hibbs JB.** 1991. Role of nitric oxide synthesis in macrophage

    antimicrobial activity. Curr Opin Immunol **3:**65-70.

7. **Luckhart S, Pakpour N, Giulivi C.** 2015. Host-pathogen interactions in malaria:

    cross-kingdom signaling and mitochondrial regulation. Curr Opin Immunol **36:**73-

    79.

8. **Koshiba T.** 2013. Mitochondrial-mediated antiviral immunity. Biochim Biophys

    Acta **1833:**225-232.

9. **Li T, Xu Y, Liu L, Huang M, Wang Z, Tong Z, Zhang H, Guo F, Chen C.** 2016.

    Brucella Melitensis 16M Regulates the Effect of AIR Domain on Inflammatory

    Factors, Autophagy, and Apoptosis in Mouse Macrophage through the ROS

    Signaling Pathway. PLoS One **11:**e0167486.

10. **Stavru F, Bouillaud F, Sartori A, Ricquier D, Cossart P.** 2011. *Listeria*

monocytogenes transiently alters mitochondrial dynamics during infection. Proc

Natl Acad Sci U S A **108:**3612-3617.

11. **Owen OE, Kalhan SC, Hanson RW.** 2002. The Key Role of Anaplerosis and

Cataplerosis for Citric Acid Cycle Function. J Biol Chem **277:**30409-30412.

12. **Xavier MN, Winter MG, Spees AM, den Hartigh AB, Nguyen K, Roux CM,**

Silva TMA, Atluri VL, Kerrinnes T, Keestra AM, Monack DM, Luciw PA,

Eigenheer RA, Bäumler AJ, Santos RL, Tsolis RM. 2013. PPARγ-Mediated

Increase in Glucose Availability Sustains Chronic *Brucella abortus* Infection in

Alternatively Activated Macrophages. Cell Host Microbe **14:**159-170.

13. **Roop RM, Caswell CC.** 2013. Bacterial Persistence: Finding the "Sweet Spot".

Cell Host Microbe **14:**119-120.

14. **Eisele NA, Ruby T, Jacobson A, Manzanillo PS, Cox JS, Lam L, Mukundan**

L, Chawla A, Monack DM. 2013. *Salmonella* Require the Fatty Acid Regulator

PPARδ for the Establishment of a Metabolic Environment Essential for Long-

Term Persistence. Cell Host Microbe **14:**171-182.

15. **Cardaci S, Desideri E, Ciriolo MR.** 2012. Targeting aerobic glycolysis: 3-

bromopyruvate as a promising anticancer drug. J Bioenerg Biomembr **44:**17-29.

16. **Ko YH, Smith BL, Wang Y, Pomper MG, Rini DA, Torbenson MS, Hullihen J,**

Pedersen PL. 2004. Advanced cancers: eradication in all cases using 3-

bromopyruvate therapy to deplete ATP. Biochem Biophys Res Comm **324:**269-

275.

17. **Granchi C, Calvaresi EC, Tuccinardi T, Paterni I, Macchia M, Martinelli A,**

Hergenrother PJ, Minutolo F. 2013. Assessing the differential action on cancer

cells of LDH-A inhibitors based on the N-hydroxyindole-2-carboxylate (NHI) and

malonic (Mal) scaffolds. Org Biomol Chem **11:**10.1039/c1033ob40870a.

18. **Calvaresi EC, Granchi C, Tuccinardi T, Di Bussolo V, Huigens RW, Lee HY,**

Palchaudhuri R, Macchia M, Martinelli A, Minutolo F, Hergenrother PJ. 2013.

Dual Targeting of the Warburg Effect with a Glucose-Conjugated Lactate

Dehydrogenase Inhibitor. Chembiochem **14:**2263-2267.

19. **Wick AN, Drury DR, Nakada HI, Wolfe JB.** 1957. Localization of the primary

metabolic block produced by 2-deoxyglucose. J Biol Chem **224:**963-969.

20. **Bochner BR, Siri M, Huang RH, Noble S, Lei XH, Clemons PA, Wagner BK.**

2011. Assay of the multiple energy-producing pathways of mammalian cells.

PLoS One **6:**e18147.

21. **He Y, Reichow S, Ramamoorthy S, Ding X, Lathigra R, Craig JC, Sobral BW,**

Schurig GG, Sriranganathan N, Boyle SM. 2006. *Brucella melitensis* triggers

time-dependent modulation of apoptosis and down-regulation of mitochondrion-

associated gene expression in mouse macrophages. Infect Immun **74:**5035-

5046.

22. **Lobet E, Letesson JJ, Arnould T.** 2015. Mitochondria: A target for bacteria.

Biochem Pharmacol **94:**173-185.

23. **Ma C, Wickham ME, Guttman JA, Deng W, Walker J, Madsen KL, Jacobson**

K, Vogl WA, Finlay BB, Vallance BA. 2006. *Citrobacter rodentium* infection

causes both mitochondrial dysfunction and intestinal epithelial barrier disruption

in vivo: role of mitochondrial associated protein (Map). Cell Microbiol **8:**1669-

1686.

24. **Lyamzaev KG, Pletjushkina OY, Saprunova VB, Bakeeva LE, Chernyak BV,**

Skulachev VP. 2004. Selective elimination of mitochondria from living cells

induced by inhibitors of bioenergetic functions. Biochem Soc Trans **32:**1070-

1071.

25. **Abu Kwaik Y, Bumann D.** 2015. Host Delivery of Favorite Meals for Intracellular

Pathogens. PLoS Pathog **11:**e1004866.

26. **Mehrotra P, Jamwal SV, Saquib N, Sinha N, Siddiqui Z, Manivel V,**

Chatterjee S, Rao KV. 2014. Pathogenicity of *Mycobacterium tuberculosis* is

expressed by regulating metabolic thresholds of the host macrophage. PLoS

Pathog **10:**e1004265.

27. **Moulder JW.** 1970. Glucose Metabolism of L Cells Before and After Infection

with *Chlamydia psittaci*. J Bacteriol **104:**1189-1196.

28. **Mehta M, Sonawat HM, Sharma S.** 2005. Malaria parasite-infected erythrocytes

inhibit glucose utilization in uninfected red cells. FEBS Letters **579:**6151-6158.

29. **El-Bacha T, Menezes MMT, Azevedo e Silva MC, Sola-Penna M, Da Poian**

AT. 2004. Mayaro virus infection alters glucose metabolism in cultured cells

through activation of the enzyme 6-phosphofructo 1-kinase. Mol Cell Biochem

266:191-198.

30. **Anderson JD, Smith H.** 1965. The Metabolism of Erythritol by *Brucella Abortus*.

J Gen Microbiol **38:**109-124.

31.　**Smith H, Tang CM, Exley RM.** 2007. Effect of Host Lactate on Gonococci and

Meningococci: New Concepts on the Role of Metabolites in Pathogenicity. Infect

Immun **75:**4190-4198.

32.　**El Sayed SM, El-Magd RM, Shishido Y, Chung SP, Diem TH, Sakai T,**

Watanabe H, Kagami S, Fukui K. 2012. 3-Bromopyruvate antagonizes effects

of lactate and pyruvate, synergizes with citrate and exerts novel anti-glioma

effects. J Bioenerg Biomembr **44:**61-79.

33.　**Barban S, Schulze HO.** 1961. The Effects of 2-Deoxyglucose on the Growth and

Metabolism of Cultured Human Cells. J Biol Chem **236:**1887-1890.

34.　**Czyz DM, Jain-Gupta N, Shuman HA, Crosson S.** 2016. A dual-targeting

approach to inhibit Brucella abortus replication in human cells. Sci Rep **6:**35835.

35.　**Herrou J, Czyż DM, Willett JW, Kim HS, Chhor G, Babnigg G, Kim Y,**

Crosson S. 2016. WrpA Is an Atypical Flavodoxin Family Protein under

Regulatory Control of the *Brucella abortus* General Stress Response System. J

Bacteriol **198:**1281-1293.

36.　**Kim HS, Caswell CC, Foreman R, Roop RM, Crosson S.** 2013. The *Brucella*

abortus General Stress Response System Regulates Chronic Mammalian

Infection and Is Controlled by Phosphorylation and Proteolysis. J Biol Chem

288:13906-13916.

FIG 1. Mitochondrial function is perturbed in THP-1 cells infected with *B. abortus.*

(A) Mammalian Phenotype MicroArray (PM) plate designed to measure mitochondrial function. The ability of THP-1 cells infected with *B. abortus* at increasing multiplicity of infection (MOI) to utilize metabolic substrates was assessed by measuring the reduction of a tetrazolium dye (see Materials and Methods). **(B)** An MTT cell viability assay was performed in the presence of glucose to assess the effect of *B. abortus* infection on human cell survival at the conclusion of the mammalian PM assay. Dye reduction by **(C)** uninfected and **(D)** *Brucella*-infected human cells (MOI 125) measured as a time-dependent absorbance readout. Cells were cultured in minimal medium supplemented either with glucose (control) or central metabolic substrates. Brackets indicate substrates (α-ketoglutaric acid, β-hydroxybutyric acid, and pyruvic acid), which enter the

584   mitochondria, thereby providing a direct readout of mitochondrial function. Bars along

585   the X axes represent time (0–150 min) in 15-min increments. Dotted lines mark the

586   highest reading for utilization of glucose and pyruvic acid in uninfected THP-1 cells.

| DIC | Mitochondria | *B. abortus* | Nuclei | Overlay |

FIG 2. Intracellular *B. abortus* induces morphological changes of mitochondria in HeLa cells.

Two fields showing mitochondrial staining (MitoTracker, green), *B. abortus*-Tn7mCherry (mCherry fluorescence, red), nuclei (Hoechst stain, blue), and the corresponding image with differential interference contrast (DIC) microscopy. Red arrows indicate cells with visible intracellular *B. abortus*, white arrows indicate cells for which no infection is evident. Cells not infected with *B. abortus* have typical morphology and mitochondrial structure. Scale bar = 50 µm.

FIG 3. Host cell utilization of amino acids as a carbon source before and after *B.*

abortus **infection.**

Heatmap representing host cell metabolism as measured colorimetrically via a redox

dye (see Materials and Methods). Time-dependent metabolic change of non-infected

versus *B. abortus*-infected THP-1 cells in the presence of amino acids are plotted. Color

is an arbitrary scale from black (no reduction of the dye, 0.00) to red (maximum

reduction of the dye, 0.50).

FIG 4. *B. abortus* **induces host cell lactate production and utilizes lactate as a**
carbon and energy source as efficiently as glucose and erythritol.

(A) Lactate levels in THP-1 cells upon infection with *B. abortus* at different multiplicities
of infection. **(B)** Metabolism of *B. abortus* measured on carbon/energy sources lactate,
erythritol, or glucose. Data show an average of two independent experiments ±
standard deviation.

FIG 5. *B. abortus* **L-lactate dehydrogenase deletion strain Δ***lldD* **cannot utilize lactic acid in axenic culture and is attenuated in a macrophage infection model.**

(A) Phenotype MicroArray carbon utilization measurements of wild-type (WT) *B. abortus* and L-lactate dehydrogenase (LDH) mutant (Δ*lldD*) strains. Metabolic activities for growth on 95 distinct carbon substrates (see Materials and Methods) are plotted against each other; the dotted line marks a 99% prediction band of a linear regression. Correlation values above the line represent a gain-of-function phenotype and values below the line represent a loss-of-function phenotype. L-lactic acid and D-malic acid are marked as loss-of-function and gain-of-function, respectively **(B)** Graph representing time-dependent utilization of lactic acid as a carbon energy source, and measured by a metabolic indicator dye (see Materials and Methods). **(C)** Graph representing axenic growth of the WT and Δ*lldD* strains in glucose-containing *Brucella* broth. **(D)** Intracellular

628 replication of the WT and Δ*lldD* strains in macrophage cell infection model. Intracellular

629 Brucellae were quantified as $\log_{10}$ colony forming units (CFU) per well at 1, 24, and 48 h

630 post-infection. Lines represent WT (●), Δ*lldD* mutant (■), and complement (▲). n=3,

631 Error bars represent standard deviation.

FIG 6. Pharmacological disruption of the host glycolytic pathway and lactate production inhibits replication of intracellular *B. abortus*.

(A) Model depicting *Brucella*-dependent modulation of carbon metabolism and activation of aerobic glycolysis, resulting in the production of lactate. *B. abortus* utilizes lactate as efficiently as preferred substrates glucose and erythritol. Pharmacological inhibitors of glycolysis, 3-Bromopyruvic acid (3-BPA), NHI-2, and 2-deoxy-D-glucose (2-dG), inhibit intracellular growth of *B. abortus*: **(B)** Graphs represent the effect of 3-BPA, NHI-2, and 2-dG on intracellular replication of *B. abortus* as quantified by enumerating fluorescent bacteria per THP-1 nucleus; **(C)** cytotoxicity of inhibitory compounds to THP-1 cells, and **(D)** effect of metabolic inhibitors on *B. abortus* axenic growth/metabolism. The effect of anti-glycolytic compounds on infection was assessed by measuring the average ratio of bacteria-to-nuclei in THP-1 cells treated with each of the compounds. Statistical significance was evaluated from four biological replicates (10 images each) using one-way analysis of variance (ANOVA) followed by Dunnett's post test **$p< 0.01$, ***$p<0.0001$). Error bars represent ± standard deviation from the mean; an XTT assay was used to measure compound cytotoxicity to THP-1 cells; *Brucella* axenic growth/metabolism was assayed by measuring tetrazolium-based dye reduction at 630 nm **(E)** Fluorescence images of nuclei (blue) and intracellular *B. abortus* (pseudocolored green) reflect the efficacy of 3-BPA, NHI-2, and 2-dG to inhibit *B. abortus* replication in the intracellular niche. Scale bars = 10 μm (full view) and 50 μm (magnified insets, marked with white box in full view). Dotted line = half maximal inhibitory concentration (IC50).

FIG 7. Model of how *B. abortus* exploits the shift to Warburg-like metabolism in macrophages.

Data presented in this study support a model in which inhibition of tricarboxylic acid (TCA) cycle metabolism upon infection biases the host cell toward the production of lactic acid. **(A)** Uninfected macrophages exhibit typical central carbon metabolism in which pyruvate enters the TCA cycle. **(B)** *B. abortus* infection biases the macrophage toward aerobic glycolysis and lactate production. *B. abortus* efficiently utilizes lactic acid via the lactate dehydrogenase (LDH) enzyme, LldD. A strain lacking *lldD* is compromised in axenic lactic acid metabolism and in intracellular replication in a THP-1 macrophage model. Pharmacological inhibition of host glycolysis or LDH-A inhibits *B. abortus* replication.