Individual heterogeneity and capture-recapture models: what, why and how?

Olivier Gimenez[1], Emmanuelle Cam[2] and Jean-Michel Gaillard[3]

Olivier Gimenez (olivier.gimenez@cefe.cnrs.fr), CEFE UMR 5175, CNRS, Université de Montpellier, Université Paul-Valéry Montpellier, EPHE, 1919 Route de Mende, 34293 Montpellier Cedex 5, France. – **Emmanuelle Cam**, Evolution et Diversité Biologique, UMR 5174, CNRS-ENSFEA-IRD, Université Toulouse 3 Paul Sabatier – Bâtiment 4R1, 118 route de Narbonne, 31062 Toulouse Cedex 9, France. – **Jean-Michel Gaillard**, Laboratoire de Biométrie et Biologie Evolutive, UMR 5558, Université de Lyon, Université Lyon 1, 69622 Villeurbanne, France.

Variation among and within individuals in life history traits is ubiquitous in natural

populations. When affecting fitness-related traits like survival or reproduction, individual

heterogeneity plays a key role in population dynamics and life history evolution. However, it

is only recently that properly accounting for individual heterogeneity when studying

population dynamics of free-ranging populations has been made possible through the

development of appropriate statistical models. We aim here to review case studies of

individual heterogeneity in the context of capture-recapture models for the estimation of

population size and demographic parameters with imperfect detection. First, we define what

individual heterogeneity means and clarify the terminology used in the literature. Second, we

review the literature and illustrate why individual heterogeneity is used in capture-recapture

studies by focusing on the detection of life-history trade-offs, including senescence. Third, we

explain how to model individual heterogeneity in capture-recapture models and provide the

code to fit these models (https://github.com/oliviergimenez/indhet_in_CRmodels). The

distinction is made between situations in which heterogeneity is actually measured and

situations in which part of the heterogeneity remains unobserved. Regarding the later, we

outline recent developments of random-effect models and finite-mixture models. Eventually,

we discuss several avenues for future research.

Key words: actuarial senescence; Arnason-Schwarz model; Cormack-Jolly-Seber model;

frailty; hidden Markov models; individual covariates; life-history trade-offs; mark-recapture

models; mixed models; mixture models; multievent models; multistate models; random-effect

models; survival estimation.

Introduction

Individual variation is at the core of the evolution of traits by the means of natural selection and exists within any population of living organisms. Individual variation occurs in virtually all traits, including fitness components such as reproduction and survival (Clutton-Brock 1988, Newton 1989). However, the amount of individual variation in a given trait in a given population varies a lot both within and across species. Among-individual differences in phenotypic attributes such as age (Caughley 1966, Emlen 1970), sex (Short and Balaban 1994), body mass (Sauer and Slade 1987), or personality (Dingemanse and Dochtermann 2013), in genotype (Coulson et al. 2011), in habitat use or selection such as home range size or quality (Mcloughlin et al. 2007), or in prey selection (Estes et al. 2003) have all been reported to affect most life history traits. More recently, both current and early-life environmental conditions encountered by individuals throughout their life have been shown to generate individual differences in life history traits (Douhard et al. 2014, Berger et al. 2015).

The potential role of individual heterogeneity in terms of population ecology has been pointed out more than 30 years ago (Lomnicki 1978, Johnson et al. 1986) and repeatedly reported since (Bolnick et al. 2011, Kendall et al. 2011). Thanks to the increasing availability of high quality data collected during long-term individual monitoring of vertebrate populations (Clutton-Brock and Sheldon 2010), assessing the magnitude of individual heterogeneity, identifying its origin, and quantifying its consequences has become a specific objective in many population studies.

From these studies (reviewed in Table 1), we can conceptually envision three broad patterns of individual heterogeneity when considering a set of life-history traits, e.g. demographic parameters, independently of any methodological approach used to model these demographic tactics.

2   [Table 1 around here]

We will retain Stearns (1976)'s definition of a tactic, "a set of co-adapted

[demographic] traits designed, by natural selection, to solve particular ecological problems".

In the simplest case, individual heterogeneity corresponds to the random variation observed

independently in each of the traits. In that case, there is no covariation among demographic

traits and no life history tactic can emerge. Life history tactics can show up in response to

marked differences among individuals in terms of constraints (of genetic, developmental, or

environmental origin). The axis of demographic variation will thus involve a low-high

continuum of performance opposing individuals weakly constrained that will perform

extremely well in terms of both survival and reproduction, to individuals subjected to high

constraints (sometimes called "runt", Koenig et al. 1995) that will perform extremely poorly.

Individual heterogeneity in that case will lead the axis of demographic variation to correspond

to a low-high continuum of individual fitness, and is often designated as a continuum of

individual quality (Wilson and Nussey 2010). Alternatively, individual heterogeneity can be

associated with a set of different life history tactics so that each tactic is characterized by the

same mean individual fitness. For instance, some individuals will allocate a lot to

reproduction and pay a cost in terms of decreased survival, whereas others will allocate a lot

to avoid mortality risks and pay a cost in terms of reduced reproduction, leading the negative

co-variation between survival and reproduction expected from the allocation principle (Cody

1966) to show up.

Owing to the multiplicity of factors that shape individual heterogeneity, it is

impossible to measure empirically its amount. For a given trait, we can thus distinguish a

measured individual heterogeneity using for instance phenotypic attributes such as age, sex,

and size from an *unmeasured* individual heterogeneity that includes all the remaining

variation for given age, sex and size (Plard et al. 2015). Until recently, this *unmeasured*

individual heterogeneity was most often neglected. Assessing unmeasured individual

heterogeneity is especially tricky when studying survival (or mortality) because this trait

simply corresponds to a state shift for a given individual. Thus, an individual dying at 5 years

of age will have survived over the first five years in a row and then will have died at 5 years,

leading the standard mixed model approach generally used for assessing individual

heterogeneity in most traits (van de Pol and Verhulst 2006, van de Pol and Wright 2009) not

directly applicable. However, CR models do provide a general and flexible framework for

estimating and modeling both population size and demographic parameters (including

survival, dispersal and recruitment) in the face of imperfect detection that is inherent to

populations in the wild (Gimenez et al. 2008). These methods rely on the longitudinal

monitoring of individuals that are marked (or identifiable) ideally at birth, and then

encountered (*i.e.* recaptured or seen) on subsequent occasions. The first CR methods that dealt

with individual heterogeneity were developed with the aim to get unbiased estimates of

population size in presence of differential individual responses to trapping (Otis et al. 1978).

The context has changed in recent years, and CR studies now often focus on the process of

individual heterogeneity per se in order to assess the diversity of life history trajectories

within populations, to test for the existence of different life history tactics within populations,

or to assess the differential susceptibility of individuals to environmental insults.

Here we aim at providing a review of individual heterogeneity in the context of CR

studies. We first define *what* we mean by individual heterogeneity by examining the landmark

papers on the subject and clarifying the terminology with regard to more recent uses of the

concept. Then we review the literature and illustrate *why* individual heterogeneity is used in

CR studies by focusing on the detection of life-history trade-offs, including senescence. In a

third section, we explain *how* to model individual heterogeneity in CR models. The distinction is made between situations in which heterogeneity can be explicitly handled through the use of states (e.g. breeding or disease states) or individual (time-varying or not) covariates (e.g., age, or phenotype) and situations in which part of the heterogeneity remains unobserved. Regarding the later, we outline recent developments of random-effect models and finite-mixture models. Eventually, we discuss several avenues for future research.

What is individual heterogeneity?

A bit of history and some definitions

In CR modeling, consideration of heterogeneity among individuals of a population in demographic parameters (e.g., survival or the probability of successful reproduction) has a long history: "For predictive or modeling purposes, [...], heterogeneity can lead to seriously misleading conclusions, particularly if the product of two or more parameters is involved, and heterogeneity affects both of them" (Johnson et al. 1986). Early work emphasized the distinction between situations where members of populations differ with respect to some measurable attribute (e.g., sex, age), and situations where "heterogeneity is not clearly identified with a measurable variable" (Johnson et al. 1986). In the latter situation, developing methods that account for heterogeneity among individuals to estimate demographic parameters is more difficult. Early efforts toward this end echo their contemporary studies of heterogeneity in mortality risk and of aging in human demography (Vaupel et al. 1979, Manton et al. 1981, Hougaard 1984): "Unrecognized heterogeneity can lead to biased inferences, especially in time or age effects in cohort studies" (Johnson et al. 1986). In wildlife ecology, one of the earliest studies that investigated the consequences of unmeasured (and sometimes impossible to measure) variation among individuals on survival probability

concerned the estimation of nest success probability using longitudinal data from nest activity

(Green 1977, Johnson 1979). The authors of early papers on CR modeling were aware of the

contribution of human demographers to the development of models taking heterogeneity

among individuals into account to estimate changes in mortality risk throughout life (papers

cited in e.g. Johnson, Burnham & Nichols 1986 like, e.g., Keyfitz & Littman 1979, Vaupel,

Manton & Stallard 1979b and Manton, Stallard & Vaupel 1981). However, according to

Johnson et al. (1986), "the impact of such heterogeneity has been recognized only

occasionally in animal ecology, possibly because it is difficult to deal with, and it is often

relatively unimportant in many estimation problems."

Clearly, the issue raised in early CR studies is that the assumption of homogeneity of

populations can lead to flawed inferences about identified parameters like survival probability

or population size. In situations where survival probability varies with age, this issue has

sometimes been called a Simpson's paradox in statistics, or an "ecological fallacy" (Kramer

1983), and has been illustrated by Cohen (1986) as follows: "The crude death rate of

population A may be less than that of country B even if every age-specific death rate of

country A is greater than each corresponding one of country B". If populations are stratified

according to variables that have not been considered yet, inequality of rates can be reversed,

and any demographic parameter can be concerned. Papers by Green (1977), Johnson (1979),

Johnson et al. (1986) and the very influential paper by Vaupel & Yashin (1985) have all used

analogous examples of situations with two groups to explain the consequences of

unrecognized heterogeneity on inferences about mortality in wildlife ecology and human

demography, respectively.

In the context of closed populations (i.e. assuming a population with no immigration,

no emigration, no recruitment and no mortality), CR models are restricted to the estimation of

population size and can account for individual heterogeneity in the probability of being

detected (e.g. Burnham & Overton 1979, Pollock 1980, Pollock *et al.* 1990, Link 2004,

Pledger 2005). In such a situation, "Individuals with high detection probabilities would tend

to appear in the encountered sample in greater proportion than they occur in the population"

(White and Cooch 2017). In closed population models, data are collected several times during

the period when assumptions characterizing closed populations hold. In such models, the

probability of detection is assumed to vary among individuals in relation to, e.g., sex or age-

related behavioral differences and more recently to space (e.g. Borchers 2012). In some

studies, populations are not assumed to be composed of clusters (e.g. sex, age-classes) with

different detection probabilities, but each individual is "assumed to have its own unique

capture probability which remains constant over all the sampling times" (Pollock 1980,

Pollock et al. 1990). Although applied to detection or recapture probability in these studies,

there is a striking analogy between the above definition by Pollock and one of the oldest

definitions of individual frailty in human demography, where populations are assumed to be

composed of members who differ in their individual susceptibility to death (Vaupel et al.

1979b, Manton et al. 1981). In CR studies where heterogeneity in the probability of detecting

an individual cannot be identified using measured variables, investigators can assume that

there is a distribution of individual detection probabilities and use models with individual

random effects (White and Cooch 2017). In such situations, early approaches have used

Jackknife estimators (Burnham and Overton 1979). Mixture models have also been used later

(e.g. Norris & Pollock 1996, Pledger 2000, Morgan & Ridout 2008), where populations are

assumed to be composed of several hidden groups with different detection probabilities.

Again, the latter approach echoes early studies of heterogeneity in mortality risk in human

populations (Vaupel and Yashin 1985).

In the context of open populations (i.e. allowing immigration and emigration and/or

recruitment and mortality to occur), models are used to estimate a variety of demographic

parameters (e.g. survival rates, transition probabilities among reproductive states in

successive years). Early papers (Cormack 1964, Jolly 1965, Seber 1965) have assumed

homogeneous populations. Stratification according to age-classes (Manly and Parr 1968,

Pollock 1981) and groups (Lebreton et al. 1992) was one of the first attempts to accommodate

variation among individuals in survival probability. Arnason (1972, 1973), Schwarz et al.

(1993), Hestbeck et al. (1991), and Nichols et al. (1994) laid the ground for models

accounting for the fact that individuals might not belong to the same cluster during their entire

life (Lebreton and Pradel 2002), but might move among states (e.g. locations, breeding status)

in a stochastic manner (as opposed to movement among age-classes in a deterministic

manner).

Naturally, early papers drew the same distinction as for closed populations between

situations where individuals differ with respect to some measurable attribute (e.g., sex, age)

and situations where "heterogeneity is not clearly identified with a measurable variable"

(Pollock 1980, Johnson et al. 1986). Because open-population models can also be used to

estimate population size, early work focused on the consequences of heterogeneity in

detection probability on population size estimation (Carothers 1973). Other early studies

focused on the consequences of heterogeneity in mortality risk on nest success estimation

(Johnson 1979). Just like for detection probability, two approaches have been used to account

for unobserved sources of heterogeneity in survival probability (Johnson et al. 1986): finite

mixture models (Pledger et al. 2003) and random effects models (Royle 2008, Gimenez and

Choquet 2010). In both classes of models, individual heterogeneity refers to permanent

differences among individuals in demographic parameters. In mixture models, the population

is assumed to be composed of groups of individuals that cannot be identified using observable

criteria. This is consistent with early work in human demography about variation in mortality

risk over life (Vaupel and Yashin 1985). In early CR studies, individual heterogeneity was

defined as "variation in survival probabilities among individuals after taking into account

variability due to age, sex, or time" (Rexstad and Anderson 1992), and describes a situation

where individuals are assumed to have their own unique survival probability that remains

constant over all the sampling times (Conner and White 1999). This definition matches

exactly the concept of unobserved individual frailty proposed by Vaupel, Manton & Stallard

(1979) where frailty designates the risk of a given individual (that is constant throughout its

lifespan) to die at a given age relative to the average risk of all individuals in the population to

die at this age. Early studies assuming a continuous distribution in survival probability in CR

models have also highlighted the methodological difficulties encountered in the 1990's

(Rexstad and Anderson 1992, Burnham and Rexstad 1993).

Individual heterogeneity in contemporary CR studies

Demographic parameters are the target parameters to estimate in CR models (Lebreton et al.

1992, 2009). Other parameters such as detection probability are required to estimate

demographic parameters from CR models; detection probability is relevant to sampling in

wild populations, but is not a demographic parameter. However, the impact of individual

heterogeneity on every type of parameter has consequences on demographic parameter

estimation. CR models allow the estimation of all types of parameters.

To address individual heterogeneity in demographic parameters, the survival process

(alive vs. dead), or the reproductive process (e.g., breeder vs. non-breeder or success vs.

failure) is treated as a random variable. Individual heterogeneity measures differences among

individuals in model parameters (e.g., Marzolin, Charmantier & Gimenez 2011). Today, a

broad range of approaches is used to account for individual heterogeneity in CR studies.

Individual heterogeneity is understood as any source of variation among individuals in

demographic parameters that cannot be accounted for by temporal or spatial heterogeneity

alone, with a particular focus on the fate of individuals, their early development conditions,

their ontogeny, or their past allocation to reproduction as experienced breeders. Historically

dominating views of what 'heterogeneity' means in CR modeling depended on the class of

models used. For example, 'heterogeneity' models in papers focusing on closed populations

referred to models where "each animal has its own unique capture probability" (Pollock

1980). Conversely, for open population models, early work on 'heterogeneity' considered any

degree of stratification of populations, from discrete groups, or age-structured populations, to

a distinct survival probability value for each individual (Johnson et al. 1986). However,

'individual heterogeneity' has rapidly been reserved for situations where accounting for age-,

sex- or time-related variation in survival probability was insufficient (Rexstad and Anderson

1992). Advances in statistical methods over the past 40 years have progressively enlarged the

scope of models accounting for 'individual heterogeneity'. In CR studies, the sample scheme

involves attempting to detect individuals on a discrete time basis. Contrary to continuous-time

models used in human demography (Allison 1982), this specific feature must have eased the

development of models in the CR arena.

Different grain sizes of heterogeneity lead to different perceptions of 'heterogeneity',

but the current view of 'individual heterogeneity' incorporates a large range of biologically

relevant situations. At the lowest grain size individuals have their own unique demographic

value, like in the frailty context but with a possible variation during life (e.g. Johnson,

Burnham & Nichols 1986, Conner & White 1999, Enki, Noufaily & Farrington 2014). As

emphasized by Vaupel (2002) "All populations are heterogeneous. Two individuals of the

same age and sex in the same population can face very different chances of death. Some

individuals are frailer than others and the frail tend to die first". At a broader grain size,

heterogeneity corresponds to differences in demographic parameters among identifiable

categories of individuals (Cooch et al. 2002, Nichols 2002). Once included in the study,

individuals belong to the same cluster definitely (e.g. sex, hidden class; Péron *et al.* 2010b) or

temporarily (e.g. age, body condition, breeding state; Nichols *et al.* 1994). This broad grain

size heterogeneity also includes populations composed of several unobserved, latent groups of

individuals with different values of demographic parameters (i.e., when heterogeneity is

accounted for using mixture models; Péron *et al.* 2010b). Finally, contemporary studies still

draw a distinction between situations where investigators can identify heterogeneity using

measurable variables (e.g., identifiable groups; Drummond, Rodriguez & Oro 2011), or not

(mixture of hidden classes, Péron *et al.* 2010b, random effects models; Marzolin, Charmantier

& Gimenez 2011; hidden states, Johnson *et al.* 2016, see Table 1). What 'heterogeneity'

covers in CR models inherently depends on model specification. In all cases, populations are

considered as being heterogeneous.

From a biological viewpoint, individual heterogeneity in life history traits is often

considered to include two components.

(i)   Cases in which differences among individuals are shaped early in life and are

15   permanent during the course of the life correspond to a fixed heterogeneity. In such

16   cases, individual heterogeneity can be accounted for by using measurable

17   covariates a priori assumed to capture much of the individual heterogeneity (e.g.

18   rank of offspring in birds' clutch, Drummond, Rodriguez & Oro 2011 or body mass

19   at the end of the maternal care period in large mammals, Hamel *et al.* 2009).

20   Whether measurable individual features are translated into differences in estimates

21   of demographic parameters, and with which method, is part of the statistical

22   exercise. When measurable variables are missing, or insufficient to account

23   satisfactorily for heterogeneity (Hougaard 1991), investigators can assume a

24   discrete or continuous distribution of demographic parameters (Royle 2008, Péron

25   et al. 2010). In agreement with the concept of frailty, investigators assume that

there are differences in demographic parameters among individuals that cannot be

associated with measurable covariates and use latent variables to quantify it (e.g.

Hougaard 1995, Yashin *et al.* 2008, Cam *et al.* 2013, Hamel, Yoccoz & Gaillard

2014, Cam, Aubry & Authier 2016).

(ii)   However, not all the amount of individual heterogeneity is fixed. Individual

differences in a given trait at a given time are subjected not only to the influence of

early-life conditions, but also to current conditions both at that time and between

early life and that time. For instance, harsh environmental conditions occurring at

any time during the lifetime of a cohort are expected to increase individual

heterogeneity in life history traits, whereas favorable environmental conditions are

expected to decrease it (Lomnicki 1978). As above, in some situations, observable

variables are available to account for variation in individuals' demographic

parameters over life (e.g. age: Pollock 1981; group: Lebreton *et al.* 1992; state:

Nichols *et al.* 1994). However, when such observable variables are missing or

inefficient at capturing most individual heterogeneity (Hougaard 1991),

unobserved, latent traits changing over life can be used. Such cases correspond to

'dynamic frailty' (Pennell and Dunson 2006, Duffie et al. 2009, Chambert et al.

2013, Cam et al. 2013).

Why individual heterogeneity in a CR context?

Individual heterogeneity seen as a nuisance

The existence of individual heterogeneity has stimulated a large number of works in the early

CR literature, because failure to account for heterogeneity in detection rates can lead to biases

in estimate of abundance and demographic parameters. For example, in the context of closed

populations, models including variable detection rates among individuals have been

developed (Otis et al. 1978). As Eberhardt (1969) pointed out, "various sets of data indicate

that the equal-probability-of capture assumption is not fulfilled." In addition, unequal

detection rates may also lead to biases in survival probability or population growth in the

context of open populations (e.g. Carothers 1973, Schwarz 2001). This question is still

receiving attention in contemporary studies (Crespin et al. 2008, Cubaynes et al. 2010, Pradel

et al. 2010, Marescot et al. 2011, Oliver et al. 2011, Fletcher et al. 2012, Abadi et al. 2013),

and new methods are being developed to address heterogeneity, such as multievent models

(Pradel 2009).

Individual heterogeneity as a biological process

Looking backwards, methodological developments in the area of CR models have

considerably increased the relevance of CR studies to address questions not only in wildlife

ecology, but also in evolutionary ecology. A key feature of this development is the increased

ability of CR models to account for variation among individuals in demographic parameters

as well as detection probability at the scale assumed to be relevant in the studied context. This

scale can be the individual, rather than permanent groups (e.g. sex), or temporary aggregates

of individuals (e.g., reproductive states). The development of evolutionary biology as a

powerful conceptual and methodological framework for biological disciplines (e.g.

Dobzhansky 1973) has brought a new perspective on individual heterogeneity. In CR studies,

instead of being addressed because of potential biases in estimates of abundance, survival

probability or population growth rate (e.g., Crespin *et al*. 2008, Cubaynes *et al*. 2010, Pradel

et al. 2010, Oliver *et al*. 2011, Abadi, Botha & Altwegg 2013), individual heterogeneity has

become the focus of studies because of its biological relevance. For evolutionary biologists,

the individual level can be relevant to address natural selection if heritable variation is

expressed at this level (Chambert et al. 2014). In addition, variation among individuals in demographic parameters is relevant to population ecology and dynamics, whether it concerns traits that are heritable, or not, and whether it can be accounted for using observed variables, or not (e.g. Kendall & Fox 2002). Indeed, both population extinction risk and viability depend on the degree and structure of individual heterogeneity in survival probability and reproductive parameters (Conner and White 1999, Stover et al. 2012).

Starting from classes of models where demographic parameters varied with time (Jolly 1965), group of individuals, or age, the development of software programs to build multistate models (Arnason 1972, 1973) in the 1990's has considerably increased the attractiveness of CR models. These models indeed allow biologists to address questions about a large range of factors structuring a population, which determine individual sequences of states among which individuals move in a stochastic manner (Nichols et al. 1994, Nichols and Kendall 1995). These models triggered studies of life histories using CR models (e.g., Cam *et al*. 1998, Hadley, Rotella & Garrott 2007). A recently developed class of approaches, multievent models (Pradel 2005), has also helped biologists address questions about the influence of 'state' on demographic parameters (Sanz-Aguilar et al. 2011). Indeed, one of the difficulties in CR studies is that 'state' may not be observed with certainty, or even not be observed at all (Desprez et al. 2013). Moreover, the development of user-friendly software to build models with individual covariates has also stimulated work in evolutionary ecology using CR data (e.g. Gimenez, Grégoire & Lenormand 2009). The question of time-specific individual covariates with missing values is still a current issue (Bonner and Schwarz 2006). The ability of CR models to accommodate variation among individuals in demographic parameters raises the issue of methods of inference about model parameters in CR studies (e.g. Pledger & Schwarz 2002, Royle 2008, Gimenez & Choquet 2010). This issue is tightly linked to the level of stratification of populations, or of aggregation of observations (Cooch et al. 2002). As

emphasized by Nichols (2002): "If we view an individual organism's fate or behavior at any

point in space and time as a unique event not capable of informing us about the likelihood of

the event for other individuals or points in space and time, then generalization and prediction

become impossible". To allow formal statistical inferences about variation in demographic

parameters and detection probability in populations, CR models rely on assumptions, notably

regarding unobserved heterogeneity (e.g., a distribution of random effects; Royle 2008, or a

mixture model; Péron *et al.* 2010b, Marescot *et al.* 2011).

Assessing senescence in the wild: an increasingly popular focus of CR studies

The process of senescence, which can be interpreted in the context of the allocation principle

as the trade-off opposing performance during early life and performance in late life (Baudisch

and Vaupel 2012, Lemaître et al. 2015), has been the focus of a large number of empirical

studies during the last decade (see Nussey *et al.* 2013 for a review). As imperfect detection of

individuals is the rule in free-ranging populations (Gimenez et al. 2008), CR has become the

gold standard to measure reliably actuarial senescence in the wild (e.g. Loison *et al.* 1999,

Bouwhuis *et al.* 2012). The question of level of inference has recently emerged as a critical

point in CR studies of senescence (Péron et al. 2010, Marzolin et al. 2011). Based on early

work by human demographers addressing heterogeneity in mortality risk (e.g. Vaupel,

Manton & Stallard 1979, Vaupel & Yashin 1985), ecologists have often used the concept of

frailty. However, Vaupel & Yashin (1985) considered the case of a heterogeneous population

with two classes of individuals, frail and robust ones. As time passes and individuals age,

there is a disjunction between the variation of the mean survival probability (i.e., when

pooling frail and robust individuals) with age, and the variation in survival probability with

age within each group (Figures 1, 2). Ignoring heterogeneity in mortality risk may lead to

flawed inferences about aging rate (Vaupel and Yashin 1985, Zens and Peart 2003), a

phenomenon documented in wild animals using CR models (Nussey et al. 2008, Péron et al.

2016). This phenomenon has long been acknowledged in wildlife studies for life stages other

than senescence (e.g., nest mortality; Green 1977 and Johnson, Burnham & Nichols 1986, and

Burnham & Rexstad 1993 in the context of CR studies). The consequences of ignoring

individual heterogeneity in survival probability have also been investigated in CR studies

using special datasets with perfect detection of individuals (e.g., Cam *et al.* 2002a, 2013,

Wintrebert *et al.* 2005, Fox *et al.* 2006, Aubry *et al.* 2011, Knape *et al.* 2011).

      The requirement of accounting for heterogeneity in survival studies was raised by

human demographers very early (which distribution to use to account for individual

heterogeneity in mortality risk, Manton et al. 1986) and has become a key topic in ecology.

Some demographers argued that observable criteria may not account for individual

heterogeneity in a satisfactory manner, and developed mixed models or mixture models for

time to event data (Kannisto 1991, Abbring and Van Den Berg 2007). The debate about the

appropriate distribution to consider is also taking place in ecology (e.g. Gimenez et al. 2010,

Péron et al. 2010, 2016). However, in CR studies, biologists use discrete data (e.g., mixed

binomial models for survival), which may lead to fewer issues with parameter identifiability

and assumptions than hazard models with frailty (Wienke 2010). Moreover, to some extent,

the idea of addressing heterogeneity using a distribution of latent demographic traits is

coherent with approaches to quantify variation in populations that are familiar to evolutionary

biologists, namely, variances in traits in quantitative genetics (Lynch and Walsh 1998,

Chambert et al. 2013, 2014). Recently capture-recapture animal models (CRAM) have been

developed to estimate heritability of demographic parameters (Papaïx *et al.* 2010).

Detection of trade-offs between life history traits

Trade-offs are one of the cornerstones of the theory of life histories (Roff 1992). They are

based on the principle of allocation (Cody 1966) and express the idea that individuals face

with limited amount of energy and have thereby to share energy among various functions so

that individuals allocating a lot into current reproduction cannot allocate as much into survival

or future reproduction (Roff 1992). However, empirical analyses have often failed to detect

trade-offs in the wild because of individual heterogeneity in resource acquisition. van

Noordwijk & de Jong (1986) indeed demonstrated that positive associations between current

reproduction and future survival or reproduction occur when individual heterogeneity in

resource acquisition is greater than individual heterogeneity in resource allocation. The

development of multistate models has attracted evolutionary biologists to study trade-offs

within the CR arena (Cam et al. 1998, Yoccoz et al. 2002). Individuals are assumed to make

allocation decisions according to their own state (McNamara and Houston 1996).

Consequently, any unobserved feature of 'state' may explain why trade-offs are not detected.

Experimental approaches may help unveil trade-offs (Reznick 1985), but may also butt

against heterogeneity (e.g. Festa-Bianchet, Gaillard & Jorgenson 1998, Yoccoz *et al.* 2002).

At the extreme, each individual can be assumed to be in a unique 'state' that cannot be

measured, and trade-offs might not be detected. Some observational CR studies have provided

evidence of trade-offs after identifying traits that reliably described changes in individual state

(e.g. social rank, mass, etc., Hamel *et al.* 2009), by taking advantage of unfavorable

conditions (e.g. Descamps *et al.* 2009) or by distinguishing direct from indirect effects

(Cubaynes et al. 2012a). The development of hierarchical CR models with individual

heterogeneity has allowed investigators to assume a distribution of latent life history traits in

populations (Royle 2008, Gimenez and Choquet 2010). In particular, Buoro et al. (2010,

2011) have been successful at detecting trade-offs using this type of approach.

How to infer individual heterogeneity in CR models?

In this section, we provide details about the CR models used in the case studies reviewed

above. Specifically, we focus on multistate, random-effect and finite-mixture CR models

possibly including individual covariates because these are currently the most commonly used

tools to incorporate individual heterogeneity and deal with detectability less than one. We

focus on survival and open populations in the following tutorial, but the methods are

applicable to other CR model parameters (e.g. Matechou et al. 2016), including the detection

probability, and in other contexts like closed populations. For the sake of illustration, we

simulate data in R that we analyze i) in a frequentist framework using maximum likelihood

methods with program MARK (White and Burnham 1999) called from R using the package

RMark (Laake 2013) and E-SURGE (Choquet et al. 2009) and ii) in a Bayesian framework

using Markov chain Monte Carlo methods with program JAGS (Plummer 2003) called from

R using the package R2Jags. Below we present results from the frequentist approach only.

The code to simulate data and fit CR models is available in the Supplementary materials and

from GitHub (https://github.com/oliviergimenez/indhet_in_CRmodels).

Measured **individual heterogeneity: individual covariates and multistate CR models**

Individual covariates

We start with a simple example of an individual i with the encounter history $h_i = 101$ where

'1' is for detected and '0' for non-detected. Here, individual i was detected on the first

sampling occasion, then missed and eventually detected again on the last sampling occasion.

We consider the Cormack-Jolly-Seber model for open populations and assume that both

1 survival probability ϕ *between* two sampling occasions and detection probability p *at* a

2 sampling occasion do not vary among individuals. Then, the contribution of individual i to the

3 model likelihood is:

$$\Pr(h_i) = \phi(1-p)\phi p \tag{1}$$

4   Now let us assume that we are able to measure individual heterogeneity under the

5 form of an individual covariate, say x_i, which takes a specific value for individual i (Pollock

6 2002). Then, individual variation in the survival probability (or the detection probability) can

7 be partly explained by this covariate through:

$$\text{logit}(\phi_i) = \beta_0 + \beta_1 x_i \tag{2}$$

8 where ϕ_i is the survival probability for individual $\underline{i}$, $\text{logit}(u) = \log\left(\dfrac{u}{1-u}\right)$ is the logit function

9 and is used here as a constraint to make sure that survival is estimated between 0 and 1, and

10 the β's are regression coefficients to be estimated (e.g. North & Morgan 1979). Assuming

11 now a model with individual-specific survival, Eqn (1) becomes:

$$\Pr(h_i) = \phi_i(1-p)\phi_i p \tag{3}$$

12 We do not estimate survival for each individual, but instead the regression coefficients β's in

13 Eqn (2) by first using the reciprocal logit function $\text{logit}^{-1}(v) = \dfrac{\exp(v)}{1+\exp(v)} = \dfrac{1}{1+\exp(-v)}$ in Eqn

14 (2) and plugging in the result in Eqn (3):

$$\Pr(h_i) = \frac{1}{1+\exp\left[-(\beta_0 + \beta_1 x_i)\right]}(1-p)\frac{1}{1+\exp\left[-(\beta_0 + \beta_1 x_i)\right]}p \tag{4}$$

15 The covariate x_i may be continuous like body mass or discrete like sex. If the covariate is

16 discrete, it is usually referred to as a *group* in the CR literature (Lebreton et al. 1992).

17   So far, we have assumed that this covariate does not vary over time, in other words

18 that an individual i has the same value x_i of the covariate whatever the sampling occasion (i.e.

matching with the concept of frailty *sensu stricto*). When dealing with time-varying

individual covariates, which matches the concept of dynamic frailty (for unobserved

heterogeneity), we then need to distinguish between discrete and continuous covariates.

Discrete time-varying individual covariates and multistate CR models

Discrete time-varying individual covariates are referred to as *states* in the CR literature (e.g.

breeder/non-breeder or infected/non-infected), and are analyzed with so-called multistate CR

models (Schwarz et al. 1993, Lebreton et al. 2009). Let us assume that we measure a time-

varying individual covariate with two levels, A and B, and that individual *i* has now the

encounter history $h_i = $ A0B with obvious interpretation. Two things might have happened on

the second sampling occasion at which the individual was not detected: either it stayed in

state A or it made a transition to state B. The transition event immediately calls for the

introduction of additional parameters, namely the transition probability ψ^{AB} from state A to

state B and ψ^{BA} from state B to state A. The probability of staying in state A (or B) is

obtained as the complementary probability $\psi^{AA} = 1 - \psi^{AB}$ (or $\psi^{BB} = 1 - \psi^{BA}$). The two events

'being alive in state A' and 'being alive in state B' at the second sampling occasion cannot

occur together: these are mutually exclusive. As a result, the contribution of individual *i* to the

model likelihood has two components depending on the actual underlying encounter history

AAB or ABB:

$$\Pr(h_i) = \phi_A (1 - \psi^{AB})(1 - p)\phi_A \psi^{AB} p + \phi_A \psi^{AB} (1 - p)\phi_B (1 - \psi^{BA}) p \qquad (5)$$

20   Note that *p* does not depend on state for simplicity, but this does not need to be the case. An

21   example of the use of multistate CR models to detect life-history trade-offs in the presence of

22   individual heterogeneity is provided in Table 2.

1        [Table 2 around here]

3 *Continuous time-varying individual covariates*

4 Continuous time-varying individual covariates are difficult to deal with. Ideally, we have:

$$\Pr\left(h_i\right) = \phi_{i1}\left(1-p\right)\phi_{i2}p \tag{6}$$

5 with

$$\mathrm{logit}\left(\phi_{it}\right) = \beta_0 + \beta_1 x_{it} \tag{7}$$

6 where ϕ_{it} is the survival probability for individual i between sampling occasions t and $t+1$

7 and x_{it} is the value of covariate for individual i at occasion t. However, this only corresponds

8 to the 'ideal' situation because when an individual is not detected at a particular sampling

9 occasion, then the value of the covariate is generally unknown, which makes it impossible to

10 form Eqn (7). A first possibility is to omit individuals with missing values or to replace the

11 missing values by, for example, the mean of all covariate values observed for an individual.

12 These ad-hoc approaches result in a loss of information and bias in parameter estimates and

13 should be avoided (Kendall et al. 2003, Lee et al. 2016). A more formal approach consists in

14 imputing missing covariate values from an underlying distribution that is used to model the

15 change in covariate values over time, typically a first-order Markov process like a random

16 walk (Bonner & Schwarz 2006, Langrock & King 2013; see also Worthington, King &

17 Buckland 2015). A second possibility involves the discretization of the covariate in two or

18 more levels so that multistate CR models can be used (Nichols et al. 1992). Lastly, inference

19 can be based on a conditional likelihood approach using only the observed covariate values –

20 the so-called trinomial approach (Catchpole et al. 2010). Several studies have compared the

21 statistical performances of these methods (Bonner et al. 2010, Langrock and King 2013) and

22 found that imputation methods were sensitive to the covariate model and that all methods

23 were sensitive to the detection probability and the amount of missing values.

***Unmeasured* individual heterogeneity: random-effect and finite-mixture CR models**

If for some reasons, heterogeneity cannot be measured, or there is a reason to believe that

individual covariates do not capture the relevant variation, it can yet be incorporated using

two approaches.

Random-effect CR models

The usual random-effect approach has been adapted to CR models (Coull and Agresti 1999,

Royle 2008, Gimenez and Choquet 2010). We write:

$$\text{logit}\left(\phi_i\right) = \beta_0 + \varepsilon_i \tag{8}$$

where the ε_i's are normally distributed with mean 0 and variance σ^2 to be estimated. To be

plugged in Eqn (3) using reciprocal logit function. More complex structure in the random

effects can be considered (heritability: Papaïx *et al.* 2010a; nested effects: Choquet *et al.*

2013). An example of the use of random-effect CR models to detect senescence in the

presence of individual heterogeneity is provided in Figure 1.

[Figure 1 around here]

Finite-mixture CR models

Another avenue to handle with unobserved individual heterogeneity is to use finite-mixture

models (Pledger et al. 2003, 2010, Pledger 2005, Pledger and Phillpot 2008). These models

assume that individuals can be categorized into a finite number of heterogeneity classes

(hidden states). More explicitly, an individual may be alive in class C_1 or class C_2. Then,

$$\Pr(h_i) = \pi \; \phi^{C_1}(1-p)\phi^{C_1} \; p + (1-\pi)\phi^{C_2}(1-p)\phi^{C_2} \; p \qquad (9)$$

1   where π (resp. $1-\pi$) denotes the proportion of newly marked individuals in class C_1 (resp.

2   C_2). Transition between classes can be considered (Pradel 2009; see below). An example of

3   the use of finite-mixture CR models to detect senescence in the presence of individual

4   heterogeneity is provided in Figure 2.

6   [Figure 2 around here]

Hidden-Markov modeling framework

CR models can be fruitfully expressed as state-space models in which the biological process

(survival for example) is explicitly distinguished from the observation process (detection)

(Gimenez et al. 2007, 2012, Royle 2008, King 2012). In particular, multistate CR models

incorporating uncertainty in state assignment – multievent CR models – have been formulated

as hidden-Markov models (HMM; Pradel 2005, review in Gimenez *et al.* 2012), a particular

case of state-space models in which the states are Markovian (i.e. the next state depends only

on the current state and not on the sequence of states that occurred before). An advantage of

the HMM formulation of CR models is that it provides high flexibility in the way individual

heterogeneity is modeled. For example, the HMM formulation of finite-mixture CR models

can easily be extended to consider transitions between classes of heterogeneity (Pradel 2009,

Cubaynes et al. 2010). Let us define the states alive in class 1 ('C_1'), alive in class 2 ('C_2')

and dead ('D'). The individuals can go undetected ('0') or detected ('1'). Initially, the state of

an individual is driven by the vector of initial state probabilities:

23                   $\Pi = \begin{bmatrix} \pi & 1-\pi & 0 \end{bmatrix}$

1 where the states C_1, C_2 and D are in columns in that order. Then the observation process at

2 first capture applies through:

$$B_1 = \begin{bmatrix} 0 & 1 \\ 0 & 1 \\ 1 & 0 \end{bmatrix}$$

4 where the states are in rows and the observations (or events) are in columns (0 and 1 in that

5 order). Now that the fate of individuals at first capture occasions is modeled, the survival and

6 observation processes occur successively at the subsequent occasions. The survival process is

7 governed by:

$$\Phi = \begin{bmatrix} \phi^{C_1} & 0 & 1-\phi^{C_1} \\ 0 & \phi^{C_2} & 1-\phi^{C_2} \\ 0 & 0 & 1 \end{bmatrix}$$

9 where the states at t are in rows and the states in $t+1$ are in columns. Individuals can be

10 allowed to move from a heterogeneity class to the other by multiplying the survival matrix by

11 a transition matrix:

$$\Phi = \begin{bmatrix} \phi^{C_1} & 0 & 1-\phi^{C_1} \\ 0 & \phi^{C_2} & 1-\phi^{C_2} \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1-\psi^{C_1 \rightarrow C_2} & \psi^{C_1 \rightarrow C_2} & 0 \\ \psi^{C_2 \rightarrow C_1} & 1-\psi^{C_2 \rightarrow C_1} & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

13 The observation process at occasion t is handled with:

$$B_2 = \begin{bmatrix} 1-p & p \\ 1-p & p \\ 1 & 0 \end{bmatrix}$$

15 where states are in columns and observations in columns. The likelihood in Eqn (9) can be

16 written as the product of the matrices above (Pradel 2005).

Is individual heterogeneity statistically relevant?

Testing the statistical relevance of individual heterogeneity can be done in two ways. First, the quality of fit of models with heterogeneity to CR data can be assessed using goodness-of-fit tests. An ad-hoc procedure was proposed in the context of finite-mixture models by considering specific combinations of components of the goodness-of-fit test for homogeneous models (Péron et al. 2010). A more formal approach is being developed (Jeyam, McCrea, Pradel, unpublished results) based on methods used in behavioral sciences. Second, models with and without heterogeneity can be compared using hypothesis testing or model selection. For multistate models, model selection using the Akaike Information Criterion (AIC; Burnham & Anderson 2002) is usually favored as illustrated in Table 2. In the Bayesian context, several methods have been used and we refer to review papers for guidelines (O'Hara and Sillanpää 2009, Tenan et al. 2014, Hooten et al. 2015). In random-effect models, the question boils down to testing whether the variance of the random effect is zero, which can be addressed using likelihood ratio tests (Gimenez and Choquet 2010) but may difficult to do in a model selection framework (Bolker et al. 2009). We refer to O'Hara & Sillanpää (2009) for Bayesian methods (see also Royle 2008, Chambert, Rotella & Higgs 2014). In finite-mixture models, standard tools from the model selection framework, namely the AIC, can be used (Cubaynes et al. 2012b), although it may fail in the context of detecting senescence (Supplementary Material A in Péron *et al.* 2016). In a Bayesian context, the Deviance Information Criterion (DIC; Spiegelhalter *et al.* 2014) is known to perform poorly on mixture models, and the Watanabe-Akaike information criteria (wAIC; Gelman, Hwang & Vehtari 2014) holds promise in this context, although it is yet to be used with CR models.

Discussion and research perspectives

Random effects vs. mixture? In studies based on CR models built in the context of closed or open population CR models, as well as in human demography, there is currently a debate about the distribution to consider to account for individual heterogeneity in demographic parameters (Yashin et al. 2001, Péron et al. 2010). This debate sometimes focuses on the biological justification of continuous distributions *vs.* mixture models (e.g. Péron *et al.* 2010). The debate also focuses on alternatives to distributions that might be unrealistic or inadequate (Péron *et al.* 2010). This question is not specific to CR modeling (Hamel et al. 2016). Clusters of individuals sharing values of latent traits can be identified using mixture models. Recently, (Hamel et al. 2016) addressed the question of the identification of reproductive and growth tactics in long-lived mammals using mixture models. They also used simulations and showed that in many cases the number of clusters can be chosen using an information theoretic or a bootstrap approach. Alternatively, infinite mixture models could be developed for CR data (Rasmussen 2000, Ohlssen et al. 2007, Raman et al. 2010), where the number of clusters is *a priori* very large, but the number of clusters including at least one individual is estimated and can range from 1 to a large number; the latter situation leads a distribution of demographic parameters that approaches a continuous one (Ohlssen et al. 2007). Rather than comparing models that vary in complexity using for instance an information criterion, Bayesian nonparametric approaches "fit a single model that can adapt its complexity to the data (Gershman & Blei 2012; see Ford, Patterson & Bravington 2015 for an application to CR models). Moreover, the question of how to account for heterogeneous detection probability in CR models designed to estimate population size has a very long history (e.g. Carothers 1973, Link 2003, Ghosh & Norris 2005). Carothers (1973) investigated the consequences of violations of the assumption of equal detection probability on estimates from the Jolly-Seber

model. He concluded that "any distribution is, from the point of view of investigating bias, as

good as any other with the same [mean detection probability] and [coefficient of variation],

and it is therefore justifiable to select a distribution on the grounds of computational

convenience alone". However, the number of classes can be itself of interest, perhaps to

identify clusters of individuals with different behavior (e.g. Marescot *et al.* 2011). In the

framework of closed populations though, there is no straightforward means of determining the

number of components of a mixture model for detection probability (Link 2003).

Change in latent values of demographic parameters over lifetime. In standard models for

longitudinal data with individual heterogeneity, an independent subject-specific random effect

is assumed to be constant over time for each subject (Vaupel & Missov 2014), which matches

early version of the concept of frailty (Vaupel et al. 1979). Generally, in CR studies using

mixture models, individuals are also assumed to be a member of a latent class when they enter

the study, and they do not change class. Mortality risk or breeding success propensity at time

0 (when the individual enters the study) is assumed to be perfectly correlated with the risk

later in life (Wienke 2010). However, this assumption does not necessarily hold, and models

accommodating changes in individual latent vital rate values may offer an interesting basis to

test biological hypotheses. An alternative approach allowing individuals to experience

(reversible) changes in latent vital rates could be based on the ontogenetic view of individual

differences (Senner et al. 2015). This can be achieved with 'dynamic frailty' models (Manda

and Meyer 2005, Putter and Van Houwelingen 2014), hidden Markov models, Latent Class

transition models or mixture models, in which individuals are allowed to change latent class

over time (e.g. Kaplan 2008). Hidden Markov models are now commonly used in CR studies,

but specific applications to change in latent demographic parameters are still rare (Pradel

2005, Cubaynes *et al.* 2010).

Initial conditions. An overlooked issue in CR studies using multistate models is the 'initial

conditions' one. Before estimating the parameters of a model accounting for a stochastic

process with dependence among consecutive states (e.g., breeding states), one has to think

about how the process was "initialized". Studies of reproduction necessarily start recording

breeding outcomes at the first breeding event (recruitment, or first observed breeding

attempt). More generally, studies modeling reproductive outcomes from recruitment onwards

(e.g., Cam et al. 1998, Yoccoz et al. 2002) assume that the start of the process generating the

observed reproductive states coincides with the start of reproductive life for each individual

(Wooldridge 2005). Nevertheless, the process governing the first breeding outcome can be the

same as the process generating the subsequent observations in the individual lifetime

trajectory (Skrondal and Rabe-hesketh 2014). Such a process can include unobserved

determinants of breeding outcome. In dynamic models of reproduction incorporating the

effect of past breeding outcome at time t on the probability of breeding successfully at time

$t+1$ (e.g. multistate CR models), the outcome of the first reproductive attempt (at time t) is not

considered as the realization of a random process, because there is no reproduction at time $t-1$.

Nevertheless, failure to incorporate unobserved factors governing breeding success

probability at recruitment can translate into overestimation of transition probabilities among

subsequent reproductive states (Heckman 1981, Prowse 2012). This is particularly

problematic in studies of changes in reproductive costs throughout the lifetime (e.g. Sanz-

Aguilar *et al.* 2008), or of experience-specific variation in breeding outcome (e.g. Nevoux,

Weimerskirch & Barbraud 2007). Interestingly, Sanz-Aguilar et al. (2008) have interpreted

evidence of higher reproductive costs of reproduction at recruitment as a consequence of

within-cohort mortality selection, with frailer individuals incurring higher reproductive costs

than robust ones. The initial conditions problem can be overcome using a joint modeling

approach of the processes governing reproductive success at recruitment and subsequent

breeding occasions (Skrondal and Rabe-hesketh 2014). If CR data are available from the pre-

breeding period, then unobserved and observed determinants of breeding state can be

considered simultaneously (e.g. Fay et al. 2016b).

Inference about individual heterogeneity. A series of recent papers have revived interest in

unobserved heterogeneity in demographic parameters in CR studies (Steiner et al. 2010,

Orzack et al. 2011). More specifically, these papers have drawn attention to the approaches

used to discriminate among hypotheses about sources of variation in CR histories. In CR

studies, an influential book by Burnham & Anderson (2002) has promoted the use of

multimodel inference such as information criteria to address non-mutually exclusive

biological hypotheses about the processes governing mortality, or the arrangement of

reproductive states over lifetime trajectories of animals. For example, models accounting for

state-dependence in survival or reproduction can be considered (multistate or multievent

models; e.g., Sanz-Aguilar *et al.* 2008), models accounting for unobserved heterogeneity in

these demographic parameters too (e.g. Royle 2008, Marzolin et al. 2011), as well as models

accounting for both sources of variation in survival and reproduction (e.g. Fay et al. 2016a).

This contrasts with approaches based on a single model (namely, state-dependence) and

evaluation of the degree of consistency of observed individual CR histories with metrics

summarizing key features of histories simulated using parameter estimated with the model in

question (Steiner et al. 2010, Orzack et al. 2011).

By definition, variation in individual trajectories simulated using parameters estimated

with a multistate CR models is not caused by fixed, unobserved heterogeneity among

individual demographic parameters (Tuljapurkar et al. 2009, Steiner and Tuljapurkar 2012).

The variation in arrangements of states in simulated data stems from the realization of random

variables governed by probabilities; the resulting pattern is called 'dynamic heterogeneity'

(Tuljapurkar et al. 2009), or 'individual stochasticity' (Caswell 2009). Several papers have

provided evidence that there is a good match between observed and simulated features of

individual histories (Steiner et al. 2010, Orzack et al. 2011, Steiner and Tuljapurkar 2012).

These studies suggest that stochastic demographic processes have been overlooked in life

history studies, and that latent, unobserved heterogeneity in demographic parameters might

have been overstated in studies of longitudinal data from animals, whether detection

probability is lower than one or not (Cam et al. 2002a, 2013, Steiner et al. 2010, Orzack et al.

2011). From a conceptual viewpoint, these studies attempt to caution biologists against the

temptation of adaptationism when interpreting amounts of unobserved individual

heterogeneity in demographic parameters ("biologists commonly argue that large differences

in fitness components are likely adaptive, resulting from and driving evolution by natural

selection" Steiner & Tuljapurkar 2012, Cam, Aubry & Authier 2016). However, they have

moved away from one of the dominating statistical inference approaches in the CR area,

namely multimodel inference and information criteria (Burnham and Anderson 2002).

Current research is addressing the question of whether simulations based on multistate CR

models or simply models with state-dependence used for longitudinal data analysis allow

discriminating among alternative hypotheses about the processes generating variability in

individual histories (Plard et al. 2012, Bonnet and Postma 2016, Cam et al. 2016).

Importantly, they have overlooked notes of caution from other areas of research also

using multistate models for inferences about longitudinal data concerning possible biases in

estimates of 'state-dependence' (e.g. Heckman 1981, Ahmad 2014). A key issue in

discriminating among processes generating variation in individual histories is that a Markov

process (i.e. the basis of multistate models) and unobserved individual heterogeneity (for

instance a random effect model, Royle 2008) can create similar patterns in arrangements of

states along individual trajectories (Ahmad 2014, Authier et al. 2017). This issue has

stimulated a large body of work in econometrics (Heckman 1981, Ahmad 2014, Skrondal and

Rabe-hesketh 2014, Andriopoulou and Tsakloglou 2015). The hypothesis of a

'communicating vessels' phenomenon between sources of variation in CR histories should be

considered in wild animal populations, like in econometrics studies (Ahmad 2014, Plum and

Ayllón 2015, Cam et al. 2016). Interestingly, several CR studies have hypothesized that their

results obtained using multistate models partly reflect phenotypic within-cohort mortality

selection (i.e., the change in the composition of a heterogeneous cohort including individuals

with different baseline survival probabilities; Cam *et al.* 2002a, Barbraud & Weimerskirch

2004, Nevoux, Weimerskirch & Barbraud 2007, Sanz-Aguilar *et al.* 2008). That is, they have

hypothesized that their result may be caused by unobserved individual heterogeneity, a

question now being addressed in studies of senescence (Péron et al. 2010, 2016). This

suggests that CR models with both a Markovian structure (for observable, partially

observable, or unobservable states) and unobserved individual heterogeneity might perform

well with some datasets from wild animal populations (Fay et al. 2016a, b).

Conclusion

Our review, although not exhaustive, demonstrates that the tremendous advances in CR

modeling accomplished over the past 40 years provide investigators with a reliable way to

address multiple facets of the process of individual heterogeneity in demographic parameters.

Pioneer works by quantitative wildlife biologists focused on individual heterogeneity in

recapture or survival probability to avoid biased estimates of population size. Afterwards, the

emergence of more general questions such as cause-specific sources of mortality in game- and

non-game species (Koons et al. 2014) and the need for accurate assessment of the impact of

global change on the demography of structured populations (Gullett et al. 2014) have

triggered collaborations between biologists and statisticians to make efficient use of data,

robust inferences about demographic parameters, and achieve an increasing degree of realism

in both the sampling and ecological processes handled by CR models. As emphasized by

Conroy (2009), the nature of questions that can be addressed nowadays has been greatly

expanded to include evolutionary ecology, whose cornerstone is variation in demographic

parameters among individuals both within and among populations. The relevance of dealing

with individual heterogeneity to study eco-evolutionary processes has placed the topic of

individual heterogeneity at the core of many empirical investigations using CR data (Table 1).

Provided appropriate sampling design and sufficient data are available, the flexibility of

modern CR models now allows assessing reliably the role of individual heterogeneity in

ecology and evolutionary processes in the wild.

Acknowledgements –EC was supported by the Laboratoire d'Excellence LabEx TULIP

(ANR-10-LABX-41). OG was supported by ANR-16-CE02-0007.

References

Abadi, F. et al. 2013. Revisiting the Effect of Capture Heterogeneity on Survival Estimates in

Capture-Mark-Recapture Studies: Does It Matter? - PLoS One 8: e62636.

Abbring, J. H. and Van Den Berg, G. J. 2007. The unobserved heterogeneity distribution in

duration analysis. - Biometrika 94: 87–99.

Ahmad, N. 2014. State dependence in unemployment. - Int. J. Econ. Financ. Issues 4: 93.

Allison, P. D. 1982. Discrete-time methods for the analysis of event histories. - Sociol.

Methodol. 13: 61–98.

Andriopoulou, E. and Tsakloglou, P. 2015. Once poor, always poor? Do initial conditions

matter? Evidence from the ECHP. - In: Measurement of Poverty, Deprivation, and

Economic Mobility. Emerald Group Publishing Limited, pp. 23–70.

Arnason, A. N. 1972. Parameter estimates from mark-recapture experiments on two populations subject to migration and death. - Res. Popul. Ecol. 13: 97–113.

Arnason, A. N. 1973. The estimation of population size, migration rates and survival in a stratified population. - Res. Popul. Ecol. 15: 1–8.

Aubry, L. M. et al. 2011. Drivers of age-specific survival in a long-lived seabird: Contributions of observed and hidden sources of heterogeneity. - J. Anim. Ecol. 80: 375–383.

Authier, M. et al. 2017. Wolf in sheep's clothing: Model misspecification undermines tests of the neutral theory for life histories. - Ecol. Evol.: In press.

Barbraud, C. and Weimerskirch, H. 2004. Modelling the effects of environmental and individual variability when measuring the costs of first reproduction. - Anim. Biodivers. Conserv. 27: 109–111.

Barbraud, C. et al. 2005. Environmental conditions and breeding experience affect costs of reproduction in blue petrels. – Ecology 86: 682-692.

Barbraud, C. et al. 2013. Fisheries Bycatch as an Inadvertent Human-Induced Evolutionary Mechanism. - Plos One 8: e60353.

Baudisch, A. and Vaupel, J. W. 2012. Evolution. Getting to the root of aging. - Science 338: 618–9.

Beauplet, G. et al. 2006. Age-specific survival and reproductive performances in fur seals: evidence of senescence and individual quality. – Oikos 112: 430-441.

Berger, V. et al. 2015. Early and adult social environments have independent effects on individual fitness in a social vertebrate. - Proc. Biol. Sci. 282: 20151167.

Blomberg, E. J. et al. 2013. Seasonal reproductive costs contribute to reduced survival of female greater sage-grouse. - J. Avian Biol. 44: 149-158.

Blums, P. et al. 2005. Individual quality, survival variation and patterns of phenotypic

    selection on body condition and timing of nesting in birds. - Oecologia 143: 365-376.

Bolker, B. M. et al. 2009. Generalized linear mixed models: a practical guide for ecology and

    evolution. - Trends Ecol. Evol. 24: 127–135.

Bolnick, D. I. et al. 2011. Why intraspecific trait variation matters in community ecology. -

    Trends Ecol. Evol. 26: 183–192.

Bonenfant, C. et al. 2009. Age-dependent relationship between horn growth and survival in

    wild sheep. – J. Anim. Ecol. 78: 161-171.

Bonner, S. J. and Schwarz, C. J. 2006. An extension of the Cormack-Jolly-Seber model for

    continuous covariates with application to Microtus pennsylvanicus. - Biometrics 62:

    142–149.

Bonner, S. J. et al. 2010. Continuous covariates in mark-recapture-recovery analysis: a

    comparison of methods. - Biometrics 66: 1256–1265.

Bonnet, T. and Postma, E. 2016. Successful by Chance? The Power of Mixed Models and

    Neutral Simulations for the Detection of Individual Fixed Heterogeneity in Fitness

    Components. - Am. Nat. 187: 60–74.

Borchers, D. 2012. A non-technical overview of spatially explicit capture-recapture models. -

    J. Ornithol. 152: 435–444.

Bouwhuis, S. et al. 2012. The forms and fitness cost of senescence: Age-specific recapture,

    survival, reproduction, and seproductive value in a wild bird population. - Am. Nat. 179:

    E15–E27.

Briggs, C. W. et al. 2011. Correlates of survival in Swainson's hawks breeding in Northern

    California. - J. Wildl. Manage. 75: 1307-1314.

Buoro, M. et al. 2010. Investigating evolutionary trade-offs in wild populations of atlantic

    salmon (salmo salar): incorporating detection probabilities and individual heterogeneity.

- Evolution. 64: 2629–42.

Buoro, M. et al. 2011. Assessing adaptive phenotypic plasticity by means of conditional

strategies from empirical data: The latent environmental threshold model. - Evolution.

66: 996–1009.

Burnham, K. P. and Overton, W. S. 1979. Robust estimation of population size when capture

probabilities vary among animals. - Ecology 60: 927–936.

Burnham, K. P. and Rexstad, E. A. 1993. Modeling heterogeneity in survival rates of banded

waterfowl. - Biometrics 49: 1194–1208.

Burnham, K. P. and Anderson, D. R. 2002. Model selection and multimodel inference: A

practical information-theoretic approach. - Springer.

Cam, E. et al. 1998. Are adult nonbreeders prudent parents? The Kittiwake model. - Ecology

79: 2917–2930.

Cam, E. and Monnat, J. Y. 2000a. Stratification based on reproductive state reveals

contrasting patterns of age-related variation in demographic parameters in the kittiwake.

– Oikos 90: 560-574.

Cam, E. and Monnat, J. Y. 2000b. Apparent inferiority of first-time breeders in the kittiwake:

the role of heterogeneity among age classes. – J. Anim. Ecol. 69: 380-394.

Cam, E. et al. 2002a. Individual covariation in life-history traits: Seeing the trees despite the

forest. – Am. Nat. 159: 96-105.

Cam, E. et al. 2002b. Influence of behavioural tactics on recruitment and reproductive

trajectory in the kittiwake. – L. Appl. Stat. 29: 163-185.

Cam, E. et al. 2003. Long-term fitness consequences of early conditions in the kittiwake. – J.

Anim. Ecol. 72: 411-424.

Cam, E. et al. 2013. Looking for a needle in a haystack: inference about individual fitness

components in a heterogeneous population. - Oikos 122: 739–753.

Cam, E. et al. 2016. The Conundrum of Heterogeneities in Life History Studies. - Trends
    Ecol. Evol. 31: 872–886.

Carothers, A. D. 1973. The Effects of Unequal Catchability on Jolly-Seber Estimates. -
    Biometrics 29: 79–100.

Caswell, H. 2009. Stage, age and individual stochasticity in demography. - Oikos 118: 1763–
    1782.

Catchpole, E. A. et al. 2010. A new method for analysing discrete life history data with
    missing covariate values. - J. R. Stat. Soc. - Ser. B Stat. Methodol. 70: 445–460.

Caughley, G. 1966. Mortality patterns in mammals. - Ecology 47: 906–918.

Chambert, T. et al. 2013. Individual heterogeneity in reproductive rates and cost of
    reproduction in a long-lived vertebrate. - Ecol. Evol. 3: 2047–2060.

Chambert, T. et al. 2014. Use of posterior predictive checks as an inferential tool for
    investigating individual heterogeneity in animal population vital rates. - Ecol. Evol. 4:
    1389–1397.

Chambert, T. et al. 2015. Female Weddell seals show flexible strategies of colony attendance
    related to varying environmental conditions. – Ecology 96: 479-488.

Choquet, R. et al. 2009. Program E - SURGE : A Software Application for Fitting Multievent
    Models. - In: Thomson, D. L. et al. (eds), Environmental and Ecological Statistics.
    Springer US, pp. 845–865.

Choquet, R. et al. 2013. Estimating demographic parameters from capture-recapture data with
    dependence among individuals within clusters. - Methods Ecol. Evol. 4: 474–482.

Clutton-Brock, T. H. 1988. Reproductive Success. Studies of Individual Variation in
    Contrasting Breeding Systems. - University of Chicago Press.

Clutton-Brock, T. and Sheldon, B. C. 2010. Individuals and populations: the role of long-
    term, individual-based studies of animals in ecology and evolutionary biology. - Trends

Ecol. Evol. 25: 562–573.

Cody, M. L. 1966. A general theory of clutch size. - Evolution. 20: 174–184.

Cohen, J. 1986. An uncertainty principle in demography and the unisex issue. - Am. Stat. 40:

32–39.

Conner, M. M. and White, G. C. 1999. Effects of individual heterogeneity in estimating the

persistence of small populations. - Nat. Resour. Model. 12: 109–127.

Conroy, M. J. 2009. Application of capture–recapture to addressing questions in evolutionary

ecology. In Thomson et al. (eds) *Modeling demographic processes in marked*

populations, Environmental and Ecological Statistics (pp. 131-152). Springer US.

Cooch, E. G. et al. 2002. Occam's shadow: levels of analysis in evolutionary ecology - where

to next? - J. Appl. Stat. 29: 19–48.

Cormack, R. M. 1964. Estimates of survival from the sighting of marked animals. -

Biometrika 51: 429–438.

Coull, B. and Agresti, A. 1999. The use of mixed logit models to reflect heterogeneity in

capture–recapture studies. - Biometrics 55: 294–301.

Coulson, T. et al. 2011. Modeling Effects of Environmental change on wolf population

dynamics, trait evolution, and life history. - Science. 334: 1275–1278.

Crespin, L. et al. 2008. Is heterogeneity of catchability in capture-recapture studies a mere

sampling artifact or a biologically relevant feature of the population? - Popul. Ecol. 50:

247–256.

Cubaynes, S. et al. 2010. Importance of Accounting for Detection Heterogeneity When

Estimating Abundance: the Case of French Wolves. - Conserv. Biol. 24: 621–626.

Cubaynes, S. et al. 2012a. Testing hypotheses in evolutionary ecology with imperfect

detection: Structural equation modeling of mark-recapture data. - Ecology in press.

Cubaynes, S. et al. 2012b. Assessing individual heterogeneity using model selection criteria:

how many mixture components in capture-recapture models? - Methods Ecol. Evol. 3: 564–573.

Descamps, S. et al. 2009. Costs of reproduction in a long-lived bird: large clutch size is associated with low survival in the presence of a highly virulent disease. - Biol. Lett. 5: 278–281.

Desprez, M. et al. 2013. Known unknowns in an imperfect world: incorporating uncertainty in recruitment estimates using multi-event capture-recapture models. - Ecol. Evol. 3: 4658–4668.

Dingemanse, N. J. and Dochtermann, N. A. 2013. Quantifying individual variation in behaviour: Mixed-effect modelling approaches. - J. Anim. Ecol. 82: 39–54.

Dobzhansky, T. 1973. Nothing in Biology Makes Sense except in the Light of Evolution. - Am. Biol. Teach. 35: 125–129.

Douhard, M. et al. 2014. Fitness consequences of environmental conditions at different life stages in a long-lived vertebrate. - Proc. R. Soc. B 281: 20140276.

Drummond, H. et al. 2011. Natural "poor start" does not increase mortality over the lifetime. - Proc. R. Soc. B Biol. Sci. 278: 3421–3427.

Duffie, D. et al. 2009. Frailty correlated default. - J. Finance 64: 2089–2123.

Eberhardt, L. L. 1969. Population Estimates from Recapture Frequencies. - J. Wildl. Manage. 33: 28–39.

Emlen, J. M. 1970. Age Specificity and Ecological Theory. - Ecology 51: 588.

Enki, D. G. et al. 2014. A time-varying shared frailty model with application to infectious diseases. - Ann. Appl. Stat. 8: 430–447.

Estes, J. A. et al. 2003. Individual variation in prey selection by sea otters: Patterns, causes and implications. - J. Anim. Ecol. 72: 144–155.

Fay, R. et al. 2016a. Variation in the age of first reproduction: different strategies or

individual quality? - Ecology 97: 1842–1851.

Fay, R. et al. 2016b. Paternal but not maternal age influences early-life performance of

offspring in a long-lived seabird. - Proc. R. Soc. B 283: 20152318.

Festa-Bianchet, M. et al. 1998. Mass- and density-dependent reproductive success and

reproductive costs in a capital breeder. - Am. Nat. 152: 367–379.

Fletcher, D. et al. 2012. Bias in estimation of adult survival and asymptotic population growth

rate caused by undetected capture heterogeneity. - Methods Ecol. Evol. 3: 206–216.

Ford, J. H. et al. 2015. Modelling latent individual heterogeneity in mark-recapture data with

Dirichlet process priors. arXiv:1511.07103v1. https://arxiv.org/abs/1511.07103

Fox, G. A. et al. 2006. Consequences of heterogeneity in survival probability in a population

of Florida scrub-jays. - J. Anim. Ecol. 75: 921–927.

Garnier, A. et al. 2016. What shapes fitness costs of reproduction in long-lived iteroparous

species? A case study on the Alpine ibex. - Ecology 97: 205-214.

Gelman, A. et al. 2014. Understanding predictive information criteria for Bayesian models. -

Stat. Comput. 24: 997–1016.

Gershman, S. J. and Blei, D. M. 2012. A tutorial on Bayesian nonparametric models. - J.

Math. Psychol. 56: 1–12.

Ghosh, S. K. and Norris, J. L. 2005. Bayesian capture-recapture analysis and model selection

allowing for heterogeneity and behavioral effects. - J. Agric. Biol. Environ. Stat. 10: 35–

49.

Gimenez, O. and Choquet, R. 2010. Individual heterogeneity in studies on marked animals

using numerical integration: capture-recapture mixed models. - Ecology 91: 951–957.

Gimenez, O. et al. 2007. State-space modelling of data on marked individuals. - Ecol. Modell.

206: 431–438.

Gimenez, O. et al. 2008. The risk of flawed inference in evolutionary studies when

detectability is less than one. - Am. Nat. 172: 441–8.

Gimenez, O. et al. 2009. Estimating and visualizing fitness surfaces using mark-recapture data. - Evolution. 63: 3097–105.

Gimenez, O. et al. 2012. Estimating demographic parameters using hidden process dynamic models. - Theor. Popul. Biol. 82: 307–16.

Green, R. F. 1977. Do more birds produce fewer young? A comment on Mayfield's measure of nest success. - Wilson Bull. 89: 173–175.

Guéry, L. et al. 2017. Hidden survival heterogeneity of three common eider populations in response to climate fluctuations. - J. Anim. Ecol. (in press).

Gullett, P., Evans, K. L., Robinson, R. A., and Hatchwell, B. J. 2014. Climate change and annual survival in a temperate passerine: partitioning seasonal effects and predicting future patterns. - Oikos 123: 389-400.

Hadley, G. L. et al. 2007. Evaluation of reproductive costs for Weddell seals in Erebus Bay, Antarctica. - J. Anim. Ecol. 76: 448–458.

Hamel, S. et al. 2009. Individual variation in reproductive costs of reproduction: High-quality females always do better. - J. Anim. Ecol. 78: 143–151.

Hamel, S. et al. 2014. A standardized approach to estimate life history tradeoffs in evolutionary ecology. - Oikos 123: 151–160.

Hamel, S. et al. 2016. Cohort variation in individual body mass dissipates with age in large herbivores. - Ecol. Monogr. 86: 517–543.

Hartson, R. B. et al. 2015. Competitive release modifies the impacts of hydrologic alteration for a partially migratory stream predator. - Ecol. Fresh. Fish. 24: 276-292.

Heckman, J. J. 1981. The Incidental Parameters Problem and the Problem of Initial Conditions in Estimating: A Discrete Time-discrete Data Stochastic Process and Some Monte Carlo Evidence. - Graduate School of Business and Department of Economics,

University of Chicago, 1979.

Hernandez-Matias, A. et al. 2011. Effects of age, territoriality and breeding on survival of

Bonelli's Eagle *Aquila fasciata*. – Ibis 153: 846-857.

Hestbeck, J. B. et al. 1991. Estimates of movement and site fidelity using mark-resight data of

wintering canada geese. - Ecology 72: 523–533.

Hileman, E. T. et al. 2015. Recouping lost information when mark-recapture data are pooled:

A case study of milksnakes (*Lampropeltis triangulum*) in the Upper Midwestern United

States. - J. Herpetol. 49: 428-436.

Hooten, M. B. et al. 2015. A guide to Bayesian model selection for ecologists. - Ecol.

Monogr. 85: 3–28.

Horswill, C. et al. 2014. Survival in macaroni penguins and the relative importance of

different drivers: individual traits, predation pressure and environmental variability. - J.

Anim. Ecol. 83: 1057-1067.

Hougaard, P. 1984. Life table methods for heterogeneous populations: Distributions

describing the heterogeneity. - Biometrika 71: 75–83.

Hougaard, P. 1991. Modelling Heterogeneity in Survival Data. - J. Appl. Probab. 28: 695–

701.

Hougaard, P. 1995. Frailty models for survival data. - Lifetime Data Anal. 1: 255–273.

Hua, D. et al. 2015. Use of PIT tags to assess individual heterogeneity of laboratory-reared

juveniles of the endangered Cumberlandian combshell (*Epioblasma brevidens*) in a

mark-recapture study. - Ecol. Evol. 5: 1076-1087.

Johnson, D. H. 1979. Estimating Nest Success: The Mayfield Method and an Alternative. -

Auk 96: 651–661.

Johnson, D. H. et al. 1986. The role of heterogeneity in animal population dynamics. - Proc.

Thirteen. Int. Biometrics Conf. Sess. 5, Seattle, June 1985

Johnson, D. S. et al. 2016. Multivariate State Hidden Markov Models for Mark-Recapture

Data. - Stat. Sci. 31: 233–244.

Jolly, G. M. 1965. Explicit estimates from capture-recapture data with both death and

immigration-stochastic model. - Biometrika 52: 225–247.

Kannisto, V. 1991. Frailty and survival. - Genus 47: 101–118.

Kaplan, D. 2008. An overview of Markov chain methods for study of stage-sequential

developmental processes. - Dev. Psychol. 44: 457–467.

Kendall, B. E. and Fox, G. A. 2002. Variation among individuals and reduced demographic

stochasticity. - Conserv. Biol. 16: 109–116.

Kendall, W. L. et al. 2003. Adjusting multistate capture–recapture models for

misclassification bias: manatee breeding proportions. - Ecology 84: 1058–1066.

Kendall, B. E. et al. 2011. Demographic heterogeneity, cohort selection, and population

growth. - Ecology 92: 1985–1993.

Kennamer, R. A. et al. 2016. Effects of current reproductive success and individual

heterogeneity on survival and future reproductive success of female Wood Ducks. – Auk

133: 439-450.

Keyfitz, N. and Littman, G. 1979. Mortality in a heterogeneous population. - Popul. Stud. 33:

333–342.

King, R. 2012. A Review of Bayesian State-Space Modelling of Capture-Recapture-Recovery

Data. - Interface Focus 2: 190–204.

Knape, J. et al. 2011. Individual heterogeneity and senescence in Silvereyes on Heron Island.

- Ecology 92: 813–820.

Koenig, W. D. et al. 1995. Patterns and consequences of egg destruction among joint-nesting

acorn woodpeckers. - Anim. Behav. 50: 607–621.

Koons, D. N. et al. 2014. Methods for studying cause-specific senescence in the wild. - Meth.

Ecol. Evol. 5: 924-933.

Kovach, R. P. et al. 2010. Strong influence of microhabitat on survival for an intertidal snail,

Nucella lima. – Hydrobiologia 652: 49-56.

Kramer, G. H. 1983. The Ecological Fallacy Revisited: Aggregate- versus Individual-level

Findings on Economics and Elections, and Sociotropic Voting. - Am. Polit. Sci. Rev. 77:

92–111.

Laake, J. L. 2013. RMark: An R interface for analysis of capture-recapture data with MARK.

- AFSC Process. Rep. 2013-01 2013–1: 25.

Langrock, R. and King, R. 2013. Maximum likelihood estimation of mark–recapture–

recovery models in the presence of continuous covariates. 7: 1709–1732.

Le Bohec, C. et al. 2007. Population dynamics in a long-lived seabird: I. Impact of breeding

activity on survival and breeding probability in unbanded king penguins. – J. Anim.

Ecol. 76: 1149-1160.

Lebreton, J.-D. and Pradel, R. 2002. Multistate recapture models: Modelling incomplete

individual histories. - J. Appl. Stat. 29: 353–369.

Lebreton, J.-D. et al. 1992. Modeling survival and testing biological hypotheses using marked

animals: a unified approach with case studies. - Ecol. Monogr. 62: 67–118.

Lebreton, J.-D. et al. 2009. Modeling Individual Animal Histories with Multistate Capture-

Recapture Models. - Adv. Ecol. Res. 41: 87–173.

Lee, D. E. 2011. Effects of environmental variability and breeding experience on northern

elephant seal demography. – J. Mamm. 92: 517-526.

Lee, S. et al. 2016. Estimation in Closed Capture – Recapture Models When Covariates Are

Missing at Random. - Biometrics 72: 1294–1304

Lemaître, J.-F. et al. 2015. Early-late life trade-offs and the evolution of ageing in the wild. -

Proc. Biol. Sci. 282: 20150209.

Lescroël, A. et al. 2009. Effects of individual quality, reproductive success and environmental

variability on survival of a long-lived seabird. - J. Anim. Ecol. 78: 798-806.

Lindberg, M. S. et al. 2013. Individual heterogeneity in black brant survival and recruitment

with implications for harvest dynamics. – Ecol. Evol. 3: 4045-4056.

Link, W. A. 2003. Nonidentifiability of population size from capture-recapture data with

heterogeneous detection probabilities. - Biometrics 59: 1123–1130.

Link, W. A. 2004. Individual heterogeneity and identifiability in capture-recapture models. -

Anim. Biodivers. Conserv. 27: 87–91.

Link, W. A. and Hesed, K. M. 2015. Individual heterogeneity in growth and age at sexual

maturity: A Gamma process analysis of Capture-Mark-Recapture data. - J. Agric. Biol.

Environ. Stat. 20: 343-352.

Loison, A. et al. 1999. Age-specific survival in five populations of ungulates: Evidence of

senescence. - Ecology 80: 2539–2554.

Lomnicki, A. 1978. Individual differences between animals and the natural regulation of their

numbers. - J. Anim. Ecol. 47: 461–475.

Lynch, M. and Walsh, B. 1998. Genetics and analysis of quantitative traits.

Manda, S. O. M. and Meyer, R. 2005. Bayesian inference for recurrent events data using

time-dependent frailty. - Stat. Med. 24: 1263–1274.

Maniscalco, J. M. et al. 2010. High natality rates of endangered Steller sea lions in Kenai

fjords, Alaska and perceptions of population status in the Gulf of Alaska. - Plos One 5:

e10076.

Manly, B. F. J. and Parr, M. J. 1968. A new method for estimating population size,

survivorship, and birth rate from capture-recapture data. - Trans. Soc. Br. Entomol. 18:

81–89.

Manton, K. G. et al. 1981. Methods for comparing the mortality experience of heterogeneous

populations. - Demography 18: 389–410.

Manton, K. G. et al. 1986. Alternative models for the heterogeneity of mortality risks among the aged. - J. Am. Stat. Assoc. 81: 635–644.

Marescot, L. et al. 2011. Capture – recapture population growth rate as a robust tool against detection heterogeneity for population management. - Ecol. Appl. 21: 2898–2907.

Marzolin, G. et al. 2011. Frailty in state-space models: application to actuarial senescence in the Dipper. - Ecology 92: 562–567.

Matechou, E. et al. 2016. Bayesian analysis of Jolly-Seber type models Incorporating heterogeneity in arrival and departure. - Environ. Ecol. Stat. 23: 531–547.

Mcloughlin, P. D. et al. 2007. Lifetime reproductive success and composition of the home range in a large herbivore. - Ecology 88: 3192–3201.

McNamara, J. M. and Houston, A. I. 1996. State-dependent life histories. - Nature 380: 215–221.

Millon, A. et al. 2010. Pulsed resources affect the timing of first breeding and lifetime reproductive success of tawny owls. - J. Anim. Ecol. 79: 426-435.

Morano, S. et al. 2013. Life-history strategies of North American elk: trade-offs associated with reproduction and survival. - J. Mamm. 94: 162-172.

Morgan, B. J. T. and Ridout, M. S. 2008. A new mixture model for capture heterogeneity. - J. R. Stat. Soc. Ser. C Appl. Stat. 57: 433–446.

Moyes, K. et al. 2011. Individual differences in reproductive costs examined using multi-state methods. - J. Anim. Ecol. 80: 456-465.

Nevoux, M. et al. 2007. Environmental variation and experience-related differences in the demography of the long-lived black-browed albatross. - J. Anim. Ecol. 76: 159–167.

Newton, I. 1989. Lifetime reproduction in birds. - Academic Press.

Nichols, J. D. 2002. Discussion comments on: "Occam' s shadow: levels of analysis in

evolutionary ecology – where to next?" by Cooch, Cam and Link. - J. Appl. Stat. in

press.

Nichols, J. D. and Kendall, W. L. 1995. The use of multi-state capture-recapture models to

address questions in evolutionary ecology. - J. Appl. Stat. 22: 835–846.

Nichols, J. D. et al. 1992. Estimating transition probabilities for stage-based population

projection matrices using capture–recapture data. - Ecology 73: 306–312.

Nichols, J. D. et al. 1994. Estimating breeding proportions and testing hypotheses about costs

of reproduction with capture-recapture data. - Ecology 75: 2052–2065.

Nicolai, C. A. and Sedinger, J. S. 2012. Trade-offs between offspring fitness and future

reproduction of adult female black brent. - J. Anim. Ecol. 81: 798–805.

Norris, J. L. and Pollock, K. H. 1996. Nonparametric MLE under two closed capture

recapture models with heterogeneity. - Biometrics 52: 639–649.

North, P. M. and Morgan, B. J. T. 1979. Modeling heron survival using weather data. -

Biometrics 35: 667–681.

Nussey, D. H. et al. 2008. Measuring senescence in wild animal populations: Towards a

longitudinal approach. - Funct. Ecol. 22: 393–406.

Nussey, D. H. et al. 2013. Senescence in natural populations of animals: Widespread evidence

and its implications for bio-gerontology. - Ageing Res. Rev. 12: 214–225.

O'Hara, R. B. and Sillanpää, M. J. 2009. A review of bayesian variable selection methods:

What, how and which. - Bayesian Anal. 4: 85–118.

Ohlssen, D. I. et al. 2007. Flexible random-effects models using Bayesian semi-parametric

models: Applications to institutional comparisons. - Stat. Med. 26: 2088–2112.

Oliver, L. J. et al. 2011. Individual heterogeneity in recapture probability and survival

estimates in cheetah. - Ecol. Modell. 222: 776–784.

Orzack, S. H. et al. 2011. Static and dynamic expression of life history traits in the northern

fulmar Fulmarus glacialis. - Oikos 120: 369–380.

Otis, D. L. et al. 1978. Statistical Inference from Capture Data on Closed Animal Populations. - Wildl. Monogr. 62: 3–135.

Papaïx, J. et al. 2010. Combining capture-recapture data and pedigree information to assess heritability of demographic parameters in the wild. - J. Evol. Biol. 23: 2176–2184.

Pennell, M. L. and Dunson, D. B. 2006. Bayesian semiparametric dynamic frailty models for multiple event time data. - Biometrics 62: 1044–1052.

Péron, G. et al. 2010. Capture-recapture models with heterogeneity to study survival senescence in the wild. - Oikos 119: 524–532.

Péron, G. et al. 2016. Evidence of reduced individual heterogeneity in adult survival of long-lived. - Evolution. 70: 2909–2914.

Pirotta, E. et al. 2015. Estimating spatial, temporal and individual variability in dolphin cumulative exposure to boat traffic using spatially explicit capture-recapture methods. Anim. Cons. - 18: 20-31.

Pistorius, P. A. et al. 2008. Seasonal survival and the relative cost of first reproduction in adult female southern elephant seals. – J. Mamm. 89: 567-574.

Plard, F. et al. 2012. Modeling reproductive trajectories of roe deer females: Fixed or dynamic heterogeneity? - Theor. Popul. Biol. 82: 317–328.

Plard, F. et al. 2015. Quantifying the influence of measured and unmeasured individual differences on demography. - J. Anim. Ecol. 84: 1434–45.

Pledger, S. 2000. Unified maximum likelihood estimates for closed capture-recapture models using mixtures. - Biometrics 56: 434–442.

Pledger, S. 2005. The performance of mixture models in heterogeneous closed population capture-recapture. - Biometrics 61: 868–873.

Pledger, S. and Schwarz, C. J. 2002. Modelling heterogeneity of survival in band-recovery data using mixtures. - J. Appl. Stat. 29: 315–327.

Pledger, S. and Phillpot, P. 2008. Using mixtures to model heterogeneity in ecological capture-recapture studies. - Biometrical J. 50: 1022–1034.

Pledger, S. et al. 2003. Open Capture-Recapture Models with Heterogeneity: I. Cormack-Jolly-Seber Model. - Biometrics 59: 786–794.

Pledger, S. et al. 2010. Open capture-recapture models with heterogeneity: II. Jolly-Seber model. - Biometrics 66: 883–890.

Plum, A. and Ayllón, S. 2015. Heterogeneity in unemployment state dependence. - Econ. Lett. 136: 85–87.

Plummer, M. 2003. JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling (K Hornik, F Leisch, and A Zeileis, Eds.). - Proc. 3rd Int. Work. Distrib. Stat. Comput. March 2022: 0.

Pollock, K. H. 1980. Capture-recapture models: a review of current methods, assumptions and experimental design. - No. 1308

Pollock, K. H. 1981. Capture-recapture models allowing for age-dependent survival and capture rates. - Biometrics 37: 521–529.

Pollock, K. H. 2002. The use of auxiliary variables in capture-recapture modelling: an overview. - J. Appl. Stat. 29: 85–102.

Pollock, K. H. et al. 1990. Statistical Inference for Capture-Recapture Experiments. - Source Wildl. Monogr.: 3–97.

Pradel, R. 2005. Multievent: an extension of multistate capture-recapture models to uncertain states. - Biometrics 61: 442–7.

Pradel, R. 2009. The Stakes of Capture–Recapture Models with State Uncertainty. - In: Modeling Demographic Processes In Marked Populations. pp. 781–795.

Pradel, R. et al. 2010. Estimating population growth rate from capture-recapture data in presence of capture heterogeneity. - J. Agric. Biol. Environ. Stat. 15: 248–258.

Pradel, R. et al. 2012. Breeding Experience Might Be a Major Determinant of Breeding Probability in Long-Lived Species: The Case of the Greater Flamingo. - Plos One 7: e51016.

Prowse, V. 2012. Modeling Employment Dynamics With State Dependence and Unobserved Heterogeneity. - J. Bus. Econ. Stat. 30: 411–431.

Putter, H. and Van Houwelingen, H. C. 2014. Dynamic frailty models based on compound birth-death processes. - Biostatistics 16: 550–564.

Raman, S. et al. 2010. Infinite mixture-of-experts model for sparse survival regression with application to breast cancer. - BMC Bioinformatics 11 Suppl 8: S8.

Rasmussen, C. E. C. E. 2000. The infinite Gaussian mixture model. - Adv. Neural Inf. Process. Syst. 12: 554–560.

Reichert, B. E. et al. 2012. Extreme weather and experience influence reproduction in an endangered bird. – Ecology 93: 2580-2589.

Reid, J. M. et al. 2003. Age-specific reproductive performance in red-billed choughs *Pyrrhocorax pyrrhocorax*: patterns and processes in a natural population. – J. Anim. Ecol. 72: 765-776.

Reid, J. M. et al. 2010. Parent age, lifespan and offspring survival: structured variation in life history in a wild population. - J. Anim. Ecol. 79: 851-862.

Rexstad, E. A. and Anderson, D. R. 1992. Heterogeneous Survival Rates of Mallards (Anas-Platyrhynchos). - Can. J. Zool. 70: 1878–1885.

Reznick, D. 1985. Costs of Reproduction: An Evaluation of the Empirical Evidence. - Oikos 44: 257–267.

Robert, A. et al. 2012. The interaction between reproductive cost and individual quality is

mediated by oceanic conditions in a long-lived bird. – Ecology 93: 1944-1952.

Roff, D. A. 1992. The evolution of life histories: theory and analysis. - Chapman and Hall.

Rouan, L. et al. 2009. Estimation of lifetime reproductive success when reproductive status cannot always be assessed. - Model. Demogr. Process. Mark. Popul.: 867–879.

Roulin, A. et al. 2003. Female colour polymorphism covaries with reproductive strategies in the tawny owl *Strix aluco*. – J. Avian Biol. 34: 393-401.

Royle, J. A. 2008. Modeling individual effects in the Cormack-Jolly-Seber model: A state-space formulation. - Biometrics 64: 364–370.

Sanz-Aguilar, A. et al. 2008. The cost of reproduction and experience-dependent vital rates in a small petrel. - Ecology 89: 3195–3203.

Sanz-Aguilar, A. et al. 2011. Studying the reproductive skipping behavior in long-lived birds by adding nest inspection to individual-based data. - Ecol. Appl. 21: 555–564.

Sauer, J. R. and Slade, N. A. 1987. Uinta ground squirrel demography: is body mass a better categorical variable than age? - Ecology 68: 642–650.

Schwarz, C. J. 2001. The Jolly-seber model: More than just abundance. - J. Agric. Biol. Environ. Stat. 6: 195–205.

Schwarz, C. J. et al. 1993. Estimating migration rates using tag-recovery data. - Biometrics 49: 177–193.

Seber, G. A. F. 1965. A note on the multiple-recapture census. - Biometrika 52: 249–259.

Sedinger, J. S. et al. 2008. Fidelity and breeding probability related to population density and individual quality in black brent geese *Branta bernicla nigricans*. – J. Anim. Ecol. 77: 702-712.

Senner, N. et al. 2015. An ontogenetic perspective on individual differences. - Proc. R. Soc. B 282: 20151050.

Short, R. V. and Balaban, E. 1994. The Differences Between the Sexes. - Cambridge

University Press.

Skrondal, A. and Rabe-hesketh, S. 2014. Handling initial conditions and endogenous

covariates in dynamic / transition models for binary data with unobserved heterogeneity.

- J. R. Stat. Soc. 63: 211–237.

Souchay, G. et al. 2014. To breed or not: a novel approach to estimate breeding propensity

and potential trade-offs in an Arctic-nesting species. - Ecology 95: 2745-2756.

Spiegelhalter, D. J. et al. 2014. The deviance information criterion: 12 years on. - J. R. Stat.

Soc. Ser. B Stat. Methodol. 76: 485–493.

Stearns, S. C. 1976. Life-History Tactics: A Review of the Ideas. - Q. Rev. Biol. 51: 3–47.

Stearns, S. C. 1992. The evolution of life histories. - Oxford University Press.

Steiner, U. K. and Tuljapurkar, S. 2012. Neutral theory for life histories and individual

variability in fitness components. - Proc. Natl. Acad. Sci. U. S. A. 109: 4684–4689.

Steiner, U. K. et al. 2010. Dynamic heterogeneity and life history variability in the kittiwake.

- J. Anim. Ecol. 79: 436–444.

Stoelting, R. E. et al. 2015. Life-history tradeoffs and reproductive cycles in Spotted Owls. –

Auk 132: 46-64.

Stover, J. P. et al. 2012. Demographic heterogeneity impacts density-dependent population

dynamics. - Theor. Ecol. 5: 297–309.

Tenan, S. et al. 2014. Bayesian model selection: The steepest mountain to climb. - Ecol.

Modell. 283: 62–69.

Tuljapurkar, S. et al. 2009. Dynamic heterogeneity in life histories. - Ecol. Lett. 12: 93–106.

van de Pol, M. and Verhulst, S. 2006. Age-dependent traits: a new statistical model to

separate within- and between-individual effects. - Am. Nat. 167: 766–73.

van de Pol, M. and Wright, J. 2009. A simple method for distinguishing within-versus

between-subject effects using mixed models. - Anim. Behav. 77: 753–758.

van Noordwijk, a. J. and de Jong, G. 1986. Acquisition and Allocation of Resources: Their
Influence on Variation in Life History Tactics. - Am. Nat. 128: 137–142.

Vaupel, J. W. 2002. Life expectancy at current rates vs. current conditions: A reflection
stimulated by Bongaarts and Feeney's "How Long Do We Live?" - Demogr. Res. 7:
365–377.

Vaupel, J. W. and Yashin, A. I. 1985. Heterogeneity's ruses: some surprising effects of
selection on population dynamics. - Am. Stat. 39: 176–185.

Vaupel, J. W. and Missov, T. I. 2014. Unobserved population heterogeneity: A review of
formal relationships. - Demogr. Res. 31: 659–686.

Vaupel, J. W. et al. 1979. The impact of heterogeneity in individual frailty on the dynamics of
mortality. - Demography 16: 439–454.

Weladji, R. B. et al. 2008. Heterogeneity in individual quality overrides costs of reproduction
in female reindeer. – Oecologia 156: 237-247.

White, G. C. and Burnham, K. P. 1999. Program MARK: survival estimation from
populations of marked animals. - Bird Study 46: 120–139.

White, G. C. and Cooch, E. G. 2017. Population abundance estimation with heterogeneous
encounter probabilities using numerical integration. - J. Wildl. Manage.: In press. DOI:
10.1002/jwmg.21199

Wienke, A. 2010. Frailty models in survival analysis. - CRC Press.

Wilson, A. J. and Nussey, D. H. 2010. What is individual quality? An evolutionary
perspective. - Trends Ecol. Evol. 25: 207–214.

Wintrebert, C. M. A. et al. 2005. Joint modelling of breeding and survival in the kittiwake
using frailty models. - Ecol. Modell. 181: 203–213.

Wooldridge, J. M. 2005. Simple solutions to the initial conditions problem in dynamic,
nonlinear panel data models with unobserved heterogeneity. - J. Appl. Econom. 20: 39–

54.

Worthington, H. et al. 2015. Analysing Mark-Recapture-Recovery Data in the Presence of

Missing Covariate Data Via Multiple Imputation. - J. Agric. Biol. Environ. Stat. 20: 28–

46.

Yashin, A. I. et al. 2001. Hidden Frailty: Myths and Reality. http://bit.ly/2nclmx7

Yashin, A. I. et al. 2008. Model of hidden heterogeneity in longitudinal data. - Theor. Popul.

Biol. 73: 1–10.

Yoccoz, N. G. et al. 2002. Costs of reproduction in common eiders (Somateria mollissima):

an assessment of relationships between reproductive effort and future survival and

reproduction based on observational and experimental studies. - J. Appl. Stat. 29: 57–64.

Zens, M. S. and Peart, D. R. 2003. Dealing with death data: Individual hazards, mortality and

bias. - Trends Ecol. Evol. 18: 366–373.

Zheng, C. Z. et al. 2007. Age-dependent survival analyzed with Bayesian models of mark-

recapture data. – Ecology 88: 1970-1976.

Table 1: Case studies[1] reporting an analysis of individual heterogeneity in demographic

parameters (i.e. survival and reproductive traits) within a CR context. The table lists the

reference, the studied species, the main outcome as explicitly stated in the paper, and how

individual heterogeneity was assessed. Individual heterogeneity corresponded either to the

total amount of heterogeneity ("a priori" cases) or to heterogeneity measured using some

metrics ("a posteriori" cases). In these latter cases, the metrics used are provided.

Authors	Studied species	Main finding	Metric of individual heterogeneity
Guéry et al. 2017	Common eider *Somateria mollissima*	"Survival of the two migrating arctic populations was impacted directly by changes in the NAO, whereas the subarctic resident population was affected by the NAO with time lags of 2– 3 years. Moreover, we found evidence for intra-population heterogeneity in the survival response to the winter NAO in the Canadian eider population, where individuals migrate to distinct wintering areas"	A priori: Two classes of heterogeneity (finite mixture models)
Péron et al. 2016	Birds (5 species) and Mammals (4 species)	"Individual heterogeneity in survival was higher in species with short-generation time (< 3 years) than in species with long generation time (> 4 years)"	A priori: Two classes of heterogeneity (finite mixture models) and Continuous distribution of heterogeneity
Kennamer et al. 2016	Wood Duck *Aix sponsa*	"Strong positive relationship between survival and the number of successful nests" "Body mass was not a good proxy of quality"	A posteriori: Early incubation body mass
Fay et al. 2016a	Wandering albatross *Diomedea exulans*	"Age at first reproduction is negatively related to both reproductive performance and adult survival"	A priori: Two classes of heterogeneity (finite mixture models)
Garnier et al. 2016	Alpine ibex *Capra ibex*	"Adverse environmental conditions, such as disease outbreaks, may lead to survival costs of reproduction in long-lived species"	A priori: Two classes of heterogeneity (finite mixture models)

Hileman et al. 2015	Milksnake *Lampropeltis trinagulum*	"estimate adult survival (0.72±0.16) and abundance (N = 85±35.2)"	A posteriori: Observed maximum detection frequency
Link and Hesed 2015	Red-backed Salamander *Plethodon cinereus*	"Female *P. cinereus* mature earlier and grow more quickly than males"	A priori: Continuous distribution of heterogeneity
Hartson et al. 2015	Steelhead Trout *Oncorhynchus mykiss*	"Negative relationship between density and specific growth rate over a wide range of densities, but reductions in survival only at the highest densities"	A posteriori: Body length
Hua et al. 2015	Cumberlandian Combshell *Epioblasma brevidens*	"The overall mean detection probability and survival rate of released individuals reached 97.8 to 98.4% and 99.7 to 99.9% (per month)"	A priori: Continuous distribution of heterogeneity
Chambert et al. 2015	Weddell Seal *Leptonychotes weddellii*	"The probability of being absent from colonies was higher (1) in years when the extent of local sea ice was larger, (2) for the youngest and oldest individuals, and (3) for females with less reproductive experience"	A priori: Continuous distribution of heterogeneity
Pirotta et al. 2015	Bottlenose Dolphin *Tursiops troncatus*	"There were marked inter-individual differences in the predicted amount of time dolphins spent in the presence of boats, and individuals tended to be consistently over- orunderexposed across summers"	A priori: Continuous distribution of heterogeneity
Stoelting et al. 2015	California Spotted Owl *Strix occidentalis occidentalis*	"Breeding reduced the likelihood of reproducing in the subsequent year by 16% to 38%, but had no influence on subsequent survival"	A priori: Continuous distribution of heterogeneity
Souchay et al. 2014	Greater Snow Goose *Chen caerulescens atlantica*	"Cost of reproduction on breeding propensity in the next year, but once females decide to breed, nesting success is likely driven by individual quality"	A posteriori: Previous breeding status
Koons et al. 2014	Wild Boar *Sus scrofa* and Lesser Snow Goose *Chen caerulescens caerulescens*	"Senescence can be severe for natural causes of mortality in the wild, while being largely non-existent for anthropogenic causes"	A posteriori: Cause-specific mortality
Horswill et al. 2014	Macaroni Penguin *Eudyptes chrysolophus*	"Survival of macaroni penguins is driven by a combination of individual quality, top-down	A posteriori: Body mass

		predation pressure and bottom-up environmental forces"	
Lindberg et al. 2013	Pacific Black Brant *Branta bernicla nigricans*	"Annual survival of individuals marked as goslings was heterogeneous among individuals and year specific […]. Adult survival (0.85±0.004) was homogeneous and higher than survival of both groups of juveniles. The annual recruitment probability was heterogeneous for brant >1-year-old"	A priori: Two classes of heterogeneity
Chambert et al. 2013	Weddell Seal *Leptonychotes weddellii*	"Existence of a latent individual heterogeneity in the population, with robust individuals expected to produce twice as many pups as 'frail' individuals"	A priori: Continuous distribution of heterogeneity
Barbraud et al. 2013	Wandering Albatross *Diomedea exulans*	"Strong evidence for heterogeneity in survival with one group of individuals having a 5.2% lower annual survival probability than another group"	A priori: Two classes of heterogeneity
Blomberg et al. 2013	Greater Sage Grouse *Centrocercus urophasianus*	"Evidence for heterogeneity among females with respect to reproductive success; compared with unsuccessful females, females that raised a brood successfully in year t were more than twice as likely to be successful in year t+1"	A posteriori: Previous breeding status
Morano et al. 2013	North American Elk *Cervus elaphus*	"No difference in survival probabilities between pregnant and nonpregnant individuals or as a function of recruiting an offspring […and] negative effect of recruiting an offspring in the current year on becoming pregnant the following year"	A posteriori: Lactation status and body condition
Pradel et al. 2012	Greater Flamingo *Phoenicopterus roseus*	"Breeding probability varied within three levels of experience. [… and] With random effects, the advantage of experience was unequivocal only after age 9 while in young having > 1 experience was penalizing"	A priori: Continuous distribution of heterogeneity and A posteriori: Experience
Reichert et al. 2012	Florida Snail Kite *Rostrhamus sociabilis plumbeus*	"experience is an important factor determining whether or not individuals attempt to breed during harsh environmental conditions"	A posteriori: Experience

Robert et al. 2012	Monteiro's Storm Petrel *Oceanodroma monteiroi*	"reproductive costs act on individuals of intermediate quality and are mediated by environmental harshness"	A posteriori: successful vs. unsuccessful breeders
Hernandez-Matias et al. 2011	Bonelli's eagle *Aquila fasciata*	"4-year-old and older successful breeders were more likely to breed the following year than failed adult breeders (0.869 vs. 0.582), suggesting that the cost of reproduction is small in comparison with the variation in quality among individuals or their territories "	A posteriori: successful vs. unsuccessful breeders
Briggs et al. 2011	Swainson's hawks *Buteo swainsoni*	"Adult survival was inversely correlated with average reproductive output, with individuals producing >2 offspring having decreased survival [… and] reproduction in any year was positively correlated with survival"	A posteriori: Average annual nest productivity
Lee 2011	Northern elephant seals *Mirounga angustirostris*	"Primiparous breeders did not suffer more than experienced breeders during years of environmental stress. [… and] Lower variances in survival of multiparous breeders suggest that primiparous adults constitute a more heterogeneous portion of the population"	A posteriori: inexperienced vs. experienced breeders
Moyes et al. 2011	Red deer *Cervus elaphus*	"The probability of reproducing unsuccessfully after a successful year is relatively low and varies very little, but is highest in young individuals with low PARE. […and] Reproduction costs increase with declining PARE"	A posteriori: Proportional age-specific reproductive effort (PARE)
Marzolin et al. 2011	Dipper *Cinclus cinclus*	"Strong evidence for actuarial senescence with an onset of senescence estimated at about 2.3 years"	A priori: Continuous distribution of heterogeneity
Buoro et al. 2010	Atlantic salmon *Salmo salar*	"Cost of reproduction on survival for fish staying in freshwater and a survival advantage associated with the "decision" to migrate"	A priori: Continuous distribution of heterogeneity
Kovach et al. 2010	Intertidal snail *Nucella lima*	"Survival estimates from the best-fit model were different between habitat types"	A posteriori: Microhabitat use, individual color and length

Reid et al. 2010	Red-billed choughs *Pyrrhocorax pyrrhocorax*	"The negative correlation between offspring survival and maternal lifespan was strongest when environmental conditions meant that offspring survival was low across the population"	A posteriori: Longevity
Maniscalco et al. 2010	Steller sea lion *Eumetopias jubatus*	"Females which gave birth had a higher probability of surviving and giving birth in the following year compared to females that did not give birth"	A posteriori: Give birth vs. do not give birth
Millon et al. 2010	Tawny owl *Strix aluco*	"Females who postponed reproduction to breed for the first time at age 3 during an Increase phase, produced more recruits, even when accounting for birds that may have died before reproduction. No such effects were detected for males"	A posteriori: Age at first breeding
Lescroël et al. 2009	Adélie penguin *Pygoscelis adeliae*	"Adult survival ranged from 64-79%, with BQI accounting for 91% of variability in the entire study population, but only 17% in experienced breeders"	A posteriori: Breeding quality (BQI)
Bonenfant et al. 2009	Bighorn sheep *Ovis canadensis*	"In all age classes, natural survival was either weakly related to (lambs, adult rams) or positively associated (yearling rams) with early horn growth"	A posteriori: Early horn growth
Sedinger et al. 2008	Black brent geese *Branta bernicla nigricans*	"Individuals with a higher probability of breeding in one year also had a higher probability of breeding the next year"	A posteriori: Bred in the previous years vs. did not
Pistorius et al. 2008	Southern elephant seal *Mirounga leonina*	"Mean postbreeding (pelagic phase between breeding and molting, about 62 days) survival of primiparous females was 0.830 compared to 0.912 for more-experienced females"	A posteriori: Reproductive experience
Weladji et al. 2008	Reindeer *Rangifer tarandus*	"Successful breeders had higher survival and subsequent reproductive success than experienced non-breeders and unsuccessful breeders, independently of the age at primiparity. […and] Successful females at early primiparity remained successful throughout their life"	A posteriori: Age at primiparity
Le Bohec et	King penguin	"Failed breeders in year t have a	A posteriori:

al. 2007	*Aptenodytes patagonicus*	lower probability to reproduce successfully in year t + 1 than non-breeders in year t [… and] successful breeders showed higher survival probability"	successful vs. unsuccessful breeders
Zheng et al. 2007	Glanville fritillary butterfly *Melitaea cinxia*	"We found that mortality rate increased with age, that mortality rate was much higher during the day than during the night, and that the life span of females originating from newly established populations was shorter than the life span of females from old populations"	A priori: Continuous distribution of heterogeneity
Hadley et al. 2007	Weddell seal *Leptonychotes weddellii*	"Presence of reproductive costs to survival (mean annual survival probability was 0.91 for breeders vs. 0.94 for nonbreeders) [… and] Reproductive costs to subsequent reproductive probabilities were also present for first-time breeders (mean probability of breeding the next year was 31.3% lower for first-time breeders than for experienced breeders)"	A posteriori: Breeding experience
Beauplet et al. 2006	Subantarctic fur seal *Arctocephalus tropicalis*	"Survival was lower for non-breeders than for breeders, among both prime-aged (0.938 vs 0.982) and older (0.676 vs 0.855) females [… and] non-breeders exhibited higher probabilities of being non-breeders the following year than did breeders (0.555 vs 0.414)"	A posteriori: Breeders vs. non-breeders
Blums et al. 2005	Tufted duck *Aythya fuligula* Common pochard *Aythya ferina* Northern shoveler *Anas clypeata*	"For all three species, females that tended to nest earlier than the norm exhibited the highest survival rates, but very early nesters experienced reduced survival and late nesters showed even lower survival. For shovelers, females in average body condition showed the highest survival, with lower survival rates exhibited by both heavy and light birds. For common pochard and tufted duck, the highest survival rates were associated with birds of slightly above-average condition, with	A posteriori: Relative body condition and relative time of nesting

		somewhat lower survival for very heavy birds and much lower survival for birds in relatively poor condition"	
Barbraud et al. 2005	Blue petrel *Halobaena caerulea*	"Survival of first-time breeders was lower than that of inexperienced nonbreeders [...]. Survival of inexperienced individuals (both breeders and nonbreeders), but not of experienced ones, was negatively affected by poor environmental oceanographic conditions [... and] Survival and the probability of breeding in the next year for experienced birds were higher for breeders than for nonbreeders"	A posteriori: Breeding experience
Wintrebert et al. 2005	Kittiwake *Rissa tridactyla*	"Survival is positively correlated with breeding indicating that birds with greater inclination to breed also had higher survival"	A priori: Continuous distribution of heterogeneity
Roulin et al. 2003	Tawny owls *Strix aluco*	"The proportion of all breeding females that were reddish-brown was greater in years when the breeding density was lower [... and] greyish females bred less often than reddish-brown females, although their survival probability was similar"	A posteriori: Colour polymorphism
Reid et al. 2003	Red-billed choughs *Pyrrhocorax pyrrhocorax*	"Females that ultimately reached the greatest ages had laid small clutches and fledged few offspring during their first few breeding attempts. Females that were more productive when they were young had relatively shorter lives"	A priori: Lifespan
Cam et al. 2003	Kittiwake *Rissa tridactyla*	"Individuals with shorter rearing periods had lower local survival during the first winter [... and] The length of the rearing period had long-term consequences on reproductive performance [... and] negative influence of rank on survival before recruitment and recruitment probability"	A posteriori: Rank and length of the rearing period
Cam et al. 2002a	Kittiwake *Rissa tridactyla*	"Birds that were more likely to survive were also more likely to breed, given that they survived"	A priori: Continuous distribution of heterogeneity

Cam et al. 2002b	Kittiwake *Rissa tridactyla*	"Squatters have a higher survival and recruitment probability, and a higher probability of breeding successfully in the first breeding attempt in all age-classes where this category is represented"	A posteriori: Squatters vs. non-squatters
Cam and Monnat 2000a	Kittiwake *Rissa tridactyla*	"The influence of age on survival and future breeding probability is not the same in nonbreeders and breeders"	A posteriori: Yearly reproductive state
Cam and Monnat 2000b	Kittiwake *Rissa tridactyla*	"First-time breeders have a lower probability of success, a lower survival and a lower probability of breeding in the next year than experienced breeders [… and] neither experienced nor inexperienced breeders have a lower survival or a lower probability of breeding in the following year than birds that skipped a breeding opportunity. [… and] When age and breeding success are controlled for, there is no evidence of an influence of experience on survival or future breeding probability"	A posteriori: Breeding experience and breeding status

[1] The literature survey was performed in ISI Web of Knowledge by looking for references corresponding to the following "Topic" keywords: (((("individual variability" OR "individual heterogeneity" OR "individual quality") AND ("capture-recapture" OR "mark-recapture" OR "capture-mark-recapture"))). A total of 162 references were recovered. We read all summaries and only selected case studies looking for individual heterogeneity in demographic parameters estimated from CR models.

Table 2. Detection of a trade-off between reproduction and survival using multistate capture-recapture models after individual heterogeneity is accounted for. We simulated multistate capture-recapture data with two states, non-breeding (NB) and breeding (B). To mimic individual heterogeneity, we considered robust individuals with survival $\phi_{NB} = 0.7$ and $\phi_B = 0.8$ and frail individuals with survival $\phi_{NB} = 0.7$ and $\phi_B = 0.6$, the only difference being in the survival of frail breeders that is much lower than that of robust breeders. For each group, we simulated the fate of 100 newly marked individuals in each year of a 6-year experiment. We report parameter estimates from two multistate models in which i) we ignored individual heterogeneity (column '*Ignoring* individual heterogeneity) and ii) we explicitly incorporated an individual covariate to handle this source of heterogeneity (column '*Incorporating* individual heterogeneity'). The parameters we used to simulate the data are given in the column 'Truth'. We refer to the Appendix for more details. **The cost of breeding on survival is detected only in frail individuals after accounting for individual heterogeneity through quality** ($\Delta AICc = 120$).

Parameter	*Ignoring* Individual heterogeneity	*Incorporating* individual heterogeneity	Truth
Survival of frail non-breeders	0.69 [0.67; 0.72]	0.70 [0.66; 0.73]	0.7
Survival of frail breeders	0.70 [0.68; 0.72]	0.58 [0.55; 0.61]	0.6
Survival of robust non-breeders	0.69 [0.67; 0.72]	0.69 [0.65; 0.72]	0.7

Survival of robust breeders	0.70 [0.68; 0.72]	0.80 [0.77; 0.82]	0.8
Transition from non-breeding to breeding	0.78 [0.75; 0.80]	0.78 [0.75; 0.80]	0.8
Transition from breeding to non-breeding	0.31 [0.29; 0.33]	0.31 [0.28; 0.33]	0.3
Detection	0.90 [0.88; 0.91]	0.90 [0.88; 0.91]	0.9

1
2

Figure 1. Senescence is masked when individual heterogeneity is not accounted for:

random-effect capture-recapture model. We simulated the fate of 500 individuals (in grey)

from a single cohort with survival decreasing as they age over a 20-year study. We also added

a frailty for each individual $logit(\phi_i(a)) = \beta_0 + \beta_1 a + \varepsilon_i$ where $\varepsilon_i \sim N(0, \sigma^2)$. We used

$\beta_0 = 1, \beta_1 = -0.05$ and $\sigma = 1$. We considered the same detection probability $p = 0.5$ for all

individuals. We report the age-specific survival patterns from two models in which i) we

ignored individual heterogeneity (in blue) and ii) we incorporated an individual random effect

to handle with this source of heterogeneity (in green), both to be compared to the actual trend

that we used to simulate the data (in red). Clearly, ignoring individual heterogeneity obscures

senescence in survival. We refer to the Appendix for more details.

Figure 2. Senescence is masked when individual heterogeneity is not accounted for:

finite-mixture capture-recapture model. We simulated the fate of 1000 individuals from a

single cohort that were split into a group of robust individuals in proportion π with constant

high survival ϕ_R and a group of frail individuals with survival ϕ_F that aged over the 20 years

of the study according to the relationship $logit(\phi_F(a)) = \beta_0 + \beta_1 a$. We used $\pi = 0.3$,

$\phi_R = 0.85$, $\beta_0 = 0$ and $\beta_1 = -0.07$. We considered the same detection probability $p = 0.5$

for all individuals. We report the age-specific survival patterns from two models in which i)

we ignored individual heterogeneity (in blue) and ii) we used mixture models with two hidden

classes of individuals to handle with heterogeneity (in green), both to be compared to the

actual trend that we used to simulate the data (in grey). Clearly, ignoring individual

heterogeneity obscures senescence in survival. We refer to the Appendix for more details.