REGULATION OF LIFE SPAN BY THE GUT MICROBIOTA IN THE SHORT-LIVED AFRICAN

TURQUOISE KILLIFISH

Patrick Smith[1,5], David Willemsen[1,5], Miriam Popkes[1,5], Franziska Metge[1], Edson

Gandiwa[2], Martin Reichard[3], and Dario Riccardo Valenzano[1,4, *]

[1] Max Planck Institute for Biology of Ageing, Cologne, Germany

[2] Chinhoyi University of Technology, Chinhoyi, Zimbabwe

[3] Institute of Vertebrate Biology, Czech Academy of Sciences, Brno, Czech Republic

[4] CECAD, University of Cologne, Cologne, Germany

[5] These authors equally contributed to the paper

[*] Correspondence: dvalenzano@age.mpg.de

ABSTRACT

Gut bacteria occupy the interface between the organism and the external environment, contributing to homeostasis and disease. Yet, the causal role of the gut microbiota during host aging is largely unexplored. Here, using the African turquoise killifish (*Nothobranchius furzeri*), a naturally short-lived vertebrate, we show that the gut microbiota plays a key role in modulating vertebrate life span. Recolonizing the gut of middle-age individuals with bacteria from young donors resulted in life span extension and delayed behavioral decline. This intervention prevented the decrease in microbial diversity associated with host aging and maintained a young-like gut bacterial community, characterized by overrepresentation of the key genera *Exiguobacterium, Planococcus, Propionigenium* and *Psychrobacter*. Our findings demonstrate that the natural microbial gut community of young individuals can causally induce long-lasting beneficial systemic effects that lead to life span extension in a vertebrate model.

INTRODUCTION

Life expectancy of different species in nature is regulated by a complex combination of genetic and non-genetic factors. Genetic manipulations in model organisms have revealed key conserved molecular pathways, including the insulin-IGF1 and the mTOR pathways, which regulate aging and life span across several species, spanning from yeast to mammals (Kapahi et al., 2010; Kenyon et al., 1993; Lapierre and Hansen, 2012). Environmental interventions such as temperature and dietary manipulations have also been importantly associated with life span modulation in several species. Among these, lower temperatures (Conti et al., 2006; Miquel et al., 1976; Valenzano et al., 2006a; Van Voorhies and Ward, 1999) and reduced nutrient intake (Fontana et al., 2010; Mair and Dillin, 2008) are key environmental factors that have been associated with prolonged life span.

Complex microbial communities covering external surfaces live at the interface between organisms and the external environment – from roots and leaves in plants, to skin, mucosal surfaces and gut in animals. These microbial communities participate in a wide range of key biological processes, including nutrient absorption (Semova et al., 2012), development (Sommer and Backhed, 2013), metabolism (Nicholson et al., 2012), immune modulation (Geva-Zatorsky et al., 2017), defence against pathogens (Kamada et al., 2013; Schuijt et al., 2016) and disease (Sampson et al., 2016).

Individual gut microbiota (GM) composition changes dramatically in various diseases (Baumgart and Carding, 2007; Garrett, 2015; Sokol et al., 2008) and during aging in flies, mice and humans (Clark et al., 2015; Langille et al., 2014; O'Toole and Jeffery, 2015). Healthy GM is typically characterized by large bacterial taxonomic diversity, whereas frailty is associated with loss of diversity and expansion of more pathogenic bacterial species (Claesson et al., 2012). Following antibiotic treatment, pathogenic bacterial species, such as *Clostridium*

difficile and *Enterococcus faecalis*, can restructure the human GM and cause severe chronic

conditions that pose a major threat for public health (Backhed et al., 2012; Cox and Blaser,

2015). Studies across different human age cohorts have shown that large changes in the

abundance of subdominant bacterial taxa in the gut are a hallmark of aging; moreover,

exceptionally long-lived individuals – including supercentenarians – are characterized by the

persistence of bacterial taxa associated with a more healthy status (Biagi et al., 2016). While

diversity-associated microbial taxa often decline during age, specific bacterial taxa, such as

Clostridiales, are associated with malnutrition and increased frailty (O'Toole and Jeffery,

2015). In flies, reducing GM dysbiosis by improving immune homeostasis promotes longer

life span (Guo et al., 2014; Li et al., 2016). Manipulating the GM towards a healthy state has

the therapeutic potential to improve health in specific diseases (Dodin and Katz, 2014; Kunde

et al., 2013). However, due to the lack of suitable short-lived vertebrate experimental models,

it is not known whether age-associated gut microbial community changes causally affect the

aging process and whether resetting a young GM in middle-age individuals can improve long-

term health and affect individual life span in normal aging individuals.

In this study, we develop the turquoise killifish (*Nothobranchius furzeri*), a naturally short-

lived vertebrate species with a life span of a few months in captivity (Valenzano et al., 2015),

as a new model organism to study aging in the host gut and microbiota. We show that

turquoise killifish (TK) have a complex GM (both in the wild and in captivity), similar in

taxonomic diversity to that of mammals. We also show that during aging the overall microbial

diversity in the TK gut decreases, with increased over-representation of pathogenic

Proteobacteria. By acutely recolonizing middle age individuals with GM from young donors,

we developed an intervention that enabled fish to live significantly longer, remain more active

at old age, and maintain a highly diverse GM. Transcriptome analysis additionally revealed

that gut aging is associated with increased inflammation and reduced proliferation. Here we

provide the first evidence that acute gut microbiota transfer in the context of normal aging can

significantly prolong life span in a vertebrate, becoming a novel candidate life span enhancing

intervention. Our study also promotes the turquoise killifish as a highly suitable vertebrate

model system to study the crosstalk between intestine and gut-microbiota during host aging.

RESULTS

The TK is a short-lived vertebrate with a complex GM

TK is a naturally short-lived vertebrate, whose genome sequence has become available (Reichwald et al., 2015; Valenzano et al., 2015) and that is amenable to genetic manipulations via transgenesis or genome editing (Harel et al., 2015; Valenzano et al., 2011). Remarkably, this species is characterized by a broad spectrum of aging phenotypes, including cancer, neurodegeneration, and behavioral decline (Harel and Brunet, 2015; Kim et al., 2016). TK are adapted to living in conditions of intermittent availability of water and to surviving during brief rainy seasons and long dry seasons (Cellerino et al., 2015). In captivity, it lives between four to eight months, depending on the strain (Valenzano et al., 2015). However, nothing is known about its associated commensal microbes and whether its gut microbial complexity and taxonomic richness matches that of short-lived invertebrate model organisms, such as worms and flies, or that of longer-lived vertebrate model organisms, such as mice and zebrafish. To determine the TK's gut microbial composition, we sequenced the hyper-variable V3/V4 regions of the 16S rRNA gene amplicon from intestines of captive TK (N = 11, Materials and Methods). We found that TK are characterized by a gut microbial taxonomic diversity similar to other, longer-lived vertebrates, including zebrafish, mice and humans (Kostic et al., 2013; Qin et al., 2010; Zac Stephens et al., 2016) (Figure 1A). This bacterial taxonomic diversity is an order of magnitude higher than the gut microbial diversity present in invertebrate model organisms such as worms (Cabreiro and Gems, 2013) and flies (Buchon et al., 2013) in the laboratory (Figure 1A). Remarkably, the four most abundant bacterial phyla present in the TK's gut, i.e. Proteobacteria, Firmicutes, Actinobacteria and Bacteroidetes, are also the four most abundant human gut bacterial divisions (Zoetendal et al., 2006) (Figure 1B and

106   Supplementary file 1), although in different proportions (Figure 1A). The unique combination

107   of a complex gut

FIGURE 1

Figure 1. TK as a model to study aging of the GM

(A) Above: relative bacterial OTU composition (pie charts) and diversity (y-axis of upper plot) across different

aging model organisms. Below: maximum life span (logarithm of days) in six model organisms, data from the

AnAge longevity database. (B) Heatmap of ranked relative abundance of OTU composition at the phylum-level

for different model organisms (Supplementary file 1). Bacterial phyla are ordered by their relative abundance in

the TK. The ranked abundance is color-coded, with higher ranks (red) indicating greater relative abundance and

lower ranks (blue) indicating lower relative abundance. White cells mark phyla that are not present in the

respective model organism.

microbial composition, similar to that of other vertebrate aging model organisms, and its

naturally short life span, combined with a wide spectrum of aging phenotypes, makes the

turquoise killifish an ideal system to study the role of the gut microbiota during vertebrate

aging.

Wild and captive TK populations share a core GM

To assess whether the GM of captive TK was representative of the microbial communities

associated with wild populations, we sequenced 16S rRNA gene amplicons from individual

fish that we collected from different localities in the natural habitat of this species, ranging

from the Gonarezhou National Park in Zimbabwe to the Gaza region in Mozambique (Figure

2A, Figure 2 – figure supplements 1B, Supplementary file 2A, Materials and Methods).

Sequencing these wild populations confirmed that Proteobacteria was the dominant phylum

also in most wild populations, similar to laboratory fish (Figure 2B, Figure 2 – figure

supplements 1C). Individuals from all wild populations had a more diverse GM than

laboratory fish (Figure 2C, Shannon alpha diversity, Dunn Kruskal-Wallis test, BH-adjusted p

values < 0.05). While standard frequency-based diversity measures of observed bacterial

taxonomic units (OTUs) (Simpson's and phylogenetic alpha diversity across the whole tree,

OTU abundance, Figure 2 – figure supplements 1A) were higher in wild populations, Chao1

alpha diversity, which gives more weigh to rare bacterial OTUs, was higher in laboratory fish

(Figure 2 – figure supplements 1A). These results possibly reflect the fact that laboratory fish

are dominated by few, high-abundance OTUs, hence resulting in having more "rare" bacterial

taxonomic units compared to wild populations. Differences in OTU abundance between

laboratory-raised and wild fish might reflect ecological differences between the standardized

laboratory conditions and the more heterogeneous wild environment, characterized by

fluctuations in temperature, nutrients as well as other biotic and abiotic factors (Blažek et al.,

2017). To test whether differences in ecology across distinct wild fish populations influenced

bacterial diversity

FIGURE 2

Figure 2. Core microbiota is conserved between wild and captive TK

(A) Map location of wild populations collected in Zimbabwe (Z) and Mozambique (M1 and M2). ZW:

Zimbabwe; MZ: Mozambique; SA: South Africa. (B) Relative abundance of bacterial phyla in wild populations

and laboratory fish. (C) Shannon Index alpha diversity in laboratory and wild fish (L = laboratory, derived from

individuals originally collected in Zimbabwe). (D) PCoA of the Weighted UniFrac beta diversity distance for

wild and laboratory fish. Adonis test: L vs. all groups: p value < 0.001, M2 vs. M1: p value < 0.001, Z vs. M1

and M2: p value < 0.05. (E) Dotplot of the Weighted UniFrac distance values to visualize beta diversity within

(purple background) and between (green background) the different populations. Single dots represent

comparisons between individual fish. Red horizontal line indicates the median for each comparison. (F) Heatmap

of the Weighted UniFrac distance among wild populations and laboratory fish.

levels, we ran a regression model between diversity indexes and different recorded ecological

factors, including altitude, pond size, maximum pond depth, vegetation, water conductivity,

water temperature and water turbidity (Supplementary file 2B). None of the recorded factors

was significantly correlated to alpha diversity levels (data not shown). This suggests that other

parameters, such as food availability, fish genetics, presence of parasites or additional biotic or

abiotic factors might be the key determinants of fish GM composition in the wild.

Although microbiota diversity in wild populations was higher than in laboratory-raised fish,

the microbial composition of laboratory fish was not separated from the species-specific

bacterial composition in wild fish, and was contained within the natural microbial variation of

the species (Figure 2D). Importantly, one wild population (M1) had a divergent composition

from all other populations, possibly due to over-representation of Planctomycetes (Figure 2B

and D, Figure 2 – figure supplements 1C and E). Between-individuals diversity measures (beta

diversity) showed that, while laboratory fish's bacterial diversity was lower than in the wild

populations, between-group bacterial diversity was higher between population M1 and all the

other populations, including laboratory fish (Figure 2E-F). These results indicate that

population M1 had a more divergent microbial diversity of all the tested populations, and that

the laboratory fish were not an outlier group (Figure 2D). Based on this, laboratory fish share a

core microbiota with wild populations, despite the differences in culturing conditions between

the laboratory and the wild, similar to what is seen in zebrafish (Roeselers et al., 2011). Thus,

while wild turquoise killifish populations differ from one another in terms of microbial

composition – possibly in association with ecological differences among different localities –

laboratory fish recapitulate the core gut microbiota of wild populations, making them ideally

suited to study how a complex microbiota influences host physiology.

Aging in the TK is associated with loss of microbial diversity and decreased expression of genetic markers of gut health

An important question is what changes in the gut microbial population occur during aging in vertebrates. To gain insight into the changes in GM composition occurring throughout aging in the TK, we performed 16S rRNA gene amplicon surveys in the gastrointestinal tract of six-week-old young-adult and 16-week-old individuals raised in captivity (Figure 3A, n = 16 young and 14 old fish). We found that bacterial taxonomic diversity (alpha diversity) significantly decreased between 6-week-old and 16-week-old fish, while bacterial abundance was not changed (Figure 3B, Figure 3 – figure supplements 1A and B). This indicates that, similar to humans (Claesson et al., 2012), aging in the TK is characterized by a significant reduction in gut bacterial richness. Bacterial composition was also significantly altered between young and old fish guts (Figure 3C), including an age-dependent decrease in Firmicutes (Kruskal-Wallis test, p value = 0.002) and Actinobacteria (Kruskal-Wallis test, p value = 0.013). Young and old fish GM had significantly different community structures (Figure 3C, UniFrac p value < 0.001; Bray-Curtis p value < 0.001, Adonis test, Figure 3 – figure supplements 1C). While individual fish GM diversity was higher in young than in old fish (Figure 3B), differences among old fish GM were more pronounced compared to young fish (Unweighted UniFrac beta diversity, Bonferroni-corrected p value = 6.7E-06; Bray-Curtis distance p value = 0.003; Figure 3 – figure supplements 1D and E). This indicates that although aging in the TK was associated with decreased bacterial taxonomic diversity within each fish's gut, distinct old fish guts had highly divergent bacterial communities. While young fish guts were significantly enriched for Bacteroidetes, Firmicutes and Actinobacteria, old fish were dominated by Proteobacteria (Figure 3D). We next investigated the predicted functional metagenome biomarkers associated with young and old fish's guts using PICRUSt

FIGURE 3

Figure 3. Changes in GM and gut transcriptome between young and old fish

(A) Representative 6-week-old (young) and 16-week-old (old) male TK. (B) Alpha diversity changes in observed OTUs in young (6 weeks) and old (16 weeks) TK. N = 16 six-week-old fish and 14 16-week-old fish. The groups are compared using the Mann Whitney U test; ** indicates a p value < 0.001. (C) Beta diversity microbiota analysis separates samples based on age. Above: Bray-Curtis analysis, below: Unweighted UniFrac analysis. TW: tank water control. Adonis test, p value < 0.001 in both comparisons between 6-week and 16-week-old fish. (D) Cladogram representing microbial taxa enriched in young (green) versus old (red) individuals. (E) Predicted metagenome function in young (green) and old (red) groups (LEfSe), representing functions with p value < 0.001. The x-axis indicates the Linear Discriminant Analysis score for all the significant metabolic functions. (F) Expression heatmap for the twenty top differentially expressed genes (DEGs) between young and old fish (n = 4 for each group). Top 20 genes are highly expressed in old fish, the bottom 20 are highly expressed in young fish. (G) Top 20 Gene Ontology (GO) terms of the DEGs between young and old fish. Enrichment values (bars) and the negative natural logarithm of p values (black dots) are shown.

*HAMP is the best protein blast hit in *Danio rerio* of the TK gene NFURG05812010005.

(Langille et al., 2013) and LEfSe (Segata et al., 2011). While young fish had GM strongly

associated with glycolysis and polysaccharide metabolism, old fish's GM was depleted of

bacteria associated with carbohydrate, nucleotide and amino acid metabolism, and was

enriched for bacteria associated with pathogenesis, transport and catabolism (Figure 3E). In

particular, bacterial motility and flagellar assembly was strongly increased in GM from old

fish. These terms are associated with increased virulence in bacteria (Josenhans and

Suerbaum, 2002), supporting that old fish had a higher prevalence of potentially pathogenic

bacteria.

To examine whether metagenomic biomarkers from bacterial taxonomic diversity (i.e. OTU)

data were consistent with host responses, we asked which were the transcriptional changes in

the gut associated with young and old status. To this end, we performed an RNA-Seq analysis

of whole gut in four 6-week old and four 16-week old fish (Figure 3F and G, Figure 3 – figure

supplements 2 and Materials and Methods). Young fish had a distinct expression signature of

active proliferation (Figure 3G, Figure 3 – figure supplements 2), consistent with the bacterial

metagenomic signature of replication. Old fish, on the other hand, had significant Gene

Ontology terms associated with immune and defense responses against pathogens as well as

inflammation, consistent with the bacterial metagenome signatures associated with host

disease and overall virulence. Together, these results strongly support that in old individuals

both changes in bacterial composition and gut transcriptome are consistent with a markedly

pathological gut environment, while young fish are characterized by a molecular signature of

healthy gut and a commensal bacterial community.

Young GM transfer prolongs life span and delays age-dependent motor decline

Interventions aimed at directly modifying the complex microbial composition in experimental

organisms and patients have been mostly focused on treating diseases such as *Clostridium*

difficile infections (Dodin and Katz, 2014; Lee et al., 2016), as well as obesity and type two diabetes (Kootte et al., 2012; Turnbaugh et al., 2006). However, the impact of a young GM in modulating aging and life span has not been explored to date in vertebrates (Clark et al., 2015). To test whether resetting a young-like GM in middle age could impact aging and affect life span, we treated middle-age fish (9.5 week-old) overnight with an antibiotic cocktail (VMNA, i.e. vancomycin, metronidazole, neomycin, ampicillin) (Figure 4A, Figure 4 – figure supplements 1A and Materials and Methods). The antibiotic treatment significantly reduced gut microbial content compared to pre-treatment levels (Figure 4 – figure supplements 1B). Antibiotic-treated fish were then exposed for 12 hours to the following conditions: six-week-old donor fish gut content (Ymt), 9.5-week-old fish gut content (Omt) and sham (Abx) (Figure 4A and Figure 4 – figure supplements 1A). After antibiotic treatment and 12-hour acute gut recolonization, fish were reintroduced in the water recirculation system in individual tanks and were subjected to regular feeding (Materials and Methods). Their survival under the different experimental conditions was then scored (Supplementary file 3A). Ymt fish underwent dramatic life span prolongation compared to three control groups, which received: (i) antibiotic-only (Abx) (21% life span increase in median life span, Logrank test p value = 5.89E-05), (ii) antibiotics and same-age (i.e. 9.5 weeks) gut content (Omt) (41% increase in median life span, Logrank test p value = 5.08E-06), or (iii) no-treatment (wt) (37% increase in median life span, Logrank test p value = 4.04E-09) (Figure 4B and Figure 4 – figure supplements 1C-F). Noteworthy, acute antibiotic treatment alone was sufficient to increase fish life span compared to the wt group (14% median life span increase, Logrank test p value = 0.0129) (Figure 4B). However, Omt fish did not live longer than the control, wt group (Figure 4B). Since Abx outlived Omt and wild-type fish, while Ymt fish outlived Abx fish, it is plausible that middle-age GM composition might be primed to induce damage in the host and that its removal is therefore beneficial. However, as the recolonization of middle-age

279   individuals with young fish gut content after antibiotic treatment prolongs life span even more,

280   this implies that young GM, *per se*, has beneficial effects on host physiology that are additive

281   to the effects of the antibiotic treatment.

FIGURE 4

A

Figure 4. Transfering young GM to adult fish prolongs life span and delays motor

decline

(A) Schematic representation of microbial transfer experiment (Materials and Methods). Experimental group

legend, Abx: fish receiving only antibiotic treatment at 9.5 weeks without direct recolonization. Omt: fish

receiving same-age GM transfer after antibiotic treatment at 9.5 weeks. Wt: wild-type, untreated fish. Ymt: fish

receiving 6-week-old fish GM transfer after antibiotic treatment at 9.5 weeks. VMNA: antibiotic cocktail of

vancomycin, metronidazole, neomycin and ampicillin. (B) Survival analysis. Statistical significance is calculated

by Logrank test. * indicates a p value < 0.05; *** indicates a p value < 0.001. (C) Exploratory behavior in

different treatments. Young and old wild-types are compared with a Kruskal-Wallis test (left), the remaining

groups are compared using a Dunn Kruskal-Wallis test for multiple comparisons, and the p values are adjusted

based on BH correction. Statistical significance: * indicates a p value < 0.05; ** indicates a p value < 0.01; ***

indicates a p value < 0.001.

We then asked whether treating young fish with old fish GM could also affect life span. Compared to fish that received either the same-age GM or to untreated control fish, six-week-old fish receiving 16-week-old fish's GM after antibiotic treatment did not have a different life span (Figure 4 – figure supplements 1G). Additionally, unlike middle-aged fish treated with antibiotics, young fish receiving antibiotic treatment did not live longer than untreated control fish (Figure 4 – figure supplements 1G). These results suggest that the timing of GM transfer is critical to inducing systemic effects and modulating life span.

It was shown in previous work that spontaneous exploratory behavior in TK decreases with age (Genade et al., 2005; Valenzano et al., 2006b). We therefore asked whether treating middle-age fish with young fish GM after antibiotic treatment could improve exploratory behavior performance, considered as an integrated measure of individual health. Using an automatic video-tracking system (Materials and Methods), we assayed spontaneous locomotor activity. Young, six-week-old fish, were significantly more active than 16-week-old fish (Supplementary file 3B and Figure 4C, Kruskal-Wallis chi-squared = 10.752, df = 1, p value = 0.00104). Remarkably, Ymt were more active at 16 weeks of life than Omt and wt fish at the same age, resembling younger fish performance (Figure 4C, Dunn Kruskal-Wallis multiple comparison test, BH-adjusted p value = 0.004). Additionally, all groups except Ymt underwent a significant decrease in spontaneous locomotor activity from a week post transfer to 16 weeks of life. This suggests that the transfer of young fish gut content had long-lasting effects on a global measure of physiological health, influencing individual survival and spontaneous exploratory behavior. Thus, depleting middle-age individuals from their resident GM resulted beneficial when acutely recolonized by young-associated GM (Ymt), and in part also when it was not followed by any acute recolonization (Abx). On the other hand, acutely recolonizing the gut with same age GM after antibiotic treatment did not lead to differences

compared to the untreated control group. These results establish gut microbial recolonization as a powerful life span enhancing intervention, which leads to significant effects also on behavioral performance.

Acute GM transfer affects microbial composition at old age

To assess the extent to which one acute transfer reset the GM in recipient fish after VMNA antibiotic treatment, we performed 16S rRNA gene amplicon surveys in fish that underwent microbial transfer at 9.5 weeks of age. Untreated 6-week-old (6wk) and 16-week-old (16wk) fish were also included in the analysis. One week post transfer there were no significant differences in alpha diversity metrics among treatment groups (Figure 5 – figure supplements 1A and B). However, Ymt and Omt fish had already significantly different gut microbial population diversity (UniFrac and Bray-Curtis distances, Figure 5 – figure supplements 1C and D and Supplementary file 4). Seven weeks after the microbial transfer (16 weeks of age, corresponding to the median life span for this species in captivity), Ymt fish had significantly higher bacterial richness compared to wild-type, 16-week-old fish (Figure 5A, Dunn Kruskal-Wallis test, BH-adjusted p value = 0.009). This shows that one acute transfer had long-lasting effects on GM diversity. Bacterial OTU abundance at 16 weeks was higher in Ymt fish compared to Omt fish (Figure 5A), but lower than 6-week old wild-type fish (6wk).

GM community structure in Ymt fish was also significantly altered compared to Omt and 16wk (Figure 5B and Figure 5 – figure supplements 1E); however, it did not statistically differ from young wild-type fish, i.e. the 6wk group. Furthermore, based on hierarchical clustering, Ymt fish clustered preferentially with 6wk fish (Figure 5C), showing that the GM-transfer from young donors significantly reset a young-like GM.

Young fish (6wk), as well as fish treated with young GM (Ymt), were more enriched with members of the Bacteroidetes and Firmicutes (Figure 5D) and with the genera

FIGURE 5

Figure 5. **Microbiota transfer at 9.5 weeks influences microbial composition at 16 weeks**

(A) Alpha diversity measured by number of observed OTUs in experimental groups at 16 weeks (16wk, Abx,

Omt, Ymt) and young controls (6wk). Dunn Kruskal-Wallis test for multiple comparisons. The p values are

adjusted based on BH correction. Statistical significance: ** indicates a p value < 0.01. (B) Microbiota

community analysis using Bray-Curtis (above) and Unweighted UniFrac (below) separates samples based on

young vs. old GM treatment (significant p values are shown, Adonis test). (C) Hierarchical clustering on OTU

data. Significant clusters are highlighted by red rectangles. P values: au = approximately unbiased; bp: bootstrap

probability. (D) Cladogram of bacterial taxa enriched in young wild-type (6wk) and Ymt groups versus 16wk and

Omt. (E) Boxplots of selected bacterial genera relative abundance in experimental groups. Bacterial genera are

those represented in Figure 5D, filtered by r-square value larger than 0.8, based on the correlation between

bacterial genus abundance vs median life span in each individual group (Supplementary file 5).

Carnobacterium, *Arthrobacter*, *Exiguobacterium*, *Planococcus*, *Psychrobacter*, *Enterococcus*

and *Halomonas* (Figure 5E). Additionally, analyzing the correlation between bacterial genera

abundance and the group-specific median lifespan, we identified a set of bacterial genera

whose abundance is highly correlated with group-specific longevity (Supplementary file 5).

We then used predicted functional metagenome analysis from GM composition to compare

experimental fish groups receiving same-age GM transfer and no-treatment controls to fish

receiving young gut content. The latter group was enriched for increased saccharolytic

potential, DNA repair and recombination among other functions (Figure 5 – figure

supplements 1F), which are functional terms associated with a younger, healthy-like

physiological state. Together, these results show that gut microbial recolonization from young

killifish resulted in increased microbial diversity and a persistence of a young-like bacterial

community until old age, whose composition could be at least in part responsible for the life

span prolongation.

Microbial transfer determines bacterium-to-bacterium connectivity at old age

To investigate whether gut recolonization affected bacterium-to-bacterium association in

different experimental groups, we harnessed OTU co-occurrence to generate bacterial

connectivity networks (Agler et al., 2016; Biagi et al., 2016). To assess the bacterium-to-

bacterium connectivity in each experimental group, we built a network model for each

experimental group based on OTU co-occurrence. Significantly co-occurring genera within

each group composed a network, whose nodes were single bacterial genera. Bacterium-to-

bacterium connections (edges) were established based on r-square values (Materials and

Methods). Using this analysis, we found that Proteobacteria had higher connectivity in the

shorter-lived 16wk and Omt groups compared to 6wk, Ymt and Abx (Figure 6A). Overall,

6wk and Ymt fish had the largest networks and a higher number of highly connected nodes

FIGURE 6

Figure 6. Bacterial co-occurrence connectivity

(A) Phyla composition of the bacterial co-occurrence networks in experimental groups at 16 weeks (16wk, Ymt, Omt Abx) and at 6 weeks (6wk) (Materials and Methods). (B) Cumulative sum of degree distribution in co-occurrence networks of all analyzed TK experimental groups (black: 6wk; dashed black line: 16wk; green: Ymt; red: Omt; blue: Abx) and a mouse cohort from (Langille et al., 2014) (Inlet, green: young mice; orange: middle-aged mice; red: old mice). X-axis shows the degree count and y-axis the cumulative sum of nodes with the corresponding degree. C) Visualization of co-occurrence networks in Fruchtermann-Reingold layout of 6wk, 16wk, Ymt, Omt, and Abx. Circle size increases with degree count and circle color corresponds to 6wk hubs (green) or 16wk hubs (orange).

(hub nodes), showing that a few genera of bacteria significantly co-occurred with a large set

of other bacterial genera (Figure 6B and C).

We then tested whether the relation between age of the host and bacterial connectivity

observed in TK guts was also conserved in mammals. We analyzed a published mouse 16S

amplicon survey from an aging cohort (Langille et al., 2014). Similarly to what we found in

the TK, young mice GM had more significantly connected nodes and a higher number of

bacterial hubs compared to middle age and old mice (Figure 6B, inset).

The identification of wild-type specific young and old-related hub bacterial genera enabled us

to study whether Ymt, Omt and Abx shared hub bacterial genera with either young or old fish

groups. Two hub bacterial clusters were identified in Ymt fish (Materials and Method).

Strikingly, one was composed of bacterial genera that were also hub-bacteria in young wild-

type fish (6wk) and included *Exiguobacterium*, *Planococcus*, *Propionigenium* and

Psychrobacter (Figure 6C, Ymt network, green nodes), while the other was composed of hub

bacteria from old wild-type fish (16wk) hub and included *Propionibacterium*, *Delftia*, and

Citrobacter (Figure 6C and Supplementary file 6). Remarkably, the bacterial hubs identified in

Omt and Abx overlapped exclusively with the old wild-type group (16wk) (Figure 6C, orange

nodes, and Supplementary file 6). These results support that bacterial network topology

reflects host age both in fish and mice, with younger biological age associated with larger

networks. Acute microbiota transfers significantly affect gut microbial population topology

and subsets of bacterial genera are identified as hub nodes in young-like and old-like microbial

communities. Therefore, not only bacterial composition, but also bacterium-bacterium co-

occurrence carries a key signature of host life span and network topology depends on a few

key bacterial hubs.

Microbial transfer affects expression of host genes associated with defence to bacteria, Tor pathway and extracellular matrix

Because acute microbial transfer dramatically changed GM composition by resetting a young-like microbial community in Ymt fish, we asked whether young GM transfer could also reset the host transcriptome towards a young-like status. To this end, we performed host intestine RNA-Sequencing in wild-type young (6wk) and old (16wk) fish, as well as in Omt and Ymt fish (Figure 7A and B, Figure 7 – figure supplements 1 and Materials and Methods). Interestingly, while young GM transfer reset a young-like GM environment in aged fish, host gene transcripts separated based on fish age, with all the transcriptomes obtained from 16-week old fish groups (16wk, Omt, Ymt) clustering together (Figure 7A). However, comparing transcriptomes in Omt and Ymt revealed a distinct signature of defence response to bacteria in Ymt and increased expression of genes associated with hyaluronic acid metabolism in Omt (Figure 7B and C). Comparing the expression levels of all groups to the 6wk group, we identified a set of genes whose expression in the gut was significantly changed comparing 16wk and Omt with 6wk; comparing 16wk and Ymt, but unchanged between Ymt and 6wk. These were genes associated with the TOR-pathway (DEPTOR) and with cell adhesion and extracellular matrix composition (DSCAM, OLFML2A, PAPLNB) (Figure 7D), suggesting a potential difference in cell adhesion and gut permeability between 16wk and Omt from one side, and Ymt and 6wk on the other side. Since we generated OTU abundance tables for individual fish and we also sequenced intestinal transcripts for the same individuals, we then generated an OTU-to-host trancripts correlation matrix (Data S1). Taking advantage of this resource, we could identify the transcripts that were highly correlated with the hub genera that are shared between Ymt and young fish (6wk) networks (Figure 6C, Figure 7E and Figure 7 – figure supplements 2). Remarkably, these genes include Jak2, an important gene involved in cellular proliferation and differentiation, and several genes encoding S and L- ribosomal

FIGURE 7

Figure 7. Transcriptional changes in host intestine after acute GM transfer

(A) Cluster analysis of gut RNA-Seq from intestines from six-week-old fish (6wk) and 16-week-old fish (16wk,

Omt (red) and Ymt (green)). (B) Expression heatmap for the 20 top differentially expressed genes (DEGs)

between Ymt and Omt fish (n = 4 for each group). The top 20 genes are highly expressed in Ymt fish, the bottom

20 are highly expressed in Omt fish. *HAMP is the best protein blast hit in *Danio rerio* of the TK gene

NFURG05812010005. (C) Gene Ontology (GO) analysis of the DEGs between Ymt and Omt fish. Enrichment

values (bars) and the negative natural logarithm of p values (black dots) are shown. (D) Relative gene expression

levels compared to six-week-old fish (6wk) of genes significantly differentially expressed between 16wk (gray),

Omt (red) and 6wk (black), but not between Ymt (green) and 6wk. These genes are also significantly

differentially expressed between 16wk and Ymt. (E) Based on the correlation matrix between OTU abundance

and transcript levels in individual fish (Data S1), we identified the gene transcripts whose expression correlates

more significantly with the hub bacteria shared between Ymt and 6wk fish (full analysis in Figure 7 – figure

supplements 2). Displayed are the top hits (chosen from an r-square value larger than 0.65 on both axes) shared

between *Psychrobacter* and *Exiguobacterium*, which have the highest OTU-transcript correlation trend based on

Figure 7 – figure supplements 2. RPLs include RPL7A, RPL8, RPL12, RPL37, RPL14, MRPL35; while RPS

include RPS3A, RPS7, RPS11, RPS24.

proteins (RPS and RPL genes) (Figure 7 – figure supplements 2), whose expression in other

species has been strongly associated with aging and longevity (Steffen et al., 2008). Thus,

while transfer of young GM to middle age fish maintained a young-like GM community

throughout old age, the gene-expression signature associated with host gut at 16 weeks of age

did not indicate overall rejuvenation. However, enhanced defense response against bacteria in

Ymt compared to Omt is compatible with an increased capacity to resist to the attack of

pathogenic gut bacteria, which could provide the basis for longer life span. Remarkably,

hyaluronic acid metabolism, altered between the experimental groups receiving young and old

gut contents, has been associated with increased inflammation, deregulated immune response

and risk for cancer (Cho et al., 2017; Tian et al., 2013), all of which could provide the basis for

life span modulation. Finally, we provide a dataset that allows to associating OTU levels in

individual fish to intestinal transcripts, enabling to investigate on how GM and host gene

expression are mutually regulated.

DISCUSSION

Key aims of research on aging are to understand the mechanisms behind the phenotypic changes that occur during aging and to identify novel life span enhancing interventions. Using the TK as a naturally short-lived vertebrate model system, we report the characterization of the changes in GM composition occurring during aging and the discovery of a novel life span enhancing intervention achieved by acutely transferring young GM to middle-age individuals after antibiotic treatment. This intervention resulted in the maintenance of an overall healthier physiological status, a highly diverse and young-like gut microbial community at late age and in an enhanced transcriptional signature of defense responses to bacteria.

Our results show that TK are characterized by a complex GM community, more species-rich than worms and flies and of the same order of magnitude of mammals, both in abundance and composition. Indeed, the four most abundant bacterial phyla observed in the TK are also the four most abundant phyla found in humans and mice. However, unlike mammals, Proteobacteria is the most abundant gut bacterial phylum found in killifish, similarly to other aquatic species, such as zebrafish (Roeselers et al., 2011). Since it is the shortest-lived vertebrate to date reproduced in captivity, the TK can become an ideal model to dissect the links between GM diversity and host aging.

Intriguingly, we show that while the gut bacterial diversity of captive-raised TK is lower than that of wild-caught populations, still captive fish recapitulate the core microbial diversity of wild TK. Laboratory-raised fish are not represented by distinct gut bacterial communities, but are rather dominated by a few high-abundance bacterial taxa that are already present in wild killifish populations. This supports the conclusion that fish raised in captivity have a GM

community that is representative for the species in nature. Additionally, we found that wild

populations of TK show large between-population gut bacterial diversity, with the emergence

of population-specific high-abundance taxa. This is plausibly caused by locality-specific

differences in ecological conditions, including climate, soil composition, parasites and

diversity of food sources (Nezhybová et al., 2017; Reichard et al., 2017).

Analyzing the changes in GM composition between young and old fish, we found that young

fish are characterized by a large taxonomic bacterial diversity. Old fish are less OTU-rich,

similar to what is observed in human cohorts from different age classes (Claesson et al., 2012).

Intriguingly, while individual bacterial diversity (alpha diversity) is higher in young

individuals, old fish are more dissimilar from one another, i.e. while each individual old fish

have a more homogeneous GM composition than young fish, any two old individuals have

more divergent bacterial communities compared to young individuals. This result raises the

possibility that the changes in composition and relative abundance in GM communities from

young to old individuals could be a function of i) initial individual GM composition, ii)

differences in individual immune system composition and function or iii) a combination of

initial individual GM composition and host immune function.

Aging in this experimental model was associated not only with reduced bacterial species

richness, but more specifically with loss of bacterial taxonomic units involved in carbohydrate,

nucleotide and amino acid metabolism, which in mice are associated with aging (Langille et

al., 2014). These same changes have been associated in humans with unhealthy aging

(Claesson et al., 2012; Rampelli et al., 2013), as well as with chronic conditions such as

obesity, type II diabetes and insulin resistance (Neis et al., 2015). The shift in microbial

composition between young and old fish was consistently characterized by a higher prevalence

of Proteobacteria in old individuals, while young individuals were significantly more enriched

in Firmicutes, Actinobacteria and Bacteroidetes. Additionally, functional metagenome

analysis showed that young fish had GM associated with carbohydrate metabolism, replication

and repair, as well as DNA repair, indicating the young GM's ability to protect itself against

assault and thus maintain homeostasis. On the other hand, old GM resulted more enriched in

pathogenic bacteria, associated with dysbiosis. Bacterial pathogenicity in the gut is associated

with the accumulation of mutations over time, induced by the failed capacity to repair them

(Leimbach et al., 2013). Bacterial communities present in old fish guts were indeed more

pathogenic and our functional metagenome analysis associated them with host disease.

Consistently, while young gut status was associated with high expression of genes involved in

cell cycle activity, likely associated with proliferation and differentiation, old gut status was

associated with host immune responses to pathogenic bacteria, reflecting the prevalence of

more pathogenic bacterial taxa.

Although GM transfers from young, healthy donors have found applications in the clinic to

treat acute gut infections such as those associated with *Clostridium difficile* (Lee et al., 2016)

and have been proposed to treat obesity, metabolic syndrome and even neurodegenerative

diseases (Marotz and Zarrinpar, 2016; Xu et al., 2015), the application of this methodology as

an anti-aging intervention has not been explored to date. Remarkably, despite single

associations of different bacterial diets have shown to significantly affect life span in

invertebrate model systems such as *Caenorhabditis elegans* (Zhao et al., 2013), a functional

test of the role of a complex GM community associated with young age as an intervention

aimed at modulating the recipient's life span has not yet been carried out to date. By acutely

exposing middle age individuals to young fish GM content – after antibiotic treatment – we

could prolong life span and retard the age-dependent decline in exploratory behavior.

Noteworthy, our results exclude that the effects of the interventions depend on repopulating

the intestine with any GM community or that antibiotic treatment alone was sufficient to

explain the full extent of life span increase achieved via transfer of young GM.

Additionally, life span was not affected in young fish exposed to GM from old, young, and

sham control fish after antibiotic treatment. These results are compatible with a scenario where

the age-associated decline of immune function might be responsible for the progressively

decreased capacity of the host to i) maintain the healthy portion of the GM community and ii)

counteract the proliferation of potentially pathogenic gut bacteria.

Fish treated with young GM after depletion of their own resident GM community not only

maintained a more diverse microbial community at old age compared to wild-type, age

matched control fish, but their microbial community remained more similar to that of young

fish. This raises the possibility that bacterial consortia associated with young fish can

contribute to increased life span and enhanced individual health status. Based on functional

metagenomic analysis, young fish and fish treated with young GM were enriched for bacteria

associated with carbohydrate metabolism and DNA repair, both importantly associated with

host metabolism, health and longevity.

Young fish, as well as fish treated with young GM, had a high number of bacterial taxa that

frequently co-occurred with one another, de facto contributing to a young-associated bacterial

network. On the other end, old wild-type controls (16wk), as well as old fish treated with same

age GM (Omt), had smaller bacterial networks, possibly associated with the higher inter-

individual variation in GM composition associated with these groups. Remarkably, applying

our analyses to a published mouse cohort (Langille et al., 2014), we extended this finding to

mammals, confirming that networks built on GM OTU abundance are associated with host's

chronological age. Our network analysis enabled us to identify a subset of highly frequent taxa

associated with a young-like status and with prolonged life span in fish treated with young

GM. These involved the genera *Exiguobacterium*, *Planococcus*, *Propionigenium* and

Psychrobacter, which are key bacterial genera responsible for structuring a healthy GM

community in TK. Interestingly, species belonging to each of these genera have been

associated with energy metabolism and potential health benefits. Specifically, species of

Exiguobacterium and *Propionigenium* are able to metabolize cellulose and ferment

carbohydrates to produce short chain fatty acids, which are known anti-inflammatory

mediators and can modulate the immune system. *Planococcus* species can hydrolyze gelatin to

produce essential amino acids for use by the host and certain *Psychrobacter* species are

capable of producing omega-fatty acids. Taken together, these key bacterial genera can

produce metabolites capable of maintaining immune system health and having anti-

inflammatory effects on the host, both of which have been associated with longevity.

While GM transfers significantly affected the GM composition of experimental fish, their

overall gut transcriptional profile showed that old fish clustered together regardless of the

treatment. This could be due to the lack of transcriptional programs associated with growth in

16-week-old fish. However, transcripts involved in defense against pathogens, extracellular

matrix components and the Tor pathway, are dramatically different among experimental

groups receiving young or same-age gut content, suggesting that these key aspects might

ultimately be fundamental modulators of organismal life span and health. Generating a

correlation matrix between bacterial abundance and gut transcripts, we could isolate host

transcripts whose expression was significantly correlated with specific OTUs. In particular,

bacterial genera associated with a healthier and longer life span, such as *Psychrobacter* and

Exiguobacterium, were strongly associated with genes importantly associated with aging

modulation.

Our results indicate that improving the ecological diversity of the GM in old individuals helps

to restore health and prolongs life span. Our approach could provide a key to slowing aging

and retarding the onset of age-associated diseases by specifically targeting the GM. Given its

large bacterial taxonomic diversity and the shortest life span for a vertebrate species raised in

captivity, the TK could become a key experimental species which will help to shed light on

the functional connection between GM dynamics and aging in vertebrates.

MATERIALS AND METHODS

Wild fish samples

The wild fish samples derived from Zimbabwe were collected during the expedition in Gonarezhou National Park in 2015 (Permit No.: 23(1) (C) (II) 30/2015). Intestines were collected at each location and preserved in pure ethanol. Sampling locations coordinates are listed in Supplementary file 2.

Fish husbandry and survival scoring

Fish used for microbiota analysis and scored for survival were individually housed from week 4 post-hatching in single 2.8L tanks connected to a water recirculation system receiving 12h of light and 12hr of dark every day. Water temperature was set to 28˚C and fish were fed blood worm larvae and brine shrimp nauplii twice a day during the week and once a day during the weekend. Dead fish were removed daily from the tanks, weighed and stored in 95% ethanol.

DNA extraction, 16S rRNA gene amplification and sequencing

All dissected intestinal samples were collected at the same time prior to morning feeding and flash frozen in liquid nitrogen. For DNA isolation, frozen intestines were placed into autoclaved 2ml screw caps tubes containing 1ml of lysis buffer (80mM EDTA, 200mM Tris (pH 8.0) and 0.1M NaCl in PBS) and 0.4g of a mixture of 01.mm zirconia/silica and 1.4mm stainless steel beads (Biospec Products). Samples were bead-beaten for 3 minutes at 30 Hz (TissueLyzer II, Qiagen). Following the bead beating step, SDS (10% final concentration) and RNase A (PureLink, Invitrogen) were added and samples were incubated for 30 minutes at 55°C. DNA was then isolated using phenol:chloroform:isoamyl alcohol (Invitrogen) as per manufacturer's instructions with an additional chloroform step to remove excess phenol.

DNA was then used in one of two, two-step PCR methods designed to target the V3-V4 region

of the 16 rRNA gene (Klindworth et al., 2013). For the first method, initial primers consisted

of (5' to 3') a primer pad and linker (Caporaso et al., 2012) and the V3/V4 gene specific

forward or reverse primer sequences. The second step PCR used primers complementary for

the primer pad and linker followed by standard Illumina adaptors. The reverse primer for the

second step also contained a 12bp Golay barcode (Caporaso et al., 2012). For the second

method, initial primers consisted of (5' to 3') Illumina overhang adaptor sequences and the

V3/V4 gene specific primer sequences followed by a second step PCR using the Nextera XT

Index kit (Illumina). For both amplification methods cycling conditions were the same and the

first round of PCR was performed in triplicate with approximately equal amounts of DNA

template (250ng/reaction). PCR reactions were carried out with a two-minute denaturation

step at 98°C, followed by 25 cycles of 98°C for 30 seconds, 55°C for 30 seconds and 72°C for

30 seconds. Triplicate reactions were then pooled and cleaned using the Wizard SV Gel and

PCR Clean-up Kit (promega). The cycling conditions for the second step PCR were the same

as the first step, except annealing was performed at 60°C with only 8 cycles. Both PCR steps

used KAPA HiFi Hotstart ReadyMix (KAPA Biosystems) and 1μm of primers in 25μl total

volume. 2nd step PCR products were run on a 1.2% agarose gel and DNA products between

500-700 base pairs were excised and cleaned up as in step 1. PCR products were quantified by

Qubit (Life Technologies), diluted to 4nm and combined in equal volumes. The combined

amplicon libraries were then sequenced on the Illumina MiSeq, V3 reagents, 2x 300bp paired-

end reads.

Antibiotic treatment and fish microbiota transfer

Microbiota transfer experiments were based on those developed in zebrafish (Pham et al.,

2008). Recipient fish were removed from main water recirculating system and housed in 9L

tanks at a density of 10 fish per tank. Recipient fish (6wk or 9.5wk) were treated overnight

with a combination of Vancomycin (0.01g/L), Metronidazole (0.5g/L), Neomycin (0.5g/L)

and Ampicillin (0.5g/L) to diminish the resident bacterial community. Following antibiotic

treatment, recipient fish were washed twice for 10 minutes with autoclaved tank water.

Concurrently, whole intestines were isolated from donor fish and placed into 10 cm petri

dishes containing sterile PBS on ice. Intestines were then opened longitudinally, the intestinal

contents scraped out and then further cut into 0.5 cm pieces to facilitate the release of bacteria.

The collected intestinal contents were washed once in cold PBS and added to the fish tanks

containing autoclaved tank water and recipient fish at a ratio of 1 donor fish intestine/2

recipient fish. Fish were incubated overnight with the donor fish intestinal contents before

being returned to the main recirculating system and individually housed, where they were

regularly fed according to standard husbandry.

Microbial community analysis

Fastq files from paired end reads were joined, demultiplexed and subjected to quality filtering

with QIIME 1.8 (Q≥20) as previously described (Caporaso et al., 2012). For microbial

community analysis, QIIME was used to identify OTUs by open-reference picking using

UCLUST (97% similarity) and taxonomy was assigned with the Greengenes 13.8 database

(DeSantis et al., 2006; Edgar, 2010). A minimum OTU count of 5 was used to minimize

spurious OTUs. Representative OTU sequences were aligned with PYNAST (Caporaso et al.,

2010a) and FastTree 2 (Price et al., 2010) was used to build a phylogenetic tree. For diversity

analyses, OTU tables were rarefied to at least 5000 sequences, which allowed the majority of

samples to be kept. Qiime and R were used to calculate alpha and beta diversity metrics and

generate plots. To identify bacteria associated with specific groups, OTU tables were further

filtered for presence in at least 25% of samples with a collective abundance of greater than 100

reads. Significant changes in relative OTU abundance were identified with linear discriminant analysis effect size (LEfSe) and visualized using GraPhlAn (Asnicar et al., 2015). For metagenomics analysis, rarefied OTU tables were generated by closed reference picking and the PICRUSt tool was used to normalize by 16S copy number and predict the metagenome content of samples from 16S rRNA profiles. KEGG pathway functions were then categorized at level 2 or 3 and LEfSe was used to identify significant changes among classes. The accuracy of PICRUSt predictions was determined by nearest sequenced taxon index (NSTI).

Bacterial diversity among model organisms

To compare the overall bacterial diversity among different species, sequences were downloaded from the European nucleotide archive (ENA), the NCBI Short Read Archive (SRA) or MG-RAST and subjected to closed reference OTU picking. Taxa summary and alpha diversity measures were computed using QIIME 1.8. Samples were chosen which used Illumina sequencing of the V4 region of the 16 rRNA gene. Accession numbers. Human: ERR561021UK, ERR560915UK, ERR560902UK, ERR560855UK, mgm4489670, mgm4489628, mgm4489516, mgm4489454, mgm4538473, mgm4538468, mgm4538264, mgm4538259, mgm4538238. Mouse: SRR1820108, SRR1820074, SRR1820073, SRR1820072, SRR1820071, ERR706143, ERR706142, ERR706141. Zebrafish: SRR1581750, SRR1581753, SRR1581759, SRR1581763, SRR1581766, SRR1581889, SRR1581890, SRR1581891. Drosophila: SRR989472, SRR989473, SRR989474, SRR989469, SRR989467, SRR952981.

Bacterial load quantification

To determine bacterial load, DNA was isolated from freshly collected faecal pellets as described above. Real-time quantitative PCR was performed with DyNAmo Color Flash

705   SYBR green master mix (Thermo Scientific) and run with the BioRad CFX384 Real-Time

706   System. Samples were normalized to amount of input faecal material and graphed as 1/Ct

707   value. Primers targeting the 16s gene were described previously (Caporaso et al., 2010b). For:

708   5' TCCTACGGGAGGCAGCAGT 3' and Rev: 5' GGACTACCAGGGTATCTAATCCTGTT

709   3'.

Quantitative analysis of spontaneous locomotor activity

712   To measure total swimming distance, fish were placed in rectangular (16 x 90 cm) tanks filled

713   with 8 liters of water and allowed to acclimate for 30 minutes. The room and water

714   temperature and water quality were kept as similar to the fish's' normal environment as

715   possible. The fish were then filmed for 20 minutes with an overhead mounted camera and data

716   were analyzed using EthoVision XT11 (Noldus Information Technologies).

Co-occurrence networks

The OTU table was filtered by a minimum total count of 5509 per individual, based on the

rarefaction value (see Microbial community analysis). Subsequently, the table was divided

into subsets corresponding to the treatment groups. Normalization was achieved through

dividing the OTU sample count by the total sample count, scaled to 1000.

Co-occurrence networks were produced by applying the SparCC program (Friedman and Alm,

2012), a network inference tool specifically developed for analysis of correlations in

compositional data, such as 16S microbiome analyses. In order to avoid unreliable correlations

for very rare OTUs, the OTU tables were filtered for bacteria that were present in at least 25%

of samples of each group.

The SparCC pipeline was used to first calculate OTU-OTU correlations averaged over 20

iterations (Friedman and Alm, 2012). We then tested the significance of these correlations by

computing pseudo-p-values against 1000 bootstrap simulations, applying the same parameters.

All OTU-OTU correlations with a p-value lower than 0.05 were considered significant and

were included into network preparation. Nodes without any edges after filtering by significant

p-values were removed from the network. To analyze the network properties we used the

igraph package in R, version 1.0.1 (Csardi and Nepusz, 2006). For each group we generated an

undirected network, weighted by correlation magnitude. For biological interpretation we only

focused on positive correlations. To identify network size and composition, we calculated the

negative cumulative degree distribution and the percentage of genus present in the network

belonging to different phyla. The main clusters of each network were identified with two

different clustering algorithms of the igraph package (Csardi and Nepusz, 2006). The first

algorithm 'k-core' clusters based on the degree, with each member of the maximal subgraph

has at least a degree count of k. The second algorithm 'infomap community' searches for

community structures that minimize in the length of a random walker trajectory. We used the

overlap of the largest clusters within the two cluster algorithms to identify the main clusters in

each group. The members of clusters from young fish (wk6) and old fish (wk16) were used to

identify young-like or old-like clusters in the main clusters of the transfer groups (Ymt, Omt

and Abx).

In order to compare our findings to a different model organism, we used the data set from

Langille et al. (2014) (Langille et al., 2014). We scaled the relative frequencies from young,

middle and old mice to 1000 and excluded bacteria not present in at least 25% of samples.

Additionally, we excluded sample Y7.August14, because we included the same sample from

15[th] of August (Y7.August15). Identically to the previous analysis we calculated the negative

cumulative degree distribution.

RNA extraction

Intestines from four young (6 weeks old), four old (16 weeks old), four old received six-week-old donor fish gut content (Ymt) and four old fish received 9.5-week-old fish gut content (Omt) fish were dissected. Trizol (15596018, Thermo Fisher Scientific, USA) was used to isolate total RNA, following manufacturer's instructions. Residual genomic DNA was removed by DNaseI treatment (AM2222, Thermo Fisher Scientific, USA) and 1.5 μg of total RNA was used as a template for reverse transcription (11754050, Thermo Fisher Scientific, USA).

RNA sequencing

We sequenced four individuals of groups 6wk, 16wk, Ymt and Omt. Each sample sequenced with 100bp paired end with 30 million reads. All samples were mapped to the African Turquoise Killifish Genome (Valenzano et al., 2015) using STAR version 2.4.1c (Dobin et al., 2013) with the default parameters, except for splice junction and chimeric junction parameters set to 15 base-pairs, as well as allowing for only 5 mismatches over the whole fragment and removing all reads that map to more than two locations.

Differential gene expression

The differential gene expression was performed using the Cufflinks suite (Trapnell et al., 2010) version 2.2.1 and the edgeR package (Robinson et al., 2010) version 3.16.5. cuffquant was used to generate feature abundances using the mapped reads and the annotation file from (Valenzano et al., 2015) for each file independently. Subsequently cuffnorm was used to obtain normalized feature counts of all genes of samples. The edgeR package was used to identify differentially expressed genes between any of the groups resulting in six pairwise comparisons. Additionally, we calculated the differentially expressed genes between the old

microbiome group (Omt and 16wk) and young microbiome group (Ymt and 6wk), as well for

the comparison of normal-lived group (Omt and 16wk) versus long-lived group (Ymt). For

these comparisons we excluded genes that were differentially expressed within pairwise

comparisons of each combined group. All edgeR runs were performed using TMM

normalization, followed by tagwise dispersion and an exact test (Robinson et al., 2010). We

considered only genes that had a counts-per-million value above 0.7 in at least 50% of

compared samples. The resulting p-values were multiple testing corrected with the Benjamini-

Hochberg method. Only genes with a FDR smaller or equal than 0.1 were considered as

differentially expressed. Gene enrichment analyses were performed using the GOrilla software

(Eden et al., 2009).

Functional analysis of target genes

To gain insights into the function of genes differing between young and old fish as well as

genes differing between fish of long or normal lifespan we used the GOrilla (Eden et al., 2009)

to perform enrichment tests. We used the zebrafish database for annotation. Therefore we used

the available reciprocal blast information (Valenzano et al., 2015). If top differentially

expressed genes were not found to have a zebrafish ortholog in this resource, we used blastp

(Altschul et al., 1990) to identify the corresponding zebrafish ortholog (Table S7). First all

genes were mapped to a GO-Term. The statistical enrichment test uses a hypergeometric

distribution to test for enrichment in the ratios of target genes associated with a GO-Term, and

the ratio of background genes associated with the same GO-Term to identify pathways or GO-

Terms that have a significantly larger amount of target genes than expected. Here, we used all

genes that were found to be expressed in the intestine of the experimental groups as

background for all analysis. The number of genes differentially expressed between long and

normal-lifespan fish was too low to reach significance for any GO-Term after adjustment for

multiple hypothesis testing (Bonferroni correction). The GOrilla results were visualized with

the ggplot2 package in R.

Correlation between individual OTU and gut transcripts

Individual fish OTU tables and log2-transformed transcripts were used as a source to identify

significant correlations using Pearson's correlation.

The python code used to perform this analysis was saved on github:

https://github.com/dvalenzano/uBiome_Network/blob/master/Notes_20-04-2016.ipynb

AUTHOR CONTRIBUTIONS

P.S. and D.R.V. conceived the study and planned all the experiments. P.S. performed all the

experiments. P.S., D.W., M.L.P. and D.R.V. performed the statistical analyses. D.W. and F.M.

analyzed the RNA-Seq data, D.W. performed the network analysis. M.L.P analyzed the wild

fish GM data and helped with the network analysis. E.G. provided intellectual expertise and

logistic support for the fielwork. D.R.V. and D.W. conducted the fieldwork. M.R. contributed

with the wild specimens collection. D.R.V. designed, coordinated the study and wrote the

manuscript.

ACKNOWLEDGMENTS

We thank the Cologne Center for Genomics (CCG) and Paul Higgins for technical support with the sequencing, all the members of the Valenzano lab for their scientific input to the project. We are thankful to Adam Antebi and Linda Partridge for discussions, Anne Brunet, Anthony Zannas and Jenny Regan for critically reading the manuscript. We thank the Research Council of Zimbabwe, Zimbabwe Parks and Wildlife Management Authoriry, Patience Gandiwa, Daphine Madhlamoto, Itamar Harel, Radim Blažek, Matej Polačik, Tamuka Nhiwatiwa, Hugo and Elsabe van der Westhuizen, Evious Mpofu, Power Mupunga and all the rangers of the Gonarezhou National Park for their support with the fieldwork. All animal experiments of this study have been examined and approved by the competent authority (Landesamt für Natur, Umwelt und Verbraucherschutz Nordrhein-Westfalen; AZ: 84-02.04.2015.A377).

Fieldwork was supported by the Max Planck Society, by the Max Planck Institute for Biology of Ageing and by the CSF project 16–00291S. The experimental part was supported by the Max Planck Society and by the Max Planck Institute for Biology of Ageing.

843 **COMPETING INTEREST**

844 There are no financial and non-financial competing interests for all the authors.

REFERENCES

Agler, M.T., Ruhe, J., Kroll, S., Morhenn, C., Kim, S.T., Weigel, D., and Kemen, E.M. (2016). Microbial Hub Taxa Link Host and Abiotic Factors to Plant Microbiome Variation. PLoS Biol *14*, e1002352.

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990). Basic local alignment search tool. J Mol Biol *215*, 403-410.

Asnicar, F., Weingart, G., Tickle, T.L., Huttenhower, C., and Segata, N. (2015). Compact graphical representation of phylogenetic data and metadata with GraPhlAn. PeerJ *3*, e1029.

Backhed, F., Fraser, C.M., Ringel, Y., Sanders, M.E., Sartor, R.B., Sherman, P.M., Versalovic, J., Young, V., and Finlay, B.B. (2012). Defining a healthy human gut microbiome: current concepts, future directions, and clinical applications. Cell Host Microbe *12*, 611-622.

Baumgart, D.C., and Carding, S.R. (2007). Inflammatory bowel disease: cause and immunobiology. Lancet *369*, 1627-1640.

Biagi, E., Franceschi, C., Rampelli, S., Severgnini, M., Ostan, R., Turroni, S., Consolandi, C., Quercia, S., Scurti, M., Monti, D.*, et al.* (2016). Gut Microbiota and Extreme Longevity. Curr Biol *26*, 1480-1485.

Blažek, R., Polačik, M., Kačer, P., Cellerino, A., Řežucha, R., Methling, C., Tomášek, O., Syslová, K., Terzibasi Tozzini, E., Albrecht, T.*, et al.* (2017). Repeated intra-specific divergence in lifespan and ageing of African annual fishes along an aridity gradient. . Evolution.

Buchon, N., Broderick, N.A., and Lemaitre, B. (2013). Gut homeostasis in a microbial world: insights from Drosophila melanogaster. Nat Rev Microbiol *11*, 615-626.

Cabreiro, F., and Gems, D. (2013). Worms need microbes too: microbiota, health and aging in Caenorhabditis elegans. EMBO Mol Med *5*, 1300-1310.

Caporaso, J.G., Bittinger, K., Bushman, F.D., DeSantis, T.Z., Andersen, G.L., and Knight, R. (2010a). PyNAST: a flexible tool for aligning sequences to a template alignment. Bioinformatics *26*, 266-267.

Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Pena, A.G., Goodrich, J.K., Gordon, J.I.*, et al.* (2010b). QIIME allows analysis of high-throughput community sequencing data. Nat Methods *7*, 335-336.

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S.M., Betley, J., Fraser, L., Bauer, M.*, et al.* (2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J *6*, 1621-1624.

Cellerino, A., Valenzano, D.R., and Reichard, M. (2015). From the bush to the bench: the annual Nothobranchius fishes as a new model system in biology. Biol Rev Camb Philos Soc.

Cho, S., Roh, K., Park, J., Park, Y.S., Lee, M., Cho, S., Kil, E.J., Cho, M.J., Oh, J.S., Byun, H.S.*, et al.* (2017). Hydrolysis of Hyaluronic Acid in Lymphedematous Tissue Alleviates Fibrogenesis via TH1 Cell-Mediated Cytokine Expression. Sci Rep *7*, 35.

Claesson, M.J., Jeffery, I.B., Conde, S., Power, S.E., O'Connor, E.M., Cusack, S., Harris, H.M., Coakley, M., Lakshminarayanan, B., O'Sullivan, O.*, et al.* (2012). Gut microbiota composition correlates with diet and health in the elderly. Nature *488*, 178-184.

Clark, R.I., Salazar, A., Yamada, R., Fitz-Gibbon, S., Morselli, M., Alcaraz, J., Rana, A., Rera, M., Pellegrini, M., Ja, W.W.*, et al.* (2015). Distinct Shifts in Microbiota Composition during Drosophila Aging Impair Intestinal Function and Drive Mortality. Cell Rep *12*, 1656-1667.

Conti, B., Sanchez-Alavez, M., Winsky-Sommerer, R., Morale, M.C., Lucero, J., Brownell, S., Fabre, V., Huitron-Resendiz, S., Henriksen, S., Zorrilla, E.P.*, et al.* (2006). Transgenic mice with a reduced core body temperature have an increased life span. Science *314*, 825-828.

Cox, L.M., and Blaser, M.J. (2015). Antibiotics in early life and obesity. Nat Rev Endocrinol
11, 182-190.

Csardi, G., and Nepusz, T. (2006). The igraph software package for complex network
research. InterJournal, Complex Systems *1695*.

DeSantis, T.Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E.L., Keller, K., Huber, T.,
Dalevi, D., Hu, P., and Andersen, G.L. (2006). Greengenes, a chimera-checked 16S rRNA
gene database and workbench compatible with ARB. Appl Environ Microb *72*, 5069-5072.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*,
15-21.

Dodin, M., and Katz, D.E. (2014). Faecal microbiota transplantation for Clostridium difficile
infection. Int J Clin Pract *68*, 363-368.

Eden, E., Navon, R., Steinfeld, I., Lipson, D., and Yakhini, Z. (2009). GOrilla: a tool for
discovery and visualization of enriched GO terms in ranked gene lists. BMC Bioinformatics
10, 48.

Edgar, R.C. (2010). Search and clustering orders of magnitude faster than BLAST.
Bioinformatics *26*, 2460-2461.

Fontana, L., Partridge, L., and Longo, V.D. (2010). Extending healthy life span--from yeast to
humans. Science *328*, 321-326.

Friedman, J., and Alm, E.J. (2012). Inferring Correlation Networks from Genomic Survey
Data. Plos Comput Biol *8*.

Garrett, W.S. (2015). Cancer and the microbiota. Science *348*, 80-86.

Genade, T., Benedetti, M., Terzibasi, E., Roncaglia, P., Valenzano, D.R., Cattaneo, A., and
Cellerino, A. (2005). Annual fishes of the genus Nothobranchius as a model system for aging
research. Aging Cell *4*, 223-233.

Geva-Zatorsky, N., Sefik, E., Kua, L., Pasman, L., Tan, T.G., Ortiz-Lopez, A., Yanortsang,
T.B., Yang, L., Jupp, R., Mathis, D.*, et al.* (2017). Mining the Human Gut Microbiota for
Immunomodulatory Organisms. Cell *168*, 928-943 e911.

Guo, L., Karpac, J., Tran, S.L., and Jasper, H. (2014). PGRP-SC2 promotes gut immune
homeostasis to limit commensal dysbiosis and extend lifespan. Cell *156*, 109-122.

Harel, I., Benayoun, B.A., Machado, B., Singh, P.P., Hu, C.K., Pech, M.F., Valenzano, D.R.,
Zhang, E., Sharp, S.C., Artandi, S.E.*, et al.* (2015). A platform for rapid exploration of aging
and diseases in a naturally short-lived vertebrate. Cell *160*, 1013-1026.

Harel, I., and Brunet, A. (2015). The African Turquoise Killifish: A Model for Exploring
Vertebrate Aging and Diseases in the Fast Lane. Cold Spring Harb Symp Quant Biol.

Josenhans, C., and Suerbaum, S. (2002). The role of motility as a virulence factor in bacteria.
Int J Med Microbiol *291*, 605-614.

Kamada, N., Chen, G.Y., Inohara, N., and Nunez, G. (2013). Control of pathogens and
pathobionts by the gut microbiota. Nat Immunol *14*, 685-690.

Kapahi, P., Chen, D., Rogers, A.N., Katewa, S.D., Li, P.W.L., Thomas, E.L., and Kockel, L.
(2010). With TOR, Less Is More: A Key Role for the Conserved Nutrient-Sensing TOR
Pathway in Aging. Cell Metabolism *11*, 453-465.

Kenyon, C., Chang, J., Gensch, E., Rudner, A., and Tabtiang, R. (1993). A C. elegans mutant
that lives twice as long as wild type. Nature *366*, 461-464.

Kim, Y., Nam, H.G., and Valenzano, D.R. (2016). The short-lived African turquoise killifish:
an emerging experimental model for ageing. Dis Model Mech *9*, 115-129.

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., and Glockner, F.O.
(2013). Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-
generation sequencing-based diversity studies. Nucleic Acids Res *41*, e1.

Kootte, R.S., Vrieze, A., Holleman, F., Dallinga-Thie, G.M., Zoetendal, E.G., de Vos, W.M., Groen, A.K., Hoekstra, J.B., Stroes, E.S., and Nieuwdorp, M. (2012). The therapeutic potential of manipulating gut microbiota in obesity and type 2 diabetes mellitus. Diabetes Obes Metab *14*, 112-120.

Kostic, A.D., Howitt, M.R., and Garrett, W.S. (2013). Exploring host-microbiota interactions in animal models and humans. Genes Dev *27*, 701-718.

Kunde, S., Pham, A., Bonczyk, S., Crumb, T., Duba, M., Conrad, H., Jr., Cloney, D., and Kugathasan, S. (2013). Safety, tolerability, and clinical response after fecal transplantation in children and young adults with ulcerative colitis. J Pediatr Gastroenterol Nutr *56*, 597-601.

Langille, M.G., Meehan, C.J., Koenig, J.E., Dhanani, A.S., Rose, R.A., Howlett, S.E., and Beiko, R.G. (2014). Microbial shifts in the aging mouse gut. Microbiome *2*, 50.

Langille, M.G., Zaneveld, J., Caporaso, J.G., McDonald, D., Knights, D., Reyes, J.A., Clemente, J.C., Burkepile, D.E., Vega Thurber, R.L., Knight, R.*, et al.* (2013). Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. Nat Biotechnol *31*, 814-821.

Lapierre, L.R., and Hansen, M. (2012). Lessons from C. elegans: signaling pathways for longevity. Trends Endocrinol Metab *23*, 637-644.

Lee, C.H., Steiner, T., Petrof, E.O., Smieja, M., Roscoe, D., Nematallah, A., Weese, J.S., Collins, S., Moayyedi, P., Crowther, M.*, et al.* (2016). Frozen vs Fresh Fecal Microbiota Transplantation and Clinical Resolution of Diarrhea in Patients With Recurrent Clostridium difficile Infection: A Randomized Clinical Trial. JAMA *315*, 142-149.

Leimbach, A., Hacker, J., and Dobrindt, U. (2013). E. coli as an all-rounder: the thin line between commensalism and pathogenicity. Curr Top Microbiol Immunol *358*, 3-32.

Li, H., Qi, Y., and Jasper, H. (2016). Preventing Age-Related Decline of Gut Compartmentalization Limits Microbiota Dysbiosis and Extends Lifespan. Cell Host Microbe *19*, 240-253.

Mair, W., and Dillin, A. (2008). Aging and survival: the genetics of life span extension by dietary restriction. Annu Rev Biochem *77*, 727-754.

Marotz, C.A., and Zarrinpar, A. (2016). Treating Obesity and Metabolic Syndrome with Fecal Microbiota Transplantation. Yale J Biol Med *89*, 383-388.

Miquel, J., Lundgren, P.R., Bensch, K.G., and Atlan, H. (1976). Effects of temperature on the life span, vitality and fine structure of Drosophila melanogaster. Mech Ageing Dev *5*, 347-370.

Neis, E.P., Dejong, C.H., and Rensen, S.S. (2015). The role of microbial amino acid metabolism in host metabolism. Nutrients *7*, 2930-2946.

Nezhybová, V., Reichard, M., Blažek, R., and Ondračková, M. (2017). Metazoan parasites of African annual killifish (Nothobranchiidae): abundance, diversity and their environmental correlates. . Biotropica.

Nicholson, J.K., Holmes, E., Kinross, J., Burcelin, R., Gibson, G., Jia, W., and Pettersson, S. (2012). Host-gut microbiota metabolic interactions. Science *336*, 1262-1267.

O'Toole, P.W., and Jeffery, I.B. (2015). Gut microbiota and aging. Science *350*, 1214-1215.

Pham, L.N., Kanther, M., Semova, I., and Rawls, J.F. (2008). Methods for generating and colonizing gnotobiotic zebrafish. Nat Protoc *3*, 1862-1875.

Price, M.N., Dehal, P.S., and Arkin, A.P. (2010). FastTree 2--approximately maximum-likelihood trees for large alignments. PLoS One *5*, e9490.

Qin, J., Li, R., Raes, J., Arumugam, M., Burgdorf, K.S., Manichanh, C., Nielsen, T., Pons, N., Levenez, F., Yamada, T.*, et al.* (2010). A human gut microbial gene catalogue established by metagenomic sequencing. Nature *464*, 59-65.

Rampelli, S., Candela, M., Turroni, S., Biagi, E., Collino, S., Franceschi, C., O'Toole, P.W., and Brigidi, P. (2013). Functional metagenomic profiling of intestinal microbiome in extreme ageing. Aging (Albany NY) *5*, 902-912.

Reichard, M., Janáč, M., Polačik, M., Blažek, R., and Vrtílek, M. (2017). Community assembly in Nothobranchius annual fishes: nested patterns, environmental niche and biogeographic history. Ecology and Evolution.

Reichwald, K., Petzold, A., Koch, P., Downie, B.R., Hartmann, N., Pietsch, S., Baumgart, M., Chalopin, D., Felder, M., Bens, M.*, et al.* (2015). Insights into Sex Chromosome Evolution and Aging from the Genome of a Short-Lived Fish. Cell *163*, 1527-1538.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics *26*, 139-140.

Roeselers, G., Mittge, E.K., Stephens, W.Z., Parichy, D.M., Cavanaugh, C.M., Guillemin, K., and Rawls, J.F. (2011). Evidence for a core gut microbiota in the zebrafish. ISME J *5*, 1595-1608.

Sampson, T.R., Debelius, J.W., Thron, T., Janssen, S., Shastri, G.G., Ilhan, Z.E., Challis, C., Schretter, C.E., Rocha, S., Gradinaru, V.*, et al.* (2016). Gut Microbiota Regulate Motor Deficits and Neuroinflammation in a Model of Parkinson's Disease. Cell *167*, 1469-1480 e1412.

Schuijt, T.J., Lankelma, J.M., Scicluna, B.P., de Sousa e Melo, F., Roelofs, J.J., de Boer, J.D., Hoogendijk, A.J., de Beer, R., de Vos, A., Belzer, C.*, et al.* (2016). The gut microbiota plays a protective role in the host defence against pneumococcal pneumonia. Gut *65*, 575-583.

Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S., and Huttenhower, C. (2011). Metagenomic biomarker discovery and explanation. Genome Biol *12*, R60.

Semova, I., Carten, J.D., Stombaugh, J., Mackey, L.C., Knight, R., Farber, S.A., and Rawls, J.F. (2012). Microbiota regulate intestinal absorption and metabolism of fatty acids in the zebrafish. Cell Host Microbe *12*, 277-288.

Sokol, H., Pigneur, B., Watterlot, L., Lakhdari, O., Bermudez-Humaran, L.G., Gratadoux, J.J., Blugeon, S., Bridonneau, C., Furet, J.P., Corthier, G.*, et al.* (2008). Faecalibacterium prausnitzii is an anti-inflammatory commensal bacterium identified by gut microbiota analysis of Crohn disease patients. Proc Natl Acad Sci U S A *105*, 16731-16736.

Sommer, F., and Backhed, F. (2013). The gut microbiota--masters of host development and physiology. Nat Rev Microbiol *11*, 227-238.

Steffen, K.K., MacKay, V.L., Kerr, E.O., Tsuchiya, M., Hu, D., Fox, L.A., Dang, N., Johnston, E.D., Oakes, J.A., Tchao, B.N.*, et al.* (2008). Yeast life span extension by depletion of 60s ribosomal subunits is mediated by Gcn4. Cell *133*, 292-302.

Tian, X., Azpurua, J., Hine, C., Vaidya, A., Myakishev-Rempel, M., Ablaeva, J., Mao, Z., Nevo, E., Gorbunova, V., and Seluanov, A. (2013). High-molecular-mass hyaluronan mediates the cancer resistance of the naked mole rat. Nature *499*, 346-349.

Trapnell, C., Williams, B.A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M.J., Salzberg, S.L., Wold, B.J., and Pachter, L. (2010). Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. Nat Biotechnol *28*, 511-515.

Turnbaugh, P.J., Ley, R.E., Mahowald, M.A., Magrini, V., Mardis, E.R., and Gordon, J.I. (2006). An obesity-associated gut microbiome with increased capacity for energy harvest. Nature *444*, 1027-1031.

Valenzano, D.R., Benayoun, B.A., Singh, P.P., Zhang, E., Etter, P.D., Hu, C.K., Clement-Ziza, M., Willemsen, D., Cui, R., Harel, I.*, et al.* (2015). The African Turquoise Killifish Genome Provides Insights into Evolution and Genetic Architecture of Lifespan. Cell *163*, 1539-1554.

Valenzano, D.R., Sharp, S., and Brunet, A. (2011). Transposon-Mediated Transgenesis in the
Short-Lived African Killifish Nothobranchius furzeri, a Vertebrate Model for Aging. G3
(Bethesda) *1*, 531-538.
Valenzano, D.R., Terzibasi, E., Cattaneo, A., Domenici, L., and Cellerino, A. (2006a).
Temperature affects longevity and age-related locomotor and cognitive decay in the short-
lived fish Nothobranchius furzeri. Aging Cell *5*, 275-278.
Valenzano, D.R., Terzibasi, E., Genade, T., Cattaneo, A., Domenici, L., and Cellerino, A.
(2006b). Resveratrol prolongs lifespan and retards the onset of age-related markers in a short-
lived vertebrate. Curr Biol *16*, 296-300.
Van Voorhies, W.A., and Ward, S. (1999). Genetic and environmental conditions that increase
longevity in Caenorhabditis elegans decrease metabolic rate. Proc Natl Acad Sci U S A *96*,
11399-11403.
Xu, M.Q., Cao, H.L., Wang, W.Q., Wang, S., Cao, X.C., Yan, F., and Wang, B.M. (2015).
Fecal microbiota transplantation broadening its application beyond intestinal disorders. World
J Gastroenterol *21*, 102-111.
Zac Stephens, W., Burns, A.R., Stagaman, K., Wong, S., Rawls, J.F., Guillemin, K., and
Bohannan, B.J. (2016). The composition of the zebrafish intestinal microbial community
varies across development. ISME J *10*, 644-654.
Zhao, Y., Zhao, L., Zheng, X., Fu, T., Guo, H., and Ren, F. (2013). Lactobacillus salivarius
strain FDB89 induced longevity in Caenorhabditis elegans by dietary restriction. J Microbiol
51, 183-188.
Zoetendal, E.G., Vaughan, E.E., and de Vos, W.M. (2006). A microbial world within us. Mol
Microbiol *59*, 1639-1650.
1065
1066