Title: Natural selection can favor ratchet robustness over mutational

robustness

Authors:

Yinghong Lan[1*], Aaron Trout[2], Daniel Michael Weinreich[1], Christopher Scott Wylie[1]

Institutional affiliations:

1: Department of Ecology and Evolutionary Biology, and Center for Computational

Molecular Biology, Brown University, Providence, RI, USA

2: Department of Mathematics, Chatham University, Pittsburgh, PA, USA

Corresponding author:

*: yinghong_lan@brown.edu

Abstract

The vast majority of fitness-affecting mutations are deleterious. How natural populations evolve

to cope is a question of fundamental interest. Previous studies have reported the evolution of

mutational robustness, that is, natural selection favoring populations with less deleterious

mutations. By definition, mutational robustness provides a short-term fitness advantage.

However, this overlooks the fact that mutational robustness decreases finite asexual populations'

ability to purge recurrent deleterious mutations. Thus, mutational robustness also results in

higher risk of long-term extinction by Muller's ratchet. Here, we explore the tension between

short- and long- term response to deleterious mutations. We first show that populations can resist

the ratchet if either the selection coefficient or the ratio of beneficial to deleterious mutations

increases as fitness declines. We designate these properties as ratchet robustness, which

fundamentally reflects a negative feedback between mutation rate and the tendency to

accumulate more mutations. We also find in simulations that populations can evolve ratchet

robustness when challenged by deleterious mutations. We conclude that mutational robustness

cannot be selected for in the long term, but it can be favored in the short-term, purely because of

temporary fitness advantage. We also discuss other potential causes of mutational robustness in

nature.

Keywords: natural selection; ratchet robustness; mutational robustness; fitness landscape;

Muller's ratchet.

Introduction

The vast majority of fitness-affecting mutations are deleterious. How natural populations evolve

to cope is a question of fundamental interest. One widespread hypothesis proposes the evolution

of mutational robustness, which would mean that deleterious mutations evolve to become less

deleterious(Krakauer and Plotkin 2001; Wilke and Adami 2003). Intuitively, it may seem that

organisms with higher mutational robustness will have a fitness advantage over those with lower

mutational robustness. Taking that intuition one step further, organisms with higher deleterious

mutation rate, i.e., more frequent occurrences of deleterious mutations, might be expected to

evolve greater mutational robustness. And indeed, there exists a substantial body of literature

consistent with that idea (**Box 1**).

Box 1: Literature on the evolution of mutational robustness

Two computational studies (van Nimwegen et al. 1999; Wilke et al. 2001) have addressed the

possibility of evolution of mutational robustness. Using Avida, a platform for conducting in

silico evolutionary experiments (Ofria et al. 2009), Wilke et al. (2001) evolved populations

founded with the same ancestor under high and low mutation rates. Then they competed the two

evolved lineages against each other under different mutation rates. They found that the lineage

evolved under high mutation rate, albeit having lower fitness, wins under higher mutation rate.

Their interpretation was that that the lineage adapted to high mutation rate has higher mutation

robustness. Two experimental studies using a viroid (Codoner et al. 2006) or an RNA virus

(Sanjuan et al. 2007) similarly found that in competition at high mutation rate, populations

evolved at high mutation rate displaced those evolved at low mutation rate.

Working in the neutral network framework, van Nimwegen et al. (1999) explored a model in

which mutations are either neutral or lethal. In this case, mutational robustness of each genotype

is measured by its proportion of neutral mutations. They found that as mutation rate increases,

populations will evolve to be composed of genotypes with fewer lethal mutations, i.e., with

higher mutational robustness.

However, despite the intuitive appeal of the foregoing, it overlooks important components of

population genetics theory. Firstly, in infinite populations, where genetic drift can be neglected,

equilibrium mean fitness depends only on deleterious mutation rate (Haldane 1937; Kimura and

Maruyama 1966). Thus, mutational robustness, i.e., the reduced effect of deleterious mutations,

does not confer any long-term fitness advantage. Moreover, finite asexual populations risk

Muller's ratchet (Muller 1964; Haigh 1978), meaning that deleterious mutations accumulate at a

rate so fast that selection is overwhelmed. In other words, besides the immediate fitness cost of

deleterious mutations, Muller's ratchet represents a longer-term challenge to evolving

populations. A previous study (Goyal et al. 2012) provides the quantitative parametric boundary

above which populations persist at Mutation-Selection-Drift Equilibrium (MSDE) (Wager and

Gabriel 1990; Schultz and Lynch 1997; Poon and Otto 2000), but below which they succumb to

the ratchet. In doing so, this study demonstrates that finite populations are less susceptible to

Muller's ratchet when mutation rate is lower, population size is greater, selection coefficient is

higher, or the ratio of beneficial to deleterious mutation rates is larger.

A fundamentally important question then arises as to how populations evolve in the face of both

short- and long- term challenges brought by deleterious mutations? Trivially, the evolution of

lower mutation rate will alleviate both concerns. Therefore, throughout this paper we hold

mutation rate as an extrinsic factor and observe other responses of evolving populations.

Furthermore, we hold population size constant, i.e., we assume soft selection. This approach is

conservative, because decreasing population size renders the population more prone to Muller's

ratchet, ramifications of which have already been studied in the context of mutational meltdown

(Lynch et al. 1993; Bull et al. 2005). Consequently, we focus only on the influence of selection

coefficient and the ratio of beneficial to deleterious mutation rates. As a complement to the

phrase "mutational robustness", we correspondingly designate increases in either of these

properties as conferring "ratchet robustness." (Recently Labar and Adami 2016 introduced the

phrase drift robustness for a similar phenomenon. However, we prefer ratchet robustness, since it

more precisely describes the underlying population genetic risk against which robustness is

conferred.)

The fitness landscape provides a unifying framework for exploring both mutational robustness and ratchet robustness. First define genotype space, where spatially adjacent genotypes are mutationally adjacent, and then project the fitness of each genotype over the genotype space. Since mutational robustness means deleterious mutations have smaller selection coefficients, regions of the landscape with higher mutational robustness are "flatter". By contrast, "steeper" regions of the landscape have higher ratchet robustness, since selection coefficients are larger. These considerations seem to suggest an intrinsic tension between mutational robustness and ratchet robustness. Ratchet robustness can also be realized by increased ratio of beneficial to deleterious mutation rates, represented on the fitness landscape by the increased probability of going "uphill" rather than "downhill" when locally exploring the genotype space via mutations (Goyal et al. 2012).

The goal of this paper is to identify those features of the fitness landscape that are favored by natural selection as a consequence of the fact that most mutations are deleterious. We investigate, in isolation, the effect of selection coefficient, and the influence of ratio of beneficial to deleterious mutation rates. We find that ratchet robustness protects populations from extinction due to deleterious mutations and should therefore be favored in the long term, although in the short term mutational robustness can also be selected for.

Methods

Evolutionary model

For all simulations, we implement discrete-time Wright-Fisher evolutionary model with custom

Python code. Within populations of fixed size N, each individual's genotype is solely identified

by the number of deleterious mutations it has, which is denoted by i, $i = 0, 1 \ldots, L$. L represents

the genome length, which is also the maximum number of deleterious mutations possible.

Populations are represented by vectors V of length $L + 1$, where each element V_i records the

number of individuals with i deleterious mutations so that $\sum_i V_i = N$. Individuals with i

mutations have fitness denoted by

$$W_i = e^{-s i^{1-\epsilon}} \tag{1}$$

with selection coefficient s and epistasis parameter ϵ. $\epsilon = 0$ in the absence of epistasis (**Fig. 1**), ϵ

< 0 with negative epistasis (**Fig. 2A & B**: ϵ = -0.25), and $\epsilon > 0$ with positive epistasis (**Fig. 2C &**

D: ϵ = 0.25). Fitness defined by **Eqn. 1** depends only on the number of deleterious mutations in

the genome and not their identity. We designate such fitness landscapes as isotropic.

Each generation of evolution consists of selection and mutation. The number of offspring with i

mutations is a random variable proportional to V_i, i.e., number of parents carrying i mutations,

times W_i, i.e., fitness of these parents. Then, mutations are imposed by sampling from Poisson

distribution with parameter U_{del} to generate number of deleterious mutations, and from Poisson

distribution with parameter U_{ben} to generate number of beneficial mutations. Finally, a new

vector V' is constructed, where V_i' records number of individuals with i mutations in the new

generation. Simulations are performed for a prespecified number of generations, usually 10,000,

to ensure the result is unaffected by transient effects. We hold population size constant in all our

simulations, i.e., we assume soft selection, and characterize populations in which all individuals

are carrying $i = L$ deleterious alleles as being at extinction.

Infinite genome

For the first part of the results, we hold constant the ratio of beneficial to deleterious mutation

rates ($U_{ben}/U_{del} = 0.01$). Conceptually this is equivalent to the well-known infinite sites model

(Kimura 1969).

Simulations in **Fig. 1** and **Fig. 2** are performed on simple fitness landscapes with $L = 300$, with

only one isotropic peak defined by **Eqn. 1**.

We characterize the critical U_{del}, i.e., the highest U_{del} under which populations can resist

Muller's ratchet with reference to the time variance in fitness. In **Fig. S2**, we present the variance

in fitness across time at equilibrium for the simulations conducted in **Fig. 1A**. Notably, the

critical U_{del}, i.e., the highest U_{del} under which populations can resist Muller's ratchet is seen to

overlap with the U_{del} under which time variance in fitness reaches maximum. This is reasonable,

because when U_{del} is well below or well above the critical U_{del}, populations are either at MSDE

or extinction and thus exhibit low time variance in fitness. In contrast, near the critical U_{del}, the

"tug-of-war" between the two possible states maximizes stochasticity.

We utilize the above observation to approximately locate the critical U_{del} in **Fig. 1B**, across

different population size N and selection coefficient s. For each N and s combination, we record

the time variance in fitness under different U_{del}, and report the U_{del} associated with highest time

variance as the critical U_{del}. There are three sources of uncertainty in this analysis. First, it is

impossible to conduct simulations under every possible U_{del} value. In practice, we sample U_{del}

at intervals of 0.025, meaning that the "true" critical U_{del} may fall in between two adjacent

sampled U_{del} values. Second, because of this intrinsic granularity, the detected maximum time

variance may not be the true maximum. We measure such uncertainty as the range of U_{del} values

that show time variance above half of the detected maximum. Third, there is inevitable variation

in recorded fitness variance across replicated simulations, due to the stochastic nature of the

simulation. However, we find that the second source of variation dominates the other two by at

least one order of magnitude (not shown). Therefore, we portray only the second type of

uncertainty as error bars in **Fig. 1B**.

For simulations in **Fig. 3**, as shown in **Fig. 3A**, we fuse two isotropic peaks together: on the

positive epistasis side of the intervening fitness valley, $\epsilon = 0.25$ and $L = 270$; on the negative

epistasis side, $\epsilon = -0.25$ and $L = 29$. (These values of L were chosen so that fitness values on both

sides of the floor of the valley would match.) Mutation on both sides of the valley were biased as

described above, i.e. $U_{ben}/U_{del} = 0.01$.

For simulations in **Fig. 4**, we define a hybrid fitness landscape consisting of two domains, one

exhibiting positive epistasis and the other exhibiting negative epistasis. As shown in **Fig. 4A**, a

fraction p of the first mutations put individuals in the positive epistasis domain, where $\epsilon = 0.25$

and $L = 300$, while the remaining fraction $1 - p$ of first mutations put individuals at the negative

epistasis domain where $\epsilon = -0.25$ and $L = 300$. If a lineage returns to the peak (by beneficial

mutations), the same p and $1 - p$ apply, i.e., lineages can "travel" between the two domains

through the peak.

Finite genome

For the second part of the results, we allow U_{ben}/U_{del} to evolve with the proportion of

deleterious mutation in the genome. Specifically, U_{ben} equals proportion of deleterious

mutations times the total mutation rate U, while U_{del} equals $U - U_{ben}$. Consequently, U_{ben} and

U_{del} are unique for individuals in different V_i, and we can only specify total mutation rate. In **Fig.**

5, simulations are conducted on an isotropic non-epistatic fitness landscape with $L = 10$ and $s =$

0.4.

Results

We use individual-based simulations of finite populations of asexual organisms to study the

ability of selection and evolving ratios of beneficial to deleterious mutation rates to protect

populations from deleterious mutations. We first assume a constant ratio of beneficial to

deleterious mutation rates and examine evolutionary behavior as a function of fitness landscape

topography. We then allow the ratio of beneficial to deleterious mutation rates itself to evolve

while holding other aspects of the fitness landscape constant.

Effect of fitness landscape topology

We simulate evolution on four kinds of fitness landscapes with increasing complexity, while

holding the ratio of beneficial to deleterious mutation (U_{ben}/U_{del}) equal to 0.01. We define a

fitness peak as isotropic if the fitness of any genotype only depends on its number of deleterious

mutations, but not the identity of these mutations. We start by examining evolutionary behavior

on fitness landscapes with a single peak where every deleterious mutation shares the same effect

regardless of the current genome (**Eqn. 1**: $\epsilon = 0$, **Fig. 1A** inset). This is, by definition, an

isotropic peak in the absence of epistasis. Next, we allow the effect of deleterious mutations to

increase (**Eqn. 1**: $\epsilon < 0$, **Fig. 2A**) or decrease (**Eqn. 1**: $\epsilon > 0$, **Fig. 2C**) with the number of

deleterious mutations in the current genome. In other words, we are adding epistasis to the

isotropic peak, distinguishing between negative (**Fig. 2A**) and positive (**Fig. 2C**) epistasis. We

then examine evolutionary behavior on fitness landscapes consisting of mutationally adjacent

isotropic peaks with negative and positive epistasis (**Fig. 3A**). Finally, we relax our assumption

of isotropic peaks, and model populations evolving on hybrid fitness peaks characterized by

domains of positive and negative epistasis (**Fig. 4A**).

In the absence of epistasis, populations residing on steeper fitness landscapes survive under

higher deleterious mutation rate

We first confirm the effect that stronger selection is better able to protect populations from

Muller's ratchet, or equivalently, that mutational robustness can enable the ratchet (Goyal et al.

2012). To do so, we construct two fitness landscapes, each composed of one isotropic peak but

with different steepness (selection coefficients s) in the absence of epistasis (**Fig. 1A** inset). We

simulate the evolution of finite asexual populations on such landscapes under different

deleterious mutation rates, and record their fitness at equilibrium (**Fig. 1A**). As mentioned above,

we implement soft selection, and we define extinction as occurring when all individuals in the

population attain the minimum possible fitness.

Under low mutation rates, selection is able to purify deleterious mutations, and drift (and thus,

the ratchet) is irrelevant on both flat and steep landscapes. This is because the stochastic loss of

fittest class is extremely rare: purifying selection is strong relative to mutation, and if lost, the

fittest class is quickly restored by beneficial mutations. Consequently, equilibrium fitness lies

close to the well-known infinite population expectation $e^{-U_{del}}$ (black dashed line in **Fig. 1A**,

Haldane 1937; Kimura and Maruyama 1966). However, after U_{del} exceeds some critical value

(at this population size, $U_{del} \approx 0.4$ for $s = 0.1$ and $U_{del} \approx 1.9$ for $s = 0.4$, see **Fig. 1A**), rapidly

accumulating deleterious mutations overwhelm selection, leading to extinction by Muller's

ratchet. We name the highest U_{del} under which populations are able to resist Muller's ratchet the

"critical U_{del}". Our simulation results for the critical U_{del} under different N and s are consistent

with analytical results based on **Eqn. 7** in Goyal et al. 2012 (see **Fig. 1B**). Given a fixed

population size (N) and constant U_{ben}/U_{del}, populations on steeper landscapes are able to

survive under higher U_{del}, i.e., they demonstrate a higher critical U_{del} (**Fig. 1A & B**, Goyal et al.

2012). In other words, in the long term, increased mutational robustness increases susceptibility

to Muller's ratchet.

Fitness landscapes with negative epistasis protect populations from Muller's ratchet while

ones with positive epistasis make populations more vulnerable to the ratchet

Epistasis means the dependence of mutational fitness effects on the genetic background. Without

epistasis (**Fig. 1**), there is no evolution of selection coefficients, since each mutation will always

have the same effect. Put differently, epistasis determines how selection changes as deleterious

mutations accumulate, shaping both local mutational and ratchet robustness experienced by an

evolving population. In order to understand the role of epistasis, we consider the two simplest

cases: one where selection strength monotonically increases with the number of deleterious

mutations, and the other where selection strength monotonically decreases with the number of

deleterious mutations. In other words, we consider a single isotropic peak with either only

negative epistasis (**Fig. 2A**) or only positive epistasis (**Fig. 2C**), respectively.

When epistasis is negative, as deleterious mutations accumulate, the local strength of purifying

selection increases, and consequently subsequent deleterious mutations become less likely to

accumulate. This hints at a negative feedback between the accumulation of deleterious mutations

and the tendency to accumulate more, which potentially could halt Muller's rachet (as previously

seen in Kondrashov 1994). On the other hand, when epistasis is positive, as deleterious

mutations accumulate, local selection weakens, and deleterious mutations can more easily

accumulate. This suggests a positive feedback between the aforementioned two forces, which

may render populations more vulnerable to the ratchet than those examined above. Importantly,

the critical U_{del} will thus vary across the fitness landscape in response to this variation in the

strength of selection

Our simulations are consistent with this intuition (**Fig. 2B, Fig. 2D**). In the presence of negative

epistasis, when U_{del} is lower than the critical U_{del} at the peak, it is lower than the critical U_{del}

anywhere on the landscape. Therefore, selection will drive the population back toward the peak

regardless of where on the landscape it is initialized (**Fig. 2B**). Even if U_{del} is higher than the

critical U_{del} at the peak, there exists a point on the landscape where selection exactly offsets such

U_{del}, because purifying selection increases monotonically from the peak (see also **S1 Text**). If a

population is initialized below this point, selection locally will be strong enough to push

populations upward until this point is reached. If instead, a population is initialized above this

point, mutation will be strong enough to push the population downward to this point, but no

further. Therefore, even under high U_{del}, landscapes with negative epistasis will halt Muller's

ratchet and establish stable MSDE.

Conversely, on the landscape with positive epistasis, when U_{del} is lower than the critical U_{del} at

the peak, populations initiated at the peak can maintain MSDE. However, positive epistasis

means that purifying selection decreases monotonically from the peak, and consequently there

exists a point on the landscape where selection exactly offsets such U_{del}. Populations initiated

above this point will experience selection stronger than required to offset U_{del} and evolve to the

peak, while populations initiated below this point suffer from selection weaker than needed to

offset U_{del}, and will succumb to Muller's ratchet (**Fig. 2D**). Therefore, we refer to this point as

"point of no return". Importantly, even if parameter values are such that populations can

equilibrate at the peak, stochastic fluctuations will eventually take them across this point of no

return, after which they will succumb to the ratchet. Moreover, as U_{del} increases, the selection

required to oppose mutation naturally increases as well. Correspondingly, as U_{del} increases, the

point of no return migrates towards the peak. This imposes a greater danger of succumbing to

Muller's ratchet for populations in the vicinity of the peak via stochastic fluctuations. Finally,

once U_{del} is greater than the critical U_{del} at the peak, the point of no return overlaps the peak,

and mutation overwhelms selection everywhere on the fitness landscape.

Populations converge to the peak with negative epistasis on multi-peak fitness landscape

We next demonstrate that populations can indeed evolve to occupy regions of the fitness

landscape with negative epistasis when challenged by deleterious mutations. To do so, we

construct a fitness landscape composed of two mutationally adjacent isotropic peaks featuring

opposite signs of epistasis (**Fig. 3A**). Populations finding themselves below the point of no return

on the positive epistasis side will experience selection weaker than needed to offset mutation and

their fitness will decline to the valley, similar to populations declining to the bottom of the

landscape in **Fig. 2D**. Moreover, populations above the point of no return will nevertheless

experience stochastic fluctuations and will be eventually carried over the point of no return, at

which point they will also evolve to the valley. However, upon arrival at the valley, strongly

beneficial mutations become available, drawing populations onto the negative epistasis side of

the valley, after which they quickly climb to MSDE (**Fig. 3B**). (Note that $U_{ben} = 0.01 U_{del}$ on

both sides of the valley, and consequently this behavior is driven entirely by differences in the

local strength of natural selection.) Since the two peaks share identical selection coefficients at

the peak, the positive epistasis side has uniformly higher mutational robustness but uniformly

lower ratchet robustness. This extends our understanding of the tension between mutational and

ratchet robustness and demonstrates that ratchet robustness, instead of mutational robustness, is

likely to evolve in response to high deleterious mutation rates.

Mutational robustness and ratchet robustness on hybrid peaks

Biologically realistic landscape peaks are unlikely to have only negative or positive epistasis. As

a first attempt to investigate the complexity introduced by heterogeneity in epistasis, we now

allow the sign of epistatic effects of mutations to be dependent on the first mutation away from

the peak. Specifically, a certain fraction (p) of the first mutations now place the evolving

population on a region of the landscape exhibiting positive epistasis, while the remaining fraction

$(1 - p)$ place the evolving population on a region of the landscape exhibiting negative epistasis

(**Fig. 4A**). Concretely, among all the mutational paths leaving the peak, a proportion p of them

show positive epistasis, while the remaining fraction $1 - p$ show negative epistasis. We assume

that all first mutations share identical fitness effects, so that there is no immediate fitness

advantage to lineages entering the region with negative or positive epistasis. However, for all

genotypes with greater than one deleterious mutation, fitness is necessarily higher in regions of

the landscape with positive epistasis than in regions with negative epistasis (**Eqn. 1**). In other

words, regions of the landscape with positive epistasis have both higher fitness and higher

mutational robustness (but lower ratchet robustness) compared with ones with negative epistasis.

Note that the only mutational path between regions is through the peak. We evolved populations

on such hybrid peak with $p = 0.5$ and report the proportions of populations residing exactly at

the peak, in the negative epistasis region, and in the positive epistasis region as a function of U_{del}

(**Fig. 4B**).

When U_{del} is so low that majority of the individuals carry zero or one mutation (here $U_{del} \approx 0.2$,

Fig. 4B, **Fig. S3A**), the proportion of the population on the negative epistasis region is very close

to that on the positive epistasis region. This merely reflects the fact that p equals 0.5, since the

fitness cost of the first mutation is the same. However, as U_{del} rises to moderate level (here ~0.2

$< U_{del} <$ ~1.1, **Fig. 4B, Fig. S3B**), the proportion of the population on the negative epistasis

region begins to drop below 0.5. This reflects the selective enrichment of the subpopulation

experiencing positive epistasis: individuals with two or more mutations on the positive epistasis

region are fitter than are those with the same number of mutations on the negative epistasis

region. Nevertheless, in this intermediate range of values of U_{del}, a subpopulation on the

negative epistasis region is still sustained despite lower fitness, thanks to net mutational inflow

from the subpopulation at the peak (**Table S1**). In essence, the subpopulation on the negative

epistasis region is at mutation-selection balance: constantly being purified by selection but being

regenerated by mutation from the peak.

However, after U_{del} increases enough that the peak can no longer be sustained (here, $U_{del} \approx 1.1$),

the proportion of the population at the peak become negligible (**Fig. 4B, Fig. S3C**). As a result,

the subpopulation residing on the negative epistasis region of the fitness landscape is

mutationally disconnected from the peak and is quickly wiped out by selection. The remaining

population now occupies only the positive epistasis region. However, since U_{del} has

overwhelmed selection at the peak, it necessarily does so also at every other point in the positive

epistasis region, and the population quickly succumbs to Muller's ratchet. This threshold

recapitulates results seen when the point of no return reaches the peak on isotropic fitness peaks

with positive epistasis region (**Fig. 2C & 2D**, although the numeric value of the critical mutation

rate differs here, reflecting its dependence on the fitness landscape).

More importantly, the observed pattern does not depend on the particular value $p = 0.5$: even

when there is only a very small fraction of paths leaving the peak with positive epistasis,

subpopulations on the positive epistasis region of the landscape will be favored due to their

short-term fitness advantage (**Fig. 4C**). Such an advantage is amplified by higher U_{del}, so long as

it remains less than the critical U_{del} at the peak, i.e., the U_{del} under which individuals at the peak

are lost to Muller's ratchet. Below this threshold, the two subpopulations accumulate more

mutations, and thus experience increased fitness differences. Eventually, once U_{del} exceeds the

critical U_{del} at the peak, the mutational connection between subpopulations on the fitness

landscape is again extinguished. At this point both subpopulations are doomed. The lower-fitness

subpopulation on the region of the landscape with negative epistasis will lose its mutational input

and go extinct. At the same time, the remaining higher-fitness subpopulation on the region with

positive epistasis will succumb to Muller's ratchet.

Effect of ratio of beneficial to deleterious mutation rates

In order to isolate the role that topographic features of the fitness landscape play in an evolving

population's response to deleterious mutations, we have thus far assumed that the ratio of

beneficial to deleterious mutation rates remains fixed (see **Methods**). However, as deleterious

mutations accumulate, this ratio can itself evolve, due to reversion mutations. Moreover,

compensatory mutations (mutations that are only beneficial in the presence of deleterious

mutations at other loci) amplify this effect. For example, one survey estimated that each

deleterious mutation gives rise to approximately 12 compensatory mutations (Poon and Chao

2005).

Increasing ratio of beneficial to deleterious mutation rates with decreasing fitness protects

populations from Muller's ratchet

We therefore modify our simulations to allow deleterious mutations to revert, thereby causing

U_{ben}/U_{del} to increase with the accumulation of deleterious mutations. Biologically, this

approach is conservative because it neglects compensatory mutations induced by deleterious

mutation (e.g., Poon and Chao 2005). Since increasing U_{ben}/U_{del} reduces the rate at which new

deleterious mutations occur, this suggests that another negative feedback may exist between the

accumulation of deleterious mutations and the tendency to accumulate more (Goyal et al. 2012).

This is reminiscent of the one due to negative epistasis seen above (**Fig. 2A**). To gain more

insight into this effect, we compare equilibrium fitness in populations evolved when U_{ben}/U_{del}

is allowed to change with that seen in populations evolved under fixed U_{ben}/U_{del} (**Fig. 5**). As U

increases and deleterious mutations accumulate, we see that populations with varying U_{ben}/U_{del}

enjoy the benefit of lower U_{del} and equilibrate at higher fitness than do populations with fixed

U_{ben}/U_{del}. Eventually, under sufficiently high U, populations with fixed U_{ben}/U_{del} succumb to

Muller's ratchet, as seen above. In contrast, populations with varying U_{ben}/U_{del} halts the ratchet,

as previously described (Goyal et al. 2012).

Discussion

Negative feedbacks halt Muller's ratchet

Our results highlight the importance of negative feedbacks in halting Muller's ratchet. The

mechanisms responsible for such negative feedback may be varied, realized via negative

epistasis (**Fig. 2 & 3**), or increasing U_{ben}/U_{del} (**Fig. 5**). We note that many surveys of biological

fitness landscapes find extensive evidence for negative epistasis (Bershtein et al. 2006; Costanzo

et al. 2010; Khan et al. 2011; Steinberg and Ostermeier 2016). Additionally, U_{ben}/U_{del} tends to

increase as deleterious mutations accumulate due to reversions and compensatory mutations.

More importantly, both negative epistasis and increasing U_{ben}/U_{del} can be seen as representative

cases of changing distribution of fitness effects (DFE) (Eyre-Walker and Keightley 2007) with

decreasing fitness. Negative epistasis influences the magnitude of mutational effects, while

increasing U_{ben}/U_{del} influences the ratio between beneficial and deleterious mutations. While

outside the scope of our simulations, we suggest that as fitness goes down, both aspects of the

DFE can simultaneously evolve, resulting in a wide range of potential DFE changes that confer

ratchet robustness.

In fact, even in the presence of positive epistasis, it's plausible that increases in U_{ben}/U_{del} could

still halt Muller's ratchet. Conversely, even if U_{ben}/U_{del} decreases as deleterious mutations

accumulate, sufficiently strong negative epistasis could in principle protect populations from

deleterious mutations. (Although as noted above, there is little reason to believe that U_{ben}/U_{del}

should decline with fitness.) At this moment, we remain ignorant about the general rules driving

changing DFE as a function of fitness. In the future, theoretical models that parameterize the

DFE could be built to quantify the relative importance of negative epistasis and increasing

U_{ben}/U_{del} in protecting evolving populations from deleterious mutation. Experimental studies

could also be conducted to compare DFE at different fitness levels for various biological systems.

Notably, a previous experimental study using bacteriophage has observed that as fitness declines,

the rate of beneficial mutations changes much more drastically than does the size of mutational

effects (Silander et al. 2007).

Widespread existence of mutational robustness may result from selection for

environmental robustness

Box 2: Reconciliation with the literature on the evolution of mutational robustness

On close inspection, Fig. 2 in Wilke et al. 2001 (reproduced here as **Fig. 6**) may suggest the

evolution of ratchet robustness rather than of mutational robustness**.** Specifically, the fitness

distribution of genotypes in the mutational neighborhood of the lineage adapted under high

mutation rate appears to contain more lethal mutations than do those mutationally adjacent to the

lineage adapted to low mutation rate. Furthermore, the lineage adapted under low mutation rate

seems more prone to mutation accumulation than does the lineage adapted under high mutation

rate. Both observations suggest that the lineage adapted under low mutation rate experiences

weaker purifying selection than does the one adapted under high mutation rate. Thus, we

interpret the data in Wilke et al. to suggest that the lineage adapted under high mutation rate

actually has lower mutational robustness, and consequently higher ratchet robustness, consistent

with our results.

Unfortunately, the two studies (Codoner et al. 2006; Sanjuan et al. 2007) attempting to

experimentally validate conclusions in Wilke et al. failed to fully characterize the different

lineages' fitness landscapes. Specifically, only non-lethal genotypes were considered, while

lethal mutations were ignored. Thus, we are unable to assess whether the lineage evolved under

high mutation rate has higher mutational robustness than the lineage evolved under low mutation

rate, since only a fraction of the DFE was measured.

Findings in van Nimwegen et al. (1999) are actually in agreement with our conclusions here, as

previously noted (Wylie and Shakhnovich 2011). Specifically, by construction Muller's ratchet

is impossible in van Nimwegen et al.'s model, since any non-neutral mutation is lethal and so is

instantaneously eliminated. Put another way, under this model evolving populations enjoy the

Our results suggest that mutational robustness cannot be selected for the long term, and we

address previous theoretical studies on the evolution of mutational robustness in **Box**

2. Nevertheless, mutational robustness is seen at many levels of biological systems (Wagner

2013). We note however that the existence of mutational robustness need not imply selection for

mutational robustness (Hermisson and Wagner 2004; Proulx and Phillips 2005; Siegal and Leu

2014). Following others (de Visser et al. 2003), we suggest instead that mutational robustness

may often evolve as a correlated consequence of selection for environmental robustness, i.e., an

organism's ability to sustain fitness against environmental perturbations. Theoretically and

empirically, environmental robustness has been shown to give rise to mutational robustness

(Wagner et al. 1997; Ancel and Fontana 2000; Meiklejohn and Hartl 2002; de Visser et al. 2003;

Szollosi and Derenyi 2007; Goldsmith and Tawfik 2009; Domingo-Calap et al. 2010).

Why can natural selection favor the evolution of environmental but not mutational robustness?

The key distinction is that mutational robustness requires tolerating heritable perturbations,

which inevitably alters the "starting point" of future generations. Such heritable decay is intrinsic

to Muller's ratchet. By contrast, selection for environmental robustness entails non-heritable

environmental perturbation. Consequently, the short-term advantage of environmental robustness

is not offset by any long-term cost, accounting for the absence of an "environmental ratchet". In

summary, while mutational robustness may be widespread in nature, we suggest one alternative

interpretation for its evolution: namely as a correlated consequence of selection for

environmental robustness (de Visser et al. 2003). We address another possibility next.

Long-term and short-term fate of mutational and ratchet robustness

Our findings also illustrate that mutational robustness can be selected for in the short term,

despite the danger it imposes in the long term (**Fig. 4**). This suggests another mechanism for the

existence of mutational robustness in natural populations. How populations overcome this

shortsightedness remains an open question. One possibility described in a previous study

(O'Fallon et al. 2007) is that subdivision can protect populations from myopic selection for

mutational robustness. Because selection is more effective at purging deleterious mutations in

demes dominated by ratchet robust individuals, in that study net dispersal rates were higher from

those demes, and the population in total was thus enriched for such individuals in spite of the

short-term disadvantage. We predict that any population structure capable of hindering rapid

fixation of mutational robustness will help natural selection favor ratchet robustness. However, a

detailed survey of possible mechanisms is outside the scope of this study.

Conclusion

It is of fundamental interest to understand how natural populations evolve to cope with the fact

that almost all mutations are deleterious. Intuitively, natural selection may favor mutational

robustness, as deleterious mutations of lesser effects seem to pose less danger. However, we

have extended work of others to demonstrate another peril of deleterious mutations: long term

extinction via Muller's ratchet. Indeed, we show that mutational robustness increases the risk of

Muller's ratchet. We then demonstrate that both negative epistasis and increasing U_{ben}/U_{del}

confer ratchet robustness and can protect populations from deleterious mutations in the long term.

Overall, our findings indicate that natural selection can favor ratchet robustness over mutational

robustness, and we define important areas of future theoretical and experimental work.

References

Ancel, L. W., and W. Fontana. 2000. Plasticity, Evolvability, and Modularity in RNA. J. Exp.
Zool. 288:242–283.
Bershtein, S., M. Segal, R. Bekerman, N. Tokuriki, and D. S. Tawfik. 2006. Robustness-epistasis
link shapes the fitness landscape of a randomly drifting protein. Nature 444:929–932.
Bull, J. J., L. A. Meyers, and M. Lachmann. 2005. Quasispecies made simple.
Codoner, F. M., J.-A. Daros, R. V. Sole, and S. F. Elena. 2006. The Fittest versus the Flattest□:
Experimental Confirmation of the Quasispecies Effect with Subviral Pathogens. Plos
Pathog. 2:e136.
Costanzo, M., A. Baryshnikova, J. Bellay, Y. Kim, E. D. Spear, C. S. Sevier, H. Ding, J. L. Y.
Koh, K. Toufighi, S. Mostafavi, J. Prinz, R. P. St Onge, B. VanderSluis, T. Makhnevych, F.
J. Vizeacoumar, S. Alizadeh, S. Bahr, R. L. Brost, Y. Chen, M. Cokol, R. Deshpande, Z. Li,
Z.-Y. Lin, W. Liang, M. Marback, J. Paw, B.-J. San Luis, E. Shuteriqi, A. H. Y. Tong, N.
van Dyk, I. M. Wallace, J. A. Whitney, M. T. Weirauch, G. Zhong, H. Zhu, W. A. Houry,
M. Brudno, S. Ragibizadeh, B. Papp, C. Pál, F. P. Roth, G. Giaever, C. Nislow, O. G.
Troyanskaya, H. Bussey, G. D. Bader, A.-C. Gingras, Q. D. Morris, P. M. Kim, C. A.
Kaiser, C. L. Myers, B. J. Andrews, C. Boone, S. J. Dixon, M. Costanzo, A. Baryshnikova,
B. Andrews, C. Boone, L. Hartwell, A. H. Tong, A. H. Tong, R. Mani, R. P. St. Onge, J. L.
Hartman, G. Giaever, F. P. Roth, R. P. S. Onge, D. Segrè, A. Deluna, G. M. Church, R.
Kishony, A. Huber, E. J. Chen, C. A. Kaiser, M. C. Jonikas, M. B. Metzger, S. Michaelis, S.
Leidel, P. B. Rahl, C. Z. Chen, R. N. Collins, A. Esberg, B. Huang, M. J. Johansson, A. S.
Byström, T. Naumanen, L. D. Johansen, E. T. Coffey, T. Kallunki, L. D. Johansen, S. F.
Levy, M. L. Siegal, A. Levchenko, P. M. Kim, L. J. Lu, Y. Xia, M. B. Gerstein, H. B.
Fraser, D. P. Wall, A. E. Hirsh, C. Pál, B. Papp, L. D. Hurst, P. M. Kim, A. Sboner, Y. Xia,
M. Gerstein, B. Lehner, C. Crombie, J. Tischler, A. Fortunato, A. G. Fraser, M. E.
Hillenmeyer, M. J. Dunham, A. C. Gavin, N. J. Krogan, H. Yu, K. Tarassov, J. A. de Visser,
C. D. Meiklejohn, D. L. Hartl, A. B. Parsons, and C. S. Sevier. 2010. The genetic landscape
of a cell. Science 327:425–31.
Domingo-Calap, P., M. Pereira-Gomez, and R. Sanjuan. 2010. Selection for thermostability can
lead to the emergence of mutational robustness in an RNA virus. J. Evol. Biol. 23:2453–
2460.

Eyre-Walker, a, and P. Keightley. 2007. The distribution of fitness effects of new mutations. Nat.
Rev. Genet. 8:610–8.
Goldsmith, M., and D. S. Tawfik. 2009. Potential role of phenotypic mutations in the evolution
of protein expression and stability. Proc. Natl. Acad. Sci. U. S. A. 106:6197–202.
Goyal, S., D. J. Balick, E. R. Jerison, R. A. Neher, B. I. Shraiman, and M. M. Desai. 2012.
Dynamic mutation-selection balance as an evolutionary attractor. Genetics 191:1309–1319.
Haigh, J. 1978. The accumulation of deleterious genes in a population-Muller's Ratchet. Theor.
Popul. Biol. 14:251–267.
Haldane, J. B. S. 1937. The effect of Variation on Fitness. Am. Nat. LXXI:337–349.
Hermisson, J., and P. Wagner. 2004. The Population Genetic Theory of Hidden Variation and
Genetic Robustness. Genetics 168:2271–2284.
Khan, A. I., D. M. Dinh, D. Schneider, R. E. Lenski, and T. F. Cooper. 2011. Negative epistasis
between beneficial mutations in an evolving bacterial population. Science (80-.).
332:1193–1196.
Kimura, M. 1969. The number of heterozygous nucleotide sites maintained in a finite population
due to steady flux of mutations. Genetics 61:893–903.
Kimura, M., and T. Maruyama. 1966. The mutational load with epistatic gene interactions in
fitness. Genetics 54:1337–1351.
Kondrashov, A. S. 1994. Muller's ratchet under epistatic selection. Genetics 136:1469–1473.
Krakauer, D. C., and J. B. Plotkin. 2001. Redundancy, antiredundancy, and the robustness of
genomes. Proc. Natl. Acad. Sci. U. S. A. 99:1405–1409.
Labar, T., and C. Adami. 2016. Evolution of Drift Robustness in Small Populations of Digital
Organisms. bioRxiv 1–17.
Lynch, M., D. Butcher, R. Bürger, and W. Gabriel. 1993. The mutational meltdown in asexual
populations. J. Hered. 84:339–344.
Meiklejohn, C. D., and D. L. Hartl. 2002. A single mode of canalization. Trends Ecol. Evol.
17:468–473.
Muller, H. J. 1964. The relation of recombination to mutational advance. Mutat. Res. Mol. Mech.
Mutagen. 1:2–9.
O'Fallon, B. D., F. R. Adler, and S. R. Proulx. 2007. Quasi-species evolution in subdivided
populations favours maximally deleterious mutations. Proc. Biol. Sci. 274:3159–3164.
Ofria, C., D. M. Bryson, and C. O. Wilke. 2009. Avida: A software platform for research in
computational evolutionary biology. Pp. 3–35 *in* Artificial Life Models in Software (Second
Edition).
Poon, A., and L. Chao. 2005. The rate of compensatory mutation in the DNA
bacteriophage ??X174. Genetics 170:989–999.
Poon, a, and S. P. Otto. 2000. Compensating for our load of mutations: freezing the meltdown of
small populations. Evolution (N. Y). 54:1467–1479.
Proulx, S. R., and P. C. Phillips. 2005. The Opportunity for Canalization and the Evolution of
Genetic Networks. Am. Nat. 165:147–162.
Sanjuan, R., J. M. Cuevas, V. Furio, E. C. Holmes, and A. Moya. 2007. Selection for Robustness
in Mutagenized RNA Viruses. Plos Genet. 3:e93.
Schultz, S. T., and M. Lynch. 1997. Mutation and Extinction : The Role of Variable Mutational
Effects , Synergistic Epistasis , Beneficial Mutations , and Degree of Outcrossing. Evolution
(N. Y). 51:1363–1371.
Siegal, M. L., and J.-Y. Leu. 2014. On the Nature and Evolutionary Impact of Phenotypic

Robustness Mechanisms. Annu Rev Ecol Evol Syst. 45:496–517.

Silander, O. K., O. Tenaillon, and L. Chao. 2007. Understanding the evolutionary fate of finite
populations: The dynamics of mutational effects. PLoS Biol. 5:922–931.

Steinberg, B., and M. Ostermeier. 2016. Shifting Fitness and Epistatic Landscapes Reflect Trade-
offs along an Evolutionary Pathway. J. Mol. Biol. 428:2730–2743. Elsevier Ltd.

Szollosi, G. J., and I. Derenyi. 2007. Congruent Evolution of Genetic and Environmental
Robustness in Micro-RNA. Mol. Biol. Evol. 26:867–874.

van Nimwegen, E., J. P. Crutchfield, and M. Huynen. 1999. Neutral evolution of mutational
robustness. Proc. Natl. Acad. Sci. U. S. A. 96:9716–9720.

Visser, J. A. G. M., J. Hermisson, G. P. Wagner, L. A. Meyers, H. Bagheri-Chaichian, J. L.
Blanchard, L. Chao, J. M. Cheverud, S. F. Elena, W. Fontana, G. Gibson, T. F. Hansen, D.
Krakauer, R. C. Lewontin, C. Ofria, S. H. Rice, G. von Dassow, A. Wagner, and M. C.
Whitlock. 2003. Perspective: Evolution and Detection of Genetic Robustness. Evolution (N.
Y). 57:1959–1972.

Wager, G. P., and W. Gabriel. 1990. Quantitative Variation in Finite Parthenogenetic
Populations: What Stops Muller's Ratchet. Evolution (N. Y). 44:715–731.

Wagner, A. 2013. Robustness and evolvability in living systems.

Wagner, G. P., G. Booth, H. Bagheri-chaichian, G. P. Wagner, G. Booth, and H. Bagheri-
chaichian. 1997. A Population Genetic Theory of Canalization. Evolution (N. Y). 51:329–
347.

Wilke, C. O. 2001. Adaptive Evolution on Neutral Networks. 715–730.

Wilke, C. O., and C. Adami. 2003. Evolution of mutational robustness.

Wilke, C. O., J. L. Wang, C. Ofria, R. E. Lenski, and C. Adami. 2001. Evolution of digital
organisms at high mutation rates leads to survival of the flattest. Nature 412:331–333.

Wylie, C. S., and E. I. Shakhnovich. 2011. A biophysical protein folding model accounts for
most mutational fitness effects in viruses. Proc. Natl. Acad. Sci. U. S. A. 108:9916–21.

<u>Figure 1: Under a fixed ratio of beneficial to deleterious mutation rates, populations residing on</u>

<u>steeper fitness landscapes survive under higher deleterious mutation rate</u>

A. Equilibrium fitness of finite populations under different deleterious mutation rates (U_{del}) on

fitness landscapes with two different selection coefficients (s). Dashed black line: theoretical

prediction for equilibrium fitness in infinite populations ($e^{-U_{del}}$). Populations on the steeper

fitness landscape (blue), i.e., the one with a larger s, maintain MSDE under higher U_{del} than do

populations on the shallower landscape (red). Beneficial mutation rate $U_{ben} = 0.01 U_{del}$.

Population size $N = 1000$. Average fitness is recorded after evolving for 10,000 generations

(see **Methods**). Error bars: standard deviation over 50 replicates. Inset: comparison of the two

fitness landscapes, each of which is composed of one isotropic peak without epistasis.

B. Critical U_{del} values from simulations (points) and analysis (lines, derived numerically from

Eqn. 7 in Goyal et al. 2012), for landscapes with different selection coefficients in the absence of

epistasis. Error bars: uncertainty associated with locating the critical U_{del} from simulation results

(see **Methods**). Regardless of population size, populations on steeper landscapes always survive

under higher U_{del}.

<u>Figure 2: Fitness landscapes with negative epistasis protect populations from Muller's ratchet</u>

<u>while ones with positive epistasis make populations vulnerable to the ratchet</u>

A. Cartoon of evolutionary dynamics of populations on fitness landscape composed of one

isotropic peak with negative epistasis. Curved dashed line: fitness landscape. Straight tangent

dashed line: fitness landscape without epistasis that shares the same selection coefficient at the

peak. Note that the selection coefficient (the slope of the fitness landscape) increases with the

number of deleterious mutations. Golden vertical line and horizontal dashed line: stable MSDE

at the peak $e^{-U_{del}}$ (see main text, also S1 Text).

B. Time course of average number of deleterious mutations in 35 populations initialized at

random points on isotropic fitness peak with negative epistasis during 10,000 generations

($N = 1000, s = 0.2, U_{del} = 1.0, U_{ben} = 0.01 U_{del}, \varepsilon = -0.25$). Golden vertical line: same as

in A. All populations converge to the MSDE regardless of where they are initiated (green traces).

C. Cartoon of evolutionary dynamics of populations on fitness landscape composed of one

isotropic peak with positive epistasis. Curved dashed line: fitness landscape. Straight tangent

dashed line: fitness landscape without epistasis that shares the same selection coefficient at the

peak. Note that selection coefficient (equal to the slope of the fitness landscape) decreases with

the number of deleterious mutations. Cyan vertical line and horizontal dashed line: unstable

MSDE at the peak $e^{-U_{del}}$ (see main text). Violet vertical line: predicted point of no return

(derived numerically from **Eqn.7** in Goyal et al. 2012).

D. Time course of average number of deleterious mutations in 100 populations initialized at

random points on an isotropic fitness peak with positive epistasis during 10,000 generations

($N = 1000, s = 0.5, U_{del} = 0.5, U_{ben} = 0.01 U_{del}, \varepsilon = 0.25$). Cyan and violet vertical lines:

same as in C. Populations initiated above the point of no return tend to evolve to the peak (blue

traces), whereas ones initialized below it tend to succumb to Muller's ratchet (pink traces).

Realizations that fluctuate across the point of no return are colored in brighter blue and pink.

655 <u>Figure 3: Populations converge to peak with negative epistasis on multi-peak fitness landscape</u>

656 A. Cartoon of evolutionary dynamics of populations on multi-peak fitness landscape. Both

657 fitness peaks are isotropic, one with only positive epistasis (left) and the other with only negative

658 epistasis (right). Curved dashed lines on both sides represent the fitness landscape. Straight

659 tangent dashed line on both sides again represent the slope (i.e. the selection coefficient at the

660 peaks); this quantity is equal at both peaks. Horizontal dashed line: $e^{-U_{del}}$ on both sides. Cyan

661 vertical line: $e^{-U_{del}}$ on the positive epistasis side. Violet vertical line: numerically derived point

662 of no return on the positive epistasis side (**Eqn.7** in Goyal et al. 2012). Black vertical line: valley

663 of the landscape. Golden vertical line: $e^{-U_{del}}$ on the negative epistasis side.

664 B. Time course of average number of deleterious mutations in 100 populations initiated at

665 random, uniformly distributed points on multi-peak landscape over the course of 10,000

666 generations ($N = 10,000, s = 0.3, U_{del} = 0.5, U_{ben} = 0.01U_{del}$, positive epistasis side

667 $\varepsilon = 0.25$, negative epistasis side $\varepsilon = -0.25$). Cyan, violet, black and golden vertical lines:

668 same as in B. Populations are color coded based on their starting point (blue traces: above the

669 point of no return on the positive epistasis side, pink traces: below the point of no return on the

670 positive epistasis side, bright pink and blue traces: realizations that fluctuate across the point of

671 no return on the positive epistasis side, green traces: populations initialized on the negative

672 epistasis side). The vast majority of populations below the point of no return rapidly cross the

673 valley and converge to MSDE on the negative epistasis side. Due to uniformly stronger selection

674 on the negative epistasis side of the valley, populations there exhibit less stochasticity in their

675 trajectories. Simulations in smaller population size ($N = 1000$) show qualitatively the same

676 results, although it takes much longer to converge to the negative epistasis side (data not shown).

<u>Figure 4: Mutational robustness and ratchet robustness on hybrid peaks</u>

A. Cross-section through a hybrid fitness peak. While all first mutations in the highest fitness

genotype share the same fitness effect, a fraction p of them cause subsequent mutations to

exhibit positive pairwise epistasis, and the remaining fraction $1 - p$ cause subsequent mutations

to exhibit negative epistasis. Populations are always initiated at the peak. Here $p = 0.5$.

B. Equilibrium proportions of individuals at the peak (black), in the negative epistasis region

(blue), and in the positive epistasis region (pink), as a function of U_{del} ($U_{ben} = 0.01U_{del}$,

$p = 0.5, N = 1000, s = 0.35$, negative epistasis region: $\varepsilon = -0.25$, positive epistasis

region: $\varepsilon = 0.25$), evolved for 10,000 generations, error bars: standard deviation across 50

replicates. Under U_{del} less than the critical U_{del} at the peak (here ~1.1; see main text), the

subpopulation on the negative epistasis region relies on continual mutational input from

subpopulation on the peak. After U_{del} exceeds the critical U_{del} at the peak, the subpopulation on

the peak goes extinct, and with it, the subpopulation on the negative epistasis region. At this

point, the remaining population finds itself beyond the point of no return on the positive epistasis

region (see main text), and it succumbs to Muller's ratchet.

C. Equilibrium proportions of individuals at the peak (black), the negative epistasis region (blue),

and the positive epistasis region (pink), as a function of U_{del} (parameters are exactly the same as

panel B except $p = 0.01$), evolved for 10,000 generations, error bars: standard deviation across

50 replicates. Population dynamics resemble $p = 0.5$ (see main text).

<u>Figure 5: Increasing U_{ben}/U_{del} with decreasing fitness protects populations from Muller's</u>

<u>ratchet</u>

Equilibrium fitness as a function of total mutation rates (U) in finite asexual populations with

evolving and fixed U_{ben}/U_{del}. Dashed black line: theoretical prediction for equilibrium fitness in

infinite populations ($e^{-U_{del}}$) with fixed U_{ben}/U_{del} ($U_{ben} = 0.01U_{del}$). Population size $N =$

1000. Selection coefficient $s = 0.4$. Average fitness is recorded after evolving for 10,000

generations (see **Methods**). Populations in which U_{ben}/U_{del} evolve by virtue of reversion

mutations (blue) survive under high U, while ones with fixed U_{ben}/U_{del} (red) succumb to

Muller's ratchet. (These latter results are identical to those shown for $s = 0.4$ in **Fig. 1**.) Error

bars: standard deviation over 50 replicates.

<u>Figure 6: Annotated Fig. 2 from Wilke et al. (2001) suggests the evolution of ratchet robustness</u>

<u>in populations of digital organisms.</u> Populations in Avida (Ofria et al. 2009) were evolved for

1,000 generations at low and high mutation rates (μ = 0.5 and 2.0, respectively). In this figure,

the fitness landscapes in the mutational neighborhoods of these two populations were examined

by propagating new populations seeded with one representative individual from each for 15

generations. Top panels: "A" designates new populations seeded with a representative individual

adapted under low mutation rate. Bottom panels: "B" designates new populations seeded with a

representative individual adapted under high mutation rate. μ denotes the mutation rate during

the subsequent 15 generation experiment. Each bar indicates the frequency of genotypes with the

corresponding number of mutations at the end of these 15 generations; colors represent the

fitness of these genotypes. The left two panels illustrate the fitness landscape in the nearby

mutational neighborhood of A or B (small μ = 0.5). Importantly, the descendants of the

individual from the B population include many more lethal mutants (black) than do those from

the individual from the A population (red arrows). The right two panels depict the fitness

landscape across a broader mutational neighborhood of A or B (large μ = 2.0). Notably, the

viable descendants of the individual from the B population carry fewer deleterious mutations

than do those from the A population (red dashed lines represent the approximate mutational limit

among viable offspring: upper panel skews towards more mutations than does the lower panel).

Both observations suggest that selection against deleterious mutation is stronger in the

mutational neighborhood occupied by the B population than it is in the neighborhood of the A

population, consistent with the evolution of ratchet robustness rather than mutational robustness

in response to elevated mutation rate.

A, μ = 0.5   A, μ = 2.0

B, μ = 0.5   B, μ = 2.0

Frequency

Number of deleterious mutations

Fitness Heatmap

1.0

0.0