*Biological sciences-biochemistry*

# A shortened version of SecA (SecA^N) functions as the protein-conducting channel for nascent β-barrel outer membrane proteins

**Feng Jin[1,2], Zengyi Chang[1*]**

[1]State key Laboratory of Protein and Plant Gene Research, School of Life Sciences, Center for Protein Science, Peking University, Beijing, China
[2]Peking-Tsinghua Center for Life Sciences, Peking University, Beijing, China
[*]Correspondence: changzy@pku.edu.cn (Z.C.)

**Key words: β-barrel outer membrane protein (OMP) biogenesis, translocation, SecYEG translocon, SecA, SecA^N translocon**

During their biogenesis in cells, many proteins have to be translocated across biological membranes by protein-conducting channels. In Gram-negative bacteria, proteins destined to inner membrane, periplasm or outer membrane are synthesized in the cytoplasm and most of them are believed to be translocated across the inner membrane by the SecYEG translocon. Nevertheless, this perception, largely formed on the basis of genetic and *in vitro* studies, has hardly been proved by *in vivo* studies. Here, our *in vivo* photocrosslinking analysis mediated by genetically incorporated unnatural amino acid pBpA proved a direct interaction between the nascent periplasmic proteins and SecY, but strikingly, not between the nascent β-barrel outer membrane proteins (OMPs) and SecY. We then demonstrated that precursors of OMPs but not of periplasmic proteins are effectively processed in a SecY-defective strain. Subsequently, we discovered that a shortened form of SecA that lacks its C-terminal region, designated as SecA$^N$, apparently functions as the translocon for nascent OMPs. In support of this, we discovered that SecA$^N$ directly interacts with BamA protein of the β-barrel assembly machine, and that SecA$^N$ is able to form a ternary complex with BamA and nascent OmpF in living cells. These observations strongly implicate that, in contrast to the common perception, the SecYEG only functions as the translocon for the nascent periplasmic proteins, not for the OMPs.

## Introduction

Many proteins have to be translocated across biological membranes during their biogenesis in living cells (*1-6*). Such translocation processes are carried out by membrane-integrated protein-conducting channels (*7-11*). In Gram-negative bacteria, both outer membrane proteins and periplasmic proteins are synthesized in the cytoplasm and translocated across the inner membrane through the co- or post-translational pathway (*10, 12*). The SurA protein has been identified as the primary chaperone for escorting nascent β-barrel outer membrane proteins (henceforth abbreviated as OMPs) across the periplasm (*13-16*). The BamA protein, as the major subunit of the essential β-barrel assembly machine (Bam) complex, is integrated into the outer membrane via its C-terminal β-barrel structure and extends into the periplasm via its N-terminal POTRA domains (*17-19*).

According to the currently prevailing perception, protein translocation across the inner membrane in Gram-negative bacteria is accomplished by the *sec* transcolon, whose membrane integrated SecYEG complex acts as a protein-conducting channel while the peripheral SecA ATPase functions as the molecular motor (*7, 8, 11, 20*). The homologue of SecYEG in eukaryotic cells is designated as Sec61 and is integrated into the membrane of the endoplasmic reticulum (*4, 7, 21-23*) . Initially, the *secY* gene was identified as one whose suppressor mutations restored the export of OMPs or periplasmic proteins that have a defective signal peptide (*24, 25*), as well as one whose defect resulted in a slow processing for precursors of both periplasmic proteins and OMPs (*26, 27*). The SecY and SecE proteins were found to act together with the SecA protein in supporting translocation of the precursor form of an outer membrane protein (*28*). The protein-conducting channel is believed to be largely formed by the SecY protein in light of its crystal structure (*29, 30*) as well as *in vitro* chemical crosslinking analysis (*31, 32*).

The *secA* gene was initially identified as one whose defect resulted in accumulation of precursors of both OMPs and periplasmic proteins (*33, 34*). The SecA protein was later found to be essential for translocation of both OMPs and periplasmic proteins (*35, 36*). SecA was subsequently characterized as a cytoplasmic as well as a peripheral and integrated membrane protein (*37-40*), and was demonstrated to be exposed to the periplasm (*41-43*). Nevertheless, the subcellular localization of SecA remains an issue of debate. Although the SecA protein was proposed to undergo ATP driven cycles of membrane insertion and deinsertion (*44, 45*), it was reported that a significant fraction of the SecA protein is permanently embedded in the inner membrane (*39*). In contrast to the conventional view, it has also been reported that SecA alone, without SecYEG, is able to promote translocation of OMPs across inner membrane vesicles or artificial lipid bilayers under *in vitro* conditions (*46-51*).

In regards to the biogenesis of OMPs and periplasmic proteins, as well as the function of SecYEG and SecA in living cells, many issues remain unresolved. For example, if the prevailing perception on SecYEG is correct, then how can a single SecYEG channel translocate the nascent polypeptides of such dramatically different clients as OMPs, periplasmic proteins and even inner membrane proteins? How can SecA exist in three different subcellular localizations and what does each form exactly do in living cells? In this study, mainly via unnatural amino acid mediated *in vivo* photocrosslinking (*52*), we first unexpectedly observed that although nascent polypeptides of periplasmic proteins directly interacted with SecY, those of OMPs did not. Further, we observed

that precursors of OMPs, but not of periplasmic proteins, could be effectively processed when SecY becomes defective. Subsequently, we demonstrated that a shortened version of SecA, designated as SecA$^N$, directly interacts with the periplasmic part of BamA. These observations strongly implicate that SecA$^N$ functions as the translocon for translocating the nascent OMPs, while SecYEG functions as the translocon for nascent periplasmic proteins in living bacterial cells.

## Results

**Nascent polypeptides of periplasmic proteins but not of OMPs directly interact with SecY**

To clarify whether nascent polypeptides of both periplasmic proteins and OMPs are translocated across the inner membrane through the SecYEG translocon as commonly percepted, we first performed *in vivo* photocrosslinking analysis by individually introducing the unnatural amino acid pBpA (*p*-benzoyl-L-phenylalanine) (*52*) at 21 randomly selected residue positions across the polypeptide of OmpF (an OMP) or 10 positions of SurA (a periplasmic protein). The pBpA variants of OmpF or SurA were then each expressed in the *ompF*-deleted or *surA*-deleted LY928 bacterial strain, whose genome was modified to encode the amino-acyl tRNA synthetase and the tRNA needed for pBpA incorporation as we reported before (*16*). Strikingly, our blotting analysis against SecY failed to detect any photocrosslinked product between OmpF and SecY (**Fig. 1A**; displayed are results of 12 pBpA variants of OmpF), though successfully detected the photocrosslinked product between SurA and SecY when pBpA was introduced at signal peptide residue positions 8 and 12 in SurA (lanes 2 and 4, **Fig. 1B**). It should be pointed out that we did not detect any photocrosslinked SurA-SecY product when pBpA was introduced at the mature part of SurA protein (lanes 8-11, **Fig. 1B**). Collectively, these observations indicate that the nascent OMPs, unlike the periplasmic proteins, apparently do not directly interact with the SecYEG translocon in living cells.

**Figure 1. Nascent polypeptides of periplasmic proteins but not of OMPs directly interact with SecY.**

(**A, B**) Blotting results for detecting the *in vivo* photocrosslinked products of the indicated pBpA variants of OmpF (**A**) or of SurA (**B**) expressed in the LY928-Δ*ompF* or LY928-Δ*surA* cells, probed with the Streptavidin-alkaline phosphatase (AP) conjugate against the AVI tag added on the SecY protein.

Residue positions are numbered by including the signal peptides. Protein samples were resolved by SDS-PAGE before subjecting to the blotting. Samples of cells expressing wild type OmpF or SurA (no pBpA incorporation) were analyzed as negative controls (lanes 1, **A** and **B**). Indicated on the right of each gel are the position of SecY, SecA, photocrosslinked SurA-SecY or OmpF-SecY (The SecY protein in lanes 2-5 in panel **B** mobilized at the bottom edge of the gel, thus hardly visible), and on the left are positions of molecular weight markers.

**Precursors of OMPs but not of periplasmic proteins can be effectively processed when SecY is defective.**

To further clarify the role of the SecYEG translocon in translocating OMPs and periplasmic proteins, we employed the cold sensitive mutant strain, SecY39, in which protein export was found to be partially retarded at the permissive temperature of 37°C and dramatically retarded at the non-permissive temperatures of 20°C (*53*) . We observed a far less defect in the processing of precursor of OMPs in comparing with that of periplasmic proteins when this strain was cultured at the non-permissive temperature (**Fig. 2A**). This is reflected by a much lower level of accumulation of precursors for OMPs (OmpA and OmpF) than for periplasmic proteins (SurA and MBP), as shown in lane 2 of **Fig 2A**, as well as by a significantly more efficient processing of the accumulated precursors of OMPs than of periplasmic proteins in the 1 hour chasing period after adding chloramphenicol (that inhibits protein synthesis in cells (*54*)), as shown in lanes 2-4 of **Fig 2A**. In comparison, when the SecY39 cell was cultured at the permissive temperature of 37°C, the differences in both precursor accumulation and processing are far less significant than those observed at the non-permissive temperature of 20°C (**Fig. 2B**). Collectively, these observations indicate that the processing of the precursors for OMPs relies on the SecYEG translocon in a degree far less than that for periplasmic proteins. Our observation here is somehow consistent with what was reported before that the *secY39* mutation affected MBP export more strongly than it affected OmpA export (*53*).

**Figure 2. Precursors of OMPs but not of periplasmic proteins are effectively processed when SecY is defective.**
Immunoblotting results for detecting the precursor, mature and folded forms of the indicated OMPs (OmpA and OmpF) and periplasmic proteins (SurA and MBP) in the SecY39 cells cultured either at the non-permissive temperature (**A**), or permissive temperature (**B**); cells were treated with chloramphenicol (Chl) for the indicated length of time. Protein samples of SecY39 cells were not boiled for the semi-native SDS-PAGE analysis (lanes 2-4 in **A** and **B**). Protein samples from wild type cells (WT) were boiled for indicating positions of mature forms (that

mobilizes with a slightly higher rate than the precursors) of OMPs and periplasmic proteins (lanes 1 in **A** and **B**), or not boiled for indicating positions of folded forms of OMPs (lane 5 in **A**). The EF-Tu protein was analyzed here to indicate an equal loading of samples in lanes 2-5. Indicated on the right of each gel are positions of different forms of OMPs (precursor, mature and folded) and periplasmic proteins (precursor and mature), and on left are positions of the molecular weight markers.

**The POTRA 2 domain of BamA directly interacts with a form of SecA that lacks its C-terminal region**

Our observations described above suggest that the SecYEG translocon does not directly translocate nascent OMPs across the inner membrane. We then addressed whether SecA functions to translocate nascent OMPs across the inner membrane. We initially failed to detect any direct interaction either between OmpF and SecA (one representative data is shown in lane 3, **Fig. 4A**) or between SurA and SecA when the pBpA variants were expressed in the LY928 strain, likely due to the transient nature of these interactions in living cells. We then observed such interactions when the in vivo photocrosslinking was performed under a SecA defective condition with the LY928-SecA81 strain that we particularly constructed for this study (*55*) (**Fig S1**) and meanwhile observed that precursors of neither OMPs nor periplasmic proteins were effectively processed within this cell (**Fig. S2**).

We next examined whether SecA interacts with protein factors that are known to participate in OMP biogenesis but localized outside the inner membrane, such as SurA and BamA (*13-19*). For this purpose, we first tried to find out whether pBpA individually introduced at six residue positions of SecA (at 47, 300, 530, 863, 868 or 896) that were reported to be exposed to the periplasm (*43*) was able to mediate photocrosslinking with SurA or BamA. Our immunoblotting analysis revealed no photocrosslinked SecA-SurA product for all these six variants, but a putative SecA-BamA product band of about 145 kD, being significantly lower than the expected combined molecular mass of BamA (95 kD) and SecA (102 kD) for one pBpA variant of SecA (lane 1, **Fig. 3A**).

To verify this putative interaction between SecA and BamA, we then performed a reciprocal photocrosslinking by introducing pBpA in the periplasm-located part of BamA. Remarkably, our immunoblotting analysis revealed the same BamA-SecA product, also of about 145 kD, when pBpA was introduced at the POTRA 2 domain of BamA (lane 1, **Fig. 3B**). It is noteworthy that this putative BamA-SecA band could also be clearly detected when probed with the streptavidin-AP conjugate against the AVI tag of BamA (lane 3, **Fig. 3B**).

We next jsolated this putative photocrosslinked BamA-SecA product and subjected it to mass spectrometry analysis. Although the results revealed presence of both BamA and SecA, interestingly, the score for SecA is rather low and that the two identified peptide fragments of SecA were both derived from the N-terminal region of SecA (**Fig. 3C**), indicating that SecA in the photocrosslinked SecA-BamA product is possibly a shortened form. We then tried to clarify this by performing immunoblotting analysis with antibodies against either the N-terminal region (fragment 1-209) or the C-terminal region (fragment 665-820) of SecA. Results presented in **Fig. 3D** demonstrate that this putative SecA-BamA product could be detected only by antibodies against the N-terminal region (lanes 1 and 2 vs lanes 3 and 4) of SecA. In light of these observations, it is conceivable that the SecA in the photocrosslinked SecA-BamA product is

composed of its N-terminal region of about 50 kD. A careful look of our immunoblotting results probed with anti-SecA antibodies (lanes 1 and 2 in **Fig. 3B**; **Fig. 3D**) identified a dense but apparently non-specific band at the 50 kD position (given its far smaller size than the SecA protein).

In an effort to unveil this shortened version of SecA, we then resolved the protein samples using a separating gel that resolves smaller proteins with a higher resolution (*56*). With this modification, our immunoblotting analysis then did reveal this shortened version of SecA that could only be detected by antibodies against the N-terminal region (lane 1 vs lane 3, **Fig 3E**). However, the level of this form of SecA is apparently much lower than that of full length SecA (lane 1, **Fig 3E**). This is somehow consistent with previous observations that only about one sixth of the total amount of SecA is integrated into the inner membrane (*37, 39*). In retrospect, the reason for this shortened SecA form to be ignored before is most likely due to the co-existence of the nearby non-specific protein bands (as shown in **Fig. 3B** and **3D**). Collectively, these observations strongly indicate that a shortened form of SecA interacts with BamA in living cells. We henceforth designate this form of SecA as SecA[N].

**Figure 3. The POTRA 2 domain of BamA directly interacts with a shortened form of SecA.**

(**A**) Immunoblotting results of *in vivo* photocrosslinked products of the indicated pBpA variants of SecA expressed in LY928 cells, probed with antibodies against SurA (top) or BamA (bottom). (**B**) Blotting results of *in vivo* photocrosslinked product of BamA-V121pBpA expressed in LY928 cells, using antibodies against SecA (lanes 1 and 2) or streptavidin-AP conjugate against the AVI tag of BamA (lanes 3 and 4). (**C**) Results of mass spectrometry analysis of the putative photocrosslinked BamA-SecA product generated from BamA-V121pBpA. Shown are the protein scores for BamA and SecA, as well as the two peptide sequences identified for SecA. (**D**) Immunoblotting results of the *in vivo* photocrosslinked product of BamA-V121pBpA using antibodies against the N- (lanes 1 and 2) or C-terminal (lanes 3 and 4) region of SecA. (**E**) Immunoblotting results for detecting the shortened version of SecA after resolving on a high resolution gel, also using antibodies against the N- (lane 1) or C-terminal (lane 3) region of SecA. Protein samples were resolved by SDS-PAGE before subjecting to immunoblotting. The polymerized separating gel in panel **A**, **B**, **C** and **D** is 5%, while in panel **E** is 10%. Indicated on the right of each gel are positions of the indicated forms of proteins and on the left are Positions of molecular weight markers. Cells expressing wild type SecA (no pBpA introduced) were analyzed as negative controls (lane 7, panel **A**).

**A BamA-OmpF-SecA$^N$ ternary complex is formed in living cells**

In light of our aforementioned observations that nascent OmpF interacts with SecA (**Fig. S1**) and BamA interacts with SecA (**Fig. 3**), we then addressed whether a ternary complex could be formed among BamA, nascent OmpF and SecA$^N$. For this purpose, we simultaneously introduced pBpA into both BamA (at residue position of V121) and OmpF (at residue position of G18). We then performed this dual *in vivo* photocrosslinking within the LY928 cells co-expressing BamA-V121pBpA and OmpF-G18pBpA. Blotting analysis using antibodies against the N-terminal region of SecA or streptavidin-AP conjugate against the AVI tag of BamA, remarkably, indeed revealed the presence of a photocrosslinked ternary complex of BamA-OmpF-SecA$^N$ and a binary complex of BamA-SecA$^N$ (lanes 1 and 7, **Fig. 4**). Consistent with results presented in lane 3, **Fig. 3D**, such a ternary BamA-OmpF-SecA$^N$ could not be detected by antibodies against the C-terminal region of SecA (lane 5, **Fig. 4**). Collectively, these data strongly suggest that the SecA$^N$ protein, by interacting with the BamA protein, translocates nascent OMPs across the inner membrane.

**Figure 4. A BamA-OmpF-SecA$^N$ ternary complex is formed in living cells.**

Blotting results of the photocrosslinked products formed in the LY928 cells expressing the indicated pBpA variants of BamA and OmpF, using antibodies against the N-terminal (lanes 1-4) or C-terminal region of SecA (lanes 5 and 6), or the streptavidin-AP conjugate against the AVI tag of BamA (lanes 7 and 8).

Protein samples were resolved by SDS-PAGE before subjecting to the blotting. Indicated on the right of the gel are positions of BamA, SecA, as well as the photocrosslinked BamA-OmpF-SecA$^N$ and BamA-SecA$^N$ complexes, on the left are positions of the molecular weight markers.

**Discussion**

This study was performed in an initial attempt to clarify whether both OMPs and periplasmic proteins are translocated across the inner membrane through the SecYEG translocon in living cells, as commonly percepted (*7, 8, 26*). Our *in vivo* photocrosslinking analysis mediated by genetically incorporated unnatural amino acid, though proved a direct interaction between nascent periplasmic proteins and SecY, strikingly, did not reveal any direct interaction between nascent OMPs and SecY (**Fig. 1**). We then demonstrated that precursors of OMPs but not of

periplasmic proteins could be effectively processed in a SecY defective strain (**Fig. 2A**). Afterwards, we revealed that a shortened form of SecA that lacks its C-terminal region, SecA[N], most likely functions as the translocon for nascent OMPs in living cells, as supported by the following observations. First, SecA[N] directly interacts with BamA in living cells (**Fig. 3A**). Second, reciprocally, BamA directly interacts with SecA[N] (**Fig. 3B**). Third, a BamA-OmpF-SecA[N] ternary complex was detected (**Fig. 4**). Fourth, processing of precursors of OMPs relies largely on ATP but hardly on the transmembrane proton gradient, while that of periplasmic proteins relies on both as energy sources (**Fig. S3**).

Our observations described here, in combination with revelations reported before (*13-19, 57*), strongly implicate that SecA[N] is responsible for translocating nascent OMPs across the inner membrane. Here, as illustrated in **Fig. 6**, we emphasize the following points. First, as the translocon of OMPs, SecA[N] (whose oligomeric status remains to be resolved) is integrated into the inner membrane by interacting with a SecYEG protomer, which in turn interacts with a second SecYEG protomer that likely functions as the translocon for nascent periplasmic proteins (*58, 59*). Second, the full length form of SecA in the cytoplasm delivers the signal peptide-containing nascent OMPs and nascent periplasmic proteins to SecA[N] and SecYEG traslocons, respectively. Third, a supercomplex containing SecA, SecA[N], SecYEG, SurA, PpiD (*16, 60*) and BamA, spanning the cytoplasm, the inner membrane, the periplasm and the outer membrane functions for the biogenesis of OMPs in living cells.

**Figure 6. A model illustrating the translocation of nascent OMPs, through SecA[N], and nascent periplasmic proteins, through SecYEG, across the inner membrane in Gram-negative bacteria.**

We searched the literatures in an attempt to seek for earlier hints on the presence of SecA[N] in bacterial cells. Interestingly, we did notice the presence of a SecA form with a size very close to SecA[N] that could either be immunoprecipitated by antibodies against the N-terminal region of

SecA (as shown in lanes 7 and 8, Fig. 2, in (*61*)) or be detected in the inner membrane fraction of *E. coli* cells by antibodies against the full length SecA (Fig. 4 in (*43*)). Moreover, it was also reported that the N-terminal fragment of about 240 residues of the SecA protein from either *E coli* (*37*) or *B. Subtilis* (*62*) is integrated into the membrane. Many issues remain to be resolved for SecA$^N$. For example, how SecA$^N$ forms the protein conducting channel in the inner membrane? How SecA$^N$ is generated from the full length SecA? How SecA$^N$ interacts and works with full length SecA in the cytoplasm?

In retrospect, the perception that SecYEG is directly responsible for translocating both OMPs and periplasmic proteins has been mainly derived from genetic and *in vitro* reconstitution studies (*26, 28*). Our study reported here represents an attempt to clarify this perception by performing experiments in living cells. As observed before, our data also indicate that SecYEG does play an important role for translocating nascent OMPs across the inner membrane in living cells (**Fig. 2**). This can be explained as follows. First, SecYEG is responsible for translocating periplasm-located quality control factors for OMP biogenesis such as SurA (*13-16*). Second, SecYEG most likely anchors SecA$^N$ in the inner membrane. This is indicated by such reported observations as that an overproduction of SecYEG resulted in an increased level of membrane-integrated SecA (*42*), and that the additional of SecYEG significantly increased the efficiency of SecA in functioning as protein-conducting channels in liposomes (*50*). Unresolved issues regarding the relationship between SecYEG and SecA$^N$ include the following. Is SecA$^N$ indeed anchored in the inner membrane by SecYEG as we presumed in our model? Do SecA$^N$ and SecYEG have any functional coordination in translocating OMPs and periplasmic proteins? How does the cytoplasmic SecA effectively partition nascent OMPs and nascent periplasmic proteins to SecA$^N$ and SecYEG, respectively? The molecular mechanism for SecA$^N$ to function as a protein-conducting channel, as well as the structural and functional interaction between SecA$^N$ and SecYEG undoubtedly merit further investigations.

## Methods

### Bacteria strains and plasmid constructions

All bacteria strains and plasmids used in this research are respectively listed in **Tables S1** and **S2**. The pYLC-OmpF, pYLC-BamA, pYLC-SurA or pYLC-SecA plasmid was constructed by isolating the encoding gene (including its promoter) via PCR, using *E. coli* genomic DNA as template before the DNA fragment was inserted into the pYLC vector through restriction enzyme free cloning (*63*). The pYLC is a low copy plasmid that we derived from the pDOC plasmid (*16*). The site-specific mutagenesis for introducing the TAG amber codon into the specific gene in the plasmids to generate pBpA variant proteins was performed using the phusion site-directed mutagenesis kit (New England Biolabs, Massachusetts, USA).

### pBpA mediated *in vivo* photo-crosslinking

The pYLC-OmpF or pYLC-SurA plasmids carrying the TAG mutations were respectively expressed

in LA928-Δ*ompF* and LA928-Δ*surA* cells for *in vivo* photo-crosslinking analysis. The LA928 strain was generated by us for pBpA incorporation purposes as we described before (*16*). The cells were cultured at 37°C in LB medium containing pBpA (200 μm), grown to the mid-log phase, before irradiated with UV (365 nm) for 10 min in a Hoefer UVC-500 crosslinker. The UV irradiated cells were harvested by centrifugation, re-suspended in 100 μL SDS-PAGE loading buffer before boiled for 5 min. Protein samples were separated by SDS-PAGE before subjected to blotting analyses.

**Purification and mass spectrometry analysis of *in vivo* photocrosslinked product**

Photo-crosslinked product of BamA-V121pBpA was purified by affinity chromatography using streptavidin resin. The eluted sample was resolved by SDS-PAGE before the gel was subjected to either blotting analysis or Coomassie Blue staining. The protein band at around 145 kD was then excised from the gel and applied for Liquid chromatography-tandem mass spectrometry (LC-MS/MS) identification as we described before (*16*).

**Chloramphenicol treatment of the secY39 and secA81 cells**

A single colony of either the SecY39 or SecA81 mutant strain was inoculated in 3 mL of LB medium containing 50 μg/mL Kanamycinstrain, then cultured at the permissive temperature of 37°C to the mid-log phase (for the control), or first incubated at the non-permissive temperature (20°C for SecY39, 42°C for SecA81) for 1 hour, then treated with chloramphenicol (34 μg/mL) for 30 min or 1 hour. Treated cells were then collected by centrifugation, re-suspended in loading buffer before incubated at 37°C for 10 min for semi native SDS-PAGE analysis. Protein samples were separated by SDS-PAGE before subjected to blotting analysis.

## Acknowledgements

We thank the Keio Collections for providing us the wild type *E. coli* strain. We thank Professor Koreaki Ito from Kyoto Sangyo University for providing us the SecY39 mutant strain. We thank Dr. Wen Zhou at the Mass spectrometry Facility of the National Center for Protein Sciences at Peking University for assistance in performing mass spectrometry analysis. We thank Mr. Yang Liu and Ms. Jiayu Yu for providing the LY928 strain and the pYLC plasmid vector. We thank Ms. Jimei Zhao for constructing the plasmids expressing the pBpA variants of SurA. We thank Ms. Pan Zou for providing us the plasmid expressing SecY-Avi. This study was supported by research grants from the National Natural Science Foundation of China (No. 31670775 and No. 31470766 to ZYC) and the national Basic Research Program of China (973 Program) (No. 2012CB917300 to ZYC).

## Conflict of Interest

We declare that we have no conflict of interest to this work.

**Author Contributions**

Feng Jin designed and performed the experiments, analyzed the data and prepared the manuscript. Zengyi Chang supervised this study and prepared the manuscript.

**References**

1.    Kreil, G. (1981) Transfer of proteins across membranes, *Annual review of biochemistry 50*, 317-348.

2.    Michaelis, S., and Beckwith, J. (1982) Mechanism of incorporation of cell envelope proteins in Escherichia coli, *Annual review of microbiology 36*, 435-465.

3.    Walter, P., Gilmore, R., and Blobel, G. (1984) Protein translocation across the endoplasmic reticulum, *Cell 38*, 5-8.

4.    Rapoport, T. A., Jungnickel, B., and Kutay, U. (1996) Protein transport across the eukaryotic endoplasmic reticulum and bacterial inner membranes, *Annual review of biochemistry 65*, 271-303.

5.    Schatz, G., and Dobberstein, B. (1996) Common principles of protein translocation across membranes, *Science (New York, N.Y.) 271*, 1519-1526.

6.    Chang, Z. (2016) Biogenesis of Secretory Proteins. In: Ralph A Bradshaw and Philip D Stahl (Editor-in-Chief), Encyclopedia of Cell Biology, Vol 1, Waltham, MA: Academic Press, 535-544.

7.    Osborne, A. R., Rapoport, T. A., and van den Berg, B. (2005) Protein translocation by the Sec61/SecY channel, *Annual review of cell and developmental biology 21*, 529-550.

8.    Wickner, W., and Schekman, R. (2005) Protein Translocation Across Biological Membranes, *Science (New York, N.Y.) 310*, 1452-1456.

9.    Papanikou, E., Karamanou, S., and Economou, A. (2007) Bacterial protein secretion through the translocase nanomachine, *Nat Rev Micro 5*, 839-851.

10.   Driessen, A. J., and Nouwen, N. (2008) Protein translocation across the bacterial cytoplasmic membrane, *Annual review of biochemistry 77*, 643-667.

11.   Tsirigotaki, A., De Geyter, J., Sostaric, N., Economou, A., and Karamanou, S. (2017) Protein export through the bacterial Sec pathway, *Nature reviews. Microbiology 15*, 21-36.

12.   Mitra, K., Frank, J., and Driessen, A. (2006) Co- and post-translational translocation through the protein-conducting channel: analogous mechanisms at work?, *Nature structural & molecular biology 13*, 957-964.

13.   Lazar, S. W., and Kolter, R. (1996) SurA assists the folding of Escherichia coli outer membrane proteins, *Journal of bacteriology 178*, 1770-1773.

14.   Rouviere, P. E., and Gross, C. A. (1996) SurA, a periplasmic protein with peptidyl-prolyl isomerase activity, participates in the assembly of outer membrane porins, *Genes & development 10*, 3170-3182.

15.   Sklar, J. G., Wu, T., Kahne, D., and Silhavy, T. J. (2007) Defining the roles of the periplasmic chaperones SurA, Skp, and DegP in Escherichia coli, *Genes & development 21*, 2473-2484.

16.   Wang, Y., Wang, R., Jin, F., Liu, Y., Yu, J., Fu, X., and Chang, Z. (2016) A Supercomplex Spanning the Inner and Outer Membranes Mediates the Biogenesis of beta-Barrel Outer

Membrane Proteins in Bacteria, *The Journal of biological chemistry 291*, 16720-16729.

17. Voulhoux, R., Bos, M. P., Geurtsen, J., Mols, M., and Tommassen, J. (2003) Role of a Highly Conserved Bacterial Protein in Outer Membrane Protein Assembly, *Science (New York, N.Y.) 299*, 262-265.

18. Wu, T., Malinverni, J., Ruiz, N., Kim, S., Silhavy, T. J., and Kahne, D. (2005) Identification of a multicomponent complex required for outer membrane biogenesis in Escherichia coli, *Cell 121*, 235-245.

19. Hagan, C. L., Silhavy, T. J., and Kahne, D. (2011) beta-Barrel membrane protein assembly by the Bam complex, *Annual review of biochemistry 80*, 189-210.

20. Rapoport, T. A. (2007) Protein translocation across the eukaryotic endoplasmic reticulum and bacterial plasma membranes, *Nature 450*, 663-669.

21. Deshaies, R. J., and Schekman, R. (1987) A yeast mutant defective at an early stage in import of secretory protein precursors into the endoplasmic reticulum, *The Journal of cell biology 105*, 633-645.

22. Gorlich, D., Prehn, S., Hartmann, E., Kalies, K. U., and Rapoport, T. A. (1992) A mammalian homolog of SEC61p and SECYp is associated with ribosomes and nascent polypeptides during translocation, *Cell 71*, 489-503.

23. Stirling, C. J., Rothblatt, J., Hosobuchi, M., Deshaies, R., and Schekman, R. (1992) Protein translocation mutants defective in the insertion of integral membrane proteins into the endoplasmic reticulum, *Molecular biology of the cell 3*, 129-142.

24. Emr, S. D., Hanley-Way, S., and Silhavy, T. J. (1981) Suppressor mutations that restore export of a protein with a defective signal sequence, *Cell 23*, 79-88.

25. Shultz, J., Silhavy, T. J., Berman, M. L., Fiil, N., and Emr, S. D. (1982) A previously unidentified gene in the spc operon of Escherichia coli K12 specifies a component of the protein export machinery, *Cell 31*, 227-235.

26. Ito, K., Wittekind, M., Nomura, M., Shiba, K., Yura, T., Miura, A., and Nashimoto, H. (1983) A temperature-sensitive mutant of E. coli exhibiting slow processing of exported proteins, *Cell 32*, 789-797.

27. Shiba, K., Ito, K., Yura, T., and Cerretti, D. P. (1984) A defined mutation in the protein export gene within the spc ribosomal protein operon of Escherichia coli: isolation and characterization of a new temperature-sensitive secY mutant, *The EMBO journal 3*, 631-635.

28. Brundage, L., Hendrick, J. P., Schiebel, E., Driessen, A. J. M., and Wickner, W. (1990) The purified E. coli integral membrane protein SecYE is sufficient for reconstitution of SecA-dependent precursor protein translocation, *Cell 62*, 649-657.

29. Berg, B. v. d., Clemons, W. M., Collinson, I., Modis, Y., Hartmann, E., Harrison, S. C., and Rapoport, T. A. (2004) X-ray structure of a protein-conducting channel, *Nature 427*, 36-44.

30. Li, L., Park, E., Ling, J., Ingram, J., Ploegh, H., and Rapoport, T. A. (2016) Crystal structure of a substrate-engaged SecY protein-translocation channel, *Nature 531*, 395-399.

31. Joly, J. C., and Wickner, W. (1993) The SecA and SecY subunits of translocase are the nearest neighbors of a translocating preprotein, shielding it from phospholipids, *The EMBO journal 12*, 255-263.

32. Cannon, K. S., Or, E., Clemons, W. M., Jr., Shibata, Y., and Rapoport, T. A. (2005) Disulfide bridge formation between SecY and a translocating polypeptide localizes the translocation pore to the center of SecY, *The Journal of cell biology 169*, 219-225.

33.    Oliver, D. B., and Beckwith, J. (1981) E. coli mutant pleiotropically defective in the export of secreted proteins, *Cell 25*, 765-772.

34.    Oliver, D. B., and Beckwith, J. (1982) Identification of a new gene (secA) and gene product involved in the secretion of envelope proteins in Escherichia coli, *Journal of bacteriology 150*, 686-691.

35.    Cabelli, R. J., Chen, L., Tai, P. C., and Oliver, D. B. (1988) SecA protein is required for secretory protein translocation into E. coli membrane vesicles, *Cell 55*, 683-692.

36.    Cunningham, K., Lill, R., Crooke, E., Rice, M., Moore, K., Wickner, W., and Oliver, D. (1989) SecA protein, a peripheral protein of the Escherichia coli plasma membrane, is essential for the functional binding and translocation of proOmpA, *The EMBO journal 8*, 955-959.

37.    Cabelli, R. J., Dolan, K. M., Qian, L. P., and Oliver, D. B. (1991) Characterization of membrane-associated and soluble states of SecA protein from wild-type and SecA51(TS) mutant strains of Escherichia coli, *The Journal of biological chemistry 266*, 24420-24427.

38.    Ulbrandt, N. D., London, E., and Oliver, D. B. (1992) Deep penetration of a portion of Escherichia coli SecA protein into model membranes is promoted by anionic phospholipids and by partial unfolding, *The Journal of biological chemistry 267*, 15184-15192.

39.    Chen, X., Xu, H., and Tai, P. C. (1996) A significant fraction of functional SecA is permanently embedded in the membrane. SecA cycling on and off the membrane is not essential during protein translocation, *The Journal of biological chemistry 271*, 29698-29706.

40.    Eichler, J., and Wickner, W. (1997) Both an N-terminal 65-kDa domain and a C-terminal 30-kDa domain of SecA cycle into the membrane at SecYEG during translocation, *Proceedings of the National Academy of Sciences of the United States of America 94*, 5574-5581.

41.    Young Jae, K., Rajapandi, T., and Oliver, D. (1994) SecA protein is exposed to the periplasmic surface of the E. coli inner membrane in its active state, *Cell 78*, 845-853.

42.    van der Does, C., den Blaauwen, T., de Wit, J. G., Manting, E. H., Groot, N. A., Fekkes, P., and Driessen, A. J. (1996) SecA is an intrinsic subunit of the Escherichia coli preprotein translocase and exposes its carboxyl terminus to the periplasm, *Molecular microbiology 22*, 619-629.

43.    Ramamurthy, V., and Oliver, D. (1997) Topology of the Integral Membrane Form of Escherichia coli SecA Protein Reveals Multiple Periplasmically Exposed Regions and Modulation by ATP Binding, *Journal of Biological Chemistry 272*, 23239-23246.

44.    Economou, A., and Wickner, W. (1994) SecA promotes preprotein translocation by undergoing ATP-driven cycles of membrane insertion and deinsertion, *Cell 78*, 835-843.

45.    Economou, A., Pogliano, J. A., Beckwith, J., Oliver, D. B., and Wickner, W. (1995) SecA membrane cycling at SecYEG is driven by distinct ATP binding and hydrolysis events and is regulated by SecD and SecF, *Cell 83*, 1171-1181.

46.    Yang, Y. B., Lian, J., and Tai, P. C. (1997) Differential translocation of protein precursors across SecY-deficient membranes of Escherichia coli: SecY is not obligatorily required for translocation of certain secretory proteins in vitro, *Journal of bacteriology 179*, 7386-7393.

47.    Lin, B. R., Gierasch, L. M., Jiang, C., and Tai, P. C. (2006) Electrophysiological studies in Xenopus oocytes for the opening of Escherichia coli SecA-dependent protein-conducting channels, *J Membr Biol 214*, 103-113.

48.    Hsieh, Y. H., Zhang, H., Lin, B. R., Cui, N., Na, B., Yang, H., Jiang, C., Sui, S. F., and Tai, P. C. (2011) SecA alone can promote protein translocation and ion channel activity: SecYEG

increases efficiency and signal peptide specificity, *The Journal of biological chemistry 286*, 44702-44709.

49. Lin, B.-R., Hsieh, Y.-H., Jiang, C., and Tai, P. C. (2012) Escherichia coli Membranes Depleted of SecYEG Elicit SecA-Dependent Ion-Channel Activity but Lose Signal Peptide Specificity, *The Journal of Membrane Biology 245*, 747-757.

50. Hsieh, Y.-h., Zhang, H., Wang, H., Yang, H., Jiang, C., Sui, S.-f., and Tai, P. C. (2013) Reconstitution of functionally efficient SecA-dependent protein-conducting channels: Transformation of low-affinity SecA-liposome channels to high-affinity SecA-SecYEG-SecDF·YajC channels, *Biochemical and biophysical research communications 431*, 388-392.

51. Zhang, H., Hsieh, Y. H., Lin, B. R., Yu, L., Yang, H., Jiang, C., Sui, S. F., and Tai, P. C. (2013) Specificity of SecYEG for PhoA precursors and SecA homologs on SecA protein-conducting channels, *Biochemical and biophysical research communications 437*, 212-216.

52. Chin, J. W., Martin, A. B., King, D. S., Wang, L., and Schultz, P. G. (2002) Addition of a photocrosslinking amino acid to the genetic code of Escherichiacoli, *Proceedings of the National Academy of Sciences of the United States of America 99*, 11020-11024.

53. Baba, T., Jacq, A., Brickman, E., Beckwith, J., Taura, T., Ueguchi, C., Akiyama, Y., and Ito, K. (1990) Characterization of cold-sensitive secY mutants of Escherichia coli, *Journal of bacteriology 172*, 7005-7010.

54. Rosenbaum, M., Halvorson, H., Preston, W. S., and Levinthal, C. (1955) Developmental of phage T2 desoxyribosenucleic acid in the absence of net protein synthesis, *Journal of bacteriology 69*, 228-229.

55. Nakatogawa, H., Mori, H., Matsumoto, G., and Ito, K. (2000) Characterization of a mutant form of SecA that alleviates a SecY defect at low temperature and shows a synthetic defect with SecY alteration at high temperature, *Journal of biochemistry 127*, 1071-1079.

56. Schagger, H. (2006) Tricine-SDS-PAGE, *Nature protocols 1*, 16-22.

57. Yang, Y. B., Yu, N., and Tai, P. C. (1997) SecE-depleted membranes of Escherichia coli are active. SecE is not obligatorily required for the in vitro translocation of certain protein precursors, *The Journal of biological chemistry 272*, 13660-13665.

58. Bessonneau, P., Besson, V., Collinson, I., and Duong, F. (2002) The SecYEG preprotein translocation channel is a conformationally dynamic and dimeric structure, *The EMBO journal 21*, 995-1003.

59. Breyton, C., Haase, W., Rapoport, T. A., Kuhlbrandt, W., and Collinson, I. (2002) Three-dimensional structure of the bacterial protein-translocation complex SecYEG, *Nature 418*, 662-665.

60. Dartigalongue, C., and Raina, S. (1998) A new heat-shock gene, ppiD, encodes a peptidyl-prolyl isomerase required for folding of outer membrane proteins in Escherichia coli, *The EMBO journal 17*, 3968-3980.

61. Oliver, D. B., and Beckwith, J. (1982) Regulation of a membrane component required for protein secretion in Escherichia coli, *Cell 30*, 311-319.

62. Takamatsu, H., Nakane, A., Oguro, A., Sadaie, Y., Nakamura, K., and Yamane, K. (1994) A Truncated Bacillus subtilis SecA Protein Consisting of the N-Terminal 234 Amino Acid Residues Forms a Complex with Escherichia coli SecA51(ts) Protein and Complements the Protein Translocation Defect of the secA51 Mutant, *The Journal of Biochemistry 116*,

1287-1294.

63.   Lee, D. J., Bingle, L. E., Heurlier, K., Pallen, M. J., Penn, C. W., Busby, S. J., and Hobman, J. L. (2009) Gene doctoring: a method for recombineering in laboratory and pathogenic Escherichia coli strains, *BMC microbiology 9*, 252.