Response of microbial community function to fluctuating

geochemical conditions within a legacy radioactive waste trench

environment

Xabier Vázquez-Campos[1,2], Andrew S. Kinsela[1,3], Mark W. Bligh[1,3], Jennifer J. Harrison[3],

Timothy E. Payne[3] & T. David Waite[1*]

[1]UNSW Water Research Centre and School of Civil and Environmental Engineering, The

University of New South Wales, Sydney, New South Wales 2052, Australia

[2]NSW Systems Biology Initiative, School of Biotechnology and Biomolecular Sciences, The

University of New South Wales, Sydney, New South Wales 2052, Australia

[3]Institute for Environmental Research, Australian Nuclear Science and Technology

Organisation, Locked Bag 2001, Kirrawee DC, New South Wales 2232, Australia

17 Conflict of interest: The authors declare no conflict of interest.

19 *Corresponding author: Professor T.D. Waite; Email: d.waite@unsw.edu.au

Abstract

During the 1960s, small quantities of radioactive materials were co-disposed with chemical waste at the Little Forest Legacy Site (Sydney, Australia) in three-metre-deep, unlined trenches. Chemical and microbial analyses, including functional and taxonomic information derived from shotgun metagenomics, were collected across a six-week period immediately after a prolonged rainfall event to assess how changing water levels impact upon the microbial ecology and contaminant mobility. Collectively, results demonstrated that oxygen-laden rainwater rapidly altered the redox balance in the trench water, strongly impacting microbial functioning as well as the radiochemistry. Two contaminants of concern, plutonium and americium, were shown to transition from solid-iron-associated species immediately after the initial rainwater pulse, to progressively more soluble moieties as reducing conditions were enhanced. Functional metagenomics revealed the potentially important role that the taxonomically-diverse microbial community played in this transition. In particular, aerobes dominated in the first day followed by an increase of facultative anaerobes/denitrifiers at day four. Towards the mid-end of the sampling period, the functional and taxonomic profiles depicted an anaerobic community distinguished by a higher representation of dissimilatory sulfate reduction and methanogenesis pathways. Our results have important implications to similar near-surface environmental systems in which redox cycling occurs.

Importance

The role of chemical and microbiological factors in mediating the biogeochemistry of groundwaters from trenches used to dispose of radioactive materials during the 1960s is examined in this study. Specifically, chemical and microbial analyses, including functional and taxonomic information derived from shotgun metagenomics, were collected across a six-week period immediately after a prolonged rainfall event to assess how changing water levels influence microbial ecology and contaminant mobility.

Results demonstrate that oxygen-laden rainwater rapidly altered the redox balance in the trench water, strongly impacting microbial functioning as well as the radiochemistry. Two contaminants of concern, plutonium and americium, were shown to transition from solid-iron-associated species immediately after the initial rainwater pulse, to progressively more soluble moieties as reducing conditions were enhanced. Functional metagenomics revealed the important role that the taxonomically-diverse microbial community played in this transition. Our results have important implications to similar near-surface environmental systems in which redox cycling occurs.

Keywords: shotgun metagenomics, radionuclides, plutonium, americium, functional profile.

Introduction

The rapid expansion of an emerging nuclear industry immediately following World War II resulted in substantial volumes of low-level radioactive waste (LLRW) being generated from nuclear fuel cycle, weapons production, medical radioisotope and radiochemical research activities. Although there was no consensus at this time, low-level waste (and in some cases more active material) was commonly disposed of by burial in shallow trenches, as evidenced in the United States at Maxey Flats (1), Oak Ridge (2) and Hanford (3, 4); Canada at Chalk River (5); in the United Kingdom at Harwell (6) and LLRW (7); in Lithuania at Maišiagala (8); and more recently in Ukraine at Chernobyl (9) to name but a few. This was also the case for Australia's only nuclear (research) reactor at Lucas Heights. Known as the Little Forest Legacy Site (LFLS), radioactive materials, including minor amounts of $^{239+240}$Pu and ^{241}Am were placed in narrow (0.6 m), 3 m deep, unlined trenches from 1960 to 1968 (10–12). Large volumes of contaminated non-radioactive materials and equipment were also disposed of in these trenches (10).

The LFLS trenches were excavated within undisturbed geological matrices of red-brown and grey clay (primarily kaolinite and illite/smectite) derived from the underlying weathered shale, interspersed with minor phases of hematite and goethite (11, 13). The site was chosen in part due to the low hydraulic conductivities of these materials (~ 9 to 66 mm/day) in order to isolate the LLRW from waters associated with the local hydrology cycle (10–12). However, an unintended consequence of this (also experienced at other disposal sites) has been that periodic intense rainfall and prolonged dry conditions can facilitate complete saturation and desaturation of the more permeable waste material via surface infiltration and evapo(transpi)ration/leakage mechanisms, respectively. Elaborating further, this frequently results in infiltrating water filling up the more porous trenches to the surface (akin to a 'bathtub'), which has been shown to be a primary mechanism for the dispersion of

contaminants $^{239+240}$Pu and ^{241}Am (10). Furthermore, this has allowed for redox cycling to

occur unabated in the LLRW trenches since their construction, potentially promoting redox

tolerant plasticity in the microbial communities present (14).

With actinide mobility, in many instances, strongly dependent on their oxidation state

(15), microbial communities can potentially play decisive roles in determining the fate and

mobility of such elements in the environment. In general, microbial communities can

influence actinide chemistry by partaking in redox (16–18), dissolution (19, 20), precipitation

(21, 22), sorption (23) and/or methylation (24) reactions, which may either enhance or retard

contaminant mobility.

Despite this, much of the scientific research to date has focussed on isolated individual

species (such as *Geobacter*, *Shewanella* and *Clostridium*) and their impact on actinide,

particularly uranium, behaviour. The challenge remains as to how best to identify the role of

microorganisms and their functioning within the wider microbial community which may be

undergoing external, environmental changes (25). Furthermore, subsurface environments

such as aquifers and shallow groundwaters, aside from being poorly studied, have been

shown to be havens for microbial novelty not dominated by the well-characterised organisms

listed above (26). Culture-independent techniques, such as metagenomics, in which genomic

sequences capture the aggregate microbial ecology of a sample (27, 28) have the potential to

enhance our understanding of these complex biogeochemical systems.

The question remains at LFLS, and in similar contaminated redox-cycling

environments, as to what function microbial communities may be performing in the direct or

indirect mobilisation and/or retention of legacy radionuclides. As the establishment of

causality between the legacy contaminants and microbial communities cannot be achieved in

such a non-invasive environmental study, the aim of this research was to understand the role

that periodic inundation from a large rainfall event, and presumably oxygen penetration, had

on the concomitant changes to chemistry-radiochemistry and microbial communities in a

LLRW trench environment. The two contaminants of major concern, Pu and Am, were the

focus of this research with shotgun metagenomics used to examine the microbial systems

function and taxonomy.

Materials and methods

Sample collection

Trench water samples were collected on five separate occasions across a two-month

period from LFLS. They were obtained from a screened PVC pipe which extends 1.55 m

below the ground surface. This pipe provides the only point of access into the legacy trenches

and was opportunistically installed during the partial collapse of the trench surface. Further

details of the trench sampler have been previously described in the literature (10, 29). Briefly,

a peristaltic pump operating under low flow rate was used to purge the borehole until

chemical parameters, particularly pH and oxidation/reduction potential (ORP), became stable

(Supplementary File 1). Measurements were made using a multi-probe system (YSI 556

MPS) coupled to a 100 mL flow cell. Stability usually took between 30 and 60 min, after

which sample collection began. Chemical parameters, along with the water depth, were

monitored over the sampling period (typically some hours) on the day of sampling to ensure

the water was representative of that particular period. The stability of flow cell parameters

and the fact that the water level did not decrease during the collection of 4–5 L of sample

material provides assurance that the samples were representative of the particular event rather

than anomalous local conditions.

The sample collection period commenced (on 23[rd] April 2015) immediately after a

substantial rainfall event, during which ~220 mm of rain fell over 3 days (Figure S1). Trench

water samples were subsequently collected on the 27[th] and 29[th] April, 14[th] May and 9[th] June

(or days 0, 4, 6, 21 and 47, post-rain, Figure S1).

Samples for chemical analyses were collected both with and without an in-line filtration

membrane (0.45 µm PES, Waterra FHT-Groundwater) in pre-washed and trench water rinsed

HDPE bottles except samples for organic carbon analysis which were collected in glass

bottles. Samples for radiochemical and cation analyses were acidified using double-distilled

HNO_3. All samples were then stored at 4 °C until analysis.

Samples for microbial analyses were collected by directly filtering the trench water

using a sterile PES 0.22 µm syringe filter (Sterivex-GP, Merck Millipore) with the volumes

of filtrate passing through the membranes recorded (410.1±82.2 mL per replicate, mean ±

SD). Filters were capped on site before being stored at -18 °C until analysis. Microbial

sampling was performed in triplicate to assess inherent variation.

Due to regulatory protocols governing sampling at the site, no solid phase material

could be extracted from the trenches. As the surrounding geological materials into which the

trenches were excavated provide little resemblance both physically and chemically to the

deposited waste within the trenches, they were discarded as providing useful information for

this study. As such all sampling within the trenches pertains to the extraction of soluble or

suspended solid-phases from the screened borehole.

Chemical analyses of trench waters

The primary radiochemical contaminants, $^{239+240}$Pu and ^{241}Am, were separated from

other radiochemical fractions using TEVA™ and TRU™ resin cartridges (30) before

measuring activity by alpha spectroscopy using a Canberra Alpha Analyst system coupled

with Passive Implanted Planar Silicon detectors (Canberra).

Cations (including Si, S and P) were measured by either ICP-AES or ICP-MS

depending on relative concentration. Anions (F^-, Cl^-, Br^-, I^-, PO_4^{3-} and NO_3^-) were determined

by ion chromatography (Dionex DX-600 IC System).

Non-purgeable dissolved organic carbon (0.45 µm filtered) was measured on freshly-

collected replicate samples by combustion catalytic oxidation (TOC-5000A, Shimadzu) with

concentration determined by a five-point calibration using newly-prepared potassium

hydrogen phthalate solutions.

Ferrous iron concentrations from 0.45 µm filtered samples were preserved in the field

with ammonium acetate-buffered phenanthroline solution (31). The absorbance of Fe(II) was

measured at 510 nm (USB4000, Ocean Optics) calibrated against a freshly-prepared

ammonium ferrous sulfate solution.

In field measurements of dissolved sulfide were attempted using methylene blue

colorimetry, but were consistently below detection limits (~1 µM) at all sampling time points.

All chemical analyses were performed in triplicate.

DNA extraction and sequencing

The PowerLyzer® PowerSoil® DNA Isolation kit (MO BIO) was used to extract DNA

from Sterivex filters following the modified protocol of Jacobs et al (32). Bead beating was

performed singularly in a PowerLyzer® 24 Bench Top Bead-Based Homogenizer (MO BIO)

at 2000 rpm for 5 min.

Concentration and DNA quality was evaluated with LabChip® GX (PerkinElmer).

Libraries were prepared with the TruSeq Nano DNA Library Preparation Kit and sequenced

in a NextSeq 500 (Illumina) with a 2×150 bp high output run. Output consisted of paired-end

reads with a median insert size of ~480 nt. Reads were deposited at the European Nucleotide

Archive (PRJEB14718/ERG009353).

Data processing and functional analyses

Raw   sequencing   reads   were   pre-processed   with   Trim   Galore!

(http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) to remove residual

adapters as well as short and low quality sequences (Phred33 $\geq$ 20) while keeping unpaired

reads.

Taxonomic profiles were obtained from shotgun metagenomic data by extracting 16S

rDNA gene sequences with GraftM v0.10.1 (https://github.com/geronimp/graftM) and

mapping them with the 97% clustered GreenGenes database provided by GraftM developers

(package 4.39).

Functional analyses of the metagenomic data were performed with HUMAnN2 v0.6.0

(33). Gene families were grouped by MetaCyc reactions (RXN), keeping unmapped and

ungrouped reads for calculating copies per million and relative abundances. MetaCyc RXN

groups were filtered and only those with at least two replicates from the same sampling time-

point with a relative abundance of 10^{-4} or higher were analysed (33). MetaCyc RXN

notations have been used as is within Results and Discussion to provide an exact match with

the database. RXNs can be considered equivalent to the more traditionally used KO terms

from the KEGG database.Grouping by GO terms, specifically performed to evaluate virus

presence, indicated values under the threshold levels for all the virus-specific GO terms

identified. As such, viruses were not taken in further consideration.

Analysis of variance was performed with the software package STAMP v2.1.3 (34)

using the Benjamini-Hochberg False Discovery Rate approach (35) for correction of *p*-

values.

Results and Discussion

Water level changes and chemical analyses

The initial 220 mm rainfall event resulted in trenches filling to capacity and discharging

from the surface or porous near-surface (0–0.2 m) in the 'bathtub' mechanism described by

Payne et al (10) (Figure S1). Despite the subsequent 47-day sampling period receiving a

further 72 mm of rainfall, which periodically increased trench water levels, an overall decline

in trench water levels was observed across the sampling period (Figure S1).

Across the sampling period and as the water level declined, the pH was observed to

increase from 6.30 to 6.60, whereas the E_h values decreased from 247 to 147 mV (Figure 1).

Of the cations and anions analysed (Figure 2), iron displayed one of the greatest variations in

concentration, increasing from 0.42 mM at day 0, to 1.00 mM by day 47. The elevated

concentrations of iron are unsurprising, given that the LLRW at LFLS was disposed of in-

part within ~760 steel drums (36), and buried within a highly weathered shale geological

matrix (11). The continuous presence of Fe(II), even at day 0, confers reducing conditions in

excess of E_h values recorded, casting uncertainty over the absolute values supplied by the E_h

probe. The increase in Fe(II) concentrations with time was likely due to Fe(III) oxyhydroxide

reduction, an observation supported by the concurrent liberation of Si and P, two elements

that are typically co-associated with Fe(III)-oxyhydroxide at LFLS (37). The constant

concentrations of expected conservative elements Cl⁻ and K suggest that the drop in the

trench water level was due to subsurface outflow rather than evaporative processes across the

course of sampling. Other important elemental transitions included a 10-fold decrease in

dissolved sulfur concentrations (presumed to be sulfate), from an initial concentration of 62.4

to 6.2 µM at day 47 (Figure 2). Nitrate concentrations doubled between days 0 and 6 (from

0.24 to 0.55 µM) after which they decreased to below detection limits at days 21 and 47

(Figure 2).

The total (unfiltered) ^{241}Am activity increased from 15.7 to 27.8 Bq/L (1.8-fold

increase) while the soluble (filtered) fraction increased 3.4-times (7.2 to 24.7 Bq/L) between

days 0 and 47 (Figure 3). Although the activity of the total $^{239+240}$Pu showed a proportionally

smaller increase than ^{241}Am across the sampling period, from 30.4 to 45.6 Bq/L (1.5 times),

the soluble fraction increased from 0.21 at day 0 and reached a maximum of 0.80 (35.1 Bq/L)

by day 21. Despite the low-flow sampling conditions, the majority of ^{241}Am and $^{239+240}$Pu in

the trench water was solid-associated at Day 0. This implies that the solid-associated

actinides are relatively stable or easily (re)mobilised, and also that during the rapid influx of

rainwater when the trenches are filling up (and potentially overspilling), colloidal transport is

likely to be the major form of contaminant movement at this site. Previous redox state

measurements at LFLS found that Pu was present in the Pu(IV) (64.5%) and colloidal/Pu(III)

(35.5%) states whilst Am was present exclusively as Am(III) (29). The vast quantities of

Fe(II) as well as the results of ancillary measurements (DO, ORP) would suggest that Pu(IV)

and Am(III) were again the dominant oxidation states of the trench water contaminants. The

results of all water quality parameters measured are provided in Table 1.

Community composition

Community profiles showed that *Bacteria* dominated over *Archaea* in the trench water

across the entire sampling period. A total of 70 phyla out of the 85 included in the

GreenGenes database were detected at some point and 11 phyla with values over 1% of the

total community at all times. Eukarya were found in low (~1% to ~2%), nearly constant

abundance trough the samples. Similarly, eukaryotic-specific RXNs were all below our

threshold significance level. As such they are not considered in further detail nor for the taxa

abundance numbers throughout the manuscript.

Archaea oscillated between 2.6 and 10.6% of the classified reads with minimum and

maximum at days 4 and 47, respectively. *Micrarchaeota* and *Parvarchaeota* (DPANN) were

the most abundant phyla at all times with a combined 45.5–55.9% of *Archaea*, while

remaining sequences were shared in variable proportions between the TACK superphylum

and *Euryarchaeota* (Figure 4). Although the fraction of TACK decreased over time from

24.8% at day 0 to 12.4% at day 47, *Euryarchaeota* reached a maximum at day 47

contributing 37.2% of all *Archaea*. These changes were derived mainly from variations in the

SAGMA-X family (*Thaumarchaeota*), related to the ammonia oxidising archaeon '*Ca.*

Nitrosotalea' (38, 39) and, in *Bathyarchaeota* (MCG), which includes the only potential non-

euryarchaeal methanogens (39, 40) and/or one of the few acetogenic *Archaea* (41). The

increase in total *Archaea* abundance over time was related to changes to and ANME-2d,

Methanoregulaceae and *Methanosaetaceae* families, implicated in methane metabolism (42–

44) as well as the *Micrarchaeota* and *Parvarchaeota* phyla.

Only 8 bacterial phyla grouped more than 5% of the total bacterial reads at any one

sampling point, and amongst them, only *Chloroflexi*, OD1 (*Parcubacteria*), OP11

(*Microgenomates*), OP3 (*Omnitrophica*) and *Proteobacteria* were within the 10 most

abundant bacterial phyla at all times (Figure 4). The principal changes observed over time

refer to *Proteobacteria*, TM7 (*Saccharibacteria*), OP3, *Acidobacteria* and OD1. While the

proportion of TM7 and *Acidobacteria* decreased from day 0, both OP3, and OD1 reached

their maximum by day 47 (18.9% and 14.7%). The *Proteobacteria*, often the most abundant

phylum, showed a conspicuous increase, mainly due to *Betaproteobacteria*, comprising

48.7% of all classified reads at day 4. The order *Burkholderiales* was identified as being

primarily responsible for this change (42.6% of total), esp. the genus *Ralstonia* (20.3%)

within *Oxalobacteraceae* (34.0%). These bacteria were also responsible for the altered

functional profile observed at day 4.

Several studies focusing on the microbiology of groundwater or subsurface ecosystems

have demonstrated the existence of organisms able to pass through standard 0.22 µm filters

(45–47). Despite this, a substantial portion of the Little Forest trench water community

corresponded to taxa known to be able to pass through 0.22 µm filters, i.e. OD1, OP11 and

DPANN. We suggest this could be derived from the fast precipitation of iron in our extracted

trench waters, effectively entrapping cells, or that they were attached to colloidal particles or

to other cells. Still, it is possible that the numbers for these 'ultra-small' taxa could be

underestimated in our results.

A complete taxonomic profile of all sample replicates at each individual time point is

provided in Supplementary Information. An important aside to be noted with this data is the

inherent reproducibility between sample triplicates. This finding provides a level of

reassurance with regards to the aggregate nature of these water samples, often neglected in

similar studies.

Carbon cycling

Initial time points were characterised by a significantly ($p \leq 0.05$) higher relative

abundance of RXNs using molecular dioxygen as substrate such as: cytochrome-c oxidase

(CYTOCHROME-C-OXIDASE-RXN, EC:1.9.3.1, Figure 5A), quinol–cytochrome-c

reductase (1.10.2.2-RXN) and several mono- and dioxygenases. This was combined with a

significantly ($p \leq 0.05$) lower relative abundance of methanogenesis-related RXNs,

nitrogenase (ferredoxin, NITROGENASE-RXN), rubisco (RIBULOSE-BISPHOSPHATE-

CARBOXYLASE-RXN), or heterolactic fermentation (PHOSPHOKETOLASE-RXN).

Catalase (CATAL-RXN, EC:1.11.1.6/21, see Reactive oxygen species detoxification in SI)

gene relative abundance was highest at day 4 along with several ABC transporters and

phosphotransferases, e.g. D-ribopyranose, D-xylose, spermidine or L-arginine, as well as

assimilatory sulfur and nitrogen pathway RXNs (refer to SI for details). The higher relative

abundances of all these genes at day 4 suggest a metabolism more dependent on oxygen, i.e.

aerobic, or at least microaerophilic (see Nitrogen cycling). It also confers a higher

dependence on available organic carbon (viz. heterotrophy), or on the presence of decaying

complex organic matter (48). These interpretations are collectively supported by the

geochemical measurements including the higher E_h values, decreasing concentrations of

sulfur and absence of nitrate, along with lower iron concentrations across the first sampling

points.

Regarding the source of the organic fraction, the results suggest that soil particles and

associated organo-chemicals were mobilised from material above the trenches via advective

transport mechanisms during and immediately after rainfall. This was evidenced by the

increased proportion of soil-associated *Actinobacteria* at day 0, which diminished to 0.71%

by day 4 (Figure 4). Furthermore, the presence of genes encoding for enzymes representing

chitin degradation, such as chitobiase (RXN-12625, EC:3.2.1.52), diminished progressively

until day 47 indicating a potential one-off provision of chitin, matching the leaching

hypothesis (Figure S2). In addition, numerous facultative and strict anaerobes can degrade

chitin and/or chitobiose (49–51), which lessens the likelihood of the alternative hypothesis

that chitin depletion could be derived from the intrinsic lack of chitobiase in anaerobes.

The other primary biopolymer found in soil is cellulose (52). The relative abundance of

cellulase (RXN-2043, EC:3.2.1.4) showed no significant differences between days 0, 21 and

47 ($p = 0.172$) suggesting a consistent supply of cellulose (Figure 5B). The exception, noted

at day 4, can be explained by the proliferation of *Burkholderiales*. The constant abundance of

cellulose is unsurprising given the historical records which indicate a range of organic

(particularly cellulose-based) compounds were co-disposed of in the LFLS trenches (11).

This hypothesis accords with previous stable isotope measurements, which identified the

degradation of legacy organic matter as being responsible for enriched δ^{13}C (inorganic)

values measured near the trench (11).

Lignin is present to a lesser or greater extent in all plant biomass. From the list of

enzymes in the literature capable of degrading lignin (53, 54), only below-threshold levels of

certain genes were detected: versatile peroxidase (RXN0-267, EC:1.11.1.16), dye peroxidase

(RXN-11813, EC:1.11.1.19) and laccase (LACCASE-RXN, EC:1.10.3.2). The overall lack of

genes for proteins capable of lignin degradation strengthens the hypothesis that trench waste

material is the source of cellulosic material rather than being derived from the degradation of

plant material from the topsoil.

The two fundamental genes representing the glyoxylate pathway, isocitrate lyase

(ISOCIT-CLEAV-RXN, EC:4.1.3.1, Figure S3) and malate synthase (MALSYN-RXN,

EC:4.1.3.2, Figure 5C) showed between 3–4 times higher relative abundance in days 0 and 4

when compared to day 47. This pathway facilitates the use of short chain carbon compounds

(C2–C3) for anabolic purposes, compounds especially generated during anaerobic processes.

Although the glyoxylate pathway genes have been previously detected in organisms living in

anaerobic environments (55), it was absent from strict anaerobic organisms based on the

results from EggNOG (56), KEGG (57) and UniProtKB (58) databases (as of 24/05/2017).

This finding supports the hypothesis that trench waters became anoxic as the water level

declined, promoting conditions which favour the development of strict anaerobes.

Furthermore, the aerobic community which developed immediately after the rainfall event

likely utilised (and benefited from) the short chain carbon compounds generated by preceding

anaerobic conditions.

Both acetoclastic and hydrogenotrophic methanogenesis pathways were evident from

the functional profile analysis. Similar maximum relative abundances, about $1.4 \cdot 10^{-4}$, of 5-

methyltetrahydrosarcinapterin:corrinoid/iron-sulfur protein *Co*-methyltransferase (RXN-

12908, EC:2.1.1.245, Figure 5D) and coenzyme F_{420}-dependent hydrogenase (COENZYME-

F420-HYDROGENASE-RXN, EC:1.12.98.1, Figure S4), which are specific RXNs for the

methanogenesis from acetate and H_2 and CO_2, were observed at day 47. The presence of

methanogenesis-specific RXNs is supported by previous isotopic measurements, showing

δ^2H enrichment, relative to $\delta^{18}O$, in the vicinity of the trenches (11). Concurrently, the

taxonomy revealed the presence of the ANME-2d division contributing up to 13.8% of all

Archaea at day 47 (1.5% of the community). The ANME-2d division was initially linked to

the anaerobic oxidation of methane using NO_3^- (42) or even NO_2^- (59), constituting one of the primary methane sinks in anaerobic conditions. However, recent research has shown that some of the members of this taxa are able to utilise Fe(III) in place of NO_2^- or NO_3^- (44, 60, 61). This provides a more reasonable explanation given the limited nitrate present and abundance of iron during the later sampling days.

Sulfur cycling

The relative abundance of sulfite cytochrome *c* reductase (SULFITE-DEHYDROGENASE-RXN, EC:1.8.2.1, Figure S5), involved in the assimilatory reduction of sulfate, peaked at day 4, with ~10 times higher relative abundance compared to day 0. At the same time, dissimilatory sulfate reduction RXNs, e.g. dissimilatory sulfite reductase (HYDROGENSULFITE-REDUCTASE-RXN, EC:1.8.99.3), followed the opposite trend, with a maximum at day 47 (Figure 5E). The combined relative abundance of all the putative sulfate reducing bacteria (SRB) taxa detected (mainly *Syntrophobacterales*, *Thermodesulfovibrionaceae* and *Desulfobacterales*) also increased over time, from 2.3% of the total prokaryotes at day 0 to 4.9% at day 47. Dissimilatory sulfate reduction is one of the major redox processes in both natural (62) and artificial (63) anaerobic environments. The ten-fold reduction in the sulfur concentration is thought to result from the loss of dissolved sulfate via reduction to sulfide and subsequent precipitation of insoluble heavy metal sulfides, particularly FeS. This interpretation is supported by the lack of measurable sulfides in the trench water (see Chemical analyses of trench waters), along with previous reports showing sulfate reduction based on isotopic fractionation and severe (10 to 100-fold) depletion in sulfate concentrations in trench waters relative to surrounding wells (11). The potential contribution of sulfate- and nitrite-dependent anaerobic methane oxidation (S-DAMO and N-DAMO respectively) was discounted due to the near complete absence of ANME taxa (*Archaea*) aside from ANME-2d, and NC10 (*Bacteria*) (see Nitrogen cycling) (44, 59, 64,

378   65).

Nitrogen cycling

The (inorganic) nitrogen cycle was mostly represented by RXNs related to

denitrification and assimilatory nitrogen reactions, primarily dissimilatory nitrate reductase

(RXN-16471, EC:1.7.5.1, see Figure S6 for a complete profile of the N-associated RXNs),

although nitrogen fixation (NITROGENASE-RXN, EC:1.18.6.1) was the predominant N-

associated RXN. All other prominent N-associated RXNs peaked at day 4 with the exception

of hydroxylamine reductase (HYDROXYLAMINE-REDUCTASE-RXN, EC:1.7.99.1) and

ammonia oxygenase (AMONITRO-RXN, EC:1.14.99.39). The increase in nitrate

concentrations observed between days 4 and 6 (0.236 and 0.552 µM respectively) might be

derived from the oxidation of NO by nitric oxide dioxygenase (R621-RXN, EC:1.14.12.17).

This surge in the relative abundance of nitrogen metabolism RXNs (at day 4) coincided with

an increased dominance of *Betaproteobacteria*, mostly *Burkholderiales*. *Betaproteobacteria*

have been linked to denitrification processes in a uranium-polluted aquifer in Rifle (USA)

(66) and they may play a similar ecological role at LFLS. However, the increase in nitrate

concentrations could equally be indicative of NO or NO_2^- oxidation, originally produced by

the aerobic oxidation of ammonium by the *Thaumarchaeota* (67, 68).

The inorganic nitrogen cycling results provide important information regarding the

transition from aerobic to anaerobic conditions as the water level declines. The competing

requirements of nitrate respiration, which necessitates oxygen-limiting conditions at a

minimum (69), and NO dioxygenase needing molecular oxygen, collectively suggest that at

day 4 the trench waters were no longer aerobic, but more likely microaerophilic/hypoxic.

However, many facultative anaerobes are capable of using nitrate as an alternate electron

acceptor when oxygen is not available. As such, the high relative abundance of nitrate

reducing enzyme genes at day 4 (viz. dissimilatory nitrate reductases) could be a confounding

effect associated with the increased abundance of *Burkholderiales*. This was confirmed by

searching the denitrifying pathway in KEGG (map00910) (as of 08/07/2016) (57).

Iron cycling

Temporal increases in Fe(II) concentrations, such those experienced in the later anoxic

periods of the sampling, may be associated with the use of Fe(III) during anaerobic

respiration, previously attributed to decaheme *c*-type cytochromes from the OmcA/MtrC

family present in iron-reducing bacteria (FeRB) such as *Shewanella* spp. and *Geobacter* spp.

(70). However, their representation in the trench water metagenome data was scarce (relative

abundance $\leq 10^{-6}$). It is known that for respiration to occur, FeRB often require direct

physical contact with Fe(III) solids (71). As only aqueous/suspended phase sampling was

permissible at LFLS, it is possible the under representation of a FeRB community was due to

our sampling regime in which the solid-phase was largely excluded.

The genes associated with RXNs specific to the Fe(II) oxidation pathway (via

rusticyanin, PWY-6692): ubiquinone-cytochrome-*bc1* reductase (EC:1.10.2.2, RXN-15829)

and *aa3*-type cytochrome oxidase (EC:1.9.3.1, RXN-15830) were present, despite no iron-

rusticyanin reductase (EC:1.16.9.1, RXN-12075), *cyc2*, being found under the established

threshold (maximum value $\sim 3 \times 10^{-6}$) (72). The presence of both cytochromes could be

because HUMAnN2 could not distinguish between the 'generic' quinol–cytochrome-c

reductase (RXN-15816, EC:1.10.2.2), and cytochrome *c* oxidase (CYTOCHROME-C-

OXIDASE-RXN) respectively, particularly when analysing short reads.

The presence of 1.9% of *Gallionellaceae* (over the total community composition) at

day 0 and 2.9% of *Crenothrix*, a non-FeOB usually associated with Fe-mineralising biofilms

(73), suggest the existence of transient microaerophilic iron oxidising activity in the trench

water despite the extremely low relative abundance of *cyc2*. Waste disposal records from

LFLS indicate that numerous steel drums (~760) were co-disposed of in the trenches (36).

Based on the active sulfate respiration observed in the trench waters, we would expect that

iron/steel materials may have suffered microbially induced corrosion (MIC) (74), potentially

contributing to the elevated Fe(II) concentrations. However, the main iron MIC product is

likely FeS (74), which can be utilised by *Gallionella ferruginea* (*Gallionellaceae*), an iron

oxidising bacterium (FeOB) (75). Furthermore, this process would also provide an

explanation for a persistent source of Fe(III) (oxy)hydroxides and their ongoing interaction

with the S-cycle.

Previous research on soils experiencing fluctuating redox conditions and active iron

cycling have shown they frequently lack 'typical' Fe(III)-respiring bacteria (76), being

outcompeted by sulfate respiration (77). Despite Fe(III) being a more thermodynamically

favourable electron acceptor it is commonly acknowledged that sulfate respiration often

outcompetes Fe(III) respiration in both high and low sulfate environments (77, 78). High

levels of sulfate reduction in low-sulfate environments, akin to the LFLS trenches, has been

previously observed to occur in the presence of crystalline Fe(III) oxyhydroxides that

partially re-oxidise sulfide generated by SRB to elemental sulfur (78). This mechanism

provides a plausible explanation for our observations at LFLS. The large concentration of

dissolved Fe(II) in the trench waters would likely contribute to the Fe(II)-catalysed

transformation of amorphous ferrihydrite to more crystalline, thermodynamically-favourable

forms (79, 80), although this would be limited to some extent by the large concentrations of

dissolved organic matter and silica (81). Furthermore, the abiotic reduction of crystalline

Fe(III) oxyhydroxides would severely limit the energy acquisition by FeRB and therefore the

FeRB would only outcompete SRB when sulfate and other reducible sulfur compounds were

totally consumed.

Alternatively, fermentative bacteria coupling the oxidation of a range of organic

compounds to the reduction of ferric iron (82–84) may play an integral role in maintaining

the elevated Fe(II) concentrations observed within the trenches. One example is the organism

Propionibacterium freudenreichii which has previously been observed to reduce Fe(III) by

using humic substances as an electron mediator (85). In this regard, both heterolactic and

propionic acid fermentation pathways showed high relative abundances based on

phosphoketolase (PHOSPHOKETOLASE-RXN, EC:4.1.2.9) and methylmalonyl-CoA

decarboxylase (RXN0-310, EC:4.1.1.41), respectively (Figure S7). Their decrease in relative

abundance (at day 4) could be explained by the lack of fermentative pathways present in the

genomes associated with the population shift, i.e. increase in *Burkholderiales*. This was

confirmed by searching (as of 08/06/2016) methylmalonyl-CoA decarboxylase and

phosphoketolase in the KEGG maps (map00640 and map00030, respectively).

Synthesis of trench processes and radiochemical mobilisation implications

The biogeochemistry of this dynamic system is conceptually described by the elemental

cycling schematic shown in Figure 6. Immediately following the rainfall event, the E_h

reached its most oxidising value (247 mV) with the microbial community characterised by

higher relative abundances of pathways related to aerobic (or at least microaerophilic)

heterotrophic metabolism and one capable of chitin degradation. Although conditions were

not strongly oxidising, this pulse of oxic water would be sufficient to induce the abiotic

oxidation of part of the large store of dissolved Fe(II), likely forming a combination of the

reactive Fe(III) oxyhydroxides ferrihydrite and silica-ferrihydrite, along with the more

crystalline oxyhydroxide lepidocrocite, as has previously been observed in these trench

waters (37). As both Am(III) and Pu(III)/(IV) are known to strongly sorb to sediments and

iron oxides (15), it is of little surprise to observe the greatest proportions of (suspended)

solid-associated actinides at this time point (day 0, Figure 3). Note that even though most Am

(54.3%) and particularly Pu (78.8%) were associated with a solid-fraction >0.45 µm, they

were still extracted from the trench under our low-flow sampling method. This finding

implies that colloid-associated Pu, and to a lesser extent Am, remains mobile within the

trench waters, congruent with observations from other legacy radioactive waste sites (86).

Interestingly though, the Pu migration distance away from source at LFLS in groundwater

has been shown to be much smaller (~1 order of magnitude less) compared to other legacy

locations such as the Nevada Test Site (87), Rocky Flats (88) and Mayak (89). We attribute

this to a low permeability soil matrix, inhibiting downward migration to the

connected/permanent water table, coupled with the biogeochemical conditions within the

trenches themselves. The high concentrations of Fe(II), circumneutral pH and proliferation of

aerobic heterotrophs drives the rapid formation of large quantities of Fe(III) oxides. The

quantity of Fe(III) oxides that form upon oxic rainwater intrusion is evidently sufficient to

contain the bulk of contaminants within the trenches during 'bathtub' overflow events. The

rate of Fe(II) oxidation is likely to be crucial for the ongoing attenuation of Pu and Am at

LFLS (37).

By day 4, as water levels rapidly decrease, an increase of *Burkholderiales*

(*Betaproteobacteria*) generates a functional profile disturbance and, while the aerobic profile

is maintained, it is also the time-point at which nitrogen cycling (e.g. nitrate respiration)

becomes most active (Figure 6; ▯). Therefore, day 4 represents a potential transition away

from oxic conditions derived from rain infiltration, though this sequence of events is

somewhat confounded by the microbial growth lag phase associated with aerobic respirators.

Over the following weeks (days 21 and 47), the microbial community transitions to a

functional profile dominated by carbon fixation, methanogenesis and sulfate respiration

pathways (Figure 6; ▯▯▯▯). The increase in anaerobicity correlated with increasing

concentrations of soluble Fe and soluble radionuclides and a depletion of sulfate and nitrate.

In the case of Am, although the total activity increased gradually as the water level in the

trench declined, the soluble fraction increased by a greater proportion (Figure 3). This indicates that either a desorption or dissolution process occurred. The concomitant increase in the soluble Am fraction and Fe(II) concentrations point towards reductive dissolution of Fe(III) oxyhydroxides as a major driver behind Am solubilisation.

The Pu behaviour in the trenches presents a more complex temporal dynamic due, in part, to its more varied redox chemistry. As conditions become more reducing over time, one would expect to observe soluble Pu activities increase, through both the dissolution of Fe(III) oxides and, to a lesser extent, from Pu(IV) to Pu(III) reduction. In an interesting point of difference to Am, our results show that Pu proportionally remained in the particle-associated fraction for a substantially longer period, even as the dissolved Am(III) activity and Fe(II) concentration increased. The reason for the differing solution/solid-phase partitioning observed for Pu and Am is not clear based on the evidence to hand. The inability to sample solid materials from within the trenches has limited more conclusive understanding. However, we suggest the differing Am and Pu associations are likely due to different redox states, based on the previous measurements at LFLS, which showed a dominant Pu oxidation state of +4 (29). Electrostatically, Pu(IV), as the neutral hydrolysed cation $Pu(OH)_4(aq)$ or as an intrinsic colloid, is more likely sorbed or incorporated into positively charged pseudo-colloids such as Fe(III) oxides, compared to Am^{3+}. This is supported by research showing the highly reversible nature of Am(III) sorption onto poor crystalline iron colloids (90) and association with carbonate- and exchangeable sites on clays (91).

Conclusions

The inability to comprehensively access and sample within legacy radioactive waste environments hampers our ability to comprehend co-occurring elemental cycling and microbial metabolism, potentially curtailing our ability to effectively manage and remediate

such sites. The trench-sampling point at LFLS is therefore a particularly useful resource for

such research. In this study, our coupled use of metagenomics and chemical analyses has

provided a previously unattainable level of understanding for the LFLS trench water,

highlighting the responsiveness of the microbial community to external changes and dynamic

nature of the resulting chemistry. The combined results show that the trench waters contain a

taxonomically diverse microbial community, which has likely evolved in response to

variations in energy sources supplied by frequent redox fluctuations. When combined with

the complex nature of the waste-form, a myriad of micro-environments has developed within

the trenches, allowing for simultaneous O, N, Fe, S and C elemental cycling, as shown by co-

occurring metabolic reactions in the aggregate water samples. Consequently, it can be

inferred that Pu and Am are subject to persistent reducing conditions (as evident from active

iron oxide dissolution, sulfate reduction and methanogenesis) when the water level is low

between rainfall events. These reductive processes maintain Pu and Am solubility, despite the

occasional onset of oxidising conditions associated with rainfall events. Ultimately however,

the high concentrations of Fe present and the tendency of Fe(II) to be relatively rapidly

oxidised to strongly sorbing Fe(III) (oxy)hydroxide solids on exposure to oxic conditions

results in limited transport of Pu and Am.

Although the findings described above are intrinsically linked to the specific site under

investigation, they provide important generic insights into the dynamic biogeochemical

behaviour of iron-rich, redox-cycling environments. Of particular interest is the rapid

response of the microbial community to dynamic redox conditions and the potential impact

upon persistent contaminant solubility and enhanced mobility.

Acknowledgements

549 ANSTO staff members are thanked for their valuable contributions: Kerry Wilsher

550 (fieldwork, laboratory analyses), Sangeeth Thiruvoth (actinides), Brett Rowling (fieldwork,

551 inorganics), Henri Wong (inorganics), Chris Vardanega (inorganics) and Dioni Cendón

552 (discussion). The Australian Bureau of Meteorology is acknowledged for rainfall data.

Author contribution

554   Study concept and experimental design: XVC, ASK, TDW. Fieldwork: ASK. Chemical

555 and radiochemical analyses: ASK, JJH. Molecular biology and bioinformatics analyses:

556 XVC. Contribution of reagents/materials/analysis tools: TEP, TDW. Manuscript writing:

557 XVC, ASK. Revision and review of the manuscript: XVC, ASK, MWB, TDW.

References

1. Cleveland JM, Rees TF. 1981. Characterization of plutonium in Maxey Flats radioactive trench leachates.
Science 212:1506–1509.

2. Saunders JA, Toran LE. 1995. Modeling of radionuclide and heavy metal sorption around low- and high-
pH waste disposal sites at Oak Ridge, Tennessee. Appl Geochem 10:673–684.

3. Cantrell KJ, Felmy AR. 2012. Plutonium and americium geochemistry at Hanford: a site-wide review.
RPT-DVZ-AFRI-001 PNNL-21651. Pacific Northwest National Laboratory, Washington (USA).

4. Gerber MS. 1992. Past practices technical characterization study — 300 area — Hanford site. WHC-MR-
0388. UC-702. Westinghouse Hanford Company, Richland, WA.

5. Killey RWD, Rao RR, Eyvindson S. 1998. Radiocarbon speciation and distribution in an aquifer plume
and groundwater discharge area, Chalk River, Ontario. Appl Geochem 13:3–16.

6. Fellingham L, Graham A, Stiff S. 2003. Characterisation and remediation of beryllium waste pits in the
Southern storage area at Harwell, p. 1041–1049. *In* ASME 2003 9th International Conference on
Radioactive Waste Management and Environmental Remediation. American Society of Mechanical
Engineers, Oxford, England.

7. Wilkins MJ, Livens FR, Vaughan DJ, Lloyd JR, Beadle I, Small JS. 2010. Fe(III) reduction in the
subsurface at a low-level radioactive waste disposal site. Geomicrobiol J 27:231–239.

8. Gudelis A, Gvozdaitė R, Kubarevičienė V, Druteikienė R, Lukoševičius Š, Šutas A. 2010. On radiocarbon
and plutonium leakage to groundwater in the vicinity of a shallow-land radioactive waste repository. J
Environ Radioact 101:443–445.

578 9. Bugai D, Kashparov V, Dewiére L, Khomutinin Y, Levchuk S, Yoschenko V. 2005. Characterization of
579   subsurface geometry and radioactivity distribution in the trench containing Chernobyl clean-up wastes.
580   Environ Geol 47:869–881.

581 10. Payne TE, Harrison JJ, Hughes CE, Johansen MP, Thiruvoth S, Wilsher KL, Cendón DI, Hankin SI,
582   Rowling B, Zawadzki A. 2013. Trench 'bathtubbing' and surface plutonium contamination at a legacy
583   radioactive waste site. Environ Sci Technol 47:13284–13293.

584 11. Cendón DI, Hughes CE, Harrison JJ, Hankin SI, Johansen MP, Payne TE, Wong H, Rowling B, Vine M,
585   Wilsher K, Guinea A, Thiruvoth S. 2015. Identification of sources and processes in a low-level
586   radioactive waste site adjacent to landfills: groundwater hydrogeochemistry and isotopes. Aust J Earth Sci
587   62:123–141.

588 12. Hughes CE, Cendón DI, Harrison JJ, Hankin SI, Johansen MP, Payne TE, Vine M, Collins RN, Hoffmann
589   EL, Loosz T. 2011. Movement of a tritium plume in shallow groundwater at a legacy low-level
590   radioactive waste disposal site in eastern Australia. J Environ Radioact 102:943–952.

591 13. Isaacs SR, Mears KF. 1977. A study of the burial ground used for radioactive waste at the Little Forest
592   area near Lucas Heights New South Wales. AAEC/E--427. Australian Atomic Energy Commission
593   Research Establishment, Lucas Heights, N.S.W. (AU).

594 14. Pett-Ridge J, Firestone MK. 2005. Redox fluctuation structures microbial communities in a wet tropical
595   soil. Appl Environ Microbiol 71:6998–7007.

596 15. Choppin GR. 2007. Actinide speciation in the environment. J Radioanal Nucl Chem 273:695–703.

597 16. Kimber RL, Boothman C, Purdie P, Livens FR, Lloyd JR. 2012. Biogeochemical behaviour of plutonium
598   during anoxic biostimulation of contaminated sediments. Mineral Mag 76:567–578.

599 17. Lloyd JR, Sole VA, Praagh CVGV, Lovley DR. 2000. Direct and Fe(II)-mediated reduction of technetium
600   by Fe(III)-reducing bacteria. Appl Environ Microbiol 66:3743–3749.

601 18. Renshaw JC, Butchins LJC, Livens FR, May I, Charnock JM, Lloyd JR. 2005. Bioreduction of uranium:
602   Environmental implications of a pentavalent intermediate. Environ Sci Technol 39:5657–5660.

603 19. Francis AJ, Dodge CJ, Gillow JB. 2008. Reductive dissolution of Pu(IV) by *Clostridium* sp. under
604   anaerobic conditions. Environ Sci Technol 42:2355–2360.

605 20. Renshaw JC, Law N, Geissler A, Livens FR, Lloyd JR. 2009. Impact of the Fe(III)-reducing bacteria
606   *Geobacter sulfurreducens* and *Shewanella oneidensis* on the speciation of plutonium. Biogeochemistry
607   94:191–196.

608 21. Fomina M, Charnock JM, Hillier S, Alvarez R, Gadd GM. 2007. Fungal transformations of uranium
609   oxides. Environ Microbiol 9:1696–1710.

610 22. Wilkins MJ, Livens FR, Vaughan DJ, Beadle I, Lloyd JR. 2007. The influence of microbial redox cycling
611   on radionuclide mobility in the subsurface at a low‐level radioactive waste storage site. Geobiology
612   5:293–301.

23. Vázquez-Campos X, Kinsela AS, Collins RN, Neilan BA, Aoyagi N, Waite TD. 2015. Uranium binding
mechanisms of the acid-tolerant fungus *Coniochaeta fodinicola*. Environ Sci Technol 49:8487–8496.

24. Amachi S, Fujii T, Shinoyama H, Muramatsu Y. 2005. Microbial influences on the mobility and
transformation of radioactive iodine in the environment. J Nucl Radiochem Sci 6:21–24.

25. Bertin PN, Heinrich-Salmeron A, Pelletier E, Goulhen-Chollet F, Arsène-Ploetze F, Gallien S, Lauga B,
Casiot C, Calteau A, Vallenet D, Bonnefoy V, Bruneel O, Chane-Woon-Ming B, Cleiss-Arnold J, Duran
R, Elbaz-Poulichet F, Fonknechten N, Giloteaux L, Halter D, Koechler S, Marchal M, Mornico D,
Schaeffer C, Smith AAT, Van Dorsselaer A, Weissenbach J, Médigue C, Le Paslier D. 2011. Metabolic
diversity among main microorganisms inside an arsenic-rich ecosystem revealed by meta- and proteo-
genomics. ISME J 5:1735–1747.

26. Castelle CJ, Hug LA, Wrighton KC, Thomas BC, Williams KH, Wu D, Tringe SG, Singer SW, Eisen JA,
Banfield JF. 2013. Extraordinary phylogenetic diversity and metabolic versatility in aquifer sediment. Nat
Commun 4:2120.

27. Aguiar-Pulido V, Huang W, Suarez-Ulloa V, Cickovski T, Mathee K, Narasimhan G. 2016.
Metagenomics, metatranscriptomics, and metabolomics approaches for microbiome analysis. Evol
Bioinform 12:5–16.

28. Allen EE, Banfield JF. 2005. Community genomics in microbial ecology and evolution. Nat Rev Micro
3:489–498.

29. Ikeda-Ohno A, Harrison JJ, Thiruvoth S, Wilsher K, Wong HKY, Johansen MP, Waite TD, Payne TE.
2014. Solution speciation of plutonium and americium at an Australian legacy radioactive waste disposal
site. Environ Sci Technol 48:10045–10053.

30. Harrison JJ, Zawadzki A, Chisari R, Wong HKY. 2011. Separation and measurement of thorium,
plutonium, americium, uranium and strontium in environmental matrices. J Environ Radioact 102:896–
900.

31. APHA, AWWA, WEF. 1998. Standard methods for the examination of water and wastewater20th ed.
APHA-AWWA-WEF, Washington, D.C.

32. Jacobs J, Rhodes M, Sturgis B, Wood B. 2009. Influence of environmental gradients on the abundance
and distribution of *Mycobacterium* spp. in a coastal lagoon estuary. Appl Environ Microbiol 75:7378–
7384.

33. Abubucker S, Segata N, Goll J, Schubert AM, Izard J, Cantarel BL, Rodriguez-Mueller B, Zucker J,
Thiagarajan M, Henrissat B, White O, Kelley ST, Methé B, Schloss PD, Gevers D, Mitreva M,
Huttenhower C. 2012. Metabolic reconstruction for metagenomic data and its application to the human
microbiome. PLoS Comput Biol 8:e1002358.

34. Parks DH, Tyson GW, Hugenholtz P, Beiko RG. 2014. STAMP: statistical analysis of taxonomic and
functional profiles. Bioinformatics 30:3123–3124.

35. Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. J Roy Stat Soc B 57:289–300.

36. Payne TE. 2012. Background report on the Little Forest Burial Ground legacy waste site. ANSTO / E-
780. Institute for Environmental Research, Australian Nuclear Science and Technology Organisation,
Lucas Heights, N.S.W. (AU).

37. Kinsela AS, Jones AJ, Bligh MW, Pham AN, Collins RN, Harrison JJ, Wilsher KL, Payne TE, Waite TD.
2016. Influence of dissolved silicate on Fe(II) oxidation. Environ Sci Technol 50:11663–11671.

38. Jiang L, Zheng Y, Chen J, Xiao X, Wang F. 2011. Stratification of archaeal communities in shallow
sediments of the Pearl River Estuary, Southern China. Antonie van Leeuwenhoek 99:739–751.

39. Meng J, Xu J, Qin D, He Y, Xiao X, Wang F. 2014. Genetic and functional properties of uncultivated
MCG archaea assessed by metagenome and gene expression analyses. ISME J 8:650–659.

40. Evans PN, Parks DH, Chadwick GL, Robbins SJ, Orphan VJ, Golding SD, Tyson GW. 2015. Methane
metabolism in the archaeal phylum Bathyarchaeota revealed by genome-centric metagenomics. Science
350:434–438.

41. He Y, Li M, Perumal V, Feng X, Fang J, Xie J, Sievert SM, Wang F. 2016. Genomic and enzymatic
evidence for acetogenesis among multiple lineages of the archaeal phylum Bathyarchaeota widespread in
marine sediments. Nat Microbiol 16035.

42. Haroon MF, Hu S, Shi Y, Imelfort M, Keller J, Hugenholtz P, Yuan Z, Tyson GW. 2013. Anaerobic
oxidation of methane coupled to nitrate reduction in a novel archaeal lineage. Nature 500:567–570.

43. Karakashev D, Batstone DJ, Trably E, Angelidaki I. 2006. Acetate oxidation is the dominant
methanogenic pathway from acetate in the absence of methanosaetaceae. Appl Environ Microbiol
72:5138–5141.

44. Leu A, Hu S, Cai C, Yuan Z, Orphan VJ, Tyson GW. 2016. Understanding the metabolic potential of
ANME-2d and its role in anaerobic methane oxidation coupled to metal reduction. Montreal, Canada.

45. Luef B, Frischkorn KR, Wrighton KC, Holman H-YN, Birarda G, Thomas BC, Singh A, Williams KH,
Siegerist CE, Tringe SG, Downing KH, Comolli LR, Banfield JF. 2015. Diverse uncultivated ultra-small
bacterial cells in groundwater. Nat Commun 6:6372.

46. Baker BJ, Tyson GW, Webb RI, Flanagan J, Hugenholtz P, Allen EE, Banfield JF. 2006. Lineages of
acidophilic Archaea revealed by community genomic analysis. Science 314:1933–1935.

47. Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A, Wilkins MJ, Wrighton KC, Williams
KH, Banfield JF. 2015. Unusual biology across a group comprising more than 15% of domain Bacteria.
Nature 523:208–211.

48. Peng R-H, Xiong A-S, Xue Y, Fu X-Y, Gao F, Zhao W, Tian Y-S, Yao Q-H. 2008. Microbial
biodegradation of polyaromatic hydrocarbons. FEMS Microbiol Rev 32:927–955.

49.   Riemann L, Azam F. 2002. Widespread *N*-acetyl-D-glucosamine uptake among pelagic marine bacteria
and its ecological implications. Appl Environ Microbiol 68:5554–5562.

50.   Sorokin DY, Rakitin AL, Gumerov VM, Beletsky AV, Damsté S, S J, Mardanov AV, Ravin NV. 2016.
Phenotypic and genomic properties of *Chitinispirillum alkaliphilum* gen. nov., sp. nov., a haloalkaliphilic
anaerobic chitinolytic bacterium representing a novel class in the phylum *Fibrobacteres*. Front Microbiol
7.

51.   Zavarzina DG, Tourova TP, Kolganova TV, Boulygina ES, Zhilina TN. 2009. Description of
Anaerobacillus alkalilacustre gen. nov., sp. nov.—Strictly anaerobic diazotrophic bacillus isolated from
soda lake and transfer of *Bacillus arseniciselenatis*, *Bacillus macyae*, and *Bacillus alkalidiazotrophicus* to
Anaerobacillus as the new combinations *A. arseniciselenatis* comb. nov., *A. macyae* comb. nov., and *A.*
alkalidiazotrophicus comb. nov. Microbiology 78:723–731.

52.   Beier S, Bertilsson S. 2013. Bacterial chitin degradation—mechanisms and ecophysiological strategies.
Front Microbiol 4:149.

53.   de Gonzalo G, Colpa DI, Habib MHM, Fraaije MW. 2016. Bacterial enzymes involved in lignin
degradation. J Biotechnol 236:110–119.

54.   Pollegioni L, Tonin F, Rosini E. 2015. Lignin-degrading enzymes. FEBS J 282:1190–1213.

55.   Burow LC, Mabbett AN, Blackall LL. 2008. Anaerobic glyoxylate cycle activity during simultaneous
utilization of glycogen and acetate in uncultured *Accumulibacter* enriched in enhanced biological
phosphorus removal communities. ISME J 2:1040–1051.

56.   Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, Rattei T, Mende DR,
Sunagawa S, Kuhn M, Jensen LJ, von Mering C, Bork P. 2016. eggNOG 4.5: a hierarchical orthology
framework with improved functional annotations for eukaryotic, prokaryotic and viral sequences. Nucl
Acids Res 44:D286–D293.

57.   Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2016. KEGG as a reference resource for
gene and protein annotation. Nucl Acids Res 44:D457–D462.

58.   Magrane M, Uniprot Consortium. 2011. UniProt Knowledgebase: a hub of integrated protein data.
Database 2011:bar009.

59.   Cui M, Ma A, Qi H, Zhuang X, Zhuang G. 2015. Anaerobic oxidation of methane: an "active" microbial
process. MicrobiologyOpen 4:1–11.

60.   Timmers PHA, Welte CU, Koehorst JJ, Plugge CM, Jetten MSM, Stams AJM. 2017. Reverse
methanogenesis and respiration in methanotrophic Archaea. Archaea 2017:e1654237.

61.   Ettwig KF, Zhu B, Speth D, Keltjens JT, Jetten MSM, Kartal B. 2016. Archaea catalyze iron-dependent
anaerobic oxidation of methane. PNAS 113:12792–12796.

62.   Grein F, Ramos AR, Venceslau SS, Pereira IAC. 2013. Unifying concepts in anaerobic respiration:
Insights from dissimilatory sulfur metabolism. Biochim Biophys Acta Bioenerg 1827:145–160.

63. Paulo LM, Stams AJM, Sousa DZ. 2015. Methanogens, sulphate and heavy metals: a complex system.
Rev Environ Sci Biotechnol 14:537–553.

64. Scheller S, Yu H, Chadwick GL, McGlynn SE, Orphan VJ. 2016. Artificial electron acceptors decouple
archaeal methane oxidation from sulfate reduction. Science 351:703–707.

65. He Z, Cai C, Wang J, Xu X, Zheng P, Jetten MSM, Hu B. 2016. A novel denitrifying methanotroph of the
NC10 phylum and its microcolony. Sci Rep 6:32241.

66. Wrighton KC, Castelle CJ, Wilkins MJ, Hug LA, Sharon I, Thomas BC, Handley KM, Mullin SW,
Nicora CD, Singh A, Lipton MS, Long PE, Williams KH, Banfield JF. 2014. Metabolic interdependencies
between phylogenetically novel fermenters and respiratory organisms in an unconfined aquifer. ISME J
8:1452–1463.

67. Könneke M, Schubert DM, Brown PC, Hügler M, Standfest S, Schwander T, Borzyskowski LS von, Erb
TJ, Stahl DA, Berg IA. 2014. Ammonia-oxidizing archaea use the most energy-efficient aerobic pathway
for CO_2 fixation. PNAS 111:8239–8244.

68. Pester M, Schleper C, Wagner M. 2011. The Thaumarchaeota: an emerging view of their phylogeny and
ecophysiology. Curr Opin Microbiol 14:300–306.

69. Fischer M, Falke D, Pawlik T, Sawers RG. 2014. Oxygen-dependent control of respiratory nitrate
reduction in mycelium of *Streptomyces coelicolor* A3(2). J Bacteriol 196:4152–4162.

70. Lovley DR, Holmes DE, Nevin KP. 2004. Dissimilatory Fe(III) and Mn(IV) reduction. Adv Microb
Physiol 49:219–286.

71. Weber KA, Achenbach LA, Coates JD. 2006. Microorganisms pumping iron: anaerobic microbial iron
oxidation and reduction. Nat Rev Micro 4:752–764.

72. Kato S, Ohkuma M, Powell DH, Krepski ST, Oshima K, Hattori M, Shapiro N, Woyke T, Chan CS. 2015.
Comparative genomic insights into ecophysiology of neutrophilic, microaerophilic iron oxidizing
bacteria. Front Microbiol 6.

73. Emerson D, Fleming EJ, McBeth JM. 2010. Iron-oxidizing bacteria: an environmental and genomic
perspective. Ann Rev Microbiol 64:561–583.

74. Anandkumar B, George RP, Maruthamuthu S, Parvathavarthini N, Mudali UK. 2016. Corrosion
characteristics of sulfate-reducing bacteria (SRB) and the role of molecular biology in SRB studies: an
overview. Corrosion Rev 34:41–63.

75. Lütters-Czekalla S. 1990. Lithoautotrophic growth of the iron bacterium *Gallionella ferruginea* with
thiosulfate or sulfide as energy source. Arch Microbiol 154:417–421.

76. DeAngelis KM, Silver WL, Thompson AW, Firestone MK. 2010. Microbial communities acclimate to
recurring changes in soil redox potential status. Environ Microbiol 12:3137–3149.

77. Cozzarelli IM, Weiss JV. 2007. Biogeochemistry of Aquifer Systems, p. 843–859. *In* Hurst, CJ,
Crawford, RL, Garland, JL, Lipson, DA, Mills, AL, Stetzenbach, LD (eds.), Manual of environmental
microbiology. American Society of Microbiology.

78. Hansel CM, Lentini CJ, Tang Y, Johnston DT, Wankel SD, Jardine PM. 2015. Dominance of sulfur-
fueled iron oxide reduction in low-sulfate freshwater sediments. ISME J 9:2400–2412.

79. Boland DD, Collins RN, Miller CJ, Glover CJ, Waite TD. 2014. Effect of solution and solid-phase
conditions on the Fe(II)-accelerated transformation of ferrihydrite to lepidocrocite and goethite. Environ
Sci Technol 48:5477–5485.

80. Pedersen HD, Postma D, Jakobsen R, Larsen O. 2005. Fast transformation of iron oxyhydroxides by the
catalytic action of aqueous Fe(II). Geochim Cosmochim Acta 69:3967–3977.

81. Jones AM, Collins RN, Rose J, Waite TD. 2009. The effect of silica and natural organic matter on the
Fe(II)-catalysed transformation and reactivity of Fe(III) minerals. Geochim Cosmochim Acta 73:4409–
4422.

82. Lovley D. 2013. Dissimilatory Fe(III)- and Mn(IV)-Reducing Prokaryotes, p. 287–308. *In* Rosenberg, E,
DeLong, EF, Lory, S, Stackebrandt, E, Thompson, F (eds.), The Prokaryotes. Springer Berlin Heidelberg,
Berlin, Heidelberg.

83. Lovley DR, Phillips EJP. 1988. Novel mode of microbial energy metabolism: Organic carbon oxidation
coupled to dissimilatory reduction of iron or manganese. Appl Environ Microbiol 54:1472–1480.

84. Straub KL, Benz M, Schink B. 2001. Iron metabolism in anoxic environments at near neutral pH. FEMS
Microbiol Ecol 34:181–186.

85. Benz M, Schink B, Brune A. 1998. Humic acid reduction by *Propionibacterium freudenreichii* and other
fermenting bacteria. Appl Environ Microbiol 64:4507–4512.

86. Kersting AB. 2013. Plutonium transport in the environment. Inorg Chem 52:3533–3546.

87. Kersting AB, Efurd DW, Finnegan DL, Rokop DJ, Smith DK, Thompson JL. 1999. Migration of
plutonium in ground water at the Nevada Test Site. Nature 397:56–59.

88. Santschi PH, Roberts KA, Guo L. 2002. Organic nature of colloidal actinides transported in surface water
environments. Environ Sci Technol 36:3711–3719.

89. Novikov AP, Kalmykov SN, Utsunomiya S, Ewing RC, Horreard F, Merkulov A, Clark SB, Tkachev VV,
Myasoedov BF. 2006. Colloid transport of plutonium in the far-field of the Mayak Production
Association, Russia. Science 314:638–641.

90. Schäfer T, Artinger R, Dardenne K, Bauer A, Schuessler W, Kim JI. 2003. Colloid-borne americium
migration in Gorleben groundwater: Significance of iron secondary phase transformation. Environ Sci
Technol 37:1528–1534.

91. Lujaniené G, Beneš P, Štamberg K, Ščiglo T. 2012. Kinetics of plutonium and americium sorption to
natural clay. J Environ Radioact 108:41–49.

Figure captions

Figure 1. pH and E_h (Standard Hydrogen Electrode corrected) measurements from the trench water across the sampling period.

Figure 2. Temporal changes to element/ion concentrations in the trench water. Bars show the individual concentrations of each element/ion over the five sampling days 0, 4, 6, 21 and 47.

Figure 3. Activity of filtered and unfiltered radionuclides measured in the trench water. Filtered fractions (<0.45 µm, solid fill) are equivalent to soluble and smaller colloidal particles. Unfiltered fractions (full bar) are as considered total (soluble plus all suspended solids) concentrations. Error bars show the standard deviation of triplicate measurements.

Figure 4. Taxa relative abundances over time. Only taxa with an average relative abundance >5% of the total sequences in at least a single sampling day are shown. Taxa with <5% of all sequences are grouped under 'other'. Phyla with <5% of relative abundance at all times are grouped under 'other phyla', except Archaea.

Figure 5. Changes in the relative abundances of selected RXNs over time. A, Cytochrome C oxidase; B, cellulase; C, malate synthase; D, 5-methyltetrahydrosarcinapterin:corrinoid/iron-sulfur protein Co-methyltransferase (CH3-HSPT:Fe-S protein Co-MT); E: sulfite reductase; and F, superoxide dismutase.

Figure 6. Hypothetical global scheme of the processes at LFLS. Orange details on barrels reading Fe(III) represent solid Fe(III) minerals, mainly (oxy)hydroxides. Colour of arrows represent the time at which each process takes place: red, aerobic/after rain; brown, microaerophilic; blue, anaerobic/dry phase; and black arrows indicate processes that seem to be independent from the sampling time. Dashed arrows indicate transport. Abbreviations used in the figure: C_{org}, organic carbon; MIC, microbially induced corrosion; CH_3-HSPT:Fe-S protein *Co*-MT, 5-methyltetrahydrosarcinapterin:corrinoid/iron-sulfur protein Co-methyltransferase.

Figure 3

Figure 5