Production and purification of human Hsp90β in *Escherichia coli*

Martina Radli[1], Dmitry B. Veprintsev[2,3], Stefan G. D. Rüdiger[1*]

[1]Cellular Protein Chemistry, Bijvoet Center for Biomolecular Research, Science for Life, Utrecht University, Paadualaan 8, 3584 CH Utrecht, The Netherlands

[2]Laboratory of Biomolecular Research, Paul Scherrer Institut, 5232 Villigen PSI, Switzerland

[3]Department of Biology, ETH Zürich, 8093 Zürich, Switzerland

*Corresponding author:
E-mail: s.g.d.rudiger@uu.nl (SGDR)

Running title: Production and purification of Hsp90

Keywords: Molecular chaperones, Protein folding, Protein chemistry, Hsp90, Protein production

Abstract

The molecular chaperone Hsp90 is an essential member of the cellular proteostasis system. It plays an important role in the stabilisation and activation of a large number of client proteins and is involved in fatal disease processes e.g. Alzheimer disease, cancer and cystic fibrosis. This makes Hsp90 a crucial protein to study. Mechanistic studies require large amounts of protein but the production and purification of recombinant human Hsp90 in *E. coli* is challenging and laborious. Here we identified conditions that influence Hsp90 production and optimised a fast and efficient purification protocol. We found that the nutrient value of the culturing medium and the length of induction had significant effect on Hsp90 production in *Escherichia coli*. Our fast, single-day purification protocol resulted in a stable, well-folded and pure sample that was resistant to degradation in a reproducible manner. We anticipate that our results provide a useful tool to produce higher amount of pure, well-folded and stable recombinant human Hsp90β in *Escherichia coli* in an efficient way.

Introduction

The cellular proteostasis system evolved to maintain cellular health and stability and to protect cells from continuously occurring stress by a tight control of protein production, quality control, folding, trafficking, aggregation and degradation (1,2). Chaperones are crucial elements of the proteostasis system, they prevent protein misfolding and aggregation by various mechanisms and thereby contribute to cellular integrity (3).

The molecular chaperone Hsp90 is one of the most important element of the proteostasis system (4-7). It is involved in client protein folding, maturation, stabilisation, activation and assembly of large protein complexes (8,9). Typically, Hsp90 assists at late folding processes (10,11).

Hsp90 interacts with up to 10 % of the cellular proteome. Its main clients include transcription factors, kinases and hormone receptors (Didier Picard. Table of Hsp90 interactors. Available from: https://www.picard.ch/downloads/Hsp90interactors.pdf). Moreover, it has unconventional partners such as the disordered α-synuclein and Tau (5,10). Hsp90 plays a crucial role in the progression of several diseases (such as cancer, cystic fibrosis and Alzheimer's disease) (12-14).

The Hsp90 chaperone machinery is extensively studied to understand its complex working mechanism and involvement with multiple fatal diseases. Typically, *in vitro* biochemical and biophysical experiments require high amount of recombinant protein with excellent sample purity and stability. Significant overproduction of the recombinant human Hsp90β is challenging since it is a

large, multi-domain protein. Moreover, Hsp90 is particularly sensitive to the

proteolytic cleavage of the flexible, unfolded linkers between its domains,

therefore a time consuming purification protocol may compromise sample quality.

We systematically tested several parameters to find the best conditions for

the overproduction of recombinant human Hsp90β. Furthermore, we developed a

fast and efficient purification protocol that results in pure and stable sample with

reproducible quality. We found that the nutrient value of the medium and the

length of induction time had significant effect on Hsp90 overproduction, whereas

the concentration of induction agent (isopropyl β-D-1-thiogalactopyranoside

(IPTG)), temperature and optical density (OD_{600}) at induction (within the tested

intervals) did not influence the process. The protein samples we purified with the

optimised protocol were resistant to proteolysis upon incubation at physiological

temperature up to one day. We showed that the folding, sedimentation and

molecular weight of our Hsp90 sample corresponded to earlier results.

Materials and Methods

Wild type full length Hsp90 construct

80 pet23a+ expression vector (Novagen) was used for His$_6$-tagged wild type human

81 Hsp90 production in Rosetta 2 *E. coli* strain (Novagen). The sequence of wild

82 type Hsp90:

```
HHHHHHMPEEVHHGEEEVETFAFQAEIAQLMSLIINTFYSNKEIFLRELISNASDALDKIRYESLTDPSKLD

SGKELKIDIIPNPQERTLTLVDTGIGMTKADLINNLGTIAKSGTKAFMEALQAGADISMIGQFGVGFYSAYL

VAEKVVVITKHNDDEQYAWESSAGGSFTVRADHGEPIGRGTKVILHLKEDQTEYLEERRVKEVVKKHSQFIG

YPITLYLEKEREKEISDDEAEEEKGEKEEEDKDDEEKPKIEDVGSDEEDDSGKDKKKKTKKIKEKYIDQEEL

NKTKPIWTRNPDDITQEEYGEFYKSLTNDWEDHLAVKHFSVEGQLEFRALLFIPRRAPFDLFENKKKKNNIK

LYVRRVFIMDSCDELIPEYLNFIRGVVDSEDLPLNISREMLQQSKILKVIRKNIVKKCLELFSELAEDKENY

KKFYEAFSKNLKLGIHEDSTNRRRLSELLRYHTSQSGDEMTSLSEYVSRMKETQKSIYYITGESKEQVANSA

FVERVRKRGFEVVYMTEPIDEYCVQQLKEFDGKSLVSVTKEGLELPEDEEEKKKMEESKAKFENLCKLMKEI

LDKKVEKVTISNRLVSSPCCIVTSTYGWTANMERIMKAQALRDNSTMGYMMAKKHLEINPDHPIVETLRQKA

EADKNDKAVKDLVVLLFETALLSSGFSLEDPQTHSNRIYRMIKLGLGIDEDEVAAEEPNAAVPDEIPPLEGD

EDASRMEEVD
```

Protein overproduction test - cell culturing

Rosetta 2 cells containing pet23a+ vector with wild type full length Hsp90 were

inoculated into 100 ml 2x yeast tryptone extract (2x YT; 12.8 g Bacto Tryptone

(Merck), 8 g Bacto Yeast Extract (Merck), 4 g NaCl (Merck) in 800 ml demi water

(Millipore)) medium supplemented with 34 mg/l final concentration of

chloramphenicol (Sigma-Aldrich) and 100 mg/l final concentration of ampicillin

(Sigma-Aldrich). The cells were grown over night at 37^{o}C, shaking with 220 rpm.

Next morning 100 (for low OD_{600}) or 200 (for high OD_{600}) μl culture was

inoculated into 4 ml lysogeny broth medium (LB; 8 g Bacto Tryptone, 4 g Bacto

Yeast Extract, 4 g NaCl, 20 g tymine (Sigma-Aldrich) in 800 ml demi water), 2x

YT or terrific broth media (TB; 12 g Bacto Tryptone, 24 g Bacto Yeast Extract, 4

ml glycerol in 800 ml demi water)) supplemented with 34 mg/l final concentration

of chloramphenicol and 100 mg/l final concentration of ampicillin. The cultures

were grown at $37^{o}C$, shaking with 180 rpm. The OD_{600} was monitored every 1.5

hours by using a 1:10 dilution of the culture in an Ultrospec 3000 pro UV/Visible

Spectrophotometer (GE Healthcare). The cultures were induced at low (0.6-0.9)

or high (1-1.3) OD_{600} with 0.1/0.25/0.5 mM IPTG (Thermo Fisher Scientific). The

cultures were incubated at $16^{o}C$ or $18^{o}C$, the SDS-PAGE samples were taken

before induction and 1/3/5 day(s) after induction.

SDS-PAGE

For the SDS-PAGE samples 0.5 ml of cell culture was taken and centrifuged for 1

minute by 13,300 rpm. The supernatant was discarded and the pellet was

resuspended in 200 μl of 1x sample buffer (0.625 M Tris (Sigma-Aldrich), 12.5 %

glycerol (CARL ROTH), 1 % SDS (Bio-Rad), 0.005 % Bromophenol Blue (Bio-

Rad), 5 mM freshly added β-mercaptoethanol (Sigma-Aldrich) 2x diluted with

demi water (Millipore)) and homogenised by using a syringe of a very tiny

diameter (BD Micro-Fine). This last step was essential for loading the samples on

the gel.

15 % SDS gels were prepared (Separation buffer: 0.38 M Tris pH 8.8, 15

% acrylamide (National Diagnostics), 0.1 % SDS, 0.1 % APS (Sigma-Aldrich),

0.04 % TEMED (Sigma-Aldrich), Stacking buffer: 0.125 M Tris pH 6.8, 4 %

acrylamide, 0.1 % SDS, 0.075 % APS, 0.1 % TEMED) and ran in 1x Laemmli

buffer (0.025 M Tris base, 0.152 M glycine (SERVA Electrophoresis GmbH) , 0.1

% SDS, diluted from 10x stock). The gels stained with Coomassie staining

solution (0.2 % Coomassie Brilliant Blue (SERVA Electrophoresis GmbH), 45 %

methanol (Interchema Antonides-Interchema), 10 % acetic acid (Biosolve) and 55

% demi water and destained using destaining solution (30 % methanol, 10 %

acetic acid and 60 % demi water).

The gels were scanned in a Epson Perfection V700 Photo scanner.

Quantification of the lane profiles was done by using ImageJ. The lane profiles of

the induced samples were normalised to the lane profile of the uninduced sample

of the same gel using the intrinsic *E. coli* protein called EF-Tu. Subsequently, the

lane profiles of induced samples were aligned to the uninduced sample using the

first two bands in the beginning of the lanes. Finally, the uninduced lane profile

was deducted from the induced lane profiles and the sum of each Hsp90 peak

area was divided by the sum of the normalised EF-Tu peak area of the same lane

to be able to compare the gels with each other.

Overproduction and purification of wild type full length Hsp90

Rosetta 2 cells containing pet23a+ vector with wild type full length Hsp90 were

inoculated into 200 ml 2x YT medium supplemented with 34 mg/l final

concentration of chloramphenicol and 100 mg/l final concentration of ampicillin.

The cells were grown over night at 37^{o}C, with shaking at 220 rpm. Next morning

200 ml culture was inoculated into 6x800 ml 2x YT supplemented with 34 mg/l

final concentration of chloramphenicol and 100 mg/l final concentration of

ampicillin. The cultures were grown at 37^{o}C with shaking at 180 rpm and induced

with 0.5 mM IPTG at OD_{600} = 1. After induction the cells were incubated at 18^{o}C

for 5 days with shaking at 180 rpm.

The cells were harvested in an Avanti J-26 XP centrifuge (Beckman

Coulter) using the JLA-8.1 rotor at 4^{o}C at 4500 rpm for 30 minutes. The

supernatant was discarded and the pellet was resuspended in ice cold

resuspension-buffer (50 mM Na-phosphate pH 7.2 (Sigma-Aldrich), 150 mM

NaCl, 150 mM KCl (CARL ROTH)) and centrifuged in an MSE Harrier 18/80

Refrigerated Benchtop Centrifuge at 4^{o}C at 5000 rpm for 30 minutes. The

supernatant was discarded and the pellet was stored at -20^{o}C until further usage.

The pellet was resuspended in ice cold lysis buffer (12.5 mM Na-

phosphate pH 7.2. 75 mM NaCl, 5 mM β-mercaptoethanol, EDTA-free protease

inhibitor (1 tablet/50 ml) (Roche)). The cells were disrupted by an EmulsiFlex-C5

(Avestin) cell disruptor. The lysate was centrifuged in Avanti J-26 XP centrifuge

using JA-25.5 rotor at 21,000 rpm for 45 minutes at 4^{o}C. The lysate was filtered

by 22 μm polypropylene filter (VWR) to remove the cell debris and insoluble

aggregates. The purification was done using an AKTA Purifier (GE Healthcare).

Wild type full length Hsp90 was first purified on an IMAC POROS 20MC

(Thermo Fischer Scientific) affinity purification column (solutions connected to

pump A1 and A2: 50 mM Na-phosphate buffer pH 8.0 with 300 mM NaCl, B1:

demi water with 10 mM β-mercaptoethanol, B2: 1 M imidazole (Sigma-Aldrich)).

The eluted sample was diluted 4-fold with dilution buffer (25 mM Na-phosphate

buffer pH 7.2, 5 mM DTT (Sigma-Aldrich), complete protease inhibitor (1 tablet/50

ml) (Roche)). Next the sample was loaded on a POROS 20HQ anion exchange

column (Thermo Fischer Scientific) (solutions connected to pump A1 and A2: 50

mM Na-phosphate pH 7.2, B1: demi water with 10 mM DTT, B2: 2 M KCl). The

eluted sample was diluted 10-fold with dilution buffer (25 mM Na-phosphate

buffer pH 7.2, 5 mM DTT, complete protease inhibitor (1 tablet/100 ml)). Finally,

the sample was loaded on a HiTrap heparin affinity chromatography column (GE

Healthcare) (solutions connected to pump A1 and A2: 25 mM Na-phosphate pH

7.2, B1: demi water with 10 mM DTT, B2: 2 M KCl). The eluate was concentrated

and buffer exchanged to Hsp90 storage buffer (25 mM Na-phosphate pH 7.2, 150

mM NaCl, 150 mM KCl), 5 mM DTT, complete protease inhibitor (1 tablet/100

ml)) using a Vivaspin 20 column (50 kDa MWCO) (GE Healthcare) at 4^{o}C at 5000

rpm for 15-15 minutes until above 100 µM protein concentration. The protein

concentration was determined with ND-1000 program on an ND-1000

Spectrophotometer type NanoDrop using 57760 M^{-1}cm^{-1} extinction coefficient.

The sample was aliquoted, frozen in liquid N$_2$ and stored at -80^{o}C. Throughout

the purification procedure samples were taken from the steps of the purification

that were run on SDS-PAGE to analyse sample purity.

Protein stability

10 µM concentration Hsp90 was incubated in Hsp90-buffer ((25 mM Na-phosphate pH 7.2, 150 mM NaCl, 150 mM KCl, 5 mM DTT, complete protease inhibitor (1 tablet/100 ml)) for 24 hours at 4^oC, room temperature ($\sim21^oC$) and 37^oC. SDS-PAGE samples were taken and mixed with 2x sample buffer (1.25 M Tris, 25 % glycerol, 2 % SDS, 0.01 % Bromophenol Blue, 5 mM freshly added DTT) at 0/3/18/24 hours. The samples were analysed on SDS-PAGE.

Silver staining

The Coomassie-stained and destained gels were fixed for 30 minutes in fixation solution (30 % ethanol (Interchema Antonides-Interchema), 10 % acetic acid, 60 % demi water). The gel was washed 3-times for 20 minutes in 50 % ethanol, then pre-treated for 1 minute with 0.02 % $Na_2S_2O_3$ (Scharlau Chemicals) [100x diluted from stock (stock: 2 g into $Na_2S_2O_3$ 100 ml water (Milli-Q))] and quickly rinsed 4 times with demi water. The gels were impregnated for 20 minutes with freshly prepared staining solution (2 g/l $AgNO_3$ (Merck) and 0.75 ml/l formaldehyde (Calbiochem)). Next they were quickly rinsed 4 times with demi water, then developed till the desired result in developer solution (60 g/l Na_2CO_3 (Sigma-Aldrich), 0.5 ml/liter formaldehyde and 0.0004 % $Na_2S_2O_3$ stock). The reaction was stopped in fixation solution (30 % ethanol, 10 % acetic acid, 60 % demi water) for 10 minutes. The gel was stored in 1 % acetic acid solution and scanned by Epson Perfection V700 Photo scanner.

CD spectroscopy

The protein was centrifuged on $4^\circ C$ for 15 minutes at 13,300 rpm in Heraeus Pico 17 centrifuge (Thermo Scientific). The concentration was determined with ND-1000 program on ND-1000 Spectrophotometer type NanoDrop using 57760 $M^{-1}cm^{-1}$ extinction coefficient. The protein was diluted to 0.1 g/l concentration with CD-buffer (25 mM Na-phosphate pH 7.2 buffer, 150 mM NaF (Sigma-Aldrich)), loaded into a Teflon-sealed, polarimetrically checked quartz glass cuvette with an optical path length of 1 mm and a volume of 350 µl (Hellma Analytics). Far-UV CD spectrum was measured with the Spectra Manager (Jasco) program on a Jasco J-810 Spectropolarimeter instrument (Jasco). Experimental parameters included a wavelength increment of 1 nm, a scan speed of 20 nm/min, a temperature of $20^\circ C$). The data were analysed by MS Excel.

SEC-MALLS

The protein sample was centrifuged and its concentration was measured as described in the *CD spectroscopy* part. 10 µl of ~29 g/l sample was run with Shimadzu on a Superdex 200 column with 0.35 ml/min speed in running buffer (25 mM Na-phosphate pH 7.2, 150 mM NaCl, 5 mM freshly added DTT). The 220 nm absorption was detected by SPD-20A UV detector (Shimadzu), light scattering by Wyatt COMET^{TM} light scattering detector (Wyatt Technology) and the refractive index by the RID-10A refractive index detector (Shimadzu). The data were analysed by program Astra 6.

Analytical Ultracentrifugation (AUC)

The protein sample was centrifuged and its concentration was measured as

described in the *CD spectroscopy* part. The protein sample was diluted to 7.2 µM

concentration with Hsp90-buffer (25 Na-phosphate pH 7.2, 150 mM NaCl, 150

mM KCl, complete protease inhibitor (1 tablet/50 ml), 5 mM freshly added DTT).

The sample was centrifuged on 20^{o}C for 16 hours at 42,000 rpm in a Beckman

XL-I ultracentrifuge using An60Ti rotor. We used absorbance detection optics for

our experiment. The data was analysed by SedFit (Schuck, 2000).

Results

Systematic testing of recombinant human Hsp90β overproduction

We set out to systematically test the effect of key parameters on recombinant human Hsp90β production in Rosetta 2 cells and to identify conditions resulting in high protein levels.

We outlined our study as follows: We cultured *E. coli* cells containing human Hsp90β open reading frame on a pET23a+ vector and tested media with different nutrient values and different temperature of induction to modulate the metabolism of *E. coli* cells. We induced the cultures with different final concentration of IPTG affecting the level of T7 polymerase. We also varied the length of induction and the OD_{600} value at induction, because in case of certain proteins these are essential parameters for successful overproduction.

At given time points we took samples from the cultures and ran them on SDS-gels **(Figure 1A)**. We loaded protein marker in the first, uninduced sample in the second and purified Hsp90 in the last lanes of the SDS-gels. These samples helped to identify the Hsp90 bands that appeared after induction. Lane 3-14 contained the induced samples of the different conditions tested. The most abundant intrinsic *E. coli* band that we later used for quantification (EF-Tu) and Hsp90 bands are marked on the right of the gels (**Figure 1A**) (15).

Figure 1: Experimental set up and quantification scheme for Hsp90 overproduction. *(A) Schematic overview of the protein production experiment.*

E. coli cells containing human Hsp90β open reading frame were inoculated from

glycerol stock and cultured overnight at 37°C. The culture was divided into small

tubes and after induction the tubes were incubated in different conditions. Gel

samples were taken at day 1, 3 and day 5 and were run on SDS-PAGE. (B)

Schematic representation of the quantification of Hsp90 overproduction. The

intensity profiles of uninduced (orange) and induced lanes (green) were

determined and the induced lane profile were normalised to the uninduced lane

profile using the EF-Tu peak of the uninduced lane as reference (100 %) (top

panel). Subsequently, the induced lane profiles were aligned to the uninduced

lane profile using the first two left peaks (stars) of the uninduced lane profile as

reference points (middle panel). The uninduced lane profile was deducted from

the normalised and aligned induced lane profiles and the sum of difference at the

height of the Hsp90 peak was calculated (red area) (lower panel). Finally, the

Hsp90 peak area was divided by the normalised and aligned EF-Tu peak area.

We determined the intensity profile of the uninduced (green) and induced lanes

(orange) (**Figure 1B,** top panel) and normalised the induced lane profile to the

uninduced lane profile using the EF-Tu peak of the uninduced lane as reference

(100 %). Subsequently, we aligned the induced lane profile (green) to the

uninduced lane profile (orange) using the first two left peaks (stars) of the

uninduced lane profile as reference points **(Figure 1B,** middle panel**)**. We

deducted the uninduced lane profile (orange) from the normalised and aligned

induced lane profile (green) and calculated the sum of difference at the height of

the Hsp90 peak (red area) (**Figure 1B,** lower panel). Finally, we divided the

Hsp90 peak area by the area of the normalised and aligned EF-Tu peak.

The yield of recombinant human Hsp90β protein in E. coli using LB medium

was insufficient for preparative purposes

We tested Hsp90 overproduction in LB medium varying the length of induction,

the temperature, the OD_{600} at induction and the IPTG concentration to measure if

any of these parameters modulated the levels of the chaperone.

After one and three days of induction no Hsp90 band appeared in the

induced samples, none of the conditions led to sufficient Hsp90 overproduction

(**Figure 2A and B**, lane 3-14). When induced for five days an Hsp90 band was

apparent in several conditions (**Figure 2C**, lane 4, 5, 11 and 12). However, the

yield was still insufficient for preparative purposes and the production levels were

variable.

Figure 2*: Hsp90 overproduction in LB medium lead to insufficient protein*

yield*. Hsp90 overproduction after 1 day (A), 3 days (B), 5 days (C) of induction.*

Lane 1: protein marker, lane 2: uninduced sample, lane 3-8: protein production at

$16^{o}C$, lane 9-14: protein production at $18^{o}C$, lane 3-5 and 9-11: induction at low

OD_{600}, lane 6-8 and 12-14: induction at high OD_{600}, lane 3, 6, 9, 12: induction with

0.1 mM IPTG, lane 4, 7, 10, 13: induction with 0.25 mM IPTG, lane 5, 8, 11, 14:

induction with 0.5 mM IPTG, lane 15: purified Hsp90 sample. EF-Tu intrinsic E.

coli protein band was used for quantification of Hsp90 overproduction.

2x YT medium improved the yield of Hsp90 production

Since protein production in LB resulted in poor yields we hypothesised that the nutrient value of the medium may be a critical parameter in case of Hsp90. Therefore, we repeated the overproduction experiment in the richer 2x YT medium. We varied the length of induction, the temperature, the OD_{600} at induction and the IPTG concentration to determine if any these parameters affect Hsp90 production in 2x YT.

After one and three days of induction a new band appeared at the height of Hsp90 in the induced samples, suggesting that we could produce the chaperone (**Figure 3A and B**). The newly appearing Hsp90 band is stronger after three days of induction compared to one. Hsp90 production in 2x YT resulted in higher yields than in LB within the interval of the parameters we varied.

Figure 3: Hsp90 overproduction was improved but still insufficient in 2x YT compared to LB. Hsp90 overproduction after 1 day (A), 3 days (B), 5 days (C) of induction. Lane 1: protein marker, lane 2: uninduced sample, lane 3-8: protein production at $16^{o}C$, lane 9-14: protein production at $18^{o}C$, lane 3-5 and 9-11: induction at low OD_{600}, lane 6-8 and 12-14: induction at high OD_{600}, lane 3, 6, 9, 12: induction with 0.1 mM IPTG, lane 4, 7, 10, 13: induction with 0.25 mM IPTG, lane 5, 8, 11, 14: induction with 0.5 mM IPTG, lane 15: purified Hsp90 sample. EF-Tu intrinsic E. coli protein band was used for quantification of Hsp90 overproduction.

When induced for five days, we observed significant overproduction of the

chaperone **(Figure 3C)** in certain cases (lane 7-9), whereas other conditions led

to similar results to those observed on previous gels (lane 10, 12). Hsp90

production levels were variable in the different conditions. We concluded that

Hsp90 overproduction resulted in higher yields in 2x YT after five days of

induction compared to shorter induction times and culturing in LB medium, but

the yields were still insufficient for preparative purposes.

Recombinant human Hsp90β production had the best yield in the E. coli

cells in TB medium

Since the richer 2x YT medium had a positive effect on Hsp90 levels we decided

to test the protein production in TB medium which is higher in nutrients than 2x

YT. We varied the same parameters as described previously.

After one day of induction, a strong band appeared at the height of Hsp90

in several conditions (**Figure 4A**, lane 3 and 11) indicating that overproduction in

TB was notable. The conditions that resulted in high protein amount, however,

were variable. Already after three days of induction in TB medium resulted all the

conditions tested in a strong Hsp90 band (**Figure 4B**). We observed similar

Hsp90 levels with five days of induction (**Figure 4C**). Thus, TB medium resulted

in higher yields compared to both LB and 2x YT media, and after a shorter period

of time.

Figure 4: Hsp90 overproduction lead to the highest yields in TB medium.

Hsp90 overproduction after 1 day (A), 3 days (B), 5 days (C) of induction. Lane 1:

protein marker, lane 2: uninduced sample, lane 3-8: protein production at 16^oC,

lane 9-14: protein production at 18^oC, lane 3-5 and 9-11: induction at low OD_{600},

lane 6-8 and 12-14: induction at high OD_{600}, lane 3, 6, 9, 12: induction with 0.1

mM IPTG, lane 4, 7, 10, 13: induction with 0.25 mM IPTG, lane 5, 8, 11, 14:

induction with 0.5 mM IPTG, lane 15: purified Hsp90 sample. EF-Tu intrinsic E.

coli protein band was used for quantification of Hsp90 overproduction.

Quantification of overproduction gels revealed that the richness of the

medium and the length of induction are the most significant parameters for

Hsp90

To compare the protein production results of the different gels we estimated

Hsp90 overproduction by gel densitometry as described in Figure 1B. In LB

medium we observed low level of Hsp90 overproduction in each condition

(Figure 5A). The tendency improved with the length of induction but even after

five days the yields were insufficient for preparative purposes in all conditions.

Figure 5: Quantification of Hsp90 overproduction in different media. (A)

Summary of the quantification of Hsp90 overproduction in LB, after 1 day (white),

3 days (grey) and 5 days (black) of induction. Average of results of all conditions

(1^{st} column group), average of results on 16^oC and 18^oC (2^{nd} and 3^{rd} column

groups), average of results at low or high OD_{600} at induction (4^{th} and 5^{th} column

groups), average of results at 0.1, 0.25 and 0.5 mM IPTG concentration (6th to 8th

column groups). The length of induction is the only variable that influences Hsp90

production. (B) Summary of the quantification of Hsp90 overproduction in 2x YT.

The length of induction is the only variable that influences Hsp90 production. The

yield is higher in 2x YT compared to LB. The structure of the chart is as in (A). (C)

Summary of the quantification of Hsp90 overproduction in TB. The length of

induction is the only variable that has a significant effect on Hsp90 production.

The yield is higher in TB compared to LB and 2x YT. The structure of the chart is

as in (A).

We noted similar trends in case of 2x YT medium, the level of protein

overproduction was insufficient and had similar yields after one and three days of

induction **(Figure 5B)**. After five days of induction in 2x YT, however, Hsp90

overproduction was still insignificant. Just like in LB, the other varied conditions

did not modulate Hsp90 levels in this medium. In 2x YT medium the yield was

higher in every condition compared to LB.

Hsp90 overproduction in TB was significantly higher than in the other two

media. The length of induction had a significant effect on the Hsp90 yields, and

here we reached good yields already after three days. The other tested

parameters did not influence the Hsp90 yield in TB.

Overall we concluded that the quantification of the gels in Figure 2, 3 and 4

confirmed our observations about the overproduction gels (**Figure 5A, B and C,**

respectively**).** We found that of the tested parameters, richness of medium used

for culturing *E. coli* cells and the length of induction were the only two that had a

significant effect on Hsp90 overproduction and should be considered in the

future. Temperature, OD_{600} at induction and IPTG concentration used for

induction did not significantly modulate the yields of the chaperone within the

tested intervals.

Recombinant human Hsp90β purified in one day

After successful protein overproduction, we set out to optimise a fast, efficient

and trustworthy purification protocol to ensure excellent sample quality. This was

necessary because like other multi-domain proteins, human Hsp90β is also prone

to the degradation. Its accessible, flexible regions, especially the charged linker

between the N-terminal domain and the middle domain is often targeted by

proteases, causing N-terminal degradation. Widely used purification protocols

include time-consuming purification steps (such as dialysis, size exclusion

chromatography or both) that slow down the procedure which may have

detrimental effects on protein quality (16-19).

To avoid degradation that might occur as a consequence of long

purification procedures and eventual freeze-thaw steps, we set out to develop a

condensed protocol for purification of Hsp90 in one day that results in highly pure

and stable sample. We purified N-terminally His$_6$-tagged Hsp90 protein by Ni-

affinity chromatography, anion exchange and heparin affinity chromatography

(**Figure 6A**). To ensure sample quality and avoid degradation we carried out the

experiment in the presence of protease inhibitors at low temperature (0-4^{o}C). To

check for protein quality throughout the purification process we ran samples of

the peak fractions of each column on sodium dodecyl sulfate polyacrylamide gel

electrophoresis (SDS-PAGE) (**Figure 6B**). We observed a gradual gain in purity

and loss in degradation products and impurities. After the last step the new

Hsp90 purification protocol resulted in a protein sample free of any significant

contaminations detectable by Coomassie staining. The combination of the three

columns was necessary for high sample purity and reproducibility.

*Figure 6: **The Hsp90 sample was free of degradation products and***

***impurities after the three-step purification procedure**. (A) Schematic overview*

of the Hsp90 purification procedure. After cell disruption and centrifugation, the

lysate was loaded first on a POROS 20MC Ni-column followed by a POROS

20HQ anion exchange column. Next, the sample was loaded on a heparin affinity

chromatography column. Finally, the eluted protein sample was concentrated,

buffer-exchanged and flash frozen. The time frame of the experiment is indicated

on the bottom of the figure. The SDS-PAGE panels on the arrows show the purity

of the samples after each purification step. (B) SDS-PAGE shows the purity of the

Hsp90 sample after each purification step. Lane 1: peak fraction sample after

POROS 20MC affinity chromatography purification step, lane 2: peak fraction

sample after POROS 20HQ anion exchange purification step, lane 3: peak

fraction sample after heparin affinity chromatography purification step.

Certain biochemical and biophysical experiments are carried out at a higher,

physiological temperature (37^{o}C) and for elevated time intervals (days or

months). In case of proteins that are sensitive to proteolysis, incubation at high

temperature for longer time can be detrimental for sample quality. To test the

stability of the Hsp90 sample purified by the new protocol we incubated the

protein at 4, 21 and 37^{o}C and ran samples taken at 3, 18 and 24 hours on SDS-

PAGE. We visualised Hsp90 in the gels first with Coomassie (**Figure 7A**) and

subsequently, with more sensitive silver staining (**Figure 7B**).

Figure 7: Purified Hsp90 sample incubated on physiological temperature for

one day remained free of degradation products. (A) SDS-PAGE about the

stability of the purified Hsp90 sample. Protein marker (lane 1), Hsp90 incubated

for 0 hours (lane 2), Hsp90 incubated at 4^{o}C (lane 3-5), 21^{o}C (lane 6-8), 37^{o}C

(lane 9-11), Hsp90 incubated for 3 hours (lane 3, 6, 9), 18 hours (lane 4, 7, 10),

24 hours (lane 5, 8, 11). (B) Silver staining about the stability of the purified

Hsp90 sample. Protein marker (lane 1), Hsp90 incubated for 0 hours (lane 2),

Hsp90 incubated at 4^{o}C (lane 3-5), 21^{o}C (lane 6-8), 37^{o}C (lane 9-11), Hsp90

incubated for 3 hours (lane 3, 6, 9), 18 hours (lane 4, 7, 10), 24 hours (lane 5, 8,

11).

No additional degradation products appeared in the incubated samples compared

to the starting sample within the time frame of the experiment on the Coomassie-

stained gels **(Figure 7A)**. Therefore we concluded that Hsp90 purified using the

new purification protocol was stable if incubated at 37^{o}C up to 24 hours. Since

Coomassie staining has limited sensitivity especially in the range of low molecular

weight, we further examined silver staining to further examine the stability of the

sample by silver staining **(Figure 7B)**. Here, we observed impurities and/or

degradation products in every sample but we did not see a systematic increase of

any bands upon incubation at 37^{o}C up to 24 hours.

This newly optimised, reproducible purification protocol enabled us to

prepare a highly pure, stable and homogeneous sample within one day.

Hsp90 purified by our method was properly folded, had the correct

molecular weight and sedimentation coefficient

To reveal the folding status of proteins, we to analysed our Hsp90 samples with

circular dichroism spectroscopy (CD) that can potentially reveal the secondary

structure composition and folding status of the chaperone. The CD spectrum of

Hsp90 revealed that the chaperone is mainly composed of α-helices **(Figure 8A)**.

We observed two minima at ~209 nm and at ~ 222 nm and a maximum at ~ 194

nm in the spectrum which is in agreement with the previous findings (Prodromou

et al, 1999).

Figure 8: Purified Hsp90 sample folded, sedimented correctly and its

molecular mass was appropriate*. (A) CD spectrum of Hsp90 at 1.2 µM*

monomer concentration. The overall structure of the protein typically consists of

α-helices. (B) The AUC spectrum of Hsp90 at 7.2 µM monomer concentration.

Hsp90 dimers sediment with 5.6 S, whereas Hsp90 tetramers appear at 10.6 S.

(C) SEC-MALLS spectra of the purified Hsp90 sample at 2.2 µM monomer

concentration. The protein has the molecular weight of the dimer Hsp90 is 167.5

± 13.3 kDa. The peak maximum is at 10.07 ml. Red line: light scattering, blue

line: refractive index, green line: UV absorption at 220 nm, black dotted line: fit for

molecular mass.

To ensure that our Hsp90 sample had the correct molecular weight and did not

degrade during the purification process we measured its molecular weight using

size exclusion chromatography - multi angle laser light scattering (SEC-MALLS)

(**Figure 8B**). We observed a homogeneous peak at 10.07 ml that was fitted to

167.5 ± 13.3 kDa. This result corroborates with the expected size of two Hsp90

molecules (168.2 kDa), confirming that our sample is in a dimeric state at the

concentration used in the experiment and also in agreement with data published

earlier (Lepvrier et al, 2015, Moullintraffort et al, 2010). Taken together with the

outcome of the stability experiments SEC-MALLS results suggested that our

sample contained the intact full length Hsp90 dimer without degradation.

We measured the sedimentation coefficient of our Hsp90 sample to

analyse sedimentation properties of the Hsp90 dimer. In the sedimentation profile

of the chaperone we observed a peak at 5.6 S that corresponded to the dimer

and a smaller peak at 10.6 S that was probably the tetramer molecule (dimer of

dimers) (**Figure 8C**). The increasing fraction above 14 S suggested that the

sample aggregated to some extent (but this is a typical phenomenon in case of

analytical ultrancentrifugation (AUC) samples). Below 1 S we observed a small

peak, that could have originated from impurities or degradation products.

Discussion

Human Hsp90β is often produced in baculovirus system in insect Sf9 cells that is expensive, requires long preparation steps and is often difficult to scale-up (20-22). Our optimised recombinant human Hsp90β production protocol in *E. coli* provides an attractive alternative for the baculovirus system.

First, we systematically tested the production of recombinant human Hsp90β in *E. coli* cells to identify parameters that modulate the yield of the chaperone. We showed that the length of induction and the nutrient level of the medium had significant effect on Hsp90 overproduction, whereas temperature, IPTG concentration and OD_{600} at induction (within the tested intervals) did not significantly modify the yields of Hsp90. Induction for three days in TB medium resulted in good yield of the chaperone.

Moreover, we purified Hsp90 using a protocol that allowed us to finish within one day and resulted in well-folded, stable and pure sample. Since Hsp90 is extremely sensitive to proteolysis because of its flexible linkers, fast preparation increases sample quality, in addition to being cheaper and time-efficient. Throughout the purification procedure impurities and degradation products disappeared from the sample and final product resisted further degradation upon incubation on elevated temperature for long time intervals.

Previous Hsp90 purification protocols often contained time consuming preparation steps such as dialysis, size exclusion chromatography or both that may affect protein quality (16-19). Using the combination of Ni-affinity chromatography, anion exchange and the Hsp90-specific heparin column allowed

us to get rid of degradation products and impurities in only three purification

steps. The resulting sample was sufficiently pure and resistant to proteolytic

degradation upon incubation on physiological temperature for up to one day, and

showed expected biophysical properties (CD, SEC-MALLS).

In our AUC experiments human Hsp90 sedimented predominantly as a

dimer with S=5.6 and a small fraction as a tetramer with S=10.6. It is notable that

human Hsp90 is known to be in extended conformation (D_{max}=26 nm) in the

absence of nucleotide and co-chaperone p23 (10). Similarly to our results, earlier

findings showed that in the absence of nucleotides, consequently in its extended

conformation, yeast Hsp90 also had the sedimentation coefficient of 5.6 (23).

These results indicate that these two proteins sediment similarly when in

extended conformation. However, in the presence of ATP that stabilises compact

conformation, yeast Hsp90 sedimented with the coefficient of 6.8, suggesting that

structural rearrangements altered its sedimentation properties (23). In similar

experiments Hsp90 sedimented with the coefficient of 6.1 (24). The variations

between these results may be explained by the differences between the origin of

the samples (porcine brain Hsp90 that has different post-translational

modification pattern and was a mixture of α and β isoform vs Hsp90), moreover,

by buffer, salt and reducing agent conditions.

In summary, we optimised Hsp90 overproduction and a fast and efficient

purification protocol for the protein. The resulting sample was pure, properly

folded and resistant to degradation, consequently it is suitable for biochemical

and biophysical experiments.

Acknowledgements

We are grateful to Ineke Braakman for continuous support. We thank Jonas M. Dörr for his help with CD experiments and data analysis. We thank Camilla de Nardis, Nadia O. Leloup and Remco N. P. Rodenburg, Deniz Ugurlar and Revina C. van Scherpenzeel for their help in setting up SEC-MALLS experiments and their support in data interpretation. We are grateful to Priyanka Sahasrabudhe for her comments on the manuscript.

Author contributions

Contribution to manuscript:

The concept of the study originates from M. Radli and S.G.D. Rüdiger.

M. Radli drafted the first version of the manuscript and prepared all the figures in the chapter. M. Radli and S.G.D. Rüdiger wrote the manuscript.

Contribution to experiments:

M. Radli tested Hsp90 production, ran the SDS-PAGE gels and quantified Hsp90 overproduction.

M. Radli optimised the purification of Hsp90 and purified all the protein used for the experiments in this chapter.

M. Radli tested the stability of Hsp90 on different temperatures.

M. Radli performed and analysed the analytical ultracentrifugation (AUC), SEC-MALLS and circular dichroism (CD) experiments.

591   D.B. Veprintsev supervised the AUC experiments, helped analyzing the data and

592   interpreting the results.

593   S.G.D. Rüdiger supervised the experiments.

References

(1) Balch WE, Morimoto RI, Dillin A, Kelly JW. Adapting proteostasis for disease intervention. Science 2008 Feb 15;319(5865):916-919.

(2) Tyedmers J, Mogk A, Bukau B. Cellular strategies for controlling protein aggregation. Nat Rev Mol Cell Biol 2010 Nov;11(11):777-788.

(3) Kim YE, Hipp MS, Bracher A, Hayer-Hartl M, Hartl FU. Molecular chaperone functions in protein folding and proteostasis. Annu Rev Biochem 2013;82:323-355.

(4) Buchberger A, Bukau B, Sommer T. Protein quality control in the cytosol and the endoplasmic reticulum: brothers in arms. Mol Cell 2010 Oct 22;40(2):238-252.

(5) Taipale M, Jarosz DF, Lindquist S. HSP90 at the hub of protein homeostasis: emerging mechanistic insights. Nature Reviews Molecular Cell Biology 2010 Jul;11(7):515-528.

(6) Li J, Buchner J. Structure, function and regulation of the hsp90 machinery. Biomed J 2013 May-Jun;36(3):106-117.

(7) Pearl LH. Review: The HSP90 molecular chaperone-an enigmatic ATPase. Biopolymers 2016 Aug;105(8):594-607.

(8) Wiech H, Buchner J, Zimmermann R, Jakob U. Hsp90 chaperones protein folding in vitro. Nature 1992;358(6382):169-170.

(9) Buchner J. Hsp90 & Co. - a holding for folding. Trends Biochem Sci 1999 Apr;24(4):136-41.

(10) Karagöz GE, Duarte AMS, Akoury E, Ippel H, Biernat J, Morán Luengo T, et al.

Hsp90-Tau complex reveals molecular basis for specificity in chaperone action.

Cell 2014;156:963-974.

(11) Kirschke E, Goswami D, Southworth D, Griffin PR, Agard DA. Glucocorticoid

receptor function regulated by coordinated action of the Hsp90 and Hsp70

chaperone cycles. Cell 2014 Jun 19;157(7):1685-1697.

(12) Workman P. Combinatorial attack on multistep oncogenesis by inhibiting the Hsp90

molecular chaperone. Cancer Lett 2004 Apr 8;206(2):149-57.

(13) Neckers L. Heat shock protein 90: the cancer chaperone. J Biosci 2007

Apr;32(3):517-530.

(14) Hipp MS, Park SH, Hartl FU. Proteostasis impairment in protein-misfolding and -

aggregation diseases. Trends Cell Biol 2014 Jun 16.

(15) Pedersen S, Bloch PL, Reeh S, Neidhardt FC. Patterns of protein synthesis in E.

coli: a catalog of the amount of 140 individual proteins at different growth rates.

Cell 1978 May;14(1):179-190.

(16) Jakob U, Meyer I, Bugl H, Andre S, Bardwell JC, Buchner J. Structural organization

of procaryotic and eucaryotic Hsp90. Influence of divalent cations on structure and

function. The Journal of biological chemistry 1995;270(24):14412-14419.

(17) Prodromou C, Siligardi G, O'Brien R, Woolfson DN, Regan L, Panaretou B, et al.

Regulation of Hsp90 ATPase activity by tetratricopeptide repeat (TPR)-domain co-

chaperones. EMBO J 1999;18(3):754-762.

(18) Richter K, Muschler P, Hainzl O, Buchner J. Coordinated ATP hydrolysis by the

Hsp90 dimer. The Journal of biological chemistry 2001;276(36):33689-33696.

(19) Southworth DR, Agard DA. Client-loading conformation of the Hsp90 molecular

chaperone revealed in the cryo-EM structure of the human Hsp90:Hop complex.

Mol Cell 2011 Jun 24;42(6):771-781.

(20) Freeman BC, Morimoto RI. The human cytosolic molecular chaperones hsp90,

hsp70 (hsc70) and hdj-1 have distinct roles in recognition of a non-native protein

and protein refolding. EMBO J 1996 Jun 17;15(12):2969-2979.

(21) Buchner J, Bose S, Mayr C, Jakob U. Purification and characterization of

prokaryotic and eukaryotic Hsp90. Methods Enzymol 1998;290:409-418.

(22) Verba KA, Wang RY, Arakawa A, Liu Y, Shirouzu M, Yokoyama S, et al. Atomic

structure of Hsp90-Cdc37-Cdk4 reveals that Hsp90 traps and stabilizes an

unfolded kinase. Science 2016 Jun 24;352(6293):1542-1547.

(23) Hessling M, Richter K, Buchner J. Dissection of the ATP-induced conformational

cycle of the molecular chaperone Hsp90. Nat Struct Mol Biol 2009 Mar;16(3):287-

293.

(24) Garnier C, Barbier P, Devred F, Rivas G, Peyrot V. Hydrodynamic properties and

quaternary structure of the 90 kDa heat-shock protein: effects of divalent cations.

Biochemistry 2002 Oct 1;41(39):11770-11778.

A

Lane 1
Protein
marker

Lane 2
Uninduced
sample

Lane 3-14
Inducted
samples

Lane 15
Purified
Hsp90

Day 1   Day 3   Day 5

Hsp90

EF-Tu

B

Lane profiles before adjustments

🟠 Uninduced lane profile
🟢 Induced lane profile

EF-Tu

Hsp90

Normalisation and alignment of induced lane profile

🟠 Uninduced lane profile
🟢 Normalised, aligned
   induced lane profile

EF-Tu

Hsp90

Surface area of the Hsp90 peak

🔴 Difference between uninduced
   and inducaed lane profiles

Hsp90

A

Cell lysis,
centrifugation

20MC

20HQ

Heparin

Concentration

90'   60'   30'   120'   120'

B

After column   20MC   20HQ   Heparin

Hsp90 ——