1   <u>Title</u>: Genetic costs of domestication and improvement
2
3

4   <u>Authors</u>: Brook T. Moyers[1], Peter L. Morrell[2], and John K. McKay[1]
5
6
7   <u>Affiliations</u>: [1] Bioagricultural Sciences and Pest Management, Colorado State
8   University, Fort Collins, CO, USA 80521. [2] Department of Agronomy and Plant
9   Genetics, University of Minnesota, St. Paul, MN, USA 55108.
10
11
<u>Address for correspondence</u>:
Brook T. Moyers
Bioagricultural Sciences and Pest Management
Colorado State University
Fort Collins, CO, USA 80521
Phone: 707-235-1237
Email: brook.moyers@gmail.com

19
20
21   <u>Running title</u>: Cost of domestication
22
23
24

25 <u>ABSTRACT</u>
26
The 'cost of domestication' hypothesis posits that the process of domesticating

wild species increases genetic load by increasing the number, frequency, and/or

proportion of deleterious genetic variants in domesticated genomes. This cost

may limit the efficacy of selection and thus reduce genetic gains in breeding

programs for these species. Understanding when and how genetic load evolves

can also provide insight into fundamental questions about the interplay of

demographic and evolutionary dynamics. Here we describe the evolutionary

forces that may contribute to genetic load during domestication and

improvement, and review the available evidence for 'the cost of domestication' in

animal and plant genomes. We identify gaps and explore opportunities in this

emerging field, and finally offer suggestions for researchers and breeders

interested in addressing genetic load in domesticated species.
39

40 <u>Keywords</u>: genetic load, deleterious mutations, crops, domesticated animals

INTRODUCTION

Recently, we have seen a resurgence of evolutionary thinking applied to domesticated plants and animals (e.g. Walsh 2007; Wang *et al*. 2014; Gaut, Díez, and Morrell 2015; Kono *et al*. 2016). One particular wave of this resurgence proposes a general 'cost of domestication': that the evolutionary processes experienced by lineages during domestication are likely to have increased genetic load. This 'cost' was first hypothesized by Lu *et al*. (2006), who found an increase in nonsynonymous substitutions, particularly radical amino acid changes, in domesticated compared to wild lineages of rice. These putatively deleterious mutations were negatively correlated with recombination rate, which the authors interpreted as evidence that these mutations hitchhiked along with the targets of artificial selection (Figure 1B). Lu *et al*. (2006) conclude that: "The reduction in fitness, or the genetic cost of domestication, is a general phenomenon." Here we address this claim by examining the evidence that has emerged in the last decade on genetic load in domesticated species.

The processes of domestication and improvement potentially impose a number of evolutionary forces on populations (Box 1), starting with mutational effects. New mutations can have a range of effects on fitness, from lethal to beneficial. Deleterious variants constitute the mostly directly observable, and likely the most important, source of genetic load in a population. The shape of the distribution of fitness effects of new mutations is difficult to estimate, but theory predicts that a large proportion of new mutations, particularly those that occur in coding portions of the genome, will be deleterious at least in some proportion of the environments that a species occupies (Ohta 1972; 1992; Gillespie 1994). These predictions are supported by experimental responses to artificial selection and mutation accumulation experiments, and molecular genetic studies (discussed in Keightley and Lynch 2003).

The rate of new mutations in eukaryotes varies, but is likely at least 1×10^{-8} / base pair / generation (Baer, Miyamoto, and Denver 2007). For the average

eukaryotic genome, individuals should thereby be expected to carry a small

number of new mutations not present in the parent genome(s) (Agrawal and

Whitlock 2012). The realized distribution of fitness effects for these mutations will

be influenced by inbreeding and by effective population size (Figure 2A; Gillespie

1999; Whitlock 2000; Arunkumar *et al*. 2015; Keightley and Eyre-Walker 2007).

Generally, for a given distribution of fitness effects for new mutations, we expect

to observe relatively fewer strongly deleterious variants and more weakly

deleterious variants in smaller populations and in populations with higher rates of

inbreeding (Figure 2A).

Domestication and improvement often involve increased inbreeding. Allard

(1999) pointed out that many important early cultigens were inbreeding, with

inbred lines offering morphologically consistency in the crop. In some cases,

domestication came with a switch in mating system from highly outcrossing to

highly selfing (e.g. in rice; Kovach, Sweeney, and McCouch 2007). The practice

of producing inbred lines, often through single-seed descent, with the specific

intent to reduce heterozygosity and create genetically 'stable' varieties, remains a

major activity in plant breeding programs. Reduced effective population sizes

(N_e) during domestication and improvement and artificial selection on favorable

traits also constitute forms of inbreeding (Figure 1). Even in species without the

capacity to self-fertilize, inbreeding that results from selection can dramatically

change the genomic landscape. For example, the first sequenced dog genome,

from a female boxer, is 62% homozygous haplotype blocks, with an N50 length

of 6.9 Mb (versus heterozygous region N50 length 1.1 Mb; Lindblad-Tor *et al*.

2005). This level of homozygosity drastically reduces the effective recombination

rate, as crossover events will essentially be switching identical chromosomal

segments. The likelihood of a beneficial allele moving into a genomic background

with fewer linked deleterious alleles is thus reduced.

Linked selection, or interference among mutations, has much the same effect.

For a trait influenced by multiple loci, interference between linked loci can limit

the response to selection (Hill and Robertson 1966). Deleterious variants are

more numerous than those that have a positive effect on a trait and thus should

constitute a major limitation in responding to selection (Felsenstein 1974). This

forces selection to act on the net effect of favorable and unfavorable mutations in

linkage (Figure 1B). As inbreeding increases, selection is less effective at purging

moderately deleterious mutations and new slightly beneficial mutations more

likely to be lost by genetic drift, which will shift the distribution of fitness effects for

segregating variants and result in greater accumulation of genetic load over time

(Figure 2A–C).

Overall, the cost of domestication hypothesis posits that compared to their wild

relatives domesticated lineages will have:

1. Deleterious variants at higher frequency, number, and proportion

2. Enrichment of deleterious variants in linkage disequilibrium with loci

subject to strong, positive, artificial selection

With domestications, these effects may differ between lineages that experienced

domestication followed by modern improvement (e.g. 'elite' varieties and

commercial breeds) and those that experienced domestication only (e.g.

landraces and non-commercial populations). Generally speaking, the process of

domestication is characterized by a genetic bottleneck followed by a long period

of relatively weak and possibly varying selection, while during the process of

improvement, intense selection over short time periods is coupled with limited

recombination and an additional reduction in N_e, often followed by rapid

population expansion (Figure 1A; Yamasaki, Wright, and McMullen 2007). Gaut,

Díez, and Morrell (2015) propose that elite crop lines should harbor a lower

proportion of deleterious variants relative to landraces due to strong selection for

yield during improvement, but the opposing pattern could be driven by increased

genetic drift due to limited recombination, lower N_e, and rapid population

expansion. At least one study shows little difference in genetics load between

landrace and elite lines in sunflower (Renaut and Rieseberg 2015). It is likely that

relative influence of these factors varies dramatically across domesticated
systems.

136
____
Box 1. Domesticated lineages may experience:
• **Reduced efficacy of selection:**
○ Deleterious variants that are in linkage disequilibrium with a target
of artificial selection can rise to high frequency through genetic
hitchhiking, as long as the their fitness effects are smaller than the
effect of the targeted variant (Figure 1B).
○ Domesticated lineages can experience reduced effective
recombination rate as inbreeding increases and heterozygosity
drops (Figure 1B). Increased linkage disequilibrium allows larger
portions of the genome to hitchhike, and negative linkage
disequilibrium across advantageous alleles on different haplotypes
can cause selection interference (Felsenstein 1974).
○ The loss of allelic diversity through artificial selection, genetic drift,
and inbreeding (Figure 1B) can reduce trait variance (Eyre-Walker,
Woolfit, and Phelps 2006), which reduces the efficacy of selection.
○ **However**: any significant increase in inbreeding, particularly the
transition from outcrossing to selfing, **allows selection to more**
efficiently purge recessive, highly deleterious alleles, as these
alleles are exposed in homozygous genotypes more frequently
(Arunkumar *et al*. 2015). This shifts the more deleterious end of the
distribution of fitness effects for segregating variants towards
neutrality (Figure 2A).
160
• **Increased genetic drift:**
○ Domestication involves one or more genetic bottlenecks (Figure
1A), where N_e is reduced. In smaller populations, genetic drift is
stronger relative to the strength of selection, and deleterious
variants can reach higher frequencies.

o N_e varies across the genome (Charlesworth 2009). In particular,
when regions surrounding loci under strong selection (e.g.
domestication QTL) rapidly rise to high frequency, the N_e for that
haplotype rapidly drops, and so drift has a relatively stronger effect
in that region.
o The rapid population expansion coupled with long-distance
migration typical of many domesticated lineages' demographic
histories can result in the accumulation of deleterious genetic
variation known as 'expansion load' (Peischl *et al*. 2013;
Lohmueller 2014). This occurs when serial bottlenecks are followed
by large population expansions (e.g., large local carrying
capacities, large selection coefficients, long distance dispersal;
Peischl *et al*. 2013). This phenomenon is due to the accumulation
of new deleterious mutations at the 'wave front' of expanding
populations, which then rise to high frequency via drift regardless of
their fitness effects (i.e. 'gene surfing', or 'allelic surfing' Klopfstein,
Currat, and Excoffier 2006; Travis *et al*. 2007).

• **Increased mutation load**
o Domesticated and wild lineages may differ in basal mutation rate,
although it is unclear if we would expect this to be biased towards
higher rates in domesticated lineages.
o More likely, deleterious mutations **accumulate** at higher rates in
domesticated lineages due to the reduced efficacy of selection.
Mutations that would be purged in larger N_e or with higher effective
recombination rates are instead retained. This is reflected in a shift
in the distribution of fitness effects for segregating variants towards
more moderately deleterious alleles (Figure 2A).
o Smaller population sizes decrease the likelihood of beneficial
mutations arising, as these mutations are likely relatively rare and

196   for a given mutation rate fewer individuals mean fewer opportunities

197   for mutation.

____

<u>Genetic Load in Domesticated Species</u>

The increasing number of genomic datasets has made the study of genome-wide

genetic load increasingly feasible. Since Lu *et al*. (2006) first hypothesized

genetic load as a 'cost of domestication' in rice, similar studies have examined

evidence for such costs in other crops and domesticated animals.

One approach for estimating genetic load is to compare population genetic

parameters between two lineages, in this case domesticated species versus their

wild relatives. If domestication comes with the cost of genetic load, we might

expect domesticated lineages to have accumulated more nonsynonymous

mutations or, if the accumulation of genetic load is due to reduced effective

population sizes, to show reduced allelic diversity and effective recombination

when compared to their wild relatives. While relatively straightforward, this

approach only indirectly addresses genetic load by assuming that: (A)

nonsynonymous mutations are on average deleterious, and (B) loss of diversity

and reduced recombination indicate a decreased efficacy of selection. This

approach also assumes that wild relatives of domesticated species have not

themselves experienced population bottlenecks, shifts in mating system, or any

of the other processes that could affect genetic load relative to the ancestors.

Another approach for estimating genetic load is to assay the number and

proportions of putatively deleterious variants present in populations of

domesticated species, using various bioinformatic approaches. This approach

directly addresses the question of genetic load, but does not tell us whether the

load is the result of domestication or other evolutionary processes (unless wild

relatives are also assayed, again assuming that their own evolutionary trajectory

has not simultaneously been affected). It also requires accurate, unbiased

algorithms for identifying deleterious variants from neutral variants (see below).

Studies taking these approaches in domesticated species are presented Table 1.

We searched the literature using Google Scholar with the terms "genetic load"

and "domesticat*", and in many cases followed references from one study to the

next. To the best of our knowledge, the studies in Table 1 represent the majority

of the extant literature on this topic. We excluded studies examining only the

mitochondrial or other non-recombining portions of the genome. In a few cases

where numeric values were not reported, we extracted values from published

figures using relative distances as measured in the image analysis program

ImageJ (Schneider, Rasband, and Eliceiri 2012), or otherwise extrapolated

values from the provided data. Where exact values were not available, we

provide estimates. Please note that no one study (or set of genotypes)

contributed values to all columns for a particular domestication event, and that in

many cases methods used or statistics examined varied across species.

Recombination and linkage disequilibrium

The process of domestication may increase recombination rate, as measured in

the number of chiasmata per bivalent. Theory predicts that recombination rate

should increase during periods of rapid evolutionary change (Otto and Barton

1997), and in domesticated species this may be driven by strong selection and

limited genetic variation under linkage disequilibrium (Otto and Barton 2001).

This is supported by the observation that recombination rates are higher in many

domesticated species compared to wild relatives (e.g., chicken, Groenen *et al*.

2009; honey bee, Wilfert *et al.* 2007; and a number of cultivated plant species,

Ross-Ibarra 2004). In contrast, recombination in some domesticated species is

limited by genomic structure (e.g., in barley, where 50% of physical length and

many functional genes are in (peri-)centromeric regions with extremely low

recombination rates; The International Barley Genome Sequencing Consortium

2012), although this structure may be shared with wild relatives. Even when

actual recombination rate (chiasmata per bivalent) increases, effective

recombination may be reduced in many domesticated species due to an increase

in the physical length of fixed or high frequency haplotypes, or linkage

disequilibrium (LD). In other words, chromosomes may physically recombine, but

if the homologous chromosomes contain identical sequences then no "effective

recombination" occurs and the outcome is no different than no recombination. In

a striking example, the parametric estimate of recombination rate in maize

populations, has decreased by an average of 83% compared to the wild relative

teosinte (Wright *et al.* 2005). We note that Mezmouk and Ross-Ibarra (2014)

found that in maize deleterious variants are not enriched in areas of low

recombination (although see McMullen *et al.* 2009; Rodgers-Melnick *et al.* 2015).

Strong directional selection like that imposed during domestication and

improvement can reduce genetic diversity in long genetic tracts linked to the

selected locus (Maynard Smith and Haigh 1974). These regions of extended LD

are among the signals used to identify targets of selection and reconstruct the

evolutionary history of domesticated species (e.g., Tian, Stevens, and Buckler

2009). A shift in mating system from outcrossing to selfing, or an increase in

inbreeding, will have the same effect across the entire recombining portion of the

genome, as heterozygosity decreases with each generation (Charlesworth 2003).

Extended LD has consequences for the efficacy of selection: deleterious variants

linked to larger-effect beneficial alleles can no longer recombine away, and

beneficial variants that are not in LD with larger-effect beneficial alleles may be

lost. We see evidence for this pattern in our review of the literature: LD decays

most rapidly in the outcrossing plant species in Table 1 (maize and sunflower),

and extends much further in plant species propagated by selfing and in

domesticated animals. In all cases where we have data for wild relatives, LD

decays more rapidly in wild lineages than in domesticated lineages (Table 1).

While we primarily report mean LD in Table 1, linkage disequilibrium also varies

among varieties and breeds of the same species. For example, in domesticated

pig breeds the length of the genome covered by runs of homozygosity ranges

from 13.4 to 173.3 Mb, or 0.5–6.5% (Traspov *et al*. 2016). Similarly, among dog

breeds average LD decay (to $r^2 \leq 0.2$) ranges from 20 kb to 4.2 Mb (Gray *et al*.

2009). In both species, the decay of LD for wild individuals occurs over even

shorter distances. This suggests that patterns of LD in these species have been

strongly impacted by breed-specific demographic history (i.e., the process of

improvement), in addition to the shared process of domestication.

Genetic diversity and dN/dS

In the species we present in Table 1, we see consistent loss of genetic diversity

when 'improved' or 'breed' genomes are compared to domesticated 'landrace' or

'non-commercial' genomes, and again when domesticated genomes are

compared to the genomes of wild relatives. This ranges from ~5% nucleotide

diversity lost between wolf populations and domesticated dogs (Gray *et al*. 2009)

to 77% lost between wild and improved tomato populations (Lin *et al*. 2014). The

only case where we see a gain in genetic diversity is in the Andean

domestication of the common bean, where gene flow with the more genetically

diverse Mesoamerican common bean is likely an explanatory factor (Schmutz *et*

al. 2013). This pattern is consistent with a broader review of genetic diversity in

crop species: Miller and Gross (2011) found that annual crops had lost an

average of ~40% of the diversity found in their wild relatives. This same study

found that perennial fruit trees had lost an average of ~5% genetic diversity,

suggesting that the impact of domestication on genetic diversity is strongly

influenced by life history (see also Gaut, Díez, and Morrell 2015). Given a

relatively steady evolutionary trajectory for wild populations, loss of genetic

diversity in domesticated populations can be attributed to: genetic bottlenecks,

increased inbreeding, artificial selection. On an evolutionary timescales, even the

oldest domestications occurred quite recently relative to the rate at which new

mutations can recover the loss. This is especially true for non-recombining

portions of the genome like the mitochondrial genome or non-recombining sex

chromosomes. Most modern animal breeding programs are strongly sex-biased,

with few male individuals contributing to each generation. In horses, for example,

this has likely led to almost complete loss of polymorphism on the Y chromosome

via genetic drift (Lippold *et al*. 2011). Loss of allelic diversity can reduce the

efficacy of selection by reducing trait variation within species (Eyre-Walker,

Woolfit, and Phelps 2006). However, a loss of genetic diversity alone does not

necessarily signal a corresponding increase in the frequency or proportion of

deleterious variants, and so is not sufficient evidence of genetic load.

In six of seven species, the domesticated lineage shows an increase in genome-

wide nonsynonymous to synonymous substitution rate or number compared to

the wild relative lineage (Table 1). The exception is in soybean, where the

domesticated *Glycine max* and wild *G. soja* genomes contain approximately the

same proportion of nonsynonymous to synonymous single nucleotide

polymorphisms (Lam *et al*. 2010). These differences in nonsynonymous

substitution rate are likely driven by differences in N_e (Eyre-Walker and Keightley

2007; Woolfit 2009). If most nonsynonymous mutations are deleterious, as theory

and empirical data suggest, then an increase in nonsynonymous substitutions

will reduce mean fitness (Figure 2B-C). The comparisons in Table 1 suggest that

this has occurred in domesticated species. This result differs from Moray,

Lanfear, and Bromham (2014), who examined rates of mitochondrial genome

sequence evolution in domesticated animals and their wild relatives and found no

such consistent pattern. This difference may be attributable to the focus of each

review (genome-wide versus mitochondria) or, as the authors speculate, to

genetic bottlenecks in some of the wild relatives included in their study.

The ratio of nonsynonymous to synonymous substitutions may not be a good

estimate for mutational load. For one, nonsynonymous substitutions are

particularly likely to be phenotype changing (Kono *et al*. *bioRxiv*) and contribute

to agronomically important phenotypes (Kono *et al*. 2016). Thus artificial

selection during domestication and improvement is likely to drive these variants

to higher frequencies (or fixation) in domesticated lineages. Thus variants that

annotate as deleterious based on sequence conservation (see below), can in

some case, contribute to agronomically important phenotypes. In addition,

estimates of the proportion of nonsynonymous sites with deleterious effects

range from 0.03 (in bacteria; Hughes 2005) to 0.80 (in humans; Fay *et al*. 2001;

The Chimpanzee Sequencing and Analysis Consortium 2005). It follows that

nonsynonymous substitution rate may have a poor correlation with genetic load,

at least at larger taxonomic scales. However, the consensus proportion of

deleterious variants in Table 1 is between 0.05 and 0.25, which spans a smaller

range. Finally, dN/dS may in itself be a flawed estimate of functional divergence

because it relies on the assumptions that dS is neutral and mean dS can control

for substitution rate variation, which may not hold true, especially for closely

related taxa (Wolf *et al*. 2009; see also Kryazhimskiy and Plotkin 2008).

Deleterious variants

An increase in genetic load can come from: (1) increased frequency of

deleterious variants, (2) increased number of deleterious variants, and (3)

enrichment of deleterious variants relative to total variants. The first can be

assessed by examining shared deleterious variants between wild and

domesticated lineages, and all three by comparing both shared and private

deleterious variants across lineages. Looking at deleterious variants in high

frequency across all domesticated varieties may provide insight into the early

processes of domestication, while looking at deleterious variants with varying

frequencies among domesticated varieties may provide insight into the

processes of improvement. We recommend that researchers studying

deleterious variants present their results per genome, and then compare across

genomes and among lineages. Many, but not all, of the studies in Table 1 take

this approach. The total number, frequency, or proportion of deleterious variants

within a population will necessarily depend on the size of that population, and so

sufficient sampling is important before values can be compared across

populations. Values per genome are therefore easier to compare across most

available genomic datasets.

Renaut and Rieseberg (2015) found a significant increase in both shared and

private deleterious mutations in domesticated relative to wild lines of sunflower,

and similar patterns in two additional closely-related species: cardoon and globe

artichoke (Table 1). This pattern also holds true for *japonica* and *indica*

domesticated rice (Liu *et al*. 2017) and for the domesticated dog (Marsden *et al*.

2016) compared to their wild relatives (Table 1). In horses, deleterious mutation

load as estimated using Genomic Evolutionary Rate Profiling (GERP) appears to

be higher in both domesticated genomes and the extant wild relative

(Przewalski's horse, which went through a severe genetic bottleneck in the last

century) compared to an ancient wild horse genome (Schubert *et al*. 2014).

These four studies, spanning a wide taxonomic range, suggest that an increase

in the number and proportion of deleterious variants may be a general

consequence of domestication. However, the other studies we present that

examined deleterious variants in domesticated species did not include any

sampling of wild lineages. Without sufficient sampling of a parallel lineage that

did not undergo the process of domestication, it is difficult to assess whether the

'cost of domestication' is indeed general.

Identifying dangerous hitchhikers

The effect size of a deleterious mutation is negatively correlated with its

likelihood of increasing in frequency through any of the mechanisms we discuss

here. That is, variants with a strongly deleterious effect are more likely to be

purged by selection than mildly deleterious variants, regardless of the efficacy of

selection. Similarly, mutations that have a consistent, environmentally-

independent deleterious effect are more likely to be purged than mutations with

environmentally-plastic effects. In the extreme case, we would never expect

mutations that have a consistently lethal effect in a heterozygous state to

contribute to genetic load, as these would be lost in the first generation of their

appearance in a population. However, mutations with consistent, highly

deleterious effects are likely rare relative to those with smaller or environment-

dependent effects, especially in inbred populations (Figure 2A; Arunkumar *et al*.
2015). When thinking about genetic load, it is therefore important to recognize
that the effect of any particular mutation can depend on context, including
genomic background and developmental environment. This complexity likely
makes these classes of deleterious mutations more difficult to identify, and we
might not expect these variants to show up in our bioinformatic screens
(discussed below). Furthermore, all of these expectations are modified by
linkage: when deleterious variants are in LD with targets of artificial selection,
they are more likely to evade purging even with consistent, large, deleterious
effects (Figure 1B).
422
We searched the literature for examples of specific deleterious variants that
hitchhiked along with targets of selection during domestication and improvement.
We did not find many such cases, so we describe each in detail here. The best
characterized example comes from rice, where an allele that negatively affects
yield under drought (*qDTY1.1*) is tightly linked to the major green revolution
dwarfing allele *sd1* (Vikram *et al*. 2015). Vikram *et al*. (2015) found that the
qDTY1.1 allele explained up to 31% of the variance in yield under drought across
three RIL populations and two growing seasons. Almost all modern elite rice
varieties carry the *sd1* allele (which increases plant investment in grain yield),
and as a consequence are drought sensitive. The discovery of the *qDTY1.1*
allele has enabled rice breeders to finally break the linkage and create drought
tolerant, dwarfed lines.
435
In sunflower, the *B* locus affects branching and was a likely target of selection
during domestication (Bachlava *et al*. 2010). This locus has pleiotropic effects on
plant and seed morphology that, in branched male restorer lines, mask the effect
of linked loci with both 'positive' (increased seed weight) and 'negative' (reduced
seed oil content) effects (Bachlava *et al*. 2010). To properly understand these
effects required a complex experimental design, where these linked loci were
segregated in unbranched (*b*) and branched (*B*) backgrounds. Managing these

effects in the heterotic sunflower breeding groups has likely also been

challenging.

A similarly complex narrative has emerged in maize. The gene *TGA1* was key to

evolution of 'naked kernels' in domesticated maize from the encased kernels of

teosinte (Wang *et al*. 2015). This locus has pleiotropic effects on kernel features

and plant architecture, and is in linkage disequilibrium with the gene *SU1*, which

encodes a starch debranching enzyme (Brandenburg *et al*. 2017). *SU1* was

targeted by artificial selection during domestication (Whitt *et al.* 2002), but also

appears to be under divergent selection between Northern Flints and Corn Belt

Dents, two maize populations (Brandenburg *et al*. 2017). This is likely because

breeders of these groups are targeting different starch qualities, and this work

may have been made more difficult by the genetic linkage of *SU1* with *TGA1*.

In the above cases, the linked allele(s) with negative agronomic effects are

unlikely to be picked up in a genome-wide screen for deleterious variation, as

they are segregating in wild or landrace populations and are not necessarily

disadvantageous in other contexts. One putative 'truly' deleterious case is in

domesticated chickens, where a missense mutation in the thyroid stimulating

hormone receptor (*TSHR*) locus sits within a shared selective sweep haplotype

(Rubin *et al*. 2010). However, Rubin *et al*. (2010) argue this is more likely a case

where a 'deleterious' (i.e. non-conserved) allele was actually the target of artificial

selection and potentially contributed to the loss of seasonal reproduction in

chickens.

In the Roundup Ready (event 40-3-2) soybean varieties released in 1996, tight

linkage between the transgene insertion event and another allele (or possibly an

allele created by the insertion event itself) reduced yield by 5-10% (Elmore *et al*.

2001). This is not quite genetic hitchhiking in the traditional sense, but the yield

drag effect persisted through backcrossing of the transgene into hundreds of

varieties (Benbrook 1999). This effect likely explains, at least in part, why

transgenic soybean has failed to increase realized yields (Xu *et al*. 2013). A
second, independent insertion event in Roundup Ready 2 Yield® does not suffer
from the same yield drag effect (Horak *et al*. 2015).
477
It is likely that 'dangerous hitchhiker' examples exist that have either gone
undetected by previous studies (possibly due to low genomic resolution, limited
phenotyping, or limited screening environments), have been detected but not
publicized, or are buried among other results in, for example, large QTL studies.
It is also possible that the role of genetic hitchhiking has not been as important in
shaping genome-wide patterns of genetic load as previously assumed.
484
<u>Gaps and Opportunities</u>
For major improvement alleles, how much of the genome is in LD?
Artificial selection during domestication targets a clear change in the optimal
multivariate phenotype. This likely affects a significant portions of the genome:
available estimates include 2–4% of genes in maize (Wright *et al*. 2005) and 16%
of the genome in common bean (Papa *et al*. 2007) targeted by selection during
domestication. For crops, traits such as seed dormancy, branching,
indeterminate flowering, stress tolerance, and shattering are known to be
selected for different optima under artificial versus natural selection (Takeda and
Matsuoka 2008; Gross and Olsen 2010). In some cases, we know the loci that
underlie these domestication traits. One well studied example is the green
revolution dwarfing gene, *sd1* in rice. *sd1* is surrounded by a 500kb region (~13
genes) with reduced allelic diversity in *japonica* rice (Asano *et al*. 2011). Another
example in rice is the *waxy* locus, where a 250 kb tract shows reduced diversity
consistent with a selective sweep in temperate *japonica* glutinous varieties
(Olsen *et al*. 2006). The difference in the size of the region affected by these two
selective sweeps may be because the strength of selection on these two traits
varied, with weaker selection at *waxy* then *sd1*. Unfortunately, the relative
strength of selection on domestication traits is largely unknown, and other factors
can also influence the size of the genomic region affected by artificial selection.

The physical position of selected mutation can have large effect on this via gene

density and local recombination rate (e.g. in rice, Flowers *et al*. 2012). This

explanation has been invoked in maize (Wright *et al*. 2005), where the extent of

LD surrounding domestication loci is highly variable. For example a 1.1 Mb

region (~15 genes) lost diversity during a selective sweep on chromosome 10 in

maize (Tian, Stevens, and Buckler 2009), but only a 60–90 kb extended

haplotype came with the *tb1* domestication allele (Clark *et al*. 2004). While these

case studies provide examples of sweeps resulting from domestication, they also

show that the size of the affected region is highly variable, and we don't yet know

how this might impact genetic load. This is compounded by the fact that even in

highly researched species we don't always know what were the genomic targets

of selection during domestication (e.g., in maize; Hufford *et al*. 2012), especially if

any extended LD driven by artificial selection has eroded or the intensity or mode

of artificial selection has changed over time.

What's "worse" in domesticated species: hitchhikers, drifters, or inbreds?

We do not have a clear sense of which evolutionary processes contribute most to

genetic load, and sometimes see contrasting patterns across species. In maize,

relatively few putatively deleterious alleles are shared across all domesticated

lines (hundreds vs. thousands; Mezmouk and Ross-Ibarra 2014), which points to

a larger role for the process of improvement than for domestication in driving

genetic load. In contrast, the increase in dog dN/dS relative to wolf populations

appears to not be driven by recent inbreeding (i.e., improvement) but by the

ancient domestication bottleneck common to all dogs (Marsden *et al*. 2016).

Of the deleterious alleles segregating in more than 80% of maize lines, only 9.4%

show any signal of positive selection (Mezmouk and Ross-Ibarra 2014). This

suggests that hitchhiking during domestication played a relatively small role in

the evolution of deleterious variation in maize. The same study found little

support for enrichment of deleterious SNPs in areas of reduced recombination

(Mezmouk and Ross-Ibarra 2014). However, Rodgers-Melnick *et al*. (2015)

present contrasting evidence supporting enrichment of deleterious variants in

regions of low recombination, and the authors argue that this difference is due to

the use of a tool that does not rely on genome annotation (Genomic Evolutionary

Rate Profiling, or GERP; Cooper *et al*. 2005). This complex narrative has

emerged from just one domesticated system, and it is likely that each species will

present new and different complexities.

Specific differences

Examining general differences between wild and domesticated lineages ignores

species-specific demographic histories and changes in life history, which may be

important contributors to genetic load. Although we found some general patterns

(e.g., loss of genetic diversity, Table 1), we also see clear exceptions (e.g., the

Andean common bean). We can attribute these exceptions to particular

demographic scenarios (e.g., gene flow with Mesoamerican common bean

populations), assuming we have sufficient archeological, historical, or genetic

data. One clear problem is our inability to sample ancestral (pre-domestication)

lineages, and our subsequent reliance on sampling of current wild relative

lineages that have their own unique evolutionary trajectories. Sequencing ancient

DNA can provide some insight into the history of these lineages and their

ancestral states (e.g., in horses; Schubert *et al*. 2014). Currently, we know very

little about most domesticated species' histories. We are still working towards

understanding dynamics since domestication: even in highly-researched species

like rice, the number of domestication events and subsequent demographic

dynamics are hotly contested (Kovach, Sweeney, and McCouch 2007; Gross and

Zhao 2014; Chen, Huang, and Han 2016; Choi *et al*. 2017).

A second challenge, briefly mentioned above, is understanding the relative

importance of each of the factors in any particular domestication event. For

example, in rice the shift to selfing from outcrossing during domestication

appears to have played a larger role than the domestication bottleneck in

shaping genetic load (Liu *et al*. 2017). This is useful in understanding rice

domestication and its impact, but similar studies would need to be conducted

across domesticated species to understand the generality of this dynamic. It is

possible that general patterns may be drawn from subsets of domesticated

species (e.g., vertebrates versus vascular plants, short-lived versus long-lived,

outcrossing versus selfing versus clonally propagated etc.). For example, there

may be general differences between annual and perennial crops, including less

severe domestication bottlenecks and higher levels of gene flow from wild

populations in perennials (Miller and Gross 2011; Gaut, Díez, and Morrell 2015).

Predictive algorithms

The identification of individual deleterious variants typically relies on sequence

conservation. That is, if a particular nucleotide site or encoded amino acid is

invariant across a phylogenetic comparison of related species, a variant is

considered more likely to be deleterious. More advanced approaches use

estimates of synonymous substitution rates at a locus to improve estimates of

constraint on a nucleotide site (Chun and Fay 2009). The majority of 'SNP

annotation' approaches are intended for the annotation of amino acid changing

variants, although at least one (GERP++; Davydov *et al*. 2010) can be applied to

noncoding sequences in cases where nucleotide sequences can be aligned

across species. This estimation of phylogenetic constraint is heavily dependent

on the sequence alignment used, and so new annotation approaches have

sought to use more consistent sets of alignments across loci. Both GERP++ and

MAPP permit users to provide alignments for SNP annotation (Davydov *et al*

2010; Stone and Sidow 2005). The recently reported tool BAD_Mutations (Kono

et al. 2016; Kono *et al*. bioRxiv) permits the use of a consistent set of alignments

for the annotation of deleterious variants by automating the download and

alignment of the coding portion of plant genomes from Phytozome and Ensembl

Plants, which include 50+ sequenced angiosperm genomes (Goodstein *et al*.

2010; Kersey *et al*. 2016). Currently, the tool is configured for use with

angiosperms, but could be applied to other organisms.

Using phylogenetic conservation may be problematic in domesticated species, as
relaxed, balancing, or diversifying selection in agricultural environments could lift
constraint on sites under purifying or stabilizing selection in wild environments
(assuming some commonalities within these environment types). In one such
case, two AGPase small subunit paralogs in maize appear to be under
diversifying and balancing selection, respectively, even though these subunits
are likely under selective constraint across flowering plants (Georgelis, Shaw,
and Hannah 2009; Corbi *et al*. 2010). Similarly, broadly 'deleterious' traits may
have been under positive artificial selection in domesticated species. The loci
underlying these traits could flag as deleterious in bioinformatic screens despite
increasing fitness in the domesticated environment. For example, the *fgf4*
retrogene insertion that causes chrondrodysplasia (short-leggedness) in dogs
would likely have a strongly deleterious effect in wolves, but has been positively
selected in some breeds of dog (Parker *et al*. 2009). Finally, bioinformatic
approaches that rely on phylogenetic conservation are likely to miss variants with
effects that are only deleterious in specific environmental or genomic contexts
(plastic or epistatic effects), or which reduce fitness specifically in agronomic or
breeding contexts. Specific knowledge of the phenotypic effects of putatively
deleterious mutations is necessary to address these issues, but as we discuss
below these data are challenging to obtain.
618
Information from the site frequency spectrum, or the number of times individual
variants are observed in a sample, can provide additional information about
which variants are most likely to be deleterious. In resequencing data from many
species, nonsynonymous variants typically occur at lower average frequencies
than synonymous variants (cf. Nordborg *et al*. 2005; Ross-Ibarra *et al*. 2009;
Günther and Schmid 2010). Mutations that annotate as deleterious are
particularly likely to occur at lower frequencies than other classes of variants
(Marth *et al*. 2011; Kono *et al*. 2016; Liu *et al*. 2017) and may be less likely to be
shared among populations (Marth *et al*. 2011).
628

Tools for the annotation of potentially deleterious variants continue to be

developed rapidly (see Grimm *et al*. 2015 for a recent comparison). This includes

many tools that attempt to make use of information beyond sequence

conservation, including potential effects of variant on protein structure or function

(Adzhubei *et al*. 2010) or a diversity of genomic information intended to improve

prediction of pathogenicity in humans (Kircher *et al*. 2014). This contributes to the

issue that the majority of SNP annotation tools are designed to work on human

data and may not be applicable to other organisms (Kono *et al*. *bioRxiv*). There is

the potential for circularity when an annotation tool is trained on the basis of

pathogenic variants in humans and then evaluated on the basis of a potentially

overlapping set of variants (Grimm *et al*. 2015). Even given these limitations,

validation outside of humans is more challenging because of a paucity of

phenotype-changing variants. To address this issue, Kono et al. (*bioRxiv*) report

a comparison of seven annotation tools applied to a set of 2,910 phenotype

changing variants in the model plant species *Arabidopsis thaliana*. The authors

find that all seven tools more accurately identify phenotyping changing variants

likely to be deleterious in *Arabidopsis* than in humans, potentially because the

larger N_e in *Arabidopsis* relative to human populations may allow more effective

purifying selection.

No one is perfect, not even the reference

Bioinformatic approaches can suffer from reference bias (Simons *et al*. 2014;

Kono *et al*. 2016). One particular challenge in the identification of deleterious

variants is that with reference-based read mapping, variants are typically

identified as differences from reference, then passed through a series of filters to

identify putatively deleterious changes. For example, most annotation

approaches single out nonsynonymous differences from references. Because a

reference genome, particularly when based on an inbred, has no differences

from itself, the reference has no nonsynonymous variants to annotate and thus

appears free of deleterious variants. A more concerning type of reference bias is

that individuals that are genetically more similar to the reference genome will

have fewer differences from reference and thus fewer variants that annotate as

deleterious. This pattern is observed by Mezmouk and Ross-Ibarra (2014), who

find fewer deleterious variants in the stiff-stalk population of maize (to which the

reference genome B73 belongs) than in other elite maize populations. Along

similar lines, because gene models are derived from the reference genome,

more closely related lines with more similar coding portions of genes will appear

to have fewer disruptions of coding sequence. Finally reference bias can

contribute to under-calling of deleterious variants, either when different alleles fail

to properly align to the reference or if the reference is included in a phylogenetic

comparison. Variants that are detected as a difference from reference may then

be compared against the reference in an alignment testing for conservation at a

site. This conflates diversity within a species with the phylogenetic divergence

that is being tested in the alignment. In cases where the reference genome

contains the novel (or derived) variant relative to the state in related species, the

presence of the reference variant in the alignment will cause the site to appear

less constrained. This last problem is easily resolved by leaving the species

being tested out of the phylogenetic alignment used to annotate deleterious

variants.

Effect size

The evidence we present supports the cost of domestication hypothesis, namely

that domesticated lineages carry heavier genetic loads than their wild relatives

(Table 1). However, the distribution of fitness effects of variants is important with

the regard to the total load within an individual or population (Henn *et al*. 2015).

In other words, it is the cumulative effect of the variants carried by an individual

that make up its genetic load, not simply what proportion of those variants is

'deleterious'. As we describe above, current bioinformatic approaches that rely

on phylogenetic conservation may identify a number of false positives (driven by

new fitness optima under artificial selection) or false negatives (with specific

epistatic, dominance, or environmentally-dependent effects). Functional and

quantitative genetics approaches provide means of assessing the phenotypic

effect of genetic variants, but there are practical limitations that limit the quantity

of evaluations that can be conducted.

One potential issue is that the phenotypic effect of a genetic variant often

depends on genomic background, through both dominance (interaction between

alleles at the same locus) and epistasis (interaction among alleles at different

loci). Evaluating the effect of these variants is consequently a complex task,

requiring the creation and evaluation of multiple classes of recombinant

genomes. Similarly, the environment is an important consideration in thinking

about effect size. As we saw with the yield under drought *qDTY1.1* allele in rice

(*Identifying dangerous hitchhikers*, above; Vikram *et al*. 2015), the effect of a

genetic variant can depend on developmental or assessment environment.

These kinds of variants might be expected to be involved in local adaptation in

wild populations (e.g., Monroe *et al*. 2016), and so would not show up in a screen

based on phylogenetic conservation. Nevertheless, these variants may be

important in breeding programs that target general-purpose genotypes with, for

example, high mean and low variance yields. Identifying these alleles requires

assessment in the appropriate environment(s) and large assessment populations

(as the power for detecting genotype-phenotype correlations generally scales

with the number of genotypes assessed). Addressing effect size in the

appropriate context(s) therefore involves challenges of scale. Fortunately, new

types of mapping populations (e.g., MAGIC) and high-throughput phenotyping

platforms that can enable this work are increasingly available across

domesticated systems. Given these tools and the resultant data, we should soon

be able to parameterize genomic selection and similar models with putatively

deleterious variants and test their cumulative effects.

Heterosis

In systems with hybrid production, complementation of deleterious variants

between heterotic breeding pools may contribute substantially to heterosis (e.g.,

in maize; Hufford *et al*. 2012; Mezmouk and Ross-Ibarra 2014; Yang *et al*.

bioRxiv). If this is broadly true, hybrid production may be an interesting solution

to the cost of domestication, as long as deleterious variants are segregating

rather than fixed in domesticated species. Theory suggests that the deleterious

variants that contribute to heterosis between populations with low levels of gene

flow are likely to be of intermediate effect, and may not play a large role in

inbreeding depression (Whitlock, Ingvarsson, and Hatfield 2000). Since fitness in

these contexts is evaluated in hybrid individuals rather than inbred parents,

parental populations are likely to retain a higher proportion of slightly and

moderately deleterious variants than even selfing populations (Figure 2A). As

long as hybrid crosses are the primary mode of seed production, these alleles

may not be of high importance to breeders even though they contribute to

genetic load more broadly. Evaluating effect size in this context requires

assessing both parental and hybrid populations, and is therefore that much more

difficult. However, assuming that complementation of deleterious variants is a

substantial component of heterosis, quantifying these variants should improve

the ability of breeding programs to predict trait values in hybrid crosses. Further,

the question of whether genetic load limits genetic gains in hybrid production

systems remains open.

<u>CONCLUSIONS</u>

Research on the putative costs of domestication is still relatively new, and there

remain many open questions. However, our review of the literature suggests that

genetic loads are generally heavier in domesticated species compared to their

wild relatives. This pattern is likely driven by a number of processes that

collectively act to reduce the efficacy of selection relative to drift in domesticated

populations, resulting in increased frequency of deleterious variants linked to

selected loci and greater accumulation of deleterious variants genome-wide. We

encourage further research across domesticated species on these processes,

and recommend that researchers: (1) Sample domesticated and wild lineages

sufficiently to assess diversity within as well as between these groups and (2)

present deleterious variant data per genome and as proportional as well as

absolute values. We also strongly encourage researchers and breeders to think

about deleterious variation in context, both genomic and environmental.

FUNDING
This work was supported by the US National Science Foundation (award
numbers 1523752 to B.T.M.; and 1339393 to P.L.M.).

ACKNOWLEDGEMENTS
We thank Greg Baute and Kathryn Turner for comments on an earlier version of
this manuscript, and Brandon Gaut, Loren Rieseberg, and Greg Owens for
helpful discussion.

REFERENCES

Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, *et al.*
(2010). A method and server for predicting damaging missense mutations.
Nat Methods **7**: 248–249.

Agrawal AF, Whitlock MC (2012). Mutation load: the fitness of individuals in
populations where deleterious alleles are abundant. *Annu Rev Ecol Evol Syst*
43: 115–135.

Allard RW (1999). History of plant population genetics. *Annu Rev Genet* **33**: 1–
27.

Amaral AJ, Megens HJ, Crooijmans RPMA, Heuven HCM, Groenen MAM
(2008). Linkage disequilibrium decay and haplotype block structure in the pig.
Genetics **179**: 569–579.

Arunkumar R, Ness RW, Wright SI, Barrett SCH (2015). The evolution of selfing
is accompanied by reduced efficacy of selection and purging of deleterious
mutations. *Genetics* **199**: 817–829.

Asano K, Yamasaki M, Takuno S, Miura K, Katagiri S, Ito T, *et al.* (2011).
Artificial selection for a green revolution gene during japonica rice
domestication. *PNAS* **108**: 11034–11039.

Bachlava E, Tang S, Pizarro G, Schuppert GF, Brunick RK, Draeger D, *et al.*
(2009). Pleiotropy of the branching locus (B) masks linked and unlinked
quantitative trait loci affecting seed traits in sunflower. *Theor Appl Genet* **120**:
829–842.

Baer CF, Miyamoto MM, Denver DR (2007). Mutation rate variation in
multicellular eukaryotes: causes and consequences. *Nat Rev Genet* **8**: 619–
631.

Benbrook C (1999). *Evidence of the magnitude and consequences of the*
Roundup Ready soybean yield drag from university-based varietal trials in
1998. Ag BioTech InfoNet.
802

803   Bosse M, Megens H-J, Madsen O, Crooijmans RPMA, Ryder OA, Austerlitz F, *et*
804   *al.* (2015). Using genome-wide measures of coancestry to maintain diversity
805   and fitness in endangered and domestic pig populations. *Genome Research*
806   **25**: 970–981.
807

808   Bosse M, Megens H-J, Madsen O, Paudel Y, Frantz LAF, Schook LB, *et al.*
809   (2012). Regions of homozygosity in the porcine genome: consequence of
810   demography and the recombination landscape. *PLoS Genet* **8**: e1003100.
811

812   Brandenburg J-T, Mary-Huard T, Rigaill G, Hearne SJ, Corti H, Joets J, *et al.*
813   (2017). Independent introductions and admixtures have contributed to
814   adaptation of European maize and its American counterparts. *PLoS Genet*
815   **13**: e1006666.
816

817   Charlesworth B (2009). Fundamental concepts in genetics: Effective population
818   size and patterns of molecular evolution and variation. *Nat Rev Genet* **10**:
819   195–205.
820

821   Charlesworth D (2003). Effects of inbreeding on the genetic diversity of
822   populations. *Phil Trans Roy Soc B* **358**: 1051–1070.
823

824   Chen E, Huang X, Han B (2016). How can rice genetics benefit from rice-
825   domestication study? *National Science Review* **3**: 278–280.
826   Choi JY, Platts AE, Fuller DQ, Hsing Y-I, Wing RA, Purugganan MD (2017). The
827   rice paradox: Multiple origins but single domestication in Asian rice. *Mol Biol*
828   *Evol*: msx049.
829

830   Choi Y, Sims GE, Murphy S, Miller JR, Chan AP (2012). Predicting the
831   Functional Effect of Amino Acid Substitutions and Indels. *PLoS ONE* **7**:
832   e46688.
833

834   Chun S, Fay JC (2009). Identification of deleterious mutations within three
835   human genomes. *Genome Research* **19**: 1553–1561.
836

837   Clark RM, Linton E, Messing J, Doebley JF (2004). Pattern of diversity in the
838   genomic region near the maize domestication gene *tb1*. *PNAS* **101**: 700–707.
839

840   The Chimpanzee Sequencing and Analysis Consortium (2005). Initial sequence
841   of the chimpanzee genome and comparison with the human genome. *Nature*
842   **437**: 69–87.
843

844   The International Barley Genome Sequencing Consortium (2013). A physical,
845   genetic and functional sequence assembly of the barley genome. *Nature* **491**:
846   711–716.
847
848   Cooper GM, Stone EA, Asimenos G, Green ED, Batzoglou S, Sidow A (2005).
849   Distribution and intensity of constraint in mammalian genomic sequence.
850   *Genome Research* **15**: 901–913.
851
852   Corbi J, Debieu M, Rousselet A, Montalent P, Le Guilloux M, Manicacci D, *et al.*
853   (2010). Contrasted patterns of selection since maize domestication on
854   duplicated genes encoding a starch pathway enzyme. *Theor Appl Genet* **122**:
855   705–722.
856
857   Cruz F, Vila C, Webster MT (2008). The legacy of domestication: accumulation of
858   deleterious mutations in the dog genome. *Mol Biol Evol* **25**: 2331–2336.
859
860   Davydov EV, Goode DL, Sirota M, Cooper GM, Sidow A, Batzoglou S (2010).
861   Identifying a high fraction of the human genome to be under selective
862   constraint using GERP. *PLoS Comput Biol* **6**: e1001025.
863
864   Elmore RW, Roeth FW, Nelson LA (2001). Glyphosate-resistant soybean cultivar
865   yields compared with sister lines. *Agron J* **93**: 408–412.
866
867   Eyre-Walker A, Keightley PD (2007). The distribution of fitness effects of new
868   mutations. *Nat Rev Genet* **8**: 610–618.
869
870   Eyre-Walker A, Woolfit M, Phelps T (2006). The distribution of fitness effects of
871   new deleterious amino acid mutations in humans. *Genetics* **173**: 891–900.
872
873   Fay JC, Wyckoff GJ, Wu C-I (2001). Positive and Negative Selection on the
874   Human Genome. *Genetics* **158**: 1227–1234.
875
876   Felsenstein J (1974). The evolutionary advantage of recombination. *Genetics* **78**:
877   737–756.
878
879   Flowers JM, Molina J, Rubinstein S, Huang P, Schaal BA, Purugganan MD
880   (2012). Natural selection in gene-dense regions shapes the genomic pattern
881   of polymorphism in wild and domesticated rice. *Mol Biol Evol* **29**: 675–687.
882
883   Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, *et al.*
884   (2002). The structure of haplotype blocks in the human genome. *Science*
885   **296**: 2225–2229.
886
887   Gaut BS, Díez CM, Morrell PL (2015). Genomics and the contrasting dynamics of
888   annual and perennial domestication. *TIG* **31**: 709–719.
889

890 Georgelis N, Shaw JR, Hannah LC (2009). Phylogenetic analysis of ADP-
891   glucose pyrophosphorylase subunits reveals a role of subunit interfaces in
892   the allosteric properties of the enzyme. *Plant Physiol* **151**: 67–77.
893
894 Gheyas AA, Boschiero C, Eory L, Ralph H, Kuo R, Woolliams JA, *et al.* (2015).
895   Functional classification of 15 million SNPs detected from diverse chicken
896   populations. *DNA Research* **22**: 205–217.
897
898 Gillespie JH (1994). Substitution processes in molecular evolution. II.
899   Exchangeable models from population genetics. *Genetics* **138**: 943–952.
900
901 Gillespie JH (1999). The role of population size in molecular evolution. *Theor Pop*
902   *Biol* **55**: 145–156.
903
904 Giorgi D, Pandozy G, Farina A, Grosso V, Lucretti S, Gennaro A, *et al.* (2016).
905   First detailed karyo-morphological analysis and molecular cytological study of
906   leafy cardoon and globe artichoke, two multi-use Asteraceae crops. *CCG* **10**:
907   447–463.
908
909 Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, *et al.* (2011).
910   Phytozome: a comparative platform for green plant genomics. *Nucleic Acids*
911   *Res* **40**: D1178–D1186.
912
913 Gore MA, Chia JM, Elshire RJ, Sun Q, Ersoz ES, Hurwitz BL, *et al.* (2009). A
914   First-Generation Haplotype Map of Maize. *Science* **326**: 1115–1117.
915
916 Gray MM, Granka JM, Bustamante CD, Sutter NB, Boyko AR, Zhu L, *et al.*
917   (2009). Linkage Disequilibrium and Demographic History of Wild and
918   Domestic Canids. *Genetics* **181**: 1493–1505.
919
920 Gregory TR (2017). Animal Genome Size Database.
921 http://www.genomesize.com.
922
923 Grimm DG, Azencott C-A, Aicheler F, Gieraths U, MacArthur DG, Samocha KE,
924   *et al.* (2015). The Evaluation of Tools Used to Predict the Impact of Missense
925   Variants Is Hindered by Two Types of Circularity. *Human Mutation* **36**: 513–
926   523.
927
928 Groenen MAM, Wahlberg P, Foglio M, Cheng HH, Megens HJ, Crooijmans
929   RPMA, *et al.* (2008). A high-density SNP-based linkage map of the chicken
930   genome reveals sequence features correlated with recombination rate.
931   *Genome Research* **19**: 510–519.
932
933 Gross BL, Olsen KM (2010). Genetic perspectives on crop domestication. *Trends*
934   *Plant Sci* **15**: 529–537.
935

936 Gross BL, Zhao Z (2014). Archaeological and genetic insights into the origins of
937   domesticated rice. *PNAS* **111**: 6190–6197.
938
939 Günther T, Schmid KJ (2010). Deleterious amino acid polymorphisms in
940   *Arabidopsis thaliana* and rice. *Theor Appl Genet* **121**: 157–168.
941
942 Henn BM, Botigué LR, Bustamante CD, Clark AG, Gravel S (2015). Estimating
943   the mutation load in human genomes. *Nat Rev Genet* **16**: 333–343.
944
945 Hill WG, Robertson A (1966). The effect of linkage on limits to artificial selection.
946   *Genetics Research* **8**: 269–294.
947
948 Horak MJ, Rosenbaum EW, Kendrick DL, Sammons B, Phillips SL, Nickson TE,
949   *et al.* (2015). Plant characterization of Roundup Ready 2 Yield® soybean,
950   MON 89788, for use in ecological risk assessment. *Transgenic Res* **24**: 213–
951   225.
952
953 Huang X, Kurata N, Wei X, Wang Z-X, Wang A, Zhao Q, *et al.* (2012). A map of
954   rice genome variation reveals the origin of cultivated rice. *Nature* **490**: 497–
955   501.
956
957 Hufford MB, Xu X, van Heerwaarden J, Pyhäjärvi T, Chia J-M, Cartwright RA, *et*
958   *al.* (2012). Comparative population genomics of maize domestication and
959   improvement. *Nat Genet* **44**: 808–811.
960
961 Hughes AL (2005). Evidence for Abundant Slightly Deleterious Polymorphisms in
962   Bacterial Populations. *Genetics* **169**: 533–538.
963
964 Keightley PD, Eyre-Walker A (2007). Joint Inference of the Distribution of Fitness
965   Effects of Deleterious Mutations and Population Demography Based on
966   Nucleotide Polymorphism Frequencies. *Genetics* **177**: 2251–2261.
967
968 Keightley PD, Lynch M (2003). Toward a realistic model of mutations affecting
969   fitness. *Evolution* **57**: 683–685.
970
971 Kersey PJ, Allen JE, Armean I, Boddu S, Bolt BJ, Carvalho-Silva D, *et al.* (2016).
972   Ensembl Genomes 2016: more genomes, more complexity. *Nucleic Acids*
973   *Res* **44**: D574–D580.
974
975 Kim S, Plagnol V, Hu TT, Toomajian C, Clark RM, Ossowski S, *et al.* (2007).
976   Recombination and linkage disequilibrium in Arabidopsis thaliana. *Nat Genet*
977   **39**: 1151–1155.
978
Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J (2014). A
general framework for estimating the relative pathogenicity of human genetic
variants. *Nat Genet* **46**: 310–315.

Klopfstein S, Currat M, Excoffier L (2005). The Fate of Mutations Surfing on the

Wave of a Range Expansion. *Mol Biol Evol* **23**: 482–490.

Koenig D, Jiménez-Gómez JM, Kimura S, Fulop D, Chitwood DH, Headland LR,

et al. (2013). Comparative transcriptomics reveals patterns of selection in

domesticated and wild tomato. *PNAS* **110**: E2655–E2662.

Kono TJY, Fu F, Mohammadi M, Hoffman PJ, Liu C, Stupar RM, *et al.* (2016).

The Role of Deleterious Substitutions in Crop Genomes. *Mol Biol Evol* **33**:

2307–2317.

Kono TJY, Lei L, Shih C-H, Hoffman PJ, Morrell PL, Fay JC Comparative

genomics approaches accurately predict deleterious variants in plants.

bioRxiv.

Kovach MJ, Sweeney MT, McCouch SR (2007). New insights into the history of

rice domestication. *TIG* **23**: 578–587.

1000

1001　Kryazhimskiy S, Plotkin JB (2008). The Population Genetics of dN/dS. *PLoS*

1002　　　*Genet* **4**: e1000304.

1003

1004　Lam H-M, Xu X, Liu X, Chen W, Yang G, Wong F-L, *et al.* (2010). Resequencing

1005　　　of 31 wild and cultivated soybean genomes identifies patterns of genetic

1006　　　diversity and selection. *Nat Genet* **42**: 1053–1059.

1007

1008　Lau AN, Peng L, Goto H, Chemnick L, Ryder OA, Makova KD (2009). Horse

1009　　　Domestication and Conservation Genetics of Przewalski's Horse Inferred

1010　　　from Sex Chromosomal and Autosomal Sequences. *Mol Biol Evol* **26**: 199–

1011　　　208.

1012

1013　Lin T, Zhu G, Zhang J, Xu X, Yu Q, Zheng Z, *et al.* (2014). Genomic analyses

1014　　　provide insights into the history of tomato breeding. *Nat Genet* **46**: 1220–

1015　　　1226.

1016

1017　Lindblad-Toh K, Wade CM, Mikkelsen TS, Karlsson EK, Jaffe DB, Kamal M, *et*

1018　　　*al.* (2005). Genome sequence, comparative analysis and haplotype structure

1019　　　of the domestic dog. *Nature* **438**: 803–819.

1020

1021　Lippold S, Knapp M, Kuznetsova T, Leonard JA, Benecke N, Ludwig A, *et al.*

1022　　　(2011). Discovery of lost diversity of paternal horse lineages using ancient

1023　　　DNA. *Nat Commun* **2**: 450–6.

1024

1025　Liu A, Burke JM (2006). Patterns of Nucleotide Diversity in Wild and Cultivated

1026　　　Sunflower. *Genetics* **173**: 321–330.

1027

1028 Liu Q, Zhou Y, Morrell PL, Gaut BS (2017). Deleterious variants in Asian rice and
1029   the potential cost of domestication. *Mol Biol Evol*: msw296.
1030
1031 Lohmueller KE (2014). The Impact of Population Demography and Selection on
1032   the Genetic Architecture of Complex Traits. *PLoS Genet* **10**: e1004379.
1033
1034 Lu J, Tang T, Tang H, Huang J, Shi S, Wu C-I (2006). The accumulation of
1035   deleterious mutations in rice genomes: a hypothesis on the cost of
1036   domestication. *TIG* **22**: 126–131.
1037
1038 Mandel JR, Dechaine JM, Marek LF, Burke JM (2011). Genetic diversity and
1039   population structure in cultivated sunflower and a comparison to its wild
1040   progenitor, *Helianthus annuus* L. *Theor Appl Genet* **123**: 693–704.
1041
1042 Marsden CD, Ortega-Del Vecchyo D, O'Brien DP, Taylor JF, Ramirez O, Vilà C,
1043   *et al.* (2016). Bottlenecks and selective sweeps during domestication have
1044   increased deleterious genetic variation in dogs. *PNAS* **113**: 152–157.
1045
1046 Marth GT, Yu F, Indap AR, Garimella K, Gravel S, Leong WF, *et al.* (2011). The
1047   functional spectrum of low-frequency coding variation. *Genome Biology* **12**:
1048   R84.
1049
1050 Maynard Smith J, Haigh J (1974). The hitch-hiking effect of a favourable gene.
1051   *Genetical research* **23**: 1–13.
1052
1053 Megens H-J, Crooijmans RP, Bastiaansen JW, Kerstens HH, Coster A, Jalving
1054   R, *et al.* (2009). Comparison of linkage disequilibrium and haplotype diversity
1055   on macro- and microchromosomes in chicken. *BMC Genet* **10**: 86.
1056
1057 Mezmouk S, Ross-Ibarra J (2014). The Pattern and Distribution of Deleterious
1058   Mutations in Maize. *G3* **4**: 163–171.
1059
1060 Michaelson MJ, Price HJ, Ellison JR, Johnston JS (1991). Comparison of plant
1061   DNA contents determined by Feulgen microspectrophotometry and laser flow
1062   cytometry. *Am J Bot* **78**: 183–188.
1063
1064 Miller AJ, Gross BL (2011). From forest to field: Perennial fruit crop
1065   domestication. *Am J Bot* **98**: 1389–1414.
1066
1067 Miyata T, Miyazawa S, Yasunaga T (1979). Two types of amino acid
1068   substitutions in protein evolution. *J Molec Evol* **12**: 219–236.
1069
1070 Monroe JG, McGovern C, Lasky JR, Grogan K, Beck J, McKay JK (2016).
1071   Adaptation to warmer climates by parallel functional evolution of *CBF* genes
1072   in *Arabidopsis thaliana*. *Mol Ecol* **25**: 3632–3644.
1073

1074   Moray C, Lanfear R, Bromham L (2014). Domestication and the mitochondrial
1075   genome: comparing patterns and rates of molecular evolution in
1076   domesticated mammals and birds and their wild relatives. *Genome Biol Evol*
1077   **6**: 161–169.
1078
1079   Morrell PL, Buckler ES, Ross-Ibarra J (2011). Crop genomics: advances and
1080   applications. *Nat Rev Genet* **13**: 85–96.
1081
1082   Muir WM, Wong G, Zhang Y, Wang J, Groenen MAM, Crooijmans RPMA, *et al.*
1083   (2008). Genome-wide assessment of worldwide chicken SNP genetic
1084   diversity indicates significant absence of rare alleles in commercial breeds.
1085   *PNAS* **105**: 17312–17317.
1086
1087   Nabholz B, Sarah G, Sabot F, Ruiz M, Adam H, Nidelet S, *et al.* (2014).
1088   Transcriptome population genomics reveals severe bottleneck and
1089   domestication cost in the African rice (*Oryza glaberrima*). *Mol Ecol* **23**: 2210–
1090   2227.
1091
1092   Ng PC, Henikoff S (2003). SIFT: predicting amino acid changes that affect
1093   protein function. *Nucleic Acids Res* **31**: 3812–3814.
1094
1095   Nordborg M, Hu TT, Ishino Y, Jhaveri J, Toomajian C, Zheng H, *et al.* (2005).
1096   The Pattern of Polymorphism in *Arabidopsis thaliana*. *PLoS Biol* **3**: e196.
1097
1098   Ohta T (1972). Population size and rate of evolution. *J Molec Evol* **1**: 305–314.
1099
1100   Ohta T (1992). The nearly neutral theory of molecular evolution. *Annu Rev Ecol*
1101   *Syst* **23**.
1102
1103   Olsen KM (2006). Selection Under Domestication: Evidence for a Sweep in the
1104   Rice *Waxy* Genomic Region. *Genetics* **173**: 975–983.
1105
1106   Otto SP, Barton NH (2001). Selection for recombination in small populations.
1107   *Evolution* **55**: 1921–1931.
1108
1109   Otto SP, Whitlock MC (1997). The probability of fixation in populations of
1110   changing size. *Genetics* **146**: 723–733.
1111
1112   Papa R, Bellucci E, Rossi M, Leonardi S, Rau D, Gepts P, *et al.* (2007). Tagging
1113   the Signatures of Domestication in Common Bean (*Phaseolus vulgaris*) by
1114   Means of Pooled DNA Samples. *Ann Bot–London* **100**: 1039–1051.
1115
1116   Parker HG, vonHoldt BM, Quignon P, Margulies EH, Shao S, Mosher DS, *et al.*
1117   (2009). An expressed *Fgf4* retrogene is associated with breed-defining
1118   chondrodysplasia in domestic dogs. *Science* **325**: 995–998.
1119

1120   Peischl S, Dupanloup I, Kirkpatrick M, Excoffier L (2013). On the accumulation of
1121   deleterious mutations during range expansions. *Mol Ecol* **22**: 5972–5982.
1122

1123   Renaut S, Rieseberg LH (2015). The Accumulation of Deleterious Mutations as a
1124   Consequence of Domestication and Improvement in Sunflowers and Other
1125   Compositae Crops. *Mol Biol Evol* **32**: 2273–2283.
1126

1127   Rice A, Glick L, Abadi S, Einhorn M (2015). The Chromosome Counts Database
1128   (CCDB)–a community resource of plant chromosome numbers. *New Phytol*
1129   **206**: 19–26.
1130

1131   Rodgers-Melnick E, Bradbury PJ, Elshire RJ, Glaubitz JC, Acharya CB, Mitchell
1132   SE, *et al.* (2015). Recombination in diverse maize is stable, predictable, and
1133   associated with genetic load. *PNAS* **112**: 3823–3828.
1134

1135   Ross-Ibarra J (2004). The evolution of recombination under domestication: a test
1136   of two hypotheses. *Am Nat* **163**: 105–112.
1137

1138   Ross-Ibarra J, Tenaillon M, Gaut BS (2009). Historical Divergence and Gene
1139   Flow in the Genus *Zea*. *Genetics* **181**: 1399–1413.
1140

1141   Rubin C-J, Zody MC, Eriksson J, Meadows JRS, Sherwood E, Webster MT, *et al.*
1142   (2010). Whole-genome resequencing reveals loci under selection during
1143   chicken domestication. *Nature* **464**: 587–591.
1144

1145   Scaglione D, Reyes-Chin-Wo S, Acquadro A, Froenicke L, Portis E, Beitel C, *et*
1146   *al.* (2015). The genome sequence of the outbreeding globe artichoke
1147   constructed. *Sci Rep*: 1–17.
1148

1149   Scally A, Dutheil JY, Hillier LW, Jordan GE, Goodhead I, Herrero J, *et al.* (2012).
1150   Insights into hominid evolution from the gorilla genome sequence. *Nature*
1151   **483**: 169–175.
1152

1153   Schmutz J, McClean PE, Mamidi S, Wu GA, Cannon SB, Grimwood J, *et al.*
1154   (2014). A reference genome for common bean and genome-wide analysis of
1155   dual domestications. *Nat Genet* **46**: 707–713.
1156

1157   Schneider CA, Rasband WS, Eliceiri KW (2012). NIH Image to ImageJ: 25 years
1158   of image analysis. *Nat Methods* **9**: 671–675.
1159

1160   Schubert M, Jónsson H, Chang D, Sarkissian Der C, Ermini L, Ginolhac A, *et al.*
1161   (2014). Prehistoric genomes reveal the genetic foundation and cost of horse
1162   domestication. *PNAS* **111**: E5661–E5669.
1163

1164   Semon M, Nielsen R, Jones MP, McCouch SR (2005). The Population Structure
1165   of African Cultivated Rice *Oryza glaberrima* (Steud.): Evidence for Elevated

1166   Levels of Linkage Disequilibrium Caused by Admixture with *O. sativa* and
1167   Ecological Adaptation. *Genetics* **169**: 1639–1647.
1168
1169 Shi T, Dimitrov I, Zhang Y, Tax FE, Yi J, Gou X, *et al.* (2015). Accelerated rates
1170   of protein evolution in barley grain and pistil biased genes might be legacy of
1171   domestication. *Plant Molecular Biology* **89**: 253–261.
1172
1173 Simons YB, Turchin MC, Pritchard JK, Sella G (2014). The deleterious mutation
1174   load is insensitive to recent population history. *Nat Genet* **46**: 220–224.
1175
1176 Stone EA, Sidow A (2005). Physicochemical constraint violation by missense
1177   substitutions mediates impairment of protein function and disease severity.
1178   *Genome Research* **15**: 978–986.
1179
1180 Takeda S, Matsuoka M (2008). Genetic approaches to crop improvement:
1181   responding to environmental and population changes. *Nat Rev Genet* **9**: 444–
1182   457.
1183
1184 Tian F, Stevens NM, Buckler ES (2009). Tracking footprints of maize
1185   domestication and evidence for a massive selective sweep on chromosome
1186   10. *PNAS* **106**: 9979–9986.
1187
1188 Traspov A, Deng W, Kostyunina O, Ji J, Shatokhin K, Lugovoy S, *et al.* (2016).
1189   Population structure and genome characterization of local pig breeds in
1190   Russia, Belorussia, Kazakhstan and Ukraine. *Genetics Selection Evolution*
1191   **48**: 1–9.
1192
1193 Travis JMJ, Munkemuller T, Burton OJ, Best A, Dytham C, Johst K (2007).
1194   Deleterious Mutations Can Surf to High Densities on the Wave Front of an
1195   Expanding Population. *Mol Biol Evol* **24**: 2334–2343.
1196
1197 Vikram P, Swamy BPM, Dixit S, Singh R, Singh BP, Miro B, *et al.* (2015).
1198   Drought susceptibility of modern rice varieties: an effect of linkage of drought
1199   tolerance with undesirable traits. *Sci Rep* **5**.
1200
1201 Wade CM, Giulotto E, Sigurdsson S, Zoli M (2009). Genome Sequence,
1202   Comparative Analysis, and Population Genetics of the Domestic Horse.
1203   *Science* **326**: 865–867.
1204
1205 Walsh B (2007). Using molecular markers for detecting domestication,
1206   improvement, and adaptation genes. *Euphytica* **161**: 1–17.
1207
1208 Wang G-D, Xie H-B, Peng M-S, Irwin D, Zhang Y-P (2014). Domestication
1209   Genomics: Evidence from Animals. *Annu Rev Anim Biosci* **2**: 65–84.
1210

1211   Wang H, Studer AJ, Zhao Q, Meeley R (2015). Evidence that the origin of naked
1212   kernels during maize domestication was caused by a single amino acid
1213   substitution in *tga1*. *Genetics* **200**: 965–974.
1214
1215   Whitlock MC (2000). Fixation of new alleles and the extinction of small
1216   populations: drift load, beneficial alleles, and sexual selection. *Evolution* **54**:
1217   1855–1861.
1218
1219   Whitlock MC, Ingvarsson PK, Hatfield T (2000). Local drift load and the heterosis
1220   of interconnected populations. *Heredity* **84**: 452–457.
1221
1222   Whitt SR, Wilson LM, Tenaillon MI, Gaut BS, Buckler ES (2002). Genetic
1223   diversity and selection in the maize starch pathway. *PNAS* **99**: 12959–12962.
1224
1225   Wilfert L, Gadau J, Schmid-Hempel P (2007). Variation in genomic recombination
1226   rates among animal taxa and the case of social insects. *Heredity* **98**: 189–
1227   197.
1228
1229   Wolf JBW, Kunstner A, Nam K, Jakobsson M, Ellegren H (2009). Nonlinear
1230   Dynamics of Nonsynonymous (dN) and Synonymous (dS) Substitution Rates
1231   Affects Inference of Selection. *Genome Biol Evol* **1**: 308–319.
1232
1233   Woolfit M (2009). Effective population size and the rate and pattern of nucleotide
1234   substitutions. *Biology Letters* **5**: 417–420.
1235
1236   Wright SI, Bi IV, Schroeder SG, Yamasaki M, Doebley JF, McMullen MD, *et al.*
1237   (2005). The Effects of Artificial Selection on the Maize Genome. *Science* **308**:
1238   1310–1314.
1239
1240   Xu Z, Hennessy DA, Sardana K, Moschini G (2013). The Realized Yield Effect of
1241   Genetically Engineered Crops: U.S. Maize and Soybean. *Crop Sci* **53**: 735–
1242   745.
1243
1244   Yamasaki M, Wright SI, McMullen MD (2007). Genomic Screening for Artificial
1245   Selection during Domestication and Improvement in Maize. *Ann Bot–London*
1246   **100**: 967–973.
1247
1248   Yang J, Mezmouk S, Baumgarten A, Buckler ES, Guill KE, McMullen MD, *et al.*
1249   Incomplete dominance of deleterious alleles contributes substantially to trait
1250   variation and heterosis in maize. *bioRxiv*.
1251
1252   Zhou Z, Jiang Y, Wang Z, Gou Z, Lyu J, Li W, *et al.* (2015). Resequencing 302
1253   wild and cultivated accessions identifies genes related to domestication and
1254   improvement in soybean. *Nat Biotechnol* **33**: 408–414.
1255

1256 Zonneveld BJM, Leitch IJ, Bennett MD (2005). First Nuclear DNA Amounts in
1257   more than 300 Angiosperms. *Ann Bot–London* **96**: 229–244.

1258
1259 TABLES AND FIGURES
1260
Table 1. Evidence for genetic load across domesticated plants and animals.
Arabidopsis thaliana and *Homo sapiens* included for comparison. Plant 1C
genome sizes from the RBG Kew database (http://data.kew.org/cvalues/), except
tomato (Michaelson *et al*. 1991) and *Cynara* spp. (Giorgi *et al*. 2016). Plant
chromosome counts from the Chromosome Counts Database
(http://ccdb.tau.ac.il/). Animal 1C genome sizes and chromosome counts from
the Animal Genome Size Database (http://www.genomesize.com/, mean value
when multiple records were available). Gene numbers are high-confidence (if
available) estimates from the vertebrate and plant Ensembl databases
(http://uswest.ensembl.org/; http://plants.ensembl.org/), except common bean
(Schmutz et al. 2013), sunflower (Compositate Genome Project, unpublished),
and *Cynara* spp. (Scaglione et al. 2016). LD N50 is the approximate distance
over which LD decays to half of maximum value. Loss of genetic diversity is
calculated as 1 – (ratio of $p_{domesticated}$ to p_{wild}), or q if p not available.
1275
Figure 1. Processes of domestication and improvement. (A) Typical changes in
effective population size through domestication and improvement. Stars indicate
genetic bottlenecks. These dynamics can be reconstructed by examining
patterns of genetic diversity in contemporary wild relative, domesticated non-
commercial, and improved populations. (B) Effects of artificial selection (targeting
the blue triangle variant) and linkage disequilibrium on deleterious (red squares)
and neutral variants (grey circles, shades represent different alleles). In the
ancestral wild population, four deleterious alleles are at relatively low frequency
(mean = 0.10) and heterozygosity is high (H_O = 0.51). After domestication, the
selected blue triangle and linked variants increase in frequency (three remaining
deleterious alleles, mean frequency = 0.46), heterozygosity decreases (H_O =
0.35), and allelic diversity is lost at two sites. Recombination may change
haplotypes, especially at sites less closely linked to the selected allele (Xs). After
improvement, further selection for the blue triangle allele has: lowered
heterozygosity (H_O = 0.08), increased deleterious variant frequency (two
remaining deleterious alleles, mean = 0.55), and lost allelic diversity at six
additional sites.
1293
Figure 2. Effects of inbreeding on mutations and fitness. (A) Theoretical density
plot of fitness effects for segregating variants. In outcrossing, non-inbred
populations (solid navy line), more variants with highly deleterious, recessive
effects persist that are rapidly exposed to selection and purged in inbred
populations (dotted red line), shifting the left side of the distribution towards more
neutral effects. At the same time, the reduced effective population size created
through inbreeding causes on average higher loss of slightly advantageous
mutations and retention of slightly deleterious mutations, shifting the right side of

the distribution towards more deleterious effects. (B) Accumulation of
nonsynonymous mutations is accelerated in selfing individuals (dotted red line)
relative to outcrossing individuals (solid navy line). Synonymous mutations
accumulate at similar rates in both populations. Adapted from simulation results
in Arunkumar *et al*. 2015. (C) As a consequence of (B), mean individual fitness
drops in selfing relative to outcrossing populations. Adapted from Arunkumar *et*
al. 2015.

Species	Primary reproduction	Genome size (1C, pg)	Chromosome number (N)	Protein coding genes	Non-coding genes	LD N50	LD N50 in wild	Loss of genetic diversity	dN/dS domesticated to dN/dS wild (or Ka/Ks)*	GERP score load increase relative to wild	Genome-wide deleterious variant number	Deleterious variant proportion (of nonsynonymous variants)	Deleterious variants in wild, # (proportion)	Method(s) for deleterious variant detection	Studies
japonica rice (Oryza sativa var japonica)	selfing	0.50	12	35,679	55,401	167 kb	20 kb	0.67	1.70	NA	~6295*	0.19*	6099 (0.15)*	SIFT, PROVEAN	Lu et al. 2006; Huang et al. 2012; Liu et al. 2017
indica rice (Oryza sativa var indica)	selfing	0.50	12	40,745	45,577	123 kb	20 kb	0.25	1.75	NA	~6351*	0.18*	6099 (0.15)*	SIFT, PROVEAN	Lu et al. 2006; Huang et al. 2012; Liu et al. 2017
African rice (Oryza glaberrima)	selfing	0.83	12	33,164	38,882	~25 Mb*	NA	0.60	1.58	NA	NA	NA	NA		Semon et al. 2005; Nabholz et al. 2014
maize (Zea mays)	outcrossing	2.73	10	39,498	4,976	<1 Mb	NA	0.17	NA	NA	~1025–4056*	0.04–0.16*	NA	SIFT, MAPP, Join of these	Gore et al. 2009; Hufford et al. 2012; Mezmouk and Ross-Ibarra 2014
barley (Hordeum vulgare)	selfing	5.55	7	24,287	1,512	~2.2 Mb	~110 kb	0.13 (domestication) and 0.27 (improvement)	2.43–4.60	NA	1,006–3,400	0.06–0.19	NA	SIFT, PolyPhen2, LRT, Intersect of these	The International Barley Genome Sequencing Consortium 2012; Morrell et al. 2014; Shi et al. 2015; Kono et al. 2016
soybean (Glycine max)	selfing	1.13	20	54,174	NA	83 kb (domesticated) and 123 kb (improved)	27 kb	0.52 (domestication) and 0.64 (improvement)	1.01**	NA	784–3,881	0.03–0.13	NA	SIFT, PolyPhen2, LRT, Intersect of these	Lam et al. 2010; Zhou et al. 2015; Kono et al. 2016
common bean (Phaseolus vulgaris, Mesoamerican domestication)	selfing	0.60	11	27,197	NA	NA	NA	0.20	NA	NA	NA	NA	NA		Schmutz et al. 2013
common bean (Phaseolus vulgaris, Andean domestication)	selfing	0.60	11	27,197	NA	NA	NA	-0.50	NA	NA	NA	NA	NA		Schmutz et al. 2013
tomato (Solanum lycopersicum)	selfing	1.00	12	33,837	4,849	257 kb (domesticated) and 866 kb (improved)	9 kb	0.46 (domestication) and 0.77 (improvement)	1.37***	NA	NA	NA	NA		Koenig et al. 2013; Lin et al. 2014
sunflower (Helianthus annuus)	outcrossing	2.43	17	52,232	NA	~1 kb	200 bp	0.33	NA	NA	5,626–20,273*	0.14–0.17*	7,882–24,479 (0.12–0.14)*	PROVEAN	Liu and Burke 2006; Mandel et al. 2011; Renaut and Rieseberg 2015; L. Rieseberg unpublished
globe artichoke (Cynara cardunculus var. scolymus)	clonal propagation, outcrossing	1.20	17	26,889	NA	NA	NA	NA	NA	NA	1,537–1,889*	0.19–0.20*	1,465 (0.17)*	PROVEAN	Renaut and Rieseberg 2015
cardoon (Cynara cardunculus var. altilis)	outcrossing	1.10	17	26,889	NA	NA	NA	NA	NA	NA	1,239–1,458*	0.22–0.23*	1,900 (0.19)*	PROVEAN	Renaut and Rieseberg 2015
thale cress (Arabidopsis thaliana)	selfing	0.30	5	27,655	1,398	~3-4 kb	NA	NA	NA	NA	NA	NA	NA	SIFT, MAPP	Kim et al. 2007; Günther and Schmid 2010
chicken (Gallus gallus domesticus)	outcrossing	1.25	39	18,346	6,492	~50–250 kb	< 50 kb	~0.50 (from domesticated to 'commercial' breeds*)	NA	NA	35,889–88,655	0.20–0.49	NA	SIFT, PROVEAN, Join and Intersect of these	Muir et al. 2008; Megens et al. 2009; Gheyas et al. 2015
dog (Canis familiaris)	outcrossing	3.12	39	19,856	11,898	~100 kb	< 10 kb	0.05 (domestication) / 0.35 (breed formation)	1.54	2.1%	~5,115*	~0.14	~4,990 (~0.138)*	Miyata distance; GERP	Lindblad-Tor et al. 2005; Cruz, Vila & Webster 2008; Gray et al. 2009; Marsden et al. 2016
horse (Equus caballus)	outcrossing	3.22	32	20,449	2,142	~200 kb	NA	0.40**	NA	1.1–11%	NA	NA	NA	GERP	Lau et al. 2009; Wade et al. 2009; Schubert et al. 2014
pig (Sus scrofa)	outcrossing	3.17	19	21,630	3,124	5 kb–1 Mb	< 50 kb	0.19	NA	NA	656*	NA	NA	SIFT*	Amaral et al. 2008; Bosse et al. 2012; Bosse et al. 2015
human (Homo sapiens)	outcrossing	3.50	23	20,441	22,219	22-44 kb	NA	NA	1.03	NA	796–838*	0.08–0.15*	NA	LRT	Gabriel et al. 2002; Chun and Fay 2009; Scally et al. 2012

* likely due to population structure in sampled accessions (Semon et al. 2005)

*estimated as portion of hypothetical ancestral allele frequency distribution lost

* or versus sister species for human

* per genome

* per genome

*significantly lower in wild

** data are ratio of nonsynonymous to synonymous SNP counts rather than substitutions

*** autosomal nucleotide diversity only

*** comparison of domesticated lineage versus whole tree rates (five wild lineages and one domesticated)