Multiplex genome editing by natural transformation (MuGENT) for synthetic biology in *Vibrio natriegens*

Triana N. Dalia[1], Chelsea A. Hayes[1], Sergey Stolyar[2], Christopher J. Marx[2], James B. McKinlay[1], and Ankur B. Dalia[1,*]

[1]Department of Biology, Indiana University, Bloomington, IN 47401. [2]Department of Biological Sciences, University of Idaho, Moscow, ID 83844.

*Author for correspondence: Ankur B. Dalia, ankdalia@indiana.edu

ABSTRACT

Vibrio natriegens has recently emerged as an alternative to *Escherichia coli* for molecular

biology and biotechnology, but low-efficiency genetic tools hamper its development. Here,

we uncover how to induce natural competence in *V. natriegens* and describe methods for

multiplex genome editing by natural transformation (MuGENT). MuGENT promotes

integration of multiple genome edits at high-efficiency on unprecedented timescales. Also,

this method allows for generating highly complex mutant populations, which can be

exploited for metabolic engineering efforts. As a proof-of-concept, we attempted to

enhance production of the value added chemical poly-β-hydroxybutyrate (PHB) in *V.*

natriegens by targeting the expression of nine genes involved in PHB biosynthesis via

MuGENT. Within 1 week, we isolated edited strains that produced ~100 times more PHB

than the parent isolate and ~3.3 times more than a rationally designed strain. Thus, the

methods described here should extend the utility of this species for diverse academic and

industrial applications.

V. natriegens is the fastest growing organism known, with a doubling time of <10 min[1, 2].

With broad metabolic capabilities, lack of pathogenicity, and its rapid growth rate, it is an

attractive alternative to *E. coli* for diverse molecular biology and biotechnology

applications[3, 4]. Methods for classical genetic techniques have been developed for *V.*

natriegens, but these are relatively laborious, require multiple steps, and must be used

sequentially to generate multiple genome edits[3, 4]. The challenges of these techniques

contrast with the ease of genetics in *Vibrio* species that are naturally transformable.

Competent *Vibrio* species can take up DNA from the environment and integrate it into their

genome by homologous recombination; processes known as natural competence and

natural transformation, respectively[5-8]. The inducing cue for natural transformation in

competent *Vibrio* species is growth on the chitinous shells of crustacean zooplankton,

which are commonly found in the aquatic environment where these microbes reside[5].

Chitin induces the expression of the competence regulator TfoX[9, 10]. In fact, overexpression

of TfoX obviates the need for chitin induction, allowing competent *Vibrio* species to be

naturally transformed in rich media[5, 9].

As no reports of natural transformation existed for *V. natriegens*, we first sought to

establish whether this was possible. Unlike naturally competent *V. cholerae*, incubation on

chitin did not lead to detectable transformation in *V. natriegens* (data not shown). However,

ectopic expression of TfoX (either the endogenous *tfoX* gene or one from *Vibrio cholerae*)

on an IPTG-inducible plasmid (pMMB) supported high rates of natural transformation (**Fig.**

1a). This was tested using a linear PCR product that replaces the gene encoding the DNA

endonuclease Dns with an antibiotic resistance (Ab^R) marker. The *dns* locus was used as a

target for transformation assays throughout this manuscript because loss of this gene does

not impact growth or viability in rich medium. Under optimal conditions ~1-10% of the

population had integrated the transforming DNA (tDNA), which matches the highest rates

of transformation observed among competent species[11] (**Fig. 1a-c**). Natural transformation

of *V. natriegens* required very little transforming DNA (tDNA) (highly efficient with even 1

ng / 10^8 CFU) and was dependent on the length of homologous sequence surrounding the

mutation (**Fig. 1b and c**). This method could also be used to introduce point mutations into

V. natriegens (tested with tDNA containing an *rpsL* K43R Sm[R] allele); however, this activity

was partially suppressed by the mismatch repair system (**Fig. 1d**).

Having demonstrated *V. natriegens* is naturally competent, we sought to determine if we

could use natural transformation to perform scarless multiplex genome editing by natural

transformation (MuGENT)[12]. MuGENT operates under the premise that under competence

inducing conditions, only a subpopulation of cells is transformable. Those cells that can be

transformed, however, have the capacity to take up and integrate multiple tDNAs[12, 13]. Thus,

during MuGENT, cells are incubated with two types of linear tDNA; (1) a selected product

that introduces an antibiotic resistance marker into the genome and (2) unselected

products that introduce scarless genome edits of interest at one or more loci.

We first tested the ability of MuGENT to introduce a single unmarked genome edit (also

known as cotransformation). To facilitate measurement of cotransformation, we noted this

species forms opaque colonies on agar plates (**Fig. 2a**), which could be due to the

production of a capsular polysaccharide. Consistent with this, inactivating a homolog of the

essential capsule biosynthesis gene *wbfF*[14] resulted in the formation of transparent

colonies on agar plates and loss of expression of a high molecular weight polysaccharide

(**Fig. 2a and 2b**). Thus, to test cotransformation we used an unselected product to replace

~500 bp of the 5' end of the *wbfF* gene with a premature stop codon and scored

cotransformation via colony morphology (opaque vs. transparent) on agar plates (**Fig. 3a**).

We found that cotransformation was remarkably efficient in *V. natriegens* (up to ~80%),

even with low amounts (~25-50 ng / 10^8 CFU) of the unselected product (**Fig. 3b**). Also,

cotransformation with 1 kb flanks on the unselected product was possible, but at ~6-fold

lower frequencies than with 3 kb flanks (**Fig. 3c**).

We next tested the full multiplex genome editing capacity of MuGENT to simultaneously

cotransform multiple scarless genome edits into the genome in a single step[12, 15]. Since

there is no selection for integration of the unselected genome edits *in cis* during MuGENT,

output populations are highly heterogeneous and individual mutants contain any number

and combination of the multiplexed genome edits. Also, this process can be carried out in

multiple iterative cycles to further increase the complexity of genome edits in the

population (**Fig. 3d**)[12].

As an initial test of multiplex genome editing, we targeted 5 genes whose mutagenesis was

considered unlikely to affect viability or growth in LB. These targets included four

carbohydrate transporters (specific for mannitol, fructose, sucrose, and trehalose – all of

which are absent in LB) and the *dns* gene. All genes were targeted for inactivation by

replacing ~500 bp of the 5' end of each gene with a premature stop codon. Integration of

genome edits was determined by multiplex allele-specific colony PCR (MASC-PCR)[16] (**Fig.**

3e). Following one cycle of MuGENT, we found that ~70% of the population contained at

least 1 genome edit, with ~25% of the population containing 3-4 genome edits (**Fig. 3f**). A

quadruple mutant from this experiment was isolated and whole genome sequencing of this

strain did not reveal any off-target mutations. Thus, MuGENT rapidly generated *V.*

natriegens strains with multiple large (0.5 kb) scarless genome edits at high-efficiency

without off-target effects, and can be used to make highly complex mutant populations.

As a second demonstration of multiplex genome editing, we demonstrated its utility in

metabolic engineering by attempting to rapidly enhance production of a value-added

chemical in *V. natriegens*. This species naturally accumulates low levels of the bioplastic

precursor poly-β-hydroxybutyrate (PHB) as a storage polymer[17]. PHB is derived from the

condensation and subsequent NADPH-dependent reduction of acetyl-CoA precursors[18].

Thus, for our targets, we tuned the expression (swap out P_{native} for IPTG-inducible P_{tac}) or

inactivated genes that we hypothesized would affect NADPH and/or acetyl-CoA availability.

The targets for promoter swaps were the PHB synthesis operon (*phaBAC*), NAD kinase

(*nadK*), and two transhydrogenases (*pntAB* and *udhA*), while targets for inactivation were

phosphoglucose isomerase (*pgi*), citrate synthase (*gltA*), phosphotransacetylase (*pta*),

isocitrate lyase (*aceA*), and lactate dehydrogenase (*ldhA*) (**Fig. 4a**). Thus, there were 512

possible combinations for these 9 genome edits. We performed multiple cycles of MuGENT

to introduce these genome edits into a competent population of *V. natriegens*. At each cycle,

the selected product was designed to swap the Ab^R marker at the *dns* locus to maintain

coselection at each step. Following four cycles of MuGENT, which took just 5 days to

perform, ~50% of the population had 3 or more genome edits and ~10% contained 5+

genome edits (**Fig. 4b**). To select mutants with increased PHB production, we then plated

this output population onto media containing Nile red, which stains PHB granules[19]. Nile

red fluorescence on these plates was highly heterogeneous, suggesting that some

genotypes produced more PHB than the parent isolate (**Fig. 4c**). A number of highly

fluorescent colonies were picked and the genotypes determined by MASC-PCR. Also, PHB in

these select strains was directly measured by HPLC. Cumulatively, these analyses rapidly

revealed genotypes that produced ~100-fold more PHB than the parent and ~3.3-fold

more than a strain with just the P_{tac}-*phaBAC* mutation (**Fig. 4d**). Overexpression of the

phaBAC locus is a commonly used rational approach for enhancing PHB production [18, 20].

Thus, this result demonstrates that MuGENT can allow for rapid isolation of genotypes

associated with enhanced phenotypes (e.g. enhanced PHB production) compared to

rationally engineered strains (e.g. a P_{tac}-*phaBAC* mutant) without prior knowledge of

effective combinations of individual mutations.

While many methods for multiplex genome editing in bacterial systems have been

described[21], many of these are limited to small changes such as SNPs. MuGENT, on the

other hand, can efficiently swap, insert, or remove whole promoters or coding sequences as

demonstrated above. Furthermore, one of the major limitations to other multiplex genome

editing methods is that mutagenesis must be performed in strains lacking DNA repair

pathways to allow for high-efficiency integration of genome edits, which results in a large

number of off-target mutations[16, 21]. MuGENT in *V. natriegens* is performed in DNA repair

sufficient backgrounds, thus, little to no off target mutations are introduced during the

procedure as indicated above. Also, unlike other multiplex editing approaches, MuGENT

requires no specialized equipment and, thus, has the potential to make multiplex genome

editing commonplace.

In conclusion, this study demonstrates that MuGENT is a rapid, efficient, and simple tool for

engineering the *V. natriegens* genome. This microbe is already being developed as an

alternative to *E. coli*, and we believe that the ease and speed of MuGENT will extend the use

of *V. natriegens* as a novel chassis for diverse molecular biology and biotechnology

applications.

METHODS

Bacterial strains and culture conditions

The parent *V. natriegens* strain used throughout this study was a spontaneous rifampicin-

resistant derivative of ATCC 14048[2]. For a list of all strains used / generated in this study,

see **Table S1**. Strains were routinely grown in LB+v2 salts (LBv2)[3], which is LB Miller broth

(BD) supplemented with 200 mM NaCl, 23.14 mM $MgCl_2$, and 4.2 mM KCl. LBv2 was

supplemented with 100 µM IPTG, 50 µg/mL kanamycin (Kan), 200 µg/mL spectinomycin

(Spec), 100 µg/mL rifampicin (Rif), 100 µg/mL streptomycin (Sm), or 100 µg/mL

carbenicillin (Carb) as appropriate.

Generation of mutant strains and constructs

Mutant constructs were generated by splicing-by-overlap extension (SOE) PCR exactly as

previously described[22]. Briefly, for three-piece mutant constructs (i.e. for constructs where

a gene of interest is replaced with an Ab^R cassette or where the native promoter is swapped

for a P_{tac} promoter) segments were designated UP, MIDDLE, and DOWN and correspond to:

(1) UP = the upstream region of homology amplified with F1 and R1 primers, (2) DOWN =

the downstream region of homology amplified with F2 and R2 primers, and (3) MIDDLE =

the Ab^R marker or promoter swap fragment. For two-piece mutant constructs (i.e. for

constructs where ~501 bp of the 5' end of a gene is replaced with a stop codon), the

mutation of interest is incorporated into the R1 and F2 primers used to amplify the

upstream and downstream regions of homology, respectively. Gel purified segments were

then mixed in equal ratios and used as template for a SOE PCR reaction with the F1 and R2

primers. All mutant constructs were made using Phusion polymerase. These were

introduced into the *V. natriegens* genome via natural transformation as described below.

All primers used to generate mutant constructs are listed in **Table S2**.

Natural transformation / MuGENT assays

Strains harboring pMMB-*tfoX* (Vn *tfoX* or Vc *tfoX*) were induced to competence by growing

overnight (12-18 hours) in LBv2+100 µg/mL carbenicillin+100 µM IPTG in a rollerdrum at

30°C. Then, ~10^8 CFUs of this overnight culture (~3.5 µL) were diluted directly into 350 µL

of instant ocean medium (28 g/L; Aquarium Systems Inc.) supplemented with 100 µM

IPTG. Transforming DNA (tDNA) was then added as indicated, and reactions were

incubated statically at 30°C for 5 hours. Next, 1 mL of LBv2 was added and reactions were

outgrown at 30°C with shaking (250 rpm) for ~1-2 hrs. Then, reactions were plated for

quantitative culture onto media to select for integration of tDNA (i.e. LB+drug =

transformants) and onto nonselective media (i.e. plain LB = total viable counts).

Transformation efficiency is shown as: transformants / total viable counts.

For MuGENT, transformation assays were conducted exactly as described above. Unless

otherwise specified, ~50 ng of the selected product was incubated with cells along with

~200 ng of each unselected product. After outgrowth, 1/10th of the reaction was removed

and plated for MASC-PCR analysis (described below). If multiple cycles of MuGENT were

performed, the rest of the reaction was grown overnight in LBv2 supplemented with 100

µM IPTG, 100µg/mL carbenicillin (to maintain pMMB-*tfoX*), and the antibiotic to select for

integration of the selected product. The following day, the population was then subjected to

another round of MuGENT as described above using a selected product containing a

different Ab^R marker to maintain coselection at each cycle.

Integration of genome edits was detected via MASC-PCR exactly as previously described[12,

16]. Briefly, colonies were boiled in 50 µL of sterile water, vortexed, and then 2 µL were used

as template in a 25 µL PCR reaction. PCR was conducted with Taq polymerase (SydLabs)

using a modified 5X Taq buffer: 200 mM Tris pH 8.8, 100 mM KCl, 100 mM $(NH_4)_2SO_4$, 30

mM $MgSO_4$, and 1% Triton X-100. The total primer used in each MASC-PCR reaction

(regardless of the number of multiplexed products being detected) was 1200 nM (i.e. for

detection of 4 multiplexed genome edits, 300 nM of each genome edit-specific primer pair

was used). The cycling conditions used were: 95°C 3 min; 26 × [95°C 40s, 58°C 30s, 72°C 3

min]; 72°C 3 min; 12°C hold. Reactions were then run on 2% agarose gels and imaged with

GelGreen dye according to manufacturer's instructions (Biotium). For a list of all primers

used for MASC-PCR see **Table S2**.

Alcian blue stained gels

To prepare cell lysates, ~10^9 cells of the indicated *V. natriegens* strains were pelleted and

then resuspended in 180 μL of Buffer ATL (Qiagen). Then, 20 μL of a 20 mg/mL proteinase

K stock solution was added to each reaction and incubated at 56°C for 20 mins. Samples

were then boiled in 2X SDS PAGE sample buffer and separated on 4-12% SDS PAGE gels.

Gels were then stained with 0.1% Alcian Blue 8GX in 40% ethanol/3% acetic acid as

previously described[23]. The gel was then destained in a 40% ethanol/3% acetic acid and

imaged on a Biorad ChemiDoc MP Imaging system.

Whole genome sequencing

Genomic DNA was prepped from strains and sequencing libraries were prepped via

homopolymer-tail mediated ligation exactly as previously described[24]. Single-end 50 bp

reads were collected on the Illumina platform. Then, data was analyzed for small indels and

single nucleotide variants using CLC Genomics Workbench exactly as previously

described[15, 25].

Qualitative and quantitative assessment of PHB production

PHB was qualitatively assessed in MuGENT edited populations of *V. natriegens* by plating

onto Nile red containing medium with excess glucose as a carbon source and 100 μM IPTG

to induce P$_{tac}$-containing genome edits = recipe per L: 28 g instant ocean, 2.5 g tryptone, 1 g

yeast extract, 20 g glucose, 15 g agar, and 1 mg Nile red. Fluorescence of colonies was

detected using a PrepOne Sapphire LED blue light base (475 nm ± 30 nm) and amber filter

(530 nm long pass) (Embi Tec).

For quantitative assessment of PHB levels, the indicated strains were grown overnight in

M9 minimal medium (BD) supplemented with 2 mM MgSO$_4$, 100 μM CaCl$_2$, 200 mM NaCl,

30 μM FeSO$_4$, 100 μM IPTG, 1% tryptone, and 2% glucose. Approximately 8×10^9 cells were

then pelleted, resuspended with 50 μL water and transferred to pre-weighed glass screw-

cap tubes. Cell suspensions were dried for 5 h at 80°C and then the tubes were weighed

again to determine dry cell weights. PHB was then hydrolyzed and extracted as crotonic

acid by boiling the dried cells in 1 ml of pure sulfuric acid. Extracts were chilled on ice and

diluted with 4 ml ice-cold water. Aliquots were further diluted 10-fold with water,

centrifuged, filtered, and then crotonic acid was quantified by HPLC as described[26].

ACKNOWLEDGEMENTS

5 We would like to thank Tufts TUCF Genomics and the Indiana University CGB for assistance

6 with whole genome sequencing of strains. This work was supported by

7 US National Institutes of Health Grant AI118863 to ABD. JBM and ABD were also supported

8 by the Indiana University College of Arts and Sciences.

COMPETING FINANCIAL INTERESTS

11 MuGENT is the subject of a pending patent application.

AUTHOR CONTRIBUTIONS

14 ABD conceived the study. TND, SS, CJM, JMB, and ABD designed experiments. TND, CAH,

15 JMB, and ABD performed experiments. All authors played a role in writing and/or

16 proofreading the manuscript.

SUPPORTING INFORMATION

19 Supplementary protocol: Natural transformation / MuGENT in *V. natriegens*.

20 Supplementary Tables S1-S2: strains and primers used in this study.

REFERENCES

[1] Eagon, R. G. (1962) Pseudomonas natriegens, a marine bacterium with a generation

time of less than 10 minutes, *J Bacteriol 83*, 736-737.

[2] Payne, W. J., Eagon, R. G., and Williams, A. K. (1961) Some observations on the

physiology of Pseudomonas natriegens nov. spec, *Antonie Van Leeuwenhoek 27*, 121-

128.

[3] Weinstock, M. T., Hesek, E. D., Wilson, C. M., and Gibson, D. G. (2016) Vibrio natriegens

as a fast-growing host for molecular biology, *Nat Methods 13*, 849-851.

[4] Lee, H. H., Ostrov, N., Wong, B. G., Gold, M. A., Khalil, A., and Church, G. M. (2016) Vibrio

natriegens, a new genomic powerhouse, *bioRxiv*, doi:10.1101/058487.

[5] Meibom, K. L., Blokesch, M., Dolganov, N. A., Wu, C. Y., and Schoolnik, G. K. (2005) Chitin induces natural competence in Vibrio cholerae, *Science 310*, 1824-1827.

[6] Chen, Y., Dai, J., Morris, J. G., Jr., and Johnson, J. A. (2010) Genetic analysis of the capsule polysaccharide (K antigen) and exopolysaccharide genes in pandemic Vibrio parahaemolyticus O3:K6, *BMC Microbiol 10*, 274.

[7] Gulig, P. A., Tucker, M. S., Thiaville, P. C., Joseph, J. L., and Brown, R. N. (2009) USER friendly cloning coupled with chitin-based natural transformation enables rapid mutagenesis of Vibrio vulnificus, *Appl Environ Microbiol 75*, 4936-4949.

[8] Pollack-Berti, A., Wollenberg, M. S., and Ruby, E. G. (2010) Natural transformation of Vibrio fischeri requires tfoX and tfoY, *Environ Microbiol 12*, 2302-2311.

[9] Dalia, A. B., Lazinski, D. W., and Camilli, A. (2014) Identification of a membrane-bound transcriptional regulator that links chitin and natural competence in Vibrio cholerae, *MBio 5*, e01028-01013.

[10] Yamamoto, S., Mitobe, J., Ishikawa, T., Wai, S. N., Ohnishi, M., Watanabe, H., and Izumiya, H. (2014) Regulation of natural competence by the orphan two-component system sensor kinase ChiS involves a non-canonical transmembrane regulator in Vibrio cholerae, *Mol Microbiol 91*, 326-347.

[11] Lorenz, M. G., and Wackernagel, W. (1994) Bacterial gene transfer by natural genetic transformation in the environment, *Microbiol Rev 58*, 563-602.

[12] Dalia, A. B., McDonough, E., and Camilli, A. (2014) Multiplex genome editing by natural transformation, *Proc Natl Acad Sci U S A 111*, 8937-8942.

[13] Erickson, R. J., and Copeland, J. C. (1973) Congression of unlinked markers and genetic mapping in the transformation of Bacillus subtilis 168, *Genetics 73*, 13-21.

[14] Bik, E. M., Bunschoten, A. E., Willems, R. J., Chang, A. C., and Mooi, F. R. (1996) Genetic organization and functional analysis of the otn DNA essential for cell-wall polysaccharide synthesis in Vibrio cholerae O139, *Mol Microbiol 20*, 799-811.

[15] Hayes, C. A., Dalia, T. N., and Dalia, A. B. (2017) Systematic genetic dissection of PTS in Vibrio cholerae uncovers a novel glucose transporter and a limited role for PTS during infection of a mammalian host, *Mol Microbiol*.

[16] Wang, H. H., Isaacs, F. J., Carr, P. A., Sun, Z. Z., Xu, G., Forest, C. R., and Church, G. M.

(2009) Programming cells by multiplex genome engineering and accelerated

evolution, *Nature 460*, 894-898.

[17] Chien, C. C., Chen, C. C., Choi, M. H., Kung, S. S., and Wei, Y. H. (2007) Production of poly-

beta-hydroxybutyrate (PHB) by Vibrio spp. isolated from marine environment, *J*

Biotechnol 132, 259-263.

[18] Centeno-Leija, S., Huerta-Beristain, G., Giles-Gomez, M., Bolivar, F., Gosset, G., and

Martinez, A. (2014) Improving poly-3-hydroxybutyrate production in Escherichia

coli by combining the increase in the NADPH pool and acetyl-CoA availability,

Antonie Van Leeuwenhoek 105, 687-696.

[19] Spiekermann, P., Rehm, B. H., Kalscheuer, R., Baumeister, D., and Steinbuchel, A. (1999)

A sensitive, viable-colony staining method using Nile red for direct screening of

bacteria that accumulate polyhydroxyalkanoic acids and other lipid storage

compounds, *Arch Microbiol 171*, 73-80.

[20] Kidwell, J., Valentin, H. E., and Dennis, D. (1995) Regulated expression of the

Alcaligenes eutrophus pha biosynthesis genes in Escherichia coli, *Appl Environ*

Microbiol 61, 1391-1398.

[21] Csorgo, B., Nyerges, A., Posfai, G., and Feher, T. (2016) System-level genome editing in

microbes, *Curr Opin Microbiol 33*, 113-122.

[22] Dalia, A. B., Lazinski, D. W., and Camilli, A. (2013) Characterization of undermethylated

sites in Vibrio cholerae, *J Bacteriol 195*, 2389-2399.

[23] Mercaldi, M. P., Dams-Kozlowska, H., Panilaitis, B., Joyce, A. P., and Kaplan, D. L. (2008)

Discovery of the dual polysaccharide composition of emulsan and the isolation of

the emulsion stabilizing component, *Biomacromolecules 9*, 1988-1996.

[24] Lazinski, D. W., and Camilli, A. (2013) Homopolymer tail-mediated ligation PCR: a

streamlined and highly efficient method for DNA cloning and library construction,

Biotechniques 54, 25-34.

[25] Seed, K. D., Yen, M., Shapiro, B. J., Hilaire, I. J., Charles, R. C., Teng, J. E., Ivers, L. C., Boncy,

J., Harris, J. B., and Camilli, A. (2014) Evolutionary consequences of intra-patient

phage predation on microbial populations, *eLife 3*, e03497.

[26] Karr, D. B., Waters, J. K., and Emerich, D. W. (1983) Analysis of Poly-beta-

Hydroxybutyrate in Rhizobium japonicum Bacteroids by Ion-Exclusion High-

Pressure Liquid Chromatography and UV Detection, *Appl Environ Microbiol 46*,

1339-1344.

FIGURE LEGENDS

Fig. 1 – *Natural transformation of* V. natriegens. (**a-d**) Transformation assays of *V.*

natriegens. (**a**) *V. natriegens* strains containing a pMMB empty vector or pMMB with the

tfoX gene from either *V. natriegens* (Vn) or *V. cholerae* (Vc) were transformed with 100 ng

of a Δ*dns*::Kan^R tDNA containing 3 kb flanks of homology on both sides of the mutation (i.e.

3 kb/3 kb). Transformation assay of *V. natriegens* pMMB-*tfoX* (Vc) with (**b**) the indicated

concentration of Δ*dns*::Kan^R (3 kb/3 kb) tDNA or (**c**) 5 ng of Δ*dns*::Kan^R tDNA containing

the indicated amount of homology on each side of the mutation. (**d**) Transformation assay

in the indicated strain backgrounds with 5 ng of *rpsL* K43R Sm^R (3 kb/3 kb) or Δ*dns*::Spec^R

(3 kb/3 kb) tDNA as indicated. All strains in **d** harbor Δ*dns*::Kan^R mutations and pMMB-*tfoX*

(Vc). All data are shown as the mean ± SD and are the result of at least 4 independent

biological replicates. ** = *p*<0.01.

Fig. 2 – V. natriegens *produces a WbfF-dependent capsular polysaccharide*. (**a**) Colony

morphologies of parent (white arrow) and Δ*wbfF* (black arrow) strains, which demonstrate

the phenotypes screened for in cotransformation assays. (**b**) Cell lysates of the indicated

strains were run on a 4-12% SDS PAGE gel and stained with the carbohydrate stain Alcian

blue. The presence of a high molecular weight polysaccharide in the parent is indicated by a

red arrow.

Fig. 3 – *Cotransformation is highly efficient in* V. natriegens. (**a**) Cotransformation was

tested using a Δ*dns*::Kan^R (3 kb/3 kb) selected product and an unselected product that

deleted ∼500 bp of the 5' end of *wbfF* gene. Cotransformation assays were performed using

50ng of the Δ*dns*::Kan^R (3 kb/3 kb) selected product and (**b**) the indicated amount of the

Δ*wbfF* (3 kb/3 kb) unselected product or (**c**) 200 ng of Δ*wbfF* unselected products

containing the indicated length of homology on each side of the mutation. Data in **b** and **c**

are from at least four independent biological replicates and shown as the mean ± SD. (**d**)

Schematic of MuGENT. The selected product is indicated by a red box, while multiple

unselected genome edits are depicted by distinct gray shapes. Since there is no selection for

genome edits *in cis*, output mutants can have any number and combination of the

unselected genome edits. Circles inside cells represent the two circular chromosomes of *V.*

natriegens. (**e** and **f**) MuGENT was performed with 5 unselected genome edits. The selected

product was Δ*wbfF*::Kan^R, while the unselected products targeted four carbohydrate

transporters and *dns* for inactivation by replacing ~500 bp of the 5' end of each gene with a

premature stop codon. (**e**) A representative MASC-PCR gel of 24 colonies from the edited

population. The targets of each genome edit are indicated on the left and the presence of a

band indicates integration of the indicated genome edit. Strains containing 4 genome edits

are indicated by the green arrows. (**f**) Distribution of genome edits in the population

determined by MASC-PCR analysis of 48 random mutants.

Fig. 4 – *MuGENT rapidly enhances PHB production in* V. natriegens. (**a**) The indicated

targets were subjected to either a promoter swap (top) or inactivation by replacing

~500bp of the 5' end of each gene with a short sequence to introduce a premature stop

codon (bottom). (**b**) Distribution of the 9 genome edits in a population of cells following

four cycles of MuGENT. (**c**) Representative image of the mutant pool generated in **b** plated

on Nile red containing plates, which stain PHB granules. White arrows indicate colonies

with increased fluorescence intensity compared to the parent. (**d**) PHB content of select

MuGENT optimized strains is shown as the % of dry cell weight (DCW). The genotype of

each mutant is shown below each bar where a filled box indicates the presence of the

genome edit indicated on the left. Data are shown as the mean ± SD and are from at least 2

independent biological replicates. * = $p<0.05$.

Fig. 1 – *Natural transformation of* V. natriegens. (**a-d**) Transformation assays of *V. natriegens*. (**a**) *V. natriegens* strains containing a pMMB empty vector or pMMB with the *tfoX* gene from either *V. natriegens* (Vn) or *V. cholerae* (Vc) were transformed with 100 ng of a Δ*dns*::Kan^R tDNA containing 3 kb flanks of homology on both sides of the mutation (i.e. 3 kb/3 kb). Transformation assay of *V. natriegens* pMMB-*tfoX* (Vc) with (**b**) the indicated concentration of Δ*dns*::Kan^R (3 kb/3 kb) tDNA or (**c**) 5 ng of Δ*dns*::Kan^R tDNA containing the indicated amount of homology on each side of the mutation. (**d**) Transformation assay in the indicated strain backgrounds with 5 ng of *rpsL* K43R Sm^R (3 kb/3 kb) or Δ*dns*::Spec^R (3 kb/3 kb) tDNA as indicated. All strains in **d** harbor Δ*dns*::Kan^R mutations and pMMB-*tfoX* (Vc). All data are shown as the mean ± SD and are the result of at least 4 independent biological replicates. ** = *p*<0.01.

Fig. 2 – V. natriegens *produces a WbfF-dependent capsular polysaccharide*. (**a**) Colony morphologies of parent (white arrow) and Δ*wbfF* (black arrow) strains, which demonstrate the phenotypes screened for in cotransformation assays. (**b**) Cell lysates of the indicated strains were run on a 4-12% SDS PAGE gel and stained with the carbohydrate stain Alcian blue. The presence of a high molecular weight polysaccharide in the parent is indicated by a red arrow.

Fig. 3 – *Cotransformation is highly efficient in* V. natriegens. (**a**) Cotransformation was tested using a Δ*dns*::Kan^R (3 kb/3 kb) selected product and an unselected product that deleted ~500 bp of the 5' end of *wbfF* gene. Cotransformation assays were performed using 50ng of the Δ*dns*::Kan^R (3 kb/3 kb) selected product and (**b**) the indicated amount of the Δ*wbfF* (3 kb/3 kb) unselected product or (**c**) 200 ng of Δ*wbfF* unselected products containing the indicated length of homology on each side of the mutation. Data in **b** and **c** are from at least four independent biological replicates and shown as the mean ± SD. (**d**) Schematic of MuGENT. The selected product is indicated by a red box, while multiple unselected genome edits are depicted by distinct gray shapes. Since there is no selection for genome edits *in cis*, output mutants can have any number and combination of the unselected genome edits. Circles inside cells represent the two circular chromosomes of *V. natriegens*. (**e** and **f**) MuGENT was performed with 5 unselected genome edits. The selected product was Δ*wbfF*::Kan^R, while the unselected products targeted four carbohydrate transporters and *dns* for inactivation by replacing ~500 bp of the 5' end of each gene with a premature stop codon. (**e**) A representative MASC-PCR gel of 24 colonies from the edited population. The targets of each genome edit are indicated on the left and the presence of a band indicates integration of the indicated genome edit. Strains containing 4 genome edits are indicated by the green arrows. (**f**) Distribution of genome edits in the population determined by MASC-PCR analysis of 48 random mutants.

Fig. 4 – *MuGENT rapidly enhances PHB production in* V. natriegens. (**a**) The indicated targets were subjected to either a promoter swap (top) or inactivation by replacing ~500bp of the 5' end of each gene with a short sequence to introduce a premature stop codon (bottom). (**b**) Distribution of the 9 genome edits in a population of cells following four cycles of MuGENT. (**c**) Representative image of the mutant pool generated in **b** plated on Nile red containing plates, which stain PHB granules. White arrows indicate colonies with increased fluorescence intensity compared to the parent. (**d**) PHB content of select MuGENT optimized strains is shown as the % of dry cell weight (DCW). The genotype of each mutant is shown below each bar where a filled box indicates the presence of the genome edit indicated on the left. Data are shown as the mean ± SD and are from at least 2 independent biological replicates. * = $p < 0.05$.

Supplementary material for:

Multiplex genome editing by natural transformation (MuGENT) for synthetic biology in *Vibrio natriegens*

Triana N. Dalia, Chelsea A. Hayes, Sergey Stolyar, Christopher J. Marx, James B. McKinlay, and Ankur B. Dalia

<u>Contents:</u>

Supplementary protocol: Natural transformation / MuGENT in *V. natriegens*

Materials:
1. LBv2 = Mix 400 mL LB + 32 mL sterile 5 M NaCl + 3.4 mL sterile 1 M KCl + 18.5 mL sterile 1 M MgCl2
2. 2XIO = 28 g/L of instant ocean sea salts (www.instantocean.com)
3. Transforming DNA (tDNA) –
   a. selected product = mutant construct that has a selectable marker (antibiotic resistance cassette) that replaces the gene of interest or a neutral locus (for MuGENT). Homology on each side of the mutation can be as little as 0.5 kb. Homology of 3 kb on either side of the mutation results in the highest transformation efficiencies.
   b. Unselected product (cotransformation / MuGENT) = mutant construct that lacks any selectable marker but has a mutation of interest (deletion, point mutation, promoter swap, etc.). Unselected products should have 3 kb of homology on each side of the mutation for the highest rates of cotransformation / MuGENT.

Notes:
1. Carb100 = Carbenicillin 100 μg/mL
2. SAD1306 = *V. natriegens* Rif^R 14048 pMMB67EH-tfoX
3. The pMMB67EH-tfoX plasmid is very stable in *V. natriegens*, therefore, Carbenicillin (Carb) is not needed to maintain it throughout the transformation protocol.

Procedure:
1. Inoculate 3 mL of LBv2+Carb100+100 μM IPTG in a culture tube with SAD1306. Grow at 30°C in rollerdrum overnight (12-18 hours).
2. Next day, take culture out of incubator and measure OD_{600}. Generally, our overnights are at an OD_{600} of between 7-10.
3. For each transformation reaction, take 3.5 μL of the overnight culture and dilute into 350 μL of 2XIO+100 μM IPTG (no Carb). Invert gently to mix. Be sure to also prep a "no DNA" control reaction.
4. Add tDNA to each reaction and invert gently to mix:
   a. For a selected product (i.e. a product that has an antibiotic resistance marker) = ~5-50 ng yields thousands of colonies.
   b. For cotransformation / MuGENT = use 50 ng of a selected product and ~200 ng of each unselected product
5. Incubate reactions at 30°C statically for 4-6 hours.
6. Next, add 1mL LBv2 (no drug) to each transformation reaction and outgrow at 30°C shaking for 1-2 hours
7. To determine the transformation efficiency:
   a. Plate all reactions for quantitative culture on media to select for the transformants (i.e. on antibiotic plates that select for integration of selected

product) and on plates without any drug to determine the total CFU in the
culture.
   b. Transformation efficiency = transformants CFU / total CFU
8. For cotransformation / MuGENT:
   a. Plate onto media to select for integration of the selected product.
   b. Pick single colonies and screen by MASC-PCR to identify clones with the
      desired genome edits.
   c. To perform a subsequent round of MuGENT, ~200 μL of the outgrown
      transformation can be inoculated into 3 mL of LBv2+Carb100+100 μM
      IPTG+the antibiotic that selects for integration of the selected product (e.g. if
      Δdns::Kan^R was used in the first cycle of MuGENT, then 50 μg/mL Kan would
      be included in this overnight culture). Start at "step 2" of this procedure to
      perform the next cycle of MuGENT being sure to use a selected product with
      a distinct Ab^R marker.

Supplementary Tables

Table S1 – *Strains used in this study*

Strain name	Genotype and antibiotic resistances	Description	Reference / (strain#)
WT	Rif^R	Spontaneous Rif^R derivative *V. natriegens* ATCC14048 that is the parent isolate for all strains used in this study.	This study (SAD1304)
pMMB-tfoX (Vc)	pMMB-*tfox* (Vc) Carb^R	SAD1304 containing pMMB-*tfoX* (Vc), a vector containing the *tfoX* gene from *V. cholerae* (VC1153) under the control of an IPTG-inducible P$_{tac}$ promoter. Vector is derived from pMMB67EH and has a Carb^R gene for selection.	This study (SAD1306)
pMMB-tfoX (Vn)	pMMB-*tfox* (Vn) Carb^R	SAD1304 containing pMMB-*tfoX* (Vn), a vector containing the *tfoX* gene from *V. natriegens* (BA890_05980) under the control of an IPTG-inducible P$_{tac}$ promoter. Vector is derived from pMMB67EH and has a Carb^R gene for selection.	This study (TND0322 / SAD1495)
pMMB	pMMB empty vector Carb^R	SAD1304 containing the pMMB67EH empty vector	This study (TND0321 / SAD1496)
WT (Fig. 1D)	pMMB-*tfoX* (Vc) Carb^R, Δdns::Kan^R	SAD1306 with Δdns::Kan^R (ΔBA890_12415)	This study (SAD1313)
$\Delta mutS$	pMMB-tfoX (Vc) Carb^R, Δdns::Kan^R, $\Delta mutS$	Generated by cotransformation into SAD1306 with Δdns::Kan^R and a product to delete ~500bp of the 5' end of the *mutS* gene	This study (TND0362 / SAD1497)

		(BA890_12150).	
Δ*wbfF*	pMMB-tfoX (Vc) Carb^R, Δ*wbfF*::Kan^R	Introduced a Δ*wbfF*::Kan^R mutation (ΔBA890_01135) into the SAD1306 strain background.	This study (CAH509 / SAD1498)
MuGENT quadruple mutant	pMMB-tfoX (Vc) Carb^R, Δ*wbfF*::Kan^R, ΔBA890_01815 (mannitol transporter), ΔBA890_19540 (sucrose transporter), ΔBA890_16410 (fructose transporter), Δ*dns*	MuGENT into SAD1306 strain with 5 unselected genome edits. This quadruple mutant was whole genome sequenced and no off target mutations were identified.	This study (TND0338 / SAD1499)
Fig. 4E, second bar	pMMB-tfoX (Vc) Carb^R, Δ*dns*::Kan^R, P*tac*-*phaBAC*	MuGENT into SAD1306 to enhance PHB production. The strain contains the genome edits indicated.	This study (TND0364 / SAD1500)
Fig. 4E, third bar	pMMB-tfoX (Vc) Carb^R, Δ*dns*::Kan^R, P*tac*-*phaBAC*, P*tac*-*nadK*, Δ*ldhA*	MuGENT into SAD1306 to enhance PHB production. The strain contains the genome edits indicated.	This study (SAD1501)
Fig. 4E, fourth bar	pMMB-tfoX (Vc) Carb^R, Δ*dns*::Kan^R, P*tac*-*phaBAC*, P*tac*-*pntAB*, Δ*pta*, Δ*gltA*, Δ*aceA*	MuGENT into SAD1306 to enhance PHB production. The strain contains the genome edits indicated.	This study (SAD1502)
Fig. 4E, fifth bar	pMMB-tfoX (Vc) Carb^R, Δ*dns*::Spec^R, P*tac*-*phaBAC*, P*tac*-*nadK*, Δ*pta*, Δ*gltA*, Δ*ldhA*	MuGENT into SAD1306 to enhance PHB production. The strain contains the genome edits indicated.	This study (SAD1503)
Fig. 2E, sixth bar	pMMB-tfoX (Vc) Carb^R, Δ*dns*::Spec^R, P*tac*-*phaBAC*, P*tac*-*nadK*, Δ*pta*, Δ*gltA*, Δ*ldhA*, Δ*aceA*	MuGENT into SAD1306 to enhance PHB production. The strain contains the genome edits indicated.	This study (SAD1504)

Table S2 – *Primers used in this study*

Primer Name	Primer Sequence (5'→3')*	Description
Primers for Mutant constructs		
ABD123	ATTCCGGGGATCCGTCGAC	Amplify MIDDLE Ab^R (Kan^R, Spec^R, or Tm^R cassettes) F
ABD124	TGTAGGCTGGAGCTGCTTC	Amplify MIDDLE Ab^R (Kan^R, Spec^R, or Tm^R cassettes) R
BBC1264	CTAACATGGCTAAGCACCTG	Δ*dns* F1 (3kb)
BBC1605	GCACTTCTTCGCGAATTCGC	Δ*dns* F1 (2kb)
BBC1607	AGTGATTGGGTCACTCATTGG	Δ*dns* F1 (1kb)
BBC1609	AATGAGATTCGCCTTAACCC	Δ*dns* F1 (0.5kb)
BBC1265	gtcgacggatccccggaatAGAGAACAGGTATTTCATAGTTAAAGTC	Δ*dns* R1
BBC1266	gaagcagctccagcctacaTAATCCTCACCAATCGCGAC	Δ*dns* F2
BBC1610	TCGAGCTTTACGCCACAACG	Δ*dns* R1 (0.5kb)
BBC1608	ACACCTTGGTCGAGGTGAAG	Δ*dns* R1 (1kb)
BBC1606	ATAACGCAGTAGAAAGTATCCAC	Δ*dns* R1 (2kb)
BBC1267	ACTGGTAAGCCATAACGACC	Δ*dns* R1 (3kb)
DOG0246	AGGCTCGTGTTGCATGTGAG	Δ*dns* 501bp F1
DOG0247	gctaattcagtttaagcggccatCATAGTTAAAGTCTTTAAAAAGTA	Δ*dns* 501bp R1

	TGACTT	
DOG0248	atggccgcttaaactgaattagcATCGCTCGTACCTATCTTTATATG	Δ*dns* 501bp F2
DOG0249	TAAGGTGTCTCAAATCTCAATCTAGG	Δ*dns* 501bp R2
BBC1255	TGAGAAATTCTTTGCATCACATC	*rpsL* K43R (Sm[R]) F1
BBC1256	GAAGTGCTGAGTTAGGTTTTcTAGGTGTAGTAGTGTAAACAC	*rpsL* K43R (Sm[R]) R1
BBC1257	GTGTTTACACTACTACACCTAgAAAACCTAACTCAGCACTTC	*rpsL* K43R (Sm[R]) F2
BBC1258	GTAGTGACGAGTTGGAGTG	*rpsL* K43R (Sm[R]) R2
BBC1552	GAACTGCATGAATACGTTGTTCC	Δ*mutS* 501bp F1
BBC1553	gctaattcagtttaagcggcCACAGGTAAGTTCTTTTGTTTATTTCC	Δ*mutS* 501bp R1
BBC1554	GTGgccgcttaaactgaattagcCGCACCGCACCACGTGAG	Δ*mutS* 501bp F2
BBC1555	GAGTATCAGCAACACAGTAACC	Δ*mutS* 501bp R2
BBC1347	TAGCAACTGTTTTAGCGCTG	Δ*wbfF* F1
BBC1348	gtcgacggatccccggaatCTTTTATCATCATACTCATTCATTAAAG	Δ*wbfF* R1
BBC1349	gaagcagctccagcctacaTGATGTATAAGCGTCATTTATTCG	Δ*wbfF* F2
BBC1350	GTTCCTGTCGATAAGTATTGATC	Δ*wbfF* R2
DOG0353	AATGTCGGCCTTCTGATTAG	Δ*wbfF* 501bp F1 (3kb)
BBC1612	TAAACTTTATCAGCGACGTCAG	Δ*wbfF* 501bp F1 (2kb)
BBC1614	TTCAGGAACGATGTCGACAG	Δ*wbfF* 501bp F1 (1kb)
DOG0354	gctaattcagtttaagcggccatTATCATCATACTCATTCATTAAAGTTTTAA	Δ*wbfF* 501bp R1
DOG0355	atggccgcttaaactgaattagcACTAATAACGTCAGTGTATACGTAAAC	Δ*wbfF* 501bp F2
BBC1615	CCACGCAATGTAGTCATCAATC	Δ*wbfF* 501bp R2 (1kb)
BBC1613	GGATACGCAGCATACCTTG	Δ*wbfF* 501bp R2 (2kb)
BBC1611	TTAATTGTGCCTGAGCAAGC	Δ*wbfF* 501bp R2 (3kb)
DOG0271	AAGTAGTGATGATCCGAAGCG	ΔBA890_01815 501bp (mannitol transporter) F1
DOG0272	gctaattcagtttaagcggccatCATAACAATTCCCCGTTCGATG	ΔBA890_01815 501bp (mannitol transporter) R1
DOG0273	atggccgcttaaactgaattagcCTTGTATCAGCGCACCTTCTAC	ΔBA890_01815 501bp (mannitol transporter) F2
DOG0274	ATCGTGGTAAATATCGTCAGGTAG	ΔBA890_01815 501bp (mannitol transporter) R2
DOG0266	ATCTCGGCTTGTCTACACCAG	ΔBA890_19540 (sucrose transporter) F1
DOG0267	gctaattcagtttaagcggccatCATTGCACACCCCGATTGG	ΔBA890_19540 (sucrose transporter) R1
DOG0268	atggccgcttaaactgaattagcTATTTACCTGTTTTATTGGCGTTTTC	ΔBA890_19540 (sucrose transporter) F2
DOG0269	TGAACTGAATCCTCGCAGG	ΔBA890_19540 (sucrose transporter) R2
DOG0256	ATGCTCGTCATCCATGGGAC	ΔBA890_16410 (fructose transporter) F1
DOG0257	gctaattcagtttaagcggccatCATACTGATAACCTTCTGTTCCTTAG	ΔBA890_16410 (fructose transporter) R1
DOG0258	atggccgcttaaactgaattagcACCGCGCAAGAGATCGAAG	ΔBA890_16410 (fructose transporter) F2
DOG0259	TTGGGTGCTTTGCTTCTCG	ΔBA890_16410 (fructose transporter) R2

DOG0261	ATCTGAACTTAGGATACTCACATC	ΔBA890_03375 (trehalose transporter) F1
DOG0262	gctaattcagtttaagcggccatCATAACTTTGCCCACCCTGTATTG	ΔBA890_03375 (trehalose transporter) R1
DOG0263	atggccgcttaaactgaattagcTTCTTCCTGCCTGTTGGC	ΔBA890_03375 (trehalose transporter) F2
DOG0264	AGTCAGATGGCGATTGATGTG	ΔBA890_03375 (trehalose transporter) R2
ABD840	TTAATTGCGTTGCGCTCACTGCCCGACTCCCGTTCTGGATAATGTTTTTTGC	Amplify MIDDLE P$_{tac}$ construct F
ABD625	CTGATGAATCCCCTAATGATTTTGG	Amplify MIDDLE P$_{tac}$ construct R
BBC1536	GTAACGAACGTGTCATCAGTG	P$_{tac}$-*phaBAC* F1
BBC1540	CGGGCAGTGAGCGCAACGCAATTAATGCAAGCGCACTAATATGAC	P$_{tac}$-*phaBAC* R1
BBC1541	CAAAATCATTAGGGGATTCATCAGAAAGAATGGAGTCGTCAATGAATAAAG	P$_{tac}$-*phaBAC* F2
BBC1577	CGACATCTTCACCAACACG	P$_{tac}$-*phaBAC* R2
BBC1621	TCTGGAGAGTATGTTGGCC	P$_{tac}$-*pntAB* F1
BBC1622	cgggcagtgagcgcaacgcaattaaCCTTGTATACATATCAATTAATTAGTCCC	P$_{tac}$-*pntAB* R1
BBC1623	caaaatcattagggggattcatcagAggaggTTGCGTTTTGCAAATCGGTGTAC	P$_{tac}$-*pntAB* F2
BBC1624	AGACTACGCCAAACTATACAGC	P$_{tac}$-*pntAB* R2
BBC1616	CTTCTTCGTCTTCAAAACGACG	P$_{tac}$-*nadK* F1
BBC1617	cgggcagtgagcgcaacgcaattaaGCATTAAAGAGGCTTGAATCAGG	P$_{tac}$-*nadK* R1
BBC1618	caaaatcattagggggattcatcagaggaggtAATGCTATGAAAAATCCATGTAACG	P$_{tac}$-*nadK* F2
BBC1619	CTGCGCTGATAATAAACAAGC	P$_{tac}$-*nadK* R2
BBC1626	CACAAATAGCGAAGCTAACTG	P$_{tac}$-*udhA* F1
BBC1627	cgggcagtgagcgcaacgcaattaaTATTTGCTTAACATTGCCTTAGC	P$_{tac}$-*udhA* R1
BBC1628	caaaatcattagggggattcatcagAggaggtTACATCATGGCGCATGTAAATC	P$_{tac}$-*udhA* F2
BBC1629	GTGAAAGTATTTTCGCCTTTCG	P$_{tac}$-*udhA* R2
BBC1636	GACAAGTCAGAAAGTCCAGTCAC	Δ*pta* 501bp F1
BBC1637	gctaattcagtttaagcggccatAGACATTCGTAGAGTACCTTTGC	Δ*pta* 501bp R1
BBC1638	atggccgcttaaactgaattagcGTTATCATCAACAAGCTAAACGCAC	Δ*pta* 501bp F2
BBC1639	GATATCAACGAGTTTGCATCTG	Δ*pta* 501bp R2
BBC1646	GCTAACATCAATGCGTATGCC	Δ*pgi* 501bp F1
BBC1647	gctaattcagtttaagcggccatCAACATGGTCTTTATCCCGATG	Δ*pgi* 501bp R1
BBC1648	atggccgcttaaactgaattagcGCACTGGCACCATACAAAAAC	Δ*pgi* 501bp F2
BBC1649	CCTTTCTCAGACACTATCGACAC	Δ*pgi* 501bp R2
BBC1641	AGCCTTCTTCTACATCAAGTGTG	Δ*gltA* 501bp F1
BBC1642	gctaattcagtttaagcggccatATCCGCCATAACAATCTCCTTTG	Δ*gltA* 501bp R1
BBC1643	atggccgcttaaactgaattagcACACTGGCGGCAATGTGTTAC	Δ*gltA* 501bp F2
BBC1644	CAAGAGTACTACGAAGAGCTG	Δ*gltA* 501bp R2
BBC1651	CTTGTAACACTGCCGCTAAGAG	Δ*ldhA* 501bp F1
BBC1652	gctaattcagtttaagcggccatCATGGTTCTCTCTCGAAATCATTG	Δ*ldhA* 501bp R1
BBC1653	atggccgcttaaactgaattagcATGGAAATTCTTTGCCATGATCC	Δ*ldhA* 501bp F2
BBC1654	AGTGTGTTACTTATTTGGAGGATG	Δ*ldhA* 501bp R2
BBC1631	TGAACTGCTGGCGAAAGGAC	Δ*aceA* 501bp F1

BBC1632	GCTAATTCAGTTTAAGCGGCCATTGGTCTATCCCTCTTTATAATTTGC	Δ*aceA* 501bp R1
BBC1633	ATGGCCGCTTAAACTGAATTAGCCTAAATGCTTACGAACTGATGAAATC	Δ*aceA* 501bp F2
BBC1634	CGATTGAAGCTTGAAGAACAAGC	Δ*aceA* 501bp R2
Primers for MASC-PCR		
ABD969	ATGGCCGCTTAAACTGAATTAGC	Universal F primer for all Δ501 bp genome edits
DOG0250	TGGTTGCCTTGTACTTTGGC	R detect for Δ*dns* 501 bp (152bp product)
BBC1556	AGTGATCGAGAACAGCGG	R detect for Δ*mutS* 501bp (402bp product)
DOG0356	ATAGCTACCGCGTTCAGGG	R detect for Δ*wbfF* 501bp (165bp product)
DOG0275	AGTGACGTGGATGTTCAGAC	R detect for Δ*BA890_01815* 501bp (mannitol transporter) (750bp product)
DOG0270	AACCCAGTGATACCAGATGG	R detect for Δ*BA890_19540* (sucrose transporter) (650bp product)
DOG0260	TATTCATCAGTGCAGCGGC	R detect for Δ*BA890_16410* (fructose transporter) (352 bp product)
DOG0265	TCTTGCATTAACTGTAAATCCACG	R detect for Δ*BA890_03375* (trehalose transporter) (500 bp product)
BBC435	ACACTCTTTGGGGGCCAAAATCATTAGGGGATTCATCAG	Universal F primer to detect all P*tac* genome edits
BBC1551	GGTAAACCCTTTGCTGTTAAACC	R detect for P*tac*-*phaBAC* (170bp product)
BBC1625	CTTGAGCTCGAGAGATACG	R detect for P*tac*-*pntAB* (400bp product)
BBC1620	GATAAAATTCGTGCGGCTC	R detect for P*tac*-*nadK* (260bp product)
BBC1630	AGATAATGATATGACGAGGGTC	R detect for P*tac*-*udhA* (550bp product)
BBC1640	CGAATTGGAGAAGTGTTGAAG	R detect for Δ*pta* (140bp product)
BBC1650	AACCCAGTCCCAGAATTCAAAC	R detect for Δ*pgi* (300bp product)
BBC1645	GATGTTGACGCGTTTTGTTCG	R detect for Δ*gltA* (200bp product)
BBC1655	GGCTTCTACGTTATTTAGTGTC	R detect for Δ*ldhA* (450bp product)
BBC1656	TGTTGTGAATACCCGCTAGAG	R detect for Δ*aceA* (600bp product)

*Lower case nucleotides specify overlap regions for SOE PCR