Classification: Evolution and Genetics

Gene-centric intra- and inter-clade recombination in a context of *Esche-

***richia coli* subpopulations**

Yu Kang[1]*, Xing Shi[1,2]*, Lina Yuan[1]*†, Yanan Chu[1]*, Fei Chen[1,2]*, Zilong He[1]*, Zhancheng Gao[3]*, Xinmiao

Jia[4], Qiang Lin[1], Qin Ma[5], Jian Wang[1], Rongrong Fu[1], Jiayan Wu[1], Jingfa Xiao[1,2], Songnian Hu[1,2], Jun

Yu[1,2‡].

1.CAS Key Laboratory of Genome Sciences and Information, Beijing Institute of Genomics, Chinese Academy of Sci-

ences, 100101, Beijing, PR China.

2.University of Chinese Academy of Sciences, Beijing 100190, PR China.

3.Department of Respiratory & Critical Care Medicine, Peking University People's Hospital, Beijing, 100044, PR China

4. Central Research Laboratory, Peking Union Medical College Hospital, Peking Union Medical College & Chinese

Academy of Medical Sciences, Beijing 100730, P.R.China

5.Department of Agronomy, Horticulture, and Plant Science, South Dakota State University, Brookings, SD, 57007,

USA.

* These authors should be regarded as joint First Authors.

† Current address: Suzhou PerkinElmer Medical Laboratory Co., Ltd. No. 115, North Tai Ping Road,Taicang, Jiangsu

215400, P. R. China.

‡ To whom correspondence should be addressed. CAS Key Laboratory of Genome Sciences and Information, Beijing

Institute of Genomics, Chinese Academy of Sciences, Beijing 100101, P.R. China. Tel: +8610-84097898; Fax: +8610-

84097540; Email: junyu@big.ac.cn

Running Head: Fine-scale Map of Genome-wide Recombination

ABSTRACT（**250** words）

Recombination is one of the most important mechanisms of prokaryotic species evolution but its exact roles are still in debate. Here we try to infer genome-wide recombination events within a species utilizing a dataset of 104 complete genomes of *Escherichia coli* from diverse origins, among which 45 from world-wide animal-hosts are in-house sequenced using SMRT (single-molecular real time) technology. Two major clades are identified based on evidences of ecological and physiological characteristics, as well as distinct genomic features implying scarce inter-clade genetic exchange. By comparing the synteny of identical fragments genome-widely searched for each genome pair, we achieve a fine-scale map of recombination within the population. The recombination is rather extensive within clade, which is able to break linkages between genes but does not interrupt core genome framework and primary metabolic portfolios possibly due to natural selection for physiological compatibility and ecological fitness. Meanwhile, the recombination between clades declines drastically as the phylogenetic distance increases, generally 10-fold reduced than those of the intra-clade, which establishes genetic barrier between clades. These empirical data of recombination suggest its critical role in the early stage of speciation, where recombination rate differs according to phylogentic distance. The extensive intra-clade recombination coheres sister strains into a quasi-sexual group and optimizes genes or alleles to streamline physiological activities, whereas shapely declined inter-clade recombination split the population into clades adaptive to divergent ecological niches.

Significance Statement （**120 words**）

Roles of recombination in species evolution have been debated for decades due to difficulties in inferring recombination events during the early stage of speciation, especially when recombination is always complicated by frequent gene transfer events of bacterial genomes. Based on 104 high-quality complete *E. coli* genomes, we infer gene-centric dynamics of recombination in the formation of two *E. coli* clades or subpopulations, and recombination is found to be rather intensive in a within-clade fashion, which forces them to be quasi-sexual. The recombination events can be mapped among individual genomes in the context of genes and their variations; decreased between-clade and increased intra-clade

recombination engender a genetic barrier that further encourages clade-specific secondary metabolic port-

folios for better environmental adaptation. Recombination is thus a major force that accelerates bacterial

evolution to fit ecological diversity.

Main Text

Introduction

Roles of recombination in bacterial evolution have been debated for decades (1, 2). Asexual bac-

teria are believed to rarely recombine their genomes when passing genetic information to progenies, and

a cluster of genes is capable of taking over the entire population if they happen to be linked with an ad-

vantageous gene through genetic hitchhiking, leading to an effect of genome-wide selective sweep as pro-

posed in an "ecotype model"(3, 4). The theory is supported by a recent time-series metagenomics re-

search (5), whereas other studies reveal the counterpart where recombination breaks the linkage and

leaves only the adaptive gene to sweep a population (6, 7). The issue comes that to what extent recombi-

nation actually breaks such genetic linkages and undermines the theory of genome sweep. Another issue

relevant to recombination is its contradictory roles in speciation, where recombination produces incon-

sistent signals in clonal phylogenies (8) in one hand and hijacks similar strains into a separating subpopu-

lation (1, 9) in the other. Recent studies have postulated extensive recombination in bacterial populations

even to a quasi-sexual status(6). Will extensive recombination interrupt primary metabolic portfolios or ge-

nome organization frameworks (GOF) (10), and lead to puzzled or discordant phylogenies? These issues

can be readily settled by resolving recombination events in a genome-scale and at single gene level

within a bacterial population where subpopulations are recently diverged, making a fine-scale and pano-

ramic display of the recombination dynamics in early speciation. However, it is still unachievable due to

the difficulties in inferring recombination events between closely related strains (11).

Here, we investigate the species *E. coli*, an excellent model for studying genotype-phenotype rela-

tionships and molecular evolution mechanisms due to its collective knowledge, ecological diversity, and

availability of high-quality genomes and their annotations. Pangenomic studies have revealed great ge-

nome variability and intensive recombination in *E. coli*, which underlie discordance between phylogenies

of various orthologous genes (12). To generate recombination dynamics genome-wide and in a fine-scale,

we start with 45 in-house genome assemblies from animal hosts, using SMRT (Single-molecule real-time

sequencing) technology, and also put together a collection of 104 complete genome sequences of strains

from diverse geographic and host range from public databases. Based on genome-based phylogenies

coupled with experimentation and detailed genetic analysis, we further classify *E. coli* isolates into two ma-

jor clades or subpopulations, *Vigorous* for its fast movement and *Sluggish* for its slow movement, abbrevi-

ated as Vig and Slu. Taking advantage of our complete genome assemblies, as well as their precise

synteny of identical fragments and genes, we are able to infer recombination events for each genome pair

and to calculate the mobility (transfer among isolates within and across clades) of each gene, which are

used to address: (1) In what degree and intensity does recombination happen within a species and its

subpopulations? (2) What are molecular and physiological bases of the early separation of clades? (3)

Does recombination disrupt GOF and metabolic portfolios of a host strain?

Results

Our study begins with a thorough collection of 202 *E. coli* strains of animal hosts with broad diver-

sity in geography, climate, and host range by field scientists, and the strains have been served as repre-

sentation of the species for detailed taxonomic characterization (13). Of this collection, 45 isolates which

represent maximal genetic heterogeneity and biodiversity in terms of host diet, geographic distribution

(Fig. S1) and MLST clustering (Fig. S2), are selected for genome sequencing with both Illunima and Pac-

Bio RSII platform; and all genomes are finished to a single circular chromosome.

For sequence analysis, we retrieve 59 complete genomes of human-host strains together with 7

draft genomes of environmental strains (14) from the public databases. Based on the 111 genomes, we

extract the core genome of *E. coli*, which has collectively 1,095 genes in a total length of 1.05 Mb. The

maximal-likelihood phylogeny based on the core genome indicates clear separation between host-related

and environmental strains, whereas human- and animal-host strains appear entangled (Fig.1). Most host-

related strains cluster into two major clades; the larger one contains 56 strains of phylotype A and B1,

whereas the smaller one is composed of 31 phylotype B2 strains. The rest 17 strains, mainly in phylotype

D and E, form several minor clades and are more closely positioned to the environment strains. Overall,

this clustering pattern is largely in congruence with their phylotypes and previous reports (15, 16). As for

the two major clades, although they both contain commensals and pathogens and show overlaps in a

wide range of hosts and geographic distributions, they still exhibit skewed distributions in some ecological

and clinical niches. The larger clade includes strains that prefer tropics, carnivorous hosts, and occasion-

ally lead to pandemic infections, whereas strains in the smaller clade show the opposite preferences as

cold climate, herbivorous host, and often cause extra-intestinal and antibiotic-resistant infections. We fur-

ther explore the reasons underlying these ecological distinctions by performing physiological experiments

with our collection strains in the two clades. The strains in the larger clade show higher growth rate, espe-

cially in higher temperature, apparent vulnerability and poorer survival rates at heat stress due to faster

growth, rapid response and motility to amino acid chemotaxis, as well as slower growth in polysaccharides

carbohydrate sources when compared to the smaller clade (Fig.S3). We subsequently name the two

clades as the Vig and the Slu clades, respectively.

We examine genetic bases of the phenotypical distinctions between Vig and Slu in their genomes.

First, we find apparent differences between the two clades in their profiles of dispensable genes based in

a neighbor-joining tree (Fig. S4). By using two-sample Kolmogorov-Smirnov test to all dispensable genes,

we identify 279 and 336 orthologs that are significantly enriched in Vig and Slu, respectively (Table S2).

Half of these clade-associated-genes (CAGs) are metabolic genes according to the COG categories, and

as expected (Fig.S5A), the Vig clade enriches more genes related to motility and metabolism of purine

and amino acid than the Slu clade. Strains of the same clade share common patterns in dispensable met-

abolic genes regardless of host diet preferences (Fig.S5B). We further predict nutrient consumption and

fermentation products by constructing the genome-scale models (GSMs) of metabolism for each strain *in*

silico (17, 18), the two clades are largely separated by characteristic substrates and products, albeit a few

confusions (Fig. 1B). Given a complete rich media which contains 16279 compounds, GSMs predict that

our stains consume or produce 102 of these compounds, and 82 show differences between strains. A total

of 12 compounds show significant difference ($p<0.01$, Wilcoxon test) between the two clades, i.e., the Slu

strains uptake more plant-origin nutrients of dulcose, maltose, and cellobiose, whereas the Vig strains up-

take more taurine and xylose. The Vig strains produce more formate, glycerol, and L-rhamnose, as com-

pared to the Slu strains that produce more lactate and L-proline and are more active in xanthine metabo-

lism. For dispensable virulence genes, the Vig pathogens are rich in T3SS and other secretion systems

and the lethal Shiga-like toxins (Stx) (19), whereas the Slu pathogens have more genes for adhesion, in-

vasion and Extended-Spectrum Beta Lactamases (ESBLs) that possibly facilitate their extra-intestinal in-

fections and antibiotic-resistance (Fig.S5C,S5D). Next, we compare sequence variances between

orthologs shared by the two clades. In the core genome, 126 and 227 non-synonymous polymorphisms in

97 and 168 genes show specificity for the Vig and Slu clades, respectively, which scatter along the core

genome (Fig.S6A, Table S3). Half of the genes that contain these clade-associated variations (CAVs) are

in the COG category of metabolism, which also underpin the metabolic distinctions between the two

clades (Fig.S6B). Furthermore, homologues of the core genes involved in the metabolic pathways of

amino acids and carbohydrates show a clear separation in their maximal-likelihood phylogeny when con-

catenated (Fig. S7). Dispensable orthologs with high prevalence (present in > 80% of the 104 strains) also

show notable larger sequence distance in paired homologs between-clade than within-clade (Fig. S8).

These differences in sequence distance suggest that each clade utilizes their own preferred alleles of both

the core and dispensable genes, which are not readily traded between clades.

The obvious genetic differences between the clades suggest they exchange genetic material

scarcely. We thus infer genome-wide recombination for each genome pair to see if recombination de-

creases sharply and shows signals of genetic separation between the two clades. Taking advantage of

our complete genomes, we are able to infer recombination events by comparing synteny of aligned identi-

cal fragments. These fragments if moved to other locations are recognized as authentic recombinant frag-

ments. Our result shows that extent of intra-clade recombination is rather extensive than that of inter-clade

(Fig. 2A), in accordance with a previous report of less samples(20). For intra-clade genome pairs, the total

lengths of recombinant fragments range 0.3Mb – 2Mb, and in some cases even approaching half of the

genome, whereas those of inter-clade pairs are 10-fold shorter and never exceed 250kb (Fig. 2B). The

intensity of intra-clade recombination exhibit almost the same extent even between commensal and patho-

genic strains (Fig. 2B inlet), which implies that virulence genes can be freely transferred between com-

mensals and pathogens of the same clade and makes efforts to distinguish them extremely difficult (21).

Not only the intensity of recombination shows a clear cut-off between inter- and intra-clade recombination,

but also logarithmically correlated to phylogenetic distance based on core genome polymorphism (Fig.2C)

as previously estimated (22). Seemingly, the numbers of fragments that represent recombination fre-

quency are much greater in intra-clade pairs than inter-clade pairs (Fig. 2D) and correlate with phyloge-

netic distance as well (Fig. 2E), except in case of very close strains where recombination frequency can

be significantly underestimated due to the overlapping of recombinant fragments. The average size of re-

combinant fragments is also much longer in intra-clade genome pairs than what of inter-clade (Fig. 2F)

and correlate with phylogenetic distance (Fig. 2G). Since restriction-modification system has been demon-

strated to control recombination between species (23), we test its role in our samples with the DNA methylation information obtained from PacBio platform. However, the result shows no clade-specific restriction

ylation information obtained from PacBio platform. However, the result shows no clade-specific restriction

motif, although there are many strain-specific methylated motifs and a common motif of GATC in all *E. coli*

strains (Fig. S9).

The majority of recombinant fragments are in length of 3kb which fits in the common size of bacte-

rial operons, even some super-fragments can approach 2Mb (Fig. 3A) as long as reported in other species

(24, 25). As so efficient and frequent the recombination is in exchanging genetic materials especially

within clade, we further examine its influence on the organization and metabolic scheme of the recipient

genome. Firstly, we calculate the mobility of each single gene as its normalized frequency of being recom-

bined. The result indicates a broad range of gene mobility from 0 to 100% and a clear negative correlation

to prevalence, i.e., core genes which present in all strains are rather stable and seldom recombined,

whereas dispensable genes, especially those only present in few strains, are actively transferred between

strains (Fig. 3B). This is in accordance with the concept of Genome Organization Framework (GOF) (10),

where core genes constitute a stable genome framework, leaving the intervals filled with mobile variable

genes. Next, we examine the mobility of the clade-characteristic CAGs which show comparable overall

mobility as other dispensable genes of similar prevalence (Fig. 3B), but differ their mobility in the two

clades: Vig-CAGs are much more stable in Vig strains than in Slu strains, and those of Slu show opposite

mobility in the two clades (Fig. 3C). As for the core genes containing CAVs, they are all rarely recombined,

but still a little more stable in corresponding clades although the difference is not significant (Fig. 3D). The

stability of core genes as well as CAGs in corresponding clades suggest that the recombination has little

overall influences on genes with critical adaptive functions. Thus the extensive recombination does not

disturb the framework of genome and adaptive metabolic scheme heavily, possibly due to strong negative

selections for any changes that impair the fitness of host cell, according to the complexity hypothesis and

cellular evolution(26).

Discussion

Our study is based on a dataset of high-quality complete genomes as well as physiological features

experimentally acquired for the representative collection of *E. coli* strains. Two major clades of the popu-

lation, Vig and Slu, show distinct ecological and clinical features due to their physiological differences and

clade-featured metabolic profile. The phenotypes of the two clades are genetically underpinned by their

differences in genome content and alleles they used for orthologous which make them adaptive to respec-

tive ecological niches. The thorough inference of recombination events for each pair of genomes indicates

that recombination is rather frequent and intensive between within-clade strains but logarithmically re-

duced with phylogenetic distance, leading to genetic separation between clades. The extensive recombi-

nation, although keep cutting-and-pasting new genetic materials until they reach assortive genome loca-

tions, do not untidy GOFs and adaptive metabolic portfolios.

One of our findings is to quantitate the intensity of genetic recombination occurring within and

across the *E. coli* clades we have defined, which is not easy due to weak signals of recombined se-

quences intrinsic to draft assemblies (11, 27). For a long time, recombination has been acknowledged to

be very rare in asexual bacterial species, on which the theory of genome sweep has been established and

genes adjacent to advantageous alleles may achieve high prevalence through genetic hitchhiking. Recent

studies have questioned the theory and proposed bacteria as quasi-sexual populations where adaptive

genes instead of genome sweep the population driven by nature selection (6, 7). Our data, providing ex-

actly count on each recombination event for genome pairs, directly verify a high intra-clade recombination

rate and quasi-sexual status of microbial populations.

The poor inter-clade recombination rate and its small recombinant fragment size suggest a mech-

anistic barrier in favor of speciation (28). In simulations of evolution model, given suitable mutation rate

and enough population size, even without other involvement of nature selection or geographic separation,

populations can diverge neutrally only if recombination rate depends on homolog distance (similar to phy-

logeny distance in our work) (29). In other words, if individuals in a population recombine more intensively

with closer relatives, the disparity in recombination rate would by itself homogenize strains with similar ge-

nome sequences into a clade and build genetic barriers to others (28). As a result, the disparity of recom-

bination intensity may consistently generate new clades; some of them are deemed to be adaptive or spe-

cialized to ecological niches and selected by nature and others may just drift on and off the edge. This is

precisely the scenario of the Vig and Slu clade separation in our study, and such scenarios have been

published in other bacteria , archaea (7, 30-33) and eukaryotes (34). However, molecular mechanism of

the phylogeny-restricted recombination have yet to be described. Previous studies have proposed roles of

geographic isolation, diversity of CRISPR-Cas system (35), DNA uptake signal sequences (36), and in-

compatible transfer mechanisms due to pili (37) for the mechanisms. Recently, high-efficiency recombina-

tion mechanisms, such as co-evolving phage and distributive conjugal transfer have been reported, which

allows DNA fragments over 100kb in a single recombination event to happen but such a function deterio-

rates between remote relatives (20, 38). All these mechanisms with various transfer capacities and host

range enable strains to utilize suitable means in exchanging genetic material according to phylogenetic

distance of relatives.

The assessment of mobility for each single gene indicates that high recombination rate does not

have impact on genes with critical adaptive functions as well as GOF and primary metabolic portfolios they

built over long evolutionary time scale when genes or their alleles are constantly shuffled by the high-rate

recombination until reach assortive locations and fixed by nature selection in a manner of cut-and-try. Be-

sides niche adaptation, nature also selects for higher genetic compatibility among metabolic pathways,

genes, and even alleles, such as the CAVs in our case, because conflicts among these components may

undermine the competence of host strains in constant competitions with adjacent sister strains (39). Thus

GOF and primary metabolic portfolios and their related genes may be resistant to recombination that re-

sults in allele substitution(40) and location rearrangement(41) because phylogenies with reduced inte-

grated fitness may be eliminated by nature selections. Recent studies have shown genetic interaction and

compatibility may create fitness cost and constrain the dissemination and evolution of dispensable genes

(42, 43), which similarly direct the transfer of genes and alleles intra- and inter-clade to confer Vig and Slu

distinct ecological and clinical features. Thus recombination accelerates the optimization of the genome

content to streamlines physiological functions and metabolic portfolios of host strains. High-rate recombi-

nation is able to help adaptive genes to sweep the population, which benefit recipient strains with competi-

tive advantages and faster evolution.

In conclusion, our empirical data regarding to the separation of clades Vig and Slu in species *E.*

coli clarify the role of recombination in population diversification and evolution. The extensive intra-clade

recombination interrupts linkage between genes and leaves them independent under selections, but do

not disarranged the optimized GOF and metabolic scheme of host strains possibly due to critical selection

for streamlined physiological activities. Meanwhile, the extensive recombination within a clade coheres

strains into a quasi-sexual population, but its intensity declines sharply as phylogeny distance increases,

which builds barriers between clades and diverge the population.

Methods and Materials

Strains and MLST typing. A world-wide collection of 202 *E. coli* strains from vertebrate hosts (13) was kindly provided by Professor Shulin Liu (Genomics Research Center, Harbin Medical University, Harbin, China). A total of 149 successfully resuscitated strains were subjected to MLST sequencing and construction of a maximum-likelihood phylogeny using RaxML with model GTR and 1,000 bootstraps, together with 59 complete genomes of human-host deposited in NCBI (ftp://ftp.ncbi.nlm.nih.gov/genomes/Bacteria/) as of July 2013 and seven draft genomes published by a study of environmental strains (14) from ftp://ftp.ncbi.nlm.nih.gov/genomes/Bacteria_DRAFT/.

Genome sequencing, annotation, and phylogeny inference. The 45 selected representative strains were sequenced by using Hiseq 2000 and a paired-end sequencing protocol (2 × 100 bp run) at depth of 150×. Long reads are also added for contiguity from PacBio RS II with one or two cells per strain. After quality-filtration using the SMRT Analysis software suite (Pacific Biosciences Inc.), *de novo* assembly was performed according to the Hierarchical Genome Assembly Process (HGAP) workflow, and consensus sequences were corrected using Pilon (v1.22) with the clean reads generated from Illumina Hiseq 2000. Genomes were annotated with GeneMark.hmm and BLAST against the COG database and KEGG GENES (www.genome.jp/kegg/genes.html). Orthologs were identified by using PGAP (44), and the annotated genomes show apparent consistency when compared to the genome framework generated from GAAP (http://gaap.big.ac.cn), a synteny of core genes on genomic scale (45). Phylotypes of all strains were identified *in silicon* according to the presence of three phylotype-specific genes or fragments as previously described (46).

Measurement of growth rate, survival rate, and mobility. Refer to SI.

Genome content similarity, gene distribution, and homologue distance between the Vig and Slu clades. The Bray–Curtis dissimilarity index d was calculated for paired genomes as $1 - [2 * S_{ij}/(S_i + S_j)]$, where S_{ij} is the number of genes shared by strains i and j, and S_i, S_j as that in strains i and j, respectively. Two-sample Kolmogorov-Smirnov test was applied to select ortholog significantly enriched in Vig or Slu, when $p < 0.01$. Antibiotic-resistance and Shiga-like toxin genes were identified by BLAST against CBMAR database (http://14.139.227.92/mkumar/lactamasedb) and the protein sequences of the toxin from *E. coli* O157:H7 (lcl|AB015056.1_prot_BAA88123.1_1 for A-subunit and lcl|AB015056.1_prot_BAA88124.1_2 for B-subunit). Pairwise distance of orthologs was calculated by using Protdist in the Phylip package.

Metabolism reconstruction. The metabolic profile of each strain was reconstructed *in silico* using the genome-scale models (GSMs) as described previously (17). Briefly, we first removed the pseudogenes in annotation and used ModelSEED (http://modelseed.org/) to build the genome-scale models based on flux balance analysis (18). Here we used the condition of "gram negative template" and "complete media" which contains 16279 compounds in KBase (ftp://ftp.kbase.us/assets/KBase_Reference_Data/Biochemistry). A compound was predicted to be consumed or produced when corresponding transport, metabolism or biosynthesis reactions were available in a strain.

Recombination inference. To infer genome-wide recombination, we utilized gmos (Genome MOsaic Structure) to make local alignments between paired genomes and identify identical fragments over 600bp (47) as candidate recombinant fragments. We further used the syntenic information of these candidates which parsimoniously retain those changed locations as authentic recombination fragments. Gene mobility was calculated as the percentage of being recombined in total genome pairs the gene was present. The recombination of a gene was also identified using the syntenic information as what for recombinant fragments.

Data Availability. The genome sequences we contribute are available on the NCBI GenBank (accession No. CP012760, CP012781, CP029328, CP029327, CP028732-CP028772) and Genome Sequence Archive of GSA database (http://gsa.big.ac.cn, Accession No. GWHAACT- GWHAAEJ). Raw data of sequencing are deposited in the GSA database (PacBio, Accession No. CRR028990-CRR029033; Hiseq, Accession No. CRR019978-CRR020022).

Acknowledgements

Competing interests: The authors declare no competing interests.

Funding: This work was supported by the National Natural Scientific Foundation of China [31470180, 31471237, 31671350and 30971610], the National Key Research and Development Program of China [2016YFC0903800], Programs of Ministry of Health of the People's of China [201402018], Programs of the Chinese Academy of Sciences [QYZDY-SSW-SMC017, YZ201568, YZ201402], Programs of Beijing Municipal Science and Technology Project [Z171100001317011], Taicang Municipal Science and Technology Project [TC2016SF08].

Author contribution: J.Y. and Y.K. conceived the project and led the writing; L.Y., Y.C., R.F., and J. W.

compiled the data; and X.S., F.C., Z.H., Z.G., X.J., Q.L, Q.M., J.W., J.X., and S.H. analyzed the data. All

authors contributed to the writing and/or intellectual development of the manuscript.

References

1.   Fraser C, Hanage WP, & Spratt BG (2007) Recombination and the nature of bacterial
speciation. *Science (New York, N.Y.)* 315(5811):476-480.
2.   Shapiro BJ, Leducq JB, & Mallet J (2016) What Is Speciation? *PLoS genetics*
12(3):e1005860.
3.   Cohan FM (2001) Bacterial species and speciation. *Systematic biology* 50(4):513-524.
4.   Cohan FM & Perry EB (2007) A systematics for discovering the fundamental units of
bacterial diversity. *Current biology : CB* 17(10):R373-386.
5.   Bendall ML*, et al.* (2016) Genome-wide selective sweeps and gene-specific sweeps in
natural bacterial populations. *The ISME journal* 10(7):1589-1601.
6.   Rosen MJ, Davison M, Bhaya D, & Fisher DS (2015) Microbial diversity. Fine-scale
diversity and extensive recombination in a quasisexual bacterial population occupying a
broad niche. *Science (New York, N.Y.)* 348(6238):1019-1023.
7.   Shapiro BJ*, et al.* (2012) Population genomics of early events in the ecological
differentiation of bacteria. *Science (New York, N.Y.)* 336(6077):48-51.
8.   Doolittle WF (2012) Population genomics: how bacterial species form and why they don't
exist. *Current biology : CB* 22(11):R451-453.
9.   Fraser C, Alm EJ, Polz MF, Spratt BG, & Hanage WP (2009) The bacterial species
challenge: making sense of genetic and ecological diversity. *Science (New York, N.Y.)*
323(5915):741-746.
10.   Kang Y*, et al.* (2014) Flexibility and symmetry of prokaryotic genome rearrangement
reveal lineage-associated core-gene-defined genome organizational frameworks. *MBio*
5(6):e01867.
11.   Ravenhall M, Skunca N, Lassalle F, & Dessimoz C (2015) Inferring horizontal gene
transfer. *PLoS computational biology* 11(5):e1004095.
12.   Martinez-Castillo A & Muniesa M (2014) Implications of free Shiga toxin-converting
bacteriophages occurring outside bacteria for the evolution and the detection of Shiga
toxin-producing Escherichia coli. *Frontiers in cellular and infection microbiology* 4:46.
13.   Souza V, Rocha M, Valera A, & Eguiarte LE (1999) Genetic structure of natural
populations of Escherichia coli in wild hosts on different continents. *Applied and*
environmental microbiology 65(8):3373-3385.
14.   Luo C*, et al.* (2011) Genome sequencing of environmental Escherichia coli expands
understanding of the ecology and speciation of the model bacterial species. *Proceedings*
of the National Academy of Sciences of the United States of America 108(17):7200-
7205.
15.   Didelot X, Meric G, Falush D, & Darling AE (2012) Impact of homologous and non-
homologous recombination in the genomic evolution of Escherichia coli. *BMC genomics*
13:256.

16.   Kaas RS, Friis C, Ussery DW, & Aarestrup FM (2012) Estimating variation within the
genes and inferring the phylogeny of 186 sequenced diverse Escherichia coli genomes.
BMC genomics 13:577.
17.   Monk JM, *et al.* (2013) Genome-scale metabolic reconstructions of multiple Escherichia
coli strains highlight strain-specific adaptations to nutritional environments. *Proceedings*
of the National Academy of Sciences of the United States of America 110(50):20338-
20343.
18.   Henry CS, *et al.* (2010) High-throughput generation, optimization and analysis of
genome-scale metabolic models. *Nature biotechnology* 28(9):977-982.
19.   Tarr PI, Gordon CA, & Chandler WL (2005) Shiga-toxin-producing Escherichia coli and
haemolytic uraemic syndrome. *Lancet* 365(9464):1073-1086.
20.   Dixit PD, Pang TY, Studier FW, & Maslov S (2015) Recombinant transfer in the basic
genome of Escherichia coli. *Proceedings of the National Academy of Sciences of the*
United States of America 112(29):9070-9075.
21.   Picard B, *et al.* (1999) The link between phylogeny and virulence in Escherichia coli
extraintestinal infection. *Infection and immunity* 67(2):546-553.
22.   Roberts MS & Cohan FM (1993) The effect of DNA sequence divergence on sexual
isolation in Bacillus. *Genetics* 134(2):401-408.
23.   Pleska M, *et al.* (2016) Bacterial Autoimmunity Due to a Restriction-Modification System.
Current biology : CB 26(3):404-409.
24.   Boritsch EC, *et al.* (2016) Key experimental evidence of chromosomal DNA transfer
among selected tuberculosis-causing mycobacteria. *Proceedings of the National*
Academy of Sciences of the United States of America 113(35):9876-9881.
25.   Chen L, Mathema B, Pitout JD, DeLeo FR, & Kreiswirth BN (2014) Epidemic Klebsiella
pneumoniae ST258 is a hybrid strain. *mBio* 5(3):e01355-01314.
26.   Woese CR (2002) On the evolution of cells. *Proceedings of the National Academy of*
Sciences of the United States of America 99(13):8742-8747.
27.   Nielsen KM, Bohn T, & Townsend JP (2014) Detecting rare gene transfer events in
bacterial populations. *Frontiers in microbiology* 4:415.
28.   Dixit PD, Pang TY, & Maslov S (2017) Recombination-Driven Genome Evolution and
Stability of Bacterial Species. *Genetics* 207(1):281-295.
29.   Falush D, *et al.* (2006) Mismatch induced speciation in Salmonella: model and data.
Philosophical transactions of the Royal Society of London. Series B, Biological sciences
361(1475):2045-2053.
30.   Didelot X, *et al.* (2011) Recombination and population structure in Salmonella enterica.
PLoS genetics 7(7):e1002191.
31.   Huang CL, *et al.* (2015) Ecological genomics in Xanthomonas: the nature of genetic
adaptation with homologous recombination and host shifts. *BMC genomics* 16:188.
32.   Zwick ME, *et al.* (2012) Genomic characterization of the Bacillus cereus sensu lato
species: backdrop to the evolution of Bacillus anthracis. *Genome research* 22(8):1512-
1524.
33.   Cadillo-Quiroz H, *et al.* (2012) Patterns of gene flow define species of thermophilic
Archaea. *PLoS biology* 10(2):e1001265.
34.   Ellison CE, *et al.* (2011) Population genomics and local adaptation in wild isolates of a
model microbial eukaryote. *Proceedings of the National Academy of Sciences of the*
United States of America 108(7):2831-2836.

35.   Samson JE, Magadan AH, & Moineau S (2015) The CRISPR-Cas Immune System and
405   Genetic Transfers: Reaching an Equilibrium. *Microbiology spectrum* 3(1):Plas-0034-
406   2014.
407   36.   Frye SA, Nilsen M, Tonjum T, & Ambur OH (2013) Dialects of the DNA uptake sequence
408   in Neisseriaceae. *PLoS genetics* 9(4):e1003458.
409   37.   Cehovin A*, et al.* (2013) Specific DNA recognition mediated by a type IV pilin.
410   *Proceedings of the National Academy of Sciences of the United States of America*
411   110(8):3065-3070.
412   38.   Gray TA, Krywy JA, Harold J, Palumbo MJ, & Derbyshire KM (2013) Distributive conjugal
413   transfer in mycobacteria generates progeny with meiotic-like genome-wide mosaicism,
414   allowing mapping of a mating identity locus. *PLoS biology* 11(7):e1001602.
415   39.   Bao YJ, Shapiro BJ, Lee SW, Ploplis VA, & Castellino FJ (2016) Phenotypic
416   differentiation of Streptococcus pyogenes populations is induced by recombination-driven
417   gene-specific sweeps. *Scientific reports* 6:36644.
418   40.   Kacar B, Garmendia E, Tuncbag N, Andersson DI, & Hughes D (2017) Functional
419   Constraints on Replacing an Essential Gene with Its Ancient and Modern Homologs.
420   *MBio* 8(4).
421   41.   Ma Q*, et al.* (2013) Computational analyses of transcriptomic data reveal the dynamic
422   organization of the Escherichia coli chromosome under different conditions. *Nucleic acids*
423   *research* 41(11):5594-5603.
424   42.   San Millan A, Toll-Riera M, Qi Q, & MacLean RC (2015) Interactions between
425   horizontally acquired genes create a fitness cost in Pseudomonas aeruginosa. *Nature*
426   *communications* 6:6845.
427   43.   Porse A, Schou TS, Munck C, Ellabaan MMH, & Sommer MOA (2018) Biochemical
428   mechanisms determine the functional compatibility of heterologous genes. *Nature*
429   *communications* 9(1):522.
430   44.   Zhao Y*, et al.* (2012) PGAP: pan-genomes analysis pipeline. *Bioinformatics* 28(3):416-
431   418.
432   45.   Yuan L*, et al.* (2017) GAAP: Genome-organization-framework-Assisted Assembly
433   Pipeline for prokaryotic genomes. *BMC genomics* 18(Suppl 1):952.
434   46.   Clermont O, Bonacorsi S, & Bingen E (2000) Rapid and simple determination of the
435   Escherichia coli phylogenetic group. *Applied and environmental microbiology*
436   66(10):4555-4558.
437   47.   Domazet-Loso M & Domazet-Loso T (2016) gmos: Rapid Detection of Genome
438   Mosaicism over Short Evolutionary Distances. *PloS one* 11(11):e0166602.
439
440
441

Figures and legends

Figure 1. The two major *E. coli* clades: Vig and Slu.

(A) An ML phylogeny based on the core-genome derived polymorphisms that separate host-related strains from the environmental strains (free-living in the tidal zone) and host-related strains into two major clades and a few minor clades. Names of each strain are labelled. Clade (colored leaf), phylotype (colored branch), origin (colored inner stripe), and pathotype (colored outer stripe) are indicated for each *E. coli* strain. (B) Prediction of metabolic compounds each strain uptakes and produces based on genome-scale models (GSMs) and genome annotations for the two clades. The cluster diagram represents 82 metabolic compounds with quantitative differences among strains, and those with significant inter-clade differences ($p < 0.01$, Wilcoxon test) are highlighted in red boxes. Each column represents a single strain with name labeled in red or blue color for the Vig or Slu, respectively. The top clustering is based on the Euclidean distance between strains and Ward clustering algorithms. Compound flux is normalized by its maximal value and colored in red for consuming and blue for producing (see the right scaling bar).

Figure 2. Intra- and inter-clade Recombinations.

(A) Pair-wise assessment of recombination within and across clades. Strains are circularly displayed

and the clades of Vig (red) and Slu (blue) are on both sides and the minor clade is placed (green) in

the middle. Pathogenic and commensal strains are indicated in dark and light colors. Each line be-

tween strains indicates recombination between the two strains with thickness proportional to the total

fragment length over 5kb. (B) Recombinant length distributions. Recombinants are grouped into intra-

clade (Vig–Vig, red; Slu–Slu, blue) and inter-clade (Vig–Slu, purple), and those between commen-

sal/pathogenic pairs (dotted line) are also shown in the insets. (C) Recombinant length is plotted as a

function of phylogenetic distance. (D) A plot of recombination frequency (number of recombinant frag-

ment). (E) Recombination frequency as a function of phylogenetic distance. (F) Average recombinant

length distribution. (G) Average recombinant length as a function of phylogenetic distance.

Figure 3. Gene mobility.

(A) Recombinant size distributions. Recombinants of Vig intra-clade (Vig–Vig), Slu intra-clade (Slu–

Slu), and inter-clade (Vig–Slu) are plotted as a function of size (bp) in red, blue, and purple, respec-

tively. (B) Gene mobility (migration flexibility among genomes within and across clades) as a function

of prevalence (the number of genomes that contain the gene). The regression curve (solid line) is

based on generalized additive model with the shadow representing 95% confidence interval. CAGs

(clade-associated genes) of Vig and Slu are indicated with red and blue solid circles, respectively. (C)

Mobility of CAGs in Vig (red) and Slu (blue). (D) Mobility of CAV (clade-associated variations)-contain-

ing genes in Vig (red) and Slu (blue).