1   **The nuclear and mitochondrial genomes of the facultatively eusocial orchid**
2   **bee *Euglossa dilemma.***
3
4   **Authors:**
5

6   Philipp Brand*,†, Nicholas Saleh*,†, Hailin Pan‡, Cai Li‡, Karen M. Kapheim§, Santiago
7   R. Ramírez*
8
9   **Affiliations:**
10
11 * Department for Evolution and Ecology, Center for Population Biology, University
12 of California, Davis, California 95616
13
14 † Graduate Group in Population Biology, University of California, Davis, California
15 95616
16
17 ‡ China National Genebank, BGI-Shenzhen, Shenzhen, 518083, China
18
19 § Department of Biology, Utah State University, Logan, Utah 84322
20

21 **Data availability**
22
23 The *E. dilemma* genome assembly *Edil_v1.0*, the annotation, and the original gene set
24 *Edil_OGS_v1.0* are available for download via NCBI [XXX], Beebase [XXX], the i5k NAL
25 workspace [xxx], and the Ramirez Lab website [URL]. The raw reads are available at
26 the NCBI Sequence Read Archive [XXX]. The published raw transcriptome sequence
27 reads are available at the NCBI Sequence Read Archive [SRA: SRX765918].

47 **Running Title:**
48
49 The genomes of the orchid bee *Euglossa dilemma*
50
51 **Key Words:**
52
53 whole-genome assembly; corbiculate bee; orchid bee; invasive species;
54 mitochondrial genome
55

56 **Corresponding Author:**
57
58 Philipp Brand
59 Department of Evolution & Ecology
60 University of California, Davis
61 One Shields Ave
62 Davis, CA 95616
63
64 Office: (530) 752-7614
65 pbrand@ucdavis.edu

Abstract

Bees provide indispensable pollination services to both agricultural crops and wild plant populations, and several species of bees have become important models for the study of learning and memory, plant-insect interactions and social behavior. Orchid bees (Apidae: Euglossini) are especially important to the fields of pollination ecology, evolution, and species conservation. Here we report the nuclear and mitochondrial genome sequences of the orchid bee *Euglossa dilemma*. *Euglossa dilemma* was selected because it is widely distributed, highly abundant, and it was recently naturalized in the southeastern United States. We provide a high-quality assembly of the 3.3 giga-base genome, and an official gene set of 15,904 gene annotations. We find high conservation of gene synteny with the closely related honey bee. This genomic resource represents the first draft genome of the orchid bee genus *Euglossa,* and the first draft orchid bee mitochondrial genome, thus representing a valuable resource to the research community.

Introduction

Bees (Apoidea) are important models for the study of learning and memory (Menzel and Muller 1996), plant-insect interactions (Doetterl and Vereecken 2010) and the evolution of social behavior (Nowak *et al.* 2010; Woodard *et al.* 2011; Kapheim *et al.* 2015). Among the >20,000 bee species worldwide, lineages have evolved varied degrees of specialization on floral resources such as pollen, resins, and oils (Wcislo and Cane 2003; Michener 2007; Litman *et al.* 2011). These relationships are wide-ranging and have substantial impact on the health and function of natural and agricultural systems (Klein *et al.* 2007). Furthermore, several transitions from an ancestral solitary to a derived eusocial behavior have occurred within bees (Danforth 2002; Cardinal and Danforth 2011; Branstetter *et al.* 2017). Thus, bees provide unique opportunities to investigate the genetic underpinnings of multiple major ecological and evolutionary transitions. The repeated evolution of different behavioral phenotypes in bees, including foraging and social behavior, provides a natural experiment that allows for the determination of general as well as species-specific molecular genomic changes underlying phenotypic transitions. In order to capitalize on this potential, whole-genome sequences of a divergent array of bee species with different life histories are needed (Kapheim *et al.* 2015).

Orchid bees (Apidae; Euglossini) are among the most important pollinators of thousands of diverse neotropical plant species (Ramírez *et al.* 2002). While female orchid bees collect nectar, pollen and resin for nest construction and provisioning, male bees collect perfume compounds from floral and non-floral sources (Vogel 1966; Whitten *et al.* 1993; Eltz *et al.* 1999; Roubik and Hanson 2004). These volatile compounds are used to concoct a species-specific perfume blend that is subsequently used during courtship, presumably to attract conspecific females. This unique male scent-collecting behavior has recently been examined in a broad array of molecular ecological and evolutionary studies, focusing on phenotypic evolution,

138 chemical communication, plant-insect mutualisms, and speciation (Eltz *et al.* 2008;
139 2011; Ramírez *et al.* 2011; Brand *et al.* 2015).
140
While most of the approximately 220 species of orchid bees appear to be solitary,
several species have transitioned to living in coordinated social groups (Garófalo
1985; Pech *et al.* 2008; Augusto and Garófalo 2009). Female *Euglossa dilemma*
individuals, for example, can either form solitary nests and provision their own
brood cells, or live in small groups where daughters remain in their natal nest and
help their mother rear her offspring, instead of dispersing to found their own nest.
The social *E. dilemma* nests (similar to the closely related *E. viridissima;* Pech *et al.*
2008) exhibit true division of labor, with subordinate daughters foraging for
resources and the reproductively dominant mothers laying eggs (Saleh & Ramirez
pers. obs.). This facultative eusocial behavior represents an early stage in social
evolution and makes *E. dilemma* well-suited for studying the genomic changes
underlying the transition from solitary to eusocial behavior. While other facultative
eusocial species evolved throughout the bee lineage, orchid bees have a unique
taxonomic position (Cardinal and Danforth 2011). Orchid bees are part of the
corbiculate bees, together with the honey bees, bumblebees and stingless bees,
three obligately eusocial bee lineages (Figure 1a). As the sister group to all other
corbiculate bee lineages (Romiguier *et al.* 2015; Peters *et al.* 2017; Branstetter *et al.*
2017), orchid bees may provide key insights into the early stages of eusociality and
the possible evolutionary trajectories that led to the obligate eusocial behavior
exhibited by honey bees.
161
Here we present the draft genome of the orchid bee species *Euglossa dilemma*. Using
a combined paired-end and mate-pair library sequencing approach, we assembled
18% of the predicted 3.2Gb genome, and annotated a high-quality gene set including
15,904 genes. In addition, we reconstructed three quarters of the mitochondrial
genome with the help of transcriptome data, representing the first orchid bee
mitogenome. These genomic resources will facilitate the genetic study of
outstanding ecological and evolutionary questions, such as the evolution of resource
preferences and the evolution of eusociality. Moreover, it provides an important
genomic resource for an endangered group of crucial neotropical pollinators, that
are of specific concern for conservation biologists (Zimmermann *et al.* 2011; Suni
and Brosi 2012; Suni 2016; Soro *et al.* 2016).
173
174 **Materials and Methods**
175
176 **Genome sequencing and assembly**
177
Nuclear Genome. Sequencing of the *E. dilemma* genome was based on six haploid
male individuals collected at Fern Forest Nature Center in Broward County, FL in
February 2011. This population was chosen due to its low nucleotide diversity
resulting from a bottleneck during a single introduction to Southern Florida about
15 years ago (Skov and Wiley 2005; Pemberton and Wheeler 2006; Zimmermann *et*
al. 2011). DNA was extracted from each bee independently and used for the

construction of four paired-end (two 170bp and 500bp libraries, respectively) and
four mate-pair (two 2kb and 5kb libraries, respectively) sequencing libraries. Next,
the paired-end libraries were sequenced in 90 cycles and the mate-pair libraries for
49 cycles on an Illumina HiSeq2000. The resulting sequence data was run through
fastuniq v1.1 (Xu *et al.* 2012) to remove PCR duplicates and quality trimmed using
trim_galore v0.3.7 (Babraham Bioinformatics). Subsequently, reads were used for *de*
novo assembly with ALLPATHS-LG v51750 (Gnerre *et al.* 2011) and Soap-denovo2
(Luo *et al.* 2012) with varying settings. Gaps within scaffolds were closed using
GapCloser v1.12 (Luo *et al.* 2012) for each assembly. ALLPATHS-LG with default
settings resulted in the highest-quality assembly, based on assessments of
annotation completeness (see below). This assembly (*E. dilemma* genome assembly
v1.0) was used for all subsequent analyses. All other assemblies were excluded from
analysis, but are available upon request.
197
The pre-processed reads were used for k-mer based genome size estimates. We
used ALLPATHS-LG to produce and analyze the k-mer frequency spectrum (k=25).
Genome size was estimated on the basis of the consecutive length of all reads
divided by the overall sequencing depth as $(N \times (L - K + 1) - B)/D = G$, where N is
the total number of reads, L is the single-read length, K is the k-mer length, B is the
total count of low-frequency (frequency ≤3) k-mers that are most likely due to
sequencing errors, D is the k-mer depth estimated from the k-mer frequency
spectrum, and G is the genome size. In addition, we used the ALLPATHS-LG k-mer
frequency spectrum to predict the repetitive fraction of the genome.
207
208 The quality of the genome assembly was assessed using standard N statistics, and
209 assembly completeness as measured by the CEGMA v2.5 (Parra *et al.* 2007) and
210 BUSCO v1.1 (Simão *et al.* 2015) pipelines. CEGMA was run in default mode, whereas
211 BUSCO was run with the arthropoda_odb9 OrthoDB database (Zdobnov *et al.* 2017)
212 in genome mode.
213
214 We estimated the mean per-base genome coverage on the basis of the pre-processed
215 reads and the estimated genome size as $\frac{\sum_{i=1}^{4}(R_i * L_i)}{G} = C$, where R is the number of
216 reads and L the mean read length of sequence library i, G is the estimated genome
217 size and C the resulting per-base coverage.
218
Mitogenome. Initial attempts to reconstruct the mitochondrial genome from our
whole-genome shotgun sequencing reads were only partially successful, due to high
sequence variability of sequencing reads with similarity to mitochondrial loci (data
not shown). In addition, we have observed that the amplification of mitochondrial
DNA in standard polymerase chain reactions (PCR) leads to a high level of
polymorphic sites in *E. dilemma* and other orchid bees (Brand & Ramírez pers. obs.).
Together, this suggests the presence of nuclear copies of the mitochondrial genome
(NUMTs) that interfere with the assembly process and PCR amplification. Therefore,
we used available *E. dilemma* transcriptome assemblies in order to reconstruct the
mitochondrial genome from cDNA (Brand *et al.* 2015). In order to find

mitochondrial genes in the transcriptome assembly of Brand et al. 2015, we used
blastx with the honey bee mitochondrial genome as query (Crozier and Crozier
1993) and an E-value cutoff of 10E-12 (Altschul *et al.* 1990; Camacho *et al.* 2009).
The contigs and scaffolds that were detected with this approach were annotated
following Dietz *et al.* 2015. Briefly, we performed tblastn and tblastx searches with
protein coding genes and rRNA genes of the honey bee mitochondrial genome,
respectively. All hits were used for manual gene annotation using Geneious v8.0.5
(Biomatters Ltd. 2012). Since the recovered mitochondrial mRNA scaffolds
contained more than one gene, we searched and annotated intergenic tRNAs using
ARWEN 1.2.3 (Laslett and Canbäck 2008) and tRNAscan-SE 1.21 (Lowe and Eddy
1997).
240
241 **Genome annotation**
242
Gene annotation. Genes were annotated based on sequence homology and *de novo*
gene predictions. The homology approach was based on the recently updated high-
quality official gene set of the honey bee (OGS v3.2; Elsik *et al.* 2014). All honey bee
original gene set (OGS) proteins were used in initial tblastn searches against all *E.*
dilemma scaffolds with an E-value cutoff of 10E-4. Proteins with a hit covering ≥
50% of the honeybee protein query were selected for accurate exon-intron
boundary prediction for each scaffold using exonerate v2.42.1 (Slater and Birney
2005) with the minimum fraction of the possible optimal similarity per query set to
35%. In a second round, genes not annotated under the previous settings were
rerun with minimum similarity set to 15%. In the case of multiple annotations, we
discarded all but one annotation with the best hit to the honeybee OGS (based on
completeness and similarity). This approach proved feasible due to the close
relatedness of *E. dilemma* and the honey bee. For *de novo* gene prediction we used
Augustus (Stanke *et al.* 2008) and SNAP (Korf 2004) trained on the honey bee, with
the *E. dilemma* genome masked for repetitive regions (See below) as input. Only
genes predicted by both programs were taken into account. Gene predictions with
≥85% sequence similarity to each other were discarded, to prevent the inclusion of
putative unmasked transposable element derived genes in the official gene set. *De*
novo predictions were added to the *E. dilemma* OGS if not annotated by the
homology-based approach.
263
264 *Repetitive element annotation.* Repetitive elements including tandem repeats,
265 nuclear copies of the mitochondrial genome (NUMTs), and transposable elements
266 (TEs) were annotated using multiple methods.
267
268 *Tandem repeats.* We searched for micro- and mini-satellites (1–6 bp and 7–1000 bp
269 motif length, respectively) in all scaffolds using Phobos 3.3.12 (Mayer 2010). We
270 performed two independent runs for each class of tandem repeats with Phobos
271 parameter settings following Leese et al. 2012 (gap score and mismatch score set to
272 -4 and a minimum repeat score of 12; Leese *et al.* 2012).
273

274 *NUMTs*. We annotated NUMTs using blastn runs with the partial mitochondrial
275 genome (see above) as query and an E-value cutoff of 10E-4 as used in NUMT
276 analyses of other insect genomes (Pamilo *et al.* 2007). This approach allowed us to
277 find NUMTs with medium to high similarity to the actual transcriptome-based
278 mitochondrial genome.
279
280 *TEs*. In order to annotate TEs, we used Repeatmasker v4.0.5 (Smit et al. 2016) with
281 Crossmatch v. 0.990329 as search engine in sensitive mode. The Repbase
282 invertebrate database v21.12 (Jurka 2000; Bao *et al.* 2015) was used as TE
283 reference, due to the lack of a bee-specific database.
284
285 **Genome structure**
286
To analyze genome structure, we compared the genome wide gene synteny of *E.*
dilemma and the honey bee. We used the genomic locations of homologous genes (as
determined above) of the honey bee and *E. dilemma* scaffolds of at least 100kb
length to build haplotype blocks with a minimum length of 1kb. Haplotype blocks
included the entire gene span as well as intergenic regions whenever two or more
adjacent genes were homologous in both species. We discarded gene annotations
from downstream analysis that were recovered as homologous to multiple genomic
locations in either species. Furthermore, we excluded *E. dilemma* genes that were
recovered as homologous to honey bee scaffolds belonging to unknown linkage
groups.
297
298 **Data availability**
299
The *E. dilemma* genome assembly *Edil_v1.0*, the annotation, and the original gene set
Edil_OGSv1.0 are available for download via NCBI [XXX], Beebase [XXX] (Elsik *et al.*
2016), the i5k NAL workspace [xxx] (i5K Consortium 2013), and the Ramirez Lab
website [URL]. The raw reads are available at the NCBI Sequence Read Archive
[XXX]. The published raw transcriptome sequence reads are available at the NCBI
Sequence Read Archive [SRA: SRX765918] (Brand *et al.* 2015).
306
307 **Results and Discussion**
308
309 **Whole-genome assembly**
310
The *E. dilemma* genome assembly resulted in 22,698 scaffolds with an N50 scaffold
length of 144Kb and a total length of 588Mb (Table 1). This represents 18% of the k-
mer based estimated genome size of 3.2Gb. Of all sequence reads, 68% aligned to
the genome assembly, of which 56% aligned more than once. Further, the k-mer
frequency spectrum based on all sequencing reads was strongly positively skewed
indicating the presence of highly repetitive sequences in the read set (Figure 1b).
Based on the k-mer frequency spectrum, 87.7% of the genome was estimated to be

repetitive. This suggests that the genome of *E. dilemma* consists largely of highly
repetitive sequences, explaining the low consecutive assembly length and the high
assembly fragmentation. The mean per-base coverage was estimated to be
comparatively low in comparison to previous bee genome assemblies, with 19.7x
based on the pre-processed reads and estimated genome size (Kocher *et al.* 2013;
Kapheim *et al.* 2015). Total genomic GC content was 39.9%, and thus similar to
previously sequenced bee genomes ranging between 32.7% and 41.5% (Table 1)
(Kocher *et al.* 2013; Elsik *et al.* 2014; Kapheim *et al.* 2015).
326
Despite the fragmentation of the genome assembly representing less than 20% of
the estimated genome size, CEGMA analysis revealed complete assemblies of 231
out of 248 core eukaryotic genes (93.2% completeness). Similarly, BUSCO analysis
revealed that 1007 out of 1066 highly conserved arthropod genes were completely
assembled (94.4% completeness). Our gene prediction approach generated a
comprehensive official gene set including 15,904 protein-coding genes (Table 1). Of
these gene models, 11,139 were derived from homology-based predictions,
representing 73% of the 15,314 honey bee genes used for annotation. These
annotations are well within or exceeding previous bee genome assemblies, and are
similar to those reported for the other orchid bee genome available (Table 1)
(Kocher *et al.* 2013; Elsik *et al.* 2014; Park *et al.* 2015; Sadd *et al.* 2015; Kapheim *et*
al. 2015).
339
The CEGMA and BUSCO analysis and the gene annotation results suggest that the
gene-coding fraction of the *E. dilemma* genome was properly assembled, despite the
large estimated genome size and comparatively low per-base sequencing coverage.
Fragmentation of the assembly is thus likely to be primarily the result of highly
repetitive genomic elements, and less the result of low coverage. Overall, the results
suggest that our approach was sufficient to produce a high quality official gene set.
Due to the chosen homology-based approach, the majority of annotated genes in the
official gene set has known homology to honey bee genes (Table S1), which will
greatly facilitate genome-wide expression studies including gene ontology analyses
and comparative gene set analyses among insects.
350
Mitochondrial Genome assembly
The recently published transcriptome assembly used for the reconstruction of the
mitochondrial genome contained four scaffolds between 1,222bp and 4,188bp long
with a total consecutive length of 11,128bp (Figure 2). This corresponds to about
75% of the estimated length of the mitochondrial genome, based on other
corbiculate bee species (Crozier and Crozier 1993; Cha *et al.* 2007). The *E. dilemma*
mitogenome fragments contained 5 out of 22 tRNAs, 11 out of 13 protein coding
genes of which two were only partially recovered, and the 16S rRNA gene. Within
scaffolds all genes showed the known hymenopteran gene order and orientation,
while the orientation of the 5 tRNAs detected was identical to those in the honey bee
(Crozier and Crozier 1993; Cha *et al.* 2007). Attempts to complete the mitochondrial
genome using the nuclear genome assembly yielded no improvement of the
assembly (data not shown).

364
The high success in mitochondrial gene reconstruction is likely due to the nature of
the analyzed transcriptome data. Short intergenic regions as well as polycistronic
mitochondrial mRNA likely lead to the assembly of multiple genes into single
scaffolds. The A-T rich region is completely missing as well as the ND2 and 12S
rRNA genes flanking the region in insect mitogenomes. This unrecovered region also
contains a high number of tRNAs in the honeybee, which could explain the low
number of recovered tRNAs in *E. dilemma*. While the partial mitochondrial genome
assembly is only 75% complete, it represents the first mitogenome for the group of
orchid bees and will thus be a valuable resource for future phylogenetic analyses
within the lineage and between more distantly related bee taxa.
375
376   **Repetitive elements**
377
378   *Tandem Repeats*. We detected 76,001 microsatellite loci with a consecutive length of
379   2,291,067 bp. Minisatellites with motif lengths from 7bp to 1000bp were less
380   numerous in the genome (67,323 loci), totaling 13,343,515bp. Accordingly, tandem
381   repeats represent 3.86% of the genome assembly, suggesting that they contribute
382   only a small proportion to the overall genome size (Figure 1c).
383
NUMTs. We detected fragments with similarity to the draft mitochondrial genome
on 129 scaffolds totaling a length of 150,670 bp. The fragments had a mean length of
764.8 bp and a mean similarity of 91.5% to the mitogenome. This suggests that
these fragments are not derived from the mitochondrial genome and represent
actual NUMTs. A total of 39 scaffolds carried multiple fragments with high similarity
to the mitogenome with a concatenated length of up to 6566bp, suggesting that
respective NUMTs might have originated from larger fragments of the mitogenome.
In total, only 0.04% of the whole-genome assembly had hits to the mitogenome
(Figure 1c). This is likely an underestimate, due to the incompleteness of the
reconstructed mitochondrial genome. Nevertheless, NUMTs likely represent only a
small fraction of the whole nuclear *E. dilemma* genome. Previous analyses have
shown a high density of NUMTs in the honey bee in comparison to other insect
genomes totaling about 0.1% of the overall genome size (Pamilo *et al.* 2007).
Accordingly, given the NUMT content detected in *E. dilemma*, it is possible that a
comparatively high NUMT density is a common feature of corbiculate bee genomes.
399
TEs. In our RepeatMasker analysis we detected 47,553 interspersed repeats with a
cumulative length of 7,747,824bp. This corresponds to 1.82% of the genome
assembly (Figure 1c). This low number is surprising, since large genome sizes as in
E. dilemma have been mainly associated with TE activity and content in genomes
from unicellular eukaryotes to complex multicellular organisms including plants,
invertebrates and vertebrates (Kidwell 2002). However, TEs are fast evolving and
highly specific to their host lineages, which leads to large underestimates of genomic
TE content in previously unstudied lineages (Chalopin *et al.* 2015; Platt *et al.* 2016).
The only bee repeat content included in the Repbase database used for TE
annotation is the honey bee, a species with a comparatively small genome (0.23Gb)

and low TE diversity and content (Weinstock *et al.* 2006; Kapheim *et al.* 2015). Thus, the low percentage of TEs detected in the *E. dilemma* genome is very likely an underestimate of the actual density. *De novo* TE reconstructions using the genomic resources presented here should be performed in the future to provide a better estimate of the actual TE density in the *E. dilemma* genome.

Genome structure

Of the 22,698 *E. dilemma* scaffolds, 580 were at least 100kb in length and used for synteny analysis with the honey bee genome. A total of 356 of these scaffolds carried at least one gene annotation with known homology to the honey bee, and 329 of these *E. dilemma* scaffolds were homologous to honey bee scaffolds with known linkage group (LG) association (Table S1). Of these scaffolds, 272 (83%) showed ≥95% syntenic homology to a single honey bee LG (Figure 1d). Overall, the detected syntenic linkage blocks cover 222MB of scaffold length with homology to the honeybee, representing 85% of the 329 filtered scaffolds. Mean syntenic block length is 206,033bp for *E. dilemma* (min: 1017bp; max: 4,458,123bp; median: 47,154bp) and 143,221.4bp for the honey bee (min: 1029bp; max: 2,205,566bp; median: 41,726bp). This suggests that the genomic architecture is very similar between *E. dilemma* and the honey bee, representing a high level of conservation during the 80 million years since the two lineages diverged. Further, our results support a recent comparative analysis of the honey bee and the bumblebee genomes, which revealed high conservation of genomic synteny (Stolle *et al.* 2011). Together, these results support a general pattern of surprisingly slow genome evolution in gene coding regions in corbiculate bees, independent of the fraction of repetitive genome content.

Conclusion

The genome assembly of the orchid bee *E. dilemma* that we present here is of high quality, despite its large genome size (estimated to be 3.3Gb). The 15,904 gene annotations provide a comprehensive set of genes with known homology to the honey bee, facilitating future gene ontology and functional genomic analyses. While we were unable to annotate the mostly repetitive majority of the genome assembly with our approach, the provided sequence reads will be useful for future analyses of repetitive genetic elements in the genome. The nuclear and mitochondrial draft genomes represent a valuable genomic resource for the community of bee geneticists. This genomic resource will likely prove valuable in genetic and functional genomic analyses dealing with the ecology, evolution, and conservation of orchid bees. Furthermore, the genome of the facultatively eusocial *E. dilemma* will be helpful in the study of the evolution of eusociality, due to its taxonomic placement as the sister lineage to the three obligately eusocial corbiculate bee tribes including stingless bees, bumblebees, and honey bees.

Acknowledgements
We would like to thank Gene E. Robinson, Guojie Zhang, and the BGI for financial

456 support for genome sequencing. P.B. was supported by a fellowship of the German
457 academic exchange service (Deutscher Akademischer Austauschdienst, DAAD) for
458 parts of the project.
459

References

Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman, 1990 Basic local
alignment search tool. J. Mol. Biol. 215: 403–410.

Augusto, S. C., and C. A. Garófalo, 2009 Bionomics and sociological aspects of
Euglossa fimbriata (Apidae, Euglossini). Genet. Mol. Res. 8: 525–538.

Bao, W., K. K. Kojima, and O. Kohany, 2015 Repbase Update, a database of repetitive
elements in eukaryotic genomes. Mobile DNA 2015 6:1 6: 11.

Brand, P., S. R. Ramírez, F. Leese, J. J. Quezada-Euan, R. Tollrian *et al.*, 2015 Rapid
evolution of chemosensory receptor genes in a pair of sibling species of orchid
bees (Apidae: Euglossini). BMC Evol. Biol. 15: 176.

Branstetter, M. G., B. N. Danforth, J. P. Pitts, B. C. Faircloth, P. S. Ward *et al.*, 2017
Phylogenomic Insights into the Evolution of Stinging Wasps and the Origins of
Ants and Bees. Current Biology 27: 1019–1025.

Camacho, C., G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos *et al.*, 2009 BLAST plus
: architecture and applications. BMC Bioinformatics 10:.

Cardinal, S., and B. N. Danforth, 2011 The Antiquity and Evolutionary History of
Social Behavior in Bees (C. S. Moreau, Ed.). PLoS ONE 6:.

Cha, S. Y., H. J. Yoon, E. M. Lee, M. H. Yoon, J. S. Hwang *et al.*, 2007 The complete
nucleotide sequence and gene organization of the mitochondrial genome of the
bumblebee, Bombus ignitus (Hymenoptera: Apidae). Gene 392: 206–220.

Chalopin, D., M. Naville, F. Plard, D. Galiana, and J.-N. Volff, 2015 Comparative
Analysis of Transposable Elements Highlights Mobilome Diversity and Evolution
in Vertebrates. Genome Biol Evol 7: 567–580.

Crozier, R. H., and Y. C. Crozier, 1993 The mitochondrial genome of the honeybee
Apis mellifera: complete sequence and genome organization. Genetics 133: 97–
117.

Danforth, B. N., 2002 Evolution of sociality in a primitively eusocial lineage of bees.
Proc Natl Acad Sci USA 99: 286–290.

Dietz, L., P. Brand, L. M. Eschner, and F. Leese, 2015 The mitochondrial genomes of
the caddisflies Sericostoma personatum and Thremma gallicum (Insecta:
Trichoptera). Mitochondrial DNA 1–2.

Doetterl, S., and N. J. Vereecken, 2010 The chemical ecology and evolution of bee–
flower interactions: a review and perspectives. Canadian Journal of Zoology 88:
668–697.

Elsik, C. G., A. Tayal, C. M. Diesh, D. R. Unni, M. L. Emery *et al.*, 2016 Hymenoptera
Genome Database: integrating genome annotations in HymenopteraMine.
Nucleic Acids Res. 44: D793–D800.

Elsik, C. G., K. C. Worley, A. K. Bennett, M. Beye, F. Camara *et al.*, 2014 Finding the
missing honey bee genes: lessons learned from a genome upgrade. BMC
Genomics 15: 86.

Eltz, T., F. Fritzsch, and J. R. Pech, 2011 Characterization of the orchid bee Euglossa
viridissima (Apidae: Euglossini) and a novel cryptic sibling species, by
morphological, chemical, and genetic characters. Zool. J. Linn. Soc.

Eltz, T., W. M. Whitten, D. W. Roubik, and K. E. Linsenmair, 1999 Fragrance
Collection, Storage, and Accumulation by Individual Male Orchid Bees. J. Chem.
Ecol. 25: 157–176.

Eltz, T., Y. Zimmermann, C. Pfeiffer, J. R. Pech, R. Twele *et al.*, 2008 An olfactory shift
is associated with male perfume differentiation and species divergence in orchid
bees. Curr. Biol. 18: 1844–1848.

Garófalo, C. A., 1985 Social Structure of Euglossa cordata Nests (Hymenoptera:
Apidae: Euglossini). Entomologia Generalis 11: 77–83.

Gnerre, S., I. MacCallum, D. Przybylski, F. J. Ribeiro, J. N. Burton *et al.*, 2011 High-
quality draft assemblies of mammalian genomes from massively parallel
sequence data. Proc. Natl. Acad. Sci. U.S.A. 108: 1513–1518.

i5K Consortium, 2013 The i5K Initiative: Advancing Arthropod Genomics for
Knowledge, Human Health, Agriculture, and the Environment. Journal of
Heredity 104: 595–600.

Jurka, J., 2000 Repbase Update: a database and an electronic journal of repetitive
elements. Trends in Genetics 16: 418–420.

Kapheim, K. M., H. Pan, C. Li, S. L. Salzberg, D. Puiu *et al.*, 2015 Genomic signatures of
evolutionary transitions from solitary to group living. Science aaa4788.

Kidwell, M. G., 2002 Transposable elements and the evolution of genome size in
eukaryotes. Genetica 115: 49–63.

Klein, A.-M., B. E. Vaissière, J. H. Cane, I. Steffan-Dewenter, S. A. Cunningham *et al.*,
2007 Importance of pollinators in changing landscapes for world crops.
Proceedings of the Royal Society B: Biological Sciences 274: 303–313.

Kocher, S. D., C. Li, W. Yang, H. Tan, S. V. Yi *et al.*, 2013 The draft genome of a socially
polymorphic halictid bee, Lasioglossum albipes. Genome Biology 14: R142.

Korf, I., 2004 Gene finding in novel genomes. BMC Bioinformatics 5:.

Laslett, D., and B. Canbäck, 2008 ARWEN: a program to detect tRNA genes in
metazoan mitochondrial nucleotide sequences. Bioinformatics 24: 172–175.

Leese, F., P. Brand, A. Rozenberg, C. Mayer, S. Agrawal *et al.*, 2012 Exploring
Pandora's box: potential and pitfalls of low coverage genome surveys for
evolutionary biology. (B. J. Mans, Ed.). PLoS ONE 7: e49202.

Litman, J. R., B. N. Danforth, C. D. Eardley, and C. J. Praz, 2011 Why do leafcutter bees
cut leaves? New insights into the early evolution of bees. Proc. Biol. Sci. 278:
3593–3600.

Lowe, T. M., and S. R. Eddy, 1997 tRNAscan-SE: A program for improved detection of
transfer RNA genes in genomic sequence. Nucleic Acids Res. 25: 955–964.

Luo, R., B. Liu, Y. Xie, Z. Li, W. Huang *et al.*, 2012 SOAPdenovo2: an empirically
improved memory-efficient short-read de novo assembler. GigaScience 2016 5:1
1: 18.

Mayer, C., 2010 Phobos Version 3.3.12. A tandem repeat search program. 20 pp.

Menzel, R., and U. Muller, 1996 Learning and Memory in Honeybees: From Behavior
to Neural Substrates. Annu. Rev. Neurosci. 19: 379–404.

Michener, C. D., 2007 *The Bees of the World. 2nd*. Ed. Johns Hopkins.

Nowak, M. A., C. E. Tarnita, and E. O. Wilson, 2010 The evolution of eusociality. 466:
1057–1062.

Pamilo, P., L. Viljakainen, and A. Vihavainen, 2007 Exceptionally High Density of
NUMTs in the Honeybee Genome. Mol Biol Evol 24: 1340–1346.

Park, D., J. W. Jung, B.-S. Choi, M. Jayakodi, J. Lee *et al.*, 2015 Uncovering the novel
characteristics of Asian honey bee, Apis cerana, by whole genome sequencing.
BMC Genomics 16:.

Parra, G., K. Bradnam, and I. Korf, 2007 CEGMA: a pipeline to accurately annotate
core genes in eukaryotic genomes. Bioinformatics 23: 1061–1067.

Pech, M. E. C., W. de J May-Itzá, L. A. M. Medina, and J. J. G. Quezada-Euan, 2008
Sociality in Euglossa (Euglossa) viridissima Friese (Hymenoptera, Apidae,
Euglossini). 55: 428–433.

Pemberton, R. W., and G. S. Wheeler, 2006 Orchid bees don't need orchids: evidence

from the naturalization of an orchid bee in Florida. Ecology 87: 1995–2001.

Peters, R. S., L. Krogmann, C. Mayer, A. Donath, S. Gunkel *et al.*, 2017 Evolutionary
History of the Hymenoptera. Current Biology 0:.

Platt, R. N., L. Blanco-Berdugo, and D. A. Ray, 2016 Accurate Transposable Element
Annotation Is Vital When Analyzing New Genome Assemblies. Genome Biol Evol
8: 403–410.

Ramírez, S., R. L. Dressler, and M. Ospina, 2002 Abejas euglosinas (Hymenoptera:
Apidae) de la Región Neotropical: Listado de especies con notas sobre su
biología. 3: 7.

Ramírez, S. R., T. Eltz, M. K. Fujiwara, G. Gerlach, B. Goldman-Huertas *et al.*, 2011
Asynchronous Diversification in a Specialized Plant-Pollinator Mutualism.
Science 333: 1742–1746.

Romiguier, J., S. A. Cameron, S. H. Woodard, B. J. Fischman, L. Keller *et al.*, 2015
Phylogenomics controlling for base compositional bias reveals a single origin of
eusociality in corbiculate bees. Mol Biol Evol msv258.

Roubik, D. W., and P. E. Hanson, 2004 *Orchid bees of tropical America: biology and
field guide*. Instituto Nacional de Biodiversidad (INBio), Santo Domingo De
Heredia.

Sadd, B. M., S. M. Barribeau, G. Bloch, D. C. de Graaf, P. Dearden *et al.*, 2015 The
genomes of two key bumblebee species with primitive eusocial organization.
Genome Biology 16: 76.

Simão, F. A., R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva, and E. M. Zdobnov,
2015 BUSCO: assessing genome assembly and annotation completeness with
single-copy orthologs. Bioinformatics 31: 3210–3212.

Skov, C., and J. Wiley, 2005 Establishment of the neotropical orchid bee Euglossa
viridissima (Hymenoptera: Apidae) in Florida. Florida Entomologist 88: 225–
227.

Slater, G. S. C., and E. Birney, 2005 Automated generation of heuristics for biological
sequence comparison. BMC Bioinformatics 6: 31.

Soro, A., J. J. G. Quezada-Euan, P. Theodorou, R. F. A. Moritz, and R. J. Paxton, 2016
The population genetics of two orchid bees suggests high dispersal, low diploid
male production and only an effect of island isolation in lowering genetic
diversity. Conserv Genet 1–13.

Stanke, M., M. Diekhans, R. Baertsch, and D. Haussler, 2008 Using native and
syntenically mapped cDNA alignments to improve de novo gene finding.

Bioinformatics 24: 637–644.

Stolle, E., L. Wilfert, R. Schmid-Hempel, P. Schmid-Hempel, M. Kube *et al.*, 2011 A
second generation genetic map of the bumblebee Bombus terrestris (Linnaeus,
1758) reveals slow genome and chromosome evolution in the Apidae. BMC
Genomics 12: 48.

Suni, S. S., 2016 Dispersal of the orchid bee Euglossa imperialis over degraded
habitat and intact forest. Conserv Genet 1–10.

Suni, S. S., and B. J. Brosi, 2012 Population genetics of orchid bees in a fragmented
tropical landscape. Conserv Genet 13: 323–332.

Vogel, S., 1966 Parfümsammelnde Bienen als Bestäuber von Orchidaceen und
Gloxinia. Österr bot Z 113: 302–361.

Wcislo, W. T., and J. H. Cane, 2003 Floral Resource Utilization by Solitary Bees
(Hymenoptera: Apoidea) and Exploitation of Their Stored Foods by Natural
Enemies. 41: 257–286.

Weinstock, G. M., G. E. Robinson, R. A. Gibbs, K. C. Worley, J. D. Evans *et al.*, 2006
Insights into social insects from the genome of the honeybee Apis mellifera.
Nature 443: 931–949.

Whitten, W. M., A. M. Young, and D. L. Stern, 1993 Nonfloral sources of chemicals
that attract male euglossine bees (Apidae: Euglossini). J. Chem. Ecol. 19: 3017–
3027.

Woodard, S. H., B. J. Fischman, A. Venkat, M. E. Hudson, K. Varala *et al.*, 2011 Genes
involved in convergent evolution of eusociality in bees. Proc Natl Acad Sci USA
108: 7472–7477.

Xu, H., X. Luo, J. Qian, X. Pang, J. Song *et al.*, 2012 FastUniq: a fast de novo duplicates
removal tool for paired short reads. (D. Doucet, Ed.). PLoS ONE 7: e52249.

Zdobnov, E. M., F. Tegenfeldt, D. Kuznetsov, R. M. Waterhouse, F. A. Simão *et al.*,
2017 OrthoDB v9.1: cataloging evolutionary and functional annotations for
animal, fungal, plant, archaeal, bacterial and viral orthologs. - PubMed - NCBI.
Nucleic Acids Res. 45: D744–D749.

Zimmermann, Y., D. L. P. Schorkopf, R. F. A. Moritz, R. W. Pemberton, J. J. G. Quezada-
Euan *et al.*, 2011 Population genetic structure of orchid bees (Euglossini) in
anthropogenically altered landscapes. Conserv Genet 12: 1183–1194.

Figures

Figure 1. Genomic features. (A) Phylogeny of the four corbiculate bee tribes with orchid bees as sistergroup to honey bees, stingless bees, and bumblebees (Romiguer et al. 2015). **(B)** K-mer distribution spectrum (k=25) of genomic sequence reads. The positively skewed spectrum reveals a high abundance of a few k-mers, leading to an estimate of 87.7% repetitiveness of the *E. dilemma* genome. Red shows the k-mer spectrum before, and blue after error correction. **(C)** Genomic element density including genic and non-genic features as a fraction of the overall genome assembly length. Over 85% of the assembly could not be annotated with the selected methods. **(D)** Synteny between the *E. dilemma* and the honey bee (*Apis mellifera*) genome. In an analysis including *E. dilemma* scaffolds of ≥100kb length, 83% showed ≥95% synteny to a single honeybee scaffold. Photographs in **(A)** are reproduced from Wikimedia under the CC BY-SA 3.0 license.

Figure 2. Mitochondrial genome reconstruction. The structure of the honey bee mitochondrial genome and information of the homologous reconstructed parts of the *E. dilemma* mitochondrial genome. Non-reconstructed parts of incompletely reconstructed genes are hatched.

Figure 1

653
654 **Figure 2**

655 **Tables**

657 **Table 1.** *E. dilemma* **genome assembly statistics in comparison to previously**
658 **published bee genomes.**

659
660 **Table 1**

Species	N50	N25	Longest scaffold	Scaffolds	Assembly length	%GC	Predicted genes	Ref.
Euglossa dilemma	143,590	1,417,006	10,108,120	22,698	588,199,720	39.94	15,904	1
Eufriesea mexicana	2,427	443,231	4,677,300	3,522,543	1,031,837,970	41.38	12,022	2
Apis mellifera	997,192	1,922,192	4,736,299	5,644	234,070,657	32.70	15,314	3
Melipona quadrifasciata	68,085	1,896,322	12,087,087	38,604	507,114,161	38.88	14,257	2
Bombus impatiens	1,399,493	2,389,513	5,466,090	5,559	249,185,056	37.75	15,896	4
Lasioglossum albipes	616,426	1,130,413	3,533,895	41,433	341,616,641	41.50	13,448	5

661 N50 and N25 indicate the length of the shortest scaffold of those including 50% and 25% of the base pairs in a
662 genome assembly. References (Ref.): 1: This study, 2: Kapheim et al. 2015, 3: Elsik et al. 2014, 4: Sadd et al. 2015,
663 5: Kocher et al. 2012.