Diversity Predicts Ability of Bacterial Consortia to Mitigate a Lethal Wildlife Pathogen

Rachael E. Antwis[1,2], Xavier A. Harrison[3]

1. School of Environment and Life Sciences, University of Salford, Salford, UK

2. Unit for Environmental Sciences and Management, North-West University, Potchefstroom,

South Africa

3. Institute of Zoology, Zoological Society of London, Regent's Park, London NW1 4RY

Address for Correspondence:

Dr. Rachael Antwis, University of Salford, Room 336, Peel Building, The Crescent, Salford,

M5 4WT, UK

01612954641

r.e.antwis@salford.ac.uk

Running Title:

Diversity predicts bacterial anti-fungal activity

Conflict of Interest:

The authors declare no conflict of interest.

ABSTRACT

Symbiotic bacterial communities can protect their hosts from infection by pathogens.

Treatment of wild individuals with protective bacteria can combat the spread of emerging

infectious diseases (EIDs), but it is unclear whether the degree of bacterially-mediated host

protection is uniform across multiple isolates of globally-distributed pathogens. Here we use

the lethal fungal pathogen *Batrachochytrium dendrobatidis (Bd)* as a model to investigate the

traits predicting broad-scale *in vitro* inhibitory capabilities of both individual bacteria and

multiple-bacterial consortia. We show that inhibition of multiple pathogen isolates is rare, with

no clear phylogenetic signal at the genus level. Bacterial consortia offer stronger protection

against *B. dendrobatidis* compared to single isolates, but critically this was only true for

consortia containing multiple genera, and this pattern was not uniform across all *B.*

dendrobatidis isolates. These novel insights have important implications for the effective

design of bacterial probiotics to mitigate EIDs.

INTRODUCTION

The last 50 years have seen the emergence of several hypervirulent wildlife pathogens with broad host ranges (Tompkins et al 2015). These emerging infectious disease (EIDs) have decimated wildlife populations globally, and are a major contributor to the so-called current "biodiversity crisis" (e.g. Skerrat et al 2007; McCallum 2012). Both climate change (Cohen et al 2017) and the global trade in animals (Tompkins et al 2015) are contributing to the increased spread of EIDs, and broad-scale, effective treatments and/or prophylaxis for these pathogens in the wild are often lacking (Sleeman 2013; Garner et al 2016). Developing such treatments is often complicated by broad variation in genetic and phenotypic traits such as virulence exhibited by these pathogens (e.g. de Jong & Hien 2006; Schock et al 2010; Farrer et al 2011). Successful mitigation of EIDs in the wild demands that preventative or curative therapies demonstrate broad activity over as many genetic variants of the pathogen as possible, and developing mitigation strategies that satisfy this criterion remains a major outstanding research goal.

Most EIDs are attributed to fungal pathogens, including *Pseudogymnoascus destructans* that causes white nose syndrome in bats, and *Batrachochytrium spp.*, which causes chytridiomycosis in amphibians (Fisher et al 2012). *Batrachochytrium dendrobatidis* comprises multiple, deeply diverged lineages, and is capable of rapid evolution (Farrer et al 2011; 2013). Endemic lineages of *B. dendrobatidis* have been identified, including *Bd*CAPE (South Africa), *Bd*CH (Switzerland), *Bd*Brazil (Brazil) and a lineage from Japan (Goka et al 2009; Farrer et al 2011; Schloegel et al 2012; Rosenblum et al 2013; Rodriguez et al 2014), although there are cases where these have spread to other regions and are implicated in population declines in those regions (e.g. *Bd*CAPE in Mallorcan midwife toads, *Alytes muletensis;* Doddington et al 2013). The globally distributed and hypervirulent global panzootic lineage (*Bd*GPL) is the genetic lineage of *B. dendrobatidis* associated with phenomenal mass mortalities and rapid population declines of amphibians around the world, and is a major driver of the current "amphibian extinction crisis" (Fisher et al 2009; Farrer et

al 2011; Olson et al 2012). Isolates within this lineage exhibit enormous and unpredictable

variation in virulence, even within a single host species exposed under laboratory conditions

(Farrer et al 2011; Farrer et al 2013). There is currently no cure for this disease in the wild

(reviewed in Garner et al 2016), and given that amphibian communities may be host to

multiple *Bd*GPL genotypes (Morgan et al 2007; Rodriguez et al 2014), and the continuous

global movement of humans and wildlife that can transport the fungus, any prophylactic or

curative treatment needs to effective against multiple *B. dendrobatidis* genotypes and

isolates.

Bacterial probiotics represent a promising tool to combat major emerging fungal pathogens in

the wild, including *Pseudogymnoascus destructans* (Hoyt et al 2015), *B. dendrobatidis*, and

the closely related *B. salamandrivorans* (Martel et al 2013; 2014). Of these, probiotic

research is currently most advanced for *B. dendrobatidis* (reviewed in Bletz et al 2013 and

Rebollar et al 2016). Laboratory and field studies have shown host-associated bacterial

communities (hereafter referred to as the 'microbiome') protect amphibians from *B.*

dendrobatidis infection, and that it is possible to artificially augment the microbiome with

'probiotic' bacteria to improve survivorship in response to the pathogen (Bletz et al 2013; Jani

& Briggs 2014; Becker et al 2015; Walke et al 2015).

To date, most *in vitro Bd*GPL challenge experiments have tested the ability of candidate

probiotics to limit the growth just a single isolate of *Bd*GPL. This is problematic, because the

inhibitory capabilities of individual bacteria are not uniform across the variation presented by

*Bd*GPL (Antwis et al 2015). Previous work has found no evidence of a phylogenetic signal in

the ability of bacterial genera to inhibit a singular BdGPL isolate (Becker et al 2015), but a

major gap in our understanding concerns whether some bacterial genera are more likely to

show broad-spectrum inhibition across a range of *Bd*GPLs, allowing a more focussed search

for effective amphibian probiotics. Furthermore, both *in vivo* amphibian probiotic trials and i*n*

vitro challenges focus on the application of a singular bacterial isolate to arrest the growth of

B. dendrobatidis, yet the importance of a complex and diverse microbiome for resilience to

infection has been repeatedly demonstrated across a range of host taxa (e.g. Dillon et al

2005; Matos et al 2005; Van Elsas et al 2012; Eisenhauer at el 2013). A novel alternative

strategy involves a 'bacterial consortium' approach to probiotics, whereby multiple inhibitory

bacterial isolates are applied simultaneously. Multi-species consortia can increase the

inhibition of *Bd*GPL (Piova-Scott et al 2017), and so may offer greater inhibitory capabilities

across a wider range of *B. dendrobatidis* isolates, but the generality of this pattern across

multiple pathogen variants remains untested. Addressing the shortfall in our understanding is

critical for developing effective tools for the mitigation of EIDs in the wild.

Here we extend previous work to quantify the ability of metabolites from both individual

bacteria and co-cultured bacterial consortia to demonstrate broad-scale inhibition across a

panel of *B. dendrobatidis* isolates. First, we test 58 bacterial isolates from 10 genera for

inhibition against a suite of 10 different *Bd*GPL isolates to quantify i) variation among

bacterial genera in ability to demonstrate broad-spectrum *Bd*GPL inhibition and ii) variation

among *Bd*GPL isolates is susceptibility to inhibition. Second, we quantify the relative efficacy

of using single bacterial isolates or bacterial consortia to modify *B. dendrobatidis* growth

rates *in vitro*. Specifically, we investigate iii) whether consortia yield stronger inhibition than

single bacteria across three *B. dendrobatidis* isolates from two lineages (*Bd*GPL and

*Bd*CAPE); and iv) whether the diversity of a bacterial consortium (number of member

genera) affects inhibitory capabilities.

METHODS

Phylogeny screening

In vitro challenges were conducted for 58 bacteria isolated from wild *Agalychnis spp.* frogs in Belize (Antwis et al 2015) to screen for inhibitory capabilities against 10 *Bd*GPL isolates (Table 1, Figure 1) using an *in vitro* spectrophotometer assay method adapted from Bell et al. (2013), Woodhams et al (2014) and Becker et al (2015). Bacteria belonged to 10 genera, with 3-11 bacterial isolates per genus (Table S1). Bacteria were grown by adding 50ul of frozen stock bacteria (stored in 30% glycerol, 70% tryptone solution at -80°C) to 15ml of 1% tryptone, and incubating at 18°C for 36 hours until visibly turbid (three cultures per bacterial isolate). Bacterial densities were counted using a haemocytometer and adjusted to ~500,000 cells/ml. Bacteria were then filtered through a 0.22um sterile filter (Millipore, Ireland) to remove live cells, leaving only bacterial metabolites in solution, which were combined across cultures for a given bacterial isolate, and then kept on ice until *B. dendrobatidis* challenges were conducted. *Bd*GPL (Table 1) isolates were grown in 1% tryptone broth until maximum zoospore production was observed (~3-4 days; ~1 x 10^6 zoospores ml^{-1}). As with bacteria, three flasks per *B. dendrobatidis* isolate were grown and then combined prior to challenges to limit flask-effect. Zoospores were separated from sporangia by filtering through 20um sterile filters (Millipore, Ireland). To conduct the spectrophotometer assays, 50ul of bacterial metabolites and 50ul of *B. dendrobatidis* suspension were pipetted into 96-well plates. Each *B. dendrobatidis*-bacteria combination was run with three repeats. Positive controls were included using 50ul 1% tryptone instead of bacterial metabolites. Negative controls were included using 50ul sterile water and 50ul of heat-treated *B. dendrobatidis* for each isolate. Plate readings were taken every 24 hours for four days using a 492nm filter.

For each measurement, data were transformed using the equation Ln(OD/(1-OD)), and a regression analysis was used to gain the slope values for each sample over time. Slopes of triplicate replicates for each Bd/Bacteria combination were averaged, and total *B.*

dendrobatidis inhibition was calculated using the formula: Inhibition (%) = [1-(slope of

sample/slope of control)] x 100. A positive value represents inhibition of *B. dendrobatidis*

growth, and a negative value indicates enhanced growth of *B. dendrobatidis*.

Bacterial consortium challenges

Three bacteria were then selected from each of four genera (*Acinetobacteria,*

Chryseobacterium, Serratia, Stenotrophomonas) based on their inhibition profiles; poor to

medium inhibitors were selected to determine whether combining these bacteria would

improve their inhibitory capabilities. Bacteria were grown individually until turbid, adjusted to

~500,000 cells/ml, and added to fresh tryptone either individually (strains A, B and C of each

genus separately), or as a triple (strains A, B and C of each genus together to form single-

genus mixes, or a random combination of strains across genera to form multi-genus (Table

2)). To combine bacteria, a total of 3ml of bacteria were added to 12ml of fresh 1% tryptone

broth and left to grow together for 18 hours. The volume of each bacterium added depended

on whether the consortium contained one or three bacteria, and the volume was split evenly

between the number of bacteria added to each group. Following this, bacteria-*B.*

dendrobatidis challenges were conducted using the same methods as described above

against three *B. dendrobatidis* isolates (Table 1). Average inhibition percentages for each

consortium-*B. dendrobatidis* combination were calculated as previously described.

Statistical Analysis

All statistical analyses were conducted in the software R v.3.3.2 (R Core Team 2016). R

scripts for all analyses presented in this manuscript are provided as a R Markdown document

in supplementary information.

Phylogeny Data: To quantify differences among genera in proportion of *Bd*GPL isolates

inhibited (inhibition score >0), we fitted a Binomial GLM with the proportion of the 10 *Bd*GPL

isolates each bacterial isolate inhibited as the response, and genus as a fixed effect. We

used the quasibinomial error structure as the model was overdispersed (dispersion 6.4), and

tested the model containing a genus term with the reduced intercept-only model using a

likelihood ratio test.

To quantify differences among genera in the *degree* of inhibition (size of inhibition score), we

fitted a hierarchical model in the R package *MCMCglmm* (Hadfield 2010) with the individual

inhibition scores of each bacterial isolate (n=58) for each *Bd*GPL isolate (n=10; total n = 580)

as a Gaussian response. We fitted both *Bd*GPL isolate, and bacterial strain ID nested within

bacterial genus as random effects. We use uninformative, parameter-expanded priors for the

random effects as detailed in Hadfield (2010). We ran models for a total of 100,000 iterations

following a burn-in of 10,000 iterations and using a thinning interval of 50. Posterior model

checks indicated no significant autocorrelation within chains (all values < 0.05) and adequate

convergence using the Geweke diagnostic (Geweke 1992). Inspection of model residuals

from the frequentist analogue of this model fitted in *lme4* (Bates et al 2015) revealed

normally-distributed residuals and no evidence of heteroscedasticity. Rerunning models with

stronger priors has no effect on model results.

To calculate % variance in inhibition explained by *Bd*GPL isolate, bacterial genus, and

bacterial strain respectively, we extracted the variance components from the variance-

covariance matrix of the model above. We expressed the variance of a component *V* as a

percentage of the total variance calculated as (V_{BdGPL} + V_{genus} + V_{strain} + $V_{residual}$). We

calculated both mean and 95% credible intervals using the posterior samples from the

model. To construct Figs. 1 and 2, we extracted the marginal means and 95% credible

intervals for each bacterial strain and *Bd*GPL isolate, respectively. That is, the bacterial strain

modes are marginalised with respect to *Bd*GPL and vice versa, to quantify whether the

average scores for each *Bd*GPL or bacterial isolate are significantly different from zero.

Consortium Data: To calculate the relative mean inhibition of single-genus (SG) vs multi-

genus (MG) consortia, we fitted a mixed model in *MCMCglmm* with inhibition as a Gaussian

response, consortium type as a 2-level factor, and a random effect of *B. dendrobatidis* and

using uninformative priors. To calculate whether consortia exhibited stronger inhibition than

the mean of their individual isolates, we constructed a binary variable with outcome 1 if a

consortium's inhibition was greater than the single isolate mean and 0 if lower. We fitted this

as a response in a binary GLMM with consortium type as a fixed effect, *B. dendrobatidis* as a

random effect and using uninformative priors. Neither model exhibited signs of

autocorrelation and Geweke statistics for both models indicated convergence.

Simulated Consortium Trials: To probe the relative effectiveness of single bacteria, SG

consortia and MG consortia (hereafter 'probiotic types') for modifying the growth rates of *B.*

dendrobatidis, we ran three sets of simulations, each comprising 1000 iterations. For each

set of simulations, we calculated i) the proportion of times a MG consortium yielded higher

inhibition than a SG consortium; ii) the proportion of times a MG consortium yielded higher

inhibition than a single bacterial isolate; iii) the probability that a MG, SG or single bacterial

isolate would yield at least 50% inhibition, which we class as strong inhibition. We derived

95% confidence intervals for each test statistic by performing 10,000 bootstrap samples with

replacement from the test distributions. The three simulations were as follows:

(1) Averaged Over All *B. dendrobatidis* isolates: For each iteration, we randomly selected a

B. dendrobatidis isolate, and then randomly selected both a SG and a MG consortium. A

Single bacterial isolate score was then selected randomly from one of the members of the

MG consortium.

(2) *B. dendrobatidis* specific scores: To investigate the potential for the effectiveness of

consortia to differ depending on *B. dendrobatidis* isolate, we repeated the simulations as in

(1) but performed 1000 simulations for *each B. dendrobatidis* isolate. (3) Sequential *B.*

dendrobatidis exposure: Finally, we examined the ability of the three probiotic types to inhibit

two *B. dendrobatidis* isolates in series by randomly selecting two of the three *B.*

dendrobatidis isolates. For each iteration, we selected a random MG and SG consortium,

followed by a randomly-selected single isolate member from the MG consortium. Individual

inhibition scores for these three groups were then extracted for both selected *B.*

dendrobatidis isolates (i.e. probiotic ID was kept consistent over both pathogen isolates). We

calculated the probability that the MG consortium would yield superior inhibition to the SG

consortium and single bacterial isolate across both *B. dendrobatidis* isolates, and the

probability that all three probiotic types would yield >50% inhibition.

229
230

RESULTS

Phylogenetic Signals of *Bd*GPL Inhibition

We assayed the ability of 58 bacterial isolates from 10 genera to modify the growth rates of 10 *Bd*GPL isolates. Mean inhibition scores ranged from 100% (complete inhibition of growth) to -225% (strong facilitation of growth). At the genus level, there was no significant variation among genera in mean proportion of *Bd*GPL isolates inhibited (Binomial GLM; χ^2_9 = 6.2, p=0.72; Table 3). Six isolates from five genera showed at least weak inhibition across all 10 *Bd*GPLs, whilst seven isolates from five genera facilitated the growth of all 10 isolates (Supplementary Table S1).

Variance component analysis revealed considerably more variation in inhibition scores among bacterial strains *within* genera than among genera themselves (Fig. 1). Bacterial strain ID explained 51% [95% credible interval (CRI) 37-63%] of the variation in *Bd*GPL inhibiton scores compared to just 1.3% [0.09-6%] for bacterial genus. *Bd*GPL isolate explained 15.6% [4.8-30%] of the variation in inhibition scores and highlighted two isolates whose marginal effect sizes were significantly negative (JEL423 and AUL2), and one isolate with a significantly positive marginalised inhibition score (08MG04; Fig. 2). JEL423 and AUL2 therefore exhibit strongly enhanced growth in the presence of bacterial metabolites, whereas 08MG04 is particularly susceptible to inhibition of growth. The remaining seven *Bd*GPL isolates demonstrated no evidence of systematic susceptibility to inhibition of their growth rates across the bacteria tested (Fig. 2).

Multi-Isolate Consortia as Tools for Pathogen Mitigation

Consortia containing isolates from Multi-Genus (MG) exhibited significantly higher mean inhibition scores compared to Single-Genus (SG) consortia when marginalising with respect to *B. dendrobatidis* isolate (MG consortia mean inhibition: 36.88%; SG consortia mean:

16.9%; 95% CRI of difference 4.12 – 36.52%, p_{MCMC} = 0.02; Fig. 3). If the ability of a

consortium to inhibit *B. dendrobatidis* was simply an additive function of the inhibitory

capabilities of the individual bacteria it comprised, we would expect the consortium's

inhibition score to be equal to the mean of the individual inhibition scores, weighted by

relative abundance. Inhibition scores of consortia greater than the mean of individual isolate

scores are indicative of synergistic effects, whereby the combined pool of metabolites from

multiple bacteria inhibits *B. dendrobatidis* more strongly than the individual isolates. MG

consortia had a 61% probability of demonstrating stronger inhibition than the mean of their

single composite bacterial isolates, which was significantly higher than the corresponding

probability for SG isolates (26.6%, Mean difference 39.4% [95% Credible Interval 11.2-

65.1%], p_{MCMC} = 0.01).

in silico Probiotic Trials: Of the 1000 simulated probiotic trials, naïve application of a MG

consortium yielded higher *B. dendrobatidis* inhibition in 69.4% of cases [95% CI 66.5-72.3%]

compared to SG consortia (null expectation 50%, p_{RAND}<0.001). Moreover, MG consortia had

a 38.1% [35.1 – 41.1%] probability of yielding inhibition greater than 50% (strong inhibition),

compared to only 13.9% [11.8 – 16.1%] probability for SG consortia. MG consortia

outperformed the single isolate in 61% [58-64%] of cases (null expectation 50%,

p_{RAND}<0.001). However, by averaging over all *B. dendrobatidis* isolates, these results

masked substantial variation among *B. dendrobatidis* isolates in the relative efficacy of MG

versus SG consortia. We repeated the above simulations separately for each *B.*

dendrobatidis isolate, and found that MG consortia were superior to SG consortia and single

bacterial isolates for only two *B. dendrobatidis* isolates (*Bd*GPL MODS28 and *Bd*CAPE

TF5a1), and performed slightly worse than SG consortia for *Bd*GPL SFBC019 (Fig. 4A).

Moreover, although MG consortia have the greatest probability of yielding >50% inhibition for

*Bd*GPL MODS28 and *Bd*CAPE TF5a1, this was not the case for *Bd*GPL SFBC019, where

SG consortia had a marginally higher probability of delivering strong inhibition (Fig. 4B).

Finally, *we* tested the ability of both MG and SG consortia to inhibit the growth of two

different *B. dendrobatidis* isolates in series, as individuals in a single location may be

exposed to multiple variants of a pathogen (Goka et al 2009; Schloegel et al 2012; Rodriguez

et al 2014; Jenkinson et al 2016), or strong spatial structure of the pathogen and high host

dispersal may expose individuals to multiple pathogen variants consecutively. For a given

trial, the modelling outcomes were; i) MG consortia inhibited both *B. dendrobatidis* isolates

more strongly than SG consortia; ii) SG consortia inhibited both *B. dendrobatidis* isolates

more strongly than MG consortia; iii) MG inhibited the first *B. dendrobatidis* isolate more

strongly than SG consortia, but not the second; iv) MG inhibited the second *B. dendrobatidis*

isolate more strongly than SG consortia, but not the first. Applying the same MG consortium

to two *B. dendrobatidis* isolates in series achieved stronger inhibition than SG consortia in

49.4% [46.3 – 52.5%] of cases (i.e. modelling outcome i; null expectation 25% [0.5^2],

p_{RAND}<0.001). This compared to only 7.9% [6.4-9.6%] of cases where SG consortia exhibited

superior inhibition for both *B. dendrobatidis* isolates (i.e. modelling outcome iv). MG consortia

provided superior inhibition for only one of the *B. dendrobatidis* isolates in the remaining 43%

of cases (mean 20.3% and 22.4% of simulations with superior inhibition for the first and

second isolate respectively). MG consortia exhibited strong inhibition (>50%) for both

isolates in 14.7% [12.5-17%] of cases, compared to zero cases where SG isolates did so.

Applying a single bacterial isolate instead of a SG or MG consortium resulted in strong

inhibition for both *B. dendrobatidis* isolates in only 4% [2.9-5.3%] of cases (Fig. 4C).

DISCUSSION

The principal objectives of this study were two-fold: i) to determine the magnitude, if any, of phylogenetic signal in the ability of certain genera of bacteria to inhibit a broad range of *Bd*GPL isolates; and ii) to examine the relative effectiveness of single bacteria and bacterial consortia to inhibit several isolates of *B. dendrobatidis*. We found no evidence of variation among bacterial genera in their ability to exhibit broad-range inhibition across multiple *Bd*GPL isolates. Furthermore, our data suggested consortia provide superior *B. dendrobatidis* inhibition than individual bacteria, but critically this pattern is not uniform across pathogen isolates, and is contingent on consortium taxonomic diversity. Our results have important implications for our understanding of the factors determining *in vivo* resistance to infection in the wild, and provide novel insights into effective strategies for designing probiotic therapies to mitigate lethal cutaneous infections.

Phylogenetic Signals of Bd*GPL Inhibition*

We detected no phylogenetic signal in the ability of individual bacterial genera to inhibit multiple *Bd*GPL isolates. These data support previous work suggesting the ability to inhibit *B. dendrobatidis* is distributed widely over bacterial genera (Antwis et al 2015; Becker et al 2015); several isolates demonstrated at least weak inhibition for all 10 *Bd*GPLs but were spread across multiple genera with no clear pattern. That there is clear functional redundancy among genera in this host-protective trait suggests it is not prudent to focus on any one genus in the search for beneficial probiotics (Becker et al 2015), as highly divergent microbial communities can still possess similar functional traits (e.g. Bletz et al 2016). The principal source of variance in inhibition was among bacterial strains, with the number of isolates demonstrating broad-spectrum *facilitation* of *Bd*GPL being roughly equal to the number exhibiting broad-scale *inhibition* of the pathogen. The phenomenon of *Bd*GPL growth facilitation has been described previously for single pathogen isolates (Bell et al 2013;

Becker et al 2015), but crucially our results suggest that a bacterial strain's ability to facilitate

the growth of *B. dendrobatidis* may extend across a broad suite of pathogen isolates.

It is unclear *why* some bacterial isolates facilitate *B. dendrobatidis* growth, but one likely

explanation is that certain bacterial metabolites can act as growth substrates for fungi

(Garbaye 1994; Hardoim et al 2015), or that different bacterial metabolites alter the abiotic

environment (e.g. pH) to confer different growth rates (Romanowski et al 2011). Here we

have provided some of the first evidence that facilitation of *B. dendrobatidis* growth is not

simply a rare phenomenon arising from specific *Bd*GPL/bacterial combinations, but that this

is widespread across bacterial isolates, and different *Bd*GPL isolates differ systematically in

their growth rates when exposed to bacterial metabolites. That said, all four CORN isolates

showed similar levels of inhibition across all bacterial isolates, whereas the two AUL isolates

exhibited markedly different inhibition profiles (Figure 2). We identified one *Bd*GPL isolate

that was significantly prone to inhibition, and a further two isolates that demonstrated strong

resistance to inhibition across the 58 bacterial isolates we tested. That there is variation in

this trait among *Bd*GPL isolates is intriguing; if facilitation occurs because *B. dendrobatidis*

uses bacterial metabolites for nutrition, it may suggest some *B. dendrobatidis* variants can

use those metabolites more efficiently for growth. Data gathered from additional isolates will

allow us to formally test this hypothesis by probing whether a *Bd*GPL's susceptibility to

inhibition or facilitation correlates with virulence. Previous work has shown no among-isolate

variation in susceptibility of *B. dendrobatidis* to an echinocandin antifungal drug (Fisher et al

2009), yet our data suggest this pattern is not the same for bacterial metabolites.

Recombination among lineages of *Bd*GPL is common (Farrer et al 2011), providing a

mechanism whereby metabolic genes favouring enhanced growth may be spread following

contact among lineages. Our data have two important implications given the proclivity of *B.*

dendrobatidis for recombination. First, among-isolate variation in susceptibility to inhibition

suggests that the relative efficacy of probiotic or curative therapies in the wild will be modified

by local *B. dendrobatidis* genotype. Second, though we tend to treat bacterial inhibition

scores as fixed traits, this ignores the ability of genetic recombination among *B.*

dendrobatidis lineages to modify the relationship between bacterial metabolites and

pathogen growth rates. Even the application of probiotics themselves may represent a strong

selective pressure favouring genetic variants of *B. dendrobatidis* that lack susceptibility to

those probiotics. Although several trials have demonstrated the potential for probiotic

prophylaxis against *B. dendrobatidis* (e.g. Harris 2009; Muletz et al 2012; Kueneman et al

2016), we still lack the requisite data to measure selection caused by those trials on the

pathogen. *In vitro* experimental evolution assays between pathogen and bacteria may prove

the most powerful means for detecting such patterns.

Consortium-Based Approaches to Combatting Fungal Pathogens

Our results revealed a positive link between the taxonomic richness of a probiotic consortium

and its ability to inhibit *B. dendrobatidis* growth, but crucially this relationship was highly

dependent on *B. dendrobatidis* isolate. Multi-genus consortia outperformed both single-

genus consortia and single bacterial isolates in *B. dendrobatidis* inhibition, and were far more

likely to produce strong inhibition of 50% or greater, but only for two of the three pathogen

variants.

The general relationship between inhibition and consortium diversity was in the expected

direction; low community relatedness (i.e. high community dissimilarity) and high species

richness both increase the resistance of a bacterial community to pathogenic 'invaders' (e.g.

Jousset et al 2011; Eisenhauer et al 2012, 2013). Furthermore, previous work has linked

higher species diversity of probiotic consortia to increased *B. dendrobatidis* inhibition using a

single pathogen isolate (Piova-Scott et al 2017). Superior inhibition from consortia, rather

than single isolates, may arise as a by-product of the interference competition over resources

created by co-culture (Scheuring & Yu 2012). Thus, even bacteria that are weak inhibitors

when grown individually can increase the overall inhibitory power of a consortium by creating

a competitive environment that favours greater production of anti-fungal compounds.

Functional dissimilarity has been proposed as more important than taxonomic diversity in

predicting a community's resilience to invasion (Eisenhauer et al 2013), but may explain why

single-genus consortia did not perform as well as multi-genus consortia. In selecting for

genetic diversity, we may have been simultaneously selecting for functional diversity not

present when co-culturing three members of the same genus.

That *B. dendrobatidis* isolate can alter the strength of the relationship between consortium

diversity and inhibition is a highly novel finding. Our simulated probiotic trials revealed that for

two *B. dendrobatidis* isolates, combining bacteria into multi-genus consortia yielded

significantly better inhibition than applying one of the member bacteria in isolation. These

results provide further support for a synergistic effect of co-culture on inhibition. If multi-

genus consortia were no better at inhibition than the mean of their composite members,

Monte Carlo integration over all single isolate scores would not have recovered a significant

difference between the two groups. Yet, for *Bd*GPL MODS28 and *Bd*CAPE TF5a1, multi-

genus consortia yielded by far the highest probability of observing strong inhibition of 50% or

more. That this pattern was not conserved for *Bd*GPL SFBC019 is perhaps the most

intriguing finding. As for *Bd*GPL variants JEL423 and AUL2 in the phylogenetic trials,

SFBC019 was largely resistant to inhibition, with individual bacterial inhibition scores that

were often negative. One possible explanation for the lack of efficacy of consortia against

SFBC019 is that the when a variant of *B. dendrobatidis* is resistant to inhibition and/or there

is little variation in inhibition, co-culture fails to produce any synergistic inhibitory effects. That

is, if a pathogen is highly resistant to most bacterial metabolites in the first instance,

increases in the relative concentrations of those metabolites through co-culture-mediated

competition are unlikely to elicit any significant increases in inhibitory capability. The most

important consequence of this pattern is that for some pathogenic variants, multi-genus

consortia are highly unlikely to be able to yield high inhibition in cases where individual

isolates have failed to do so. Despite the observed variance in success of consortia across

B. dendrobatidis isolates, our simulation trials revealed that multi-genus consortia offer the

best broad-spectrum protection across multiple *B. dendrobatidis* isolates encountered in

series. This finding is important; human-mediated spread of *B. dendrobatidis* through the

amphibian trade (Fisher & Garner 2007) means we cannot assume that local populations will

be exposed to only one pathogenic variant. Successful mitigation of the pathogen in the wild

demands that we employ strategies with the highest broad-spectrum success over multiple

pathogen genotypes.

Conclusion

This study adds to a growing body of evidence suggesting that diverse, multi-species

consortia may represent powerful disease mitigation tools, offering superior probiotic

protection against disease compared to single bacterial isolates. Our work has highlighted

that different isolates of a pathogen can modify the strength of inhibition caused by the

probiotic, meaning we cannot expect probiotic effectiveness to be uniform across the genetic

landscape of the pathogen. Despite the relative merits of multi-genus consortia for mitigating

single and multiple *B. dendrobatidis* variants, it remains to be determined how readily these

consortia will be able to colonise the host skin *in vivo*. This is crucial for to being able to

quantify how applicable inhibition measures derived *in vitro* are to real-world scenarios.

Nevertheless, our data highlight the merits of a community-level approach to probiotic

mitigation of wildlife disease, which may offer more broad-spectrum host protection in the

face of large-scale heterogeneity in pathogen genotype.

ACKNOWLEDGEMENTS

This study was partly funded by a North-West University Postdoctoral Research Fellowship awarded to REA. XAH was funded by an Institute of Zoology Research Fellowship. The authors would like to thank Prof. Richard Preziosi and Dr. Trenton Garner for additional provision of consumables, and Prof. Ché Weldon, Dr. Trenton Garner and Prof. Matthew Fisher for access to *Batrachochytrium dendrobatidis* isolates used in this study.

REFERENCES

Antwis, R.E., Preziosi, R.F., Harrison, X.A., Garner, T.W. (2015). Amphibian symbiotic

bacteria do not show a universal ability to inhibit growth of the global panzootic lineage of

Batrachochytrium dendrobatidis. Applied and Environmental Microbiology, 81, 3706-3711.

Bates, D., Maechler, M., Bolker, B., Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. Journal of Statistical Software, 67, 1-48. doi:10.18637/jss.v067.i01.

Becker, M.H., Walke, J.B., Murrill, L., Woodhams, D.C., Reinert, L.K., Rollins-Smith, L.A., et

al. (2015). Phylogenetic distribution of symbiotic bacteria from Panamanian amphibians that

inhibit growth of the lethal fungal pathogen Batrachochytrium dendrobatidis. Molecular

Ecology, 24, 1628-1641.

Bell, S.C., Alford, R.A., Garland, S., Padilla, G., Thomas, A.D. (2013). Screening bacterial

metabolites for inhibitory effects against Batrachochytrium dendrobatidis using a

spectrophotometric assay. Diseases of Aquatic Organisms, 103, 77-85.

Bletz, M.C., Loudon, A.H., Becker, M.H., Bell, S.C., Woodhams, D.C., Minbiole, K.P. et al.

(2013). Mitigating amphibian chytridiomycosis with bioaugmentation: characteristics of

effective probiotics and strategies for their selection and use. Ecology letters, 16, 807-20.

Bletz, M.C., Goedbloed, D.J., Sanchez, E., Reinhardt, T., Tebbe, C.C., Bhuju, S., et al.

(2016). Amphibian gut microbiota shifts differentially in community structure but converges

on habitat-specific predicted functions. Nature Communications, 7.

Cohen, J. M., Venesky, M. D., Sauer, E. L., Civitello, D. J., McMahon, T. A., Roznik, E. A. et

al. (2017). The thermal mismatch hypothesis explains host susceptibility to an emerging

infectious disease. Ecology Letters, 20, 184–193. doi:10.1111/ele.12720

de Jong, M.D., Hien, T.T. (2006). Avian influenza A (H5N1). Journal of Clinical Virology, 35,

2-13.

Dillon, R.J., Vennard, C.T., Buckling, A., Charnley, A.K. (2005). Diversity of locust gut

bacteria protects against pathogen invasion. Ecology Letters, 8, 1291-8.

Doddington, B.J., Bosch, J., Oliver, J.A., Grassly, N.C., Garcia, G., Schmidt, B.R., et al.

(2013), Context-dependent amphibian host population response to an invading pathogen.

Ecology, 94, 1795-1804.

Eisenhauer, N., Scheu, S., Jousset, A. (2012). Bacterial diversity stabilizes community

productivity. PloS one, 7, e34517.

Eisenhauer, N., Schulz, W., Scheu, S., Jousset, A., Pfrender, M. (2013). Niche

dimensionality links biodiversity and invasibility of microbial communities. Functional

Ecology, 27, 282-8.

Farrer, R.A., Weinert, L.A., Bielby, J., Garner, T.W., Balloux, F., Clare, F., et al. (2011)

Multiple emergences of genetically diverse amphibian-infecting chytrids include a globalized

hypervirulent recombinant lineage. Proceedings of the National Academy of Sciences of the

United States of America, 108, 18732-6.

Farrer, R.A., Henk, D.A., Garner, T.W., Balloux, F., Woodhams, D.C., Fisher, M.C. (2013).

Chromosomal copy number variation, selection and uneven rates of recombination reveal

cryptic genome diversity linked to pathogenicity. PLoS Genetics, 9, p. e1003703.

Fisher, M.C., Garner, T.W.J. (2007) The relationship between the emergence of

Batrachochytrium dendrobatidis, the international trade in amphibians and introduced

amphibian species. Fungal Biology Reviews, 21, 2-9.

Fisher, M.C., Bosch, J., Yin, Z., Stead, D.A., Walker, J., Selway, L., et al. (2009) Proteomic

and phenotypic profiling of the amphibian pathogen Batrachochytrium dendrobatidis shows

that genotype is linked to virulence. Molecular ecology, 18, 415-29.

Fisher, M.C., Henk, D.A., Briggs, C.J., Brownstein, J.S., Madoff, L.C., McCraw, S.L., et al.

(2012). Emerging fungal threats to animal, plant and ecosystem health. Nature Reviews,

484, 186-194.

Garbaye, J. (1994). Helper bacteria: a new dimension to the mycorrhizal symbiosis. New

Phytologist, 128, 197-210.

Garner, T.W., Schmidt, B.R., Martel, A., Pasmans, F., Muths, E., Cunningham, A.A., et al.

(2016). Mitigating amphibian chytridiomycoses in nature. Phil. Trans. R. Soc. B, 371,

20160207.

Geweke, J. (1992) Evaluating the accuracy of sampling-based approaches to calculating

posterior moments. In *Bayesian Statistics 4* (ed JM Bernado, JO Berger, AP Dawid and AFM

Smith). Clarendon Press, Oxford, UK.

Goka, K., Yokoyama, J., Une, Y., Kuroki, T., Suzuki, K., Nakahara, M., et al. (2009).

Amphibian chytridiomycosis in Japan: distribution, haplotypes and possible route of entry into

Japan. Molecular Ecology, 18, 4757-74.

Hadfield, J.D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed Models:

The MCMCglmm R Package. Journal of Statistical Software, 33(2), 1-22. URL:

http://www.jstatsoft.org/v33/i02/.

Hardoim, P.R., van Overbeek, L.S., Berg, G., Pirttilä, A.M., Compant, S., Campisano, A., et

al. (2015). The hidden world within plants: ecological and evolutionary considerations for

defining functioning of microbial endophytes. Microbiology and Molecular Biology Reviews,

79, 293-320. doi:10.1128/MMBR.00050-14.

Harris, R.N., Lauer, A., Simon, M.A., Banning, J.L., Alford, R.A. (2009). Addition of antifungal

skin bacteria to salamanders ameliorates the effects of chytridiomycosis. Diseases of Aquatic

Organisms, 83, 11-16.

Hoyt, J.R., Cheng, T.L., Langwig, K.E., Hee, M.M., Frick, W.F., Kilpatrick, A.M. (2015).

Bacteria isolated from bats inhibit the growth of Pseudogymnoascus destructans, the

causative agent of white-nose syndrome. PLoS One, 10, e0121329.

Jani, A.J., Briggs, C.J. (2014). The pathogen Batrachochytrium dendrobatidis disturbs the

frog skin microbiome during a natural epidemic and experimental infection. Proceedings of

the National Academy of Sciences of the United States of America, 111, E5049-58.

Jenkinson, T.S., Betancourt Román, C.M., Lambertini, C., Valencia-Aguilar, A., Rodriguez,

D., Nunes-de-Almeida, C.H., et al. (2016) Amphibian-killing chytrid in Brazil comprises both

locally endemic and globally expanding populations. Molecular Ecology, 25, 2978–2996.

Jousset, A., Schmid, B., Scheu, S., Eisenhauer, N. (2011). Genotypic richness and

dissimilarity opposingly affect ecosystem functioning. Ecology Letters, 14, 537-45.

Kueneman, J.G., Woodhams, D.C., Harris, R., Archer, H.M., Knight, R., McKenzie, V.J.,

(2016) Probiotic treatment restores protection against lethal fungal infection lost during

amphibian captivity. Proc. R. Soc. B, 283, 1839.

Martel, A., Spitzen-van der Sluijs, A., Blooi, M., Bert, W., Ducatelle, R., Fisher, M.C., et al.

(2013). Batrachochytrium salamandrivorans sp. nov. causes lethal chytridiomycosis in

amphibians. Proceedings of the National Academy of Sciences, 110, 15325-15329.

Martel, A., Blooi, M., Adriaensen, C., Van Rooij, P., Beukema, W., Fisher, M.C., et al. (2014).

Recent introduction of a chytrid fungus endangers Western Palearctic

salamanders. Science, 346, 630-631.

Matos, A., Kerkhof, L., Garland, J.L. (2005). Effects of microbial community diversity on the

survival of Pseudomonas aeruginosa in the wheat rhizosphere. Microbial Ecology, 49, 257-

64.

McCallum, H. (2012) Disease and the dynamics of extinction. Philos Trans R Soc Lond B

Biol Sci 367: 2828–2839. doi: 10.1098/rstb.2012.0224. pmid:22966138

Morgan, J.A., Vredenburg, V.T., Rachowicz, L.J., Knapp, R.A., Stice, M.J., Tunstall, T., et al.

(2007). Population genetics of the frog-killing fungus *Batrachochytrium dendrobatidis*.

Proceedings of the National Academy of Sciences of the United States of America, 104,

13845-50.

Muletz, C.R., Myers, J.M., Domangue, R.J., Herrick, J.B., Harris, R.N. (2012). Soil

bioaugmentation with amphibian cutaneous bacteria protects amphibian hosts from infection

by Batrachochytrium dendrobatidis. Biological Conservation, 152, 119-126.

Olson, D.H., Aanensen, D.M., Ronnenberg, K.L., Powell, C.I., Walker, S.F., Bielby, J., et al.

(2013). Mapping the global emergence of Batrachochytrium dendrobatidis, the amphibian

chytrid fungus. PloS one, 8, e56802.

Piovia-Scott, J., Rejmanek, D., Woodhams, D. C., Worth, S. J., Kenny, H., McKenzie, V., et

al. (2017). Greater Species Richness of Bacterial Skin Symbionts Better Suppresses the

Amphibian Fungal Pathogen Batrachochytrium Dendrobatidis. Microbiology Ecology, DOI

10.1007/s00248-016-0916-4.

R Core Team. (2016). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rebollar, E.A., Antwis, R.E., Becker, M.H., Belden, L.K., Bletz, M.C., Brucker, R.M., et al.

(2016). Using "Omics" and Integrated Multi-Omics Approaches to Guide Probiotic Selection

to Mitigate Chytridiomycosis and Other Emerging Infectious Diseases. Frontiers in

Microbiology, 7, 68.

Rodriguez, D., Becker, C. G., Pupin, N.C., Haddad, C.F.B., Zamudio, K. R. (2014). Long-

term endemism of two highly divergent lineages of the amphibian-killing fungus in the Atlantic

Forest of Brazil. Molecular Ecology, 23, 774-87.

Romanowski, K., Zaborin, A., Fernandez, H., Poroyko, V., Valuckaite, V., Gerdes, S., et al.

(2011). Prevention of siderophore- mediated gut-derived sepsis due to P. aeruginosa can be

achieved without iron provision by maintaining local phosphate abundance: role of pH. BMC

Microbiology, 11: 212

Rosenblum, E.B., James, T.Y., Zamudio, K.R., Poorten, T.J., Ilut, D., Rodriguez, D., et al.

(2013). Complex history of the amphibian-killing chytrid fungus revealed with genome

resequencing data. Proceedings of the National Academy of Sciences of the United States of

America, 110, 9385-90.

Scheuring, I., Yu, D.W. (2012). How to assemble a beneficial microbiome in three easy

steps. Ecology Letters, 15, 1300-7.

620 Schock, D.M., Bollinger, T.K., Collins, J.P. (2009). Mortality rates differ among amphibian

621 populations exposed to three strains of a lethal ranavirus. EcoHealth, 6, 438-48.

623 Schloegel, L.M., Toledo, L.F., Longcore, J.E., Greenspan, S.E., Vieira, C.A., Lee, M., et al.

624 (2012) Novel, panzootic and hybrid genotypes of amphibian chytridiomycosis associated with

625 the bullfrog trade. Molecular Ecology, 21, 5162-77.

627 Skerratt, L.F., Berger, L., Speare, R., Cashins, S., McDonald, K.R., et al. (2007) Spread of

628 chytridiomycosis has caused the rapid global decline and extinction of frogs. Ecohealth 4:

629 125–134.

631 Sleeman, J.M. (2013). Has the time come for big science in wildlife health? EcoHealth, 10,

632 335-338.

634 Tompkins, D.M., Carver, S., Jones, M.E., Krkošek, M., Skerratt, L.F. (2015). Emerging

635 infectious diseases of wildlife: a critical perspective. Trends in Parasitology, 31, 149-159.

637 van Elsas, J.D., Chiurazzi, M., Mallon, C.A., Elhottova, D., Kristufek, V., Salles, J.F. (2012).

638 Microbial diversity determines the invasion of soil by a bacterial pathogen. Proceedings of

639 the National Academy of Sciences of the United States of America, 109, 1159-64.

641 Walke, J.B., Becker, M.H., Loftus, S.C., House, L.L., Teotonio, T.L., Minbiole, K.P. et al.

642 (2015). Community Structure and Function of Amphibian Skin Microbes: An Experiment with

643 Bullfrogs Exposed to a Chytrid Fungus. PloS one, 10, e0139848.

Woodhams, D.C., Brandt, H., Baumgartner, S., Kielgast, J., Küpfer, E., Tobler, U., et al.

(2014). Interacting symbionts and immunity in the amphibian skin mucosome predict disease

risk and probiotic effectiveness. *PLoS One*, *9*(4), p.e96375.

Table 1

Batrachochytrium dendrobatidis isolates used in the study.

Table 2

Composition of multi-genus consortia used in the study. Single-genus consortia comprised all

three bacterial isolates (A, B and C) for a given genus (*Acinetobacter, Chryseobacterium,*

Serratia, Stentrophomonas).

Table 3

Mean Proportion of 10 *Bd*GPL isolates for which at least weak inhibitory capability was

observed, averaged over all bacterial isolates in a genus. 95% CI: 95% confidence intervals

from an overdispersion-corrected Binomial GLM.

Figure 1. Inhibition scores of 58 bacterial strains from 10 genera when tested against 10 *Bd*GPL isolates.

Estimates are derived from a Bayesian mixed effects model with bacterial isolate nested within genus, and *Bd*GPL isolate fitted as random effects. Points are conditional modes of the individual isolate random effects, marginalised with respect to *Bd*GPL isolate. Error bars are 95% credible intervals.

Figure 2. Inhibition scores of 10 *Bd*GPL isolates

Estimates are derived from a Bayesian mixed effects model with bacterial isolate nested

within genus, and *Bd*GPL isolate fitted as random effects. Points are conditional modes of

the individual *Bd*GPL isolate random effects, marginalised with respect to bacterial isolate.

Error bars are 95% credible intervals.

688 **Figure 3.** Inhibition scores for Single-Genus and Multi-Genus Consortia across three *B.*

689 *dendrobatidis* isolates (*Bd*GPL MODS28.1, *Bd*GPL SFBC019 and *Bd*CAPE TF5a1).

690 Points have been jittered for display purposes.

Figure 4. Simulation results examining the relative efficacy of different probiotic strategies

(A) the probability of Multi-Genus Consortia (MGC) yielding higher inhibition compared to Single-Genus Consortia (SGC) or a Single Bacterial Isolate (Single); (B) the probability of MGC, SGC or Single bacteria yielding inhibition > 50% when applied to each of three *B. dendrobatidis* isolates; (C) The probability of an individual consortium type yielding >50% inhibition when applied to two randomly chosen *B. dendrobatidis* isolates in series.

Isolate	Lineage	Geographic origin	Host species	Collector	Year	Phylogeny screening
MG04	GPL	Silver Mine, Western Cape, South Africa	*Amietia fuscigula*	Trenton Garner	2010	X
CORN2.2	GPL	Penhale Farm, Cornwall, UK	*Ichthyosaurus alpestris*	Trenton Garner	2012	X
CORN2.3	GPL	Penhale Farm, Cornwall, UK	*Ichthyosaurus alpestris*	Trenton Garner	2012	X
CORN3.1	GPL	Penhale Farm, Cornwall, UK	*Ichthyosaurus alpestris*	Trenton Garner	2012	X
CORN3.2	GPL	Penhale Farm, Cornwall, UK	*Ichthyosaurus alpestris*	Trenton Garner	2012	X
AUL1.2	GPL	Lac d'Aule, France	*Alytes obstetricans*	Matthew Fisher	2010	X
AUL2	GPL	Lac d'Aule, France	*Alytes obstetricans*	Matthew Fisher	2010	X
IA2011	GPL	Ibon Acherito, Spain	*Alytes obstetricans*	Matthew Fisher	2011	X
MODS 28.1	GPL	Mont Olia, Sardinia	*Discoglossus sardus*	Trenton Garner	2010	X
JEL423	GPL	Guabal, Panama	*Agalychnis lemur*	Joyce Longcore	2004	X
SFBC019	GPL	Sellafield, Cumbria, UK	*Epidalea calamita*	Peter Minting	2010	
TF5a1	CAPE	Torrent des Ferrerets, Mallorca	*Alytes muletensis*	Matthew Fisher	2007	

Consortium challenges
X
X
X

Inter-genera mix 1	Inter-genera mix 2	Inter-genera mix 3	Inter-genera mix 4
Chryseobacterium B	*Acinetobacter C*	*Acinetobacter A*	*Acinetobacter B*
Serratia B	*Serratia C*	*Chryseobacterium A*	*Chryseobacterium C*
Stentrophomonas B	*Stentrophomonas C*	*Serratia A*	*Stentrophomonas A*

Genus	Number Isolates	Mean Proportion *BdGPL* Inhibition	95% CI
Acinetobacter	6	0.33	0.1-0.65
hryseobacteriu	8	0.5	0.24-0.76
Citrobacter	3	0.67	0.24-0.95
Comamonas	4	0.7	0.32-0.95
Enterobacter	11	0.54	0.31-0.75
Microbacterium	4	0.4	0.1-0.77
Sanguibacter	3	0.63	0.21-0.94
Serratia	6	0.47	0.19-0.76
Staphylococcus	4	0.73	0.34-0.96
enotrophomon	9	0.49	0.25-0.73

Figure 1

Figure 3

Figure 4

A

B

C