A pleiotropic chemoreceptor facilitates the coupling of pheromonal signal perception

and production

Kathleen M. Zelle[1,†], Cassondra Vernier[1,†], Xitong Liang[2], Sean Halloran[3], Jocelyn G.

Millar[3], Yehuda Ben-Shahar [1,4,*]

[1]Department of Biology, Washington University in Saint Louis, 1 Brookings Drive, Saint

Louis, MO 63130 USA.

[2]Department of Neuroscience, Washington University School of Medicine, 660 South

Euclid Avenue, St. Louis, MO 63110, USA

[3]Department of Entomology, University of California, Riverside, 900 University Avenue,

Riverside, CA 92521, USA

[4]Lead Contact

[*]Correspondence: benshahary@wustl.edu (Y.B.)

[†]Denotes equal contribution

SUMMARY

Optimal mating decisions depend on reliable and precise species-specific coupling of the production and perception of mating communication signals because any functional changes in either the signal or its perception should carry a fitness cost (Boake, 1991; Brooks et al., 2005; Hoy et al., 1977b; Shaw et al., 2011; Shaw and Lesnick, 2009; Steiger et al., 2010; Sweigart, 2010; Symonds and Elgar, 2008; Wyatt and Cambridge University Press, 2014). Previously published theoretical models predict that genetic linkage between sensory-driven mate choices and specific communication signals could provide a robust mechanism for maintaining robust and stable signal production and perception at the level of populations, while still retaining the capacity for future signal diversification (Hoy et al., 1977b; Shaw et al., 2011; Shaw and Lesnick, 2009; Wiley et al., 2011). Several experimental studies have provided empirical evidence that pleiotropy is one possible genetic solution for maintaining functional coupling between specific communication signals and the mate choice behaviors they elicit at the population level (Hoy et al., 1977b; Shaw et al., 2011; Shaw and Lesnick, 2009; Wiley et al., 2011). However, the complex characteristics of mating signals and behaviors present a major barrier for identifying the actual pleiotropic genes and molecular pathways that provide genetic linkages between the production and perception of mating signals (Chenoweth and Blows, 2006; Singh and Shaw, 2012). Here we show that in the fruit fly *Drosophila melanogaster*, some pheromone-driven mating choices are coupled to the production of specific mating signals via pleiotropic chemoreceptors. Specifically, we demonstrate that *Gr8a*, a member of the gustatory receptor family, independently contributes to the perception of inhibitory mating signals in pheromone sensing neurons, as well as in regulating the production of inhibitory mating signals in oenocytes. Together, these data provide a relatively simple molecular explanation for how coupling of pheromone perception and production, two independent physiological processes, remains robust.

Keywords: *Drosophila melanogaster;* Vinegar fly; Fruit fly; Cuticular hydrocarbons; *Gr8a*; Oenocyte.

RESULTS AND DISCUSSION

Similar to other insect species, the fruit fly *Drosophila melanogaster* has evolved cuticular hydrocarbons (CHCs), synthesized by the fat body and oenocytes (Billeter et al., 2009; Gutierrez et al., 2006; Krupp et al., 2013; Makki et al., 2014), which primarily provide a hydrophobic desiccation barrier, as well as play an important role as pheromones that regulate diverse aspects of insect behavior (Blomquist and Bagnères, 2010; Chung and Carroll, 2015; Ferveur, 2005; Howard and Blomquist, 2005). Specifically, complex blends of CHCs are often utilized by insect species to communicate sex identity and female mating status, and well as defining the behavioral reproductive boundaries between closely related species (Billeter et al., 2009; Chung and Carroll, 2015; Chung et al., 2014; Coyne et al., 1994; Dweck et al., 2015; Ng et al., 2014; Shirangi et al., 2009; Yew and Chung, 2015). In *Drosophila*, the perception of volatile CHCs is mediated by olfactory sensory neurons located in the antennae and maxillary palps (Benton et al., 2007; Kurtovic et al., 2007; Lebreton et al., 2014; van der Goes van Naters and Carlson, 2007), while less volatile CHCs, which require contact, are sensed by specialized gustatory-like receptor neurons (GRNs) in the appendages (legs and wings) and the proboscis (Koh et al., 2014; Lu et al., 2012; Lu et al., 2014; Thistle et al., 2012; Toda et al., 2012). While some of the genes and pathways that contribute to CHC synthesis in *Drosophila* are known, the molecular identities of the majority of CHC receptors remain unknown. Previous work suggested that the gene *Desat1*, which encodes a fatty acid desaturase enriched in oenocytes, might also independently contribute to CHC perception (Bousquet et al., 2012). However, due to the expression of *Desat1* in central neurons (Billeter et al., 2009) and the broad impact its mutant alleles have on the CHC profiles of both males and females (Labeur et al., 2002), whether *Desat1* directly contributes to mating signal perception remains unresolved.

Consequently, we chose to examine members of the *Gustatory receptor* (*Gr*) gene family as candidates for pleiotropic factors that might contribute directly to both perception and production of pheromonal mating signals in *Drosophila*. Because several family members have already been implicated in the detection of specific excitatory and inhibitory pheromones (Bray and Amrein, 2003; Hu et al., 2015; Miyamoto and Amrein,

2008; Moon et al., 2009; Watanabe et al., 2011), and the majority of genes that encode

family members are already known to be enriched in gustatory receptor neurons

(GRNs) (Clyne et al., 2000; Dunipace et al., 2001; Scott et al., 2001; Wang et al., 2004),

we reasoned that any pleiotropic *Gr* genes should also be expressed in abdominal

oenocytes (Billeter et al., 2009). Subsequently, an RT-PCR screen revealed that 24 out

of the 60 genes that encode *Gr* family members in the *Drosophila* genome (Clyne et al.,

2000; Dunipace et al., 2001; Robertson et al., 2003; Scott et al., 2001) exhibit enriched

expression in abdominal tissues (Table S1).

Although several members of the *Gr* receptor family, including *Gr68a, Gr32a, Gr66a,*

Gr39a, and *Gr33a*, have been previously linked to the sensory perception of mating

pheromones (Bray and Amrein, 2003; Lacaille et al., 2009; Miyamoto and Amrein, 2008;

Moon et al., 2009; Watanabe et al., 2011), none of these candidate pheromone

receptors were identified in our initial screen for *Gr* genes with enriched mRNA

expression in abdominal tissues of either males or females (Table S1). However, *Gr8a*,

which we have identified as a male-specific gene in our screen, is localized to a

chromosomal region that was previously implicated in chemoreception with enriched

expression in abdomens (Table S1) and was previously shown to specifically contribute

to the sensory detection of the non-proteinogenic amino acid L-Canavanine (Lee et al.,

2012; Shim et al., 2015). Because our initial expression screen was based on whole-

abdomen RNAs, and it was previously reported that *Gr8a* is also expressed in some

abdominal neurons (Park and Kwon, 2011), we next used a GAL4 transgenic driver to

examine the spatial expression pattern of *Gr8a* in males and females. As was

previously reported, we found that *Gr8a* is expressed in 14-16 GRNs in the proboscis

(Figure 1A) (Lee et al., 2012) as well as in two paired GRNs in the foreleg pretarsus

(Figure 1B) of males and females. In addition, the *Gr8a*-GAL4 reporter was also highly

expressed in abdominal oenocyte-like cells in males but not females (Figure 1C). The

male-specific abdominal expression pattern was further supported by qRT-PCR (Figure

1D). Using immunohistochemistry, we then demonstrated that *Gr8a* is co-expressed

with the oenocyte marker *Desat1 (Billeter et al., 2009)*, as well as in *Desat1*-negative

cells with fat-body-like morphology (Figure 1E-G).

Together, the enriched expression of *Gr8a* in foreleg GRNs, a primary pheromone

chemosensory organ (Lu et al., 2012; Lu et al., 2014), and its sexually dimorphic

expression in oenocytes suggested that in addition to its role in L-Canavanine

perception, *Gr8a* also contributes to mating behaviors. Since mate choices of both male

and female flies are determined by a complex blend of excitatory and inhibitory

pheromones (Billeter et al., 2009; Krupp et al., 2013), we anticipated that if *Gr8a* is

indeed a pleiotropic factor that independently contributes to the perception and

production of specific components of the male mating pheromone bouquet, then

disruptions of the signal production in males or its perception in females should carry

similar impacts on female mate choice behavior. To directly test this hypothesis, we first

investigated whether *Gr8a* and the GRNs that express it are required for sensory

functions associated with female mate choice by using a single-pair courtship paradigm

(Lu et al., 2012; Lu et al., 2014). We found that blocking neuronal transmission in *Gr8a*-

expressing GRNs with the transgenic expression of the tetanus toxin (TNT) in females

resulted in shorter copulation latency relative to wild-type females when courted by wild-

type males (Figure 2A). Similarly, homozygous (Figure 2B) and hemizygous (Figure 2C)

Gr8a mutant females exhibited shorter copulation latencies relative to wild-type controls,

which could be rescued by the transgenic expression of a *Gr8a* cDNA (Figure 2D). In

contrast, similar genetic manipulations of the *Gr8a* gene or *Gr8a*-expressing GRNs in

males did not affect courtship behavior as measured by courtship latency and index

towards wild-type females (Figure S1). These data indicate that *Gr8a* is required for

female mate choice via the detection of male-borne inhibitory mating signals.

Because *Gr8a* expression is specifically enriched in male oenocytes, we next tested the

hypothesis that *Gr8a* also plays a role in the production and/or release of inhibitory

mating signals by males. Similar to the effect of the *Gr8a* mutation in females, we found

that wild-type virgin females exhibited shorter copulation latencies towards *Gr8a* mutant

males relative to wild-type controls (Figure 2E). These data indicate that *Gr8a* mutant

males produce and/ or release abnormally low levels of inhibitory mating pheromones.

Previous studies showed that, in order to increase their fitness, *Drosophila* males

transfer inhibitory mating pheromones to females during copulation, which subsequently

lowers the overall attractiveness of mated females to other males and the probability of

females remating (Benton et al., 2007; Billeter et al., 2009; Datta et al., 2008; Yew et al.,

2009). Because our data indicate that *Gr8a* plays a role in the production of an inhibitory

signal in males, we next tested the hypothesis that *Gr8a* mutant males would have a

reduced ability to produce and/ or transfer inhibitory pheromones to females during

copulation. Accordingly, we found that wild-type males failed to recognize mating status

of wild-type females that were previously mated with *Gr8a* mutant males (Figure 2F).

These data indicate that *Gr8a* action is required for the production and/or transfer of

inhibitory pheromones from males to females during copulation. Because *Gr8a* is

involved in the perception of an inhibitory mating signal, we also tested the hypothesis

that *Gr8a* mutant males will exhibit abnormal capacity for identifying the post-mating

status of wild-type females. Indeed, we observed that *Gr8a* mutant males were more

likely than wild type males to court a mated female, suggesting that in addition to its role

in the production of inhibitory mating signals in males, *Gr8a* is also required in males for

the sensory recognition of mating inhibitory signals that mark the post-mating status of

females (Figure 2F). Together, these studies indicate that *Gr8a* is required in males for

the production and perception of transferrable inhibitory mating signals that advertise

post-mating status in females. These data further support the pleiotropic role of the

Gr8a gene in the production of inhibitory mating signals in males and the perception of

these signals in both males and females in sex-specific contexts.

Because our behavioral data indicate that the *Gr8a* mutation has a dramatic impact on

the production and perception of a putative inhibitory mating pheromonal signal, we next

examined whether the *Gr8a* mutation has a direct impact on qualitative and quantitative

aspects of male and mated-female CHC profiles. As predicted by our behavioral data,

analyses of male epicuticular pheromones by gas chromatography and mass

spectroscopy revealed the *Gr8a* mutation has a significant impact on the overall

qualitative characteristics of the CHC profile of males (Figure 3A). Additional analyses

revealed significant quantitative effects of the *Gr8a* mutation on the cuticular levels of

several specific components in males (Figure 3B,C and Table S2). In particular, the

Gr8a mutation affects the levels of several alkenes and methyl-branched alkanes, which

have been implicated in mate choice behaviors in diverse *Drosophila* species (Billeter et

al., 2009; Chung et al., 2014; Dyer et al., 2014; Shirangi et al., 2009). Similarly, *Gr8a*

mutation affects the expression level of several desaturase enzymes in males (Figure

3D), which are involved in the biosynthesis of alkenes (Chung et al., 2014). Additionally,

although the CHC profile of females mated with *Gr8a* mutant males did not qualitatively

differ from those mated with wild-type males (Figure 3E), one specific component of the

CHC profile, nonacosane (C_{29}), differed quantitatively between the two mated-female

groups (Figure 3F). Together, our behavioral and pheromonal data indicate that *Gr8a*

action contributes to mating decisions in females by co-regulating the female perception

and male production and transfer of an inhibitory pheromone, which is consistent with a

pleiotropic function for *Gr8a*.

The data presented here demonstrate that a pleiotropic chemoreceptor co-regulates the

perception and production of a specific pheromonal signal that plays an important role in

mate choice behaviors of both sexes. In contrast to its expected function in the

perception of chemical ligands, how *Gr8a*, a member of a canonical chemoreceptor

family might also contribute to the production of pheromonal signals is not as obvious.

In some more well-understood secretory cell types, autoreceptors are essential for the

regulation of synthesis and secretion rates. For example, dopaminergic and

serotonergic cells regulate rates of synthesis and release of their respective

neuromodulators by the action of autoreceptors, which are used as sentinels that

regulate synthesis rates via specific molecular signaling feedbacks in response to level

changes in the extracellular concentrations of the secreted molecule (Ford, 2014;

Stagkourakis et al., 2016). Therefore, we hypothesize that one possible way *Gr8a* could

regulate the synthesis and/or secretion of specific CHCs is by acting as an oenocyte-

intrinsic autoreceptor, which integrates feedback information to the complex genetic

network that regulates the epicuticular CHC profile of males (Fig. 4).

Previous work indicated that *Gr8a* is expressed in "bitter" taste neurons in the

proboscis, and specifically required for the sensory perception of the feeding deterrent

L-canavanine (Lee et al., 2012; Shim et al., 2015), but not for the detection of other

inhibitory chemicals such as caffeine, strychnine, and umbelliferone (Lee et al., 2009;

Poudel et al., 2015). Our data indicate that *Gr8a* contributes to inhibitory functions

associated with both feeding and mating decisions, as was previously reported for other

'bitter' receptors in *Drosophila* (Lacaille and Hiroi, 2007; Moon et al., 2009). *Gr8a-*

depedent perception of L-canavanine seems to depend on its heterotrimeric interaction

with *Gr66a* and *Gr98b* in bitter sensing neurons in the proboscis (Shim et al., 2015).

Although both of these *Gr* genes were also identified in our initial screen for receptors

enriched in the abdomen, *Gr66a* was expressed in both males and females while *Gr98b*

was specifically enriched in females (Table S1). These data suggest that *Gr8a*-

depedent contributions to feeding and mating decisions are independently mediated via

interactions with different *Gr* genes in each sensory context, and further point to plastic

subunit composition as an essential aspect of the mechanism for ligand specificity in

insect gustatory receptors. Although we do not know yet the specific chemical identity of

the ligand of *Gr8a*, previous studies indicated that at least two inhibitory mating

pheromones, cVA and CH503, are transferred from males to females during

copulations. Although our data suggest that the *Gr8a* mutation affects the level of cVA

expressed by males, it is not likely that either cVA or CH503 are the putative *Gr8a*

ligands because the volatile cVA acts primarily via the olfactory receptor *Or67d* (Benton

et al., 2007; Datta et al., 2008; Kurtovic et al., 2007), and CH503 has been reported to

signal via *Gr68a*-expressing neurons, which are anatomically distinct from the *Gr8a*

GRNs we describe here (Figure 1A-B) (Shankar et al., 2015; Yew et al., 2009).

Therefore, we anticipate that future studies will identify additional male CHCs that

function as inhibitory mating signals.

As populations diversify, both pheromonal signals and their receptors have to co-evolve

to maintain fitness via mate choice (Boake, 1991; Symonds and Elgar, 2008). However,

because pheromone production and perception are anatomically and physiologically

distinct biological processes, how they might be linked genetically at the molecular and

physiological levels remains unknown for most studied mating communication systems.

Several theoretical models have previously stipulated that one possible solution to this

conundrum is a genetic linkage between signal production and associated mate choice

behaviors, which supports the co-evolution of mating signals with their cognate

receptors (Arnold et al., 1996; Lande, 1981; Svensson et al., 2007), *via* mechanisms

such as pleiotropy and the co-segregation of linked alleles (Boake, 1991; Bousquet et

al., 2012; Butlin and Ritchie, 1989; Lofstedt et al., 1989; Shaw et al., 2011; Singh and

Shaw, 2012). The studies we present here do not directly address the possible

contribution of *Gr8a* to the rapid evolution of mating signals across *Drosophila*.

Nevertheless, phylogenetic analyses of *Gr8a* orthologs indicate that this is a conserved,

sexually dimorphic receptor across *Drosophila* (Figure S2A-B). Furthermore, analysis of

the *Gr8a* protein alignment revealed that in spite of the high overall sequence

conservation, the *Gr8a* protein includes at least one phylogenetically variable domain,

(Figure S2C-D).

Studies in other animal species suggest that receptor pleiotropy might also play a role in

mating communication via other sensory modalities including auditory communication in

crickets (Hoy et al., 1977a; Wiley et al., 2011) and visual communication in fish

(Fukamachi et al., 2009). While the specific genes and signaling pathways that mediate

communication in these species are mostly unknown, these data suggest that the

genetic coupling of signal-receptor pairs in mating communication systems might be

more common than previously thought. Therefore, the genetic tractability of *D.*

melanogaster, in combination with the diversity of mating communication systems in this

species-rich phylogenetic group, provide a unique opportunity for understanding the

evolution and mechanisms that drive and maintain mating communication systems at

the genetic, molecular, and cellular levels.

AUTHOR CONTRIBUTIONS

K.M.Z., C.V., J.G.M. and Y.B-S designed experiments. K.M.Z., C.V., X.L., S.H., J.G.M.

and Y.B-S collected and analyzed data. K.M.Z., C.V. and Y.B-S wrote the manuscript.

ACKNOWLEDGMENTS

We thank members of the Ben-Shahar lab for comments on earlier versions of the

manuscript. We thank the Millar laboratory (UC Riverside) for performing the CHC

analyses. We thank Nabeel Chowdhury for assistance with the RT-PCR screen, and

Paula Kiefel for technical help with generating transgenic flies. This work was supported

by NSF grant 1545778 and NIH grant NS089834 awarded to Y. B-S. Stocks obtained

from the Bloomington Drosophila Stock Center (NIH P40OD018537) were used in this

study. Wild type *Drosophila* species used in this study were obtained from the

Drosophila Species Stock Center at the University of California, San Diego.

References

Abascal, F., Zardoya, R., and Posada, D. (2005). ProtTest: selection of best-fit models of protein
evolution. Bioinformatics *21*, 2104-2105.
Arnold, S.J., Verrell, P.A., and Tilley, S.G. (1996). THE EVOLUTION OF ASYMMETRY IN SEXUAL
ISOLATION: A MODEL AND A TEST CASE. Evolution *50*, 1024-1033.
Benton, R., Vannice, K.S., and Vosshall, L.B. (2007). An essential role for a CD36-related receptor
in pheromone detection in Drosophila. Nature *450*, 289-293.
Billeter, J.C., Atallah, J., Krupp, J.J., Millar, J.G., and Levine, J.D. (2009). Specialized cells tag
sexual and species identity in Drosophila melanogaster. Nature *461*, 987-991.
Blomquist, G.J., and Bagnères, A.-G. (2010). Insect hydrocarbons : biology, biochemistry, and
chemical ecology. (Cambridge: Cambridge University Press).
Boake, C.R.B. (1991). Coevolution of senders and receivers of sexual signals: Genetic coupling
and genetic correlations. Trends in Ecology & Evolution *6*, 225-227.
Bousquet, F., Nojima, T., Houot, B., Chauvel, I., Chaudy, S., Dupas, S., Yamamoto, D., and
Ferveur, J.F. (2012). Expression of a desaturase gene, desat1, in neural and nonneural tissues
separately affects perception and emission of sex pheromones in Drosophila. Proc Natl Acad Sci
U S A *109*, 249-254.
Bray, S., and Amrein, H. (2003). A putative Drosophila pheromone receptor expressed in male-
specific taste neurons is required for efficient courtship. Neuron *39*, 1019-1029.
Brooks, R., Hunt, J., Blows, M.W., Smith, M.J., Bussière, L.F., Jennions, M.D., and Tregenza, T.
(2005). EXPERIMENTAL EVIDENCE FOR MULTIVARIATE STABILIZING SEXUAL SELECTION.
Evolution *59*, 871-880.
Butlin, R.K., and Ritchie, M.G. (1989). Genetic coupling in mate recognition systems: what is the
evidence? Biological Journal of the Linnean Society *37*, 237-246.
Chenoweth, S.F., and Blows, M.W. (2006). Dissecting the complex genetic basis of mate choice.
Nat Rev Genet *7*, 681-692.
Chung, H., and Carroll, S.B. (2015). Wax, sex and the origin of species: Dual roles of insect
cuticular hydrocarbons in adaptation and mating. BioEssays : news and reviews in molecular,
cellular and developmental biology *37*, 822-830.
Chung, H., Loehlin, D.W., Dufour, H.D., Vaccarro, K., Millar, J.G., and Carroll, S.B. (2014). A single
gene affects both ecological divergence and mate choice in Drosophila. Science *343*, 1148-1151.
Clyne, P.J., Warr, C.G., and Carlson, J.R. (2000). Candidate taste receptors in Drosophila. Science
287, 1830-1834.
Coyne, J.A., Crittenden, A.P., and Mah, K. (1994). Genetics of a pheromonal difference
contributing to reproductive isolation in Drosophila. Science *265*, 1461-1464.
Datta, S.R., Vasconcelos, M.L., Ruta, V., Luo, S., Wong, A., Demir, E., Flores, J., Balonze, K.,
Dickson, B.J., and Axel, R. (2008). The Drosophila pheromone cVA activates a sexually dimorphic
neural circuit. Nature *452*, 473-477.
Dunipace, L., Meister, S., McNealy, C., and Amrein, H. (2001). Spatially restricted expression of
candidate taste receptors in the Drosophila gustatory system. Curr Biol *11*, 822-835.
Dweck, H.K., Ebrahim, S.A., Thoma, M., Mohamed, A.A., Keesey, I.W., Trona, F., Lavista-Llanos,
S., Svatos, A., Sachse, S., Knaden, M., et al. (2015). Pheromones mediating copulation and
attraction in Drosophila. Proc Natl Acad Sci U S A *112*, E2829-2835.

Dyer, K.A., White, B.E., Sztepanacz, J.L., Bewick, E.R., and Rundle, H.D. (2014). Reproductive
character displacement of epicuticular compounds and their contribution to mate choice in
Drosophila subquinaria and Drosophila recens. Evolution *68*, 1163-1175.
Ferveur, J.F. (2005). Cuticular hydrocarbons: their evolution and roles in Drosophila pheromonal
communication. Behav Genet *35*, 279-295.
Ford, C.P. (2014). The role of D2-autoreceptors in regulating dopamine neuron activity and
transmission. Neuroscience *282*, 13-22.
Fukamachi, S., Kinoshita, M., Aizawa, K., Oda, S., Meyer, A., and Mitani, H. (2009). Dual control
by a single gene of secondary sexual characters and mating preferences in medaka. BMC
Biology *7*, 64.
Gutierrez, E., Wiggins, D., Fielding, B., and Gould, A.P. (2006). Specialized hepatocyte-like cells
regulate Drosophila lipid metabolism. Nature *445*, 275-280.
Howard, R.W., and Blomquist, G.J. (2005). ECOLOGICAL, BEHAVIORAL, AND BIOCHEMICAL
ASPECTS OF INSECT HYDROCARBONS*. Annu. Rev. Entomol. *50*, 371-393.
Hoy, R., Hahn, J., and Paul, R. (1977a). Hybrid cricket auditory behavior: evidence for genetic
coupling in animal communication. Science *195*, 82-84.
Hoy, R.R., Hahn, J., and Paul, R.C. (1977b). Hybrid cricket auditory behavior: evidence for
genetic coupling in animal communication. Science *195*, 82.
Hu, Y., Han, Y., Shao, Y., Wang, X., Ma, Y., Ling, E., and Xue, L. (2015). Gr33a modulates
Drosophila male courtship preference. Scientific reports *5*, 7777.
Koh, T.W., He, Z., Gorur-Shandilya, S., Menuz, K., Larter, N.K., Stewart, S., and Carlson, J.R.
(2014). The Drosophila IR20a clade of ionotropic receptors are candidate taste and pheromone
receptors. Neuron *83*, 850-865.
Krupp, J.J., Billeter, J.C., Wong, A., Choi, C., Nitabach, M.N., and Levine, J.D. (2013). Pigment-
dispersing factor modulates pheromone production in clock cells that influence mating in
drosophila. Neuron *79*, 54-68.
Kurtovic, A., Widmer, A., and Dickson, B.J. (2007). A single class of olfactory neurons mediates
behavioural responses to a Drosophila sex pheromone. Nature *446*, 542-546.
Labeur, C., Dallerac, R., and Wicker-Thomas, C. (2002). Involvement of desat1 gene in the
control of Drosophila melanogaster pheromone biosynthesis. Genetica *114*, 269-274.
Lacaille, F., Everaerts, C., and Ferveur, J.F. (2009). Feminization and alteration of Drosophila
taste neurons induce reciprocal effects on male avoidance behavior. Behav Genet *39*, 554-563.
Lacaille, F., and Hiroi, M. (2007). An Inhibitory Sex Pheromone Tastes Bitter for Drosophila
Males. Plos One *2*, e661.
Lande, R. (1981). Models of speciation by sexual selection on polygenic traits. Proc Natl Acad Sci
USA *78*.
Lebreton, S., Grabe, V., Omondi, A.B., Ignell, R., Becher, P.G., Hansson, B.S., Sachse, S., and
Witzgall, P. (2014). Love makes smell blind: mating suppresses pheromone attraction in
Drosophila females via Or65a olfactory neurons. Scientific reports *4*, 7119.
Lee, Y., Kang, M.J., Shim, J., Cheong, C.U., Moon, S.J., and Montell, C. (2012). Gustatory
receptors required for avoiding the insecticide L-canavanine. J Neurosci *32*, 1429-1435.
Lee, Y., Moon, S.J., and Montell, C. (2009). Multiple gustatory receptors required for the
caffeine response in Drosophila. Proc Natl Acad Sci U S A *106*, 4495-4500.

Lofstedt, C., Hansson, B.S., Roelofs, W., and Bengtsson, B.O. (1989). No linkage between genes
controlling female pheromone production and male pheromone response in the European corn
borer, Ostrinia nubilalis Hubner (Lepidoptera; Pyralidae). Genetics *123*, 553-556.
Lu, B., LaMora, A., Sun, Y., Welsh, M.J., and Ben-Shahar, Y. (2012). ppk23-Dependent
chemosensory functions contribute to courtship behavior in Drosophila melanogaster. PLoS
Genet *8*, e1002587.
Lu, B., Zelle, K.M., Seltzer, R., Hefetz, A., and Ben-Shahar, Y. (2014). Feminization of pheromone-
sensing neurons affects mating decisions in Drosophila males. Biology Open *In Press*, 152-160.
Makki, R., Cinnamon, E., and Gould, A.P. (2014). The development and functions of oenocytes.
Annual review of entomology *59*, 405-425.
Miller, M.A., Pfeiffer, W., and Schwartz, T. (2010). Creating the CIPRES Science Gateway for
inference of large phylogenetic trees. In 2010 Gateway Computing Environments Workshop
(GCE), pp. 1-8.
Miyamoto, T., and Amrein, H. (2008). Suppression of male courtship by a Drosophila
pheromone receptor. Nat Neurosci *11*, 874-876.
Moon, S.J., Lee, Y., Jiao, Y., and Montell, C. (2009). A Drosophila gustatory receptor essential for
aversive taste and inhibiting male-to-male courtship. Curr Biol *19*, 1623-1627.
Ng, S.H., Shankar, S., Shikichi, Y., Akasaka, K., Mori, K., and Yew, J.Y. (2014). Pheromone
evolution and sexual behavior in Drosophila are shaped by male sensory exploitation of other
males. Proc Natl Acad Sci U S A *111*, 3056-3061.
Park, J.H., and Kwon, J.Y. (2011). A systematic analysis of Drosophila gustatory receptor gene
expression in abdominal neurons which project to the central nervous system. Mol Cells *32*,
375-381.
Poudel, S., Kim, Y., Kim, Y.T., and Lee, Y. (2015). Gustatory receptors required for sensing
umbelliferone in Drosophila melanogaster. Insect Biochem Mol Biol *66*, 110-118.
Robertson, H.M., Warr, C.G., and Carlson, J.R. (2003). Molecular evolution of the insect
chemoreceptor gene superfamily in Drosophila melanogaster. PNAS *100*, 14537-14542.
Scott, K., Brady, R., Jr., Cravchik, A., Morozov, P., Rzhetsky, A., Zuker, C., and Axel, R. (2001). A
chemosensory gene family encoding candidate gustatory and olfactory receptors in Drosophila.
Cell *104*, 661-673.
Shankar, S., Chua, J.Y., Tan, K.J., Calvert, M.E., Weng, R., Ng, W.C., Mori, K., and Yew, J.Y. (2015).
The neuropeptide tachykinin is essential for pheromone detection in a gustatory neural circuit.
eLife *4*.
Shaw, K.L., Ellison, C.K., Oh, K.P., and Wiley, C. (2011). Pleiotropy, "sexy" traits, and speciation.
Behavioral Ecology *22*, 1154-1155.
Shaw, K.L., and Lesnick, S.C. (2009). Genomic linkage of male song and female acoustic
preference QTL underlying a rapid species radiation. Proceedings of the National Academy of
Sciences *106*, 9737-9742.
Shim, J., Lee, Y., Jeong, Y.T., Kim, Y., Lee, M.G., Montell, C., and Moon, S.J. (2015). The full
repertoire of Drosophila gustatory receptors for detecting an aversive compound. Nat Commun
6, 8867.
Shirangi, T.R., Dufour, H.D., Williams, T.M., and Carroll, S.B. (2009). Rapid evolution of sex
pheromone-producing enzyme expression in Drosophila. Plos Biol *7*, e1000168.

Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H.,
Remmert, M., Söding, J., et al. (2011). Fast, scalable generation of high☐quality protein multiple
sequence alignments using Clustal Omega. Molecular Systems Biology *7*.
Singh, N.D., and Shaw, K.L. (2012). On the scent of pleiotropy. Proc Natl Acad Sci U S A *109*, 5-6.
Stagkourakis, S., Kim, H., Lyons, D.J., and Broberger, C. (2016). Dopamine Autoreceptor
Regulation of a Hypothalamic Dopaminergic Network. Cell reports.
Steiger, S., Schmitt, T., and Schaefer, H.M. (2010). The origin and dynamic evolution of chemical
information transfer. Proceedings of the Royal Society B: Biological Sciences.
Svensson, E.I., Karlsson, K., Friberg, M., and Eroukhmanoff, F. (2007). Gender differences in
species recognition and the evolution of asymmetric sexual isolation. Curr Biol *17*, 1943-1947.
Sweigart, A.L. (2010). The genetics of postmating, prezygotic reproductive isolation between
Drosophila virilis and D. americana. Genetics *184*, 401-410.
Symonds, M.R., and Elgar, M.A. (2008). The evolution of pheromone diversity. Trends Ecol Evol
23, 220-228.
Thistle, R., Cameron, P., Ghorayshi, A., Dennison, L., and Scott, K. (2012). Contact
Chemoreceptors Mediate Male-Male Repulsion and Male-Female Attraction during Drosophila
Courtship. Cell *149*, 1140-1151.
Toda, H., Zhao, X., and Dickson, B.J. (2012). The Drosophila Female Aphrodisiac Pheromone
Activates ppk23+ Sensory Neurons to Elicit Male Courtship Behavior. Cell reports
doi:10.1016/j.celrep.2012.05.007, 599-607.
van der Goes van Naters, W., and Carlson, J.R. (2007). Receptors and neurons for fly odors in
Drosophila. Curr Biol *17*, 606-612.
Wang, Z., Singhvi, A., Kong, P., and Scott, K. (2004). Taste representations in the Drosophila
brain. Cell *117*, 981-991.
Watanabe, K., Toba, G., Koganezawa, M., and Yamamoto, D. (2011). *Gr39a*, a highly diversified
gustatory receptor in *Drosophila*, has a role in sexual behavior. Behav Genet *41*, 746-753.
Wiley, C., Ellison, C.K., and Shaw, K.L. (2011). Widespread genetic linkage of mating signals and
preferences in the Hawaiian cricket Laupala. Proceedings of the Royal Society B:
Biological Sciences.
Wyatt, T.D., and Cambridge University Press (2014). Pheromones and Animal Behavior :
Chemical Signals and Signatures. (Cambridge: Cambridge University Press,), p. 1 online
resource (426 pages).
Yew, J.Y., and Chung, H. (2015). Insect pheromones: An overview of function, form, and
discovery. Prog Lipid Res *59*, 88-105.
Yew, J.Y., Dreisewerd, K., Luftmann, H., Muthing, J., Pohlentz, G., and Kravitz, E.A. (2009). A new
male sex pheromone and novel cuticular cues for chemical communication in Drosophila. Curr
Biol *19*, 1245-1254.
Zheng, X., Valakh, V., DiAntonio, A., and Ben-Shahar, Y. (2014). Natural antisense transcripts
regulate the neuronal stress response and excitability. eLife *3*, e01849.
432

433   Figures and figure legends:

Figure 1. ***Gr8a* is a sexually dimorphic chemosensory receptor enriched in male**

oenocytes. (A) *Gr8a* promoter activity in proboscis, (B) forelegs, and (C) abdomens of

males (top panels) and females (bottom panels). (D) *Gr8a* mRNA expression. Relative

mRNA levels were measured by real-time quantitative RT-PCR. **, p<0.01 Mann

Whitney Rank Sum Test. (E) Confocal z-stack image of *Gr8a*>EGFP in abdominal cells.

(F) Confocal z-stack image of *desat1*>Luciferase in abdominal cells. (g) Co-expression

of *Gr8a* and *desat1*. Green, *Gr8a*; Red, *desat1*; Blue, nuclear DAPI stain. Orange

arrowhead, fat body cells; white arrowhead, oenocytes. Scale bar = 100μm.

Figure 2. *Gr8a* activity contributes to the perception and production of inhibitory

signal associated with mating decisions in males and females. (A) Blocking neural

activity in female *Gr8a*-expressing sensory neurons shortens copulation latency.

Homozygous (B) or hemizygous (C) *Gr8a* null females show shortened copulation

latency relative to wild-type controls. Df(1)BSC663 is a deficiency that covers the *Gr8a*

locus. Df(1)BSC754 was used as a control. (D) Expression of *Gr8a* cDNA with *Gr8a*

promoter rescues the copulation latency phenotype in *Gr8a* mutant females. (E) Wild-

type females exhibit shorter copulation latency when courted by *Gr8a* mutant relative to

wild-type males. (F) *Gr8a* mutant males do not recognize mating status of females, and

have a reduced transfer of inhibitory mating pheromones during copulations. Female,

female genotype; Sperm donor, genotype of males mated first with focal females; Focal

male, genotypes of experimental males presented with mated females. Different letters

above bars indicate statistically significant *post hoc* contrasts between groups (panels

C, D, and F; p<0.05 ANOVA, n>15/group). *, p<0.05, Mann Whitney Rank Sum Test,

n>15/group)

Figure 3. The *Gr8a* mutation affects the pheromone profiles of males and mated females. (A) Nonmetric multidimensional scaling (NMDS) plot of CHC profiles of wild-type and *Gr8a* mutant males. p<0.001, permutation MANOVA. (B-C) The effect of the *Gr8a* mutation on levels of individual CHCs in males. Only affected CHCs shown. See Table S2 for the complete list. *, p<0.05, **, p<0.001, Student's t-test or Mann Whitney Rank Sum Test. (D) The effect of Gr8a mutation on the expression level of several desaturase genes. Only affected genes shown. See Table S3 for the complete list. *, p<0.05, Student's t-test. (E) NMDS plot of CHC profiles of females mated with wild-type or *Gr8a* mutant males. p=0.570, permutation MANOVA. (F) Nonacosane (C$_{29}$) differs between females mated with wild-type and *Gr8a* mutant males. See Table S4 for complete list of mated-female CHCs. *, p<0.05, Student's t-test.

Figure 4. Model for the pleiotropic action of *Gr8a* in the perception and production of pheromones. (A) *Gr8a* functions as a chemoreceptor for an inhibitory signal in pheromone-sensing GRNs of males and females. (B) *Gr8a* also functions as a CHC autoreceptor in oenocytes, which regulates CHC secretion [1] or CHC synthesis [2] via signaling feedback loops [3].

Methods

Animals. Flies were maintained on a standard cornmeal medium under a 12:12 light-dark cycle at 25 Celsius. Unless specifically stated, the *D. melanogaster Canton-S (CS)* strain served as wild-type control animals. UAS-mCD8::GFP, UAS-myr::GFP, UAS-TNT-E, UAS-TNT-IMP-V1-A, *Gr8a-GAL4* and *Gr8a^1* fly lines were from the Bloomington Stock center. Originally in the *w^{1118}* background, the *Gr8a^1 null* allele was outcrossed for six generations into the *CS* wild-type background. *PromE(800)-GAL4* and *PromE(800)>Luciferase* were from Joel Levine at the University of Toronto. The following *Drosophila* species were obtained from the San Diego Stock Center: *D. simulans* 14011-0251.192, *D. sechellia* 14021-0248.03, *D. yakuba* 14021-0261.01, *D. erecta* 14021-0224.00, *D. ananassae* 14024-0371.16, *D. pseudoobscura* 14011-0121.104, D. *persimilis* 14011-0111.50, *D. willistoni* 14030-0811.35, *D. mojavensis* 15081-1352.23, and *D. virilis* 15010-1051.118. The UAS-*Gr8a* transgenic line was generated by cloning the *Gr8a* cDNA in the pUAST-attB vector using 5' EcoRI and 3' NotI restriction sites, followed by *ΦC31* integrase-dependent transgene integration at a Chromosome 2 *att*P landing site (2L:1476459) as previously described (Zheng et al., 2014). For genetic rescue experiments, the UAS-*Gr8a* and *Gr8a*-GAL4 lines were transgressed into the *Gr8a^1* CS genetic background.

Immunohistochemistry. To visualize the expression pattern of *Gr8a* in males and females, *Gr8a-GAL4* flies (Lee et al., 2012) were crossed to UAS-CD8::EGFP and live-imaged at 5 days old using a Nikon-A1 confocal microscope. To demonstrate *Gr8a* expression in oenocytes, abdomens from *Gr8a*-GAL4*/*UAS-*myr::GFP ; PromE(800)>Luciferase* flies were dissected and immunostained as previously described (Lu et al., 2012; Zheng et al., 2014) by using a Rabbit anti-GFP (1:1000; A-11122, Thermo Fisher Scientific) and a mouse anti-luciferase (1:100; 35-6700, Thermo Fisher Scientific) antibodies followed by AlexaFluor 488 anti-rabbit and AlexaFluor 568 anti-mouse secondary antibodies (Both at 1:1000; Thermo Fisher Scientific).

mRNA expression. Newly eclosed flies were separated by sex under CO_2 and aged for
5 days on standard cornmeal medium. On day 6, flies were placed in a -80°C freezer
until RNA extraction. To separate body parts, frozen flies were placed in 1.5ml
microcentrifuge tubes, dipped in liquid nitrogen, and then vortexed repeatedly until
heads, appendages, and bodies were clearly seperated. Total RNA was extracted using
the Trizol Reagent (Thermo Fisher Scientific) separately from heads, bodies, and
appendages for *Gr8a* expression and from bodies for desaturase enzyme genes.
cDNAs were synthesized using SuperScript II reverse transcriptase (Thermo Fisher
Scientific) with 500 ng total RNA in a 20 uL reaction. Real-time quantitative RT-PCR
was carried out as previously described with *Rp49* as the loading control gene (Lu et al.,
2012; Lu et al., 2014; Zheng et al., 2014). Primer sequences are described in Tables
S5, S6 and S7.

Courtship Behavior. Single-pair assays were performed as we have previously
published (Lu et al., 2012; Lu et al., 2014). In short, newly eclosed males were kept
individually on standard fly food in plastic vials (12 x 75mm). Newly eclosed virgin
females were kept in groups of 10 flies. All behaviors were done with 4-7 day-old
animals, which were housed under constant conditions of 25° C and a 12h:12h light-
dark cycle. Courtship was video recorded for 10 min for male courtship and 15 min for
female mating receptivity. Male courtship latency and index were measured as
previously described (Lu et al., 2012; Lu et al., 2014). Female receptivity index was
defined as the time from the initiation of male courtship until copulation was observed.
Unless otherwise indicated, assays were performed under red light conditions.

Fitness measures. Three-day old virgin male-female pairs were housed together for 24
hours after which males were discarded. Subsequently, groups of five females were
allowed to lay eggs daily on fresh agar grape plates for five consecutive days. Fitness
was measured as total number of eggs laid.

Phylogenetic analysis. Protein sequences of GR8A orthologs from the 12 sequenced *Drosophila* reference genomes were aligned by using the ClustalW algorithm in the Omega package (Sievers et al., 2011), followed by ProtTest (v2.4) to determine the best model of protein evolution (Abascal et al., 2005). Subsequently, Akaike and Bayesian information criterion scores were used to select the appropriate substitution matrix. We then used a maximum likelihood approach and rapid bootstrapping within RAxML v 7.2.8 Black Box on the Cipres web portal to make a phylogenetic tree (Miller et al., 2010). Visualizations of the bipartition files were made using FigTree v1.3.1 (http://tree.bio.ed.ac.uk/software/figtree/).

Pheromone Analysis. Virgin flies were collected upon eclosion under a light CO_2 anesthesia and kept in single-sex vials in groups of 10 with 6 biological replications for each genotype and sex. Virgin flies were aged for 5 days on standard cornmeal medium at 25°C. To collect mated flies, both females and males were aged for 3 days before single mating pairs were placed in a standard fly vial with standard cornmeal food for 24 hours. The pair was then separated for 24 hours before collection. Copulation was confirmed by the presence of larvae in the vials of mated females several days later. On the morning of day 5, flies were anesthetized under light CO_2 and groups of five flies were placed in individual scintillation vials (VWR 74504-20). To extract CHCs, each group of flies was covered by 100 uL hexane (Sigma-Aldrich #139386-500ML) containing 50μg/mL hexacosane (Sigma-Aldrich #241687-5G) and was washed for ten minutes. Subsequently, hexane washes were transferred into a new 2 ml glass vial containing a 350 uL insert (Thermo Scientific C4000-LV-1W) and were stored at -20°C until shipment to the Millar laboratory.

Analyses of CHC profiles were done by gas chromatography and mass spectroscopy (GC-MS) in the Millar laboratory at UC Riverside as previously described (Chung et al., 2014). Peak areas were measured, and data was normalized to known quantity of internal standard hexacosane (Sigma-Aldrich #241687-5G). The relative proportion of each compound in each sample was calculated and used in further statistical analysis. Qualitative data were analyzed through a permutation MANOVA using the ADONIS

function in the vegan package of R with Bray-Curtis dissimilarity measures (Oksanen et

al., 2017). Data were visualized using non-metric multidimensional scaling (vegdist

function in the vegan package of R followed by cmdscale function in the stats package

(Oksanen et al., 2017)) using Bray-Curtis dissimilarity, and either 2 or 3 dimensions in

order to minimize stress to < 0.1. Quantitative data were analyzed by using a t-test or

Mann-Whitney Rank Sum Test in R 3.3.2 (R Core Team, 2016).

Data availability. All relevant data are available from the corresponding author upon

request.

Supplemental Table Legends

Table S1. Candidate *Gr* genes in male and female abdomen. Plus sign indicates
presence of PCR product and minus signs indicates PCR product not detected. Genes
with no PCR product detected in male or female abdomens not shown.

Table S2. Male CHCs. Retention time (R.T.), compound, and percent total (% total) of
each compound as part of the total pheromonal bouquet for wild-type (wt) and *Gr8a*
mutant males.

Table S2. Male CHCs. Retention time (R.T.), compound, and percent total (% total) of
each compound as part of the total pheromonal bouquet for wild-type (wt) and *Gr8a*
mutant males.

Table S3. Desaturase gene expression. Relative mRNA expression of each desaturase
gene for wild-type (wt) and *Gr8a* mutant males.

Table S4. Mated-female CHCs. Retention time (R.T.), compound, and percent total (%
total) of each compound as part of the total pheromonal bouquet for females mated with
wild-type (wt) or *Gr8a* mutant males.

Table S5. Nucleotide sequences for qRT-PCR primers for *D. melanogaster Gr* genes.

Table S6. Nucleotide sequences for qRT-PCR primers for *D. melanogaster Gr8a*, *Rp49*
and orthologs.

Table S7. Nucleotide sequences for qRT-PCR primers for *D. melanogaster* desaturase
enzyme genes