Chemoreceptor pleiotropy facilitates the functional coupling of mating pheromone
synthesis and perception

Kathleen M. Zelle[1,†], Cassondra Vernier[1,†], Nicole Leitner[1], Xitong Liang[2], Sean Halloran[3],
Jocelyn G. Millar[3], Yehuda Ben-Shahar[1,4,*]

[1]Department of Biology, Washington University in Saint Louis, 1 Brookings Drive, Saint
Louis, MO 63130 USA.

[2]Department of Neuroscience, Washington University School of Medicine, 660 South
Euclid Avenue, St. Louis, MO 63110, USA

[3]Department of Entomology, University of California, Riverside, 900 University Avenue,
Riverside, CA 92521, USA

[4]Lead Contact

[*]Correspondence: benshahary@wustl.edu (Y.B.)

[†]Denotes equal contribution

ABSTRACT

Optimal mating decisions depend on stable, coupled signaling systems because any independent changes in either the signal or its perception could carry a fitness cost. However, since the perception and production of specific mating signals are often mediated by different tissues and cell types, the genetic and cellular mechanisms that drive their coupling on the evolutionary and physiological timescales remain a mystery for most animal species. Here we show that in *Drosophila melanogaster,* sensory perception and synthesis of an inhibitory mating pheromone is regulated by the action of *Gr8a*, a member of the *Gustatory receptor* gene family. In particular, *Gr8a* acts as a neurosensory receptor for pheromones in males and females and independently regulates their synthesis in the male fat body and pheromone-producing oenocytes. These data provide a relatively simple molecular explanation for how genetic coupling allows for the robust and stable flow of social information at the population level.

Keywords: *Drosophila melanogaster;* Vinegar fly; Cuticular hydrocarbons; *Gr8a*; Oenocyte.

INTRODUCTION

The majority of sexually-reproducing animals use intricate mating signaling systems, which rely on the robust functional coupling between the production and perception of signals since any independent changes in either the signal or the capacity to sense it would carry a fitness cost (Boake, 1991; Brooks et al., 2005; Hoy et al., 1977; Shaw et al., 2011; Shaw and Lesnick, 2009; Steiger et al., 2011; Sweigart, 2010; Symonds and Elgar, 2008; Wyatt, 2014). Previously published theoretical models have predicted that one possible solution for maintaining the functional coupling of signal production and perception is through genetic coupling, in which physically linked or pleiotropic genes control both mating signal production and perception (Boake, 1991; Butlin and Ritchie, 1989; Shaw et al., 2011; Shaw and Lesnick, 2009). Genetic coupling promotes a genetic correlation between signal and receptor (Chebib and Guillaume, 2019; Kirkpatrick and Hall, 2004; Lande, 1984, 1981, 1980), which provides a robust molecular mechanism for maintaining stable and reliable population-level communication systems while still retaining the capacity for future signal diversification, as necessitated for speciation (Hoy et al., 1977; Shaw et al., 2011; Shaw and Lesnick, 2009; Wiley et al., 2012). While previous models suggest these two genetic architectures should function similarly in maintaining signal-receptor genetic correlations (Lande, 1984), a recent model shows that pleiotropy, in which a single gene controls more than one phenotypic trait, maintains a higher signal-receptor genetic correlation than gene linkage in the face of evolutionary processes (Chebib and Guillaume, 2019). This suggests that pleiotropy may play a particularly robust and important role in maintaining stable signal-receptor coupling in animal mating systems.

Several experimental studies have provided empirical evidence that pleiotropy maintains functional coupling between mating-specific signals and the behaviors they elicit at the population level (Hoy et al., 1977; Shaw et al., 2011; Shaw and Lesnick, 2009; Wiley et al., 2012). However, the complex characteristics of mating signals and animal behaviors present a major barrier for identifying the actual pleiotropic genes and molecular pathways that provide genetic coupling between the production and perception of mating signals (Chenoweth and Blows, 2006; Singh and Shaw, 2012). Furthermore, how the perception and production of mating signals remains stably coupled is particularly

puzzling for mating pheromones, since the sensing of specific chemicals is mediated by
the action of the chemosensory peripheral nervous system, while their production is often
restricted to specialized, non-neuronal pheromone producing cells (Chung and Carroll,
2015; McKinney et al., 2015; Mucignat-Caretta, 2014). Therefore, whether pleiotropy
could function in both the production and perception of mating pheromones, a ubiquitous
means of signaling across animal mating systems (Mucignat-Caretta, 2014), remained
unknown.

Here we show that, in the vinegar fly *Drosophila melanogaster*, some pheromone-driven
mating decisions are coupled to the production of specific pheromones via pleiotropic
chemoreceptors. Specifically, we demonstrate that *Gr8a*, a member of the gustatory
receptor family, independently contributes to the perception of inhibitory mating signals in
pheromone-sensing neurons, as well as the production of inhibitory mating pheromones
in pheromone producing cells called oenocytes. Together, these data provide a relatively
simple molecular explanation for how the perception and production of specific mating
pheromones are functionally and genetically coupled.

RESULTS

Similar to other insect species, in *Drosophila,* cuticular hydrocarbons (CHCs), or long-
chain fatty acids synthesized by the fat body and oenocytes (Billeter et al., 2009; Gutierrez
et al., 2007; Krupp et al., 2013; Makki et al., 2014), provide a hydrophobic desiccation
barrier for the insect body and play an important role as pheromones in regulating diverse
behaviors, including those used in mating (Blomquist and Bagnères, 2010; Chung and
Carroll, 2015; Ferveur, 2005; Howard and Blomquist, 2005; McKinney et al., 2015).
Specifically, complex blends of CHCs are often utilized by insect species to communicate
sex identity and female mating status, as well as to define the behavioral reproductive
boundaries between closely related species (Billeter et al., 2009; Chung et al., 2014;
Chung and Carroll, 2015; Coyne et al., 1994; Dweck et al., 2015; Ng et al., 2014; Shirangi
et al., 2009; Yew and Chung, 2015). In *Drosophila*, the perception of volatile CHCs is
mediated by olfactory sensory neurons located in the antennae and maxillary palps
(Benton et al., 2007; Kurtovic et al., 2007; Lebreton et al., 2014; van der Goes van Naters

and Carlson, 2007), while less volatile CHCs, whose perception requires physical contact,

are sensed by specialized gustatory-like receptor neurons (GRNs) in the appendages

(legs and wings) and the proboscis (Koh et al., 2015; Lu et al., 2014, 2012; Thistle et al.,

2012; Toda et al., 2012). While some of the genes and pathways that contribute to CHC

synthesis in *Drosophila* are known, the molecular identities of most CHC receptors remain

unknown. Previous work suggested that the gene *Desat1*, which encodes a fatty acid

desaturase enriched in oenocytes, might also independently contribute to CHC

perception (Bousquet et al., 2011). However, due to the expression of *Desat1* in central

neurons (Billeter et al., 2009) and the broad impact its mutant alleles have on the CHC

profiles of both males and females (Labeur et al., 2002), whether *Desat1* directly

contributes to mating signal perception remains unresolved.

Consequently, we chose to examine members of the *Gustatory receptor* (*Gr*) gene family

as candidates for pleiotropic factors that might contribute directly to both the perception

and production of pheromonal mating signals in *Drosophila*. Because several family

members have already been implicated in the detection of specific excitatory and

inhibitory pheromones (Bray and Amrein, 2003; Hu et al., 2015; Miyamoto and Amrein,

2008; Moon et al., 2009; Watanabe et al., 2011), and the majority of genes that encode

family members are already known to be enriched in GRNs (Clyne et al., 2000; Dunipace

et al., 2001; Scott et al., 2001; Wang et al., 2004), we reasoned that any pleiotropic *Gr*

genes should also be expressed in abdominal oenocytes (Billeter et al., 2009).

Subsequently, an RT-PCR screen revealed that 24 out of the 60 genes that encode *Gr*

family members in the *Drosophila* genome (Clyne et al., 2000; Dunipace et al., 2001;

Robertson et al., 2003; Scott et al., 2001) exhibit enriched expression in abdominal

tissues (Table 1).

Although several members of the *Gr* receptor family, including *Gr68a, Gr32a, Gr66a,*

Gr39a, and *Gr33a*, were previously linked to the sensory perception of mating

pheromones (Bray and Amrein, 2003; Lacaille et al., 2009; Miyamoto and Amrein, 2008;

Moon et al., 2009; Watanabe et al., 2011), none of these candidate pheromone receptors

were identified in our initial screen for *Gr* genes with enriched mRNA expression in

abdominal tissues of either males or females (Table 1). However, *Gr8a*, which we

identified as having male-specific expression in the abdomen, is localized to a

chromosomal region with enriched expression in abdomens that was previously

implicated in chemoreception (Table 1) and was shown to contribute to the sensory

detection of the non-proteinogenic amino acid L-Canavanine (Lee et al., 2012; Shim et

al., 2015). Because our initial expression screen was based on whole-abdomen RNAs,

and it was previously reported that *Gr8a* is also expressed in some abdominal neurons

(Park and Kwon, 2011), we next used a GAL4 transgenic driver to examine the spatial

expression pattern of *Gr8a* in male and female flies. As was previously reported, we found

that *Gr8a* is expressed in 14-16 GRNs in the proboscis (Figure 1A) (Lee et al., 2012) as

well as in two paired GRNs in the foreleg pretarsus (Figure 1B) of male and female flies.

In addition, the *Gr8a*-GAL4 reporter was also highly expressed in abdominal oenocyte-

like cells in males but not females (Figure 1C). The male-specific abdominal expression

pattern was further supported by qRT-PCR (Figure 1D). Using immunohistochemistry, we

then demonstrated that *Gr8a* is co-expressed with the oenocyte marker *Desat1* (Billeter

et al., 2009), as well as in *Desat1*-negative cells with fat-body-like morphology (Figure

1E-G). Finally, by using immunohistochemistry on a CRISPR/Cas9 generated GFP

tagged allele of the *Gr8a* locus, we demonstrated that the GR8A protein localizes to the

fat body (Figure 1H).

Together, the enriched expression of *Gr8a* in foreleg GRNs, a primary pheromone

chemosensory organ (Lu et al., 2014, 2012), and its sexually dimorphic expression in

oenocytes and fat body suggested that in addition to its role in L-Canavanine perception

(Lee et al., 2012; Shim et al., 2015), *Gr8a* contributes to mating behaviors. Since mating

choices of both male and female flies are determined by a complex blend of excitatory

and inhibitory pheromones (Billeter et al., 2009; Krupp et al., 2013), we anticipated that if

Gr8a is indeed a pleiotropic factor that independently contributes to the perception and

production of specific components of the male mating pheromone bouquet, then

disruptions of the signal production in males or its perception in females should carry

similar impacts on female mate choice behavior. To directly test this hypothesis, we first

investigated whether *Gr8a* and the GRNs that express it are required for sensory

functions associated with female mate choice by using a single-pair courtship paradigm

(Lu et al., 2014, 2012). We found that blocking neuronal transmission in *Gr8a*-expressing

GRNs with the transgenic expression of the tetanus toxin (TNT) in females resulted in a

shorter copulation latency relative to wild-type females when courted by wild-type males

(Figure 2A). Similarly, homozygous (Figure 2B) and hemizygous (Figure 2C) *Gr8a* mutant

females exhibited shorter copulation latencies relative to wild-type controls, which could

be rescued by the transgenic expression of a *Gr8a* cDNA (Figure 2D). In contrast, similar

genetic manipulations of the *Gr8a* gene or *Gr8a*-expressing GRNs in males did not affect

courtship behavior as measured by courtship latency and index towards wild-type females

(Figure 2- figure supplement 1). These data indicate that *Gr8a* is required for female mate

choice via the detection of male-borne inhibitory mating signals.

Because *Gr8a* expression is specifically enriched in male oenocytes, we next tested the

hypothesis that *Gr8a* also plays a role in the production and/or release of inhibitory mating

signals by males. Similar to the effect of the *Gr8a* mutation in females, we found that wild-

type virgin females exhibited shorter copulation latencies towards *Gr8a* mutant males

relative to wild-type controls (Figure 2E). These data indicate that *Gr8a* mutant males

produce and/ or release abnormally low levels of inhibitory mating pheromones.

Mating decisions in *D. melanogaster* rely on a balance of excitatory and inhibitory drives

(Kallman et al., 2015). Therefore, in this case, male-borne inhibitory signals may promote

female mate choice. Additionally, previous studies showed that, in order to increase their

fitness, *Drosophila* males transfer inhibitory mating pheromones to females during

copulation, which subsequently lowers the overall attractiveness of mated females to

other males and the probability of females remating (Benton et al., 2007; Billeter et al.,

2009; Datta et al., 2008; Yew et al., 2009). Because our data indicate that *Gr8a* plays a

role in the production of an inhibitory signal in males, we reasoned that the same inhibitory

signal may be transferred from males to females during copulation. Therefore, we next

tested the hypothesis that *Gr8a* mutant males would have a reduced ability to produce

and/or transfer inhibitory pheromones to females during copulation. Accordingly, we

found that wild-type males failed to recognize the mating status of wild-type females that

were previously mated with *Gr8a* mutant males (Figure 2F). These data indicate that *Gr8a*

action is required for the production and/or transfer of inhibitory pheromones from males

to females during copulation. Because *Gr8a* is involved in the perception of an inhibitory

mating signal, we also tested the hypothesis that *Gr8a* mutant males will exhibit an

abnormal capacity for identifying the post-mating status of wild-type females. Indeed, we

observed that *Gr8a* mutant males were more likely than wild type males to court a mated
female (Figure 2F), suggesting that in addition to its role in the production of inhibitory
mating signals, *Gr8a* is also required in males for the sensory recognition of mating
inhibitory signals that mark the post-mating status of females. Together, these studies
indicate that *Gr8a* is required in males for the production and perception of transferrable
inhibitory mating signals that advertise post-mating status in females. These data further
support the pleiotropic role of the *Gr8a* gene in the production of inhibitory mating signals
in males and the perception of these signals in both males and females in sex-specific
contexts.

Because our behavioral data indicate that the *Gr8a* mutation has a dramatic effect on the
production and perception of a putative inhibitory mating pheromonal signal, we next
examined whether the *Gr8a* mutation has a direct effect on qualitative and quantitative
aspects of male and mated-female CHC profiles. As predicted by our behavioral data,
analyses of male epicuticular pheromones by gas chromatography and mass
spectroscopy revealed that the *Gr8a* mutation has a significant impact on the overall
qualitative characteristics of the CHC profile of males (Figure 3A). Additional analyses
revealed significant quantitative effects of the *Gr8a* mutation on the cuticular levels of
several specific CHCs in males (Figure 3B-3C and Table 2). In particular, the *Gr8a*
mutation affects the levels of several alkenes and methyl-branched alkanes, which have
been implicated in mate choice behaviors in diverse *Drosophila* species (Billeter et al.,
2009; Chung et al., 2014; Dyer et al., 2014; Shirangi et al., 2009). Similarly, *Gr8a* mutation
affects the expression level of two desaturase enzymes in males (Figure 3D), which are
involved in the biosynthesis of alkenes (Chung and Carroll, 2015). Additionally, although
the CHC profile of females mated with *Gr8a* mutant males did not qualitatively differ from
those mated with wild-type males (Figure 3E), one specific, lowly-expressed component
of the CHC profile, nonacosane (C_{29}), differed quantitatively between the two mated-
female groups (Figure 3F). In addition to these identified CHCs, *Gr8a* may have a direct
effect on other pheromones that are unable to be measured. Together, our behavioral
and pheromonal data indicate that *Gr8a* action contributes to mating decisions in females
by co-regulating the perception of an inhibitory mating pheromone by females, as well as
its production by males, which is consistent with a pleiotropic function for *Gr8a*.

Next, we sought to identify the *Gr8a*-associated inhibitory component of the male
Drosophila CHC profile by testing the effect of the specific CHCs impacted by the *Gr8a*
mutation on courtship behaviors. By measuring female copulation latency in a single-pair
courtship paradigm (Lu et al., 2014, 2012), we found that wild-type females did not
copulate with *Gr8a* mutant males that were first perfumed with an exaggerated amount
of the alkenes 9-C25, 7-C25, and 7-C27 (Figure 4). Similarly, we found that wild-type
males exhibited a lower courtship index and longer courtship latency when females were
perfumed with an exaggerated amount of 9-C_{25} (Figure 5A-C), and a longer copulation
latency when females were perfumed with an exaggerated amount of 7-C_{25} (Figure 5D-
5F), but no differences when females were perfumed with an exaggerated amount of 7-
C_{27} (Figure 5G-I). These data suggest that the alkenes 7-C_{25}, 9-C_{25}, and 7-C_{27} may
contribute to an inhibitory signal associated with *Gr8a*.

DISCUSSION

The data presented here demonstrate that a pleiotropic chemoreceptor co-regulates the
perception and production of a specific pheromonal signal that plays an important role in
mate choice behaviors of both sexes. In contrast to its expected function in the perception
of chemical ligands, how *Gr8a*, a member of a canonical chemoreceptor family might also
contribute to the production of pheromonal signals is not obvious. In some of the better
understood scretory cell types, autoreceptors are essential for the regulation of synthesis
and secretion rates. For example, dopaminergic and serotonergic cells regulate rates of
synthesis and release of their respective neuromodulators. This is accomplished by the
action of autoreceptors, which are used as sentinels that regulate synthesis rates via
specific molecular signaling feedbacks in response to level changes in the extracellular
concentrations of the secreted molecule (Ford, 2014; Stagkourakis et al., 2016).
Therefore, we hypothesize that one possible way *Gr8a* might regulate the synthesis
and/or secretion of specific CHCs is by acting as an oenocyte-intrinsic autoreceptor,
which integrates feedback information to the complex genetic network that regulates the
epicuticular CHC profile of males (Figure 6).

Previous work indicated that *Gr8a* is expressed in "bitter" taste neurons in the proboscis
and are specifically required for the sensory perception of the feeding deterrent L-
canavanine (Lee et al., 2012; Shim et al., 2015), but not for the detection of other inhibitory
chemicals such as caffeine, strychnine, and umbelliferone (Lee et al., 2009; Poudel et al.,
2015). Our data indicate that *Gr8a* contributes to inhibitory functions associated with
mating decisions as well, as was previously reported for other 'bitter' receptors in
Drosophila (Lacaille et al., 2007; Moon et al., 2009). Together this indicates similar
function in 'bitter' taste perception and the perception of inhibitory signals in this species.

Gr8a-dependent perception of L-canavanine seems to depend on its heterotrimeric
interaction with *Gr66a* and *Gr98b* in bitter sensing neurons in the proboscis (Shim et al.,
2015). Although both of these *Gr* genes were also identified in our initial screen for
receptors enriched in the abdomen, *Gr66a* was expressed in both males and females
while *Gr98b* was specifically enriched in females (Table 1). These data suggest that *Gr8a*-
dependent contributions to feeding and mating decisions are independently mediated via
interactions with different *Gr* genes in each sensory context, and further point to plastic
subunit composition as an essential aspect of the mechanism for ligand specificity in
insect gustatory receptors. Although we do not yet know the specific chemical identity of
the ligand of *Gr8a*, previous studies indicated that at least two inhibitory mating
pheromones, cVA and CH503, are transferred from males to females during copulations.
Although our data suggest that the *Gr8a* mutation affects the level of cVA expressed by
males, it is unlikely that either cVA or CH503 are the putative *Gr8a* ligands because the
volatile cVA acts primarily via the olfactory receptor *Or67d* (Benton et al., 2007; Datta et
al., 2008; Kurtovic et al., 2007), and CH503 has been reported to signal via *Gr68a*-
expressing neurons, which are anatomically distinct from the *Gr8a* GRNs we describe
here (Figure 1A-B) (Shankar et al., 2015; Yew et al., 2009). Our data indicate that the
alkenes 5-C_{25}, 7-C_{25}, and 7-C_{27} differ in abundance between the CHC profiles of wild-type
and *Gr8a* mutant males and likely function as inhibitory mating signals. If Gr8a is acting
as an autoreceptor, then it is possible that these alkenes are ligands of *Gr8a*. However,
future studies are needed to assess the specific relationship of these alkenes with *Gr8a*.

As populations diversify, sexual communication systems within a species should be
evolutionarily constrained, since any mismatch between signal and its perception would

result in a decrease in fitness (Boake, 1991; Symonds and Elgar, 2008). However, sexual

isolation between closely related species often relies on distinct sexual signaling systems,

suggesting that these systems must undergo frequent and quick divergence (Shaw and

Lesnick, 2009). Therefore, how mating signals and their receptors are linked at the

molecular and physiological levels but retain the ability for future diversification remains

unknown for most studied mating communication systems. Several theoretical models

have previously stipulated that one possible solution to this conundrum is a genetic

coupling between signal production and associated mate choice behaviors, which

supports the co-evolution of mating signals with their cognate receptors in the face of

speciation (Arnold et al., 2006; Boake, 1991; Butlin and Ritchie, 1989; Lande, 1981;

Lofstedt et al., 1989; Shaw et al., 2011; Shaw and Lesnick, 2009; Singh and Shaw, 2012;

Svensson et al., 2007). The studies we present here do not directly address the possible

contribution of *Gr8a* to the rapid evolution of mating signals across *Drosophila*.

Nevertheless, phylogenetic analyses of *Gr8a* orthologs indicate that this is a conserved,

sexually dimorphic receptor across *Drosophila* species (Figure 7A-B). Furthermore,

analysis of the *Gr8a* protein alignment revealed that in spite of the high overall sequence

conservation, the *Gr8a* protein includes at least one phylogenetically variable domain

(Figure 7C-D). This suggests a possible role of *Gr8a* in the divergence of mating

pheromone signaling systems across *Drosophila* species, yet more studies are needed

to elucidate this.

Studies in other animal species suggest that receptor pleiotropy might also play a role in

mating communication via other sensory modalities including auditory communication in

crickets (Hoy et al., 1977; Wiley et al., 2012) and visual communication in fish (Fukamachi

et al., 2009). While the specific genes and signaling pathways that mediate

communication in these species are mostly unknown, these data suggest that the genetic

coupling of signal-receptor pairs in mating communication systems might be more

common than previously thought. Therefore, the genetic tractability of *D. melanogaster*,

in combination with the diversity of mating communication systems in this species-rich

phylogenetic group, provide a unique opportunity for understanding the evolution and

mechanisms that drive and maintain mating communication systems at the genetic,

molecular, cellular, and evolutionary levels.

MATERIALS AND METHODS

Animals. Flies were maintained on a standard cornmeal medium under a 12:12 light-dark cycle at 25 Celsius. Unless specifically stated, the *D. melanogaster Canton-S (CS)* strain served as wild-type control animals. UAS-mCD8::GFP, UAS-myr::GFP*,* UAS-TNT-E, UAS-TNT-IMP-V1-A, *Gr8a-GAL4* and *Gr8a^1* fly lines were from the Bloomington Stock center. Originally in the *w^{1118}* background, the *Gr8a^1 null* allele was outcrossed for six generations into the *CS* wild-type background. *PromE(800)-GAL4* and *PromE(800)>Luciferase* were from Joel Levine at the University of Toronto. The following *Drosophila* species were obtained from the San Diego Stock Center: *D. simulans* 14011-0251.192, *D. sechellia* 14021-0248.03, *D. yakuba* 14021-0261.01, *D. erecta* 14021-0224.00, *D. ananassae* 14024-0371.16, *D. pseudoobscura* 14011-0121.104, D. *persimilis* 14011-0111.50, *D. willistoni* 14030-0811.35, *D. mojavensis* 15081-1352.23, and *D. virilis* 15010-1051.118. The UAS-*Gr8a* transgenic line was generated by cloning the *Gr8a* cDNA in the pUAST-attB vector using 5' EcoRI and 3' NotI restriction sites, followed by *ΦC31* integrase-dependent transgene integration at a Chromosome 2 *att*P landing site (2L:1476459) as previously described (Zheng et al., 2014). For genetic rescue experiments, the UAS-*Gr8a* and *Gr8a*-GAL4 lines were transgressed into the *Gr8a^1* CS genetic background.

The GFP-tagged allele of *Gr8a* was generated via CRISPR/*Cas9*-dependent editing using a modified "scarless" strategy (www.flyCRISPR.molbio.wisc.edu) as described in (Hill et al., 2019) using the sgRNA CGAGCAAGGCGGGAACGATT, and a donor plasmid containing 1) A backbone with ampicillin resistance; 2) Eye-specific dsRed reporter driven by the 3XP3 promoter, flanked by two *PiggyBac* transposase recognition sites; 3-4) The left and right homology arms, which consisted of 1kb genomic DNA fragments upstream and downstream of the sgRNA site respectively. Single base pair mutations were introduced into the PAM sequence at each sgRNA binding site on the homology arms to prevent Cas9-dependent cutting of the donor plasmid. The left homology arm included the GFP coding sequence in the last intracellular loop of the Gr8a protein. pDCC6 plasmids containing sgRNA and cas9 sequences (100 ng/µL) and the donor plasmid (500

ng/µL) were co-injected into eggs of the *yw* background. Subsequently, correct genomic

integration of the GFP tag was verified by screening for DsRed-positive animals, followed

by sequencing of genomic PCR fragments. Control lines with the same genetic

background were generated by selecting DsRed-negative animals. The final

tagged *Gr8a* allele was generated by removing the DsRed cassette via the introduction

of the *piggyBac* transposase (Bloomington #8285).

Immunohistochemistry. To visualize the expression pattern of *Gr8a* in males and

females, *Gr8a-GAL4* flies (Lee et al., 2012) were crossed to UAS-CD8::EGFP and live-

imaged at 5 days old using a Nikon-A1 confocal microscope. To demonstrate *Gr8a*

expression in oenocytes, abdomens from *Gr8a*-GAL4*/*UAS-*myr::GFP* ;

PromE(800)>Luciferase flies were dissected and immunostained as previously described

(Lu et al., 2012; Zheng et al., 2014) by using a Rabbit anti-GFP (1:1000; A-11122, Thermo

Fisher Scientific) and a mouse anti-luciferase (1:100; 35-6700, Thermo Fisher Scientific)

antibodies followed by AlexaFluor 488 anti-rabbit and AlexaFluor 568 anti-mouse

secondary antibodies (Both at 1:1000; Thermo Fisher Scientific). To visualize the GR8A

protein, abdomens of control flies and flies with CRISPR/Cas9 generated GFP-tagged

GR8A were dissected and immunostained as previously described (Lu et al., 2012; Zheng

et al., 2014) using a Rabbit anti-GFP antibody (1:1000; A-11122, Thermo Fisher

Scientific) followed by AlexaFluor 488 anti-rabbit secondary antibody (1:1000; Thermo

Fisher Scientific).

mRNA expression. Newly eclosed flies were separated by sex under CO_2 and aged for

5 days on standard cornmeal medium. On day 6, flies were placed in a -80°C freezer until

RNA extraction. To separate body parts, frozen flies were placed in 1.5ml microcentrifuge

tubes, dipped in liquid nitrogen, and then vortexed repeatedly until heads, appendages,

and bodies were clearly separated. Total RNA was extracted using the Trizol Reagent

(Thermo Fisher Scientific) separately from heads, bodies, and appendages for *Gr8a*

expression and from bodies for desaturase enzyme genes. cDNAs were synthesized

using SuperScript II reverse transcriptase (Thermo Fisher Scientific) with 500 ng total

RNA in a 20 uL reaction. Real-time quantitative RT-PCR was carried out as previously

described with *Rp49* as the loading control gene (Lu et al., 2014, 2012; Zheng et al.,
2014). Primer sequences are described in Tables 5, 6 and 7.

Courtship Behavior. Single-pair assays were performed as we have previously
published (Lu et al., 2014, 2012). In short, newly eclosed males were kept individually on
standard fly food in plastic vials (12 x 75mm). Newly eclosed virgin females were kept in
groups of 10 flies. All behaviors were done with 4-7 day-old animals, which were housed
under constant conditions of 25° C and a 12h:12h light-dark cycle. Courtship was video
recorded for 10 min for male courtship and 15 min for female mating receptivity. Male
courtship latency and index were measured as previously described (Lu et al., 2014,
2012). Female receptivity index was defined as the time from the initiation of male
courtship until copulation was observed. Unless otherwise indicated, assays were
performed under red light conditions.

Fitness measures. Three-day old virgin male-female pairs were housed together for 24
hours after which males were discarded. Subsequently, groups of five females were
allowed to lay eggs daily on fresh agar grape plates for five consecutive days. Fitness
was measured as total number of eggs laid.

Perfuming studies. Synthetic compounds were synthesized by J.G.M. Perfuming
studies were performed using a modified protocol from (Billeter et al., 2009). In short, 3
mg of each compound was dissolved in 6 mL hexane (Sigma-Aldrich #139386-500ML)
and 0.5 mL was pipetted into individual 2 mL glass vials fitted with 9mm PTFE lined caps
(Agilent Crosslab, Santa Clara, CA, USA). The hexane was evaporated under a nitrogen
gas flow, such that a residue of the compound was left around the bottom one-third of the
vial. Control vials were prepared using hexane without a spiked compound. Vials were
kept at -20°C until use. Flies used in these trials were collected as described above, kept
in single sex groups and aged for 4 days on standard cornmeal medium at 25°C. 24 hours
before perfuming, 20 flies of one or the other sex were placed in glass vials containing

standard cornmeal medium (12 x 75mm). To perfume the flies, these groups of 20 flies
were dumped without anesthesia into each 2 mL vial containing the compound of interest,
and were vortexed at medium-low speed for 3 pulses of 20 seconds punctuated by 20
second rest periods. Flies were transferred to new food vials and were allowed to recover
for one hour. Perfumed flies were then used in courtship behavior assays as described
above and the remaining flies were used in pheromone analyses to verify compound
transfer. The genotype of flies that were perfumed differed based upon the genotype with
the lower amount of each compound as determined in Figure 3 (B, C, F). In all cases,
compound transfer was verified by CHC extraction and GC/MS (Table 8).

Phylogenetic analysis. Protein sequences of GR8A orthologs from the 12 sequenced
Drosophila reference genomes were aligned by using the ClustalW algorithm in the
Omega package (Sievers et al., 2011), followed by ProtTest (v2.4) to determine the best
model of protein evolution (Abascal et al., 2005). Subsequently, Akaike and Bayesian
information criterion scores were used to select the appropriate substitution matrix. We
then used a maximum likelihood approach and rapid bootstrapping within RAxML v 7.2.8
Black Box on the Cipres web portal to make a phylogenetic tree (Miller et al., 2010).
Visualizations of the bipartition files were made using FigTree v1.3.1
(http://tree.bio.ed.ac.uk/software/figtree/).

Pheromone Analysis. Virgin flies were collected upon eclosion under a light CO_2
anesthesia and kept in single-sex vials in groups of 10 with 6 biological replications for
each genotype and sex. Virgin flies were aged for 5 days on standard cornmeal medium
at 25°C. To collect mated flies, both females and males were aged for 3 days before
single mating pairs were placed in a standard fly vial with standard cornmeal food for 24
hours. The pair was then separated for 24 hours before collection. Copulation was
confirmed by the presence of larvae in the vials of mated females several days later. On
the morning of day 5, flies were anesthetized under light CO_2 and groups of five flies were
placed in individual scintillation vials (VWR 74504-20). To extract CHCs, each group of
flies was covered by 100 uL hexane (Sigma-Aldrich #139386-500ML) containing 50µg/mL

hexacosane (Sigma-Aldrich #241687-5G) and was washed for ten minutes.
Subsequently, hexane washes were transferred into a new 2 ml glass vial containing a
350 uL insert (Thermo Scientific C4000-LV-1W) and were stored at -20°C until shipment
to the Millar laboratory.

Analyses of CHC profiles were done by gas chromatography and mass spectroscopy
(GC-MS) in the Millar laboratory at UC Riverside as previously described (Chung et al.,
2014). Peak areas were measured, and data was normalized to known quantity of internal
standard hexacosane (Sigma-Aldrich #241687-5G). The relative proportion of each
compound in each sample was calculated and used in further statistical analysis.
Qualitative data were analyzed through a permutation MANOVA using the ADONIS
function in the vegan package of R with Bray-Curtis dissimilarity measures (Oksanen et
al., 2017). Data were visualized using non-metric multidimensional scaling (vegdist
function in the vegan package of R followed by cmdscale function in the stats package
(Oksanen et al., 2017)) using Bray-Curtis dissimilarity, and either 2 or 3 dimensions in
order to minimize stress to < 0.1. Quantitative data were analyzed by using a t-test or
Mann-Whitney Rank Sum Test in R 3.3.2 (R Core Team, 2016).

Data availability. All relevant data are available from the corresponding author upon
request.

AUTHOR CONTRIBUTIONS

K.M.Z., C.V., J.G.M. and Y.B-S designed experiments. K.M.Z., C.V., N.L., X.L., S.H.,
J.G.M. and Y.B-S collected and analyzed data. K.M.Z., C.V. and Y.B-S wrote the
manuscript.

ACKNOWLEDGMENTS

We thank members of the Ben-Shahar lab for comments on earlier versions of the
manuscript. We thank the Millar laboratory (UC Riverside) for performing the CHC

analyses. We thank Joshua Krupp for assistance with perfuming studies, Nabeel

Chowdhury and Deanna Simon for assistance with qRT-PCR analysis, and Paula Kiefel

for technical help with generating transgenic flies. This work was supported by NSF grant

1545778 and NIH grant NS089834 awarded to Y. B-S. Stocks obtained from the

Bloomington Drosophila Stock Center (NIH P40OD018537) were used in this study. Wild

type *Drosophila* species were obtained from the National *Drosophila* Species Stock

Center at Cornell University.

REFERENCES

Abascal F, Zardoya R, Posada D. 2005. ProtTest: Selection of best-fit models of protein
evolution. *Bioinformatics* **21**:2104–2105. doi:10.1093/bioinformatics/bti263

Arnold SJ, Verrell PA, Tilley SG. 2006. The Evolution of Asymmetry in Sexual Isolation:
A Model and a Test Case. *Evolution (N Y)* **50**:1024. doi:10.2307/2410643

Benton R, Vannice KS, Vosshall LB. 2007. An essential role for a CD36-related receptor
in pheromone detection in Drosophila. *Nature* **450**:289–293.
doi:10.1038/nature06328

Billeter JC, Atallah J, Krupp JJ, Millar JG, Levine JD. 2009. Specialized cells tag sexual
and species identity in Drosophila melanogaster. *Nature* **461**:987–991.
doi:10.1038/nature08495

Blomquist GJ, Bagnères A-G. 2010. Insect Hydrocarbons: Biology, Biochemistry, and
Chemical Ecology. Cambridge: Cambridge University Press.

Boake CRB. 1991. Coevolution of senders and receivers of sexual signals: Genetic
coupling and genetic correlations. *Trends Ecol Evol* **6**:225–227. doi:10.1016/0169-
5347(91)90027-U

Bousquet F, Nojima T, Houot B, Chauvel I, Chaudy S, Dupas S, Yamamoto D, Ferveur
J-F. 2011. Expression of a desaturase gene, desat1, in neural and nonneural
tissues separately affects perception and emission of sex pheromones in
Drosophila. *Proc Natl Acad Sci* **109**:249–254. doi:10.1073/pnas.1109166108

Bray S, Amrein H. 2003. A putative Drosophila pheromone receptor expressed in male-
specific taste neurons is required for efficient courtship. *Neuron* **39**:1019–1029.
doi:10.1016/S0896-6273(03)00542-7

Brooks R, Hunt J, Blows MW, Smith MJ, Bussière LF, Jennions MD. 2005.
Experimental evidence for multivariate stabilizing sexual selection. *Evolution (N Y)*
59:871–880. doi:10.1111/j.0014-3820.2005.tb01760.x

Butlin RK, Ritchie MG. 1989. Genetic coupling in mate recognition systems: what is the
evidence? *Biol J Linn Soc* **37**:237–246. doi:10.1111/j.1095-8312.1989.tb01902.x

Chebib J, Guillaume F. 2019. Pleiotropy or linkage? Their relative contributions to the
genetic correlation of quantitative traits and detection by multi-trait GWA studies.
bioRxiv 656413. doi:10.1101/656413

Chenoweth SF, Blows MW. 2006. Dissecting the complex genetic basis of mate choice.
Nat Rev Genet **7**:681–692. doi:10.1038/nrg1924

Chung H, Carroll SB. 2015. Wax, sex and the origin of species: Dual roles of insect
cuticular hydrocarbons in adaptation and mating. *BioEssays* **37**:822–830.
doi:10.1002/bies.201500014

Chung H, Loehlin DW, Dufour HD, Vaccaro K, Millar JG, Carroll SB. 2014. A single
gene affects both ecological divergence and mate choice in Drosophila. *Science*
(80-) **343**:1148–1151.

Clyne PJ, Warr CG, Carlson JR. 2000. Candidate taste receptors in Drosophila.
Science (80-) **287**:1830–1834. doi:10.1126/science.287.5459.1830

Coyne JA, Crittenden AP, Mah K. 1994. Genetics of a pheromonal difference
contributing to reproductive isolation in Drosophila. *Science (80-)* **265**:1461–1464.
doi:10.1126/science.8073292

Datta SR, Vasconcelos ML, Ruta V, Luo S, Wong A, Demir E, Flores J, Balonze K,
Dickson BJ, Axel R. 2008. The Drosophila pheromone cVA activates a sexually
dimorphic neural circuit. *Nature* **452**:473–477. doi:10.1038/nature06808

Dunipace L, Meister S, Mcnealy C, Amrein H. 2001. Spatially restricted expression of
candidate taste receptors in the. *Curr Biol* 822–835.

Dweck HKM, Ebrahim SAM, Thoma M, Mohamed AAM, Keesey IW, Trona F, Lavista-
Llanos S, Svatoš A, Sachse S, Knaden M, Hansson BS. 2015. Pheromones
mediating copulation and attraction in Drosophila. *Proc Natl Acad Sci U S A*
112:E2829–E2835. doi:10.1073/pnas.1504527112

Dyer KA, White BE, Sztepanacz JL, Bewick ER, Rundle HD. 2014. Reproductive
character displacement of epicuticular compounds and their contribution to mate
choice in Drosophila subquinaria and Drosophila recens. *Evolution (N Y)* **68**:1163–
1175. doi:10.1111/evo.12335.REPRODUCTIVE

Ferveur JF. 2005. Cuticular hydrocarbons: Their evolution and roles in Drosophila
pheromonal communication. *Behav Genet* **35**:279–295. doi:10.1007/s10519-005-
3220-5

Ford CP. 2014. The role of D2-autoreceptors in regulating dopamine neuron activity and
transmission. *Neuroscience* **282**:13–22. doi:10.1016/j.neuroscience.2014.01.025

Fukamachi S, Kinoshita M, Aizawa K, Oda S, Meyer A, Mitani H. 2009. Dual control by
a single gene of secondary sexual characters and mating preferences in medaka.
BMC Biol **7**:64. doi:10.1186/1741-7007-7-64

Gutierrez E, Wiggins D, Fielding B, Gould AP. 2007. Specialized hepatocyte-like cells
regulate Drosophila lipid metabolism. *Nature* **445**:275–280.
doi:10.1038/nature05382

Hill AS, Jain P, Folan NE, Ben-Shahar YY. 2019. The Drosophila ERG channel seizure
plays a role in the neuronal homeostatic stress response. *PLoS Genet* **15**:1–24.
doi:10.1371/journal.pgen.1008288

Howard RW, Blomquist GJ. 2005. Ecological, Behavioral, and Biochemical Aspects of
Insect Hydrocarbons*. *Annu Rev Entomol* **50**:371–393.
doi:10.1146/annurev.ento.50.071803.130359

Hoy RR, Hahn J, Paul RC. 1977. Hybrid cricket auditory behavior: Evidence for genetic
coupling in animal communication. *Science (80-)* **195**:82–84.
doi:10.1126/science.831260

Hu Y, Han Y, Shao Y, Wang X, Ma Y, Ling E, Xue L. 2015. Gr33a modulates drosophila
male courtship preference. *Sci Rep* **5**:1–7. doi:10.1038/srep07777

Kallman BR, Kim H, Scott K. 2015. Excitation and inhibition onto central courtship
neurons biases drosophila mate choice. *Elife* **4**:1–18. doi:10.7554/eLife.11188

Kirkpatrick M, Hall DW. 2004. Sexual Selection and Sex Linkage All use subject to
https://about.jstor.org/terms. *Evolution (N Y)* **58**:683–691.

Koh T, He Z, Gorur-shandilya S, Menuz K, Larter NK, Carlson JR. 2015. Candidate
Taste and Pheromone Receptors. *Neuron* **83**:850–865.

doi:10.1016/j.neuron.2014.07.012.The

Krupp JJ, Billeter JC, Wong A, Choi C, Nitabach MN, Levine JD. 2013. Pigment-
Dispersing Factor Modulates Pheromone Production in Clock Cells that Influence
Mating in Drosophila. *Neuron* **79**:54–68. doi:10.1016/j.neuron.2013.05.019

Kurtovic A, Widmer A, Dickson BJ. 2007. A single class of olfactory neurons mediates
behavioural responses to a Drosophila sex pheromone. *Nature* **446**:542–546.
doi:10.1038/nature05672

Labeur C, Dallerac R, Wicker-Thomas C. 2002. Involvement of desat1 gene in the
control of Drosophila melanogaster pheromone biosynthesis. *Genetica* **114**:269–
274. doi:10.1023/A:1016223000650

Lacaille F, Everaerts C, Ferveur JF. 2009. Feminization and alteration of Drosophila
taste neurons induce reciprocal effects on male avoidance behavior. *Behav Genet*
39:554–563. doi:10.1007/s10519-009-9286-8

Lacaille F, Hiroi M, Twele R, Inoshita T, Umemoto D, Manière G, Marion-Poll F, Ozaki
M, Francke W, Cobb M, Everaerts C, Tanimura T, Ferveur JF. 2007. An inhibitory
sex pheromone tastes bitter for Drosophila males. *PLoS One* **2**:1–7.
doi:10.1371/journal.pone.0000661

Lande R. 1984. The genetic correlation between characters maintained by selection,
linkage and inbreeding. *Genet Res* **44**:309–320. doi:10.1017/S0016672300026549

Lande R. 1981. Models of speciation by sexual selection on polygenic traits. *Proc Natl*
Acad Sci **78**:3721–3725.

Lande R. 1980. The genetic covariance between characters maintained by pleiotropic
mutations. *Genetics* **94**:203–215.

Lebreton S, Grabe V, Omondi AB, Ignell R, Becher PG, Hansson BS, Sachse S,
Witzgall P. 2014. Love makes smell blind: Mating suppresses pheromone attraction
in Drosophila females via Or65a olfactory neurons. *Sci Rep* **4**:1–6.
doi:10.1038/srep07119

Lee Y, Kang MJ, Shim J, Cheong CU, Moon SJ, Montell C. 2012. Gustatory receptors

required for avoiding the insecticide L-canavanine. *J Neurosci* **32**:1429–1435.

doi:10.1523/JNEUROSCI.4630-11.2012

Lee Y, Moon SJ, Montell C. 2009. Multiple gustatory receptors required for the caffeine

response in Drosophila. *Proc Natl Acad Sci U S A* **106**:4495–4500.

doi:10.1073/pnas.0811744106

Lofstedt C, Hansson BS, Roelofst W, Bengtsson B 0. 1989. No Linkage Between Genes

Controlling Female Pheromone Production and Male Pheromone Response in the

European Corn Borer, Ostm'nia nubilalis Hubner (Lepidoptera; Pyralidae).

Lu B, LaMora A, Sun Y, Welsh MJ, Ben-Shahar Y. 2012. ppk23-Dependent

chemosensory functions contribute to courtship behavior in Drosophila

melanogaster. *PLoS Genet* **8**:e1002587. doi:10.1371/journal.pgen.1002587

Lu B, Zelle KM, Seltzer R, Hefetz A, Ben-Shahar Y. 2014. Feminization of pheromone-

sensing neurons affects mating decisions in Drosophila males. *Biol Open* **3**:152–

160. doi:10.1242/bio.20147369

Makki R, Cinnamon E, Gould AP. 2014. The Development and Functions of Oenocytes.

Annu Rev Entomol **59**:405–425. doi:10.1146/annurev-ento-011613-162056

McKinney RM, Vernier C, Ben-Shahar Y. 2015. The neural basis for insect pheromonal

communication. *Curr Opin Insect Sci* **12**:86–92. doi:10.1016/j.cois.2015.09.010

Miller M, Pfeiffer WT, Schwartz T. 2010. Creating the CIPRES Science Gateway for

Inference of Large Phylogenetic Trees, Proceedings of the Gateway Computing

Environments Workshop: 14 Nov 2010. doi:10.1109/GCE.2010.5676129

Miyamoto T, Amrein H. 2008. Suppression of male courtship by a Drosophila

pheromone receptor. *Nat Neurosci* **11**:874–876. doi:10.1038/nn.2161

Moon SJ, Lee Y, Jiao Y, Montell C. 2009. A Drosophila gustatory receptor essential for

aversive taste and inhibiting male-to-male courtship. *Curr Biol* **19**:1623–1627.

doi:10.1038/jid.2014.371

Mucignat-Caretta C, editor. 2014. Neurobiology of Chemical Communication. CRC

Press.

Ng SH, Shankar S, Shikichi Y, Akasaka K, Mori K, Yew JY. 2014. Pheromone evolution
and sexual behavior in Drosophila are shaped by male sensory exploitation of other
males. *Proc Natl Acad Sci U S A* **111**:3056–3061. doi:10.1073/pnas.1313615111

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR,
O'Hara RB, Simpson GL, Solymos P, Henry M, Stevens H, Szoecs E, Wagner H.
2017. vegan: Community Ecology Package.

Park JH, Kwon JY. 2011. A systematic analysis of drosophila gustatory receptor gene
expression in abdominal neurons which project to the central nervous system. *Mol*
Cells **32**:375–381. doi:10.1007/s10059-011-0128-1

Poudel S, Kim Y, Kim YT, Lee Y. 2015. Gustatory receptors required for sensing
umbelliferone in Drosophila melanogaster. *Insect Biochem Mol Biol* **66**:110–118.
doi:10.1016/j.ibmb.2015.10.010

Robertson HM, Warr CG, Carlson JR. 2003. Molecular evolution of the insect
chemoreceptor gene superfamily in Drosophila melanogaster. *Proc Natl Acad Sci U*
S A **100**:14537–14542. doi:10.1073/pnas.2335847100

Scott K, Brady R, Cravchik A, Morozov P, Rzhetsky A, Zuker C, Axel R. 2001. A
chemosensory gene family encoding candidate gustatory and olfactory receptors in
Drosophila. *Cell* **104**:661–673. doi:10.1016/S0092-8674(01)00263-X

Shankar S, Chua JY, Tan KJ, Calvert ME, Weng R, Ng WC, Mori K, Yew JY. 2015. The
neuropeptide tachykinin is essential for pheromone detection in a gustatory neural
circuit. *Elife* **4**:1–23. doi:10.7554/elife.06914

Shaw KL, Ellison CK, Oh KP, Wiley C. 2011. Pleiotropy, "sexy" traits, and speciation.
Behav Ecol **22**:1154–1155. doi:10.1093/beheco/arr136

Shaw KL, Lesnick SC. 2009. Genomic linkage of male song and female acoustic
preference QTL underlying a rapid species radiation. *Proc Natl Acad Sci U S A*
106:9737–9742. doi:10.1073/pnas.0900229106

Shim J, Lee Y, Jeong YT, Kim Y, Lee MG, Montell C, Moon SJ. 2015. The full repertoire
of Drosophila gustatory receptors for detecting an aversive compound. *Nat*
Commun **6**. doi:10.1038/ncomms9867

Shirangi TR, Dufour HD, Williams TM, Carroll SB. 2009. Rapid evolution of sex
pheromone-producing enzyme expression in Drosophila. *PLoS Biol* **7**.
doi:10.1371/journal.pbio.1000168

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation
of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst*
Biol **7**. doi:10.1038/msb.2011.75

Singh ND, Shaw KL. 2012. On the scent of pleiotropy. *Proc Natl Acad Sci* **109**:5–6.
doi:10.1073/pnas.1118531109

Stagkourakis S, Kim H, Lyons DJ, Broberger C. 2016. Dopamine Autoreceptor
Regulation of a Hypothalamic Dopaminergic Network. *Cell Rep* **15**:735–747.
doi:10.1016/j.celrep.2016.03.062

Steiger S, Schmitt T, Martin Schaefer H. 2011. The origin and dynamic evolution of
chemical information transfer. *Proc R Soc B Biol Sci* **278**:970–979.
doi:10.1098/rspb.2010.2285

Svensson EI, Karlsson K, Friberg M, Eroukhmanoff F. 2007. Gender Differences in
Species Recognition and the Evolution of Asymmetric Sexual Isolation. *Curr Biol*
17:1943–1947. doi:10.1016/j.cub.2007.09.038

Sweigart AL. 2010. The genetics of postmating, prezygotic reproductive isolation
between Drosophila virilis and D. americana. *Genetics* **184**:401–410.
doi:10.1534/genetics.109.111245

Symonds MRE, Elgar MA. 2008. The evolution of pheromone diversity. *Trends Ecol*
Evol **23**:220–228. doi:10.1016/j.tree.2007.11.009

Team RC. 2016. R: A language and environment for statistical computing.

Thistle R, Cameron P, Ghorayshi A, Dennison L, Scott K. 2012. Contact
chemoreceptors mediate male-male repulsion and male-female attraction during
drosophila courtship. *Cell* **149**:1140–1151. doi:10.1016/j.cell.2012.03.045

Toda H, Zhao X, Dickson BJ. 2012. The Drosophila Female Aphrodisiac Pheromone

Activates ppk23+ Sensory Neurons to Elicit Male Courtship Behavior. *Cell Rep*
1:599–607. doi:10.1016/j.celrep.2012.05.007

van der Goes van Naters W, Carlson JRR. 2007. Receptors and Neurons for Fly Odors
in Drosophila. *Curr Biol* **17**:606–612. doi:10.1016/j.cub.2007.02.043

Wang Z, Singhvi A, Kong P, Scott K. 2004. Taste representations in the Drosophila
brain. *Cell* **117**:981–991. doi:10.1016/j.cell.2004.06.011

Watanabe K, Toba G, Koganezawa M, Yamamoto D. 2011. Gr39a, a Highly diversified
gustatory receptor in drosophila, has a role in sexual behavior. *Behav Genet*
41:746–753. doi:10.1007/s10519-011-9461-6

Wiley C, Ellison CK, Shaw KL. 2012. Widespread genetic linkage of mating signals and
preferences in the Hawaiian cricket Laupala. *Proc R Soc B Biol Sci* **279**:1203–
1209. doi:10.1098/rspb.2011.1740

Wyatt TD. 2014. Pheromones and Animal Behavior: Chemical Signals and Signatures,
2nd ed. Cambridge: Cambridge University Press.

Yew JY, Chung H. 2015. Insect pheromones: An overview of function, form, and
discovery. *Prog Lipid Res*. doi:10.1016/j.plipres.2015.06.001

Yew JY, Dreisewerd K, Luftmann H, Müthing J, Pohlentz G, Kravitz EA. 2009. A New
Male Sex Pheromone and Novel Cuticular Cues for Chemical Communication in
Drosophila. *Curr Biol* **19**:1245–1254. doi:10.1016/j.cub.2009.06.037

Zheng X, Valakh V, DiAntonio A, Ben-Shahar Y. 2014. Natural antisense transcripts
regulate the neuronal stress response and excitability. *Elife* **3**.
doi:10.7554/elife.01849

Figures and figure legends:

Figure 1. *Gr8a* **is a sexually dimorphic chemosensory receptor enriched in male**
oenocytes and fat body. (A) *Gr8a* promoter activity in proboscis, (B) forelegs, and (C)
abdomens of males (top panels) and females (bottom panels). (D) *Gr8a* mRNA
expression. Relative mRNA levels were measured by real-time quantitative RT-PCR. **,
p<0.01 Mann Whitney Rank Sum Test, n=4/group. (E) Confocal z-stack image of
Gr8a>EGFP in abdominal cells. (F) Confocal z-stack image of *desat1*>Luciferase in
abdominal cells. (G) Co-expression of *Gr8a* and *desat1*. (H) Confocal z-stack image of
GFP-tagged GR8A protein in abdominal cells. Green, *Gr8a*; Red, *desat1*; Blue, nuclear
DAPI stain. Orange arrowhead, fat body cells; white arrowhead, oenocytes. Scale bar =
100μm.

Figure 2. *Gr8a* activity contributes to the perception and production of an inhibitory signal associated with mating decisions in males and females. (A) Blocking neural activity in female *Gr8a*-expressing sensory neurons shortens copulation latency. Homozygous (B) or hemizygous (C) *Gr8a* null females show shortened copulation latency relative to wild-type controls. Df(1)BSC663 is a deficiency that covers the *Gr8a* locus. Df(1)BSC754 was used as a control. (D) Expression of *Gr8a* cDNA with *Gr8a* promoter (Gr8a-;Gr8a^tg) rescues the copulation latency phenotype in *Gr8a* mutant females. (E) Wild-type females exhibit shorter copulation latency when courted by *Gr8a* mutant relative to wild-type males. (F) *Gr8a* mutant males do not recognize mating status of females, and have a reduced transfer of inhibitory mating pheromones during copulations. Female, female genotype; Sperm donor, genotype of males mated first with focal females; Focal male, genotypes of experimental males presented with mated females. Different letters above bars indicate statistically significant *post hoc* contrasts between groups (panels C, D, and F; p<0.05 ANOVA, n>15/group). *, p<0.05, Mann Whitney Rank Sum Test, n>15/group).

Figure 2—figure supplement 1. *Gr8a* has no effect on male Courtship Index or

Latency toward wild-type females. (A) Courtship Index and Latency (s) of control *Gr8a-*

gal4/UAS-IMP-TNT- V1A (*TNT*^{inactive}) and *Gr8a-gal4/UAS-TNT-E* mutant males

towards wild-type females. (B) Courtship index and latency (s) of wild-type (*CS*) and

Gr8a null males toward wild-type decapitated females. Mann Whitney Rank Sum Test,

not significant (p>0.05), n=15/group.

Figure 3. The *Gr8a* mutation affects the pheromone profiles of males and mated females. (A) Nonmetric multidimensional scaling (NMDS) plot of CHC profiles of wild-type and *Gr8a* mutant males. p<0.001, permutation MANOVA. **(B-C)** The effect of the *Gr8a* mutation on levels of individual CHCs in males. (B) CHCs found at high proportions in males. (C) CHCs found at low proportions in males. Only affected CHCs are shown.

See Table 2 for the complete list. *, $p<0.05$, **, $p<0.001$, Student's t-test or Mann Whitney Rank Sum Test, n=6 (Gr8a-) or 7 (wt). **(D)** The effect of *Gr8a* mutation on the expression level of several desaturase genes. Only affected genes are shown. See Table 3 for the complete list. *, $p<0.05$, Student's t-test, n=4/group. **(E)** NMDS plot of CHC profiles of females mated with wild-type or *Gr8a* mutant males. $p=0.570$, permutation MANOVA. **(F)** Nonacosane (C_{29}) differs between females mated with wild-type and *Gr8a* mutant males. See Table 4 for complete list of mated-female CHCs. *, $p<0.05$, Student's t-test, n=6/group.

Figure 4. *Gr8a*-associated alkenes inhibit normal courtship behaviors when males

are perfumed. Perfuming males with exaggerated amounts of several alkenes increases

copulation latency compared to control males. Asterisks above bars indicate statistically

significant contrasts compared to control flies, $p < 0.05$, Kruskal-Wallis Test followed by

Dunn's Test, n=15/group.

Figure 5. *Gr8a*-associated alkenes inhibit normal courtship behaviors when females are perfumed. Perfuming females with $9\text{-}C_{25}$ increases courtship latency (A), decreases courtship index (B), but does not affect copulation latency compared to control females. Perfuming females with $7\text{-}C_{25}$ does not affect courtship latency (D) or index (E), but increases copulation latency (F) compared to control females. Perfuming females with $7\text{-}C_{75}$ does not affect courtship latency (G), courtship index (H), or copulation latency (I) compared to control females. Asterisks above bars indicate statistically significant contrasts compared to control flies, $p<0.05$, Mann Whitney Rank Sum Test, n=15/group.

Figure 6. Model for the pleiotropic action of *Gr8a* in the perception and production of pheromones. (A) *Gr8a* functions as a chemoreceptor for an inhibitory signal in pheromone-sensing GRNs of males and females. (B) *Gr8a* also functions as a CHC autoreceptor in oenocytes, which regulates CHC secretion [1] or CHC synthesis [2] via signaling feedback loops [3].

Figure 7. *Gr8a* expression is sexually dimorphic across the *Drosophila* genus. (A)

Phylogenetic tree of *Drosophila Gr8a* proteins. Substitution rate = 0.2. (B) *Gr8a* mRNA

expression is enriched in males relative to females across *Drosophila*. Black, males;

white, females. *, p<0.05; **,p<0.01; Mann Whitney Rank Sum Test, n=4/group. Live *D.*

grimshawi was not analyzed because live specimens were not available at the

Drosophila Species Stock Center (DSSC). (C) Multiple aligned amino acid sequences of

Gr8a protein sequences from 12 species across *Drosophila.* Box highlights a putative

hypervariable protein domain, which is shown at a higher resolution in (D). Numbers on

top of alignment indicate amino acid number. Black, 100% identical; Dark Gray, 80-

100% similar; Light Gray, 60-80% similar; White, less than 60% similar (Blosum62 score

matrix, threshold=1). Bars below consensus represent overall level of amino acid

conservation.

Tables and table legends

Gene	Male	Female
Gr2a	-	+
Gr8a	+	-
Gr10a	+	+
Gr21a	-	+
Gr22a	+	-
Gr22e	+	+
Gr36c	+	-
Gr58c	+	+
Gr59a	+	+
Gr59b	+	+
Gr63a	+	-
Gr64a	+	-
Gr64b	+	+
Gr64c	+	+
Gr64d	+	-
Gr66a	+	+
Gr89a	+	+
Gr93a	-	+
Gr93d	+	+
Gr97a	+	+
Gr98a	+	+
Gr98b	-	+
Gr98c	+	+
Gr98d	+	+

781   **Table 1.** Candidate *Gr* genes in male and female abdomen. Plus sign indicates presence

782   of PCR product and minus signs indicates PCR product not detected. Genes with no PCR

783   product detected in male or female abdomens not shown.

R.T.	Compound	wt % total	*Gr8a* % total	p value
12.31	C_{21}	0.589	1.102	<0.001
13.24	Unknown	0.071	0.191	<0.001
14.2	C_{22}	0.893	0.949	0.339
14.34	$7\text{-}C_{22}$	0.362	0.490	<0.001
15.25	Unknown	0.128	0.226	<0.001
16.22	C_{23} & $9\text{-}C_{23}$	13.682	12.671	<0.001
16.4	$7\text{-}C_{23}$	30.201	35.478	0.002
16.53	$5\text{-}C_{23}$	2.772	2.389	<0.001
16.71	CvA	10.240	15.700	<0.001
18.03	C_{24} & $9\text{-}C_{24}$	0.578	0.367	<0.001
18.19	$8\text{-}C_{24}$	0.742	0.493	<0.001
18.27	$7\text{-}C_{24}$	0.579	0.402	<0.001
18.37	$6\text{-}C_{24}$	0.395	0.233	<0.001
18.46	$5\text{-}C_{24}$	0.040	0.047	0.943
19.09	$2Me\text{-}C_{24}$	2.426	3.240	<0.001
19.95	C_{25}	0.000	2.038	0.001
20.02	C_{25} & $9\text{-}C_{25}$	6.195	1.793	<0.001
20.18	$7\text{-}C_{25}$	14.781	4.344	<0.001
20.42	$5\text{-}C_{25}$	0.296	0.000	<0.001
22.89	$2Me\text{-}C_{26}$	6.007	6.933	0.002
23.7	C_{27}	1.127	0.719	0.052
23.94	$7\text{-}C_{27}$	0.308	0.083	<0.001
26.46	$2Me\text{-}C_{28}$	4.992	5.775	0.078
27.25	C_{29}	0.351	0.317	0.574
29.89	$2Me\text{-}C_{30}$	1.245	2.353	<0.001

785 **Table 2.** Male CHCs. Retention time (R.T.), compound, and percent total (% total) of each

786 compound as part of the total pheromonal bouquet for wild-type (wt) and *Gr8a* mutant

787 males.

Gene	wt mRNA fold difference	*Gr8a* mRNA fold difference	p value
desat1	1.282	0.931	0.037
desat2	1.413	1.270	0.506
CG8630	0.951	1.429	0.012
CG9747	0.838	0.525	0.343
CG9743	1.060	0.959	0.373
CG15331	0.774	1.000	0.21

789 **Table 3.** Desaturase gene expression. Relative mRNA expression of each desaturase

790 gene for wild-type (wt) and *Gr8a* mutant males.

R.T.	Compound	wt % total	*Gr8a* % total	p value
12.31	C_{21}	0.234	0.242	0.712
14.2	C_{22}	0.330	0.352	0.512
14.34	$7\text{-}C_{22}$	0.029	0.029	0.958
15.25	Unknown	0.238	0.200	0.128
16.09	C_{23}	6.481	6.580	0.748
16.4	$7\text{-}C_{23}$	2.454	2.729	0.149
16.39	$7,11\text{-}C_{23}$	0.473	0.426	0.208
16.53	$5\text{-}C_{23}$	0.254	0.235	0.416
16.71	CvA	1.349	1.135	0.233
17.99	C_{24}	0.660	0.732	0.209
18.19	$8\text{-}C_{24}$	0.156	0.174	0.334
19.09	$2Me\text{-}C_{24}$	1.214	1.104	0.155
19.95	C_{25}	5.404	5.683	0.336
20.02	C_{25} & $9\text{-}C_{25}$	3.141	3.851	0.699
20.18	$7\text{-}C_{25}$	3.144	2.999	0.422
20.25	$7,11\text{-}C_{25}$	1.822	1.684	0.315
20.42	$5\text{-}C_{25}$	0.526	0.555	0.375
20.47	$5,9\text{-}C_{25}$	0.686	0.704	0.589
22.89	$2Me\text{-}C_{26}$	10.010	9.247	0.1554
23.7	C_{27}	3.616	3.393	0.24
23.8	$9\text{-}C_{27}$	2.498	2.739	0.937
23.94	$7\text{-}C_{27}$	3.819	4.885	0.24
24.1	$7,11\text{-}C_{27}$	21.464	19.770	0.18
24.28	$5,9\text{-}C_{27}$	2.557	2.438	0.12
25.85	$7,11\text{-}C_{28}$	0.658	0.645	0.573
26.46	$2Me\text{-}C_{28}$	4.779	5.022	0.306
27.25	C_{29}	0.577	0.693	0.01
27.7	$7,11\text{-}C_{29}$	19.411	19.589	0.824
29.89	$2Me\text{-}C_{30}$	1.359	1.465	0.17
31.03	$7,11\text{-}C_{31}$	0.562	0.606	1

792 **Table 4.** Mated-female CHCs. Retention time (R.T.), compound, and percent total (%

793 total) of each compound as part of the total pheromonal bouquet for females mated with

794 wild-type (wt) or *Gr8a* mutant males.

Gene	Forward Primer	Reverse Primer
Gr10a	AGCGCGAGAGCTTTCGGATGC	AAAGTCCACACAGCATGGGCGG
Gr10b	GCGACTGTCCAGAGGATGTGGC	CAGCTGATCAAAGCGAATGCGC
Gr21a	AACTACCGGCTCGCCGATCG	AGGTTGCCCCGTGGTCGATGA
Gr22a	TGGGACCTTTGGCAGGGAATCG	TGCGATGACTGCAGAAGAGGGT
Gr22b	GGTTGTGCATTGCCGCATGCG	CTGCAAAGCCAAGCGAACTCGT
Gr22c	GCCGCCAGCGATCTAGCTGAG	CAATGACCGCACAGCAATGCGA
Gr22e	TGGTCTTCGTTCAAGCGCTGGT	GCCGGCGATGGCTGCAGAATAAT
Gr22f	GATCGCCTGCTGAAGCTGAGTGA	GTGGGTGCACAGCCATGCGAAT
Gr23a	AGCGATTTGTGGTGACCGCCA	CATCCGCTGCCTCGCGTTGA
Gr28a	TCGCCCTGCCGCACATCAATA	GGGTGAGAGTTCCAAGGCGCT
Gr28b	ACTGGAGGCATAGTGCACTCCCT	AGTGGTCAAGGCCCCGCTGAT
Gr2a	TTTGCGACCGAACGGCCCAG	GTTGAAGAAGCCAGCGGCGC
Gr32a	ACACGGTCACCACCAACTGCT	ACCAAGTATGTTGTGACGGCCGAA
Gr33a	ATCGCCACGCACCCCTTACC	GCTCACCACAAAGCAGGCCG
Gr36a	GAGGTCGTGTCGTTGCAGCCC	GTGCTCGCTGACGCCATCTGT
Gr36b	TGGCATGCAAGGGCTTATGGCT	CCGCAGTTGAGCCACGCGTG
Gr36c	CGGCAGAGCAAACATGTTGCACG	ACTGCAAGCCCAGTCCCAGGT
Gr39a	TACCCCGAACCGGGACTGGG	CCAGAGCGAAAAATCCATAAGCGGT
Gr39b	GGGCGCCAACTGCACATTGG	TCCAATCGAGGGATGGCAGGACA
Gr43a	GCTGGCCCTGGCTCCATATGC	AACCCCTTGTTCTTGGCTGCGGC
Gr47a	GGTCGGGCCAGAGAAATCGGC	CCCGCTTGGGATGGAGGGCC
Gr47b	GCCCTGGGATCCCGAGTACG	TGCCCGATGTCAACTGGTGATCT
Gr57a	GGCAAGGAGATGCCTTCGCCT	GCCCATCTTCTGGGTCATGGCG
Gr58a	TGTGGATGACGTGACGCGGA	ACGGATGACAACGAAGGTGATCCGA
Gr58b	ACCTCCTGCAACAACACTGGCC	CCGATTCTGCCTCAGCTGCATGG
Gr58c	ACCGCTGTAAGTGGTTGGGACT	TCCAACGATGTGCAGACGTCTCA
Gr59a	GCCAAAGTTCTTCGCCCCCGA	TCCAATCGCGTGCTGCTCTCG
Gr59b	CAGCCAAGCACTCGTGGAGTGA	TGGCAAACCACATGCTGCGATT
Gr59c	CCGAGTGCTGGCGGATCGAAC	CTGGCCAGTTGCAGTTGGAAGT
Gr59d	CGTCGATTGCTGGATCAACGCG	TCGACCAAACTCGAAAAGGCCG
Gr59e	CCTCTCTGGGAGGAGTCCGTGC	AGGGCAACGCCCATTAGCGACT
Gr59f	GCCATGCACGTGGGAAAGGTGT	ACATTGCCCTGAACCGACTTGGA
Gr5a	GCCTGCAACGTGCTCGTCCT	TTGAGCGGGACAAGGCGCAG
Gr61a	AAAGTGCGGCGCCAGAAGCA	GTGCGACCAGGAGCGAGCAA
Gr63a	CGCAAGAAGGGTGACGCGGTG	AAGGGTCCACTCAGGGAGCGC
Gr64a	CGCGTGCAATTGGCTGCCTTG	GTCCGCCGCCGTGAGGAATAC
Gr64b	GCCCGCGGTCTTTTGGACAGA	CATGGCCAGGACGAGTGAGCG
Gr64c	ATGCACTGGCGCATCAGGCA	CCGCCCTCGCAGAAGCTCTT

Gr64d	ACCGCGCCTGGATGGTGTTC	TGCACCACGGCACCTTGCAT
Gr64e	TGGCCACCACTTGCTCGCTG	ACGCCGCGTGTTAGCCGAAA
Gr64f	TCTGGCTAGGCAGTGGCCCC	GCATGGGCCACCGATGGCAT
Gr66a	GAGGACCGCAGCCTAAAGGCC	CGCGGTCAACTGGTCGGAGAC
Gr77a	CGCAGTTCAGGCCGTCTGCTT	CCACAGGCGCTGACGGCAAATA
Gr85a	CTCTCGCCTACATGGCACGC	ATAGTGACGTGCGTGTGCTGC
Gr89a	GCTGCATGGAGTTCCCCTGACC	ACATTGCCAGTGGGCGTGAGC
Gr8a	ACATACGCATCGCCGTGGACTG	ATGTAGGTGCCCACGGAACAGG
Gr92a	TGTCTCCGTTGAGGGAGCGGT	TGGGTGACATCGAATAGGCCCA
Gr93a	GGTGAGGCAGTCAGAAGTGCC	CCTCGGACGCCTCAAAGCCAC
Gr93b	ACCACGCCATTCTGCGAAGAC	GCCCAAAGGAAACACTCGCAGC
Gr93c	TGCGACGCCTGAGTCTGGAGA	CCCAAGGGTCTAACGCGAAACTCG
Gr93d	ACTCGGGAACGTGCTCTGGAT	AGAGGCCCAACAAGTTAACCCTGA
Gr97a	TGCGTCGCTTCTGCGAGTTAGC	GCGTGCTGCAGATCGAATCGC
Gr98a	GTTGAGCACACAAGGGTGGCTG	GCTGTTGCTTGCTGCTGCTCTGA
Gr98b	TTGGAGGCTGGAGGGCGATGT	ATGGTGACAAGCAACCCGCCA
Gr98c	GCCAGGATTCAGGAGCTGTGCG	CGCTGGTATTTGTCTGTAGCCGA
Gr98d	CTGGCTGGACGCATTTGGGGC	ATGTCAAAGAGGCCACCGCATGT
Gr9a	TGTCCTCTGTCTGGGCGACTGG	ACAAGGGACACGAACTGCAGGA

Table 5. Nucleotide sequences for qRT-PCR primers for *D. melanogaster Gr* genes.

Species	*Gr8a* Forward Primer	*Gr8a* Reverse Primer
D. melanogaster	TGACCATCAACATACGCATCG	CGTATATGAAGGCGGGAATCTC
D. simulans	GAACTTTTCGCTGCAACTCC	ACTTCGGTATAAACTGGATGGTG
D. sechellia	GAGATTCCCGCCTTCATATACG	GAGTTGCAGCGAAAAGTTCTG
D. erecta	CAGATTCAGAACTTTTCGCTGC	GGTGTAGATCATGTAGGTGCC
D. yakuba	TGCCTCGGACTAACAATTCTG	GTGTAGATCATGTAGGTGCCC
D. ananassae	AATGTACCGAAGTTTCCAGGG	GCGGGTATGATCAGGAAATAGTC
D. pseudoobscura	CCCGTTTCCGTGACAATATTG	ACCATCTACATATCCGTTGCC
D. persimilis	TTTCGCTTCTCCACACTGAC	AGGCGGGCAATATCAAAGAG
D. willistoni	GAAATGTTGCCCAGAATAGCC	CCCAAAGCATGTATAACCACTG
D. virilis	TCTTCAGATCCAAAACTTTTCGC	TTGGGCATCAGTTGTACGG
D. mojavensis	CATATACCCGCCTTTCTCTACAC	GTTCGTGCAGAATTTGTAGCG

Species	*Rp49* Forward Primer	*Rp49* Reverse Primer
D. melanogaster	ATCTTGGGCCTGTATGCTG	TGTGATGGGAATTCGTGGG
D. simulans	GTCGGATCGATATGCTAAGCTG	CAGATACTGTCCCTTGAAGCG
D. sechellia	CATACAGGCCCAAGATCGTG	CAGATACTGTCCCTTGAAGCG
D. erecta	GTCGGATCGATATGCTAAGCTG	CAGATACTGTCCCTTGAAGCG
D. yakuba	CATACAGGCCCAAGATCGTG	GGCATCAGATACTGTCCCTTG
D. ananassae	TACAGGCCCAAGATCGTTAAG	GTACTGACCCTTGAAGCGAC
D. pseudoobscura	CCAGCTCCAAAATGACGATTC	TCAATACCCTTAGGCTTGCG
D. persimilis	AAGCACTTCATCCGTCACC	TCAATACCCTTAGGCTTGCG
D. willistoni	AAGCACTTCATCCGTCACC	GTTGGGCATCAGATATTGGC
D. virilis	AGTCGGATCGTTATGCTAAGTTG	TGGAGGGTACGCTTGTTTG
D. mojavensis	ACCATTCGTCCAGCATACAG	TTGGCCCTTGAAGCGAC

Table 6. Nucleotide sequences for qRT-PCR primers for *D. melanogaster Gr8a*, *Rp49* and orthologs.

Gene	Forward Primer	Reverse Primer
desat1	TTCTGAACGTAACCTGGCTG	GGGAAGACGTGATGGTAGTTATG
desat2	TTTACTGGAATGAGACGCTGG	CTGGGTAGGATTCATGGTCTTG
Fad2	GTGCCTCAGTCTCAATCTCATC	GTGGTAGTTGTGGTATCCCTC
CG8630	ATGTGATGGCTAAGATCGGAC	TGGTCGGAATGGCACTATTG
CG9747	TCAAGTGGGACAAAGTCATCC	GCCGTAACTCCGAATCCG
CG9743	GGTACTTCTGGAACGAGGATC	TCAGTGGACATGAGGTTCTTG
CG15531	CTCAAGGATGTGGATATGTCGG	ATGGAGGTGGCTAGGGAG
rp49	CACCAAGCACTTCATCCG	TCGATCCGTAACCGATGT

801 **Table 7.** Nucleotide sequences for qRT-PCR primers for *D. melanogaster* desaturase

802 enzyme genes.

Compound	Genotype	Sex	ng/sample	ng/fly
C_{25}	wt	male	1350	270
C_{25}	wt	male	1170	234
C_{25}	wt	male	27730	5546
$9\text{-}C_{25}$	Gr8a-	male	67390	13478
$9\text{-}C_{25}$	Gr8a-	male	59380	11876
$9\text{-}C_{25}$	Gr8a-	male	1000	200
$7\text{-}C_{25}$	Gr8a-	male	43790	8758
$7\text{-}C_{25}$	Gr8a-	male	31330	6266
$7\text{-}C_{25}$	Gr8a-	male	24130	4826
$5\text{-}C_{25}$	Gr8a-	male	12050	2410
$5\text{-}C_{25}$	Gr8a-	male	14160	2832
$5\text{-}C_{25}$	Gr8a-	male	1870	374
C_{27}	Gr8a-	male	890	178
C_{27}	Gr8a-	male	720	144
C_{27}	Gr8a-	male	263	52.6
$7\text{-}C_{27}$	Gr8a-	male	36710	7342
$7\text{-}C_{27}$	Gr8a-	male	21250	4250
$7\text{-}C_{27}$	Gr8a-	male	15910	3182
C_{29}	wt	male	260	52
C_{29}	wt	male	340	68
C_{29}	wt	male	890	178
$9\text{-}C_{25}$	wt	female	14060	2812
$9\text{-}C_{25}$	wt	female	13830	2766
$7\text{-}C_{25}$	wt	female	23200	4640
$7\text{-}C_{25}$	wt	female	12780	2556
$7\text{-}C_{27}$	wt	female	1010	202
$7\text{-}C_{27}$	wt	female	1080	216

Table 8. Amount (ng) of each perfumed compound measured in each sample of perfumed flies (5 flies per sample).