INTRACELLULAR PASSAGE OF Na$^+$ IN AN ACTIVE STATE G-PROTEIN COUPLED RECEPTOR

Owen N. Vickery[1,2], Catarina A. Carvalheda[1,2], Saheem A. Zaidi[3], Andrei V. Pisliakov[1,2], Vsevolod Katritch[3,4], Ulrich Zachariae[1,2,*]

[1]School of Life Sciences, University of Dundee, Dundee DD1 5EH, UK.

[2]School of Science and Engineering, University of Dundee DD1 4NH, UK.

[3]Departments of Biological Sciences and [4]Chemistry, Bridge Institute, University of Southern California, Los Angeles, CA 90089, USA

*Corresponding author; email: u.zachariae@dundee.ac.uk

ABSTRACT

Playing a central role in cell signalling, GPCRs have evolved into the largest superfamily of membrane proteins and form the majority of drug targets in humans. How extracellular agonist binding triggers the activation of GPCRs and associated intracellular effector proteins remains, however, poorly understood. High resolution structural studies have recently revealed that inactive class-A GPCRs harbour a conserved binding site for Na$^+$ ions in the centre of their transmembrane domain, accessible from the extracellular space. Here, we show that the opening of a conserved hydrated channel in the activated state receptors allows the Na$^+$ ion to egress from its binding site into the cytosol. Coupled with protonation changes, this ion movement occurs without significant energy barriers, and can be driven by physiological transmembrane ion and voltage gradients. We propose that Na$^+$ ion exchange with the cytosol is a key step in GPCR activation, which locks receptors in long-lived active-state conformations.

INTRODUCTION

G-protein coupled receptors (GPCRs) mediate the transfer of external ligand binding information across the plasma membrane to activate a range of intracellular signaling pathways (Pierce, Premont, & Lefkowitz, 2002). Playing a central role in regulation of vital biological systems, including nervous, cardiovascular, immune, digestive, reproductive etc., they represent the majority of membrane proteins in humans and the largest class of present drug targets (Overington, Al-Lazikani, & Hopkins, 2006; Rask-Andersen, Masuram, & Schiöth, 2014). In recent years, a number of crystal structures have been solved to reveal conformational changes between inactive and active state receptors, including common movement in transmembrane helices and conserved microswitches (Katritch, Cherezov, & Stevens, 2013; Venkatakrishnan et al., 2013). However, despite this wealth of structural information, it is still not fully understood how ligand binding leads to activated receptors, which are able to trigger nucleotide exchange in intracellular effector G-protein complexes.

One of the major unknowns is the role of the highly conserved hydrophilic water-filled channel observed in crystal structures of class A GPCRs, which extends along the receptor axis from the external ligand binding region nearly all the way to the effector binding site. The channel is sealed towards the cytoplasm by a thin layer of hydrophobic residues in inactive state GPCRs (Fig 1A,B). Structures of high resolution, crystallized in the inactive conformation, reveal a Na^+ ion near the floor of this pocket, coordinated by water and three or four conserved residues including an acidic aspartate that is fully conserved in all ligand-sensing class A GPCRs (Christopher et al., 2013; Fenalti et al., 2014; Kruse et al., 2012; Liu et al., 2012; Miller-Gallacher et al., 2014; Pardo, Deupi, Dölker, López-Rodríguez, & Campillo, 2007; Zhang et al., 2012) ($D^{2.50}$; superscript refers to the Ballesteros and Weinstein residue numbering system) (Isberg et al., 2015). The allosteric effect of monovalent cations, in particular Na^+ ions, for GPCR function has been known for almost half a century (Pert & Synder, 1974), and the bulk of recent evidence shows that these effects are largely mediated by the ion binding at the $D^{2.50}$ site at the physiological concentration of Na^+ (140 mM and lower) (Fenalti et al., 2014; Liu et al., 2012; Massink et al., 2015). Due to the highly conserved nature of $D^{2.50}$ and other Na^+ ion coordinating residues, Na^+ ion binding at this site is likely to be a ubiquitous feature shared by the vast majority of class A GPCRs (Katritch et al., 2014).

In active receptor conformations, the ion binding site near $D^{2.50}$ shows a collapsed state,
which is likely not optimal for Na^+ ion binding (Huang et al., 2015; Kruse et al., 2013; Liu
et al., 2012; Rasmussen et al., 2011). It was therefore proposed that Na^+ ion leaves the
hydrophilic pocket upon receptor activation by a ligand or during receptor-G-protein
complex formation. However, how this movement is triggered and which pathway is
followed by the ion remains unknown.

Here, we investigated the link between ligand-induced receptor activation, the fate of the
bound Na^+ ion in class A GPCRs and its implications for transmembrane (TM) signal
transduction by equilibrium and non-equilibrium atomistic simulations on the M2
muscarinic receptor (m2r). When one addresses these questions, it is important to take
physiologically relevant electrochemical membrane conditions into consideration. Strong
TM Na^+ and K^+ gradients produce a sizable voltage across the plasma membrane of up
to -100 mV in the resting state of mammalian cells (Kandel, Schwartz, & Jessell, 2000).
Both the ionic gradients and electric field have been shown to influence the function of
GPCRs (Ben-Chaim et al., 2006; Navarro-Polanco et al., 2011; Rinne, Mobarec, Mahaut-
Smith, Kolb, & Bunemann, 2015) and are likely to impact the movement of the Na^+ ion
within the membrane region.

Our data reveal that the Na^+ ion observed in the TM domain of class A GPCRs can readily
traverse the receptor and, driven by the electrochemical gradients, migrate into the
cytoplasm in active receptor conformations. This result implies that a Na^+ ion may be
exchanged from the extracellular space to the cytoplasm as an important step in receptor
activation. Furthermore, the movement of Na^+ in the receptor, and intracellular egress, are
coupled to a protonation change of $D^{2.50}$.

RESULTS

GPCR activation opens a hydrated pathway across the receptor

We were first interested whether the conformational change from the inactive to active
receptor state renders the ion binding pocket sterically incapable of accommodating a Na^+
ion. The binding site for Na^+ appears to adopt a collapsed conformation in active crystal
structures. We started from an inactive state structure of the m2 muscarinic acetylcholine
receptor (m2r, PDB ID: 3UON) and, using a targeted molecular dynamics (MD) approach,
gently drove this conformation to the active state of this receptor (PDB ID: 4MQT) (Fig S1).

Our simulations show that the active state of m2r initially retains sufficient space for the ion.
The electrostatic attraction between the ion and the negatively charged side chain of $D69^{2.50}$
keeps the ion bound to this site during and after the transition from the inactive to the active
receptor conformation (Fig S2). However, our simulations show a widening of the
intracellular portion of the TM helices below the hydrophilic pocket during this
conformational change, which subsequently becomes fully hydrated (Fig 1B). The hydrated
pathway forms a connection between the orthosteric ligand-binding site, the hydrophilic
pocket and the G-protein binding site. The slim hydrophobic layer that delimits the
hydrophilic pocket towards the G-protein binding site in the inactive crystal structure
undergoes substantial conformational changes, which are especially evident from the
sidechain position of $Y440^{7.53}$. Our simulations show two major conformations of the
$Y440^{7.53}$ sidechain following the transition – an upward state similar to the conformation
observed in the inactive crystal structure (PDB: 3UON; Fig S3B) and a downward
configuration, which is also seen in the active crystal structure (PDB: 4MQT; Fig S3A). The
formation of a hydrated pathway connecting the receptor ligand and effector binding sites
has been reported in previous simulation studies on the $A_{2A}R$ and $5-HT_{1A}$ receptors (Yuan,
Filipek, Palczewski, & Vogel, 2014; Yuan, Peng, Palczewski, Vogel, & Filipek, 2016),
however the previous reports did not take the presence of a Na^+ ion into consideration.

Figure 1: Major structural features and internal hydration of class A GPCRs in the inactive and active state as shown by the m2r. (A) The main structural features of class A GPCRs, as exemplified by the m2r, include 7 TM helices (blue), an extracellular ligand binding site, the intracellular effector (G-protein) binding site as well as conserved and functionally important residues termed microswitches (selected ones are highlighted). The scale bar shown and all positions stated in the text use the Cα atom of D103$^{3.32}$ as reference. (B) Conformation of inactive m2r (PDB: 3UON) during the simulations showing the presence of the hydrophobic layer separating the hydrophilic pocket and effector binding site. (C) After transition to the active state (PDB: 4MQT), and further simulation, m2r displays a continuous water channel connecting the orthosteric ligand binding site, hydrophilic pocket and effector binding site. Water molecules are shown in red (surface representation), the position of the allosteric Na^{+} ion, as obtained from our initial simulations, is shown as a green sphere, and residues forming the hydrophobic layer (yellow) as well as the bound ligand (carbachol, light green) are depicted in stick representation.

The position of the internal Na$^+$ ion is coupled to protonation of D2.50

We were next interested in the interplay between the Na$^+$ ion and the key conserved titratable residue D69$^{2.50}$. A number of computational studies have explored functional implications of the protonation state of D$^{2.50}$, in particular its role in receptor activation, Na$^+$ ion binding, and interaction with the "ionic lock" motif (D$^{3.49}$R$^{3.50}$Y$^{3.51}$) in several class A family GPCRs (Miao, Caliman, & McCammon, 2015; Ranganathan, Dror, & Carlsson, 2014; Vanni, Neri, Tavernelli, & Rothlisberger, 2010). Here, we focused on a potential coupling between the position of the Na$^+$ ion within the receptor and protonation of D69$^{2.50}$. We carried out pK_a calculations on D69$^{2.50}$ using more than 800 equilibrated frames from simulations of the m2r receptor in a variety of conformations, including both the upward and downward configurations of the Y440$^{7.53}$ side chain. Due to the formation of a hydrated pathway across the receptor from the ligand to the effector binding sites in the active state simulations, we were able to evaluate the effect of the Na$^+$ ion positional changes on the D69$^{2.50}$ pK_a, where the Na$^+$ ion was shifted both in the upward (towards the extracellular face) and downward direction.

Figure 2 shows that the pK_a value and, thus, the most likely protonation state of D69$^{2.50}$ are strongly influenced by the Na$^+$ ion. If the cation is within ~3-5Å of D69$^{2.50}$, its positive charge strongly stabilises the negatively charged form of D69$^{2.50}$, leading to a pK_a value of ~3–4. However, displacement of the Na$^+$ ion to distances of 5Å and greater gives rise to a substantial pK_a shift to values between 8-12. This can be understood given the location of D69$^{2.50}$ in the middle of the transmembrane domain, surrounded by many non-polar residues. Transient movements of the internal Na$^+$ ion from its binding site, facilitated by activation-related conformational changes in the Na$^+$ pocket, can therefore be sufficient to lead to protonation of D69$^{2.50}$.

Figure 2: Proximity of the Na⁺ ion modulates protonation of D69$^{2.50}$.

Continuum electrostatics calculations of the pK_a of the D69$^{2.50}$ sidechain using a multitude of m2r
conformations obtained from our atomistic simulations in the carbachol-bound active state, both for
Y440$^{7.53}$ in the upward (left) and downward (right) conformations. The pK_a is shown as a function of
Z, the separation between the Na⁺ ion and the C$_\alpha$ atom of D103$^{3.32}$, which marks the orthosteric
ligand binding pocket, along the TM axis (see Fig 1A). The data points are in addition coloured
according to their distance to the D69$^{2.50}$ sidechain. The black continuous line, a smoothed spline fit,
indicates the approximate average pK_a for each separation for illustrative purposes, and the dashed
black line shows a pK_a of 7.

For the protonation of D$^{2.50}$, we propose that the most likely proton entry route would be

from the extracellular side, along the negative membrane potential gradient. Moreover, in

the m2r and other aminergic receptors the proton could be transferred from the conserved

D$^{3.32}$ in the orthosteric binding pocket via a short chain of water molecules (Isom &

Dohlman, 2015). In the apo state, our calculations in m2r indicate that D$^{3.32}$ is generally

protonated (pK_a = 11.2±1.7), whereas upon ligand binding the pK_a is substantially lowered

(pK_a = 7.6±1.9). A possible protonation change of D$^{3.32}$ could thus facilitate the shuttling of

protons to D$^{2.50}$. Furthermore, if a G-protein complex with a receptor is preformed before

agonist binding, D$^{2.50}$ would be readily accessible for protonation from the extracellular side

via a hydrated pathway. In this context, it has further been argued that bound agonists, but

not antagonists, may sustain the hydrated pathway past the ligand which connects the

extracellular space with the Na⁺ ion binding site upon receptor activation (Yuan et al.,

2016).

Simulations under electrochemical gradient show ion movement to the intracellular face

Next we conducted atomistic simulations with the Computational Electrophysiology

(CompEL) protocol (Kutzner et al., 2016) on the active conformation of m2r. We applied a

physiological Na^+ ion gradient of 150:10mM across the membrane from the extracellular to

the intracellular side, in addition to a small ion imbalance evoking a hyperpolarised V_m at -

250mV. Due to the wide range of pK_a values that $D69^{2.50}$ can adopt, its sidechain was

modeled both in charged and neutral forms.

Our simulations at -250 mV show that the Na^+ ion exhibits a substantial degree of mobility

even when $D69^{2.50}$ is in the charged state (Fig 3A, B, S7). The Na^+ ion is predominantly

coordinated by the residues $D69^{2.50}$, $S110^{3.39}$, $N435^{7.45}$ and $S433^{7.46}$. Under a small

membrane voltage, a bimodal distribution of distances between the ion and $D69^{2.50}$ is

observed, where larger distances of 5–6 Å are not uncommon (Fig S7C). As our pK_a

calculations showed that moderate excursions of the ion from its original binding site on

this scale likely have a major impact on the pK_a and protonation state of the $D69^{2.50}$

sidechain (Fig 2), we investigated the effect of a protonation change of $D69^{2.50}$ in the active

conformation.

Our simulations reveal that, in this receptor conformation, the Na^+ ion readily passes

through the hydrated channel into the intracellular solution. When $D69^{2.50}$ is neutral, we

observe the Na^+ ion to be expelled into the intracellular solution in three out of four

simulations at -250mV (Fig 3A, C, for complete list of trajectories see S4, 7A). At -500mV

the effect is, expectably, even more pronounced and movement into the cytoplasm is seen

in all the simulations we conducted (Fig 3B, D, for complete list of trajectories see S4, 7A).

In contrast, when $D69^{2.50}$ is charged, such a transition is observed in only in one out of

eight simulations at a raised membrane voltage (Fig 3B, S7B). The observed translocation

of Na^+ to the intracellular side occurs irrespective of the conformation adopted by $Y440^{7.53}$

(Fig 3C, D, S3).

In our simulations as well as under physiological conditions, both TM ion concentration

and voltage gradients drive ion flow across membrane pores. In the case of the Na^+ ion,

both gradients act synergistically in the resting state of the cell, driving the Na^+ ion towards

the cytoplasm. Under the conditions used in the simulations, fast ion motion through the

receptor is predominantly voltage-driven. Converted into an effective force, and using a

linear approximation to describe the gradient across the membrane (Dill & Bromberg,

2011), the influence of the concentration gradient would be about 10-fold smaller (~1.3

pN) than the driving force caused by the voltage gradient in these conditions (~13 pN).

223
224
225
Figure 3: Na$^+$ ion migration across the receptor to the intracellular side.
Z-coordinate of the Na$^+$ ion in the m2r under a hyperpolarised V_m of -250mV **(A)** and -500mV **(B)**.
Black lines denote the simulations with a charged D69$^{2.50}$; the purple, green and red lines display
simulations with a neutral D69$^{2.50}$. Here we show a selection of simulations, when D69$^{2.50}$ is neutral;
the trajectories show full passage of the ion to the intracellular side (for complete list of trajectories
see Fig S4, S7). Trajectories of the Na$^+$ ion moving from the hydrophilic pocket, accessible from the
extracellular space, into the intracellular bulk solution at **(C)** -250mV and **(D)** -500mV. The colour
of the Na$^+$ ion corresponds to panels A and B respectively. The Y440$^{7.53}$ upward and downward
conformations are shown in green. The pathways of the ion towards the intracellular side are almost
indistinguishable from each other until the ion passes Y440$^{7.53}$, thereafter the pathways diverge
somewhat due to the widened exit region to the cytoplasm.
237

Energetics of ion movement to the cytoplasm

As the initiation of fast movement of ions to the intracellular side was initially tested under slightly supra-physiological levels of V_m in our CompEL simulations of active state m2r, we next evaluated the detailed equilibrium energetics of the Na^+ ion movement on this pathway (i.e without applied gradients) to ascertain the physiological relevance of this transition. We calculated the potential-of-mean-force (PMF) for the migration of the cation in four different states: in addition to probing the influence of the $D69^{2.50}$ protonation state, we examined the role of the conformation of the $Y440^{7.53}$ sidechain, which substantially affects the width and overall shape of the formed hydrated pathway into the cytoplasm (Fig 3C,D).

When $D69^{2.50}$ is charged (Fig 4), the free energy difference between the internal the Na^+ ion binding site and the free intracellular bulk solution is ~30 kJ/mol. Accordingly, the active conformation of m2r retains a Na^+ ion at the allosteric site with relatively high affinity, as long as $D69^{2.50}$ remains unprotonated. The major barrier to migration into the cytoplasm is located near the $Y440^{7.53}$ sidechain. In its upward state, the free energy barrier amounts to ~41 kJ/mol, while it increases to ~48 kJ/mol in the downward state (Fig 4).

As our pK_a calculations showed that even a moderate displacement of the Na^+ ion away from its binding site at $D69^{2.50}$ is likely to lead to a protonation change of the aspartate, we also calculated the PMF of the Na^+ ion movement along the intracellular pathway for neutral $D69^{2.50}$. Importantly, this state no longer shows any affinity for the Na^+ ion, and ion movement into the intracellular bulk is not obstructed by any energy barrier significantly larger than the thermal energy (*kT,* ~2.5 kJ/mol) in the upward-oriented $Y440^{7.53}$ conformation. When $Y440^{7.53}$ is oriented downward, a small but readily surmountable energy barrier (on physiologically relevant timescales) of ~14 kJ/mol exists for this transition. The downward conformation of $Y440^{7.53}$ in conjunction with the neutral state of $D69^{2.50}$ also has a small influence on the shape and configuration of the ion binding site at $D69^{2.50}$, which leads to a reduction of the number of hydrogen bonds formed between the protein and the ion (Fig S5), raising the free energy of binding at this site further by ~7.5 kJ/mol (Fig 4A, B).

272
273
Figure 4: Energetics of Na⁺ translocation from the hydrophilic pocket to the intracellular side.
Equilibrium potential of mean force (PMF) profiles of the energetics of Na$^+$ translocation along the
Z-axis in m2r without any applied voltage or concentration gradients. Four relevant states were
considered: **(Left)** negatively charged D69$^{2.50}$ (black) or neutral D69$^{2.50}$ (red) with the Y440$^{7.53}$
sidechain in an upward conformation; **(Right)** negatively charged D69$^{2.50}$ (black) or neutral D69$^{2.50}$
(red) with a downward-oriented Y440$^{7.53}$ sidechain. The standard deviation of the PMF, obtained
from Bayesian bootstrap analysis, is depicted as shaded area. For each PMF, the intracellular bulk
solution was used as a reference, and the range of positions adopted by the Y440$^{7.53}$ sidechain is
denoted by blue dotted lines.

Conservation of the pocket and intracellular exit channel

Additional support for an important role of intracellular Na$^+$ egress in the activation of class
A GPCRs is provided by analysis of residue conservation along its exit pathway. As we
detailed previously (Katritch et al., 2014), there is a remarkable level of conservation for the
16 residues of the Na$^+$ binding pocket in class A GPCRs (Figure 5, Table S1), suggesting
conserved functional role of Na$^+$ in receptor activation mechanism. Interestingly, our
analysis of Na$^+$ contacts along the MD trajectories in this study shows that the residues
lining the ion exit path to the intracellular side are well conserved too. Thus, out of the 36
contact residues, 32 are 100% conserved among all five muscarinic receptors, 17 are >90%
conserved among all aminergic receptors, and 22 are consensus residues among all class A
GPCRs. Most importantly, the predicted exit pathway includes Na$^+$ contacts with the highly
conserved N$^{1.50}$ (100% and 98% conserved in aminergic and in all class A respectively),

$D^{3.49}$ (100% and 64%), Y5.58 (94% and 73%) and other residue positions generally

conserved as polar residues, including $N^{1.60}$, $T^{2.37}$ and $N^{2.39}$. Particularly, in the inactive M2

muscarinic receptor and in other inactive state GPCR structures as well, the $Y^{7.53}$ residue is

directed towards the Na^+ ion-binding pocket, and hence may play a role as first point of

polar contact outside the Na^+ ion-binding pocket for the intracellular movement of Na^+. The

Na^+ ion passage towards the cytosol may be further facilitated by other conserved polar

residues, including $D^{3.49}$, $N^{2.39}$, $N^{2.40}$ and $T^{2.37}$. Such conservation of the Na^+ ion pocket and

the path for intracellular egress of Na^+ suggests that Na^+ transfer described in this study can

occur in all muscarinic receptors and other class A GPCRs, comprising a key "irreversible"

part of the activation mechanism.

306
307
Figure 5: **Conservation of the intracellular Na^+ ion pathway.** Muscarinic M2 receptor shown in
blue cartoon representation, along with ball-and-stick representation of residues involved in the
egress of the Na^+ ion. The carbon atoms of 17 residues that are >90% conserved among aminergic
receptors are shown in green, the carbon atoms of additional 15 residues that are conserved among
the muscarinic family of receptors are shown in yellow, the carbon atoms of the 4 non-conserved
residues are shown in orange.
314

DISCUSSION

The principal role of GPCRs is to transmit information about an extracellular agonist binding event towards the cytoplasm, by catalysing GDP release from a bound intracellular G-protein complex (Pierce et al., 2002). This is known to involve conformational changes in the receptor, including conserved residue microswitches, and large scale movement of TM helices 6 and 7 in the intracellular side that open the nucleotide binding site of the $G\alpha$ protein (Dror et al., 2015; Huang et al., 2015; Mahoney & Sunahara, 2016). It has, furthermore, long been recognised that G-protein binding, and stabilisation of this conformation on the intracellular side of the receptor, increases agonist affinity on the extracellular face (DeVree et al., 2016; Maguire, Van Arsdale, & Gilman, 1976).

Na^+ ions, binding to an internal receptor site between the G-protein and the external ligand binding pockets, are known to act as powerful allosteric modulators of class A GPCR function (Katritch et al., 2014; Pert & Synder, 1974). Na^+ was found to selectively diminish the affinity of agonists, but not antagonists, to GPCR, which can be interpreted as a structural stabilisation of the inactive receptor state by the ions (Miller-Gallacher et al., 2014; Quitterer, AbdAlla, Jarnagin, & Müller-Esterl, 1996; Selley, Cao, Liu, & Childers, 2000). Accordingly, while receptor X-ray structures of sufficient resolution crystallised in the inactive state display a Na^+ ion bound to $D^{2.50}$, this binding site is collapsed in active receptor conformations, and ions are not observed (Huang et al., 2015; Katritch et al., 2014). Mutations around the Na^+ ion binding site have a major impact on receptor function in most class A GPCRs either completely abolishing G-protein activation, or resulting in constitutive ligand independent or pathway biased signaling (Fenalti et al., 2014; Liu et al., 2012; Massink et al., 2015).

Our work shows that the Na^+ ion binding pocket, which is accessible only from the extracellular face in the inactive state (Selent, Sanz, Pastor, & de Fabritiis, 2010; Vickery, Machtens, Tamburrino, Seeliger, & Zachariae, 2016), is transformed into a fully permeable, water-filled channel in the activated receptor conformation of m2r. This channel bridges the extracellular ligand and intracellular G-protein binding sites. Water access from the ligand binding site all the way to the cytoplasmic side of the receptor has previously also been observed in simulations on the $A_{2A}R$ and $5-HT_{1A}$ receptors (Yuan et al., 2014, 2016). We show here that the activated receptor state permits the Na^+ ion to cross the receptor towards the cytoplasmic side, without experiencing any major energy barriers. The high hydration level of this pathway in the active state is thereby an important factor in facilitating ion

passage. The correlation between hydration level and ion transfer has previously been
demonstrated in the case of ion channels (Beckstein et al., 2003; Dong, Fiorin, Carnevale,
Treptow, & Klein, 2013; Zhu & Hummer, 2012). In simulations of the inactive state, by
contrast, the application of substantially larger forces seems to be necessary to achieve
inward migration of Na^+, as no continuous hydrated channel is formed (Shang et al., 2014).

The inward motion of the Na^+ ion is facilitated by a protonation change of $D^{2.50}$ from the
negatively charged to the neutral form, which we show to occur even upon small
displacements of the ion from its equilibrium binding position. Neutralisation of $D^{2.50}$
substantially reduces its affinity for Na^+ ions. Migration of the ion towards the cytosol is
then driven by the negative membrane voltage and by a greater than 10-fold Na^+ gradient
across the cytoplasmic membrane under physiological conditions, both strongly attracting
Na^+ ions inwards. Indeed, we observe that moderately negative membrane voltages allow
fast escape of the allosteric Na^+ ion to the cytoplasm on 10–100 ns-timescales in our
simulations.

Figure 6: Proposed role of Na^+ translocation in GPCR activation. Key checkpoints during the
transition from the inactive (A) to active (D) state of the receptor. **(A)** The initial, inactive receptor
conformation shows no bound agonist or G-protein, and displays a Na^+ ion bound in a pocket which
is sealed towards the cytosol by a hydrophobic layer around $Y^{7.53}$. **(B)** G-protein and agonist bind to
the receptor (in undetermined order), leading to the formation of a continuous water channel across
the GPCR. The increased mobility of the Na^+ ion results in a pK_a shift and subsequent protonation of
$D^{2.50}$. **(C)** Neutralisation of $D^{2.50}$ and the presence of the hydrated pathway facilitate transfer of Na^+
to the intracellular side, which is driven by the transmembrane Na^+ gradient and the membrane
voltage. **(D)** The expulsion of Na^+ towards the cytosol results in a prolonged active state of the
receptor.

According to our results, conformational changes associated with agonist binding from the extracellular side and/or G-protein binding from the cytoplasm alter the Na^+ site conformation and the dynamics of the Na^+–$D^{2.50}$ salt bridge. This, in turn, leads to a protonation change of this residue, and subsequent egress of the Na^+ ion via a hydrated exit channel to the intracellular side. We therefore suggest that intracellular Na^+ ion transfer, facilitated by the membrane potential and Na^+ gradient, is a pivotal step during receptor activation, as it traps the receptor in the active state (Fig 6). It has been shown that, once activated, GPCRs remain in a prolonged active state, capable of signalling even when the receptors are internalized from the cytoplasmic membrane during endocytosis (Irannejad et al., 2013; Thomsen et al., 2016). The crucial role of the Na^+ ion movement within the receptor is reflected by the nearly complete conservation of the Na^+ ion binding site in class A GPCRs as well as the high conservation level of the exit pathway. The mechanism suggested here is also consistent with agonist independent basal signalling of GPCRs (Kobilka & Deupi, 2007), explaining this phenomenon as spontaneous protonation of $D^{2.50}$ and egress of the bound Na^+ ion on the intracellular pathway, leading to receptor activation.

Charge movements within membrane proteins, such as the coupled transfer of Na^+ ions and protons suggested by our MD simulations and pK_a calculations, should be sensitive to the membrane voltage. Indeed, it has been demonstrated that GPCR signalling is modulated by membrane voltage changes (Ben-Chaim et al., 2006; Mahaut-Smith, Martinez-Pinna, & Gurung, 2008; Martinez-Pinna, Tolhurst, Gurung, Vandenberg, & Mahaut-Smith, 2004; Moreno-Galindo, Alamilla, Sanchez-Chapula, Tristani-Firouzi, & Navarro-Polanco, 2016; Rinne et al., 2015; Vickery, Machtens, Tamburrino, et al., 2016). This applies both to the conformation of the receptors as well as their transmitted signal. Our findings are therefore consistent with these observations, as they suggest that movement of ions in the receptors constitute a key element in the receptor activation process. The observed voltage regulation of GPCRs is of particular relevance for receptors expressed in electrically excitable cells(Heifetz, James, Morao, Bodkin, & Biggin, 2016). In these cell types, the membrane voltage undergoes large-scale oscillations during action potentials. The transmitted receptor signal could thereby be tuned depending on the specific cell type and its excitation status (Vickery, Machtens, & Zachariae, 2016). Crucially, many GPCR drug targets are located in excitable tissue in the brain or muscle, where voltage regulation and a differential response to drugs may play an important role.

To summarise, our results suggest a model for class A GPCR activation, in which

conformational changes induced by G-protein and agonist binding are accompanied by the

intracellular transfer of an internally bound Na^+ ion. Importantly, these conformational

changes encompass rearrangement of the sidechain of $Y^{7.53}$, a conserved receptor

microswitch (Katritch et al., 2013), which in its upward state allows nearly barrier-free

intracellular permeation of Na^+ ions. This observation forms a functional link between the

major Na^+ binding site $D^{2.50}$ and $Y^{7.53}$ as the first polar point of contact on the intracellular

migration pathway of the Na^+ ion. Translocation of the ion is further facilitated by

protonation of the conserved $D^{2.50}$ residue (Fig 6), and driven by the physiological

membrane Na^+ and voltage gradients. The voltage sensitivity of GPCRs, which has been

previously reported for many receptors (Vickery, Machtens, & Zachariae, 2016), would thus

be a natural consequence of an activation mechanism which incorporates the movement of

ions as a key element. The Na^+ free receptors are likely to be trapped in an active state,

potentially explaining the prolonged mechanisms of signalling observed in many GPCRs.

METHODS

The simulation systems for the m2r were constructed using the crystal structures (PDB:

3UON, 4MQT)(Haga et al., 2012; Kruse et al., 2013). Ligands and non-GPCR subunits

were removed. The missing loop ICL3 was modelled using Modeller (v9.14)(Šali &

Blundell, 1993). For both simulation systems all internal water molecules and ions were

retained. The charged N- and C-termini were capped using acetyl and methyl moieties,

respectively. All ionisable groups were simulated with default protonation states, unless

otherwise mentioned. Each receptor was embedded into an equilibrated and hydrated 1,2-

palmitoyl-oleoyl-sn-glycero-3-phosphocholine (POPC) lipid bilayer using the GROMACS

utility g_membed (Wolf, Hoefling, Aponte-Santamaría, Grubmüller, & Groenhof, 2010)

resulting in a system size of ~92 x 88 x 97 Å. A concentration of 150 mM NaCl in the

aqueous solution was used for the single bilayer systems. During equilibration, all protein

heavy atoms were position-restrained with a force constant of 1000 kJ mol^{-1} nm^{-2} for 5-10

ns. Due to the low degree of internal hydration and medium resolution of the m2r structures

the equilibrations were extended by another 100 ns, now without position restraints, to

enable full hydration of the hydrophilic pocket. To study the active structure, the ligand

carbachol was parameterised using AMBER16, GAFF2, AM1-BCC parameters (Case et al.,

2016), and docked into the orthosteric ligand binding site using GOLD (v5.2.2). A Na^+ ion

was placed into the hydrophilic pocket in the inactive structure. We used a targeted MD

approach with the RMSD to the protein backbone of the active m2r crystal structure (PDB:

4MQT) as a reference, in order to gently enforce the transition from the inactive (PDB:

3UON) to the active state within ~250 ns. The two major conformations of $Y440^{7.53}$ we

observed during this simulation were probed systematically in the PMF calculations using

distance restraints between $N^{1.50}$-C_α and $D^{2.50}$-C_α to $Y^{7.53}$-C_ζ or dihedral restraints on the
sidechain of $Y^{7.53}$. To keep the G-protein binding site in an active conformation despite the
absence of bound G-protein, we applied a minimal set of four distance restraints to the C_α
atoms of the terminal groups of TM helices 2,5,6 and 7, namely residues 2.39-6.33, 2.39-
5.61, 2.43-7.54 and 6.36-7.54, at this interaction site (Fig S6).

For the CompEL simulations, the aforementioned active systems were duplicated along the
z axis to construct double bilayer systems. A NaCl gradient of 150mM:10mM between the
extracellular and intracellular compartments was used, along with an ion imbalance of 1 to 2
Cl^- ions to generate a V_m of $\sim$-250 to $\sim$ -500mV, as previously described (Kutzner,
Grubmüller, de Groot, & Zachariae, 2011). The V_m was determined by the GROMACS
utility gmx potential.

To calculate the PMF for the Na^+ ion within the hydrophilic pocket at neutral V_m, umbrella
sampling calculations were performed in bins of 0.25Å and analysed with the GROMACS
utility gmx wham. We used a simulation time of 50ns in each window and harmonic
potentials of 900–2000 kJ mol^{-1} nm^{-2} to restrain the Na^+ ion in the z-direction. The standard
deviation of the PMF profiles was estimated by using the Bayesian bootstrap method, as
implemented in gmx wham, with 200 runs. The free energy of the Na^+ ion in bulk solution
was set to 0. The position of the Na^+ ion (Z-coordinate) is reported relative to the $D103^{3.32}$-
C_α atom (ligand binding site).

For all simulations, the amber99sb_ildn force field was used for the protein (Lindorff-
Larsen et al., 2010), Berger parameters for lipids (Berger, Edholm, & Jähnig, 1997), which
were adapted for use with the amber99sb force field (Cordomí, Caltabiano, & Pardo, 2012),
and the SPC/E model for water molecules (Berendsen, Grigera, & Straatsma, 1987). Water
bond angles and distances were constrained by SETTLE (Miyamoto & Kollman, 1992)
while all other bonds were constrained using the LINCS method (Hess, Bekker, Berendsen,
& Fraaije, 1997). The temperature and pressure were kept constant throughout the
simulations at 310 K and 1 bar, respectively, with the protein, lipids, and water/ions coupled
individually to a temperature bath by the v-rescale method using a time constant of 0.2 ps
and a semi-isotropic Berendsen barostat (Berendsen, Postma, van Gunsteren, DiNola, &
Haak, 1984; Bussi, Donadio, & Parrinello, 2007). Employing a virtual site model for
hydrogen atoms (Feenstra, Hess, & Berendsen, 1999) allowed the use of 4-fs time steps
during the simulation. All simulations were performed with the GROMACS software
package, version 5.1.2

The pK_a calculations were performed using a continuum electrostatics method, namely the
Poisson-Boltzmann/Monte Carlo (PB/MC) approach, on multiple snapshots taken at a 2 ns
interval from different umbrella sampling simulations. PB calculations were performed
using MEAD (version 2.2.9)(Bashford & Gerwert, 1992) with a dielectric constant (ε_p) of 4
for the protein and 80 for the solvent (ε_w), in the presence of an explicit membrane. The
temperature was set to 310 K and the ionic strength to 0.145 M. The same temperature was
used for MC calculations (10^3 steps in each calculation), which were performed using

MCRP (Baptista, Martel, & Soares, 1999). Each MC step consisted of a cycle of
random choices of a state for all individual sites and pairs of sites with couplings above
2.0 pK_a units (Baptista et al., 1999), whose acceptance/rejection followed a Metropolis
criterion (Metropolis, Rosenbluth, Rosenbluth, Teller, & Teller, 1953); tautomeric
forms were not included.
495
496   The GROMACS software package, version 5.0.4 analysis toolkit was used to identify
497   residues with non-hydrogen heavy atoms within 4 Å of the sodium ion path during the
498   simulations. The residue conservation profile of the amino acids was obtained from the
499   GPCRdb server (Isberg et al., 2015).
500
501
502

ACKNOWLEDGEMENTS

This work was supported by the BBSRC (Training Grant BB/J013072/1 to U.Z.) and the
Scottish Universities' Physics Alliance (C.A.C, A.V.P. and U.Z.). This research was
partially supported by National Institute of Health grant DA035764 to V.K. We thank
Salomé Llabrés and Daniel Seeliger for fruitful discussions.

SUPPLEMENTAL INFORMATION
510
Figure S1: Backbone RMSD during a targeted MD simulation from the inactive to the active
state of m2r.
Figure S2: Na$^+$ position during a targeted MD simulation from the inactive to the active
state of m2r.
Figure S3: Y440$^{7.53}$ conformations used in pK_a and PMF calculations.
Figure S4: Backbone RMSD during MD simulations of the active state m2r under
membrane voltage.
Figure S5: Number of hydrogen bonds around the Na$^+$ binding site.
Figure S6: Depiction of the minimal set of distance restraints used to maintain the active
conformation of the m2r at the G-protein binding site.
Figure S7: Na$^+$ ion migration across the receptor to the intracellular side.
Table S1. Conservation of the residues in the transmembrane region that were observed to
be in contact (>4.5 Å) with Na$^+$ in the MD simulations. Overall, the sodium ion was
observed to come into close proximity with 34 residues.
525
526

527   **REFERENCES**
528

529   Baptista, A. M., Martel, P. J., & Soares, C. M. (1999). Simulation of Electron-Proton
530   Coupling with a Monte Carlo Method: Application to Cytochrome c3 Using
531   Continuum Electrostatics. *Biophysical Journal*, *76*(6), 2978–2998.
532   http://doi.org/10.1016/S0006-3495(99)77452-7
533   Bashford, D., & Gerwert, K. (1992). Electrostatic calculations of the pKa values of

ionizable groups in bacteriorhodopsin. *Journal of Molecular Biology*, *224*(2), 473–486. http://doi.org/10.1016/0022-2836(92)91009-E

Beckstein, O., Biggin, P. C., Bond, P., Bright, J. N., Domene, C., Grottesi, A., … Sansom, M. S. P. (2003). Ion channel gating: Insights via molecular simulations. *FEBS Letters*, *555*(1), 85–90. http://doi.org/10.1016/S0014-5793(03)01151-7

Ben-Chaim, Y., Chanda, B., Dascal, N., Bezanilla, F., Parnas, I., & Parnas, H. (2006). Movement of "gating charge" is coupled to ligand binding in a G-protein-coupled receptor. *Nature*, *444*(7115), 106–9. http://doi.org/10.1038/nature05259

Berendsen, H. J. C., Grigera, J. R., & Straatsma, T. P. (1987). The Missing Term in Effective Pair Potentials. *Journal of Physical Chemistry*, *91*(24), 6269–6271. http://doi.org/10.1021/j100308a038

Berendsen, H. J. C., Postma, J. P. M., van Gunsteren, W. F., DiNola, A., & Haak, J. R. (1984). Molecular dynamics with coupling to an external bath. *The Journal of Chemical Physics*, *81*, 3684–3690. http://doi.org/10.1063/1.448118

Berger, O., Edholm, O., & Jähnig, F. (1997). Molecular dynamics simulations of a fluid bilayer of dipalmitoylphosphatidylcholine at full hydration, constant pressure, and constant temperature. *Biophysical Journal*, *72*(May 1997), 2002–2013. http://doi.org/10.1016/S0006-3495(97)78845-3

Bussi, G., Donadio, D., & Parrinello, M. (2007). Canonical sampling through velocity rescaling. *Journal of Chemical Physics*, *126*. http://doi.org/10.1063/1.2408420

Case, D., Betz, R., Botello-Smith, W., Cerutti, D. S., Cheatham, T. E., Darden, T. A., … Kollman, P. A. (2016). Amber 2016. University of California, San Francisco.

Christopher, J. A., Brown, J., Doré, A. S., Errey, J. C., Koglin, M., Marshall, F. H., … Congreve, M. (2013). Biophysical Fragment Screening of the β 1 -Adrenergic Receptor: Identification of High Affinity Arylpiperazine Leads Using Structure-Based Drug Design. *Journal of Medicinal Chemistry*, *56*(9), 3446–3455. http://doi.org/10.1021/jm400140q

Cordomí, A., Caltabiano, G., & Pardo, L. (2012). Membrane protein simulations using AMBER force field and Berger lipid parameters. *Journal of Chemical Theory and Computation*, *8*, 948–958. http://doi.org/10.1021/ct200491c

DeVree, B. T., Mahoney, J. P., Vélez-Ruiz, G. A., Rasmussen, S. G. F., Kuszak, A. J., Edwald, E., … Sunahara, R. K. (2016). Allosteric coupling from G protein to the agonist-binding pocket in GPCRs. *Nature*, *535*(7610), 182–6. http://doi.org/10.1038/nature18324

Dill, K. A., & Bromberg, S. (2011). *Molecular Driving Force. Statistical Thermodynamics in Chemistry*. http://doi.org/10.1093/bib/4.4.382

Dong, H., Fiorin, G., Carnevale, V., Treptow, W., & Klein, M. L. (2013). Pore waters regulate ion permeation in a calcium release-activated calcium channel. *Proceedings of the National Academy of Sciences of the United States of America*, *110*, 17332–7. http://doi.org/10.1073/pnas.1316969110

Dror, R. O., Mildorf, T. J., Hilger, D., Manglik, A., Borhani, D. W., Arlow, D. H., … Shaw, D. E. (2015). Structural basis for nucleotide exchange in heterotrimeric G proteins. *Science*, *348*(6241), 1361–1365. http://doi.org/10.1126/science.aaa5264

Feenstra, K. A., Hess, B., & Berendsen, H. J. C. (1999). Improving efficiency of large time-scale molecular dynamics simulations of hydrogen-rich systems. *Journal of Computational Chemistry*, *20*(8), 786—798. http://doi.org/10.1002/(SICI)1096-987X(199906)20:8<786::AID-JCC5>3.0.CO;2-B

Fenalti, G., Giguere, P. M., Katritch, V., Huang, X.-P., Thompson, A. a, Cherezov, V., … Stevens, R. C. (2014). Molecular control of δ-opioid receptor signalling. *Nature*, *506*(7487), 191–196. http://doi.org/10.1038/nature12944

Haga, K., Kruse, A. C., Asada, H., Yurugi-Kobayashi, T., Shiroishi, M., Zhang, C., …
Kobayashi, T. (2012). Structure of the human M2 muscarinic acetylcholine receptor
bound to an antagonist. *Nature*, *482*(7386), 547–551.
http://doi.org/10.1038/nature10753

Heifetz, A., James, T., Morao, I., Bodkin, M. J., & Biggin, P. C. (2016). Guiding lead
optimization with GPCR structure modeling and molecular dynamics. *Current Opinion
in Pharmacology*, *30*, 14–21. http://doi.org/10.1016/j.coph.2016.06.004

Hess, B., Bekker, H., Berendsen, H. J. C., & Fraaije, J. G. E. M. (1997). LINCS: A linear
constraint solver for molecular simulations. *Journal of Computational Chemistry*,
18(12), 1463–1472. http://doi.org/10.1002/(SICI)1096-
987X(199709)18:12<1463::AID-JCC4>3.0.CO;2-H

Huang, W., Manglik, A., Venkatakrishnan, A. J., Laeremans, T., Feinberg, E. N., Sanborn,
A. L., … Kobilka, B. K. (2015). Structural insights into μ-opioid receptor activation.
Nature, *524*(7565), 315–321. http://doi.org/10.1038/nature14886

Irannejad, R., Tomshine, J. C., Tomshine, J. R., Chevalier, M., Mahoney, J. P., Steyaert, J.,
… von Zastrow, M. (2013). Conformational biosensors reveal GPCR signalling from
endosomes. *Nature*, *495*(7442), 534–538. http://doi.org/10.1038/nature12000

Isberg, V., de Graaf, C., Bortolato, A., Cherezov, V., Katritch, V., Marshall, F. H., …
Gloriam, D. E. (2015). Generic GPCR residue numbers – aligning topology maps while
minding the gaps. *Trends in Pharmacological Sciences*, *36*(1), 22–31.
http://doi.org/10.1016/j.tips.2014.11.001

Isom, D. G., & Dohlman, H. G. (2015). Buried ionizable networks are an ancient hallmark
of G protein-coupled receptor activation. *Proceedings of the National Academy of
Sciences*, *2015*, 201417888. http://doi.org/10.1073/pnas.1417888112

Kandel, E. R., Schwartz, J. H., & Jessell, T. M. (2000). *Principles of Neural Science*.
Mcgraw-Hill Publ.Comp. (Vol. 3). http://doi.org/10.1036/0838577016

Katritch, V., Cherezov, V., & Stevens, R. C. (2013). Structure-function of the G protein-
coupled receptor superfamily. *Annual Review of Pharmacology and Toxicology*, *53*,
531–56. http://doi.org/10.1146/annurev-pharmtox-032112-135923

Katritch, V., Fenalti, G., Abola, E. E., Roth, B. L., Cherezov, V., & Stevens, R. C. (2014,
May). Allosteric sodium in class A GPCR signaling. *Trends in Biochemical Sciences*.
Elsevier Ltd. http://doi.org/10.1016/j.tibs.2014.03.002

Kobilka, B. K., & Deupi, X. (2007). Conformational complexity of G-protein-coupled
receptors. *Trends in Pharmacological Sciences*, *28*(8), 397–406.
http://doi.org/10.1016/j.tips.2007.06.003

Kruse, A. C., Hu, J., Pan, A. C., Arlow, D. H., Rosenbaum, D. M., Rosemond, E., …
Kobilka, B. K. (2012). Structure and dynamics of the M3 muscarinic acetylcholine
receptor. *Nature*. http://doi.org/10.1038/nature10867

Kruse, A. C., Ring, A. M., Manglik, A., Hu, J., Hu, K., Eitel, K., … Kobilka, B. K. (2013).
Activation and allosteric modulation of a muscarinic acetylcholine receptor. *Nature*,
504(7478), 101–106. http://doi.org/10.1038/nature12735

Kutzner, C., Grubmüller, H., de Groot, B. L., & Zachariae, U. (2011). Computational
electrophysiology: the molecular dynamics of ion channel permeation and selectivity in
atomistic detail. *Biophysical Journal*, *101*(4), 809–17.
http://doi.org/10.1016/j.bpj.2011.06.010

Kutzner, C., Köpfer, D. A., Machtens, J., Groot, B. L. De, Song, C., & Zachariae, U. (2016).
Biochimica et Biophysica Acta Insights into the function of ion channels by
computational electrophysiology simulations. *BBA - Biomembranes*.
http://doi.org/10.1016/j.bbamem.2016.02.006

Lindorff-Larsen, K., Piana, S., Palmo, K., Maragakis, P., Klepeis, J. L., Dror, R. O., &

Shaw, D. E. (2010). Improved side-chain torsion potentials for the Amber ff99SB
protein force field. *Proteins*, *78*(8), 1950–8. http://doi.org/10.1002/prot.22711

Liu, W., Chun, E., Thompson, A. a, Chubukov, P., Xu, F., Katritch, V., … Stevens, R. C.
(2012). Structural Basis for Allosteric Regulation of GPCRs by Sodium Ions. *Science*,
337(6091), 232–236. http://doi.org/10.1126/science.1219218

Maguire, M. E., Van Arsdale, P. M., & Gilman, a G. (1976). An agonist-specific effect of
guanine nucleotides on binding to the beta adrenergic receptor. *Molecular*
Pharmacology, *12*, 335–339.

Mahaut-Smith, M. P., Martinez-Pinna, J., & Gurung, I. S. (2008, August). A role for
membrane potential in regulating GPCRs? *Trends in Pharmacological Sciences*.
http://doi.org/10.1016/j.tips.2008.05.007

Mahoney, J. P., & Sunahara, R. K. (2016). Mechanistic insights into GPCR-G protein
interactions. *Current Opinion in Structural Biology*, *41*, 247–254.
http://doi.org/10.1016/j.sbi.2016.11.005

Martinez-Pinna, J., Tolhurst, G., Gurung, I. S., Vandenberg, J. I., & Mahaut-Smith, M. P.
(2004). Sensitivity limits for voltage control of P2Y receptor-evoked Ca^{2+} mobilization
in the rat megakaryocyte. *The Journal of Physiology*, *555*(Pt 1), 61–70.
http://doi.org/10.1113/jphysiol.2003.056846

Massink, A., Gutierrez-de-Teran, H., Lenselink, E. B., Ortiz Zacarias, N. V., Xia, L.,
Heitman, L. H., … IJzerman, A. P. (2015). Sodium Ion Binding Pocket Mutations and
Adenosine A2A Receptor Function. *Molecular Pharmacology*, *87*(2), 305–313.
http://doi.org/10.1124/mol.114.095737

Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H., & Teller, E. (1953).
Equation of state calculations by fast computing machines. *Journal Chemical Physics*.
http://doi.org/http://dx.doi.org/10.1063/1.1699114

Miao, Y., Caliman, A. D., & McCammon, J. A. (2015). Allosteric Effects of Sodium Ion
Binding on Activation of the M3 Muscarinic G-Protein-Coupled Receptor. *Biophysical*
Journal, *108*(7), 1796–1806. http://doi.org/10.1016/j.bpj.2015.03.003

Miller-Gallacher, J. L., Nehmé, R., Warne, T., Edwards, P. C., Schertler, G. F. X., Leslie, A.
G. W., & Tate, C. G. (2014). The 2.1 Å Resolution Structure of Cyanopindolol-Bound
β1-Adrenoceptor Identifies an Intramembrane Na^+ Ion that Stabilises the Ligand-Free
Receptor. *PloS One*, *9*(3), e92727. http://doi.org/10.1371/journal.pone.0092727

Miyamoto, S., & Kollman, P. A. (1992). SETTLE: an analytical version of the SHAKE and
RATTLE algorithm for rigid water models. *Journal of Computational Chemistry*, *13*,
952–962. http://doi.org/10.1002/jcc.540130805

Moreno-Galindo, E. G., Alamilla, J., Sanchez-Chapula, J. A., Tristani-Firouzi, M., &
Navarro-Polanco, R. A. (2016). The agonist-specific voltage dependence of M2
muscarinic receptors modulates the deactivation of the acetylcholine-gated K+ current
(I KACh). *Pflügers Archiv - European Journal of Physiology*.
http://doi.org/10.1007/s00424-016-1812-y

Navarro-Polanco, R. a, Moreno Galindo, E. G., Ferrer-Villada, T., Arias, M., Rigby, J. R.,
Sánchez-Chapula, J. a, & Tristani-Firouzi, M. (2011). Conformational changes in the
M2 muscarinic receptor induced by membrane voltage and agonist binding. *The*
Journal of Physiology, *589*(Pt 7), 1741–1753.
http://doi.org/10.1113/jphysiol.2010.204107

Overington, J. P., Al-Lazikani, B., & Hopkins, A. L. (2006). How many drug targets are
there? *Nature Reviews. Drug Discovery*, *5*(12), 993–6. http://doi.org/10.1038/nrd2199

Pardo, L., Deupi, X., Dölker, N., López-Rodríguez, M. L., & Campillo, M. (2007). The role
of internal water molecules in the structure and function of the rhodopsin family of G
protein-coupled receptors. *ChemBioChem*. http://doi.org/10.1002/cbic.200600429

Pert, C. B., & Synder, S. H. (1974). Opiate Receptor Binding of Agonists and Antagonists
Affected Differentially by Sodium. *Molecular Pharmacology*, *10*(6), 868–879.
Pierce, K. L., Premont, R. T., & Lefkowitz, R. J. (2002). Seven-transmembrane receptors.
Nature Reviews. Molecular Cell Biology, *3*(September), 639–650.
http://doi.org/10.1038/nrm908
Quitterer, U., AbdAlla, S., Jarnagin, K., & Müller-Esterl, W. (1996). Na+ ions binding to
the bradykinin B2 receptor suppress agonist-independent receptor activation.
Biochemistry, *35*(41), 13368–77. http://doi.org/10.1021/bi961163w
Ranganathan, A., Dror, R. O., & Carlsson, J. (2014). Insights into the Role of Asp79$^{2.50}$ in
β2 Adrenergic Receptor Activation from Molecular Dynamics Simulations.
Biochemistry, *53*(46), 7283–7296. http://doi.org/10.1021/bi5008723
Rask-Andersen, M., Masuram, S., & Schiöth, H. B. (2014). The Druggable Genome:
Evaluation of Drug Targets in Clinical Trials Suggests Major Shifts in Molecular Class
and Indication. *Annual Review of Pharmacology and Toxicology*, *54*(1), 9–26.
http://doi.org/10.1146/annurev-pharmtox-011613-135943
Rasmussen, S. G. F., DeVree, B. T., Zou, Y., Kruse, A. C., Chung, K. Y., Kobilka, T. S., …
Kobilka, B. K. (2011). Crystal structure of the β2 adrenergic receptor-Gs protein
complex. *Nature*, *477*(7366), 549–55. http://doi.org/10.1038/nature10361
Rinne, A., Mobarec, J. C., Mahaut-Smith, M., Kolb, P., & Bunemann, M. (2015). The mode
of agonist binding to a G protein-coupled receptor switches the effect that voltage
changes have on signaling. *Science Signaling*, *8*(401), ra110-ra110.
http://doi.org/10.1126/scisignal.aac7419
Šali, A., & Blundell, T. L. (1993). Comparative Protein Modelling by Satisfaction of Spatial
Restraints. *Journal of Molecular Biology*, *234*(3), 779–815.
http://doi.org/10.1006/jmbi.1993.1626
Selent, J., Sanz, F., Pastor, M., & de Fabritiis, G. (2010). Induced effects of sodium ions on
dopaminergic G-protein coupled receptors. *PLoS Computational Biology*, *6*(8).
http://doi.org/10.1371/journal.pcbi.1000884
Selley, D. E., Cao, C. C., Liu, Q., & Childers, S. R. (2000). Effects of sodium on agonist
efficacy for G-protein activation in mu-opioid receptor-transfected CHO cells and rat
thalamus. *British Journal of Pharmacology*, *130*(5), 987–996.
http://doi.org/10.1038/sj.bjp.0703382
Shang, Y., Lerouzic, V., Schneider, S., Bisignano, P., Pasternak, G. W., & Filizola, M.
(2014). Mechanistic insights into the allosteric modulation of opioid receptors by
sodium ions. *Biochemistry*, *53*(31), 5140–9. http://doi.org/10.1021/bi5006915
Thomsen, A. R. B., Plouffe, B., Cahill, T. J., Shukla, A. K., Tarrasch, J. T., Dosey, A. M.,
… Lefkowitz, R. J. (2016). GPCR-G Protein-B-Arrestin Super-Complex Mediates
Sustained G Protein Signaling. *Cell*, *166*(4), 907–919.
http://doi.org/10.1016/j.cell.2016.07.004
Vanni, S., Neri, M., Tavernelli, I., & Rothlisberger, U. (2010). A Conserved Protonation-
Induced Switch can Trigger "Ionic-Lock" Formation in Adrenergic Receptors. *Journal*
of Molecular Biology, *397*(5), 1339–1349. http://doi.org/10.1016/j.jmb.2010.01.060
Venkatakrishnan, a J., Deupi, X., Lebon, G., Tate, C. G., Schertler, G. F., & Babu, M. M.
(2013). Molecular signatures of G-protein-coupled receptors. *Nature*, *494*(7436), 185–
94. http://doi.org/10.1038/nature11896
Vickery, O. N., Machtens, J.-P., Tamburrino, G., Seeliger, D., & Zachariae, U. (2016).
Structural Mechanisms of Voltage Sensing in G Protein-Coupled Receptors. *Structure*,
24(6), 997–1007. http://doi.org/10.1016/j.str.2016.04.007
Vickery, O. N., Machtens, J.-P., & Zachariae, U. (2016). Membrane potentials regulating
GPCRs: insights from experiments and molecular dynamics simulations. *Current*

734   *Opinion in Pharmacology*, *30*, 44–50. http://doi.org/10.1016/j.coph.2016.06.011

735 Wolf, M. G., Hoefling, M., Aponte-Santamaría, C., Grubmüller, H., & Groenhof, G. (2010).

736   G-membed: Efficient insertion of a membrane protein into an equilibrated lipid bilayer

737   with minimal perturbation. *Journal of Computational Chemistry*, *31*, 2169–2174.

738   http://doi.org/10.1002/jcc.21507

739 Yuan, S., Filipek, S., Palczewski, K., & Vogel, H. (2014). Activation of G-protein-coupled

740   receptors correlates with the formation of a continuous internal water pathway. *Nature*

741   *Communications*, *5*(May), 4733. http://doi.org/10.1038/ncomms5733

742 Yuan, S., Peng, Q., Palczewski, K., Vogel, H., & Filipek, S. (2016). Mechanistic Studies on

743   the Stereoselectivity of the Serotonin 5-HT 1A Receptor. *Angewandte Chemie*

744   *International Edition*, *55*(30), 8661–8665. http://doi.org/10.1002/anie.201603766

745 Zhang, C., Srinivasan, Y., Arlow, D. H., Fung, J. J., Palmer, D., Zheng, Y., … Kobilka, B.

746   K. (2012). High-resolution crystal structure of human protease-activated receptor 1.

747   *Nature*, *492*(7429), 387–92. http://doi.org/10.1038/nature11701

748 Zhu, F., & Hummer, G. (2012). Drying transition in the hydrophobic gate of the GLIC

749   channel blocks ion conduction. *Biophysical Journal*, *103*(2), 219–227.

750   http://doi.org/10.1016/j.bpj.2012.06.003