Atlantean Evolution in Darwin's Finches—Issues and Perspectives in Species Delimitation using

Phenotypic Data

CARLOS DANIEL CADENA[1*], IVÁN JIMÉNEZ[2], AND FELIPE ZAPATA[3]

[1]*Departamento de Ciencias Biológicas, Universidad de los Andes, Bogotá, Colombia; [2]Center for Conservation and*

Sustainable Development, Missouri Botanical Garden, St. Louis, MO, USA; [3]Department of Ecology and

Evolutionary Biology, University of California, Los Angeles, CA, USA.

Correspondence to be sent to: Departamento de Ciencias Biológicas, Universidad de los Andes, Carrera 1 no. 18A-

10, Bogotá, Colombia; E-mail: ccadena@uniandes.edu.co

Abstract.— Progress in the development and use of methods for species delimitation employing

phenotypic data lags behind conceptual and practical advances in molecular genetic approaches. The basic

evolutionary model underlying the use of phenotypic data to delimit species assumes random mating and

quantitative polygenic traits, so that phenotypic distributions within a species should be approximately

normal for individuals of the same sex and age. Accordingly, two or more distinct normal distributions of

phenotypic traits suggest the existence of multiple species. In light of this model, we show that analytical

approaches employed in taxonomic studies using phenotypic data are often compromised by three issues:

(1) reliance on graphical analyses of phenotypic space that do not consider the frequency of phenotypes;

(2) exclusion of characters potentially important for species delimitation following reduction of data

dimensionality; and (3) use of measures of central tendencies to evaluate phenotypic distinctiveness. We

outline approaches to overcome these issues based on statistical developments related to normal mixture

models and illustrate them empirically with a reanalysis of morphological data recently used to claim that

there are no morphologically distinct species of Darwin's ground-finches (*Geospiza*). We found negligible

support for this claim relative to taxonomic hypotheses recognizing multiple species. Although species

limits among ground-finches merit further assessments using additional sources of information, our results

bear implications for other areas of inquiry including speciation research: because ground-finches have

likely speciated and are not trapped in a process of "Sisyphean" evolution as recently argued, they remain

useful models to understand the evolutionary forces involved in speciation. Our work underscores the

importance of statistical approaches grounded on appropriate evolutionary models for species

delimitation. Approaches allowing one to fit normal mixture models without *a priori* information about

species limits offer new perspectives in the kind of inferences available to systematists, with significant

repercusions on ideas about the structure of biological biodiversity. [morphology; normal mixture model;

phenotype; principal components analysis; species limits; variable selection.]

Systematic biology seeks to discover and describe species, and to establish phylogenetic relationships

among them and among clades at higher levels. Given these two main goals of the field, reviews published

over a decade ago noted that the literature on theory and methods of phylogenetic inference and on theory

of species concepts was extensive, whereas methods for delimiting species had received much less

attention (Sites and Marshall 2003; Sites and Marshall 2004). Over the past few years, this imbalance has

been partly overcome with considerable development, application, and integration of methods for species

delimitation (Padial et al. 2010; Camargo and Sites 2013). Largely driven by increased availability of

multilocus datasets brought about by advances in DNA sequencing technology, however, much recent

progress has focused on probabilistic methods for analyses of molecular data (reviewed by Fujita et al.

2012; Carstens et al. 2013), whereas relatively little effort has been devoted to approaches using

phenotypic data to delimit species (Wiens and Servedio 2000; Ezard et al. 2010; Guillot et al. 2012;

Zapata and Jiménez 2012; Edwards and Knowles 2014; Solís-Lemus et al. 2014). Yet, because most fossil

and living species have been discovered and named based on phenotypic distinctiveness (Luckow 1995;

Mallet 2013; Miller 2016), and because genomic-based species delimitation approaches are no substitutes

for judicious assessments of other sources of information (Sukumaran and Knowles 2017), the theory and

practice of delimiting species using phenotypic data remain central to modern systematics.

Although species descriptions employing phenotypic data are often non-quantitative and although

systematists may often not be explicit about the rationale they follow to delimit species (Luckow 1995;

McDade 1995; Sangster 2014; Allmon 2016), the use of objective criteria for species diagnosis based on

phenotypic characters has a long tradition in taxonomy, rooted on evolutionary theory (Wiens and

Servedio 2000; Zapata and Jiménez 2012; Futuyma 2013). The basic evolutionary model for the

distribution of a continuous quantitative character within a species (Fisher 1918) assumes polygenic

inheritance and random mating; under these assumptions, gene frequencies are expected to be close to

Hardy-Weinberg equilibrium and phenotypic variation among individuals of a single species tends to be

normally distributed (Templeton 2006). On the other hand, if phenotypic variation is best described by

two or more distinct normal distributions, then one may conclude that there is more than one species in a

sample of individuals (Coyne and Orr 2004; Mallet 2008). This conclusion is granted under the

assumption that distinct normal distributions in polygenic traits do not reflect age- or sex-related variation,

or phenotypic plasticity. It follows that distinct normal distributions in cases in which phenotypic variation

is caused by few loci of large effect (e.g., Smith 1993) or largely driven by environmental factors (e.g.,

Moczek and Emlen 1999) do not constitute evidence of more than one species. Finally, because distinct

phenotypic distributions may represent evidence of species boundaries given a variety of criteria for

species delimitation (Luckow 1995; Zapata and Jiménez 2012), the Fisherian model described above

serves as a conceptual basis to establish species limits under multiple species definitions (sensu de

Queiroz 1998).

Despite the long tradition of this basic model for species delimitation based on quantitative phenotypic

characters, statistical tools for its formal application to empirical data were limited until recently.

Procedures allowing one to fit combinations of normal distributions to phenotypic variation among

specimens, without *a priori* knowledge of species limits, were initially developed in the late XIX century

(Pearson 1894). However, practical application only became possible following computational advances

in the 1970s (i.e., the expectation–maximization algorithm; McLachlan and Peel 2000) and software

development from the late XX century into the present (Fraley and Raftery 2002; Fraley et al. 2012).

Because these statistical approaches entered the literature on species delimitation only a few years ago

(Ezard et al. 2010), it is not surprising that even recent studies do not employ them when analyzing

phenotypic data to delimit species. Instead, systematists frequently infer species limits examining

phenotypic variation based on visual inspection of scatter plots defined by a few axes that account for

most phenotypic variance, often derived from principal components analysis (PCA). In addition,

systematists often delimit species based on differences between groups of specimens in the central

tendency of phenotypes. This is true of work on living plants and animals (reviewed by Rieseberg et al.

2006), as well as in studies of extinct taxa in the fossil record (reviewed by Allmon 2016).

Here, we show that analytical approaches commonly employed in taxonomic studies are inadequate in

light of the evolutionary model underlying species delimitation described above. It follows that if species

delimited by inadequate statistical approaches are used as units for subsequent analyses, then any mistakes

may carry on and influence views in other areas of inquiry, such as speciation research. Focusing on

Darwin's finches from the Galapagos Islands, an iconic group for the study of natural selection,

speciation, and adaptive radiation (Lack 1947; Bowman 1961; Grant 1999; Grant and Grant 2008; Grant

and Grant 2014), we provide an example of how employing statistical approaches explicitly related to the

basic evolutionary model underlying the use of phenotypic data in species delimitation may enhance

assessments of species limits and thus our understanding of evolutionary processes.

96   SISYPHEAN EVOLUTION IN DARWIN'S FINCHES?

Among Darwin's finches, the many studies of ground-finches in the genus *Geospiza* have been especially

productive in terms of insights into species formation and the role of geographic isolation, natural

selection, and hybridization in microevolutionary processes that may scale up to macroevolutionary

patterns (reviewed by Grant 1999; Grant and Grant 2008; Grant and Grant 2014). There has been

considerable disagreement in the literature about the number of species in the group (reviewed by McKay

and Zink 2015), but most modern taxonomic treatments have recognized six species of ground-finches

(Lack 1947; Rising et al. 2011). However, based on genomic evidence (Lamichhaney et al. 2015) and

some vocal and behavioral data, three subspecies were recently elevated to species rank, bringing the total

number of recognized species to nine (Remsen et al. 2017).

In a provocative recent paper, however, McKay and Zink (2015) offered an intriguing alternative

perspective on the taxonomy and evolution of ground-finches (see also Zink 2002). These authors boldly

argued that morphological evidence for the existence of multiple species of *Geospiza* is lacking, and they

presented the iconoclastic argument that different phenotypes should be considered transient ecomorphs

within a single species. Furthermore, according to these authors, ground-finches are an appropriate model

to study forces involved in geographic variation and local adaptation, but not to demonstrate the workings

of speciation because in their view speciation in the group has not occurred. Instead, incipient speciation

has been repeatedly stalled or reversed owing to shifting conditions affecting the strength and direction of

natural selection and to ongoing gene flow, a situation they wittily referred to as "Sisyphean" evolution

(McKay and Zink 2015). Because of its originality in challenging "entrenched orthodoxy regarding

speciation in Darwin's Finches", the study by McKay and Zink (2015) was duly recognized with an award

by a major ornithological organization (Cooper Ornithological Society 2016).

A central premise of the arguments by McKay and Zink (2015) was their assertion that phenotypic

discontinuities do not exist among recognized species of ground-finches (contra Lack 1947; Grant et al.

1985). Although they rightly noted that "the real test of species limits is determining the extent to which

specimens form multiple morphological clusters when *a priori* specimen identifications are ignored",

McKay and Zink (2015) did not formally conduct such a test. Instead, their approach illustrates three

problematic issues in analyses of phenotypic data for species delimitation. In the next section we describe

these issues and outline possible solutions afforded by statistical tools directly related to the basic

evolutionary model underlying the use of phenotypic data in species delimitation. We then implement

these solutions in a reanalysis of the morphological data on *Geospiza* ground-finches to revisit the

question of whether morphological evidence supports the hypothesis that there are several species in the

group.

THREE FREQUENT ISSUES IN ANALYSES OF PHENOTYPIC DATA FOR SPECIES DELIMITATION

1) *Graphical analyses may convey little information on phenotype frequencies crucial to assess evidence*

for multiple species.

Many species delimitation studies rely on visual inspection of bivariate (rarely trivariate) scatter plots of

phenotypic space to detect discontinuities and thus define phenotypic groups (e.g., Fig. 1 in McKay and

Zink 2015). These scatter plots may offer only limited insight into the structure of character variation

because visual cluttering and record overplotting hinder perception of phenotype frequencies crucial to

identify groups (McLachlan 2004). We illustrate this problem with a hypothetical example in which

specimens from a given locality seem to reveal no phenotypic discontinuities, with intermediate

phenotypes across the range of variation (Fig. 1a); accordingly, a univariate scatter plot fails to reveal

evidence of distinct phenotypic groups (Fig. 1b). The problem with scatter plots concealing crucial

information (also common in two- and three-dimensional scatter plots) is revealed by a histogram of

phenotype frequencies employing the same data, which reveals two distinct normal distributions (Fig. 1c).

Following the model for species delimitation based on continuous phenotypic characters described above,

this histogram suggests the existence of two species.

Graphical analysis of phenotype frequencies (e.g., Fig. 1c) may be effective to detect groups when few

characters are relevant (but see McLachlan and Peel 2000, page 9). However, it may be difficult to detect

distinct normal distributions in phenotypic spaces defined by more than two dimensions, where complex

covariance structures are likely (McLachlan 2004). Moreover, detection of phenotypic groups exclusively

based on graphical analysis is potentially highly subjective and difficult to replicate, or, as stated by Karl

Pearson (1894) over a century ago: "To throw the solution on the judgment of the eye in examining the

graphical results is, I feel certain, quite futile". Therefore, graphical analysis of phenotype frequencies is a

useful but limited tool for species delimitation.

Recent statistical developments allow systematists to go beyond graphical analysis by using normal

mixture models (NMMs, McLachlan and Peel 2000) as a formal approach to test for the existence of

distinct species based on multivariate phenotypic data (Ezard et al. 2010; Guillot et al. 2012; Edwards and

Knowles 2014; Kleindorfer et al. 2014). These models conceptualize phenotypic variation as a

combination (i.e., a mixture) of distinct normal distributions; a mixture may include one or more distinct

normal distributions, representing the hypothesis of one or more species, respectively. The parameters of a

NMM specifying a particular hypothesis include the means and variance-covariance matrices describing

the Gaussian phenotypic distribution of each species. These parameters can be estimated using maximum

likelihood from data on phenotypic measurements, without *a priori* knowledge of species limits,

employing the expectation–maximization algorithm (McLachlan and Krishnan 2008). Comparison of

empirical support among models representing different hypotheses is often based on the Bayesian

Information Criterion (BIC; Schwarz 1978), which evaluates the likelihood of each model while adjusting

for model complexity (Fraley and Raftery 2002).

2) *Reduction of dimensionality via PCA may exclude important characters for species delimitation*.

Species delimitation studies often begin analyses by reducing the dimensionality of phenotypic space,

typically via principal component analysis (PCA) or related procedures (McLachlan 2004; Ezard et al.

2010), and then focus attention on few principal components accounting for most of the variation in the

data. For example, McKay and Zink (2015) focused on three principal components explaining 99% of the

variation in six morphological characters of *Geospiza* ground-finches (see their Fig. 1). This use of PCA

and related procedures in taxonomy was suggested decades ago (Sneath and Sokal 1973) and is still

prescribed nowadays (e.g., Ezard et al. 2010). However, there is no reason to believe that principal

components accounting for most of the variation in a dataset are most useful for group discrimination

(Chang 1983).

To illustrate the problem of reducing dimensionality to the principal components accounting for most of

the variation, we use a hypothetical example based on two phenotypically distinct species, each

represented by a bivariate normal distribution (Fig. 2a). The first principal component of the mixture of

these two distributions explains >99% of the variation and, yet, it is useless to distinguish the two species

(Fig. 2b). In contrast, the second principal component accounts for <1% of the variation and perfectly

discriminates species (Fig. 2c). This example is bivariate for simplicity, but the statistical principle applies

to mixtures of two normal distributions in any number of dimensions (Chang 1983). We stress that the

problem at hand is not rotation of the data using PCA or related procedures, because such rotation may

serve a number of useful purposes; rather, the problem is employing the amount of phenotypic variance

explained by each principal component as a criterion to judge its usefulness to distinguish phenotypic

groups (Chang 1983).

Although alternatives to PCA and related approaches for dimensionality reduction should be regularly

considered in analysis aiming to detect groups in multivariate space (McLachlan and Peel 2000;

McLachlan 2004), they are rarely implemented in species delimitation studies. For example, one may

reduce dimensionality based on *a priori* considerations about which set of characters may be best to

diagnose particular species. In particular, when *a priori* information about specific traits separating species

is available (e.g. original species descriptions), one should favor analyzing variation in such traits; far

from being circular (McKay and Zink 2015), it is only natural that one should critically examine evidence

for species limits precisely in the dimensions in which such limits are hypothesized to exist (see also

Remsen 2010; Patten and Remsen 2017). Alternatively, one may use methods that aim to find the set of

variables (phenotypic traits) that best discriminates groups in a NMM, with no *a priori* information about

groups (Raftery and Dean 2006; Maugis et al. 2009a; Maugis et al. 2009b).

3) *Differences in central tendency are not evidence of distinct phenotypic distributions.*

Distinct normal distributions in quantitative characters constitute evidence for the existence of distinct

species, but differences in central tendency between groups of individuals do not. This issue has been

pointed out previously (e.g., Mayr et al. 1953; Luckow 1995; Patten and Unitt 2002), but seems to be

ignored when statistical procedures to investigate differences in central tendency (e.g., t-tests, analysis of

variance, Cohen's d) are advanced as potentially valid tools to evaluate species boundaries (e.g., Simpson

1951; Henderson 2006; Tobias et al. 2010). McKay and Zink (2015, page 695 and their Fig. 2) have done

as much by suggesting that statistical differences in average phenotypes between allopatric island

populations of ground-finches could be equated to distinct morphological groups which, in turn, would

have to be recognized as species.

Because this issue seems to commonly afflict assessments of species limits between allopatric forms

(Tobias et al. 2010; McKay and Zink 2015), we illustrate it with a hypothetical example of two allopatric

populations. Grouping specimens from these populations according to collection localities (e.g., two

islands) reveals statistically significant differences in the central tendency of phenotypes (Fig. 3a). Yet,

there is no evidence that phenotypic variation across specimens from the two populations is best described

by more than a single normal distribution (Fig. 3b). Therefore, there is no evidence for more than one

species in the sample of specimens regardless of differences in average phenotypes. This illustration

focuses on groups of specimens defined by geography (i.e., allopatric populations), but the issue may

affect comparisons involving groups of specimens defined by time (i.e., allochronic populations; Simpson

1951) or by any other criterion.

The solution to the problem is simple: do not treat phenotypic differences in central tendency as evidence

for the existence of distinct phenotypic groups and, therefore, distinct species. No matter how statistically

significant, even very large effect sizes are not germane in light of the basic model for species delimitation

based on quantitative phenotypic characters. In light of this model, the focus of analysis should be on

determining the number of normal distributions needed to describe phenotypic variation among

specimens, as well as on estimating the parameters of those distributions (e.g., means and variance-

covariance matrices). Indeed, strong evidence may exist for more than one distinct normal distribution in

the absence of differences in central tendency (Hennig 2010), suggesting biologically meaningful

differences between species in the variance of phenotypic traits (Supplementary Material, Appendix 1). As

explained above, NMMs are a useful tool to test for distinct normal distributions.

243   ARE THERE PHENOTYPICALLY DISTINCT GROUPS OF GROUND-FINCHES?

We examined phenotypic variation among *Geospiza* ground-finches by analyzing data from six

morphological measurements of museum specimens (wing length, tail length, tarsus length, bill length,

bill width, and bill depth) taken on adult males by H. S. Swarth for his monographic revision of the birds

of the Galapagos (Swarth 1931). These were the same data employed by McKay and Zink (2015), which

we use here with permission from the California Academy of Sciences; our sample sizes differ from those

of the earlier study (501 vs. 486 male individuals) because we excluded a few individuals that were

duplicated in the original dataset. The data we employed and the R code used to conduct the analyses

described below are available as supplementary material (Appendices 2 and 3, respectively).

We asked how many distinct groups of ground-finches exist in the Galapagos using morphological data

from specimens collected across the archipelago (total 18 islands). To define the morphological space for

this analysis, we followed McKay and Zink (2015) and used PCA on the covariance matrix of log-

transformed data. Rather than examining evidence for species limits using only the first three principal

components accounting for >99% of the variation (McKay and Zink 2015), we used the R package

clustvarsel (Scrucca and Raftery 2004) to reduce the dimensionality of the data by selecting the set of

principal components most useful for group discrimination in NMMs, without *a priori* information about

groups (Raftery and Dean 2006; Maugis et al. 2009a; Maugis et al. 2009b). We used the R package mclust

5.0 (Scrucca et al. 2016) to fit a wide range of NMMs. At one extreme, NMMs assuming one

morphological group represented the Sisyphean evolution hypothesis that there is a single species of

ground-finch (McKay and Zink 2015). Toward the opposite end, NMMs assuming up to 30 distinct

morphological groups represented hypotheses alluded to by McKay and Zink (2015) when they suggested

ground-finches may comprise "dozens of cluster species", or "1 or 6 or 30 species" (p. 695). We also

fitted NMMs specifying the six (Lack 1947) or nine (Lamichhaney et al. 2015; Remsen et al. 2017)

species recognized by alternative taxonomic treatments of *Geospiza*, using the original specimen

identifications in Swarth's data updated to reflect changes in nomenclature. We used the Bayesian

Information Criterion (BIC; Schwarz 1978) to measure empirical support for different NMMs (Fraley and

Raftery 2002) and thereby explicitly evaluated the hypothesis that there is only one species of ground-

finch (McKay and Zink 2015) relative to hypotheses that there are several species in the group (Lack

1947; Lamichhaney et al. 2015; Remsen et al. 2017).

We found the first four principal components to be most useful for group discrimination; NMMs ignoring

the fourth principal component, although it explained only 0.006% of the morphological variance, had

substantially less empirical support (ΔBIC $\geq$ 55) than those including it. Therefore, in contrast to McKay

and Zink (2015), we did not discard the fourth principal component for analysis. The models specifying

seven and eight distinct morphological groups of ground-finches received the strongest support (ΔBIC $\leq$

1.26). Support for all other models was considerably lower (ΔBIC in all cases >20; Fig. 4). In turn, the

model with the lowest support represented the Sisyphean evolution hypothesis proposing no distinct

morphological groups of ground-finches (i.e., that there is a single group; McKay and Zink 2015), which

had a 500 BIC difference to the second-worse model and > 82 BIC difference to the two best models.

Relative to the best models, those specifying groupings consistent with taxonomy recognizing six or nine

species were weakly supported (Fig. 4), considering differences in BIC scores greater than 6 are typically

regarded as strong or very strong evidence against models with lower support (Kass and Raftery 1995). In

sum, the data provided poor empirical support for the hypothesis that ground-finches consist of only one

species (McKay and Zink 2015) and strongly supported hypotheses of several morphologically distinct

groups (Fig. 5, Supplementary Fig. 1); however, those groups did not readily align with existing

taxonomic treatments of *Geospiza* (Fig. 6, Supplementary Fig. 2).

Despite their comparatively low empirical support, models specifying six or nine morphological groups

according to taxonomic treatments of *Geospiza* (Lack 1947; Remsen et al. 2017) were partially consistent

with the best models (Fig. 6 and Supplementary Fig. 2). For example, in the best models, all specimens of

two of the nine currently recognized species (*G. scandens, G. septentrionalis*) were assigned to two

respective morphological groups which included few or no specimens of other species (Fig. 6;

Supplementary Fig. 2). Discrepancies between our analysis and current taxonomy were most evident in

cases such as those of (1) *G. propinqua, G. conirostris* and *G. fortis*, which were assigned to three, three

(or two) and four morphological groups, respectively, or (2) *G. fuliginosa* and *G. acutirostris*, in which all

specimens were assigned to the same group to the exclusion of nearly all specimens of other species. In

addition, some morphological groups included specimens of multiple species (e.g., morphological group 1

contained specimens identified as *G. propinqua, G. fortis, G. magnirostris*, and *G. conirostris*). Part of the

lack of agreement between the morphological groups we detected and groups recognized by taxonomy

may be accounted for by considering that species may be told apart by phenotypic characters different

from those we considered. For example, *G. fuliginosa* and *G. acutirostris* are indistinguishable in our

analysis, but are distinct given subtle differences in bill profile and marked differences in songs (Grant and

Grant 2008). Likewise, some of the discrepancies with current taxonomy (Remsen et al. 2017) involved

cases in which species delimitation was not based on morphology, but rather resulted from recent genomic

analyses revealing that phenotypically similar populations are distantly related (Lamichhaney et al. 2015).

This likely explains why our analysis did fully discriminate some species pairs in the morphological space

we examined (*G. conirostris* vs. *G. propinqua, G. difficils* vs. *G. septentrionalis*), although they may be

more distinct in other phenotypic spaces including bill profile and song as well as behavior (Grant et al.

2000; Grant and Grant 2002). Also, we assumed that specimen identifications in the data set we analyzed

were faultless; thus, part of the apparent mismatch between morphological groups detected in our analyses

and taxonomy may reflect identification errors. Evaluating this possibility would require detailed

examinations of individual specimens beyond the scope of our work.

Geographic context is an important consideration in assessments of species limits using phenotypic traits.

Under a wide range of species definitions (sensu de Queiroz 1998), distinct phenotypic groups among

sympatric individuals are readily accepted as evidence for the existence of distinct species (Mayr 1992;

Mallet 2008), whereas distinct phenotypic groups corresponding to non-sympatric populations may be less

readily accepted as evidence of distinct species because they may reflect within-species differentiation due

to geographic isolation or local adaptation (Zapata and Jiménez 2012). The morphological groups of

ground-finches we detected (Fig. 5, Supplementary Fig. 1) cannot be interpreted to reflect within-species,

among-island variation because these groups occurred on multiple islands and were sympatric with other

groups; all of the morphological groups identified in the best NMMs were widely distributed across the

Galapagos Archipelago (median = 8.5 or 9.0, range 3-14 islands per group; Table 1 and Supplementary

Table 1) and most islands harbored several groups (up to 6 in Santiago and 7 in Santa Cruz; Fig. 7 and

Supplementary Figure 3). Importantly, almost all morphological groups co-occurred with each other in at

least one island; the only exception was morphological group 3 in one of the models, which co-occurred

with four out of the other seven groups (Table 1 and Supplementary Table 1). McKay and Zink (2015)

indicated that different morphs of ground-finches exist within islands and argued that if such morphs were

treated as species, then one would need to recognize dozens of species in the group; our analysis suggests

this is not the case given the occurrence of all morphological groups in multiple islands.

At this point we note that because the specimens we analyzed were collected several decades ago (Swarth

1931), they may not faithfully reflect patterns in morphological variation nor the geographic distributions

of morphological groups in the present. This is because over the past century, ground-finch populations

have experienced a few colonization and extinction events, changes in the degree of morphological

differentiation among populations due to natural and human-mediated hybridization, and bouts of

selection in shifting directions over multiple generations in association with environmental variation in

space and time (Harris 1973; De León et al. 2011; Grant and Grant 2014). Thus, we refrain from

additional discussions about species limits involving comparisons of historical morphological data with

contemporary evidence (e.g., genomics; Lamichhaney et al. 2015). Nonetheless, our analyses serve to

demonstrate that statistically distinct morphological groups of ground-finches existed in the past, and we

strongly suspect they still exist in the present. Accordingly, we suggest that the burden of proof for

systematists proposing to lump ground-finches into a single species based on morphological data is on

showing that distinct groups do not longer exist.

349   CONCLUSIONS; OR, ATLANTEAN EVOLUTION IN DARWIN'S FINCHES

Our reanalysis of morphological data pointed strongly to the existence of several groups of phenotypically

distinct *Geospiza* ground-finches based only on six linear morphological measurements. In addition, we

found evidence of distinct phenotypes in geographic scenarios (i.e., sympatry within islands) where one

should not expect them if populations had not achieved evolutionary independence. Specifically, because

the variation in quantitative morphological traits we examined is polygenic (Abzhanov et al. 2004;

Abzhanov et al. 2006; Lamichhaney et al. 2015; Chaves et al. 2016; Lamichhaney et al. 2016) and not

caused by differences in sex or age (we restricted analyses to adult males), the existence of distinct

phenotypic groups in areas where populations come into contact implies there are likely several species of

ground-finches. Therefore, we contend that ground-finches are not an example of Sisyphean evolution

(McKay and Zink 2015), a term that could well apply to other systems in nature (Seehausen 2006; Nosil et

al. 2009; Rudman and Schluter 2016). Instead, evolutionary forces maintaining populations of ground-

finches apart are likely in place, just as in Greek mythology Atlas prevents the merging of the Earth and

the sky with his shoulders. Ground-finches thus likely represent an example of what one might call

"Atlantean evolution". One, of course, does not need a new term to refer to speciation, but thinking of

Atlas brings to mind *atlas,* a collection of maps, which reminds one of the central role of geography in

speciation and in the basic model underlying species delimitation based on phenotypic variation.

The question of exactly how many species of Darwin's ground-finches are there remains open and

requires further attention to morphology, including careful scrutiny of discrepancies between

morphological variation and taxonomy (e.g., Fig. 6). In addition, morphological variation should be

further examined in light of biological factors including additional phenotypic characters, ecological

niches, mating behavior, population dynamics, and patterns of genetic and genomic variation among

populations (Grant 1999; Huber et al. 2007; Grant and Grant 2008; Farrington et al. 2014; Grant and Grant

2014; Lamichhaney et al. 2015; McKay and Zink 2015). Fruitful discussions about species limits in the

group would likely start by addressing some of the additional thought-provoking arguments advanced by

McKay and Zink (2015) that we did not touch on and which are beyond the scope of our work (e.g., the

extent to which morphological groups are stable lineages over time or the evidence for the existence of

distinct gene pools). Any such discussions, however, as well as discussions over species delimitation in

other organisms, should bear in mind that phenotypic evidence for species limits is best assessed using

statistical approaches appropriately grounded on evolutionary models.

380   OUTLOOK

We do not claim that the approaches used here to analyze phenotypic data for species delimitation are free

of problems. Issues such as estimation of the number of groups in NMMs (McLachlan and Peel 2000;

McLachlan and Rathnayake 2014) or how to select variables for NMM analyses of multidimensional

datasets (Poon et al. 2013) are critical areas of active research in statistics in which progress remains to be

made. Despite these issues, however, we argue that the statistical tools we used are appropriate because

they are directly related to the basic evolutionary model underlying species delimitation using phenotypic

data (Fisher 1918). Moreover, these tools allow systematists to go beyond fairly limited graphical analysis,

and to break free from problems resulting from reduction of dimensionality using PCA or related

approaches and from comparisons of measurements of central tendencies. The value of embracing

approaches with a solid theoretical basis despite limitations in their implementation in systematics is clear

considering other developments in the field in which theory predated robust methodologies that

subsequently blossomed. Such developments include the use of statistical methods to study species limits

among fossil populations (Newell 1956), the application of probabilistic models to infer phylogenetic trees

(Felsenstein 1981), time-calibration of molecular phylogenies (Kishino and Hasegawa 1990), and the

estimation of species trees from gene trees (Maddison 1997).

Practical approaches to fit NMMs without *a priori* information about species limits offer a fresh

perspective in inferences available to systematists. In the absence of these tools, it seemed reasonable to

argue that species limits should be based on fixed phenotypic differences because continuous variation

could only be subdivided using subjective criteria (Cracraft 1989; Davis 1997). Accordingly, overlap of

phenotypic ranges has been conventionally stressed as a criterion to suggest samples of individuals are

conspecific (e.g., Simpson 1951; Davis and Heywood 1963; Zink 2002; McKay and Zink 2015).

However, overlap in phenotypic ranges under the framework offered by NMMs is not relevant for species

delimitation because (1) one may find strong empirical support for models in which the phenotypic ranges

of distinct normal distributions overlap (e.g., Fig. 1 and 5), and (2) absence of range overlap does not

imply strong empirical support for models with more than a single species (e.g., Fig. 3). Critically,

absence of range overlap need not imply a phenotypic gap (i.e., a phenotypic region with low frequency of

individuals) because continuous phenotypic variation can be arbitrarily split into mutually exclusive parts

regardless of phenotype frequencies. On the other hand, although true phenotypic gaps (along with

multimodality in phenotypic distributions) are sufficient to suggest species boundaries (Zapata and

Jiménez 2012; Mallet 2013), they are not necessary to demonstrate such boundaries exist because NMMs

specifying more than one species may be strongly supported in the absence of phenotypic gaps. An

example of support for more than one normal distribution in the absence of phenotypic gaps was provided

at the inception of NMMs: Karl Pearson inferred two groups among specimens of the shore crab

(*Carcinus maenas*) from the Bay of Naples, even though the mixture of the groups was not bimodal and

therefore they were not separated by a gap (Pearson 1894). Moreover, Pearson examined the possibility of

inferring the existence of groups with different phenotypic variances but identical phenotypic means,

which are by definition not separated by a gap (Supplementary Material Appendix 1).

To conclude, we note that the criteria for species delimitation discussed above are relevant in the context

of ideas about the reality of species. In particular, it has been argued that if the hypothesis that species are

real entities in nature is correct, then biological diversity should be a patchwork of phenotypic clusters

delineated by gaps (Coyne and Orr 2004; Barraclough and Humphreys 2015). This prediction, however,

would not follow from the hypothesis that species are real if, as we argue, phenotypic gaps are not

necessary criteria for species delimitation. In other words, species may be real entities in nature even if

phenotypic gaps are not major elements structuring biological diversity. Because statistical approaches

related to NMMs now allow systematists to make unprecedented formal inferences about the existence of

species even if they overlap in phenotypic space, they constitute particularly useful tools to describe the

structure of biological diversity, a necessary step to understand the evolutionary processes that generated

it.

432   ACKNOWLEDGEMENTS

433   We thank the California Academy of Sciences (Jack Dumbacher) for allowing us to use and publish

434   morphological data from H. S. Swarth's archive. Bailey McKay kindly provided tabulated data. We thank

435   Peter Grant, Van Remsen, Elizabeth Spriggs, Peter Stevens, and members C.D. Cadena's laboratory group

436   for discussion and helpful comments on the manuscript.

439   REFERENCES

Abzhanov A, Kuo WP, Hartmann C, Grant BR, Grant PR, Tabin CJ. 2006. The calmudolin pathway and

evolution of elongated beak morphology in Darwin's finches. Nature, 442:563-567.

Abzhanov A, Protas M, Grant BR, Grant PR, Tabin CJ. 2004. *Bmp4* and morphological variation of beaks

in Darwin's finches. Science, 305:1462-1465.

Allmon WD. 2016. Studying species in the fossil record: A review and recommendations for a more

unified approach. In: Allmon WD, Yacobucci MM editors. Species and speciation in the fossil

record. Chicago, IL, USA, University of Chicago Press, p. 59-120.

Barraclough TG, Humphreys AM. 2015. The evolutionary reality of species and higher taxa in plants: a

survey of post-modern opinion and evidence. New Phytologist, 207:291-296.

Bowman R. 1961. Morphological differentiation and adaptation in the Galapagos finches. University of

California Publications in Zoology, 58:1-302.

Camargo A, Sites JW, Jr. 2013. Species delimitation: A decade after the renaissance. In: Pavlinov IY

editor. The Species Problem: Ongoing Issues, InTech DOI: 10.5772/3313, p. 225-247.

Carstens B, Pelletier TA, Reid NM, Satler JD. 2013. How to fail at species delimitation. Molecular

Ecology, 22:4369-4383.

Chang W-C. 1983. On using principal components before separating a mixture of two multivariate normal

distributions. Applied Statistics, 32:267-275.

Chaves JA, Cooper EA, Hendry AP, Podos J, De León LF, Raeymaekers JAM, Macmillan WO, Uy JAC.

2016. Genomic variation at the tips of the adaptive radiation of Darwin's finches. Molecular

Ecology, 25:5282-5295.

Cooper Ornithological Society. 2016. Katma Award 2015, to Bailey McKay and Robert Zink. Condor,

118:209-210.

Coyne JA, Orr HA. 2004. Speciation. Sunderland, Massachusetts, Sinauer Associates.

Cracraft J. 1989. Speciation and its ontology: the empirical consequences of alternative species concepts

for understanding patterns and processes of differentiation. In: Otte D, Endler JA editors. Speciation

and its consequences. Sunderland, Massachussetts, Sinauer Associates, p. 28-59.

Davis JI. 1997. Evolution, evidence and the role of species concepts in phylogenetics. Systematic Botany,

22:373-403.

Davis PH, Heywood V. 1963. Principles of angiosperm taxonomy. Princeton, NJ, USA, D. van Nostrand

Co.

De León LF, Raeymaekers JAM, Bermingham E, Podos J, Herrel A, Hendry AP. 2011. Exploring

possible human influences on the evolution of Darwin's finches. Evolution, 65:2258-2272.

de Queiroz K. 1998. The general lineage concept of species, species criteria, and the process of speciation.

In: Howard DJ, Berlocher SH editors. Endless forms: species and speciation. New York, Oxford

University Press, p. 57-75.

Edwards DL, Knowles LL. 2014. Species detection and individual assignment in species delimitation: can

integrative data increase efficacy? Proceedings of the Royal Society of London B, 281:20132765.

Ezard THG, Pearson PN, Purvis A. 2010. Algorithmic approaches to aid species' delimitation in

multidimensional morphospace. BMC Evolutionary Biology, 10:175.

Farrington HL, Lawson LP, Clark CM, Petren K. 2014. The evolutionary history of Darwin's finches:

speciation, gene flow, and introgression in a fragmented landscape. Evolution, 68:2932-2944.

Felsenstein J. 1981. Evolutionary trees from DNA sequences: a maximum likelihood approach. Journal of

Molecular Evolution, 17:368-376.

Fisher RA. 1918. The correlation between relatives on the supposition of Mendelian inheritance.

Transactions of the Royal Society of Edinburgh, 52:399-433.

Fraley C, Raftery AE. 2002. Model-based clustering, discriminant analysis, and density estimation.

Journal of the American Statistical Association, 97:611-631.

Fraley C, Raftery AE, Murphy TB, Scrucca L. 2012. mclust Version 4 for R: Normal Mixture Modeling

for Model-Based Clustering, Classification, and Density Estimation. Technical Report No. 597.

Department of Statistics, University of Washington.

Fujita MW, Leaché AD, Burbrink FT, McGuire JA, Moritz C. 2012. Coalescent-based species

delimitation in an integrative taxonomy. Trends in Ecology and Evolution, 27:480-488.

Futuyma DJ. 2013. Evolution, third edition. Sunderland, Massachusetts, Sinauer Associates.

Grant BR, Grant PR. 2002. Simulating secondary contact in allopatric speciation: an empirical test of

premating isolation. Biological Journal of the Linnean Society, 76:545-556.

Grant PR. 1999. Ecology and evolution of Darwin's finches. Princeton, New Jersey, USA, Princeton

University Press.

Grant PR, Abbott I, Schluter D, Curry RL, Abbott LK. 1985. Variation in the size and shape of Darwin's

finches. Biological Journal of the Linnean Society, 25:1-39.

Grant PR, Grant BR. 2008. How and why species multiply: the radiation of Darwin's finches. Princeton,

New Jersey, USA, Princeton University Press.

Grant PR, Grant BR. 2014. 40 Years of Evolution: Darwin's Finches on Daphne Major Island. Princeton,

New Jersey, USA, Princeton University Press.

Grant PR, Grant BR, Petren K. 2000. The allopatric phase of speciation: the sharp-beaked ground finch

(*Geospiza difficilis*) on the Galápagos islands. Biological Journal of the Linnean Society, 69:287-

317.

Guillot G, Renaud S, Ledevin R, Michaux J, Claude J. 2012. A unifying model for the analysis of

phenotypic, genetic, and geographic data. Systematic Biology, 61:897-911.

Harris MP. 1973. The Galápagos avifauna. Condor, 75:265-278.

Henderson A. 2006. Traditional morphometrics in plant systematics and its role in palm systematics.

Botanical Journal of the Linnean Society, 151:103-111.

Hennig C. 2010. Methods for merging Gaussian mixture components. Advances in Data Analysis and

Classification, 4:3-34.

Huber SK, De León LF, Hendry AP, Bermingham E, Podos J. 2007. Reproductive isoloation of sympatric

morphs in a population of Darwin's finches. Proceedings of the Royal Society of London B,

274:1709-1714.

Kass RE, Raftery AE. 1995. Bayes factors. Journal of the American Statistical Association, 90:773-795.

Kishino H, Hasegawa M. 1990. Converting distance to time: Application to human evolution. Methods in

Enzymology, 183:550-570.

Kleindorfer S, O'Connor JA, Dudaniec RY, Myers SA, Robertson J, Sulloway FJ. 2014. Species collapse

via hybridization in Darwin's tree finches. American Naturalist, 183:325-341.

Lack D. 1947. Darwin's Finches. New York, Cambridge University Press.

Lamichhaney S, Berglund J, Almén MS, Maqbool K, Grabherr M, Martinez-Barrio A, Promerová M,

Rubin C-J, Wang C, Zamani N*, et al.* 2015. Evolution of Darwin's finches and their beaks revealed

by genome sequencing. Nature, 518:371-375.

Lamichhaney S, Han F, Berglund J, Wang C, Almén MS, Webster MT, Grant BR, Grant PR, Andersson

L. 2016. A beak size locus in Darwin's finches facilitated character displacement during a drought.

Science, 352:470-474.

Luckow M. 1995. Species concepts: assumptions, methods, and applications. Systematic Botany, 20:589-

518.

Maddison WP. 1997. Gene trees in species trees. Systematic Biology, 463:523-536.

Mallet J. 2008. Hybridization, ecological races and the nature of species: empirical evidence for the ease

of speciation. Philosophical Transactions of the Royal Society of London B, 363:2971-2986.

Mallet J. 2013. Species, concepts of. In: Levin SA editor. Encyclopedia of Biodiversity. Volume 6.

Waltham, Mass, Academic Press, p. 679-691.

Maugis C, Celeux G, Martin-Magniette M-L. 2009a. Variable selection for clustering with Gaussian

mixture models. Biometrics, 65:701-709.

Maugis C, Celeux G, Martin-Magniette M-L. 2009b. Variable selection in model-based clustering: A

general variable role modeling. Computational Statistics and Data Analysis, 53:3872-3882.

Mayr E. 1992. A local flora and the biological species concept. American Journal of Botany, 79:222-238.

Mayr E, Linsley EG, Usinger RL. 1953. Methods and Principles of Systematic Zoology. New York,

McGraw-Hill.

McDade LA. 1995. Species concepts and problems in practice: insights from botanical monographs.

Systematic Botany, 20:606-622.

McKay BD, Zink RM. 2015. Sisyphean evolution in Darwin's finches. Biological Reviews, 90:689-698.

McLachlan G, Krishnan T. 2008. The EM Algorithm and Extensions, second edition. Hoboken, NJ, USA,

John Wiley and Sons.

McLachlan G, Peel DA. 2000. Finite mixture models. Hoboken, NJ, USA, Wiley Series in Probability and

Statistics, John Wiley and Sons.

McLachlan GJ. 2004. Discriminant analysis and statistical pattern recognition. Hoboken, NJ, USA, Wiley

Series in Probability and Statistics, John Wiley and Sons.

McLachlan GJ, Rathnayake S. 2014. On the number of components in a Gaussian mixture model. WIREs

Data Mining Knowledge and Discovery, 4:341-355.

Miller W. 2016. The species problem: concepts, conflicts, and patterns preserved in the fossil record. In:

Allmon WD, Yacobucci MM editors. Studying species in the fossil record: A review and

recommendations for a more unified approach. Chicago, IL, USA, University of Chicago Press, p.

28-58.

Moczek AP, Emlen DJ. 1999. Proximate determination of male horn dimorphismin the beetle

Onthophagus taurus (Coleoptera: Scarabaeidae). Journal of Evolutionary Biology, 12:27-37.

Newell ND. 1956. Fossil populations. In: Sylvester-Bradley PC editor. The species concept in

paleontology: A symposium. London, UK, London Systematics Association, p. 63-82.

Nosil P, Harmon LJ, Seehausen O. 2009. Ecological explanations for (incomplete) speciation. Trends in

Ecology and Evolution, 24:145-156.

Padial JM, Miralles A, De la Riva I, Vences M. 2010. The integrative future of taxonomy. Frontiers in

Zoology, 7:216.

Patten MA, Remsen JV, Jr. 2017. Complementary roles of phenotype and genotype in subspecies

delimitation. Journal of Heredity, in press:doi:10.1093/jhered/esx1013.

Patten MA, Unitt P. 2002. Diagnosability versus mean differences of sage sparrow subspecies. Auk,

119:26-35.

Pearson K. 1894. Contributions to the theory of mathematical evolution. Philosophical Transactions of the

Royal Society of London A, 185:71-110.

Poon LKM, Zhang NL, Liu T, Liu AH. 2013. Model-based clustering of high-dimensional data: Variable

selection versus facet determination. International Journal of Approximate Reasoning, 54:196-215.

Raftery AE, Dean N. 2006. Variable selection for model-based clustering. Journal of the American

Statistical Association, 101:168-178.

Remsen JV, Jr. 2010. Subspecies as a meaningful taxonomic rank in avian classification. Ornithological

Monographs, 67:62-78.

Remsen JV, Jr., Areta JI, Cadena CD, Claramunt S, Jaramillo A, Pacheco JF, Pérez-Emán J, Robbins MB,

Stiles FG, Stotz DF, *et al.* 2017. A classification of the bird species of South America, version

[date]. American Ornithologists' Union. http://www.museum.lsu.edu/~Remsen/SACCBaseline.htm.

Rieseberg LH, Wood TE, Baack EJ. 2006. The nature of plant species. Nature, 440:524-527.

Rising JD, Jaramillo A, Copete JL, Ryan PG, Madge SC. 2011. Family Emberizidae (Buntings and New

World Sparrows). In: del Hoyo J, Elliott A, Christie DA editors. Handbook of the Birds of the

World. Vol. 16. Tanagers to New World Blackbirds. Barcelona, Lynx Edicions, p. 428-683.

Rudman SM, Schluter D. 2016. Ecological impacts of reverse speciation in threespine stickleback. Current

Biology, 26:490-495.

Sangster G. 2014. The application of species criteria in avian taxonomy and its implications for the debate

over species concepts. Biological Reviews, 89:199-214.

Schwarz G. 1978. Estimating the dimension of a model. Annals of Statistics, 6:461-464.

Scrucca L, Fop M, Murphy TB, Raftery AE. 2016. mclust 5: Clustering, classification and density

estimation using Gaussian finite mixture models. The R Journal, 8:289-317.

Scrucca L, Raftery AE. 2004. clustvarsel: A package implementing variable selection for model-based

clustering in R. Preprint available on arXiv https://arxiv.org/abs/1411.0606.

Seehausen O. 2006. Conservation: losing biodiversity by reverse speciation. Current Biology, 16:R334-

R337.

Simpson GG. 1951. The species concept. Evolution, 5:285-298.

Sites JW, Jr., Marshall JC. 2003. Delimiting species: a Renaissance issue in systematic biology. Trends in

Ecology and Evolution, 18:462-470.

Sites JW, Jr., Marshall JC. 2004. Operational criteria for delimiting species. Annual Review of Ecology,

Evolution and Systematics, 35:199-227.

Smith TB. 1993. Disruptive selection and the genetic basis of bill size polymorphism in the African finch

Pyrenestes. Nature, 363:618-620.

Sneath PHA, Sokal RR. 1973. Numerical taxonomy. San Francisco, CA, W. H. Freeman and Co.

Solís-Lemus C, Knowles LL, Ané C. 2014. Bayesian species delimitation combining multiple genes and

traits in a unified framework. Evolution, 69:492-507.

Sukumaran J, Knowles LL. 2017. Multispecies coalescent delimits structure, not species. Proceedings of

the National Academy of Sciences of the USA, 114:1607-1612.

Swarth HS. 1931. The avifauna of the Galápagos Islands. Occasional Papers of the California Academy of

Sciences, 18:1-299.

Templeton AR. 2006. Population genetics and microevolutionary theory. Hoboken, NJ, USA, John Wiley

& Sons.

Tobias JA, Seddon N, Spottiswoode CN, Pilgrim JD, Fishpool LDC, Collar NJ. 2010. Quantitative criteria

for species delimitation. Ibis, 152:724-746.

Wiens JJ, Servedio MR. 2000. Species delimitation in systematics: inferring diagnostic differences

between species. Proceedings of the Royal Society of London B, 267:631-636.

Zapata F, Jiménez I. 2012. Species delimitation: Inferring gaps in morphology across geography.

Systematic Biology, 61:179-194.

Zink RM. 2002. A new perspective on the evolutionary history of Darwin's finches. Auk, 119:864-871.

620
Figure 1. Visual inspection of phenotypic data may yield limited insight regarding species limits. (A)
Sample of 200 museum specimens (triangles) arranged according to a morphological phenotype (triangle
size), from small in the upper left to large in the lower right. The specimens appear to form a smooth
gradient with no morphological gaps. (B) Plot of specimen measurements along a single continuous axis
representing the size of the morphological trait in A. At the resolution of the measurements, extreme
values in the sample seem to be gradually connected by intermediate phenotypes throughout. Thus, there
seem to be no obvious morphological gap, suggesting the specimens correspond to a single variable
species. (C) Two distinct normal distributions are revealed by examining the frequency (gray bars) of
specimen phenotypes in the sample, suggesting that specimens may correspond to two species. In fact, the
sample was drawn from a mixture of two normal distributions (continuous black lines).

633
634
635
Figure 2. The ability of principal components to discriminate species is not necessarily proportional to the
total phenotypic variance they explain. (A) Hypothetical example of two distinct species in the space
defined by two phenotypic traits. Each species is described by a bivariate normal distribution, shown as
ellipses covering 95% of the individuals of each species. Dotted lines represent the two principal
component axes of the normal mixture of the two species. Note that the two principal components are
orthogonal, forming a right angle that may not be apparent due to the magnification of the ordinate
relative to the abscissa. (B) Probability density of individuals of the two species along the first principal
component (PC1). This axis is useless to discriminate the phenotypic distributions of the two species,
despite the fact that it explains 99.73% of the variance. (C) Probability density of individuals of the two
species along the second principal component (PC2). The phenotypes of the two species can be readily
distinguished along this axis, even though it explains only 0.27% of the variance. Because systematists
often discard phenotypic axes accounting for small fractions of the total variance, they may miss crucial
phenotypic evidence for species limits.
649

651
652
653
Figure 3. Differences in central tendency are not evidence of distinct phenotypic distributions. (A)
Specimens from two allopatric populations (gray and striped histogram bars) differ markedly in the central
tendency of a phenotypic trait (abscissa), as described by normal probability density functions (continuous
black lines) and boxplots on top. The boxplots show the median (solid thick line), the interquartile range
(box), whiskers extending to the most extreme values within $1.5 \times$ interquartile ranges from the box, and
outliers. The difference between means (1.717 and 2.707) is statistically significant (0.99, 95% CI: 0.89 −
1.09, t-test p-value $< 2 \times 10^{-16}$). Cohen's d = 2.66, which is generally regarded as a large effect size. Note
that the phenotypic ranges of the two populations do not overlap. (B) Despite the difference in central
tendency and absence of phenotypic overlap, a single normal distribution (continuous black line) describes
phenotypic variation across all specimens (gray bars) better than two normal distributions. In particular,
empirical support for a normal mixture model assuming two distinct normal distributions (A) is
substantially lower than that for a model specifying a single normal distribution (B): a difference of 10 in
Bayesian Information Criterion (Schwarz 1978, Fraley and Raftery 2002). Thus, in light of the basic
model for species delimitation based on quantitative phenotypic characters, there are no grounds to
suggest that the specimens represent two distinct species despite marked differences in central tendency.

669
670
Figure 4. Analysis of morphological data strongly supported hypotheses that there are multiple distinct
groups of *Geospiza* ground-finches. The plot shows the empirical support (ordinate) for normal mixture
models assuming 1-30 distinct morphological groups (abscissa), and for the two models specifying
groupings of specimens reflecting taxonomic treatments recognizing six species (Lack 1947, Rising et al.
2011) or nine species (Remsen et al. 2017). Empirical support was measured as difference in Bayesian
Information Criterion relative to the best model (ΔBIC). The two models with highest empirical support
assumed seven and eight distinct morphological groups. Empirical support for the model corresponding to
the Sisyphean evolution hypothesis positing there is a single species of ground-finch (i.e. a single
morphological group; McKay and Zink 2015) was negligible (ΔBIC > 820).
680
681

Figure 5. Eight morphological groups of *Geospiza* ground-finches identified by one of the best normal
mixture models. Panels A–F show the groups in the space defined by the four principal components most
useful for group discrimination (PC1–PC4). Colored symbols represent specimens assigned to different
morphological groups and ellipses show 95% high density regions for normal distributions representing
each morphological group. Arrows in G and H display the contribution of measured morphological traits
to each principal component, gauged by the loadings of each trait on each principal component (i.e.,
elements of normalized eigenvectors). Circles show the length of arrows expected if all six traits
contributed equally to bidimensional principal component spaces; arrows exceeding this expectation
contribute most significantly and are labeled. PC1 and PC2 reflect general aspects of beak size and shape
(G), with group 2 having long, deep and wide beaks, group 7 having short, shallow and narrow beaks, and
morphological group 8 having long, shallow and narrow beaks. PC3 and PC4 reflect aspects of tail and
tarsus length (H), with group 5 having a relatively long tail, and group 6 having a relatively long tarsus.
PC4 is particularly useful to distinguish group 6 despite explaining only 0.6% of the total variance. The
morphological distribution of groups in the other well supported NMM is fairly similar to the one shown
here, the main difference being that group 3 is merged into groups 1 and 8 (Supplementary Fig. 1).

Current taxonomy (Remsen et al. 2017)

Previous taxonomy (Lack 1947, Rising et al. 2011)

Morphological group

Figure 6. Eight morphological groups of *Geospiza* ground-finches in the Galapagos Archipelago identified by one of the best normal mixture models partially correspond to the nine species recognized by current taxonomy (Remsen et al. 2017) and to the six species recognized by previous taxonomy (Lack 1947, Rising et al. 2011). Each histogram shows, for each recognized species, the number of specimens assigned to each of the eight morphological groups. Groups are colored according to the scheme in Fig. 5.

Figure 7. Eight distinct morphological groups of *Geospiza* ground-finches identified by one of the best normal mixture models have broad geographic distributions across the Galapagos Archipelago. For each island, numbers indicate individuals included in the analysis and ringplots depict the fraction of such individuals assigned to each morphological group following the color scheme in Fig. 5. The existence of distinct morphological groups in potential sympatry within islands (e.g., >4 groups in Santa Cruz, Santiago, and Pinta) suggests that such groups are unlikely to reflect within-species differentiation due to geographic isolation or local adaptation.

Table 1. Number of islands in the Galapagos Archipelago where each of the eight morphological groups
of *Geospiza* ground-finches identified by one of the best normal mixture models were found to occur
(diagonal) and co-occur with other groups (off diagonal). All groups co-occurred with each other in at
least one island, except for cases involving group 3, which did not co-occur with three other groups. Note,
however, that group 3 was not recovered as distinct in the other best model, which identified only seven
groups (Supplementary Table 1).
726

	Group 1	Group 2	Group 3	Group 4	Group 5	Group 6	Group 7	Group 8
Group 1	8	4	2	4	2	1	6	4
Group 2		9	2	4	4	3	7	5
Group 3			3	0	1	0	1	0
Group 4				11	2	3	10	7
Group 5					4	2	2	2
Group 6						3	3	3
Group 7							14	3
Group 8								9

727
728
729

Supplementary Material

Appendix 1. Biological significance of differences between species in variance of phenotypic traits despite equal phenotypic means.

In analyses of phenotypic data, normal mixture models may reveal the existence of two or more normal distributions with different variances but equal means (Pearson 1894, McLachlan and Peel 2000, Hennig 2010). However, the biological significance of such phenotypic patterns might not be readily evident (Hennig 2010). Traditionally, systematists examining evidence for species limits have emphasized differences between species in phenotypic means, which are a necessary condition for the occurrence of phenotypic gaps (i.e., phenotypic spaces with low frequency of individuals) between species. Therefore, it is reasonable to ask why might differences between species in the variance of phenotypic traits be meaningful despite equal phenotypic means. Here we present a simple numerical example to show that, in the context of the Fisherian model for species delimitation based on phenotypic data we described, differences between species in phenotypic variance may be biologically meaningful and thus relevant for species delimitation.

The example considers diploid organisms in which a polygenic trait, z, is determined by n diallelic loci lacking dominance relationships or epistasis:

$$z = \sum_{i=1}^{n} \gamma_i (X_i + X_i' - 1) \tag{1},$$

where γ_i is the allelic effect at locus i, and X_i and X_i' indicate which allele is present at locus i in two homologous chromosomes, respectively. Thus, $X_i = 1$ if the "+" allele is present in one chromosome and $X_i = 0$ otherwise. Likewise, $X_i' = 1$ if the "+" allele is present in the other chromosome and $X_i' = 0$ otherwise. Therefore, the (genetic) mean and variance of trait z are, respectively:

$$E[z] = \sum_{i=1}^{n} \gamma_i (2p_i - 1) \tag{2},$$

$$var[z] = \sum_{i=1}^{n} \gamma_i^2 p_i (1 - p_i) \tag{3},$$

where p_i is the frequency of the "+" allele in locus i (de Vladar and Barton 2014). We assume eight unlinked loci ($n = 8$) of equal allelic effect ($\gamma_i = 1$ for all i). However, many examples with varying assumptions about number of loci and allelic effects are possible.

764　Imagine two sympatric species, A and B, for which the Fisherian model for species delimitation based on

765　phenotypic data can be reasonably applied (see main text for assumptions of this model). Further imagine

766　that allele frequencies at the eight unlinked loci determining trait z in each species are as follows:

Locus	Species A	Species B
1	$p_1 = 0.5$	$p_1 = 1$
2	$p_2 = 0.5$	$p_2 = 0$
3	$p_3 = 0.5$	$p_3 = 1$
4	$p_4 = 0.5$	$p_4 = 0$
5	$p_5 = 0.9$	$p_5 = 0.9$
6	$p_6 = 1$	$p_6 = 1$
7	$p_7 = 1$	$p_7 = 1$
8	$p_8 = 1$	$p_8 = 1$

768
769

It can be seen, using equations 2 and 3, that the variance of trait z is higher in species A than in species B

(1.09 and 0.09, respectively), despite a common trait mean (3.8 in both species). Thus, the differences in

allele frequencies between species A and B are reflected in the variance of phenotypic trait z, and not in

the mean of trait z. In the context of the Fisherian model for species delimitation based on phenotypic

data, these differences in allele frequencies are biologically meaningful, because they would vanish after a

few generations of random mating (Templeton 2006). In other words, variation in allele frequencies

resulting in equal means but different variances between populations will only persist if such populations

belong to different species; hence, differences in variances that one may detect employing normal mixture

models are evidence supporting the hypothesis that there is more than one species in a sample of

individuals.

780
781　**References**
782

de Vladar, HP, Barton, N. 2014. Stability and response of polygenic traits to stabilizing selection and
mutation. Genetics, 197: 749-767.
Hennig C. 2010. Methods for merging Gaussian mixture components. Advances in Data Analysis and
Classification, 4:3-34.
McLachlan G, Peel DA. 2000. Finite mixture models. Hoboken, NJ, USA, Wiley Series in Probability and
Statistics, John Wiley and Sons.
Pearson K. 1894. Contributions to the theory of mathematical evolution. Philosophical Transactions of the
Royal Society of London A, 185:71-110.
Templeton, AR. 2006. Population genetics and microevolutionary theory. Hoboken, NJ, USA, John Wiley
& Sons.

793
794

795   **Appendix 2.** Morphological measurements and geographic provenance of male specimens of *Geospiza*

796   ground-finches employed in analyses are provided in a separate file **data.csv**. The data are from H. S.

797   Swarth's archive and were employed in this study and made available thanks to permission from the

798   California Academy of Sciences.

800   **Appendix 3.** R code employed to conduct analyses is provided in a separate file **Analysis_code.R**.

801
802

804
805
Supplementary Figure 1. Seven morphological groups of *Geospiza* ground-finches identified by one of the
best normal mixture models. Panels A–F show the groups in the space defined by the four principal
components most useful for group discrimination (PC1–PC4). Colored symbols represent specimens
assigned to different morphological groups and ellipses show 95% high density regions for normal
distributions representing each morphological group. Arrows in G and H display the contribution of
measured morphological traits to each principal component, gauged by the loadings of each trait on each
principal component (i.e., elements of normalized eigenvectors). Circles show the length of arrows
expected if all six traits contributed equally to bidimensional principal component spaces; arrows
exceeding this expectation contribute most significantly and are labeled. PC1 and PC2 reflect general
aspects of beak size and shape (G), with group 2 having long, deep and wide beaks, group 6 having short,
shallow and narrow beaks, and group 7 having long, shallow and narrow beaks. PC3 and PC4 reflect
aspects of tail and tarsus length (H), with group 4 having a relatively long tail, and group 5 having a
relatively long tarsus. PC4 is particularly useful to distinguish group 5 despite explaining only 0.6% of the
total variance. The morphological distribution of groups in the other well supported NMM is fairly similar
to the one shown here, the main difference being that some individuals from groups 1 and 7 are placed
together in an additional group (see Fig. 5 in main text).
822

Current taxonomy (Remsen et al. 2017)

Previous taxonomy (Lack 1947, Rising et al. 2011)

Morphological group

823
824
Supplementary Figure 2. Seven morphological groups of *Geospiza* ground-finches in the Galapagos
Archipelago identified by one of the best normal mixture models partially correspond to the nine species
recognized by current taxonomy (Remsen et al. 2017) and to the six species recognized by previous
taxonomy (Lack 1947, Rising et al. 2011). Each histogram shows, for each recognized species, the
number of specimens assigned to each of the eight morphological groups. Groups are colored according to
the scheme in Supplementary Fig. 1.
831

832
833
Supplementary Figure 3. Seven distinct morphological groups of *Geospiza* ground-finches identified by
one of the best normal mixture models have broad geographic distributions across the Galapagos
Archipelago. For each island, numbers indicate individuals included in the analysis and ringplots depict
the fraction of such individuals assigned to each of the eight morphological groups following the color
scheme in Supplementary Fig. 1. The existence of distinct morphological groups in potential sympatry
within islands (e.g., >4 groups in Santa Cruz, Santiago, and Pinta) suggests that such groups are unlikely
to reflect within-species differentiation due to geographic isolation or local adaptation.
841

842 Supplementary Table 1. Number of islands in the Galapagos Archipelago where each of the seven
843 morphological groups of *Geospiza* ground-finches identified by one of the best NMMs were found to
844 occur (diagonal) and co-occur with other groups (off diagonal). All groups co-occurred with each other in
845 at least one island.

	Group 1	Group 2	Group 3	Group 4	Group 5	Group 6	Group 7
Group 1	9	5	4	2	1	7	5
Group 2		9	4	4	3	7	6
Group 3			11	2	3	10	7
Group 4				4	2	2	2
Group 5					3	3	3
Group 6						14	10
Group 7							10

848