Lethal Haplotypes and Candidate Casual Mutations in Angus Cattle

Jesse L. Hoff[1], Jared E. Decker[1,2], Robert D. Schnabel[1,2] and Jeremy F. Taylor[1,*]

[1] Division of Animal Sciences, University of Missouri, Columbia, Missouri 65211, USA

[2] Informatics Institute, University of Missouri, Columbia, Missouri 65211, USA

*Corresponding author

E-mail addresses:

JLH: jlh4df@mail.missouri.edu

JED: deckerje@missouri.edu

RDS: schnabelr@missouri.edu

JFT: taylorjerr@missouri.edu

Abstract

Background: If unmanaged, high rates of inbreeding in livestock populations

adversely impact their reproductive fitness. In beef cattle, historical selection

strategies have increased the frequency of several segregating fatal autosomal

recessive polymorphisms. Selective breeding has also decreased the extent of

haplotypic diversity genome-wide. By identifying haplotypes for which

homozygotes are not observed but would be expected based on their frequency,

developmentally lethal recessive loci can be localized. This analysis comes without

the need for observation of the loss-associated phenotype (e.g., failure to implant,

first trimester abortion, deformity at birth). In this study, haplotypes were

estimated for 3,961 registered Angus individuals using 52,545 SNP loci using

findhap v2, which exploited the complex pedigree among the individuals in this

population.

Results: Seven loci were detected to possess haplotypes that were not observed in

homozygous form despite a sufficiently high frequency and pedigree-based

expectation of homozygote occurrence. These haplotypes were identified as

candidates for harboring autosomal recessive lethal alleles. Of the genotyped

individuals, 109 were resequenced to an average 27X depth of coverage to identify

putative loss-of-function alleles genome-wide and had variants called using a

custom in-house developed pipeline. For the candidate lethal-harboring haplotypes

present in these bulls, sequence-called genotypes were used to identify concordant

variants. In addition, whole-genome sequence imputation of variants was

performed into the set of 3,961 genotyped animals using the 109 resequenced

animals to identify candidate lethal recessive variants at the seven loci.

Conclusions: Selective breeding programs could utilize the predicted lethal

haplotypes associated with SNP genotypes. Sequencing and other methods for

identifying the causal variants underlying these haplotypes can allow for more

efficient methods of management such as gene editing. These two methods in total

will reduce the negative impacts of inbreeding on fertility and maximize overall

genetic gains.

Keywords: Inbreeding; Autosomal Recessives; Lethal Haplotypes; Phasing;

Imputation

Background

The implementation of a national animal evaluation system in U.S. registered Angus cattle has generated estimates of genetic merit that are used to evaluate and select elite seedstock. Selection on individual traits and indices of traits has resulted in the genetic improvement of multiple traits, such as growth rate, carcass quality and calving ease [1, 2]. At the same time, artificial insemination has increased the utilization of certain paternal lineages. Selective breeding in livestock is known to contribute to the enrichment of deleterious alleles carried by highly utilized sires and also to increase the overall levels of relatedness among individuals. Many numerically important breeds, such as Holstein, Jersey, Nordic Red, and Angus, with extensive use of artificial insemination have recently found autosomal recessive lethal loci at moderate frequencies and that significantly impact fertility [3]. In recent decades several defects that are effectively lethal such as, Neuropathic hydrocephalus, Arthrogryposis Multiplex and Osteoperosis have been propagated within international Angus populations [4, 5]. A striking feature of these defects is their high prevalence in the U.S. registered Angus population, despite their severely deleterious phenotypic presentation [4]. This suggests that it is possible for recessive loci to have major negative impacts on fertility and overall performance without their early detection [3-5]. Consequently, recessive loci causing embryonic loss early in gestation could exist at relatively high population allele frequencies. These loci can reach high frequencies due to drift following severe population bottlenecks, due to their propagation by the extensive use of popular sire lines via artificial insemination, or due to linkage to beneficial alleles at strongly selected loci

[7]. The extent of the impact that these alleles have on fertility and fitness in

livestock is unknown, particularly because most of the reproductive process is

unmonitored. However, when these impacts become large, they can be detected by

genomic analysis of the population. In this study, we examined the inheritance of

haplotypes genome-wide in U.S. registered Angus cattle to reveal candidate

haplotypes that may harbor variants that cause embryonic lethality.

The dynamics of inbreeding depression in closed populations can be

described statistically, but genomic analysis can reveal their biological basis. The

accumulation and impacts of inbreeding are a function of the effective population

size, the initial genetic load of deleterious sites, mating patterns, strength of

selection and the extent or range of linkage disequilibrium within the genome [8].

Our work focuses on sites for which fitness = 0 when homozygous for a deleterious

allele. Consequently, identity by descent at loci harboring loss of function (LOF)

alleles can significantly impact fitness in populations that are accumulating

inbreeding [9]. What is unclear, is how much inbreeding is tolerable and how

common are recessive lethal alleles in livestock. Recent investigations in inbred

human populations suggest that high levels of relatedness are needed to

significantly impact fitness. Counterintuitively, up to a certain high threshold,

parental relatedness appears to be beneficial for fitness [10]. An investigation into

the North American human Hutterite population estimated that an average of 0.29

recessive lethal variants exist per haploid human genome, and that the primary

force that removes these alleles from a population is drift [11]. Simulations by these

authors also revealed that the majority (57.4%) of recessive deleterious variants

that were segregating in the founder population were not observed in the modern

population. Of the remaining recessive variants, only 8.23% had phenotypic effects.

This result is important, because it suggests that in a population such as the U.S.

registered Angus breed, a high proportion of latent recessive lethal variants could

still be segregating without their having been detected by breeders.

A recent study in U.S. Holstein dairy cattle using genotyped trios found that

expected inbreeding coefficients of offspring (obtained by simulated matings of

actual parental genotypes) were slightly lower than the realized inbreeding

coefficients, suggesting that increased inbreeding was not a constraint on viability of

offspring [12]. This result, along with the knowledge of the nature of several existing

recessive defects in the U.S. Holstein population suggests that the frequency of

lethals may not be sufficiently high to produce an observable impact on the

population when the average inbreeding coefficient is only 3.53%. These findings

are also consistent with a recent study of dog domestication and breed formation

that found that the major factor underlying the enrichment of deleterious variation

in modern breed dogs was severe population bottlenecks, and not recent inbreeding

[13].

Given experiences in other breeds and species, we hypothesize that there

may be many more recessive lethal alleles in the U.S. registered Angus population

than have previously been detected. To identify these, we implemented a method

that was first described by VanRaden [14]. A sample of 3,961 genotyped Angus

cattle, primarily bulls extensively used in artificial insemination that were members

of a pedigree containing 117,212 identified individuals, was analyzed to identify

haplotypes that were expected to be observed but that were not actually observed

in the homozygous state. The pedigree spanned more than 60 generations with the

earliest ancestor born in 1836, and with the earliest genotyped animal born in 1955

[2]. The analysis assumed that the haplotypes harbored fully penetrant lethal alleles

that would preclude the viability and genotyping of an animal homozygous for the

haplotype. Within this pedigree, haplotypes that were identical by descent were

repeatedly sampled, allowing for the detection of autosomal recessive lethal alleles.

The adoption of high-density array-based genotyping in commercial beef and

dairy cattle populations is rapidly increasing, due to the utility of genomic selection

[15] and as a consequence a large number of trios, patrios (sire, maternal grandsire

and son) and half-sib families have now been genotyped. Whole-genome sequencing

of influential population members and the use of genotype imputation will allow the

identification of lethal alleles without the observation of the phenotype responsible

for the loss [16]. Characterizing the number and identity of these variants will

provide a deeper understanding of the biological and quantitative underpinnings of

inbreeding depression. It will also enable enhanced management of animal

reproduction, as these variants can be identified in any genotyped animal.

Methods

Genotypes and Animals

BovineSNP50 BeadChip (Illumina, San Diego, CA)[17] data for 3,993 registered

Angus animals born between 1955 and 2012 and representing 63 generations were

available for analysis. Genotypes had been filtered to retain data with a call rate of ≥

90% and minor allele frequency (MAF) ≥ 0.01 [2]. Pedigrees had also been validated

by homozygous transmission incompatibility rates estimated as the frequency of

loci for which the parent and offspring were alternate homozygotes. Pedigree

relationships were expunged when the incompatibility rate exceeded 1.5%, and

individuals who did not match their recorded parents or offspring were removed

from the pedigree. After filtering, 52,545 loci and 3,961 animals remained. These

sites were next phased and missing genotypes were imputed using findhap (v2)

based upon a combination of simple haplotype frequency sorting and the use of all

available pedigree information [18].

Population and Pedigree Based Haplotype Analyses

Haplotypes were defined by 20-marker sliding windows genome-wide. Evidence for

a lethal allele within a haplotype was evaluated using two statistics. The first was

based on population frequency. For each haplotype in each window, frequency was

estimated and the number of individuals in the population that were homozygous

for the haplotype was tallied. When there was an absence of homozygotes, the

likelihood of this occurrence was calculated under a model that assumed carriers to

be randomly mating and selective neutrality of the haplotype.

In the second analysis, the actual matings within the pedigree were used to

estimate the expected number of homozygotes for any given haplotype based upon

the number of matings involving carriers and the assumption of selective neutrality.

The pedigree for the genotyped animals was parsed to identify patrios, defined here

as families for which genotypes were available for the offspring, sire and maternal

grandsire. Maternal genotypes were rarely available as DNA was primarily sourced

from cryopreserved semen [2]. This family structure allowed the testing of

segregation distortion when both the sire and maternal grandsire were

heterozygous for the same haplotype. For each sliding window of 20 markers if a

haplotype was never observed in the homozygous state and had a sample frequency

of greater than 2%, the number of families for which the sire and maternal

grandsire were both heterozygous was counted. The probability of observing at

least one homozygote was then calculated based on the count of patrios and the

assumption of selective neutrality. The probability that no homozygous *hh*

haplotypes are observed in the progeny of C patrios when both the sire and

maternal grand-sire are *Hh* heterozygotes and the *h* haplotype does not harbor a

selected deleterious allele is $(0.875 - 0.25q)^C$ where q is the frequency of the *h*

haplotype in the population. There were 2,480 patrios represented in the sample.

Regions of the genome with an identified deficit of homozygotes for a specific

haplotype were examined for underlying genes using pybedtools and the UMD3.1

genome assembly annotation run 104 [14, 15].

Generation of Sequence Data

To further analyze the variation within the genomic regions harboring haplotypes

that were deficient for homozygotes, the whole-genome sequences of 109 animals

from this population were examined [20]. These animals were selected for

sequencing based upon their impact on the breed assessed by the expected numbers

of genome equivalents (progeny have 0.5, grandprogeny have 0.25, etc) present

within registered descendants in the population. They were sequenced with paired-

end 2 x 100 bp sequence reads to an average of 27X depth of coverage of the

UMD3.1 assembly with Illumina Genome Analyzer, GAII, HiSeq 2000 or 2500

instruments from two libraries with 350 bp and 550 bp average fragment sizes.

FastQC was used to analyze the quality of the reads [21]. Exact duplicates were

removed, and adapters were trimmed using a custom in-house Perl script. All

remaining reads were error corrected using QuorUM [22]. Newly created duplicates

(due to the trimming of low quality ends and correction of errors) and reads shorter

than 35 bp were removed and the final data set was aligned to the UMD3.1

reference genome assembly using NextGENe 2.4.1 (SoftGenetics, LLC, State College,

PA) alignment software. Reads were required to have a matching segment at least

35 bases long and 95% overall match, and a maximum of 2 bases of mismatch across

the whole alignment. Up to 1000 alignments of equal likelihood were allowed

genome-wide. NextGene 2.4.1 was also used for variant calling.

The sequence-derived variants for the 109 bulls were used for two purposes;

identification of variants shared amongst animals identified as carriers for the

putatively lethal BovineSNP50 20 SNP haplotypes, and for the imputation of the

entire genotyped population to whole-genome sequence variation.

Examining Carrier Sequence Data

Within the genomic regions identified from the marker data as containing candidate

lethal haplotypes, all variants identified in our sequenced sample were analyzed for

carrier concordance, implemented using python scripts. This involved examination

of variants that were observed as heterozygous in all bulls predicted to be carriers

of the homozygote deficient haplotype. The sample size for the sequenced animals

was not sufficient to run the population allele frequency or patrio analyses.

Imputation of BovineSNP50 Genotypes to Whole Genome Sequence Variation

We imputed the BovineSNP50 genotypes for this population to whole-genome

sequence level variation in order to identify potential lethal variants, using the 109

sequenced animals as the reference population. We selected 24,974,785 SNPs from

the full set that had been identified in these animals. We first included SNPs found in

the 109 sequenced Angus bulls that were biallelic and located within gene

boundaries (777,432), UTRs (33,379) or that were splice site variants (636,492).

These variants spanned the allele frequency spectrum but were identified at high

sequence coverage. To enable imputation we also included 23,527,482 variants that

had been independently identified in run 5 of the 1000 Bull Genomes project [23].

These variants represent filtered, high quality segregating sites identified from the

whole genome sequences of 1,578 animals from multiple taurine breeds including

Angus. Genotypes called for the 24,974,785 variable sites genome-wide in the 109

registered Angus bulls were used as the reference set for whole-genome sequence

imputation of the BovineSNP50 data using Fimpute [24].

224   The imputed genotypes were individually analyzed for the absence of

homozygotes for alleles present within each of the candidate haplotyped loci using

the frequency and pedigree approaches. We first identified high frequency variants

for which no homozygous individuals were predicted. The pedigree analysis was

also performed for all candidate variants identified in the frequency analysis.

Variants identified by either of these processes were characterized using the variant

effect predictor release 79 [25].

Results

Identification of Putative Lethal Haplotypes

We identified seven haplotypes with a pattern of inheritance in U.S. registered

Angus cattle that suggests that they each harbor an autosomal recessive lethal allele

(Table 1). Using a binomial distribution for the number of observed homozygotes in

the progeny of C patrios, we calculated the probability of observing no homozygotes

when each haplotype was selectively neutral. We selected haplotypes as putatively

harboring autosomal recessive lethals when the probability of observing no

homozygotes was less than 0.02. This threshold for statistical significance provides

considerable confidence that the lack of homozygosity for these haplotypes did not

occur by chance alone.

Sensitivity to Window Size

We evaluated the sensitivity of identification of these marker-based haplotypes to

window size and concluded that a window size of 20 contiguous BovineSNP50

markers was appropriate for capturing the haplotypic diversity within the

population (Figure 1). This window size appears to discriminate between

haplotypes that are identical by descent (IBD) and those that are identical by state

(IBS). Our analysis shows that the number of common haplotypes detected rises as

window size increases, but begins to plateau at 20 markers. Considering the

moderate marker density of the BovineSNP50 (1 SNP per 50 kb), haplotypes that

are defined by only two markers are assumed to be IBS for the purposes of analysis

but likely actually represent a number of distinct haplotypes at the level of genome

sequence. As the window size increases, the likelihood increases that two

haplotypes found in different individuals that are IBS are also IBD and are thus

concordant at the level of sequence variation. However, with large window sizes,

recombination may lead to a lethal variant being present on more than one

haplotype, thus decreasing the power of the analysis. Indeed, as window size

increases, the overall rate of individuals homozygous for any haplotype declined.

The window size selected for this study appears to achieve an appropriate balance

of genome-wide homozygosity and rate of occurrence of high frequency haplotypes

(Figure 1). Slight changes in the haplotype window size did not affect the detection

of the 7 haplotypes reported in Table 1.

We were also able to validate IBD status of these regions by an examination

of the sequence data generated for animals that were predicted to be carriers of

identical haplotypes. For each of the loci predicted to harbor recessive lethal

haplotypes (Table 1), we identified the bulls among the 109 sequenced animals that

were predicted to be carriers of each putative lethal haplotype and computed the

pairwise rate of opposing homozygous sequenced sites between all pairs of carrier

animals. For example, for the haplotype on chromosome 29, with 16 predicted

carriers we made 120 pairwise comparisons using all sequence called variant

genotypes within the haplotype coordinates (Figure 2). Amongst individuals that

share a common haplotype, the alternate homozygote rate is related to the error

rate for sequence-based genotype calls for heterozygotes. The rate observed in our

predicted carriers within our 6 testable haplotype regions was similar to the rate for

sire-son pairs. One region (Chr 1) had only one sequenced predicted carrier and

could not be evaluated. The majority of the carrier pair comparisons with rates of

opposing homozygous sequenced sites >0.01 were caused by the inclusion in the

analysis of 3 individuals that had been sequenced to averages depths of < 10X.

Within the haplotype, opposing homozygous sequenced site rates were markedly

lower for predicted carriers than for randomly selected individuals. When predicted

carriers for a lethal haplotype were analyzed for a randomly selected 20 marker

region elsewhere in the genome, they had opposing homozygote rates that were

similar to those of the unrelated sire pairs. This indicates that the haplotypes

generated from the BovineSNP50 data successfully identified genomic regions that

were IBD at the level of the genome sequence. We also observed that this method is

sensitive to the depth of sequence coverage due to the inaccurate identification of

heterozygous loci as being homozygous when the alternate allele was never

sequenced. In total, 16 animals had depths of sequence coverage of < 10X and most

of the remaining animals (70) had > 20X. For a Sire-Son pair, low sequence coverage

for at least one member of the pair led to a rate of homozygous inconsistency that

was increased by an order of magnitude.

Analysis of Sequence Variation and Candidate Genes in Bulls Predicted to be Lethal

Haplotype Carriers

To identify the causal variants underlying these putatively lethal haplotypes, we

first directly examined resequencing data from bulls that were predicted to be

carriers of the lethal haplotypes. Among the 109 sequenced bulls up to 21 animals

were predicted to be carriers of each of the BovineSNP50 putatively lethal

haplotypes reported in Table 1. Within these seven genomic windows, we identified

candidate variants that were never homozygous in the 109 sequenced bulls.

However, none were observed to be exclusively heterozygous in the predicted

carrier animals. That is, all of the alleles found to be heterozygous in all of the

predicted carriers were also observed to be heterozygous in animals that did not

carry the predicted lethal haplotype. A recessive lethal variant could be

heterozygous in animals that were not carriers of the putatively lethal haplotype if

the mutation is sufficiently old that recombination has occurred relative to the

haplotype on which the mutation occurred. Table 1 reports, for each genomic region

predicted to harbor a lethal haplotype, the number of variants that were

heterozygous in all predicted haplotype carriers and that were never found to be

homozygous in any of the 109 resequenced bulls. For instance, on chromosome 15,

an intronic variant in *SSRP1* was found in 9 of the 10 sequenced bulls that were

predicted to carry the putatively lethal haplotype, and in 25 of all 109 sequenced

Angus bulls, but was never found to be homozygous. Although the variant is

intronic, with no expected impact on splice site variation, it resides in an interesting

candidate gene as *SSRP1* is essential for mouse embryonic development [26].

This analysis was also conducted after excluding the 16 animals sequenced to

low average sequence depth of coverage (< 10X). Genotypes for these animals are

likely to contain false positive homozygotes at many true heterozygous sites. This

resulted in the detection of additional concordant loci but again none were exclusive

to the predicted carriers. The chromosome 29 locus had 12 additional variants

identified that were located in genes such as *CAPN*, *NRXN2*, *PACS1* and *PP25B*.

However, none of these mutations are in exons or had other interpretations in

Variant Effect Predictor (VEP) and all 12 were heterozygous in 18 animals, including

4 that were not predicted by their marker data to be carriers. Thus, the region

appears to comprise a large consistent haplotype with no one particular variant

being simply implicated as causal for lethality. The locus on chromosome 4 had 118

variants identified as being heterozygous in haplotype carriers following the

removal of the low sequence coverage animals, but none were predicted to alter

protein amino acid sequences.

The locus on chromosome 1 for which we predicted a lethal haplotype

contains only one gene, *GBE1* that encodes a protein that catalyzes the branching of

glycogen, the main form of energy storage in the body [24]. There was only one

sequenced animal in our population predicted to carry this disease and further

analysis of the unique heterozygotes within this gene were therefore not feasible.

Analysis of Imputed Sequence Variation in 3,961 Registered Angus

When analyzed in the 3,961 animals, 2,504 of the 24,974,785 imputed variants had

a MAF ≥ 2%, no predicted homozygotes, and were found in ≥ 30 double-carrier

patrios. Of the 24,974,785 variants, 147,764 had a deleterious consequence

predicted by their SIFT scores [30], but only 4 were among the 2,504 candidate

autosomal lethal alleles. Two of these sites were in *LOC521645*, an olfactory

receptor gene (near, but not within the chromosome 15 locus, Table 1), and another

was in *LOC100336589* on chromosome 18. The fourth was on chromosome 21 in

GEMIN2, for which a recessive lethal embryonic phenotype occurs in mouse

knockouts and is associated with the survival motor neuron complex [31]. This

allele had a frequency of 7.8% following imputation into the 3,961 animals and was

observed in 55 double-carrier patrios without producing a homozygous progeny.

Twelve variants within *GBE1*, none of which were predicted to be

deleterious, were at a MAF ≥ 2%, had no homozygotes predicted, and were found in

≥ 30 double-carrier patrios, with one variant present in 133 double-carrier patrios.

However, only one of the sequenced animals was predicted to carry this haplotype

making it difficult to use the data for the sequenced animals to exclude any

heterozygous sites within this animal's diplotype from candidacy for lethality. The

haplotype on chromosome 1 spanning *GBE1* was the only predicted lethal to contain

any of the 2,504 candidate variants identified in the analysis of the imputed whole

genome sequence data.

Discussion

Due to the repeated sampling of parental gametes and the sharing of haplotypes

across families in this extensive Angus pedigree, we were able to powerfully test the

fitness consequence of many haplotypes. The use of artificial insemination increases

selection intensity by allowing relatively few individuals to sire large numbers of

progeny and this population has historically been strongly selected for growth and

calving ease [2]. Four bulls in this dataset were each represented in over 100 of the

2,480 genotyped patrios, suggesting that intense selection can drive deleterious

alleles carried by these bulls to a relatively high frequency. In some cattle

populations, allele frequencies of up to 20% have been observed for some recessive

lethal haplotypes [6]. These common recessive lethals have the greatest impact on

population fitness, but as genotyping becomes more pervasive, rare recessive lethal

haplotypes will be detected and the available genotypes should allow for their

management.

Management Strategies

Angus breeders have historically selected against recessive lethal alleles which

manifest as fatal calf defects in this population [4]. Registered animals are tested for

known defects and registration is prevented for young animals that test to be

carriers. As new deleterious alleles including those causing early embryonic loss are

discovered, this approach is likely to be untenable. The seven new putatively

autosomal recessive lethal haplotypes may now be predicted in hundreds of

thousands of genotyped animals [32] and many more deleterious loci may be

discovered as the number of genotyped animals increases and new lineages rise to

prominence that contain yet to be detected recessive alleles. Management must shift

from registration exclusion to a means of incorporating marker diagnostics into

genomic selection and mate selection protocols.

A mate selection procedure has been implemented in the MateSel software

that applies a linear weight against either the number of recessive alleles present in

the progeny generation (LethalA) or the number of homozygous progeny (LethalG)

[33]. In a simulation with a higher count (N=100) for the number of lethal alleles

segregating in a population than identified in this study, it was not possible to

sufficiently weight in either selection scheme to eliminate embryonic mortality. This

scenario may approach reality as the number of identified deleterious recessive

alleles causing calf defects as well as embryonic loss increases. These simulations

also suggest that implementing the LethalG strategy is a more efficient means of

achieving genetic gain while reducing the frequency of recessive alleles.

Cole (2015) suggested an alternative strategy in which parent average

breeding indexes are adjusted for lost progeny by adding a cost associated with

recessive haplotypes [34]. This method could be extended to account for the joint

impact of multiple linked or unlinked segregating loci to reduce the frequency of

deleterious alleles. Counterintuitively, this study also found a zero to inverse

relationship between the embryo's realized inbreeding coefficient and its

probability of being homozygous for an allele responsible for a recessive disorder.

This suggests that the goal of reducing the long-term rate of accumulation of

inbreeding in breeding programs may not impact the rate of embryonic loss due to

the action of recessive lethal alleles. This is consistent with the observation from

studies of embryonic loss and coancestry in humans [8].

Managing False Positives

Managing selection based on these results requires certainty about the lethal

haplotype's effect. In this study, a total of 12,020 haplotypes were found genome-

wide (including partially overlapping haplotypes) that occurred at a MAF ≥ 2% but

were never found as homozygotes. Assuming random mating, a haplotype at this

frequency is expected to have, on average, only 1.5 homozygotes in our Angus

sample. However, the existence of non-random mating within this population could

substantially decrease the likelihood of observing homozygotes, and explain why so

many of these regions were observed. The patrio analysis directly incorporates the

matings that created the population to detect deviations from selective neutrality in

progeny genotypes. However, this approach is limited by the availability of

genotyped patrios. In breeds where insemination and pregnancy records are more

detailed, these have been crucial for validating putative lethal haplotypes[14].

In the absence of these data, both the MateSel approach and the index

adjustment could be adapted to account for the uncertainty of the lethality of the

predicted allele or haplotype. In addition to validated haplotypes, and the high

confidence haplotypes that we observe here, this approach could enable the

incorporation of haplotypes into the selection scheme that appear to be lethal but

that are at low frequency in the population. There are now Angus pedigrees in

which hundreds of thousands of animals have been genotyped world-wide and an

analysis of these data would likely improve the resolution of lethal haplotype

detection and could also identify many rarer variants [15]. If these lethal alleles are

individually rare but each individual carries many of them, recessive lethals could

affect a substantial portion of pregnancies.

False Negatives

Even with adequate sample sizes to ensure statistical power, there are limitations to

the methods that we have employed. The approaches employed will only detect

recessive alleles that are perfectly concordant with a BovineSNP50 haplotype. If a

recent autosomal recessive lethal mutation has occurred on a common haplotype,

the population will comprise haplotypes harboring either the lethal mutation or the

wild type allele and homozygotes that include the wild type allele will be observed.

In this case, very large sample sizes are required to detect deviations from the

number of homozygotes expected under Hardy-Weinberg equilibrium. This could

also involve restricting the analysis to different pedigree lineages for an IBS

haplotype. This may explain why we failed to identify any of the recent recessive

genetic defects found in the Angus breed [5]. In recent years, the alleles responsible

for these defects have frequencies that have ranged from 3-9% [4]. However, the

regions of the genome that harbor these alleles were not detected in the marker-

based haplotype analysis and the causal variants were not detected in the analysis

of the imputed sequence data. The likely cause of this is that the haplotypes we

examined did not always indicate the presence of the defective allele. For instance,

the mutation causing Neuropathic Hydrocephalus, originated in bull G A R Precision

1680 born in 1990 [35]. Consequently, there are both wild-type and deleterious

versions of the BovineSNP50 haplotype on which the mutation arose segregating in

the genotyped population. When larger sample sizes become available, it would be

useful to repeat these analyses and test for homozygote deficiency rather than

complete absence.

We have also not attempted to model mutations in loci with parent of origin

affects, such as imprinting associated defects [36]. SNP array genotypes identify

large heterozygous deletions as being homozygous for the alleles present on the

non-deletion chromosome, which may prevent the identification of carriers for the

deletion. This type of mutation has previously been associated with lethal recessive

diseases in cattle [7]. Additionally, lethal alleles with incomplete penetrance will not

be captured by our analyses. Other errors in genotyping, phasing or imputation also

likely contribute to reductions in power to detect lethal haplotypes.

Using sequence data

One potential solution to the limitations of array genotype data is to analyze

sequence-derived and/or imputed genotypes. These data may help with the

management of putative recessive lethal alleles. True causal variants can be tracked

more effectively than marker haplotypes. When the causative alleles have been

identified, gene editing may also present an efficient means of reducing the genetic

load of elite sires in a manner that is complementary to the current breeding system

[37].

Sequence data also has potential advantages for the detection of recessive

loci. If appropriately processed to capture SNPs, large and small indels and

structural variants, they directly represent the pool of all recessive deleterious

alleles. However, in practice, sample sizes have been small and the identification of

large indels, particularly insertions, and complex structural variants has been

challenging. Our analysis of sequence data failed to identify any candidate causative

mutations in the marker-based haplotypes that were predicted to be lethal. The

sample size for the sequenced animals was not sufficient to conduct the frequency

or pedigree analysis with the genome-wide sequence variants. Furthermore, we did

not attempt to analyze many types of complex variation, such as large indels or

structural variants [38]. Large structural variants are enriched for deleterious

variation but can be complex to analyze with short read data [39]. Alternative

analyses of the sequence data to identify these variants or the use of methods which

generate longer reads may be necessary to capture the causative variants.

In this study we did not detect a particular variant within a putative

haplotype that was likely to cause a recessive lethal phenotype. However, the gene

within the region on Chromosome 1, *GBE1* appears quite promising. Mutations in

this gene produce recessive phenotypes in mammals including horse, mouse and

human [24, 25]. In the U.S. Quarter Horse population, phenotypes created by

homozygotes for *GBE1* mutations ranged from stillbirth to early failure to thrive,

with death never occurring later than 18 weeks of age [28]. Mouse knockout

analysis revealed few visible or biochemical phenotypic effects in heterozygotes.

Monitoring of embryonic development in homozygous knockouts revealed that

deformities only occurred late in gestation and led to stillbirth or death shortly after

birth. Mice with a construct with low GBE1 activity incorporated into their genome

to replace the wild type allele demonstrated poor metabolic performance, and the

accumulation of polyglucosan [29]. None of the mice with limited GBE1 function

lived beyond 39 weeks, while all control mice survived the trial. While not all of the

recessive *GBE1* genotypes in other species have resulted in embryonic loss, the

reduced growth associated with homozygosity for these mutations makes it unlikely

that an affected animal would be selected as a sire or dam. They would therefore be

highly unlikely to be included in our genotyped sample of Angus cattle. However,

identifying homozygous calves from those produced by mating carriers and

assaying their GBE1 functionality might be possible.

Mapping Candidate Variants without Sequencing

Rather than generating expensive sequence data for identifying recessive lethals,

two strategies might be useful: assay development and imputation. Novel variants of

all classes that are detected by sequencing can readily be incorporated onto

commercial genotyping platforms. This expedites fine-mapping within a known

lethal haplotype in a commercial population. Variants that had predicted deleterious

functional impacts based on bioinformatic analysis would also be excellent

candidates for inclusion on commercial genotyping platforms.

Imputation accuracy, particularly for rare variants, may not be sufficient to

identify candidate segregating recessive lethals. We previously analyzed the

accuracy of imputation using variants from Run4 of the 1000 bull genomes project

as a reference for our Angus BovineSNP50 genotyped population using the same

imputation methods that were used in this study [20]. Our 109 sequenced animals

had their BovineSNP50 genotypes imputed to the 1000 Bull Genomes Run4

sequence reference set as well as variants called directly from their whole-genome

sequences. Comparing these two sets of genotypes revealed correlations between

genotypes in the range of 80-90% for common variants, and 60-80% for variants

with MAF ≤ 10% [20]. We would expect the causal variants underlying these lethal

haplotypes to fall in the rare, more inaccurately imputed frequency class. This

greatly complicates the utility of imputation for this application, and the results of

imputation presented here are not appropriate for application to breeding

decisions. Had an interesting candidate locus with a plausible biological mechanism

emerged, it would have been a good target for further confirmation and possibly

immediate use. More sophisticated sequence imputation methods that provide

higher imputation accuracies across the allele frequency spectrum are now

becoming available. These have been used in human studies to identify individuals

that are homozygous for rare variants with predicted deleterious effects [16].

Applying these methods to livestock populations with extensively described and

genotyped pedigrees is feasible, and may prove useful for fine-mapping within

candidate haplotypes.

Conclusions

We identified 7 potentially recessive lethal haplotypes segregating in the U.S.

registered Angus population that open opportunities for improving breeding

success and increasing the mean fitness of the population. These haplotypes have

been propagated throughout Angus lineages represented in a set of 3,961 genotyped

animals and were not observed in homozygous form. The phenotypic effects of

these haplotypes have not been directly observed, but may be inconspicuous such as

in the event of early embryonic loss or may be unreported defects leading to the loss

of the calf. Efforts to identify causal mutations with a clear molecular impact from

sequence data were unsuccessful but interesting candidate genes such as *GBE1* were

identified. Further validation of the impact of these haplotypes on fertility and the

direct observation of calves that are homozygous for these haplotypes could reveal

interesting biology. Our capacity to detect these loci will continue to improve as

increasingly large numbers of animals are genotyped and sequenced. The quality of

the reference genome assembly and methods for characterizing and imputing

structural variants are also improving and will improve the quality of this type of

analysis. Eventually, we will identify dozens of deleterious recessive loci, and can

use chip-based genotyping to manage matings, track alleles through lineages and

potentially use gene editing to remove them from elite animals.

Declarations

Ethics approval and consent to participate

The genotype data described in this manuscript was previously analyzed or

collected from commercially generated animal semen; as such no ethics or animal

welfare approval was required.

Consent for publication

Not applicable

Availability of data and material

Genotypes are available to scientists interested in non-commercial research upon

signing a Materials Transfer Agreement (MTA). All sequence data will be deposited

under NCBI Bioproject Accession PRJNA343262.

Funding

This project was supported by National Research Initiative grants number 2008-

35205-04687 and 2008-35205-18864 from the USDA Cooperative State Research,

Education and Extension Service and National Research Initiative grants number

2009-65205-05635 and 2013-68004-20364 from the USDA National Institute of

Food and Agriculture.

Authors' contributions

JLH JED and RDS JFT designed and conducted the study. JFT and RDS collected

samples and processed the genotypes. RDS processed NGS data. JLH conducted

bioinformatic analyses of genotypes and sequence variants. JLH and JFT wrote the

manuscript. JFT RDS and JED edited the manuscript.

Acknowledgements

We gratefully acknowledge the provision of semen samples from Angus breeders

and semen distributors. We appreciate the financial support of the American Angus

Association, the Australian Angus Association, The New Zealand Angus Association

and the Argentine Angus Association to sequence the genomes of bulls used in this

study.

References

1. **Angus Genetic Trends** [http://www.angus.org/nce/genetictrends.aspx]

2. Decker JE, Vasco DA, McKay SD, McClure MC, Rolf MM, Kim J, Northcutt SL, Bauck S, Woodward BW, Schnabel RD, Taylor JF: **A novel analytical method, Birth Date Selection Mapping, detects response of the Angus (Bos taurus) genome to selection on complex traits**. *BMC Genomics* 2012, **13**:606.

3. Fritz S, Capitan A, Djari A, Rodriguez SC, Barbat A, Baur A, Grohs C, Weiss B, Boussaha M, Esquerré D, Klopp C, Rocha D, Boichard D: **Detection of Haplotypes Associated with Prenatal Death in Dairy Cattle and Identification of Deleterious Mutations in GART, SHBG and SLC37A2**. *PLoS One* 2013, **8**:e65550.

4. Teseling CF, Parnell P.: **How Angus breeders have reduced the frequency of deleterious recessive genetic conditions**. In *Association of Advancement Animal Breeding and Genetics*. AAABG; 2013:20:558-561.

5. American Angus Association. **Genetic Conditions Policy** 2017 https://www.angus.org/pub/GeneticConditionPolicy.aspx Accessed 4 April 2017

6. Sonstegard TS, Cole JB, VanRaden PM, Van Tassell CP, Null DJ, Schroeder SG, Bickhart D, McClure MC: **Identification of a nonsense mutation in CWC15 associated with decreased reproductive efficiency in Jersey cattle.** *PLoS One* 2013, **8**:e54872.

7. Kadri NK, Sahana G, Charlier C, Iso-Touru T, Guldbrandtsen B, Karim L, Nielsen US, Panitz F, Aamand GP, Schulman N, Georges M, Vilkki J, Lund MS, Druet T: **A 660-Kb deletion with antagonistic effects on fertility and milk production segregates at high frequency in Nordic Red cattle: additional evidence for the common occurrence of balancing selection in livestock.** *PLoS Genet* 2014, **10**:e1004049.

8. Charlesworth D, Morgan MT, Charlesworth B: **The effect of linkage and population size on inbreeding depression due to mutational load**. *Genet Res* 2009, **59**:49.

9. Falconer D, Mackay TFC: *Introduction to Quantitative Genetics*. 4th edition. New York: Wiley; 1996.

10. Helgason a., Palsson S, Guthbjartsson DF, Kristjansson T, Stefansson K: **An Association Between the Kinship and Fertility of Human Couples**. *Science (80-)* 2008, **319**:813–816.

11. Gao Z, Waggoner D, Stephens M, Ober C, Przeworski M: **An estimate of the average number of recessive lethal mutations carried by humans.** *Genetics* 2015, **199**:1243–54.

12. Bjelland DW, Weigel KA, Coburn AD, Wilson RD: **Using a family-based structure to detect the effects of genomic inbreeding on embryo viability in Holstein cattle.** *J Dairy Sci* 2015, **98**:4934–44.

13. Marsden CD, Ortega-Del Vecchyo D, O'Brien DP, Taylor JF, Ramirez O, Vilà C,
Marques-Bonet T, Schnabel RD, Wayne RK, Lohmueller KE: **Bottlenecks and**
selective sweeps during domestication have increased deleterious genetic
variation in dogs. *Proc Natl Acad Sci U S A* 2015, **113**:152–157.

14. VanRaden PM, Olson KM, Null DJ, Hutchison JL: **Harmful recessive effects on**
fertility detected by absence of homozygous haplotypes. *J Dairy Sci* 2011,
94:6153–61.

15. Decker JE: **Agricultural Genomics: Commercial Applications Bring Increased**
Basic Research Power. *PLoS Genet* 2015, **11**:e1005621.

16. Sulem P, Helgason H, Oddson A, Stefansson H, Gudjonsson SA, Zink F, Hjartarson
E, Sigurdsson GT, Jonasdottir A, Jonasdottir A, Sigurdsson A, Magnusson OT, Kong A,
Helgason A, Holm H, Thorsteinsdottir U, Masson G, Gudbjartsson DF, Stefansson K:
Identification of a large set of rare complete human knockouts. *Nat Genet* 2015,
advance on.

17. Matukumalli LK, Lawley CT, Schnabel RD, Taylor JF, Allan MF, Heaton MP,
O'Connell J, Moore SS, Smith TPL, Sonstegard TS, Van Tassell CP: **Development and**
characterization of a high density SNP genotyping assay for cattle. *PLoS One*
2009, **4**:e5350.

18. VanRaden PM, O'Connell JR, Wiggans GR, Weigel KA: **Genomic evaluations with**
many more genotypes. *Genet Sel Evol* 2011, **43**:10.

19. Dale RK, Pedersen BS, Quinlan AR: **Pybedtools: a flexible Python library for**
manipulating genomic datasets and annotations. *Bioinformatics* 2011, **27**:3423–
4.

20. Taylor JF, LK Whitacre, JL Hoff, PC Tizioto, JW Kim JD and R: **Lessons from**
cattle genome and transcriptome sequencing. *Genet Sel Evol* .

21. Andrews S: **FastQC**. *A Qual Control tool high throughput Seq data.[h ttp//www*
bioinformatics bbsrc ac uk/projects/fastqc/] 2010.

22. Marçais G, Yorke JA, Zimin A: **QuorUM: An Error Corrector for Illumina**
Reads. *PLoS One* 2015, **10**:e0130821.

23. Daetwyler HD, Capitan A, Pausch H, Stothard P, van Binsbergen R, Brøndum RF,
Liao X, Djari A, Rodriguez SC, Grohs C, Esquerré D, Bouchez O, Rossignol M-N, Klopp
C, Rocha D, Fritz S, Eggen A, Bowman PJ, Coote D, Chamberlain AJ, Anderson C,
VanTassell CP, Hulsegge I, Goddard ME, Guldbrandtsen B, Lund MS, Veerkamp RF,
Boichard DA, Fries R, Hayes BJ: **Whole-genome sequencing of 234 bulls**
facilitates mapping of monogenic and complex traits in cattle. *Nat Genet* 2014,
46:858–865.

24. Sargolzaei M, Chesnais JP, Schenkel FS: **A new approach for efficient genotype**
imputation using information from relatives. *BMC Genomics* 2014, **15**:478.

25. McLaren W, Pritchard B, Rios D, Chen Y, Flicek P, Cunningham F: **Deriving the**
consequences of genomic variants with the Ensembl API and SNP Effect

Predictor. *Bioinformatics* 2010, **26**:2069–70.

26. Cao S, Bendall H, Hicks GG, Nashabi A, Sakano H, Shinkai Y, Gariglio M, Oltz EM,
Ruley HE: **The high-mobility-group box protein SSRP1/T160 is essential for**
cell viability in day 3.5 mouse embryos. *Mol Cell Biol* 2003, **23**:5301–7.

27. Wagner ML, Valberg SJ, Ames EG, Bauer MM, Wiseman JA, Penedo MCT, Kinde H,
Abbitt B, Mickelson JR: **Allele Frequency and Likely Impact of the Glycogen**
Branching Enzyme Deficiency Gene in Quarter Horse and Paint Horse
Populations. *J Vet Intern Med* 2006, **20**:1207–1211.

28. Ward T, Valberg S, Adelson D, Abbey C, Binns M, Mickelson J: **Glycogen**
branching enzyme (GBE1) mutation causing equine glycogen storage disease
IV. *Mamm Genome* 2004, **15**.

29. Akman HO, Sheiko T, Tay SKH, Finegold MJ, Dimauro S, Craigen WJ: **Generation**
of a novel mouse model that recapitulates early and adult onset glycogenosis
type IV. *Hum Mol Genet* 2011, **20**:4430–9.

30. Kumar P, Henikoff S, Ng PC: **Predicting the effects of coding non-synonymous**
variants on protein function using the SIFT algorithm. *Nat Protoc* 2009, **4**:1073–
81.

31. Jablonka S, Holtmann B, Meister G, Bandilla M, Rossoll W, Fischer U, Sendtner M:
Gene targeting of Gemin2 in mice reveals a correlation between defects in the
biogenesis of U snRNPs and motoneuron cell death. *Proc Natl Acad Sci U S A*
2002, **99**:10126–31.

32. Rolf MM, Decker JE, McKay SD, Tizioto PC, Branham KA, Whitacre LK, Hoff JL,
Regitano LCA, Taylor JF: **Genomics in the United States beef industry**. *Livest Sci*
2014, **166**:84–93.

33. Eenennaam AL Van, Kinghorn BP: **Use of Mate Selection Software to Manage**
Lethal Recessive Conditions in Livestock Populations. In *10th World Congress on*
Genetics Applied to Livestock Production; 2014.

34. Cole JB: **A simple strategy for managing many recessive disorders in a dairy**
cattle breeding program. *Genet Sel Evol* 2015, **47**:94.

35. Brian K. Whitlock: **HERITABLE BIRTH DEFECTS IN CATTLE**. In *Applied*
Reproductive Strategies Conference Proceedings; 2010:146–154.

36. Flisikowski K, Venhoranta H, Nowacka-Woszuk J, Mckay SD, Flyckt A, Taponen J,
Schnabel R, Schwarzenbacher H, Szczerbal I, Lohi H, Fries R, Taylor JF, Switonski M,
Andersson M: **A novel mutation in the maternally imprinted PEG3 domain**
results in a loss of MIMT1 expression and causes abortions and stillbirths in
cattle (Bos taurus). *PLoS One* 2010, **5**:1–9.

37. Jenko J, Gorjanc G, Cleveland MA, Varshney RK, Whitelaw CBA, Woolliams JA,
Hickey JM: **Potential of promotion of alleles by genome editing to improve**
quantitative traits in livestock breeding programs. *Genet Sel Evol* 2015, **47**:55.

38. Pang AW, MacDonald JR, Pinto D, Wei J, Rafiq MA, Conrad DF, Park H, Hurles ME,
Lee C, Venter JC, Kirkness EF, Levy S, Feuk L, Scherer SW: **Towards a**
comprehensive structural variation map of an individual human genome.
Genome Biol 2010, **11**:R52.

39. Li X, Kim Y, Tsang EK, Davis JR, Damani FN, Chiang C, Zappala Z, Strober BJ, Scott
AJ, Ganna A, Merker J, Hall IM, Battle A, Montgomery SB: **The impact of rare**
variation on gene expression across tissues. *bioRxiv* 2016.

Table 1: Chromosomal regions predicted to harbor lethal haplotypes identified in the analysis of the BovineSNP50 data.

Chr	Location (Mb)	Length (Mb)	Haplotype Frequency[1]	Number of Patrios[2]	Probability[3]	Sequenced Carriers	Concordant Variants	Concordant In High Coverage
1	27.7-29.0	1.3	0.023	39	0.0042	1	4	4
4	82.5-84.0	1.5	0.076	127	2.66E-09	21	9	118
8	62.0-63.0	1.0	0.023	35	0.0074	5	1	1
12	60.0-61.2	1.2	0.032	46	0.0014	12	0	0
15	82.3-83.1	0.8	0.038	31	0.011	10	1	1
17	46.5-47.5	1.0	0.045	49	0.00076	15	2	2
29	43.0-44.2	1.2	0.044	118	3.22E-08	16	3	13

706 [1]Haplotypes estimated for 20 contiguous SNP loci.

707 [2]Number of families out of 2,480 for which the sire and maternal grandsire were both heterozygotes for the haplotype.

708 [3]Probability of observing no homozygous progeny if the haplotype is selectively neutral.

Figure 1: Effect of window size on haplotypic diversity and lethal haplotype detection. A) As the size of the window expands, many more distinct haplotypes are detected genome-wide. However, fewer of the newly detected haplotypes are common as window size increases, and the number of common haplotypes that are never observed as being homozygous asymptotes. B) Rate of homozygosity, which is the percentage of individuals that are homozygous for any haplotype, is high for small window sizes but quickly declines. The assumption that phased marker homozygosity implies identity by descent underlies the population frequency and patrio tests for haplotype lethality.

Figure 2: Validating the sequence level IBD status of bulls predicted to be carriers of a BovineSNP50 lethal haplotype using sequence data. Homozygous inconsistency rates are calculated pairwise amongst predicted carriers, sire-son pairs and randomly sampled animals (not 1st degree relatives). Two different regions are compared: within the tested chromosome 29 haplotype and across the entirety of chromosome 29. The discordance rate among of pairs (shown in red) that contain one animal with low coverage (<10x) is greatly elevated.