# Estimating DSM Validity for Attention Deficit/Hyperactivity Disorder Based on Neurophysiological, Psychological, and Behavioral Correlates

Dimitri M. Abramov[1], Evelyne Vigneau[2], Saint-Clair Gomes-Junior[3], Carlos Alberto Mourão-Júnior[4], Monique Castro-Pontes[1], Carla Quero Cunha[1], Leonardo C. deAzevedo[1], Vladimir V. Lazarev[1]

1. Laboratory of Neurobiology and Clinical Neurophysiology, National Institute of Women, Children, and Adolescents Health Fernandes Figueira, Oswaldo Cruz Foundation, Rio de Janeiro, Brazil;

2. StatSC, Oniris, INRA, Nantes, France;

3. Clinical Research Unit, National Institute of Women, Children, and Adolescents Health Fernandes Figueira, Oswaldo Cruz Foundation, Rio de Janeiro, Brazil; and

4. Laboratory of Psychophysiology, Institute of Biological Sciences, Federal University of Juiz de Fora, Juiz de Fora, Brazil.

**Corresponding Author:**
Dimitri Marques Abramov
National Institute of Women, Children, and Adolescents Health Fernandes Figueira,
Oswaldo Cruz Foundation
Avenida Ruy Barbosa, 716, Flamengo, Rio de Janeiro, RJ
Postal Code: 22.250-020
Phone: +55 21 2554-1955, +55 21 2554-1805
E-mail: dimitri.abramov@iff.fiocruz.br

**Short Title:** *Estimating DSM validity for ADHD diagnosis.*

# Abstract

**Background.** The psychiatric nosology lacks objective biological foundation, as well as characteristic biomarkers for the diagnoses, which raises questions about its validity. The problem is particularlly evident concerning the Attention Defict/Hyperactivity Disorder (ADHD). The objective of this study is estimate if the "Diagnostic and Statistical Manual of Mental Disorders" (DSM) is biologically valid for ADHD diagnosis, using a multivariate analysis of a large dataset concerning neurophysiological, behavioral, and psychological variables.

**Methods:** Twenty typically developing boys and 19 boys diagnosed with ADHD, aged 10-13 years, were examined using the Attentional Network Test (ANT) with records of event-related potentials (ERPs). From 815 variables, a reduced number of latent variables (LVs) were extracted with a clustering method for the reclassification of subjects using the k-means method.

**Results:** From datasets including ANT and WISC scores and ERPs from the mid-frontal, mid-parietal, right frontal, and central channels, only eight subjects were miss-reclassified by the LVs. In the reclassification, the estimated specificity of 75% and sensivity of 85% for DSM. The kappa index between DSM and

43 behavioral/psychological/neurophysiological data was 0.72, regarded as a "substantial

44 level of agreement".

45 **Discussion:** Results suggest the biological validity of DSM for ADHD diagnosis, in

46 accordance to alteractions in fronto-striatal networks previously related to ADHD.

# Introduction

The validity of the *Diagnostic and Statistical Manual of Mental Disorders* -DSM

(American Psychiatric Association, 1994 – 2013) [1,2] is usually questioned due to its

subjective criteria and to the absence of objective tests to define nosological entities [3,

4]. Therefore, the lack of objective determinants opens the way to criticism and doubts

about the nosological criteria itself [4]. This is particularly important in controversial

mental diseases such as the Attentional Deficit/Hyperactivity Disorder – ADHD [5,6].

Similar to some other disorders, ADHD diagnosis is based on the quantification of

normal behavior characteristics using DSM scores. Thus, DSM has a taxonomy of

categories that are defined by dimensional phenomena based on a variation from

normality [7,8]. Correlations between biological alterations and ADHD are still not

enough to support the diagnosis [8, 9].

Mental disorders are multidimensional complex entities that can be mostly associated with a pool of biological and neurobehavioral variables, often with non-linear relationships [8, 10], rather than being defined by a single distinct biomarker such as an antibody or aberrant protein [11], which formally defines their complexity. A complex nosological entity such as ADHD may be defined by a set of different quantitative variables. From this standpoint, multivariate factor analysis techniques have been applied to electroencephalographic or biochemical data since the 70's in order to develop high-sensitivity and accuracy models for the diagnosis of mental disorders [12-16].

The present study is an attempt to discuss whether DSM is valid for an accurate diagnosis of the nosological entity called ADHD, analyzing a small sample of subjects where the confounding variables and the bias were well controlled. Large datasets of (1) behavioral variables, and (2) neurophysiological data (event-related potentials – ERPs, including late cognitive component P3 or P300 [17,18]), both obtained while subjects performed the Attention Network Test - ANT [19,20], as well as (3) psychological characteristics (from WISC-III) have been collected from typically developing boys (TD) and boys diagnosed with ADHD using the DSM-IV-TR criteria.

However, high-dimensional datasets from relatively few observations or subjects usually appear to be statistically questionable and hamper unambiguous interpretation. In order to tackle this problem (the curse of the dimensionality), approaches based on

feature selection in a supervised context or on feature extraction in explorative data analysis (for instance, Principal Components Analysis, Canonical Correlation Analysis) have been investigated. In particular, we have considered a strategy of unsupervised clustering of variables that makes it possible to extract synthetic/latent variables through cluster analyses [21-23].

The relationships between ADHD manifestations, evaluated using DSM-IV-TR, and the objective characteristics obtained by the above-mentioned experimental measurements were here analyzed. The aim was to discuss the taxonomic (i.e. classificatory) aspects of ADHD diagnosis using DSM, compared with a data-driven statistical approach, without considering the role of these characteristics in putative brain mechanisms underlying ADHD. This study does not address controversial issues, as if the phenomenology of ADH actually configures a mental disorder or not. Here we seek to discuss the validity of the DSM as a tool to identify this phenomenology, inferring about its sensitivity and specificity.

# Methods

## Design and Subject Selection

This transversal and exploratory study was conducted in accordance with the

Declaration of Helsinki and approved by the Ethics Committee of the National Institute of

Women, Children, and Adolescents Health Fernandes Figueira (CAAE

08340212.5.0000.5269). All the participants provided their oral consent in the presence

of their caregivers, who provided written informed consent after receiving a complete

description of the study.

Sixty boys, aged 10-13 years, were included according to DSM-IV-TR: 35 with

ADHD (coded as T001 to T035) and 25 TD (coded as C001 to C025). The exclusion

criteria were: (1) history of chronic diseases, and any suspicion of psychiatric disorders

other than ADHD (psychosis, affective, obsessive-compulsive and tic disorders, phobic

and post-traumatic stress conditions, anorexia, bulimia, encopresis or enuresis) as

screened by K-SADS-PL [23]; (2) use of any psychotropic medicines for at least 30

days; (3) estimated I.Q. equal or lower than 80; and (4) less than 6 hours of regular

sleep and (5) report of somnolence before the ANT testing.

We considered as confounding variables years in school, monthly family income

(in Brazilian Reals), sleep hours the night before, and mean weekly time which the boy

dedicated to computer activities and videogames (ranked as follows: 1, less than

2h/week; 2, 2-4h; 3, 5-8h; 5, more than 15 h/week), learning rate (see below) in the

experimental procedure, as well as age.

After applying the exclusion criteria, the behavioral analysis of accuracy (AC) and speed-accuracy tradeoff (see below), and also after excluding confounding outliers, there remained 19 ADHD and 20 TD boys in the sample, whose data were considered as experimental results.

## Clinical and Psychological Examination

Each subject was evaluated using a structured interview where their caregivers were shown the DSM-IV-TR criteria and were instructed to point out carefully whether or not each specific criterion was an exact characteristic of their children's behavior. If there was any doubt or hesitation on any item, it was disregarded. Thus, subjects were classified in accordance with the DSM-IV-TR.

Intelligence Quotient (I.Q.) was estimated by Block Design and Vocabulary subtests from the Wechsler Intelligence Scale for Children, WISC-III [24, 25]. We also performed the Arithmetic and Digit Spam subtests to the pool of variables used in the multivariate analysis.

## Experimental Procedures

The ANT version adapted for children was used [20, 26]. A forced, two-choice test was performed, where the subject was instructed to observe the horizontal orientation of a target stimulus (yellow fish), which is (or not) preceded by a cue signal (a red star).

The target appeared above or below the fixation point. There were three equiprobable

cue conditions corresponding to this signal's position or to its non-appearance: 1) at the

subsequent upper or lower position of the target - spatial cue condition; 2) at the central

fixation point - neutral cue condition; or 3) no cue condition. The subject had to promptly

press the left or right arrow key on the keyboard, according to the horizontal orientation

of the target. The test was organized in 8 test blocks, with 24 trials each, and one

preceding training block. Reaction time (RT) and accuracy (AC, rate of hits) were

recorded.

Subjects with both AC lower than 70% and speed-accuracy tradeoff (estimated as

AC × RT) lower than the mean sample value minus two standard deviations were

excluded.

## EEG Acquisition

144 During the ANT, the subject's EEG was recorded using a Nihon Kohden NK1200

145 EEG System at 20 scalp points according to the International 10/20 System (Figure 1),

146 with reference at the lateral central leads (linked C3 and C4, the physical reference of

147 the System). Impedance was below 10 kΩ, sampling rate was 1000 Hz with a 16-bit

148 resolution, and the filters were: low-pass 0.5 Hz, high-pass 100 Hz, and notch 60 Hz.

**Figure 1. Electrical potentials related to Attention Network Test.** (A) Cue- and target-related potentials and inter-stimulus voltage variation during Attention Network Test performance (averaged over all ANT conditions and all subjects). Overview of the ERPs for all channels, including 'C3 minus C4' (normalized amplitudes, µV), with waves of interest (in dotted squares) to calculate the mean square of the total area under each wave and peak amplitudes/latencies. In the grey box, C and T stand for cue and target onsets; the trigger signal is identified by dotted lines. Above the ERP graphs, scalp areas are marked with letters with even (right), odd (left) or z (midline) indices: occipital – O, parietal – P, central – C, frontal – F, frontopolar – Fp, mid-temporal – T, posterior temporal – T5 and T6, and anterior temporal – F7 and F8. Scalp scheme for 10-20 montage at the bottom left. (B) ERPs for selected channels, in different colors, showing wave amplitudes (µV).

# Data Analysis

Predictor (i.e. DSM score for ADHD as the sum for inattention and hyperactivity/impulsivity scores) and confounding variables were compared between the groups using Student's t-test, after normality and heteroscedasticity assessments were performed using the Shapiro-Wilk and Levene's tests, respectively.

Several behavioral variables were gathered from the ANT for each subject: mean AC of task performance, i.e. the percentage of correct responses; mean Reaction Time (RT) and its standard deviation, called intra-individual variation of RT (IVRT), for each cue and target condition. Additionally, data included the ANT scores of the subjects: alerting, orienting, and conflict resolution, mean RT for correct and incorrect responses,

and learning rates (the mean scores of the first ANT trial block divided by those of the

last experimental block evaluated for RTs and IVRTs).

The following ERP characteristics were estimated inside time intervals of interest

(see dotted squares in Figure 1 and their measures in S1 Table) for each subject, in

each cue and target condition and in each EEG derivation (except for C3 and C4): (1)

peak amplitude, (2) peak latency, and (3) the total amplitude under the wave (i.e. the

sum of all amplitude values, proportional to the mean amplitude). These three time

intervals included late ERPs evoked by 1) the cue, 2) the target stimuli, and 3) the

voltage variation between these two ERPs. The same ERP parameters were calculated

for the difference between signals of the two central channels (C3 minus C4) instead of

estimating the absolute signal values in each of these leads, observing the EEG

reference used here (C3+C4).

Thus, the total number of variables of interest (VOI) was 774: four scores

obtained from the above mentioned WISC-III subtests, 29 behavioral variables obtained

from ANT, and 741 neurophysiological variables (39 for each of the 18 channels and C3-

C4).

The neurophysiological variables obtained represent topographical parameters of

regional (local) cerebral activity, while the variables from ANT and WISC provide the

integral characteristics of the subject. For analysis, we assumed that the activity in

certain brain regions of the subjects performing ANT (but not all) would best correlate with the phenomenology of ADHD. For this reason, 40 different sets of channels (Table 2) were used for data mining, in an attempt to find the best agreement with DSM (if any).

The data from each investigated set of EEG channels was analyzed either with or without behavioral (ANT) and psychological (WISC) variables. They were submitted to an unsupervised method called Clustering of Variables around Latent Variables (CLV) [21,22]. The aim was to reduce the number of dimensions in the variables space by extracting latent variables (LVs), each of them associated with a cluster of the VOI variables. This clustering approach is based on correlation similarity indices between variables and aims to identify directional clusters of variables. As it is, each extracted LV, associated with a specific cluster of variables, is defined as the first (standardized) principal component of the variables assigned to this cluster [21]. In our study, the clusters of variables around one to four LVs was systematically considered and the best number of LVs was determined according to the classification of subjects.

Indeed, our ultimate purpose was to reindex the subjects into two groups according to the objective behavioral, psychological, and neurophysiological data, all of which were summarized around the extracted LVs. A k-means clustering method [27] was performed on the subjects using the LVs. Each reclassification obtained (named "output") was checked for agreement compared to the classification of subjects using the DSM-IV-TR (called "DSM agreement"). The probability that the DSM agreement could

be observed under randomness was checked using the Chi-Square test for categorical variables, with one degree of freedon, where the expected value was 20 (absolute randomness). The kappa index [28] was also evaluated to estimate the DSM agreement level (S2 method).

In order to determine reclassification consistency, we analyzed which subjects were reclassified into the other group than his/her original DSM group ("miss-reclassification"). This was done for each specified DSM agreement rate, on the basis of the outputs providing this level of agreement. By dividing the DSM agreement level and the number of miss-reclassified subjects in any output with that DSM agreement, we suggested a measure for the "classifier power".

Subsequently, we considered that the outputs with the highest classifier power were the most representative to estimate the validity of the DSM. From these most representative outputs, we suggested a sensitivity and specificity analysis for DSM criteria for ADHD

The dataset with variables used in multivariate analysis is available as supplementary material (S3 Dataset).

# Results

## Evaluating predictor and confounding variables

Control and ADHD groups showed significant differences in the quantitative

scores from DSM-IV-TR (p < 0.05) (Table 1). The significance level  of the difference

between the two groups was higher for the inattention score than for the

hyperactivity/impulsivity score. Among variables considered as confounders, the I.Q.

scores in ADHD group were lower than in TD boys (p = 0.006), although they were

always higher than 80. All other variables, including learning rate, were not significantly

different between the two groups according to the Student's t-test (normality and

homoscedasticity were not statistically rejected).

**Table 1.** Predictor and Confounding Variables Between Groups

| | Typically Developing (n = 20) | | ADHD (n = 19) | | | |
|---|---|---|---|---|---|---|
| | *M* | *SD* | *M* | *SD* | T-value[a] | *p*-value |
| Predictor Variables (DSM scores) | | | | | | |
| **Inattention** | **2.40** | **1.66** | **7.21** | **1.27** | **-10.09** | **< 0.001** |
| **Hyperactive+Impulsive** | **2.60** | **1.63** | **4.36** | **2.90** | **-2.36** | **0.023** |
| **Total** | **5.00** | **2.75** | **11.63** | **3.00** | **-7.19** | **< 0.001** |
| Confounding Variables | | | | | | |
| Age (years) | 11.30 | 0.86 | 11.52 | 1.07 | -0.73 | 471 |
| **Estimated I.Q.** | **109.35** | **13.53** | **97.36** | **12.53** | **2.86** | **0.006** |
| Hours of Sleep (last night) | 7.25 | 2.12 | 7.68 | 1.73 | -0.70 | 0.490 |
| Videogame [b] | 3.40 | 1.23 | 2.89 | 1.66 | 1.08 | 0.286 |
| Computer handling [c] | 3.30 | 1.26 | 2.94 | 1.68 | 0.74 | 0.462 |
| Years in school | 6.05 | 1.14 | 6.10 | 1.32 | -0.14 | 0.889 |
| Family Incomes [d] | 6355.00 | 5643.57 | 3978.94 | 4436.91 | 1.46 | 0.153 |
| Learning Index [e] | 1.22 | 0.34 | 1.26 | 0.30 | -0.36 | 0.721 |

## Overview of Event-Related Potentials

243     Cue and target-related potentials with late peak latency (> 200ms, corresponding

244  to parietal P3) were observed in all channels. Peak amplitude varied from the maximum

245  value of 18μV for the frontopolar target ERP to the minimum value of 1μV for the

246  difference between C3 and C4 (Fig 1).

## Data Mining and Reclassification of subjects

For all 40 channel sets and LV sets (extracting 1 to 4 LVs), with and without

behavioral variables enclosed (Table 2), we found that the DSM agreement rates

ranging from 0.51 (20 subjects, p = 0.125, $\chi^2$ = 0.00, p = 1.000) to 0.82 (32 subjects, $\chi^2$

= 7.20, p =  0.004). The *worst* mean agreement level between DSM and K-means

reclassification was 23.25 for channel set [F7, T3, T5, Fp1, F3, C3-C4, P3, O1, Fz, Pz]

($\chi^2$ = 0.43, p = 0.528); and the best mean DSM agreement was 30.62 for channel set

[C3-C4, F8, F4, Fz, Pz] $\chi^2$ = 5.63, p = 0.010).

**Table 2.** DSM Agreement (Number of Subjects) by Channel Set and Number of Latent Vectors for

Variables of Interest

| Channel Sets [a] | ERPs variables only | | | | All variables of Interest | | | | MEAN |
|---|---|---|---|---|---|---|---|---|---|
| | 1 LV | 2 LV | 3 LV | 4 LV | 1 LV | 2 LV | 3 LV | 4 LV | |
| *All Channels* | 20 | 24 | 24 | 25 | 20 | 24 | 24 | 28 | 23.63 |
| *Fz Cz Pz Oz* | 23 | 25 | 23 | 22 | 27 | 28 | 29 | 29 | 25.75 |
| *F7 T3 T5 Fp1 F3 C3-C4 P3 O1 Fz Pz* | 21 | 22 | 22 | 24 | 20 | 22 | 27 | 28 | 23.25 |
| *C3-C4 F8 T4 T6 Fp2 F4 P4 O2 Fz Pz* | 24 | 24 | 24 | 24 | 24 | 24 | 24 | 24 | 24.00 |
| *F7 T3 T5 F8 T4 T6 Fz Cz Pz* | 20 | 25 | 25 | 24 | 23 | 25 | 29 | 28 | 24.88 |
| *F7 T3 T5 C3-C4 F8 T4 T6 Fz Cz Pz* | 22 | 20 | 22 | 24 | 22 | 28 | 28 | 28 | 24.25 |
| *F7 T3 T5 Fp1 C3-C4 F8 T4 T6 Fp2 Fz Cz Pz* | 21 | 23 | 24 | 24 | 23 | 28 | 25 | 28 | 24.50 |
| *C3-C4 Fz Cz Pz* | 27 | 27 | 27 | 26 | 30 | 30 | 30 | 30 | 28.38 |
| *F7 T3 T5 C3-C4 Fz Pz* | 22 | 28 | 23 | 22 | 22 | 30 | 28 | 29 | 25.50 |
| *C3-C4 F8 T4 T6 Fz Pz* | 20 | 25 | 24 | 22 | 31 | 31 | 29 | 28 | 26.25 |
| *F7 T3 C3-C4 Fz Pz* | 23 | 27 | 20 | 31 | 26 | 29 | 31 | 29 | 27.00 |
| *C3-C4 F8 T4 Fz Pz* | 31 | 30 | 28 | 29 | 30 | 30 | 31 | 28 | 29.63 |
| *C3-C4 Fz Pz* | 30 | 30 | 29 | 30 | 29 | 28 | 30 | 29 | 29.38 |
| *F7 T3 T5 C3-C4 F8 T4 T6 Fz Pz* | 21 | 20 | 23 | 22 | 22 | 26 | 28 | 29 | 23.88 |
| *F7 C3-C4 F8 Fz Pz* | 23 | 29 | 29 | 27 | 30 | 28 | 31 | 29 | 28.25 |
| *F7 F3 F8 F4 Fz Pz* | 27 | 25 | 31 | 30 | 29 | 29 | 30 | 31 | 29.00 |
| *F7 F3 C3-C4 F8 F4 Fz Pz* | 27 | 27 | 28 | 30 | 28 | 30 | 31 | 30 | 28.88 |
| *F7 F3 F8 F4 Fz* | 25 | 26 | 27 | 27 | 27 | 29 | 29 | 28 | 27.25 |
| *F7 F3 F8 T4 F4 Fz Pz* | 28 | 25 | 29 | 29 | 28 | 29 | 30 | 29 | 28.38 |
| *F7 F3 C3-C4 F8 F4 Fz* | 27 | 26 | 23 | 28 | 27 | 29 | 29 | 28 | 27.13 |
| *F7 Fp1 F3 C3-C4 F8 Fp2 F4 Fz Pz* | 27 | 29 | 29 | 30 | 28 | 29 | 29 | 26 | 28.38 |
| *F7 Fp1 F3 F8 Fp2 F4 Fz Pz* | 29 | 29 | 29 | 29 | 28 | 29 | 29 | 28 | 28.75 |
| *F7 Fp1 F3 C3-C4 F8 Fp2 F4 Fz* | 28 | 28 | 28 | 28 | 28 | 28 | 27 | 28 | 27.88 |

| Channel sets[a] | | | | | | | | Mean |
|---|---|---|---|---|---|---|---|---|
| F7 Fp1 F3 F8 Fp2 F4 Fz | 27 | 28 | 28 | 28 | 28 | 28 | 28 | 29 | 28.00 |
| F7 F3 C3-C4 P3 F8 F4 P4 Fz Pz | 27 | 28 | 27 | 21 | 29 | 29 | 29 | 29 | 27.38 |
| F7 F3 P3 F8 F4 P4 Fz Pz | 25 | 29 | 29 | 24 | 26 | 29 | 29 | 28 | 27.38 |
| F7 F3 C3-C4 P3 Fz Pz | 23 | 22 | 20 | 25 | 27 | 25 | 30 | 27 | 24.88 |
| F7 F3 P3 Fz Pz | 26 | 21 | 25 | 22 | 28 | 23 | 28 | 24 | 24.63 |
| **C3-C4 F8 F4 P4 Fz Pz** | **28** | **31** | **29** | **27** | **30** | **32** | **31** | **29** | **29.63** |
| F8 F4 P4 Fz Pz | 27 | 29 | 27 | 24 | 28 | 27 | 27 | 27 | 27.00 |
| F7 F3 C3-C4 Fz Pz | 26 | 27 | 28 | 22 | 29 | 29 | 30 | 29 | 27.50 |
| F7 F3 Fz Pz | 25 | 28 | 27 | 27 | 29 | 28 | 31 | 29 | 28.00 |
| **C3-C4 F8 F4 Fz Pz** | **32** | **31** | **31** | **31** | **29** | **31** | **31** | **29** | **30.63** |
| F8 F4 Fz Pz | 30 | 29 | 29 | 29 | 30 | 30 | 30 | 31 | 29.75 |
| C3-C4 F8 T4 F4 P4 Fz Pz | 28 | 30 | 27 | 23 | 31 | 30 | 31 | 27 | 28.38 |
| F7 T3 F3 C3-C4 P3 Fz Pz | 23 | 22 | 26 | 26 | 23 | 25 | 29 | 29 | 25.38 |
| C3-C4 F8 T4 F4 Fz Pz | 31 | 31 | 28 | 28 | 29 | 31 | 31 | 30 | 29.88 |
| F7 T3 F3 C3-C4 Fz Pz | 25 | 27 | 26 | 28 | 26 | 30 | 30 | 28 | 27.50 |
| Fz Pz | 27 | 29 | 29 | 29 | 29 | 27 | 28 | 30 | 28.50 |
| Fz Cz Pz | 27 | 27 | 22 | 26 | 28 | 28 | 28 | 27 | 26.63 |

[a] Channel sets according to the International 10-20 system for EEG leads; C3-C4, the channel of difference between two leads.

In bold and gray shading, higher DSM agreement.

ERPs = event-related potentials; LV = latent variables.

The two channel sets that manifested the highest DSM agreement were [C3-C4, F8, F4, Fz, Pz] (using 1 LV from ERP variables alone) and [C3-C4, F8, F4, P4, Fz, Pz] (using 3 LVs from ERP and behavioral variables together). Table 2 shows all the observed agreement rates for all channel sets.

As expected, we observed an inverse relationship between the DSM agreements

(from 20 to 32, according to the trial) and the total number of miss-reclassified subjects

(Fig 2). We observed that the highest agreement rate found (32 subjects) corresponded

to the highest classifier power (4.00). Eight subjects in these outputs were miss-

reclassified:  C005, C009, C012, C020, C023 (control subjects miss-reclassified to the

ADHD group) and T010, T012, T026 (ADHD subjects miss-reclassified to the Control

group). From these results, the estimated specificity and sensitivity of the DSM-IVTR

regarding biological variables were  75% and 84%, respectively, with an accuracy of

79.48% corresponding to the kappa index of 0.72, i.e. *a substantial strength of*

*agreement* [28].

**Figure 2. Classifier Power of the MVA analysis.** Number of miss-reclassified subjects by DSM

agreement (number of subjects), in all 320 outputs, and the corresponding classifier power (numbers at

graph marks). See methods.

# Discussion

Our study tackled the possibility of examining attention defict/hyperactivity

disorderusing the DSM manual, whose criteria seemed to be biologically justified for

ADHD diagnosis with "substantial strength of agreement" in relation to a specific pool of

behavioral, psychological, and neurophysiological variables. We found in the ANT the

larger source of behavioral and neurophysiological data related to ADHD phenomenology, since this test is suitable to access in the same setting the three dimensions of attention [19, 29].

To classify subjects as normal or ADHD according to DSM criteria, we asked caregivers what criteria were well-defined characteristics of their children. In both scientific research and clinical practice, for ADHD diagnosis it is appropriate that we also use other assessment methodologies besides the DSM. Moreover,in clinical practice the manual should not be applied as a questionnaire. However, in this case, the object of evaluation is the DSM itself, thus it was the only clinical diagnostic tool used for subjects classification as ADHD or not-ADHD. It was the most objective and controlled way to obtain the scores without interference of any other clinical impression. Supporting the present methodology, we observed that the ASRS, the WHO questionnaire for adult ADHD screening, is applied through self-evaluation [30].

The most informative neurophysiological variables in this pool proved to be topographically asymmetrical and represented the binded information from ERPs in the three time intervals of interest from mid-frontal, mid-parietal, right frontal, and right anterior temporal sites (channels Fz, Pz, F4, and F8), as well as early ERP components (45 to 290ms after target onset) from the 'C3 minus C4' channel. This asymmetry has already been shown to correlate with DSM scores, mainly with hyperactivity/impulsivity [31]. It was not the focus of the present investigation to establish models for ADHD or its

neural mechanisms. However, our results revealed certain biological aspects of ADHD regarding alterations in the frontal control over executive functions [18, 32, 33]. Several studies have shown  brain asymmetries related to ADHD phenomenology, spetially in frontal-striatal network [33-35]. By spectral and coherence analyses of the resting EEG, we also have showed asymmetrical topographic patterns, with signs of relative inactivation of the frontal and *left* temporal cortex, known to be responsible for voluntary attention, and impaired in ADHD patients [36]. These observations could partially explain the leading role of ERP data from the right frontal regions in discriminating between patients and controls in the present research.

The present results suggest that DSM-IV-TR is actually an effective and biologically justified tool for ADHD diagnosis.  Although a Kappa index of 0.72 should be considered with reservations for some clinical settings, a clinical method such as DSM, with sensitivity of 84% and specificity of 75% with regard to objective behavioral, psychological, and neurophysiological measures, should be considered as particularly relevant considering the complexity of mental disorders, spetially those whose clinical phenomenology is dimensional, and without any validated biomarkers, So much so that the only psychiatric tool for diagnosis and intervention is still phenomenological examination, which must be systematically (although qualitatively) performed based on manuals such as the DSM. Furthermore, the ADHD has been regarded as a heterogeneous condition with inattentive, hyperactive, and combined subtypes, which

could be the answer for this level of agreement since these subtypes were not predicted

for reclassification. The existence of three discrete subtypes could lead to flawed

reclassification.

The study was designed with a relatively small sample size. Any subject with the

slightest suspicion of comorbidity was excluded. We monitor potentially confounding

factors commonly ignored (such as manipulating technologies that could lead to the

development of specific skills that would interfere with test performance and the

socioeconomic niche, that could determine the psychological development).. Thus, our

sample should be regarded as well consistent with and representative of ADHD. On the

other hand, in order to classify the subjects according to their mental condition (a

multidimensional phenomenological complex) it would seem reasonable to gather a

considerable number of objective variables (as many as possible). Brain processing

during the ANT test as well as ADHD neurophysiology and its behavioral correlates in

this study were considered as potentially relevant features requiring further investigation.

Thus, we had to deal with a dataset with a much smaller number of observations than

variables, since it did not seem feasible to collect a sample with thousands of subjects

evaluating hundreds of variables in an experiment with ideal control of confounders and

bias. A small but strictly controlled and well exploited sample was considered preferable,

even in conditions of high prevalence such as the ADHD. It is well established that

multivariate statistical modeling, such as Linear Regression, Discriminant Analysis, or

Factor Analysis, are not suitable for this situation. Thus,in order to tackle a frequently

encountered problem, often designated as the curse of the dimensionality, we have

adopted an explorative data analysis suitable for small samples, to identify a small

number of synthetic latent variables. Instead of the Principal Component Analysis, which

does not always provide easily interpretable latent variables (the principal components),

we have chosen an approach with clustering of variables to reduce data dimensionality

to a small number of new latent variables. The CLV method  [21,22] has been applied in

a  wide  range  of  research  domains  among  which  are  sensory  analysis  [37],

chemometrics [38], image analysis [39],  and analysis scorecards in  health sector [40].

The estimated I.Q. scores were different between the groups diagnosed using

DSM-IV-TR, but this was not regarded as bias. Literature has shown that intelligence

tests are sensitive to ADHD, and patient scores are generally lower than in controls [41].

Even taking into account the above restrictions, there is no doubt that the data obtained

seem to be quite consistent.

A  major  limitation  of  this  study  is  the  impossibility  of  testing  all  possible

combinations among variables. Since consistency of our paradigm depends on not

considering any biological / behavioral assumptions in the grouping of variables, it is

necessary to test  them randomly grouped. But there are more than one million possible

combinations of EEG channels. The number of combinations grows exponentially if we

include the other variables individually. For computational feasibility, we had arbitrarily

choose sets of variables to analize. With the eighty sets studied, we can assume that
DSM has at least "substantial level of agreement" with biological determinants. Perhaps
some untested set of variables may manifest greater agreement with the DSM, but the
results found already satisfy our objective of testing the validity of the DSM criteria for
ADHD.

In the present research, our objective was to draw attention to psychiatric
approaches based on diverse and multiple factors intervening in the biological
mechanisms of human behavior and its disorders. In this way, we have found that there
is a consistent complex of biological information which is in accordance with previous
knowledge on ADHD, that confers  biological validity for  DSM criteria for ADHD.

## 373 Aknowledgements

We would like to thank The Child´s Neurology program of the National Institute of
Women, Children, and Adolescents' Health Fernandes Figueira. We are also grateful to
prof. Daniela Tannus for proofreading the text, and to Mrs. Maria S. Santana and Mrs.
Aldenys Perez for the technical support. Our team is especially grateful to our friend,
colleague, and co-author Dr. Adailton Tadeu Pontes, children's neurologist, who passed
away during the preparation of this article, which was greatly inspired by him.

# References

**References**

1. American Psychiatric Association. Diagnostic and statistical manual of mental

disorders. 4th ed. Washington, DC: Author; 1994.

2. American Psychiatric Association. Diagnostic and statistical manual of mental

disorders. 5th ed. Arlington, VA: American Psychiatric Publishing; 2013.

3. Insel T. Director's Blog: Transforming Diagnosis. 2013 Apr 29 [cited 06 Jul 2018] In:

The National Institute of Mental Health Website. [about 2 screens] Available from:

http://www.nimh.nih.gov/about/director/2013/transforming-diagnosis.shtml.

4. Wakefield JC. DSM-5, psychiatric epidemiology and the false positives problem.

Epidemiol Psychiatr Sci. 2015 Jun;24(3):188-96. doi: 10.1017/S2045796015000116.

5. Rafalovich A. Exploring clinician uncertainty in the diagnosis and treatment of

attention deficit hyperactivity disorder.Sociol Health Illn. 2005 Apr;27(3):305-23.

6. Strauss V. An ADHD controversy in the mental health community. 2013 February 13

[cited 06 July 2018]. In: The Washington Post [Internet]. Available from:

https://www.washingtonpost.com/blogs/answer-sheet/post/an-adhd-controversy-in-the-

mental-health-community/2012/02/12/gIQAHJun9Q_blog.html.

7.Maser JD, Akiskal HS. Spectrum concepts in major mental disorders. Psychiatr Clin

North Am. 2002 Dec;25(4):xi-xiii.

8.Kirmayer LJ, Crafa D. What kind of science for psychiatry? Front Hum Neurosci. 2014

Jun 20;8:435. doi: 10.3389/fnhum.2014.00435.

9.Furman LM. Attention-deficit hyperactivity disorder (ADHD): does new research

support old concepts? J Child Neurol. 2008 Jul;23(7):775-84. doi:

10.1177/0883073808318059.

10. Freitas-Silva LR, Ortega F. Biological determination of mental disorders: a

discussion based on recent hypotheses from neuroscience. Cad Saude Publica. 2016

Aug 29;32(8):e00168115. doi: 10.1590/0102-311X00168115.

11.Scarr E, Millan MJ, Bahn S, Bertolino A, Turck CW, Kapur S, Möller HJ, Dean B.

Biomarkers for Psychiatry: The Journey from Fantasy to Fact, a Report of the 2013

CINP Think Tank. Int J Neuropsychopharmacol. 2015 Apr 21;18(10):pyv042. doi:

10.1093/ijnp/pyv042.

12. Bochkarev VK, Lazarev VV, Nikiforov AI, Paniushkina SV, Severnyĭ AA. Clinico-

electroencephalographic correlations in asthenoadynamic subdepressions.Zh

Nevropatol Psikhiatr Im S S Korsakova. 1987;87(4):564-70.

13.Lazarev VV. The relationship of theory and methodology in EEG studies of mental

activity.Int J Psychophysiol. 2006 Dec;62(3):384-93. Epub 2006 Mar 10.

14. Lazarev VV, Lebedeva IS, Tsutsul'kovskaia MIa, Severnyĭ AA, Pozharitskaia DA.

Specificity of electroencephalographic correlations of asthenic disorders in atypical

juvenile asthenic-depressive conditions. Zh Nevropatol Psikhiatr Im S S Korsakova. 1990;90(10):73-7.

15. Monakhov K, Perris C, Botskarev VK, von Knorring L, Nikiforov AI. Functional interhemispheric differences in relation to various psychopathological components of the depressive syndromes. A pilot international study.Neuropsychobiology. 1979;5(3):143-55.

16. Schwarz E, Izmailov R, Spain M, Barnes A, Mapes JP, Guest PC, Rahmoune H, Pietsch S, Leweke FM, Rothermundt M, Steiner J, Koethe D, Kranaster L, Ohrmann P, Suslow T, Levin Y, Bogerts B, van Beveren NJ, McAllister G, Weber N, Niebuhr D, Cowan D, Yolken RH, Bahn S. Validation of a blood-based laboratory test to aid in the confirmation of a diagnosis of schizophrenia. Biomark Insights. 2010 May 12;5:39-47.

17. Hruby T, Marsalek P. Event-related potentials--the P3 wave. Acta Neurobiol Exp (Wars). 2003;63(1):55-63.

18. Barry RJ, Johnstone SJ, Clarke AR. A review of electrophysiology in attention-deficit/hyperactivity disorder: II. Event-related potentials. Clin Neurophysiol. 2003 Feb;114(2):184-98.

19. Fan J, McCandliss BD, Sommer T, Raz A, Posner MI. Testing the efficiency and independence of attentional networks. J Cogn Neurosci. 2002 Apr 1;14(3):340-7.

20.Kratz O, Studer P, Malcherek S, Erbe K, Moll GH, Heinrich H. Attentional processes

in children with ADHD: an event-related potential study using the attention network test.

Int J Psychophysiol. 2011 Aug;81(2):82-90. doi: 10.1016/j.ijpsycho.2011.05.008.

21. Vigneau E, Qannari EM. Clustering of Variables Around Latent Components.

Communications in Statistics - Simulation and Computation. 2003; 32: 1131-1150.

22. Vigneau E, Chen M, Qannari E M. ClustVarLV: An R Package for the Clustering of

Variables Around Latent Variables. The R Journal. 2015 ; 7 : 134-148.

23.Matuschek T, Jaeger S, Stadelmann S, Dölling K, Grunewald M, Weis S, von Klitzing

K, Döhnert M. Implementing the K-SADS-PL as a standard diagnostic tool: Effects on

clinical diagnoses. Psychiatry Res. 2016 Feb 28;236:119-24. doi:

10.1016/j.psychres.2015.12.021.

24.Wechsler D. Wechsler Intelligence Scale for Children(WISC-III): Manual. 3th ed. San

Antonio: The Psychological Corporation; 1991.

25.Mello CB, Argollo Shayer PBM, Abreu N, AbreuII N, GodinhoIII K,  Durán P, Vargem

F, Muskat M, Miranda MC, Bueno OFA. Abbreviated version of the WISC-III: correlation

between estimated IQ and global IQ of Brazilian children, Psicologia: Teor e Pesq. 2011;

27 (2): 149-155. dx.doi.org/10.1590/S0102-37722011000200002.

26. Abramov DM, Pontes M, Pontes AT, Mourao-Junior CA, Vieira J, Quero Cunha C,

Tamborino T, Galhanone PR, deAzevedo LC, Lazarev VV. Visuospatial information

processing load and the ratio between parietal cue and target P3 amplitudes in the

Attentional Network Test. Neurosci Lett. 2017 Apr 24;647:91-96. doi:

10.1016/j.neulet.2017.03.031.

27.Gore Jr PA. Cluster Analysis, In: Tinsley HEA, Brown SD, editors. Handbook of

Applied Multivariate Statistics and Mathematical Modeling. San Diego, CA: Academic

Press; 2000. p. 297-321.

28. Landis JR, Koch GG. The measurement of observer agreement for categorical data.

Biometrics. 1977 Mar;33(1):159-74.

29. Petersen SE, Posner MI. The attention system of the human brain: 20 years after.

Annu Rev Neurosci. 2012;35:73-89. doi: 10.1146/annurev-neuro-062111-150525.

30. Kessler RC, Adler L, Ames M, Demler O, Faraone S, Hiripi E, Howes MJ, Jin R,

Secnik K, Spencer T, Ustun TB, Walters EE. The World Health Organization Adult

ADHD Self-Report Scale (ASRS): a short screening scale for use in the general

population. Psychol Med. 2005 Feb;35(2):245-56.

31. Abramov DM., Cunha CQ, Pontes MC, Galhanone PR, deAzevedo LC, Lazarev VV.

Functional Asymmetry In The Central Brain Regions In Boys With Attention Deficit

Hyperactivity Disorder Detected By Event Related Potentials During Performance Of

The Attentional Network Test. 2016. Preprint.Avaiable on BioRxiv,

https://doi.org/10.1101/118380 ; cited on 06 July 2018.

32. Barkley RA. Behavioral inhibition, sustained attention, and executive functions:

constructing a unifying theory of ADHD. Psychol Bull. 1997 Jan;121(1):65-94. doi:

10.1037/0033-2909.121.1.65.

33. Barkley RA. Attention-Deficit/Hyperactivity Disorder: handbook for diagnosis and

treatment. 3th ed. New York: The Guilford Press; 2006. p 32-33.

34. Langleben DD, Austin G, Krikorian G, Ridlehuber HW, Goris ML, Strauss HW.

Interhemispheric asymmetry of regional cerebral blood flow in prepubescent boys with

attention deficit hyperactivity disorder.  Nucl Med Commun. 2001 Dec;22(12):1333-40.

35.Dang LC, Samanez-Larkin GR, Young JS, Cowan RL, Kessler RM, Zald DH.

Caudate asymmetry is related to attentional impulsivity and an objective measure of

ADHD-like attentional problems in healthy adults. Brain Struct Funct. 2016

Jan;221(1):277-86. doi: 10.1007/s00429-014-0906-6.

36. Lazarev VV, Pontes M, Pontes AT, Vieira J, Quero-Cunha C, Tamborino T,

deAzevedo LC, Abramov DM. EEG and ERP characteristics of Attention Deficit

Hyperactivity Disorder in children and adolescents [abstract]. Int. J. Psychophysiol.

2106; 108: 75.

37. Vigneau E, Sahmer K, Qannari EM, Bertrand D. Clustering of variables to analyze

spectral data. J Chemometrics, 2005; 19: 122-128.

38.Vigneau E, Qannari EM, Navez B, Cottet V. Segmentation of consumers in

preference studies while setting aside atypical or irrelevant consumers. Food Quality

Pref. 2005; 47: 54-63.

39.Legland D, Beaugrand J. Automated clustering of lignocellulosic fibres based on

morphometric features and using clustering of variables. Industrial Crops and Products;

2013; 45: 253– 261.

40.Lovaglio PG. Model building and estimation strategies for implementing the Balanced

Scorecard in Health sector. *Quality & Quantity.* 2011*;* 45: 199–212.

41. Mackenzie GB, Wonders E. Rethinking Intelligence Quotient Exclusion Criteria

Practices in the Study of Attention Deficit Hyperactivity Disorder. Front Psychol. 2016

May 31;7:794. doi: 10.3389/fpsyg.2016.00794.

**S1 Table. Time intervals of Interest.** Windows holding Cue-related potentials (CUE),

contigent voltage variation (CVV) and Target-related potentials (TGT), for each channel

(ms), for calculation of peak amplitude, peak latency and mean amplitude. Zero ms

refers to Target onset (see Figure 1).

**S2 Method. Calculating the Kappa Index.** Inference of the magnitude of agreement

between two categorical data (in this case, the subjects' classification by DSM and by

biological/behavioral variables in the dataset).

**S3 Dataset. DSM scores and variables of interest.** *In the first sheet:* innatention,

hyperactive+impulsive and total scores for ADHD by DSM-IV-TR; *second sheet:* all data

concerning (1) scores of the WISC-III subtests, (2) ANT performance (AC: accuracy, RT:

reaction time and IVRT: intraindividual variation of reaction time, what is the standard

deviation of RT for each subject) for each experimental condition (Allcon: mean of all

conditions, NtCue: neutral Cue condition, SpCue: spatial cue condition, ConTg:

congruent target condition, IncTg: incongruent target condition) and attentional network

(alerting, orienting, and executive) plus learning indexes (IVRT or RT at the last ANT

block/first ANT block), and (3) the neurophysiological measures (peak amplitudes and

latencies, and mean amplitudes inside each time interval of interest in S1) for EEG

channel set.

**Classifier Power**

Number of miss-reclassified subjects

DSM Agreement (n. of subjects)

0.51  0.53  0.59  0.63  0.73  0.75  0.77  0.83  1.00  1.11  1.30  1.63  4.00