Estimating the Biological Validity of the DSM for Attention Deficit/Hyperactivity Disorder Using Multivariate Analysis for Small Samples

Dimitri M. Abramov[1], Evelyne Vigneau[2], Saint-Clair Gomes-Junior[3], Carlos Alberto Mourão-Júnior[4], Monique Castro-Pontes[1], Carla Quero Cunha[1], Leonardo C. deAzevedo[1], Vladimir V. Lazarev[1]

1. Laboratory of Neurobiology and Clinical Neurophysiology, National Institute of Women, Children, and Adolescents' Health Fernandes Figueira, Oswaldo Cruz Foundation, Rio de Janeiro, Brazil;

2. StatSC, Oniris, INRA, Nantes, France;

3.Clinical Research Unit, National Institute of Women, Children, and Adolescents Health Fernandes Figueira, Oswaldo Cruz Foundation, Rio de Janeiro, Brazil; and

4.Laboratory of Psychophysiology, Institute of Biological Sciences, Federal University of Juiz de Fora, Juiz de Fora, Brazil.

Corresponding Author:
Dimitri Marques Abramov
National Institute of Women, Children, and Adolescents Health Fernandes Figueira,
Oswaldo Cruz Foundation
Avenida Ruy Barbosa, 716, Flamengo, Rio de Janeiro, RJ
Postal Code: 22.250-020
Phone: +55 21 2554-1955, +55 21 2554-1805
E-mail: dimitri.abramov@iff.fiocruz.br

Keywords: Attention Deficit/Hyperactivity Disorder, DSM, diagnostic validity, Attention Network Test, event-related potentials

    Short Title: *Estimating DSM validity for ADHD diagnosis.*

Abstract

Background. Psychiatric nosology lacks objective biological foundation, as well as typical biomarkers for diagnoses, which raises questions about its validity. The problem is particularly evident concerning Attention Deficit/Hyperactivity Disorder (ADHD). The objective of this study is to estimate whether the "Diagnostic and Statistical Manual of Mental Disorders" (DSM) is biologically valid for ADHD diagnosis using a multivariate analysis for small samples from a large dataset concerning neurophysiological, behavioral, and psychological variables.

Methods: Twenty typically developing boys and 19 boys diagnosed with ADHD, aged 10-13 years, were examined using the Attentional Network Test (ANT) with records of event-related potentials (ERPs). From 815 variables, a reduced number of latent variables (LVs) were extracted with a clustering method, for further reclassification of subjects using the k-means method. This approach allowed multivariate analysis to be applied to a significantly larger number of variables than the number of cases (E. Wigneau et al., 2003, 2015)

Results: From datasets including ERPs from the mid-frontal, mid-parietal, right frontal, and central channels, only seven subjects were miss-reclassified by the LVs. An

estimated specificity of 75.00% and sensitivity of 89.47% for DSM were found in the

reclassification.   The   kappa   index   between   DSM   and

behavioral/psychological/neurophysiological data was 0.75, which is regarded as a

"substantial level of agreement".

Discussion: Results showed that CLV is a useful method for diagnostic classification

using a large dataset of small samples, suggesting the biological validity of DSM for

ADHD diagnosis, in accordance to alterations in fronto-striatal networks previously

related to ADHD.

Introduction

The validity of the *Diagnostic and Statistical Manual of Mental Disorders* -DSM

(American Psychiatric Association, 1994 – 2013) [1,2] is usually questioned due to its

subjective criteria and to the absence of objective tests to define nosological entities [3,

4]. Therefore, the lack of objective determinants makes room for criticism and raises

doubts about the nosological criteria itself [4]. This is particularly important in

controversial mental diseases such as the Attentional Deficit/Hyperactivity Disorder –

ADHD [5,6].   Similar to some other disorders, ADHD diagnosis is based on the

quantification of normal behavior characteristics using DSM scores. Thus, DSM has a

taxonomy of categories that are defined by dimensional phenomena based on a

variation from normality [7,8]. Correlations between biological alterations and ADHD are

still not enough to support the diagnosis [8, 9].

Mental disorders are multidimensional complex entities that can be mostly

associated with a pool of biological and neurobehavioral variables, often with non-linear

relationships [8, 10], rather than being defined by a single distinct biomarker such as an

antibody or aberrant protein [11]. This formally determines their complexity; a complex

nosological entity such as ADHD may be defined by a large set of different quantitative

variables. From this standpoint, multivariate factor analysis techniques have been

applied to electroencephalographic or biochemical data since the 1970's in order to

develop high-sensitivity and accuracy models for the diagnosis of mental disorders [12-

16]. However, in order to explain these complex mental conditions in terms of biological

correlates, a substantially larger sample of subjects than those set of variables is

required. Thus, high-dimensional datasets from relatively few observations or subjects

usually appear to be statistically questionable and hamper unambiguous interpretation.

In order to tackle this problem (the curse of the dimensionality), approaches based on

feature selection in a supervised context or on feature extraction in explorative data

analysis (for instance, Principal Components Analysis, Canonical Correlation Analysis)

have been investigated. Particularly, a strategy of unsupervised "clustering of variables

around latent variables" (CLV) was used, which makes it possible to extract

synthetic/latent variables through cluster analyses [17-19]. These few latent variables

represent dataset informativeness for a classifier methodology in a subsequent step

(e.g. inputs for a k-means clustering algorithm). This was the first attempt to apply the

CLV method to a large psychophysiological dataset. This method is expected to help

solve an important methodological problem of multidimensional approaches in clinical

research when recruiting large groups of patients is difficult.

The present study is an attempt to discuss whether DSM is valid for an accurate

diagnosis of the nosological entity called ADHD, analyzing a small sample of subjects

with well-controlled confounding variables and bias. Large datasets of (1) behavioral

variables, (2) neurophysiological data (event-related potentials – ERPs, including late

cognitive component, P3 or P300 [20,21]), both obtained while subjects performed the

Attention Network Test - ANT [22,23], as well as (3) psychological characteristics (from

WISC-III) have been collected from typically developing boys (TD) and boys diagnosed

with ADHD using the DSM-IV-TR criteria.

The relationships between ADHD manifestations, evaluated using DSM-IV-TR,

and the objective characteristics obtained by the above-mentioned experimental

measurements were here analyzed. The aim was to discuss the taxonomic (i.e.

classificatory) aspects of ADHD diagnosis using DSM, compared with a data-driven

statistical approach, without considering the role of these characteristics in putative brain

mechanisms underlying ADHD. This study does not address controversial issues, e.g. if

the ADHD phenomenology actually constitutes a mental disorder or not. The present

study seeks to discuss the validity of the DSM as a tool to identify this phenomenology,

making inferences about its sensitivity and specificity.

105 Methods

106 Design and Subject Selection

This transversal and exploratory study was conducted in accordance with the

Declaration of Helsinki and approved by the Ethics Committee of the National Institute of

Women, Children, and Adolescents Health Fernandes Figueira (CAAE

08340212.5.0000.5269). All the participants provided their oral consent in the presence

of their caregivers, who provided written informed consent after receiving a complete

description of the study.

Sixty boys, aged 10-13 years, were included according to DSM-IV-TR: 35 with

ADHD (coded as T001 to T035) and 25 TD (coded as C001 to C025). The exclusion

criteria were: (1) history of chronic diseases, and any suspicion of psychiatric disorders

other than ADHD (psychosis, affective, obsessive-compulsive and tic disorders, phobic

and post-traumatic stress conditions, anorexia, bulimia, encopresis, or enuresis) as

screened by K-SADS-PL [23]; (2) use of any psychotropic medicines for at least 30

days; (3) estimated I.Q. equal or lower than 80; and (4) less than 6 hours of regular

sleep and (5) report of somnolence before the ANT testing.

The following were considered as confounding variables: years in school, monthly

family income (in Brazilian Reals), sleep hours the night before, and mean weekly time

which the boy dedicated to computer activities and videogames (ranked as follows: 1,

less than 2h/week; 2, 2-4h; 3, 5-8h; 5, more than 15 h/week), learning rate (see below)

in the experimental procedure, as well as age.

After applying the exclusion criteria, the behavioral analysis of accuracy (AC) and

speed-accuracy tradeoff (see below), and after excluding confounding outliers, 19 ADHD

and 20 TD boys remained in the sample, and their data were considered as

experimental results.

Clinical and Psychological Examination

Each subject was evaluated using a structured interview where his or her

caregivers were shown the DSM-IV-TR criteria and were instructed to point out carefully

whether or not each specific criterion was an exact characteristic of their child's

behavior. If there was any doubt about or hesitation in any item, it was disregarded.

Thus, subjects were classified in accordance with the DSM-IV-TR.

Intelligence Quotient (I.Q.) was estimated by Block Design and Vocabulary

subtests of the Wechsler Intelligence Scale for Children, WISC-III [24, 25]. The

Arithmetic and Digit Spam subtests were also performed for the pool of variables used in

the multivariate analysis.

Experimental Procedures

The ANT version adapted for children was used [20, 26]. A forced, two-choice test

was performed, where the subject was instructed to observe the horizontal orientation of

a target stimulus (yellow fish), which was (or not) preceded by a cue signal (a red star).

The target appeared above or below the fixation point. There were three equiprobable

cue conditions corresponding to this signal's position or to its non-appearance: 1) at the

subsequent upper or lower position of the target - spatial cue condition; 2) at the central

fixation point - neutral cue condition; or 3) no cue condition. The subject had to promptly

press the left or right arrow key on the keyboard, according to the horizontal orientation

of the target. The test was organized in 8 test blocks, with 24 trials each, and one

preceding training block. Reaction time (RT) and accuracy (AC, rate of hits) were

recorded.

Subjects with AC lower than 70% and speed-accuracy tradeoff (estimated as AC

× RT) lower than the mean sample value minus two standard deviations were excluded.

EEG Acquisition

During the ANT, the subject's EEG was recorded using a Nihon Kohden NK1200 EEG System at 20 scalp points according to the International 10/20 System (Figure 1), with reference at the lateral central leads (linked C3 and C4, the physical reference of the System). Impedance was below 10 kΩ, sampling rate was 1000 Hz with a 16-bit resolution, and the filters were as follows: low-pass 0.5 Hz, high-pass 100 Hz, and notch 60 Hz.

Figure 1. Electrical potentials related to Attention Network Test. (A) Cue- and target-related potentials and variation in interstimulus voltage during Attention Network Test performance (averaged over all ANT conditions and all subjects). Overview of the ERPs for all channels, including 'C3 minus C4' (normalized amplitudes, µV), with waves of interest (in dotted squares) to calculate the mean square of the total area under each wave and peak amplitudes/latencies. In the grey box, C and T stand for cue and target onsets, and the trigger signal is identified by dotted lines. Above the ERP graphs, scalp areas are marked with letters with even (right), odd (left) or z (midline) indices: occipital – O, parietal – P, central – C, frontal – F, frontopolar – Fp, mid-temporal – T, posterior temporal – T5 and T6, and anterior temporal – F7 and F8. Scalp scheme for 10-20 montage at the bottom left. (B) ERPs for selected channels, in different colors, showing wave amplitudes (µV).

Data Analysis

172   Predictor (i.e. DSM score for ADHD as the sum for inattention and
173 hyperactivity/impulsivity scores) and confounding variables were compared between the
174 groups using Student's t-test, after normality and heteroscedasticity assessments were
175 performed using the Shapiro-Wilk and Levene's tests, respectively (Table 1).

176 **Table 1.** Predictor and Confounding Variables between Groups

	Typically Developing (n = 20)		ADHD (n = 19)			
	M	*SD*	*M*	*SD*	T-value[a]	*p*-value
Predictor Variables (DSM scores)						
Inattention	**2.40**	**1.66**	**7.21**	**1.27**	**-10.09**	**< 0.001**
Hyperactive+Impulsive	**2.60**	**1.63**	**4.36**	**2.90**	**-2.36**	**0.023**
Total	**5.00**	**2.75**	**11.63**	**3.00**	**-7.19**	**< 0.001**
Confounding Variables						
Age (years)	11.30	0.86	11.52	1.07	-0.73	471
Estimated I.Q.	**109.35**	**13.53**	**97.36**	**12.53**	**2.86**	**0.006**
Hours of Sleep (previous night)	7.25	2.12	7.68	1.73	-0.70	0.490
Videogame [b]	3.40	1.23	2.89	1.66	1.08	0.286
Computer handling [c]	3.30	1.26	2.94	1.68	0.74	0.462
Years in school	6.05	1.14	6.10	1.32	-0.14	0.889
Family Incomes [d]	6355.00	5643.57	3978.94	4436.91	1.46	0.153
Learning Index [e]	1.22	0.34	1.26	0.30	-0.36	0.721

[a] Inference using Student's t-test for independent samples, significant differences (*p* value < 0.05) in bold.

[b, c] Videogame and computer handling were scored as: 1 = less than 2h/week; 2 = 2-4h/wk, 3 = 5-8 h/wk, 4
= 8-15 h/wk, 5 = more than 15h/wk.

[d] in Brazilian Reals (R$), per month

[e] Learning Index in the ANT: reaction time (RT) (ms) of last trial block / mean RT (ms) of the first
experimental blocks.

Several behavioral variables were gathered from the ANT for each subject: mean

AC of task performance, i.e. the percentage of correct responses; mean Reaction Time

(RT) and its standard deviation, called intra-individual variation of RT (IVRT), for each

cue and target condition. Additionally, data included the ANT scores of the subjects:

alerting, orienting, and conflict resolution, mean RT for correct and incorrect responses,

and learning rates (the mean scores of the first ANT trial block divided by those of the

last experimental block evaluated for RTs and IVRTs).

The following ERP characteristics were estimated inside time intervals of interest

(see dotted squares in Figure 1 and their measures in Table S1) for each subject, in

each cue and target condition and in each EEG derivation (except for C3 and C4): (1)

peak amplitude, (2) peak latency, and (3) the total amplitude under the wave (i.e. the

sum of all amplitude values, proportional to the mean amplitude). These three time

intervals included late ERPs evoked by 1) the cue, 2) the target stimuli, and 3) the

voltage variation between these two ERPs. The same ERP parameters were calculated

for the difference between signals of the two central channels (C3 minus C4) instead of

estimating the absolute signal values in each of these leads, observing the EEG

reference used here (C3+C4).

Thus, the total number of variables of interest (VOI) was 774: four scores

obtained from the above-mentioned WISC-III subtests, 29 behavioral variables obtained

from ANT, and 741 neurophysiological variables (39 for each of the 18 channels and C3-

C4).

The neurophysiological variables obtained represent topographical parameters of

regional (local) cerebral activity, while the variables from ANT and WISC provide the

integral characteristics of the subject. For the analysis, the activity in certain brain

regions of the subjects performing ANT (but not all) was assumed to best correlate with

ADHD phenomenology. For this reason, 40 different sets of channels (Table 2) were

used for data mining, in an attempt to find the best agreement with DSM (if any).

Table 2. Agreement with DSM (Number of Subjects) by Channel Set and Number of Latent Vectors for
Variables of Interest

Channel Sets [a]	ERPs variables only				All variables of Interest				MEAN
	1 LV	2 LV	3 LV	4 LV	1 LV	2 LV	3 LV	4 LV	
All Channels	20	24	24	25	20	24	24	28	23.63
Fz Cz Pz Oz	23	25	23	22	27	28	29	29	25.75
F7 T3 T5 Fp1 F3 C3-C4 P3 O1 Fz Pz	21	22	22	24	20	22	27	28	23.25
C3-C4 F8 T4 T6 Fp2 F4 P4 O2 Fz Pz	24	24	24	24	24	24	24	24	24.00
F7 T3 T5 F8 T4 T6 Fz Cz Pz	20	25	25	24	23	25	29	28	24.88
F7 T3 T5 C3-C4 F8 T4 T6 Fz Cz Pz	22	20	22	24	22	28	28	28	24.25
F7 T3 T5 Fp1 C3-C4 F8 T4 T6 Fp2 Fz Cz Pz	21	23	24	24	23	28	25	28	24.50
C3-C4 Fz Cz Pz	27	27	27	26	30	30	30	30	28.38
F7 T3 T5 C3-C4 Fz Pz	22	28	23	22	22	30	28	29	25.50
C3-C4 F8 T4 T6 Fz Pz	20	25	24	22	31	31	29	28	26.25
F7 T3 C3-C4 Fz Pz	23	27	20	31	26	29	31	29	27.00
C3-C4 F8 T4 Fz Pz	31	30	28	29	30	30	31	28	29.63
C3-C4 Fz Pz	30	30	29	30	29	28	30	29	29.38
F7 T3 T5 C3-C4 F8 T4 T6 Fz Pz	21	20	23	22	22	26	28	29	23.88
F7 C3-C4 F8 Fz Pz	23	29	29	27	30	28	31	29	28.25
F7 F3 F8 F4 Fz Pz	27	25	31	30	29	29	30	31	29.00
F7 F3 C3-C4 F8 F4 Fz Pz	27	27	28	30	28	30	31	30	28.88
F7 F3 F8 F4 Fz	25	26	27	27	27	29	29	28	27.25
F7 F3 F8 T4 F4 Fz Pz	28	25	29	29	28	29	30	29	28.38
F7 F3 C3-C4 F8 F4 Fz	27	26	23	28	27	29	29	28	27.13
F7 Fp1 F3 C3-C4 F8 Fp2 F4 Fz Pz	27	29	29	30	28	29	29	26	28.38
F7 Fp1 F3 F8 Fp2 F4 Fz Pz	29	29	29	29	28	29	29	28	28.75
F7 Fp1 F3 C3-C4 F8 Fp2 F4 Fz	28	28	28	28	28	28	27	28	27.88
F7 Fp1 F3 F8 Fp2 F4	27	28	28	28	28	28	28	29	28.00

Channel set [a]									Mean
Fz									
F7 F3 C3-C4 P3 F8 F4 P4 Fz Pz	27	28	27	21	29	29	29	29	27.38
F7 F3 P3 F8 F4 P4 Fz Pz	25	29	29	24	26	29	29	28	27.38
F7 F3 C3-C4 P3 Fz Pz	23	22	20	25	27	25	30	27	24.88
F7 F3 P3 Fz Pz	26	21	25	22	28	23	28	24	24.63
C3-C4 F8 F4 P4 Fz Pz	**28**	**31**	**29**	**27**	**30**	**32**	**31**	**29**	**29.63**
F8 F4 P4 Fz Pz	27	29	27	24	28	27	27	27	27.00
F7 F3 C3-C4 Fz Pz	26	27	28	22	29	29	30	29	27.50
F7 F3 Fz Pz	25	28	27	27	29	28	31	29	28.00
C3-C4 F8 F4 Fz Pz	**32**	**31**	**31**	**31**	**29**	**31**	**31**	**29**	**30.63**
F8 F4 Fz Pz	30	29	29	29	30	30	30	31	29.75
C3-C4 F8 T4 F4 P4 Fz Pz	28	30	27	23	31	30	31	27	28.38
F7 T3 F3 C3-C4 P3 Fz Pz	23	22	26	26	23	25	29	29	25.38
C3-C4 F8 T4 F4 Fz Pz	31	31	28	28	29	31	31	30	29.88
F7 T3 F3 C3-C4 Fz Pz	25	27	26	28	26	30	30	28	27.50
Fz Pz	27	29	29	29	29	27	28	30	28.50
Fz Cz Pz	27	27	22	26	28	28	28	27	26.63

[a] Channel sets according to the International 10-20 system for EEG leads; C3-C4, the channel of difference between two leads.

In bold and shaded in gray, higher agreement with DSM.

ERPs = event-related potentials; LV = latent variables.

The data from each investigated set of EEG channels were analyzed either with or without behavioral (ANT) and psychological (WISC) variables. They were submitted to an unsupervised method called Clustering of Variables around Latent Variables (CLV) [17,18]. The aim was to reduce the number of dimensions in the variables space by extracting latent variables (LVs), each of them associated with a cluster of the VOI variables. This clustering approach is based on correlation similarity indices between

variables and aims to identify directional clusters of variables. As it is, each extracted

LV, associated with a specific cluster of variables, is defined as the first (standardized)

principal component of the variables assigned to that cluster [21]. In our study, the

clusters of variables around one to four LVs were systematically considered and the best

number of LVs was determined according to the classification of subjects.

In fact, our ultimate purpose was to reindex the subjects into two groups

according to the objective behavioral, psychological, and neurophysiological data, all of

which were summarized around the extracted LVs. A k-means clustering method [27]

was performed on the subjects using the LVs. Each reclassification obtained (named

"output") was checked for agreement compared to the classification of subjects using the

DSM-IV-TR (called "DSM agreement"). The probability that agreement with DSM could

be observed under randomness was checked using the Chi-Square test for categorical

variables with one degree of freedom, and the expected value was 20 (absolute

randomness). The kappa index [28] was also evaluated to estimate the agreement level

with DSM (S2 method).

In order to determine whether reclassification was consistent, this study analyzed

which subjects were reclassified into another group other than his/her original DSM

group ("miss-reclassification"). This was performed for each specified agreement rate

with DSM, based on the outputs that provided this level of agreement. A measure of

"classifier power" was suggested by dividing the level of agreement with DSM and the

number of miss-reclassified subjects in any output with that agreement with DSM.

Subsequently, the outputs with the highest classifier power were considered the

most representative to estimate the validity of the DSM. Among all outputs, these were

submitted to cross-validation using the Leave-One-Out (LOO) strategy, where the

consistency of agreement with DSM *versus* LV was observed across the mean values of

the 39 LOO outputs, with 38 subjects each one. The output with the highest consistency

of agreement with DSM was regarded as the most representative.

Using the most representative outputs, a sensitivity and specificity analysis was

suggested for DSM criteria for ADHD.

The dataset with variables used in the multivariate analysis is available in the

supplementary material (S3 Dataset).

254 Results

255 Evaluating predictor and confounding variables

Control and ADHD groups showed significant differences in the quantitative

scores of DSM-IV-TR ($p < 0.05$) (Table 1). The significance level of the difference

between the two groups was higher for the inattention score than for the

hyperactivity/impulsivity score. Among variables considered as confounders, the I.Q.

scores in ADHD group were lower than in TD boys (p = 0.006) although they were

always higher than 80. All other variables, including learning rate, were not significantly

different between the two groups according to the Student's t-test (normality and

homoscedasticity were not statistically rejected).

264 Overview of Event-Related Potentials

265   Cue and target-related potentials with late peak latency (> 200ms, corresponding

266 to parietal P3) were observed in all channels. Peak amplitude varied from the maximum

267 value of 18µV for the frontopolar target ERP to the minimum value of 1µV for the

268 difference between C3 and C4 (Fig 1).

269 Data Mining and Reclassification of subjects

Values of agreement with DSM ranging from 0.51 (20 subjects, p = 0.125, χ^2 =

0.00, p = 1.000) to 0.82 (32 subjects, χ^2 = 7.20, p = 0.004) were found for all 40 channel

sets and LV sets (extracting 1 to 4 LVs), with or without behavioral variables enclosed

(Table 2). The *worst* mean agreement level between DSM and K-means reclassification

was 23.25 for [F7, T3, T5, Fp1, F3, C3-C4, P3, O1, Fz, Pz] channel set (χ^2 = 0.43, p =

0.528); and the best mean agreement level with DSM was 30.62 for [C3-C4, F8, F4, Fz,

Pz] channel set (χ^2 = 5.63, p = 0.010).

The two channel sets that manifested the highest agreement with DSM were [C3-
C4, F8, F4, Fz, Pz] (using 1 LV from ERP variables alone) and [C3-C4, F8, F4, P4, Fz,
Pz] (using 3 LVs from ERP and behavioral variables together). Table 2 shows all the
observed agreement rates for all channel sets.

As expected, there was an inverse relationship between the agreements with
DSM (from 20 to 32, according to the trial) and the total number of miss-reclassified
subjects (Fig 2). The highest agreement rate found (32 subjects) corresponded to the
highest classifier power (4.00).

Figure 2. Classifier Power of the MVA analysis. Number of miss-reclassified subjects per agreement
with DSM (number of subjects) in all 320 outputs, and the corresponding classifier power (numbers at the
graph marks). See methods.

Regarding an agreement level equal to 32, the LOO cross-validation showed that
only the output from the [C3-C4, F8, F4, Fz, Pz] channel set (using 1 LV from ERP
variables alone) was valid, and therefore, it was the most representative: the mean
agreement of all 39 outputs was 31.56 subjects (the highest mean agreement among all
others, see S4 Output), while the output from [C3-C4, F8, F4, P4, Fz, Pz] (using 3 LVs
from ERP and behavioral variables together) had a mean agreement equal to 29.87

subjects. The S4 Output has the agreement data of all outputs and their respective LOO

cross-validations (mean values of all 39 outputs).

Seven subjects in the selected output ([C3-C4, F8, F4, Fz, Pz] channel set, using

1 LV from ERP variables alone) were miss-reclassified: C005, C009, C012, C020, C023

(control subjects miss-reclassified to the ADHD group) and T010, T026 (ADHD subjects

miss-reclassified to the Control group). Based on these results, the estimated specificity

and sensitivity of the DSM-IVTR regarding biological variables were 75.00% and

89.47%, respectively, with accuracy of 82.05% corresponding to the kappa index of

0.75, i.e. *a substantial strength of agreement* [28].

304 Discussion

305   Our study tackled the possibility of examining attention deficit/hyperactivity

306   disorder using the DSM manual, whose criteria seemed to be biologically justified for

307   ADHD diagnosis with "substantial strength of agreement" regarding a specific pool of

308   behavioral, psychological, and neurophysiological variables. A larger source of

309   behavioral and neurophysiological data related to ADHD phenomenology was found in

the ANT, since this test is suitable to access the three dimensions of attention in the same setting [22, 29].

In order to classify subjects as normal or ADHD according to DSM criteria, caregivers were asked which criteria were well-defined characteristics of their children. Other assessment methodologies should also be used along with DSM for ADHD diagnosis both in scientific research and in clinical practice. Moreover, the manual should not be applied as a questionnaire in clinical practice. However, in this case, the object of evaluation is the DSM itself; it was thus the only clinical diagnostic tool used for the classification of subjects as either ADHD or not-ADHD. It was the most objective and controlled way to obtain the scores without interference from any other clinical impression. Supporting the present methodology, the ASRS (the WHO questionnaire for adult ADHD screening) was self-administered [30].

The most informative neurophysiological variables in this pool proved to be topographically asymmetrical and represented the binded information from ERPs in the three time intervals of interest from mid-frontal, mid-parietal, right frontal, and right anterior temporal sites (Fz, Pz, F4, and F8 channels), as well as early ERP components (45 to 290ms after target onset) from the 'C3 minus C4' channel. This asymmetry has already been shown to correlate with DSM scores, mainly with hyperactivity/impulsivity [31]. It was not the focus of the present investigation to establish models for ADHD or its neural mechanisms. However, our results revealed certain biological aspects of ADHD

regarding alterations in the frontal control over executive functions [21, 32, 33]. Several

studies have shown brain asymmetries related to ADHD phenomenology, especially in

the frontal-striatal network [33-35]. Asymmetrical topographic patterns were shown in

our previous studies by spectral and coherence analyses of the resting EEG, with signs

of relative inactivation of the frontal and *left* temporal cortex, known to be responsible for

voluntary attention, and which are impaired in ADHD patients [36]. These observations

could partially explain the leading role of ERP data from the right frontal regions in

discriminating between patients and controls in the present study.

Our results suggest that DSM-IV-TR is actually an effective and biologically

justified tool for ADHD diagnosis. Although a Kappa index of 0.75 should be considered

with reservations for some clinical settings, a clinical method such as DSM, with

sensitivity of 89.47% and specificity of 75% with regard to objective behavioral,

psychological, and neurophysiological measures, should be considered as particularly

relevant considering the complexity of mental disorders, especially those with

dimensional clinical phenomenology and without any validated biomarkers, so much so

that the only psychiatric tool for diagnosis and intervention is still phenomenological

examination, which must be systematically (although qualitatively) performed based on

manuals such as the DSM. Furthermore, the ADHD has been regarded as a

heterogeneous condition with inattentive, hyperactive, and combined subtypes, which

could be the answer for this level of agreement since these subtypes were not predicted

for reclassification. The existence of three discrete subtypes could lead to a flawed

reclassification.

The study was designed with a relatively small sample size. Any subject with the

slightest suspicion of comorbidity was excluded. Potentially confounding factors

commonly ignored were monitored (such as manipulating technologies that could lead to

the development of specific skills that would interfere with test performance and

socioeconomic niche that could determine psychological development). Therefore, the

sample should be regarded as well consistent with and representative of ADHD. On the

other hand, in order to classify the subjects according to their mental condition (a

multidimensional phenomenological complex) it would seem reasonable to gather a

considerable number of objective variables (as many as possible). Brain processing

during the ANT test as well as ADHD neurophysiology and its behavioral correlates in

this study were considered as potentially relevant features requiring further investigation.

Since it did not seem feasible to collect a sample with thousands of subjects evaluating

hundreds of variables in an experiment with optimal control of confounders and bias, the

dataset had to have a much smaller number of observations than variables. A small but

strictly controlled and well-exploited sample was considered preferable, even in

conditions of high prevalence such as the ADHD. Multivariate statistical modeling, such

as Linear Regression, Discriminant Analysis, or Factor Analysis, are well known as

unsuitable for this situation. Thus, in order to tackle a frequently encountered

methodological problem, often referred to as the curse of the dimensionality, an explorative data analysis suitable for small samples was adopted to identify a small number of synthetic latent variables. Instead of the Principal Component Analysis, which does not always provide easily interpretable latent variables (the principal components), an approach with clustering of variables was chosen to reduce data dimensionality to a small number of new latent variables. The CLV method [17,18] has already been applied to a wide range of research domains among which are pattern analysis [37], chemometrics [38], image analysis [39], and analysis scorecards in the health sector [40]. The results seemed promising for optimizing the experimental designs of clinical studies in which it is difficult to recruit large groups of patients.

The estimated I.Q. scores were different between the groups diagnosed using DSM-IV-TR, but this was not regarded as bias. Literature has shown that intelligence tests are sensitive to ADHD, and patient scores are generally lower than controls [41]. Even taking into account the above restrictions, there is no doubt that the data obtained seem to be quite consistent.

A major limitation of this study is the impossibility of testing all possible combinations among variables. Since the consistency of our paradigm lies on the fact that no biological / behavioral assumptions were considered in the grouping of variables, it is necessary to test them randomly grouped. However, there are more than one million possible combinations of EEG channels. The number of combinations grows

exponentially if the other variables are included individually. For computational

feasibility, sets of variables were arbitrarily chosen for analysis. Considering the eighty

sets studied, it can be assumed that DSM has at least a "substantial level of agreement"

with biological determinants. Perhaps some untested set of variables may manifest

higher agreement with the DSM, but the results found already satisfy our objective of

testing the validity of the DSM criteria for ADHD.

The aim of the present study was to draw attention to psychiatric approaches

based on diverse and multiple factors intervening in the biological mechanisms of

human behavior and its disorders. Therefore, we have found that there is a consistent

complex of biological information, which is in accordance with previous knowledge on

ADHD, that confers biological validity for DSM criteria for ADHD.

401 Aknowledgements

We would like to thank The Child´s Neurology program of the National Institute of

Women, Children, and Adolescents' Health Fernandes Figueira. We are also grateful to

Mrs. Maria S. Santana and Mrs. Aldenys Perez for the technical support. Our team is

especially grateful to our friend, colleague, and co-author Dr. Adailton Tadeu Pontes,

children's neurologist, who greatly inspired this article, and passed away during its
preparation.

References

1. American Psychiatric Association. Diagnostic and statistical manual of mental
disorders. 4th ed. Washington, DC: Author; 1994.

2. American Psychiatric Association. Diagnostic and statistical manual of mental
disorders. 5th ed. Arlington, VA: American Psychiatric Publishing; 2013.

3. Insel T. Director's Blog: Transforming Diagnosis. 2013 Apr 29 [cited 06 Jul 2018] In:
The National Institute of Mental Health Website. [about 2 screens] Available from:
http://www.nimh.nih.gov/about/director/2013/transforming-diagnosis.shtml.

4. Wakefield JC. DSM-5, psychiatric epidemiology and the false positives problem.
Epidemiol Psychiatr Sci. 2015 Jun;24(3):188-96. doi: 10.1017/S2045796015000116.

5. Rafalovich A. Exploring clinician uncertainty in the diagnosis and treatment of
attention deficit hyperactivity disorder.Sociol Health Illn. 2005 Apr;27(3):305-23.

6. Strauss V. An ADHD controversy in the mental health community. 2013 February 13
[cited 06 July 2018]. In: The Washington Post [Internet]. Available from:

https://www.washingtonpost.com/blogs/answer-sheet/post/an-adhd-controversy-in-the-

mental-health-community/2012/02/12/gIQAHJun9Q_blog.html.

7.Maser JD, Akiskal HS. Spectrum concepts in major mental disorders. Psychiatr Clin

North Am. 2002 Dec;25(4):xi-xiii.

8.Kirmayer LJ, Crafa D. What kind of science for psychiatry? Front Hum Neurosci. 2014

Jun 20;8:435. doi: 10.3389/fnhum.2014.00435.

9.Furman LM. Attention-deficit hyperactivity disorder (ADHD): does new research

support old concepts? J Child Neurol. 2008 Jul;23(7):775-84. doi:

10.1177/0883073808318059.

10. Freitas-Silva LR, Ortega F. Biological determination of mental disorders: a

discussion based on recent hypotheses from neuroscience. Cad Saude Publica. 2016

Aug 29;32(8):e00168115. doi: 10.1590/0102-311X00168115.

11.Scarr E, Millan MJ, Bahn S, Bertolino A, Turck CW, Kapur S, Möller HJ, Dean B.

Biomarkers for Psychiatry: The Journey from Fantasy to Fact, a Report of the 2013

CINP Think Tank. Int J Neuropsychopharmacol. 2015 Apr 21;18(10):pyv042. doi:

10.1093/ijnp/pyv042.

12. Bochkarev VK, Lazarev VV, Nikiforov AI, Paniushkina SV, Severnyĭ AA. Clinico-

electroencephalographic correlations in asthenoadynamic subdepressions.Zh

Nevropatol Psikhiatr Im S S Korsakova. 1987;87(4):564-70.

13.Lazarev VV. The relationship of theory and methodology in EEG studies of mental

activity.Int J Psychophysiol. 2006 Dec;62(3):384-93. Epub 2006 Mar 10.

14. Lazarev VV, Lebedeva IS, Tsutsul'kovskaia MIa, Severnyĭ AA, Pozharitskaia DA.

Specificity of electroencephalographic correlations of asthenic disorders in atypical

juvenile asthenic-depressive conditions. Zh Nevropatol Psikhiatr Im S S Korsakova.

1990;90(10):73-7.

15. Monakhov K, Perris C, Botskarev VK, von Knorring L, Nikiforov AI. Functional

interhemispheric differences in relation to various psychopathological components of the

depressive syndromes. A pilot international study.Neuropsychobiology. 1979;5(3):143-

55.

16. Schwarz E, Izmailov R, Spain M, Barnes A, Mapes JP, Guest PC, Rahmoune H,

Pietsch S, Leweke FM, Rothermundt M, Steiner J, Koethe D, Kranaster L, Ohrmann P,

Suslow T, Levin Y, Bogerts B, van Beveren NJ, McAllister G, Weber N, Niebuhr D,

Cowan D, Yolken RH, Bahn S. Validation of a blood-based laboratory test to aid in the

confirmation of a diagnosis of schizophrenia. Biomark Insights. 2010 May 12;5:39-47.

17.Vigneau E, Qannari EM. Clustering of Variables Around Latent Components.

Communications in Statistics - Simulation and Computation. 2003; 32: 1131-1150.

18.Vigneau E, Chen M, Qannari E M. ClustVarLV: An R Package for the Clustering of

Variables Around Latent Variables. The R Journal. 2015 ; 7 : 134-148.

19. Matuschek T, Jaeger S, Stadelmann S, Dölling K, Grunewald M, Weis S, von

Klitzing K, Döhnert M. Implementing the K-SADS-PL as a standard diagnostic tool:

Effects on clinical diagnoses. Psychiatry Res. 2016 Feb 28;236:119-24. doi:

10.1016/j.psychres.2015.12.021.

20.Hruby T, Marsalek P. Event-related potentials--the P3 wave. Acta Neurobiol Exp

(Wars). 2003;63(1):55-63.

21. Barry RJ, Johnstone SJ, Clarke AR. A review of electrophysiology in attention-

deficit/hyperactivity disorder: II. Event-related potentials. Clin Neurophysiol. 2003

Feb;114(2):184-98.

22. Fan J, McCandliss BD, Sommer T, Raz A, Posner MI. Testing the efficiency and

independence of attentional networks. J Cogn Neurosci. 2002 Apr 1;14(3):340-7.

23.Kratz O, Studer P, Malcherek S, Erbe K, Moll GH, Heinrich H. Attentional processes

in children with ADHD: an event-related potential study using the attention network test.

Int J Psychophysiol. 2011 Aug;81(2):82-90. doi: 10.1016/j.ijpsycho.2011.05.008.

24.Wechsler D. Wechsler Intelligence Scale for Children(WISC-III): Manual. 3th ed. San

Antonio: The Psychological Corporation; 1991.

25.Mello CB, Argollo Shayer PBM, Abreu N, AbreuII N, GodinhoIII K, Durán P, Vargem

F, Muskat M, Miranda MC, Bueno OFA. Abbreviated version of the WISC-III: correlation

between estimated IQ and global IQ of Brazilian children, Psicologia: Teor e Pesq. 2011;

27 (2): 149-155. dx.doi.org/10.1590/S0102-37722011000200002.

26. Abramov DM, Pontes M, Pontes AT, Mourao-Junior CA, Vieira J, Quero Cunha C,

Tamborino T, Galhanone PR, deAzevedo LC, Lazarev VV. Visuospatial information

processing load and the ratio between parietal cue and target P3 amplitudes in the

Attentional Network Test. Neurosci Lett. 2017 Apr 24;647:91-96. doi:

10.1016/j.neulet.2017.03.031.

27.Gore Jr PA. Cluster Analysis, In: Tinsley HEA, Brown SD, editors. Handbook of

Applied Multivariate Statistics and Mathematical Modeling. San Diego, CA: Academic

Press; 2000. p. 297-321.

28. Landis JR, Koch GG. The measurement of observer agreement for categorical data.

Biometrics. 1977 Mar;33(1):159-74.

29. Petersen SE, Posner MI. The attention system of the human brain: 20 years after.

Annu Rev Neurosci. 2012;35:73-89. doi: 10.1146/annurev-neuro-062111-150525.

30. Kessler RC, Adler L, Ames M, Demler O, Faraone S, Hiripi E, Howes MJ, Jin R,

Secnik K, Spencer T, Ustun TB, Walters EE. The World Health Organization Adult

ADHD Self-Report Scale (ASRS): a short screening scale for use in the general

population. Psychol Med. 2005 Feb;35(2):245-56.

31. Abramov DM., Cunha CQ, Pontes MC, Galhanone PR, deAzevedo LC, Lazarev VV.

Functional Asymmetry In The Central Brain Regions In Boys With Attention Deficit

Hyperactivity Disorder Detected By Event Related Potentials During Performance Of

The Attentional Network Test. 2016. Preprint.Avaiable on BioRxiv,

https://doi.org/10.1101/118380 ; cited on 06 July 2018.

32. Barkley RA. Behavioral inhibition, sustained attention, and executive functions:

constructing a unifying theory of ADHD. Psychol Bull. 1997 Jan;121(1):65-94. doi:

10.1037/0033-2909.121.1.65.

33. Barkley RA. Attention-Deficit/Hyperactivity Disorder: handbook for diagnosis and

treatment. 3th ed. New York: The Guilford Press; 2006. p 32-33.

34. Langleben DD, Austin G, Krikorian G, Ridlehuber HW, Goris ML, Strauss HW.

Interhemispheric asymmetry of regional cerebral blood flow in prepubescent boys with

attention deficit hyperactivity disorder. Nucl Med Commun. 2001 Dec;22(12):1333-40.

35.Dang LC, Samanez-Larkin GR, Young JS, Cowan RL, Kessler RM, Zald DH.

Caudate asymmetry is related to attentional impulsivity and an objective measure of

ADHD-like attentional problems in healthy adults. Brain Struct Funct. 2016

Jan;221(1):277-86. doi: 10.1007/s00429-014-0906-6.

36. Lazarev VV, Pontes M, Pontes AT, Vieira J, Quero-Cunha C, Tamborino T,

deAzevedo LC, Abramov DM. EEG and ERP characteristics of Attention Deficit

Hyperactivity Disorder in children and adolescents [abstract]. Int. J. Psychophysiol. 2106; 108: 75.

37. Vigneau E, Sahmer K, Qannari EM, Bertrand D. Clustering of variables to analyze spectral data. J Chemometrics, 2005; 19: 122-128.

38. Vigneau E, Qannari EM, Navez B, Cottet V. Segmentation of consumers in preference studies while setting aside atypical or irrelevant consumers. Food Quality Pref. 2005; 47: 54-63.

39. Legland D, Beaugrand J. Automated clustering of lignocellulosic fibres based on morphometric features and using clustering of variables. Industrial Crops and Products; 2013; 45: 253– 261.

40. Lovaglio PG. Model building and estimation strategies for implementing the Balanced Scorecard in Health sector. *Quality & Quantity*. 2011*;* 45: 199–212.

41. Mackenzie GB, Wonders E. Rethinking Intelligence Quotient Exclusion Criteria Practices in the Study of Attention Deficit Hyperactivity Disorder. Front Psychol. 2016 May 31;7:794. doi: 10.3389/fpsyg.2016.00794.

S1 table. Time intervals of Interest. Windows holding Cue-related potentials (CUE), contingent voltage variation (CVV), and Target-related potentials (TGT) for each channel

(ms), to calculate peak amplitude, peak latency, and mean amplitude. Zero ms refers to

Target onset (see Figure 1).

S2 method. Calculating the Kappa Index. Inference of the magnitude of agreement

between two categorical data (in this case, subjects' classification according to DSM and

to biological/behavioral variables in the dataset).

S3 dataset. DSM scores and variables of interest. *In the first sheet:* innatention,

hyperactive+impulsive, and total scores for ADHD according to DSM-IV-TR; *second*

sheet: all data concerning (1) scores of the WISC-III subtests, (2) ANT performance (AC:

accuracy, RT: reaction time and IVRT: intraindividual variation in reaction time, which is

the standard deviation of the subject's RT) for each experimental condition (Allcon:

mean value of all conditions, NtCue: neutral cue condition, SpCue: spatial cue condition,

ConTg: congruent target condition, IncTg: incongruent target condition) and attentional

network (alerting, orienting, and executive) plus learning indexes (IVRT or RT at the last

ANT block/first ANT block), and (3) the neurophysiological measures (peak amplitudes

and latencies, and mean amplitudes inside each time interval of interest in S1) for the

EEG channel set.

S4 output. DSM versus Biological agreement. The channel sets (rows) and number

of latent vectors extracted (rows) holding ERPs variables alone or ERPs + behavioral

(ANT and WISC) variables. The first block (on the left) are outputs from all 39 subjects.

The second block (on the right) are the averages of outputs with n-1 subjects (LOO

cross-validation).

Classifier Power