Title: Cell type boundaries organize plant development

Authors: Monica Pia Caggiano[1#], Xiulian Yu[1#], Neha Bhatia[1#], André Larsson[2], Hasthi

Ram[1], Carolyn Ohno[1], Pia Sappl[1], Elliot M. Meyerowitz[3], Henrik Jönsson[2,4,5], Marcus G.

Heisler[1,6 *]

Affiliations:

[1] European Molecular Biology Laboratory, Meyerhofstrasse 1, D-69117

Heidelberg, Germany

[2] Computational Biology and Biological Physics, Department of Astronomy and

Theoretical Physics, Lund University, Sweden

[3] Division of Biology and Biological Engineering and Howard Hughes Medical Institute,

156-29, California Institute of Technology, Pasadena, CA 91125, USA.

[4] Sainsbury Laboratory, University of Cambridge, Cambridge, UK

[5] Department of Applied Mathematics and Theoretical Physics, University of Cambridge,

UK

[6] School of Life and Environmental Sciences, University of Sydney, NSW, Australia

# - these authors contributed equally to this work

* - author for correspondence: Marcus G Heisler (marcus.heisler@sydney.edu.au)

Once sentence summary: Cell type boundaries regulate plant development

Abstract: In plants the dorsoventral boundary of leaves defines an axis of symmetry

through the centre of the organ separating the future top (dorsal) and bottom (ventral)

tissues. Although the positioning of this boundary is critical for leaf morphogenesis, the

mechanism by which it is established has not been determined. Here we show that

dorsoventral boundaries in leaves derive from a boundary between Class III

homeodomain-leucine zipper and KANADI expression in the shoot apical meristem,

confirming that leaf founder cells are pre-patterned with respect to their dorsoventral axis.

Furthermore, perturbation of this boundary not only alters leaf orientation and

morphogenesis but also organ position, revealing a general role for these boundaries in

plant development. Lastly, we show that auxin stabilizes boundary position at organ

inception and that wounds disrupt this process. Overall our findings reveal a two-way

interaction between cell type boundaries and auxin that controls fundamental aspects of

plant architecture.

Introduction

Lateral organ development in plants and animals typically involves several processes

occurring in a coordinated manner. These include organ positioning, the specification of

different cell types and organ morphogenesis. Spatial cues specifying these processes are

usually provided by a molecular pre-pattern present in precursor tissues, or from

inductive signals emanating from neighboring regions. Unlike animals however, plant

organs such as leaves arise continuously in regular patterns around the shoot apical

meristem (SAM). These phyllotactic patterns are determined by a self-organizing,

iterative process based upon feedback between transcriptional targets of the plant

hormone auxin and polar auxin transport (Bhatia et al. 2016). Nevertheless, certain

features of leaves are relatively constant including the restriction of their formation to the

meristem periphery and their flattened, dorsoventral (top-bottom) orientation with respect

to the shoot apex. The spatial cues that specify these features must therefore be provided

despite the constant displacement of cells away from the SAM and varying leaf position.

Supporting this proposal, wounding experiments involving the isolation of leaf primordia

from the meristem have suggested the presence of an inductive signal from the meristem

that promotes dorsal identity within leaf primordia at the time of organ initiation

(Reinhardt et al., 2005; Sussex, 1951). A variant on this theme is the proposal that auxin

depletion in the adaxial (adjacent to the shoot axis) tissues of leaf primordia promotes

dorsal identity (Qi et al., 2014). In contrast, other studies suggest that dorsoventral

pattering is pre-established, being directly derived from central-peripheral patterning of

the shoot (Hagemann and Gleissberg, 1996; Husbands et al., 2009; Kerstetter et al.,

2001). Overall, the nature of the initial positional information that specifies leaf

dorsoventrality remains unresolved (Kuhlemeier and Timmermans, 2016).

Besides influencing leaf differentiation and shape, genes involved in leaf dorsoventrality

influence leaf position (Izhaki and Bowman, 2007). For instance, Arabidopsis plants

mutant for the KANADI (KAN) genes develop leaves ectopically from the hypocotyl and

leaf tissues while plants mutant for the Class III HD-ZIP (HD-ZIPIII) genes develop

leaves from the center of the shoot (Izhaki and Bowman, 2007). These observations

indicate that the developmental mechanisms that specify leaf dorsoventrality may also be

involved in organ positioning although how these processes relate is unclear.

In this study we investigate the origin of dorsoventral patterning in detail. We find

evidence of a developmental pre-pattern of HD-ZIPIII and KAN gene expression in the

meristem periphery which, by restricting auxin activity, specifies both the locations of

lateral organs as well as their dorsoventrality. We also show that in addition to triggering

organ initiation, high auxin levels maintain dorsal cell identity as organs initiate. These

results reconcile previous findings and reveal a more general role for HD-ZIPIII / KAN

boundaries in determining plant architecture.

Results

Genes involved in leaf dorsoventrality pre-pattern organ formation in the shoot

To better understand the establishment of dorsoventral patterning in developing organs

we examined the expression of several genes involved in specifying leaf dorsoventrality

in the shoot apical meristem (SAM) using functional fluorescent protein reporters. We

found these proteins to be localized in non-overlapping concentric patterns with the

dorsal Class III HD-ZIP protein REVOLUTA (REV) (Otsuga et al., 2001) being detected

centrally, as marked by the expression of REVp::REV-2×YPet (REV-2×YPet), and the

ventral protein KANADI1 (KAN1) (Kerstetter et al., 2001) expressed peripherally, as

marked by the expression of KAN1p::KAN1-2×GFP (KAN1-2×GFP) (Figure 1, A to E).

Imaging PIN1-CFP together with these reporters revealed regions of high PIN1-CFP

expression, marking positions of organ inception (Heisler et al., 2005; Reinhardt et al.,

2003), to be centered on a narrow region located between the dorsal and ventral domains

(Figure 1D-K). This arrangement was conserved with few cells altering their KAN1

expression status during organogenesis (Figure 1C arrowhead and Figure 2 A-F).

Ventrally expressed MIR165/166 (Kidner and Martienssen, 2004; Merelo et al., 2016;

Nogueira et al., 2007; Yao et al., 2009) also appeared active in the SAM periphery as

marked by a *MIR166Ap::GFPER* reporter and MIR165/166 biosensor (Figure 3A-I),

consistent with previous studies (Miyashima et al., 2013). Both KAN1-2×GFP and

MIR166Ap::GFPER re-established their expression in the SAM periphery after organ

outgrowth (Figure 1E and Figure 3C). As members of the WOX family of transcription

factors are expressed at dorsoventral boundaries (Nakata et al., 2012) we imaged

functional translational fusions to both PRS and WOX1 to examine their expression in

the shoot relative to leaves. We found that although WOX1-2×GFP expression was

limited to the margins and middle domain of leaves (Figure 4A-C), in the vegetative

shoot PRS-2×GFP expression extended from the leaf middle domain to surround the

SAM in a region between the KAN1 and REV expression domains (Figure 4D-H). In the

inflorescence meristem however, PRS-2×GFP expression was restricted to early floral

primordia (Figure 4I and J). In contrast to all genes described so far, a FILp::dsRED-N7

marker was expressed in both abaxial and adaxial cells at inception, consistent with a

previous study (Tameshige et al., 2013) (Figure S1). Overall these data reveal that in

many respects although not all, dorsoventral patterning within young leaf primordia,

including the middle region, corresponds with central-peripheral patterning in the SAM.

The KAN and Class III HD-ZIP genes repress organ initiation where they are

expressed

The correlation between organ positioning and KAN-HD-ZIPIII expression boundaries as

well as previous genetic data reporting ectopic organs in *kan1 kan2 kan4* and *rev phb phv*

mutants (Izhaki and Bowman, 2007) demonstrate a role for these genes in the repression

of leaf initiation. However it is possible that the HD-ZIPIII genes influence organ

initiation only indirectly by promoting SAM formation during embryogenesis (Emery et

al., 2003). To distinguish between these possibilities we induced the expression of

MIR165/166-resistant REVr-2×VENUS throughout the epidermis using the ATML1

promoter (Sessions et al., 1999) and found that it caused an arrest of organ formation and

repression of KAN1-2×GFP in both the vegetative and inflorescence meristems (Figure 5

A-E). Similar results were observed after induction of a short tandem target mimicry

construct designed to repress MIR165/166 activity (Yan et al., 2012) (Figure 5F and G)

or after epidermal induction of MIR165/166-resistant PHAVOLUTA (Figure 5H-J).

Similarly, plants expressing KAN1 ectopically in the epidermis also stopped making new

organs (Figure 5K and L) and the inflorescence meristem took on a dome shape before

eventually arresting (Figure 5M). We conclude both the KAN and Class III HD-ZIP

genes regulate organ positioning at least in part by repressing organ initiation where they

are expressed.

Expression patterns of REV and KAN1 in the shoot regulate leaf positioning and

morphogenesis

To test whether boundaries between KAN1 and Class III HD-ZIP expression in the SAM

can play an instructive role in positioning new organs and determining their subsequent

dorsoventrality, we induced KAN1-2×GFP expression ectopically at the center of the

SAM using a pOp6/LhGR two-component (Samalova et al., 2005) system and the CLV3

promoter driving LhGR. After KAN1-2×GFP induction, most seedlings initiated several

new leaves before their growth stopped. Confocal imaging of seedlings five days after

stratification (DAS) on dexamethasone (DEX) induction medium revealed that new

organs, marked by high levels of PIN1-CFP expression, formed ectopically at the

perimeter of an enlarged and irregular central domain of induced KAN1-2×GFP

expression, in which REV-2×YPet expression had been repressed (Figure 6A and B).

Although ectopic KAN1-2×GFP was only detected within or bordering organs during

their initiation, REV-2×YPet expression was often restricted during later developmental

stages (Figure 6C-H) indicating that patterns of KAN1 gene expression within the SAM

can influence subsequent organ development. In particular, we noted that the distal

margins of developing leaf primordia always correlated with boundaries of REV

expression within the epidermis, even when REV expression was abnormally restricted

(arrow heads in Figure 6C-H; movie S1). Several classes of phenotype, including leaves

with an inverted orientation, could be distinguished at maturity (Figure 6 I-O and

Supplementary Figure S2) that correlated with the patterns of REV-2×YPet expression

observed during early development. These distinct morphological classes can be

explained according to the configuration of the HD-ZIPIII-KAN boundaries at organ

inception. Specifically, we infer that the number and orientation of boundaries within

organ founder cells (as specified by high auxin levels) determines the configuration of

later leaf marginal tissue (Figure 6P-S).

Maximal auxin response is localized to HD-ZIPIII KAN boundaries

The influence of dorsoventral gene expression on organogenesis suggests that the

boundaries between their expression domains may function generally to localize auxin

response. In support of this proposal it has been reported that in the inflorescence

meristem the auxin transcriptional marker DR5 is only responsive to auxin in the shoot

periphery (de Reuille et al., 2006). We examined DR5 expression (Liao et al., 2015) in

the vegetative meristem by examining its expression both in the wild type and after 1-N-

Naphthylphthalamic acid (NPA) treatment at 4 DAS. In mock treated seedlings we found

DR5 to be expressed at the locations of incipient primordia and at the distal tip of existing

leaf primordia of (Figure 7A and B). In contrast, NPA treated seedlings expressed DR5 in

a ring encircling the SAM prior to organ emergence and in the middle domain or

dorsoventral boundary region of developing leaves (Figure 7C and D). Similar

experiments with seedlings expressing the ratiometric R2D2 intracellular auxin reporter

(Liao et al., 2015) revealed a generally broader distribution of auxin compared to the

DR5 transcriptional response, especially after NPA treatment, where signal was not

restricted to the meristem periphery or leaf tip (Figure 7E-H). Given the similarity in

expression between auxin-induced DR5 expression and PRS expression we tested

whether PRS as well as WOX1 are auxin inducible and found that both genes respond to

auxin treatment within 12 and 15 hrs respectively although their response, like DR5, was

mainly restricted to the boundary region (Figure 7I-P). Measuring transcript levels using

qPCR indicated that the auxin response for both genes occurs at the transcriptional level

(Figure 7Q).

The absence of HD-ZIPIII and KAN expression in boundary regions may mediate

boundary function

The relatively low level or absence of KAN1 and REV expression observed within

boundary regions and the finding that both genes repress organ formation where they are

expressed suggests a simple model in which organ formation is localized to boundary

regions due to the local absence of KAN and HD-ZIPIII expression in boundary cells. We

tested this proposal *in silico* by implementing a previous model for phyllotaxis that

incorporates the polarization of PIN1 towards cells with high intracellular auxin

concentrations (Bhatia et al., 2016; Jonsson et al., 2006; Smith et al., 2006). By assuming

that both KAN1 and REV repress auxin-induced transcription and by including a narrow

region of cells located between the KAN1 and REV expression domains the model was

able to self-organize the periodic formation of auxin maxima along the boundary as

predicted, compared to a broader distribution of maxima when such boundaries are not

included (Figure 8A-D; compare 8B to 8D).

Auxin is required to maintain adaxial cell identity during organ initiation

So far, our results indicate that both HD-ZIPIII and KAN1 suppress auxin-induced organ

formation in tissues where they are expressed. However this does not exclude the

possibility that auxin may play a role in patterning HD-ZIPIII and KAN expression.

Indeed, REV expression typically extends towards PIN1 polarity convergence patterns in

the meristem periphery and disappears in axil regions where auxin is depleted, suggesting

a direct or indirect role for auxin in promoting REV expression (Figure 1) (Heisler et al.,

2005). To investigate any potential regulation of REV and KAN expression by auxin we

treated inflorescence meristems with a combination of NAA and NPA and found that the

region in between the REV and KAN1 domains appeared to narrow due to a slight

expansion of REV expression (Figure 9A and B). To test whether REV depends on high

auxin levels for its usual pattern of expression we treated triple marker plants with the

TIR1 antagonist auxinole (Hayashi et al., 2012) as well as auxin synthesis inhibitors

yuccasin (Nishimura et al., 2014) and kyn (He et al., 2011). Although treatment with

these drugs individually did not lead to an obvious response, a combination of all three

led to a cessation of organ production and an expansion and restriction of KAN1 and

REV expression centrally, respectively (Figure 9C and D). KAN expression not only

expanded into those cells originally located between the REV and KAN1 domains but

also cells that had previously been expressing REV at high levels. However KAN1

expression remained excluded from the central most cells of the shoot and established

floral primordia, where REV expression still remained (Figure 9D). Extending the drug

treatment time did not lead to a further change in expression. A similar response occurred

in the vegetative meristem (Supplementary Figure S3). Monitoring the ratiometric auxin

sensor R2D2 in response to a similar combined inhibitor treatment indicated a strong

decrease in overall intracellular auxin levels, including in the central meristem regions

where REV was still expressed (Figure 9E and F). Overall these data indicate that high

levels of auxin are necessary to promote REV expression in cells entering the peripheral

zone, thereby promoting the establishment of dorsal cell types within initiating primordia.

However this influence appears restricted to those cells not already expressing KAN1.

Wounding causes auxin depletion dependent abaxialization

Since wounding causes repolarization of PIN1 away from the wound site (Heisler et al.,

2010) resulting in auxin depletion (Landrein et al., 2015) and auxin is required to

maintain REV at the expense of KAN expression in peripheral meristem tissues, we

investigated whether wounding results in ectopic KAN1 and reductions in REV

expression. Firstly, using a pulsed IR laser to ablate cells adjacent to young organ

primordia we confirmed that auxin levels decrease in the vicinity of wounds by

monitoring the expression of the R2D2 ratiometric marker (Figure 10A and B). Next, we

monitored REV, KAN and PIN1 expression in response to such wounds. We found that

KAN1 became expressed in cells adjacent to the wound on either side, regardless of

wound orientation with respect to the SAM (Figure 10C and D – figure supplement 4A-

H). Such a response argues against the possibility that ectopic KAN is the result of

interruption of a signal emanating from the meristem that promotes dorsal and represses

ventral identity (Sussex, 1955). Instead, it supports the proposal that KAN1 expression is

promoted in the vicinity of wounds in general, possibly due to low auxin levels. To test

this hypothesis, we repeated these experiments while treating the wounded meristems

with combinations of NAA and NPA over a 48 hr period and found that the induction of

KAN1 expression around wounds could be completely eliminated if NAA and NPA were

combined (Figure 10E and F; figure supplement 4I-M). Although a similar response to

wounding was found to occur in vegetative meristems, the wound response typically

involved a more substantial reorganization of meristem structure, possibly due to the

small size of the vegetative meristem relative to the wounds (figure supplement 5A-C).

When new leaves subsequently formed, they were always properly oriented with respect

to the new meristem organization.

Discussion

In this study we shed new light on a long-standing question regarding leaf dorsoventrality

in plants – when and how is it established? The early work of Sussex, based on

histological analysis and wounding experiments, suggested that initiating leaf primordia

require an inductive signal from meristem tissues to specify dorsal cell fate (Sussex,

1955). This proposal has been further supported by a more recent study in tomato

(Reinhardt et al., 2005). In contrast, other workers in the field have claimed that

dorsoventrality arises directly from radial patterning of the shoot (Hagemann and

Gleissberg, 1996). Our results reveal that organs are pre-patterned by domains of KAN

and HD-ZIPIII expression in the shoot. The effect of ectopic KAN1 expression at the

center of the shoot on subsequent leaf positioning and growth is particularly striking and

indicates that the spatial arrangement of HD-ZIPIII and KAN expression present within

organ founder cells is incorporated into organs as they initiate and directs patterns of

morphogenesis. The propagation of this pattern into initiating organs requires high auxin

levels at early stages of development. If auxin levels decrease during organ

establishment, REV expression also decreases and KAN1 expression extends towards the

meristem center. Thus, the same signal that triggers organ initiation is also required for

proper organ differentiation. Nevertheless, at a later stage of development REV

expression in both leaf and floral primordia becomes auxin independent. Overall all our

results imply that leaf dorsoventrality results from a four-step process: (i) signals during

embryonic development establish concentric patterns of Class III HD-ZIP and KAN gene

expression (Zhang et al., 2017); (ii) the boundary between these domains helps to define

sites for auxin-dependent organogenesis i.e. the meristem peripheral zone; (iii) high

levels of auxin at organ initiation sites enables the maintenance of Class III HD-ZIP

expression in adaxial (located towards the central shoot axis) primordial cells as these

cells are displaced from the central zone; (iv) patterns of HD-ZIPIII and KAN gene

expression in developing flowers and leaves are stabilized and dictate future patterns of

organ morphogenesis.

How do DV boundaries control organ position and shape? Both our data as well as

previous studies indicate this is through the regulation of auxin perception. For instance,

organ initiation requires high auxin levels but auxin can only trigger organogenesis in the

peripheral zone (Reinhardt et al., 2000) where the HD-ZIPIII/KAN boundary occurs. A

similar relationship holds for leaves where auxin application results in growth but only

from the leaf margins, again corresponding to a DV boundary (Koenig et al., 2009). We

further show that the localization of auxin response to DV boundaries applies generally,

i.e. DR5 expression appears higher at such boundaries compared to the broader predicted

auxin distribution, as marked by R2D2. How is this restriction achieved? Imaging data

reveals that the locations of PIN1 polarity convergences correspond to a region of cells in

which the expression of both HD-ZIPIII and KAN is low or absent. Given genetic and

molecular data indicating that both the HD-ZIPIII and KAN genes repress auxin activity

(Huang et al., 2014; Merelo et al., 2013; Zhang et al., 2017), we propose that the absence

of their expression in boundaries results in a local de-repression of auxin-induced

transcription. In turn, this de-repression would be expected to potentiate feedback

between auxin signalling and cell polarity including microtubule orientations that would

result in localized outgrowth at the boundary (Bhatia et al., 2016). Such growth may

either occur in the periodic fashion typical of phyllotaxis and complex leaves, or in a

more continuous manner typical of simple leaves, depending on the strength of auxin

transport or other modifications to the feedback system (Bilsborough et al., 2011; Koenig

et al., 2009).

The conclusions stated above contrast with those of another study reporting that high

levels of auxin inhibit dorsal fate and promote ventral cell fate (Qi et al., 2014). These

findings were based on auxin application experiments in tomato that resulted in the

ventralization of leaves as well as the observation that Arabidopsis *pin1* and *pid rev*

double mutants produce trumpet or rod-shaped shaped leaves. Although our results

cannot easily explain the tomato data, since high auxin levels are required to maintain

REV expression during organ establishment, we would expect that in *pin* and *pid*

mutants, lower auxin levels may well result in lower expression levels of REV and

possibly other Class III HD-ZIPs, leading to leaf ventralization as reported in these

mutant backgrounds (Qi et al., 2014). Further auxin application experiments in tomato

that include an analysis of gene expression and auxin level changes may clarify the

relationship between auxin and dorsoventral patterning in tomato compared to

Arabidopsis.

The regulation of HD-ZIPIII and KAN expression by auxin is not only relevant to

understanding wild type organ development but also for understanding the reorganization

of tissue types in response to wounds. Wounds in the meristem outer cell layer

specifically alter cell polarities such that PIN1 becomes polarized away from wounds in

adjacent cells (Heisler et al., 2010), leading to auxin depletion (Landrein et al., 2015).

Since the 1950s wounding has also been associated with changes in leaf dorsoventrality

(Sussex, 1955) and changes to organ position (Snow, 1931). Specifically, wounds located

between the meristem and initiating organ were found to promote leaf ventralization and

a loss of lamina shape suggesting that the establishment of dorsal cell fate requires an

inductive signal from the meristem, which the wound interrupted (Sussex, 1955). Our

observations now indicate rather, that during organ establishment, HD-ZIPIII expression

depends on high levels of auxin, which wounding disrupts. Thus as suggested previously,

the meristem "acts by maintaining in the leaf-forming zone some polarized micro-

structure which collapses without it" (Snow and Snow, 1959).

Materials and Methods

Plant material

324   Plants were grown on soil at 22 °C in continuous light-conditions and cultivated either on

325   soil or on GM medium (1 % sucrose, 1× Murashige and Skoog basal salt mixture, 0.05%

326   MES 2-(MN-morpholino)-ethane sulfonic acid, 0.8 % Bacto Agar, 1 % MS vitamins, pH

327   5.7 with 1 M potassium hydroxide solution).

Construction of Transgenes

Multiply transgenic lines were generated by Agrobacterium-mediated transformation into

stable transgenic lines or by genetic crossing. The *FILp::dsREDN7* and *PIN1p::PIN1-*

GFP transgenes have been described elsewhere (*16*). *pREV::REV-2×VENUS* in the T-

DNA vector *pMLBART* (Gleave, 1992) is a modification of *pREV::REV-VENUS* (Heisler

et al., 2005) that contains a translational fusion to 2 tandem copies of the fluorescent

protein VENUS (Nagai et al., 2002). *REVp::REV-2×Ypet* containing a C-terminal fusion

to the 2× Ypet (Nguyen and Daugherty, 2005) in *pMOA36* T-DNA (Barrell and Conner,

2006) was transformed into a *PINp::PIN1-CFP* line (Gordon et al., 2007). The

KAN1p::KAN1-2×GFP transgene in pMOA34 T-DNA was created by amplifying 8.7 kb

of *KAN1* (At5g16560) genomic sequences with primers KAN1g F and KAN1g R

(Supplementary Table S1) as a translational fusion to a 9 Ala linker and 2×GFP followed

by *OCS* terminator sequences. When transformed into *kan1-2 kan2-1* segregating plants,

this construct complements the mutant phenotype. The triple marker line was generated

by transforming *KAN1p::KAN1-2×GFP* into a *REVp::REV-2×Ypet; PIN1p::PIN1-CFP*

transgenic line. *KAN1p::KAN1-2×CFP* or *KAN1p::KAN1-2×Ypet* containing a fusion to

2 copies of CFP or Ypet, respectively, were constructed similarly. *KAN1p::KAN1-*

2×Ypet and *PIN1p::PIN1-GFP* were combined in T-DNA vector *BGW* (Karimi et al.,

2002) by Gateway technology (Invitrogen) for generation of a double marker transgenic

line.

The *KAN1* cDNA was amplified by PCR with primers K1 cDNA F and K1 cDNA R to

generate C-terminal translational fusion to a 9 Ala linker followed by single GFP or

2×GFP followed by pea *rbcS E9* terminator sequence (Zuo et al., 2001) and cloned into

the pOp6/LhGR two-component system (Craft et al., 2005) for dexamethasone-inducible

misexpression.

An *ATML1p::LhGR* driver containing 3.4 kb of the L1-specific *ATML1* gene

(At4g21750) fused to the chimeric LhGR transcription factor and a *6Op::KAN1-GFP*

expression construct in a *pSULT* sulfadiazine-resistant T-DNA vector (*ATML1>>KAN1-*

GFP) was generated.   The *pSULT* T-DNA vector was derived from *pMLBART* by

replacing the *NOSp::BAR* gene with *1'-2'p::*SUL (Rosso et al., 2003), a plant selectable

marker that confers resistance to sulfadiazine herbicide to create *pSULT. A*

CLV3p::LhGR driver containing 1.49 kb of upstream regulatory sequences was PCR

amplified with primers CLV3p F and CLV3p R along with 1.35 kb of downstream

regulatory sequences with primers CLV3utr F and CLV3utr R was combined with

6Op::KAN1-2×GFP in *pSULT* T-DNA vector (*CLV3>>KAN1-2×GFP*).

ATML1>>REVr-2×VENUS is a sulfadiazine-resistant T-DNA vector to misexpress

microRNA resistant REV-2×VENUS fusion, where *6Op::REVr-2×VENUS* was

constructed by cloning a 1148 bp *Bam*HI-*Xcm*I microRNA resistant REV cDNA (a gift

from J. Bowman) harbouring two previously characterized silent mutations that disrupt

the binding of MIRNA 165/166 to the coding sequence of *REV* as previously described

(Emery et al., 2003) downstream of the *6Op* and in frame with the wild type *REV-*

2×VENUS coding sequences.

miR166Ap::GFPER T-DNA construct was kindly provided by K. Nakajima (Miyashima

et al., 2011). The MIR165/166 biosensor was created based on the design presented by

Smith Z. R. et al. (Smith and Long, 2010) in the AlcR/AlcA expression system (Roslan et

al., 2001) for ethanol-inducible expression. The sequences conferring MIR165/166

sensitivity from the *REV* coding sequence (*REV*) and the sequences conferring

MIR165/166 insensitivity (REVr) were fused to *mCherry* (Shaner et al., 2004) with

endoplasmic reticulum localization sequences *mCherryER*, which was synthesized de

novo (Genscript). The MIR165/166-*mCherryER* biosensors (both biosensor and control)

were cloned as *Hind*III-*Bam*HI fragments downstream of the *AlcA* regulatory sequences

in the *UBQ10p:AlcR_BJ36* plasmid vector. *UBQ10p:AlcR* was constructed by cloning the

UBQ10 promoter 2 kb fragment upstream of *AlcR* and the *OCS* terminator. Both

UBQ10p::AlcR and *AlcA::REV-mCherryER* or *AlcA::REVr-mCherryER* components

were combined in the T-DNA vector *pMOA34*.

The *WOX1p::2×GFP-WOX1* construct in *pMLBART* T-DNA vector was generated as

follows: 2.2 kb of *WOX1 (At3g18010)* upstream promoter sequence was amplified with

primers WOX1p F and WOX1p R and cloned using restriction enzymes KpnI and

BamHI. 3.6 kb of *WOX1* coding sequence plus 1.65 kb 3'-regulatory sequences was

amplified from wild-type Col-O genomic DNA with the primers WOX1g F and WOX1g

R and cloned using restriction enzymes *Bgl*II and *Spe*I. 2 copies of GFP were inserted in

frame at the start of the *WOX1* coding sequence at the *Bam*HI and *Bgl*II sites. A double

marker was generated by transforming the *WOX1p::2×GFP-WOX1* into *a PIN1p::PIN1-*

*C*FP transgenic line.

The *PRSp::PRS-2×GFP* construct in *pMOA34* T-DNA vector was made by amplification

of 3.9 kb *PRS (At2g28610)* genomic sequence (similar to (Shimizu et al., 2009)) with

primers PRSg F and PRSg R to create a C-terminal fusion to *2×GFP* followed by *OCS*

3'regulatory sequences. Marker combinations were generated by transforming the

PRSp::PRS-2×GFP into either a *PIN1p::PIN1-CFP* transgenic line or into *REVp::REV-*

2×YPET PIN1p::PIN1-CFP line. *(PRSp::PRS-2×GFP)* and *(KAN1p::KAN1-2×CFP)*

were combined in T-DNA vector BGW (Karimi et al., 2002) by Gateway technology

(Invitrogen) for generation of a double marker transgenic lines.

A short tandem target mimic (STTM) construct to target MIR165/166 (Yan et al., 2012)

was generated in the *pOp6/LhGR* two-component system for dexamethasone-inducible

expression with a *UBQ10p::GRLh* driver. *STTM MIR165/166-88* sequence (Yan et al.,

2012) was synthesized de novo (Genscript) and cloned downstream of *6×Op* to create

6×Op::STTM 165/166. Both components were combined in a sulfadiazine T-DNA

pSULT (*UBQ10>>STTM 165/166*).

ATML1>>PHVr is a sulfadiazine-resistant T-DNA vector containing a mutated version

of *PHV* cDNA (a gift from J. Bowman) with a Gly to Asp amino acid change that

disrupts the miRNA165/166 binding site in the *PHV* gene (McConnell et al., 2001).

6×Op::PHVr was constructed by cloning a 2.6 kb *Xho*I-*Bam*HI *PHVr* cDNA downstream

of *6×Op* and upstream of *pea3A* terminator sequences. *ATML1>>PHVr* was transformed

into a *REVp::REV-2×YPet; PIN1p::PIN1-CFP* transgenic line.

DR5-3×VENUS v2 reporter gene (Liao et al., 2015) was a generous gift from D. Weijers.

The line *R2D2 PIN1p::PIN1-GFP* was described previously (Bhatia et al., 2016).

Dexamethasone induction

For inducible gene perturbations in the vegetative SAM, seeds were germinated directly

on GM medium containing 10 μM Dexamethasone (Sigma, stock solution was prepared

in Ethanol).. Seedlings were then dissected for imaging at 4 DAS, 5 DAS or 7 DAS

depending on the experiment. For DEX induction in the IM, 10 μM DEX solution

containing 0.015% Silwet L-77 was applied to the IM every second day three times.

Inflorescences were then dissected and imaged. The number of T2 inducible transgenic
lines that exhibit the presented phenotypes and the frequencies of phenotypes amongst
imaged plants is shown in Table S2 and associated caption.

Confocal Microscopy

Plants were dissected for imaging as previously described (Bhatia et al., 2016; Heisler
and Ohno, 2014) and imaged with a Leica SP5 Upright confocal microscope using an
Argon laser. The objective used was a water-immersion HCX IRAPO L25x/0.95 W
(Leica). Excitation for CFP is 458 nm, GFP is 488 nm, YFP (Ypet and VENUS) is 514
nm and tdTomato is 561 nm. Emission signal were collected at 460-480 nm for CFP,
490-512nm for GFP, 516-550 nm for YFP (YPet and VENUS), and 570-620 nm for
tdTomato. The resulting z-stacks were processed using the Dye Separation (Channel
mode or automatic mode) function available in the LAS AF program in order to separate
the GFP channel from the YFP (Ypet or VENUS) channel. Three software packages:
LAS AF from Leica, Imaris 8.0.2 by Bitplane and FIJI (https://fiji.sc) were used for data
analysis. Ratios for R2D2 were calculated as described previously (Bhatia et al., 2016).

Measurement of distance between organs

For distance measurements between oppositely positioned leaves on plants transgenic for
inducible KAN1 (*CLV3>>KAN1-2×GFP* – see above) the measurement tool from Imaris
8.0.2 (Bitplane) was used. For comparisons to control, untreated seedlings grown on GM
were compared to seedlings grown on DEX for 5 days. t-Test was performed using Excel.

Chemical treatments for auxin depletion in the inflorescence meristems

500mM stock solutions of auxinole, yucasin and L-Kynurenine were prepared separately
in DMSO. The stocks were diluted in 1mL 0.1M phosphate buffer in sterile water to
make a working solution containing all the three drugs (0.2μL of each stock) to a final
concentration of 100mM each. The final concentration of DMSO in the working solution
containing all the three drugs was 0.06%.

Treatments were carried out on the inflorescence meristems of whole plants transplanted
from soil to boxes containing GM medium supplemented with vitamins. The older

flowers were removed as described (Heisler and Ohno, 2014). The plants were chosen

such that the stem of the meristem was a few millimeters above the rosette to prevent the

drug solution from dispersing into the surrounding medium. After imaging, the meristems

were carefully dried using a thin strip of sterile filter paper to remove excess water.

Approximately 50μL of the drug solution was added directly to the meristem, drop-wise.

The meristems were treated only once in a time course of 12-18 hours.

NPA treatment on seedlings carrying R2D2 and DR5 markers

455 Seedling aged 3.5-4DAS were dissected to expose the meristem and the first leaf pair as

456 described (Bhatia et al., 2016). After imaging, seedlings were transferred to new GM

457 medium containing plates and blotted dry with thin strips of sterile paper. 5-10μL of 100

458 μM NPA in sterile water (100 mM stock in DMSO) was added directly to the dissected

459 seedlings every twenty-four hours for three days in total.

Auxin treatment on seedlings carrying PRS and WOX1 markers

461 Seedlings aged 4DAS (days after stratification) were dissected to expose the meristem

462 and the first leaf pair as described (Bhatia et al., 2016). 5mM NAA solution was prepared

463 in liquid GM medium. Seedlings were then immersed in 100μL of NAA containing

464 medium in individual wells in 96 well plate and grown under continuous light without

465 shaking for 12 hours.

Pulsed Laser ablations.

Laser ablations on the inflorescence meristems were carried out using the Mai Tai multi-

photon laser from Spectra Physics, which was controlled with LEICA SP5 confocal

software. Z-stacks were acquired prior to ablation. Single cells were targeted one after

the other using bleach point mode. Ablations were carried out at 800nm with an output

power of ~3W. Each pulse was shot for 1-15 milliseconds. Usually ablations were

accomplished within 1-3 bursts of the laser. Ablated cells could be visually identified as

their nuclei exploded resulting in unusual auto fluorescence. Z stacks were acquired

immediately after the ablations.

Auxin and auxin plus NPA combined treatments on ablated inflorescence meristems

After ablations, the meristems were carefully blotted dry using thin strips of sterile filter paper. 20μL 5mM NAA in sterile water (0.5M stock in 1M KOH) or 20μL of a solution containing 5mM NAA and 100μM NPA in sterile water (100mM NPA stock in DMSO) or mock solution were added directly to the meristems every 24 hours for 48 hours in total.

Real time PCR

4 day-old wild-type Ler seedlings were immersed into 5mM NAA solution in liquid GM medium, and grown under continuous light without shaking for 15 hours. Cotyledons, hypocotyl, and roots were removed under dissecting scope, and only shoot meristem and the first pair of leaves were collected and immediately frozen in liquid nitrogen. Each biological replicate, represents tissue from 10-15 individual seedlings. Five biological replicates were collected for both mock and NAA treatment. RNeasy Mini kit (Qiagen) was used according to manufacturer's instruction for RNA extraction. 1 microgram of RNA was used for cDNA preparation using Super script III reverse transcriptase for Q-PCR analysis. Q-PCRs were performed in a StepOne Plus Real Time PCR system thermo cycler (The applied bio systems) using 20μl of PCR reaction containing 10 μL of SYBR Green mix (Roche), 1μl of primer mix (10μm), 2μl of 1:10 diluted cDNA and 7 μl of water. Transcript levels were normalized to ACTIN2 transcript levels. Data was analyzed using the 2−**ΔΔ**CT method. A freely available online tool was used for analysis using an unpaired t-test of the RT-PCR results: http://graphpad.com/data-analysis-resource-center/. For p-value calculation, data entry format with mean, SD and N was used. Measurements and calculations for all replicates are provided in Table S4.

Model for auxin and PIN1 dynamics

We developed a computational model to understand how interplay between the HD-ZIPIII and KAN pattern and PIN1 dynamics can influence auxin transport and primordia initiation. The model introduces a dependence on KANADI and REVOLUTA into previous models describing PIN1 and auxin dynamics (Bhatia et al., 2016; Heisler et al., 2010; Heisler and Jonsson, 2006; Jonsson et al., 2006; Sahlin et al., 2009). In the model,

auxin resides in cell compartments and is able to move between cells either via active
transport mediated by PIN1 proteins or passively via a diffusion-like process. PIN1
proteins cycle between cytosol and membrane compartments and a quasi-equilibrium
model is used for determining its membrane localization at any time point. Auxin
generates a signal able to polarise PIN1 in neighboring cells, i.e. a high auxin
concentration increases the amount of PIN1 proteins in the neighboring cell membrane
facing that cell. The molecule X in our model acts as a mediator of the signalling between
auxin and PIN1, and the signal has previously been interpreted as a molecular (Jonsson et
al., 2006), or mechanical stress signal (Heisler et al., 2010). In the model, the signal X is
activated by auxin, and repressed by KAN and REV. The equations governing the
dynamics of the molecules are

515
$$\frac{dA_i}{dt} = c_A - d_A A_i + \frac{1}{V_i}\left[D\sum_{j\in\{N_i\}} a_{ij}(A_j - A_i) + T\sum_{j\in\{N_i\}} a_{ij}(P_{ji}A_j - P_{ij}A_i)\right],$$

516
$$\frac{dP_i^{[tot]}}{dt} = c_P - d_P P_i^{[tot]},$$

517
$$\frac{dX_i}{dt} = V_X \frac{A_i^{n_{XA}}}{(K_{XA}^{n_{XA}} + A_i^{n_{XA}})} \frac{K_{XR}^{n_{XR}}}{(K_{XR}^{n_{XR}} + R_i^{n_{XR}})} \frac{K_{XK}^{n_{XK}}}{(K_{XK}^{n_{XK}} + K_i^{n_{XK}})} - d_X X_i,$$

where A_i is the auxin concentration and X_i is the level of the signalling molecule in cell i.
$\{N_i\}$ is the set of cells neighboring cell i, V_i is the cell volume of cell i, and $a_{ij} = a_{ji}$ is the
cell wall area for the wall section between cells i and j. $P_i^{[tot]}$ is the total
PIN1concentration in the cytosol and membrane compartments of cell i. Membrane-
bound PIN1 appears in the equation as P_{ij}, which is the PIN1 concentration in the
membrane compartment of cell i that faces cell j. A simple linear feedback between the
signal X_j and P_{ij} is used and a quasi-stable assumption, introduced in Jönsson *et. al.*
(2006), leads to
527

528
$$P_{ij} = \frac{P_i^{[tot]}[(1 - k_p) + k_p X_j]}{f_p + \sum_{k\in\{N_i\}}[(1 - k_p) + k_p X_k]},$$

where $f_p = k_n/k_x$ is the ratio of endocytosis and exocytosis rates and k_p sets the relation

between symmetric and polarized exocytosis ($dP_{ij}/dt = k_x[(1-k_p) + k_pX_j]P_i - k_n P_{ij}$, where

P_i is the PIN1 in the cytosol compartment). The feedback between auxin and PIN1 is

identical to previous models if the KAN and REV factors in the dX_i/dt equation are 1 (e.g.

by setting KAN and REV to zero in all cells), while the polarising signal becomes

reduced in regions where KAN and/or REV are expressed. This effect tunes the

interaction between the dorsoventral patterning and PIN1/auxin dynamics (cf. Figure. 8B

and D).

The model was simulated using the in-house developed software organism/tissue

(http://dev.thep.lu.se/organism/, available upon request). Files defining the model with

parameter values (Suppl. Table S3) and initial configuration of (static) cell geometries

and KAN and REV expression domains are provided as Supplemental Information. The

simulations use a 5th order Runge-Kutta solver with adaptive step size (William H. Press,

2007), and initial auxin, PIN1 and X concentrations are set to zero in all compartments.

Generation of geometrical template

The model defined above was run on a template containing a predefined KAN/REV

pattern (provided as Supplemental material file, Figure. 8C). The geometry of the

template was generated by a combination of cell/wall growth and mechanical interactions

together with a shortest path division rule (Sahlin and Jonsson, 2010). A KAN/REV

pattern was generated by the equations

$$\frac{dK_i}{dt} = V_K \frac{r_i^{n_K}}{K_K^{n_K} + r_i^{n_K}} - d_K K_i,$$

$$\frac{dR_i}{dt} = V_R \frac{r_i^{n_R}}{K_R^{n_R} + r_i^{n_R}} - d_R R_i.$$

This system was run to equilibrium on the above-mentioned template. In the above

equations r_i is the distance of cell i from the center of the template. The parameters were

set to $V_K=V_R=d_K=d_R=1$, $K_K=30$, $K_R=25$, $n_K=n_R=20$. These parameters are set such that

two distinct domains are created, with a small overlap of low KAN and REV

concentrations in the boundary between these regions. To make the transition of KAN

concentrations between domains sharper, KAN concentrations were set to 0 (1) if $K_i \leq 0.5$

(> 0.5).

Acknowledgements

We thank M. E. Byrne for helpful feedback and ideas on the manuscript. We thank J.

Bowman for helpful discussions as well as for plasmids containing REVr and PHVr

cDNAs. We thank D. Weijers and C. Y. Liao for a plasmid containing

pDR5v2::ntdTomato; K. Nakajima for the T-DNA construct of miR166Ap::GFPER, Dr

Atsushi Miyawaki from RIKEN Brain Science Institute for the VENUS fluorescent

protein which was obtained through an MTA. The E.M.M. laboratory is supported by

funds from the Howard Hughes Medical Institute and the Gordon and Betty Moore

Foundation (through grant GBMF3406). The research leading to these results received

funding from the Australian Research Council (M.G.H.) and European Research Council

under the European Union's Seventh Framework Programme (FP/2007-2013) / ERC

Grant Agreement n. 261081 (M.G.H.), as well as the People Programme (Marie Curie

Actions) under REA grant agreement n. 255089 (P.S.). The work was also supported by:

the European Molecular Biology Laboratory (X.Y., M.P.C., C.O., P.S, N.B., H.R. and

M.G.H.); the EMBL International PhD Programme (X.Y., N.B. and M.P.C.); Gatsby

Charitable Foundation (GAT3395/PR4) (H.J) and Swedish Research Council (VR2013-

4632) (H.J). The authors declare no competing financial interests. Supplement contains

additional data and movie. We declare there are no competing interests.

References

Barrell, P. J. and Conner, A. J. (2006). Minimal T-DNA vectors suitable for
agricultural deployment of transgenic plants. *Biotechniques* **41**, 708-710.

Bhatia, N., Bozorg, B., Larsson, A., Ohno, C., Jonsson, H. and Heisler, M. G. (2016).
Auxin Acts through MONOPTEROS to Regulate Plant Cell Polarity and Pattern
Phyllotaxis. *Curr Biol* **26**, 3202-3208.
Bilsborough, G. D., Runions, A., Barkoulas, M., Jenkins, H. W., Hasson, A.,
Galinha, C., Laufs, P., Hay, A., Prusinkiewicz, P. and Tsiantis, M. (2011).
Model for the regulation of Arabidopsis thaliana leaf margin development. *Proc*
Natl Acad Sci U S A **108**, 3424-3429.
Craft, J., Samalova, M., Baroux, C., Townley, H., Martinez, A., Jepson, I., Tsiantis,
M. and Moore, I. (2005). New pOp/LhG4 vectors for stringent glucocorticoid-
dependent transgene expression in Arabidopsis. *Plant J.* **41**, 899-918.
de Reuille, P. B., Bohn-Courseau, I., Ljung, K., Morin, H., Carraro, N., Godin, C.
and Traas, J. (2006). Computer simulations reveal properties of the cell-cell
signaling network at the shoot apex in Arabidopsis. *Proc Natl Acad Sci U S A*
103, 1627-1632.
Emery, J. F., Floyd, S. K., Alvarez, J., Eshed, Y., Hawker, N. P., Izhaki, A., Baum,
S. F. and Bowman, J. L. (2003). Radial patterning of Arabidopsis shoots by class
III HD-ZIP and KANADI genes. *Curr Biol* **13**, 1768-1774.
Gleave, A. P. (1992). A Versatile Binary Vector System with a T-DNA Organizational-
Structure Conducive to Efficient Integration of Cloned DNA into the Plant
Genome. *Plant Molecular Biology* **20**, 1203-1207.
Gordon, S. P., Heisler, M. G., Reddy, G. V., Ohno, C., Das, P. and Meyerowitz, E.
M. (2007). Pattern formation during de novo assembly of the Arabidopsis shoot
meristem. *Development* **134**, 3539-3548.
Hagemann, W. and Gleissberg, S. (1996). Organogenetic capacity of leaves: The
significance of marginal blastozones in angiosperms. *Plant Syst. Evol.* **199**, 121-
152.
Hayashi, K., Neve, J., Hirose, M., Kuboki, A., Shimada, Y., Kepinski, S. and Nozaki,
H. (2012). Rational design of an auxin antagonist of the SCF(TIR1) auxin
receptor complex. *ACS chemical biology* **7**, 590-598.
He, W., Brumos, J., Li, H., Ji, Y., Ke, M., Gong, X., Zeng, Q., Li, W., Zhang, X., An,
F., et al. (2011). A small-molecule screen identifies L-kynurenine as a
competitive inhibitor of TAA1/TAR activity in ethylene-directed auxin
biosynthesis and root growth in Arabidopsis. *Plant Cell* **23**, 3944-3960.
Heisler, M. G., Hamant, O., Krupinski, P., Uyttewaal, M., Ohno, C., Jonsson, H.,
Traas, J. and Meyerowitz, E. M. (2010). Alignment between PIN1 polarity and
microtubule orientation in the shoot apical meristem reveals a tight coupling
between morphogenesis and auxin transport. *PLoS Biol* **8**, e1000516.
Heisler, M. G. and Jonsson, H. (2006). Modeling auxin transport and plant
development. *Journal of Plant Growth Regulation* **25**, 302-312.
Heisler, M. G. and Ohno, C. (2014). Live-imaging of the Arabidopsis inflorescence
meristem. In *Flower Development*, pp. 431-440: Springer.
Heisler, M. G., Ohno, C., Das, P., Sieber, P., Reddy, G. V., Long, J. A. and
Meyerowitz, E. M. (2005). Patterns of auxin transport and gene expression
during primordium development revealed by live imaging of the Arabidopsis
inflorescence meristem. *Current Biology* **15**, 1899-1911.

Huang, T., Harrar, Y., Lin, C., Reinhart, B., Newell, N. R., Talavera-Rauh, F.,
Hokin, S. A., Barton, M. K. and Kerstetter, R. A. (2014). Arabidopsis
KANADI1 acts as a transcriptional repressor by interacting with a specific cis-
element and regulates auxin biosynthesis, transport, and signaling in opposition to
HD-ZIPIII factors. *Plant Cell* **26**, 246-262.
Husbands, A. Y., Chitwood, D. H., Plavskin, Y. and Timmermans, M. C. (2009).
Signals and prepatterns: new insights into organ polarity in plants. *Genes Dev* **23**,
1986-1997.
Izhaki, A. and Bowman, J. L. (2007). KANADI and class III HD-Zip gene families
regulate embryo patterning and modulate auxin flow during embryogenesis in
Arabidopsis. *Plant Cell* **19**, 495-508.
Jonsson, H., Heisler, M. G., Shapiro, B. E., Meyerowitz, E. M. and Mjolsness, E.
(2006). An auxin-driven polarized transport model for phyllotaxis. *Proceedings of*
the National Academy of Sciences of the United States of America **103**, 1633-
1638.
Karimi, M., Inze, D. and Depicker, A. (2002). GATEWAY vectors for Agrobacterium-
mediated plant transformation. *Trends Plant Sci* **7**, 193-195.
Kerstetter, R. A., Bollman, K., Taylor, R. A., Bomblies, K. and Poethig, R. S. (2001).
KANADI regulates organ polarity in Arabidopsis. *Nature* **411**, 706-709.
Kidner, C. A. and Martienssen, R. A. (2004). Spatially restricted microRNA directs
leaf polarity through ARGONAUTE1. *Nature* **428**, 81-84.
Koenig, D., Bayer, E., Kang, J., Kuhlemeier, C. and Sinha, N. (2009). Auxin patterns
Solanum lycopersicum leaf morphogenesis. *Development* **136**, 2997-3006.
Kuhlemeier, C. and Timmermans, M. C. (2016). The Sussex signal: insights into leaf
dorsiventrality. *Development* **143**, 3230-3237.
Landrein, B., Kiss, A., Sassi, M., Chauvet, A., Das, P., Cortizo, M., Laufs, P.,
Takeda, S., Aida, M., Traas, J., et al. (2015). Mechanical stress contributes to
the expression of the STM homeobox gene in Arabidopsis shoot meristems. *eLife*
4, e07811.
Liao, C. Y., Smet, W., Brunoud, G., Yoshida, S., Vernoux, T. and Weijers, D. (2015).
Reporters for sensitive and quantitative measurement of auxin response. *Nat*
Methods **12**, 207-210, 202 p following 210.
McConnell, J. R., Emery, J., Eshed, Y., Bao, N., Bowman, J. and Barton, M. K.
(2001). Role of PHABULOSA and PHAVOLUTA in determining radial
patterning in shoots. *Nature* **411**, 709-713.
Merelo, P., Ram, H., Pia Caggiano, M., Ohno, C., Ott, F., Straub, D., Graeff, M.,
Cho, S. K., Yang, S. W., Wenkel, S., et al. (2016). Regulation of MIR165/166
by class II and class III homeodomain leucine zipper proteins establishes leaf
polarity. *Proc Natl Acad Sci U S A*.
Merelo, P., Xie, Y., Brand, L., Ott, F., Weigel, D., Bowman, J. L., Heisler, M. G. and
Wenkel, S. (2013). Genome-wide identification of KANADI1 target genes. *PLoS*
One **8**, e77341.
Miyashima, S., Honda, M., Hashimoto, K., Tatematsu, K., Hashimoto, T., Sato-
Nara, K., Okada, K. and Nakajima, K. (2013). A comprehensive expression
analysis of the Arabidopsis MICRORNA165/6 gene family during embryogenesis

reveals a conserved role in meristem specification and a non-cell-autonomous
function. *Plant Cell Physiol* **54**, 375-384.

Miyashima, S., Koi, S., Hashimoto, T. and Nakajima, K. (2011). Non-cell-
autonomous microRNA165 acts in a dose-dependent manner to regulate multiple
differentiation status in the Arabidopsis root. *Development* **138**, 2303-2313.

Nagai, T., Ibata, K., Park, E. S., Kubota, M., Mikoshiba, K. and Miyawaki, A.
(2002). A variant of yellow fluorescent protein with fast and efficient maturation
for cell-biological applications. *Nature Biotechnology* **20**, 87-90.

Nakata, M., Matsumoto, N., Tsugeki, R., Rikirsch, E., Laux, T. and Okada, K.
(2012). Roles of the middle domain-specific WUSCHEL-RELATED
HOMEOBOX genes in early development of leaves in Arabidopsis. *Plant Cell* **24**,
519-535.

Nguyen, A. W. and Daugherty, P. S. (2005). Evolutionary optimization of fluorescent
proteins for intracellular FRET. *Nat Biotechnol* **23**, 355-360.

Nishimura, T., Hayashi, K., Suzuki, H., Gyohda, A., Takaoka, C., Sakaguchi, Y.,
Matsumoto, S., Kasahara, H., Sakai, T., Kato, J., et al. (2014). Yucasin is a
potent inhibitor of YUCCA, a key enzyme in auxin biosynthesis. *Plant J* **77**, 352-
366.

Nogueira, F. T. S., Madi, S., Chitwood, D. H., Juarez, M. T. and Timmermans, M.
C. P. (2007). Two small regulatory RNAs establish opposing fates of a
developmental axis. *Genes & Development* **21**, 750-755.

Otsuga, D., DeGuzman, B., Prigge, M. J., Drews, G. N. and Clark, S. E. (2001).
REVOLUTA regulates meristem initiation at lateral positions. *Plant J* **25**, 223-
236.

Qi, J., Wang, Y., Yu, T., Cunha, A., Wu, B., Vernoux, T., Meyerowitz, E. and Jiao,
Y. (2014). Auxin depletion from leaf primordia contributes to organ patterning.
Proc Natl Acad Sci U S A **111**, 18769-18774.

Reinhardt, D., Frenz, M., Mandel, T. and Kuhlemeier, C. (2005). Microsurgical and
laser ablation analysis of leaf positioning and dorsoventral patterning in tomato.
Development **132**, 15-26.

Reinhardt, D., Mandel, T. and Kuhlemeier, C. (2000). Auxin regulates the initiation
and radial position of plant lateral organs. *Plant Cell* **12**, 507-518.

Reinhardt, D., Pesce, E. R., Stieger, P., Mandel, T., Baltensperger, K., Bennett, M.,
Traas, J., Friml, J. and Kuhlemeier, C. (2003). Regulation of phyllotaxis by
polar auxin transport. *Nature* **426**, 255-260.

Roslan, H. A., Salter, M. G., Wood, C. D., White, M. R., Croft, K. P., Robson, F.,
Coupland, G., Doonan, J., Laufs, P., Tomsett, A. B., et al. (2001).
Characterization of the ethanol-inducible alc gene-expression system in
Arabidopsis thaliana. *Plant J* **28**, 225-235.

Rosso, M. G., Li, Y., Strizhov, N., Reiss, B., Dekker, K. and Weisshaar, B. (2003). An
Arabidopsis thaliana T-DNA mutagenized population (GABI-Kat) for flanking
sequence tag-based reverse genetics. *Plant Mol Biol* **53**, 247-259.

Sahlin, P. and Jonsson, H. (2010). A modeling study on how cell division affects
properties of epithelial tissues under isotropic growth. *PLoS One* **5**, e11750.

Sahlin, P., Soderberg, B. and Jonsson, H. (2009). Regulated transport as a mechanism
for pattern generation: capabilities for phyllotaxis and beyond. *J Theor Biol* **258**,
60-70.
Samalova, M., Brzobohaty, B. and Moore, I. (2005). pOp6/LhGR: a stringently
regulated and highly responsive dexamethasone-inducible gene expression system
for tobacco. *Plant J.* **41**, 919-935.
Sessions, A., Weigel, D. and Yanofsky, M. F. (1999). The Arabidopsis thaliana
MERISTEM LAYER 1 promoter specifies epidermal expression in meristems and
young primordia. *Plant J* **20**, 259-263.
Shaner, N. C., Campbell, R. E., Steinbach, P. A., Giepmans, B. N., Palmer, A. E. and
Tsien, R. Y. (2004). Improved monomeric red, orange and yellow fluorescent
proteins derived from Discosoma sp. red fluorescent protein. *Nat Biotechnol* **22**,
1567-1572.
Shimizu, R., Ji, J., Kelsey, E., Ohtsu, K., Schnable, P. S. and Scanlon, M. J. (2009).
Tissue specificity and evolution of meristematic WOX3 function. *Plant Physiol.*
149, 841-850.
Smith, R. S., Guyomarc'h, S., Mandel, T., Reinhardt, D., Kuhlemeier, C. and
Prusinkiewicz, P. (2006). A plausible model of phyllotaxis. *Proc Natl Acad Sci*
U S A **103**, 1301-1306.
Smith, Z. R. and Long, J. A. (2010). Control of Arabidopsis apical-basal embryo
polarity by antagonistic transcription factors. *Nature* **464**, 423-426.
Snow, M. and Snow, R. (1959). The dorsiventrality of leaf primordia. *New Phytol.* **58**,
188-207.
Snow, M. S., R. (1931). Experiments on phyllotaxis. I. The effect of isolating a
primordium. *Philos Trans R Soc Lond B Biol Sci* **221**, 1-43.
Sussex, I. M. (1951). Experiments on the cause of dorsiventrality in leaves. *Nature* **167**,
651-652.
---- (1955). Morphogenesis in Solanum tuberosum I: experimental investigation of leaf
dorsiventrality and orientation in the juvenile shoot. *Phytomorphology* **5**, 286-300.
Tameshige, T., Fujita, H., Watanabe, K., Toyokura, K., Kondo, M., Tatematsu, K.,
Matsumoto, N., Tsugeki, R., Kawaguchi, M., Nishimura, M., et al. (2013).
Pattern dynamics in adaxial-abaxial specific gene expression are modulated by a
plastid retrograde signal during Arabidopsis thaliana leaf development. *PLoS*
Genet **9**, e1003655.
William H. Press, S. A. T., William T. Vetterling, Brian P. Flannery (2007).
Numerical Recipes: The art of scientific computing (3rd edn). New York, NY,
USA: Cambridge University Press
Yan, J., Gu, Y., Jia, X., Kang, W., Pan, S., Tang, X., Chen, X. and Tang, G. (2012).
Effective small RNA destruction by the expression of a short tandem target mimic
in Arabidopsis. *Plant Cell* **24**, 415-427.
Yao, X., Wang, H., Li, H., Yuan, Z., Li, F., Yang, L. and Huang, H. (2009). Two
types of cis-acting elements control the abaxial epidermis-specific transcription of
the MIR165a and MIR166a genes. *FEBS Lett* **583**, 3711-3717.
Zhang, Z., Tucker, E., Hermann, M. and Laux, T. (2017). A Molecular Framework
for the Embryonic Initiation of Shoot Meristem Stem Cells. *Dev Cell* **40**, 264-277
e264.

Zuo, J., Niu, Q. W., Moller, S. G. and Chua, N. H. (2001). Chemical-regulated, site-
specific DNA excision in transgenic plants. *Nat Biotechnol* **19**, 157-161.
772

773
774
775

Figure 1.

Organ initiation is centered on a boundary between the expression domains of genes involved in leaf dorsoventrality.

(**A to D**) Confocal projections showing REV-2×YPet (red), PIN1-CFP (blue) and KAN1-2×GFP (green) expression in a vegetative shoot apical meristem at 3 days (A), 4 days (C) and 5 days (D) after stratification (DAS), respectively. (B) Longitudinal reconstructed section of seedling shown in (A). Dashed box in (D) corresponds to close up in (F) to (H). (**E**) Expression pattern of REV-2×YPet, KAN1-2×GFP and PIN1-CFP in an inflorescence meristem. Dashed box corresponds to close ups in (I) to (K). White arrow head marks region where KAN1-2×GFP expression is being reestablished after organ emergence. (**F to K**) localized PIN1-CFP expression marking organ inception at the REV-2×YPet /KAN1-2×GFP boundary in both the vegetative meristem (F) to (H) and inflorescence meristem (I) to (K). White arrow heads mark cells in between the REV-2×YPet and KAN1-2×GFP expression domains where PIN1-CFP expression is highest. Scale bars represent 20 μm.

796
Figure 2.
The expression of KAN1-2×GFP is relatively stable with respect to the underlying

cells within initiating organs.

(A-B) Confocal projections showing an inflorescence meristem viewed from above

expressing PIN1-CFP (blue) and KAN1-2×GFP (green) at two time points (0h and 12h).

Two incipient primordia are marked I1 and I2. **(C-D)** Close up views corresponding to

primordium I1 from (A) and (B) with white arrows marking three cells at the edge of

KAN1-2×GFP expression that retain this expression over the time interval. Yellow

arrows mark two cells in which KAN1-2×GFP is absent at 12h. **(E)** Close up views

corresponding to primordium I2 from (A) and (B) with white arrows marking three cells

that retain KAN1-2×GFP expression at the adaxial (towards the meristem) edge over the

time interval. Bars represent 20 μm in (A) and (B); 10 μm in (C) to (F).

Figure 3.

Expression and activity of MIR165/166 is localized to the periphery of the shoot meristem.

(A) Expression of *MIR166Ap::GFPER* (green), PIN1-CFP (blue) and REV-2×YPet (red) in the vegetative meristem (VM) at 3.5 DAS. **(B)** Longitudinal section of meristem shown in (A). **(C)** Expression of *MIR166Ap::GFPER* (green), PIN1-CFP (blue) and REV-2×YPet (red) in the inflorescence meristem (IM). White arrow head marks the reestablishment of *MIR166Ap::GFPER* expression around the meristem after organ emergence. **(D to F)** Expression of REV-2×YPet (red) alone (D), a MIR165/166 biosensor driven by the *UBQ10* promoter (green) alone (E) and both combined in the same IM (F). (**G to I**) Corresponding control for (D to F) where the MIR165/166 biosensor has been rendered insensitive to MIRNA activity. Bars represent 20 μm.

Figure 4.

Expression patterns of 2×GFP-WOX1 and PRS-2×GFP

(**A**) Confocal projections showing PIN1-CFP (blue) and 2×GFP-WOX1 (green)

expression patterns in the vegetative meristem and leaves of seedlings at 3 DAS. (**B**)

Longitudinal section of meristem shown in (A). (**C**) An inflorescence meristem image

showing 2×GFP-WOX1 is not expressed in the IM. (**D**) Confocal projection showing

PIN1-CFP (blue), PRS-2×GFP (green) and REV-2×YPet (red) expression in the

vegetative meristem and leaves at 3.5 DAS, where PRS-2×GFP is expressed surrounding

the VM and along the leaf margins. (**E and F**) Cross sections of leaf on the right side in

(D) showing the expression of PRS-2×GFP in the middle domain of the leaf. (**G and H**)

Confocal projection and cross section showing PRS-2×GFP (green) and KAN1-2×CFP

(red) expression patterns in the vegetative meristem and leaves of seedlings at 3 DAS. (**I**

and J) PRS-2×GFP (green) is expressed in the young flower primodia in the IM,

indicated with arrowhead in (J). Bar = 20 μm.

Figure 5.

Organ initiation depends on the restriction of Class III HD-ZIP and KANADI expression in the shoot.

(**A**) Confocal projection showing wild type control seedling at 7DAS viewed from above for comparison to (C) and (F). (**B**) Macroscopic view of control seedling at 7DAS for comparison to (D), (G) and (L). (**C**) Arrest of organogenesis after ectopic expression of REVr-2×VENUS from the *ATML1* promoter in the vegetative meristem (7 DAS) after germination on DEX, KAN1-2×GFP (green) expression is down regulated and could only be detected in a few cells in the sub-epidermis. Although PIN1-CFP (blue) expression is patchy, no leaves developed. (**D**) Macroscopic view of plant in (C). (**E**) Arrest of organogenesis after ectopic expression of REVr-2×VENUS (red) from the *ATML1* in the IM after 3 DEX treatments over 6 days. Note the absence of KAN1-2×GFP signal. (**F**) Seedlings at 7DAS showing similar phenotype to (C) after induction of a short tandem target mimic (STTM) designed to down regulate MIR165/166 activity. (**G**) Macroscopic

view of plant in (F). **(H to J)** Ectopic expression of REV-2YPet (red) and arrest of organogenesis (PIN1-CFP in blue) in 4DAS seedling after induction of MIR165/166 resistant PHAVOLUTA. (H) Longitudinal view of uninduced control. Top view (I), and macroscopic view (J) of induced seedling showing arrest of organ development. **(K and L)** Confocal projection (K) and macroscopic view (L) of seedling at 7DAS after induction of KAN1-GFP (green) in the epidermis. No leaves have developed (autofluorescence shown in red). (**M**) Arrest of organogenesis after induction of KAN1-GFP (green) driven by the *ATML1* promoter in the IM after 3 DEX treatments over 6 days; autofluorescence (red). Bars represent 20 μm in A, C, E, F, H to I, K and M; 1mm in B, D, G, J and L.

Figure 6.

KANADI1 expression boundaries in the shoot specify organ position and orientation.

(**A and B**) Confocal projections showing organ initiation marked by REV-2×YPet (red) and PIN1-CFP (blue) at border of KAN1-2×GFP expression (green) in wild type (A) and after induction of KAN1-2×GFP using the *CLV3* promoter (B). Distance separating opposite organs was greater for induced (B) compared to control (A) (114.3 ±3.3 μm, n=19 vs 54.2±1.0 μm n=10 (mean±SE, p<0.05, t-test)). (**C to H**) Confocal projections (C, E and G) and longitudinal reconstructions corresponding to dashed lines (D, F and H respectively) showing restricted REV-2×YPet expression (red) after ectopic KAN1-

2×GFP induction (green). Regions in which neither REV-2×YPet nor KAN1-2×GFP

signal was detected may potentially express endogenous KAN1, which was not

monitored. Four main configurations of REV expression and morphology were observed

(labeled 1 to 4). Class 1 organs (E and F) correspond to the wild type, Class 2 (C and D)

express REV-2×YPet centrally, Class 3 (E and F) express REV-2×YPet in a reversed

orientation and Class 4 (G and H) express REV-2×YPet centrally and laterally only.

Correspondence between REV-2×YPet expression boundaries and leaf margins indicated

by arrowheads in D, F and H; m indicates meristem. Gamma value changed to highlight

PIN1-CFP expression (blue) in (C) to (H). **(I to O)** Examples of mature leaves

corresponding to Classes 1 to 4, including the WT (I), cup-shaped (J), lotus-shaped (a

variation of cup-shaped) (K), needle-like (a further decrease in extent of dorsal tissue

compare to cup-shaped) (L), inverted (M and N) and four bladed (O). "D and V"

represent "dorsal" and "ventral" respectively in (O). **(P to S)** Diagrams summarizing

proposed configurations of REV and KAN (green) gene expression in leaf founder cells

(dashed circles) (upper diagram) leading to the observed phenotypic classes of leaf shape

(numbered 1 to 4) (lower diagram) after induction of KAN1-2×GFP using the *CLV3*

promoter. (P) represents the wild type Class 1 configuration, (Q) represents Class 2, (R)

represents Class 3 and (S) represents Class 4. Scale bars = 20µm in A to H; 1 mm in I, J

and M.

Figure 7.

Auxin promotes PRS and WOX expression.

(A-D) Response of pDR5V2-3×VENUS-N7 (green) auxin transcriptional reporter to

NPA in Arabidopsis seedlings 4DAS (Days after Stratification). (A and B) Confocal

projections 45 hours (A) and 72 hours (B) after treatment with mock solution. (C and D)

Confocal projections 45 hours (C) and 72 hours (D) after treatment with 100μM NPA

solution (n=5/5). Inset in (D) shows transverse optical section through the ring-shaped

organ showing most DR5 expression localized in the center of the organ. **(E-H)**

Expression and response of R2D2 (magenta) to auxin along with PIN1-GFP expression

(green) in Arabidopsis seedlings 4DAS. (E and F) Confocal projections 48 hours (E) and

72 hours (F) after treatment with mock solution. (G and H) Confocal projections 48 hours

(G) and 72 hours (H) after treatment with 100μM NPA solution (n=4/4). **(I-L)** Expression

and response of pPRS::2×GFP-PRS to auxin in Arabidopsis seedlings. Confocal

projections and transverse optical slices of seedlings 4DAS showing pPRS::2×GFP-PRS

expression (green) 12 hours after treatment with mock solution (I and J) and 5mM NAA

(K and L) (n=5/5). **(M-P)** Expression and response of 2XGFP-WOX to auxin in

Arabidopsis seedlings. (M and N) Confocal projections (M and O) and corresponding

optical slices (N and P) of seedlings 4DAS showing pWOX1::2×GFP-WOX1 expression

(green) 12 hours after treatment with mock solution (M and N) and 5mM NAA (O and

P). Note WOX expression increases but does not expands beyond its regular expression

domain upon auxin addition (n=5/5). **(Q)** Q-PCR analysis of PRS, WOX1 and positive

control IAA20 transcripts after 5mM NAA or mock treatment on 4 days old wild-type

(Ler) seedlings. $*=p<0.05$, $***=p<0.001$. Scale bars 20μm (A-L, K); 15μm (J and L);

30μm (M-P).

925
926
Figure 8.
Computational model illustrating how dorsoventral gene expression boundaries
may restrict phyllotactic patterning to the SAM peripheral zone.
(**A**) Illustration of model interactions. Auxin is transported passively and actively via
PIN1 between cells. PIN1 is polarized towards cells with high auxin, via a signaling
pathway represented by X (previously suggested to be realized by increased stresses in
the neighboring cells due to changes in mechanical wall properties (Heisler et al., 2010).
(**B**) As shown previously (Heisler et al., 2010; Jonsson et al., 2006; Smith et al., 2006),
peaks of auxin are formed spontaneously. (**C**) A pattern of KANADI (green) and
REVOLUTA (red) is added to the template with a boundary domain in between in which
REV expression is low or absent and KAN1 expression is absent. (**D**) If KANADI and
REVOLUTA decrease the signal X in cells where they are expressed (dashed interactions
in A), the formation of auxin peaks is restricted to the boundary.
940

Figure 9.

Effect of auxin addition and depletion on the expression pattern of DV genes.

(A and B) Confocal projections of the IMs showing expression pattern of REV-2×YPet (red), KAN1-2×GFP (green) and PIN1-CFP (blue) before (A) and 36 hours after the combined application of 5mM NAA and 100μM NPA (B). Note REV expression has broadened slightly after the application of NAA and NPA combination (compare regions marked by arrow heads in (A) vs (B) (n=2/3). **(C and D)** Confocal projections of the IMs showing expression pattern of REV-2×YPet (red), KAN1-2×GFP (green) and PIN1-CFP (blue) before (C) and 48 hours after the combined application of 100μM auxinole, 100μM KYN and 100μM Yucasin (auxin depleting drugs). Note KAN-2×GFP expression

has expanded centrally at the expense of REV-2×YPet expression (compare the cells

marked by arrowheads in (C) with (D), similar colored arrowheads mark the same cells

tracked over 48 hours) (n=6/6) **(E and F)** Confocal projections of the IMs indicating the

predicted auxin distribution (magenta) based on R2D2 expression along with PIN1-GFP

expression (green) before (E) and 17 hours after the combined application of 100μM

auxinole, 100μM kyn and 100μM yucasin (auxin depleting drugs) (F). Note lack of

detectable auxin based on R2D2 expression in (F) compared to (E) after drug application

(n=3/4). Scale bars 20μm (A-D), 30μm (E and F).

Figure 10.

Wounding induces auxin depletion dependent KANADI1 expression.

(A and B) Confocal projections of IMs showing predicted auxin distribution (magenta)

based on R2D2 expression 12hours (A) and 36 hours (B) after wounding. Note low

predicted auxin levels in the cells surrounding the ablated cells. **(C and D)** Confocal

projections of the IMs showing expression pattern of REV-2×YPet (red), KAN1-2×GFP

(green) and PIN1-CFP (blue) immediately after ablation (ablated cells are marked by

white arrowheads) (C) and 48 hours after. Note KAN1 expression (green) has completely

surrounded the wounded cells 48h after the ablation (D) compared to (C). **(E and F)**

Confocal projections of the IMs showing expression pattern of REV-2×YPet (red),

KAN1-2×GFP (green) and PIN1-CFP (blue) immediately after ablation (ablated cells are

marked by white arrowheads) (E) and 48 hours after ablation and combined NAA and

NPA application (F). Note absence of KAN1 expression (green) surrounding the wound

when wounding is accompanied by the exogenous addition of auxin and NPA (compare

to (D)). Scale bars 30μm (A-F).

Supplementary Figure S1.

FILp::dsREDN7 **expression is broad during leaf initiation but is later excluded from**

dorsal tissues. **(A-F)** Confocal projections and reconstructed sections of seedlings

expressing *FILp::dsREDN7* (green), REV-2×VENUS (red) and PIN1-CFP (blue). **(A)**

Top view of seedling at 3DAS **(B to C)** Longitudinal section of seedling shown in (A).

Dashed line shows dorsoventral axis of first leaf and arrowheads mark dorsal cells

expressing both REV-2×VENUS and *FILp::dsREDN7*. **(D)** Seedlings at 3.5 DAS with

FILp::dsREDN7 expression more restricted to the developing ventral side of the leaf. **(E**

to F) Longitudinal sections of seedling shown in (D) showing a more complementary

pattern of *FILp::dsREDN7* relative to REV-2×VENUS compared to the earlier stage

shown in (A) to (C). Scale bars represent 20 μm.

Supplementary Figure S2.

Frequency of seedlings exhibiting different leaf morphologies after ectopic induction of

KAN1-2×GFP expression in the CLV3 domain (see Figure 6). Class of phenotype

corresponds to those indicated in Figure 6. I to O.

Supplementary Figure S3.

Auxin depletion results in altered patterns of dorsoventral gene expression in vegetative meristems.

(A and B) Confocal projections of the VMs showing expression pattern of REV-2×YPet (red), KAN1-2×GFP (green) and PIN1-CFP (blue) 48 hours after the combined application of 100μM auxinole, 100μM KYN and 100μM Yucasin (auxin depleting drugs). Arrowhead in (B) marks an arrested leaf primordium expressing KAN throughout. Inset in (B) shows a longitudinal optical section of the leaf primordium ectopically expressing KAN1-2×GFP. **(C and D)** Longitudinal optical sections of the VMs in (A) and (B) respectively. Note that the meristems grow as pins with no new organs initiating. Scale bars 30μm (A-D, inset in B).

1011

1012

1013

Supplementary Figure S4.

Wounding induces ectopic KAN expression in inflorescence meristems.

 (**A-H**) Confocal projections of the inflorescence meristems showing expression pattern of KAN1-2×GFP (green) and PIN1-CFP (blue) immediately after ablation (A, C, E, and G) and corresponding meristems 48 hours after ablation (B, D, F and H). Note ectopic KAN1 expression on both the sides of the ablated cells 48 hours after wounding. (**I-L**) Confocal projections of inflorescence meristems showing expression pattern of KAN1-2×GFP (green) and PIN1-CFP (blue) immediately after (I and K) and 48 hours after ablation and treatment with NAA and NPA (J and L). Note lack of ectopic KAN1

1023   expression compared to comparable untreated meristems (B, D, F and H). **(M)**

1024   Quantification of wounding induced ectopic KAN1 expression upon mock treatment

1025   (n=6/6)), NAA application (n=2/6)) and NAA + NPA combination (n= 0/6) application

1026   on the Arabidopsis IMs expressing REV-2×YPet, KAN1-2×GFP and PIN1-CFP. Scale

1027   bars 20μm (A to L).

1028

1029

1030

1031

1032

Supplementary Figure S5.

Wounding induced ectopic KAN expression and reorganization of the vegetative

meristem.

(**A-C**) Confocal projections of the vegetative meristems showing expression pattern of

KAN1-2×YPet (green) and PIN1-GFP (blue) immediately after ablation (A), 48 hours

after ablation (B) and 72hours after ablation (C). Ablated cells are marked by arrowhead

in (A). 72 hours after wounding the vegetative meristem appears split into at least two

distinct meristems with new leaves oriented normally with respect to each meristem (C).

Scale bars 20μm (A and B); 30μm (C).

1042

1043

Table S1.

Primers for cloning and Q-PCR

Gene	Primer name	Sequence
KAN1 (At5g16560)	KAN1g F	ACGCGTTGTTTGGATGTATGACATTAAGTAAGCTAT
KAN1 (At5g16560)	KAN1g R	GGATCCGCTTTCTCGTGCCAATCTGGTCTGCCTAA
KAN1 (At5g16560)	K1cDNA F	AGATCTAACAATGTCTATGGAAGGTGTTTTCCTAGAGAAAAC
KAN1 (At5g16560)	K1 cDNA R	GGATCCGCTTTCTCGTGCCAATCTGGTCTGCCTAA
CLV3 (At2g27250)	CLV3p F	GGAATTCCGGATTATCCATAATAAAAAC
CLV3 (At2g27250)	CLV3p R	CTGCAGGTTTTAGAGAGAAAGTGACTGAGTGA
CLV3 (At2g27250)	CLV3utr F	AAACCTGCAGGGATCCGCGGC
CLV3 (At2g27250)	CLV3p R	ATAGAATATCACTAGTTAATTATCATTGGTTTAAAGTTATAG
WOX1 (At3g18010)	WOX1p F	ggtaccTCAAAACCGGTTTTTATACGACAAGAC-
WOX1 (At3g18010)	WOX1p R	ggatccTTTGGTGTGTACTTAATTTATATGTATG
WOX1 (At3g18010)	WOX1g F	gcggcagcaagatctATGTGGACGATGGGTTACAACGAAG
WOX1 (At3g18010)	WOX1g R	atagaatatcactagtACGTCACTGATGATATACTACG
PRS (At2g28610)	PRSg F	agatctGCGTACGTGTGTACGTGAATGAAAT
PRS (At2g28610)	PRSg R	ggatccAGTTTGGTACTGTCTTGTTTGGAGT
PRS (At2g28610)	PRS FP (Q-PCR)	CAACTCCAAACAAGACAGTACCA
PRS (At2g28610)	PRS RP (Q-PCR)	ACATGAATGAAACACCTGCAGA
WOX1 (At3g18010)	WOX1 FP (Q-PCR)	GCCTCCTTCGTTGTAACCCA
WOX1 (At3g18010)	WOX1 RP (Q-PCR)	GCTGTCTCTCTCCCTTCTCC
IAA20 (At2g46990)	IAA20 FP (Q-PCR)	ATGTGCAATGAGAAGAGTCACG
IAA20 (At2g46990)	IAA20 RP (Q-PCR)	TCACAGTAGACAAGAACATCTCC
ACT2 (At3g18780)	ACT2 FP (Q-PCR)	CCTGTTCTTCTTACCGAGGC
ACT2 (At3g18780)	ACT2 RP (Q-PCR)	AATTTCCCGCTCTGCTGTTG

1044

1045

Table S2.
Frequencies of phenotypes amongst transgenic plant lines

Transgenic plants	N° of T2 lines	T2 line phenotypes
ATML1>>REVr-2×VENUS	31	15 arrest or delay of organogenesis 8 partially or completely dorsalized leaves only 8 no phenotype
UBQ10>>STTM 165/166	26	6 arrested organogenesis 20 partially or completely dorsalized leaves
ATML1>>PHVr	50	4 arrest or delay of organogenesis 31 partially or completely dorsalized leaves 15 no phenotype
ATML1>>KAN1-GFP	17	11 meristem arrest 6 partially or fully radialized organs only
CLV3>>KAN1-2×GFP	32	10 mild change in organ position 10 meristem arrest only 12 leaf morphology change and meristem arrest

For *ATML1>> REVr-2×VENUS* (n=20, number of imaged specimens) and *UBQ10>>STTM 165/166* (n=12) transgenic plants, we used T3 generation plants for imaging that exhibited the reported phenotypes at a frequencies ranging from 70 to 90%. For *ATML1>>KAN1GFP* transgenic plants, we imaged a particular T2 line that exhibited meristem arrest after induction at a frequency of approximately 98% (n=10). For *CLV3>>K2G* transgenic lines, among 12 different T2 lines that showed leaf morphology changes and meristem arrest, we used a particular line that produced more than two leaves after meristem arrest at a frequency of 96.8% (n=65). For *ATML1>>PHVr* transgenic lines, we imaged a line that exhibited arrested organogenesis at a frequency of 33% (n=12). An absence of phenotype was generally associated with low levels of induced transgene expression.

1047

1048

1049

1050

Table S3
List of parameter values used in simulations. We have used the values from Heisler and
Jönsson (2006), which are based on experimental estimates where applicable.

Symbol	Value	Description
c_A	0.001	Auxin production
d_A	0.001	Auxin degradation
T	1.3	Active transport of auxin (PIN1-dependent)
D	0.002	Passive transport of auxin
c_P	0.001	PIN1 production
d_P	0.001	PIN1 degradation
V_X	10.0	Maximal production rate of polarising signal X
K_{XA}	10.0	Hill constant for auxin activating X
n_{XA}	1	Hill coefficient for auxin activating X
K_{XR}, K_{XK}	0.1	Hill constants for REV/KAN repressing X
n_{XR}, n_{XR}	2	Hill coefficients for REV/KAN repressing X
d_X	1.0	Degradation of polarising signal X
k_p	0.9	Relation of symmetric vs polarized PIN1
f_p	0.3	Ratio between PIN1 endo/exocytosis

1075　**Table S4 Statistical details for RT PCR experiment**
1076

	PRS	WOX1	IAA20
Rep1 FC	1.678900384	1.478510137	2.679542279
Rep2 FC	3.216883524	1.991086008	3.602300305
Rep3 FC	1.735102699	3.763611201	2.629479589
Rep4 FC	3.016883524	1.665693641	2.923774058
Rep5 FC	1.335102699	3.099714601	2.479542279
Mean	2.196574566	2.399723118	2.862927702
SD	0.856885899	0.988002492	0.44313845
p-value	0.0142	0.0132	0.0001
Median	1.735102699	1.991086008	2.679542279
SEM	0.383211024	0.441848147	0.198177539
N	5	5	5
95% confidence intervals			
Upper limit	2.947668173	3.265745486	3.251355679
Lower limit	1.445480959	1.53370075	2.474499725

1077
1078

1079

Movie S1

Movie shows confocal 3D projection of a single vegetative seedling corresponding to that

shown in Fig. 6, G and H. Several leaf-like organs have formed on the boundary of

ectopic KAN1 expression driven by the CLV3 promoter (green). These organs express

REV (red) in restricted patterns corresponding to the three classes described in Fig. 6

(labeled Class 2, 3 and 4). Note developing leaf margins, marked by high PIN1

expression (blue) correlate with REV expression boundaries in the epidermis.

1087

1088 **Computational model files (separate file as one zipped archive)**

1089 Archive containing files used for the computational model. Files will be extracted to the

1090 directory modelFiles, and further information is found in the file README.txt in this

1091 directory.

1092

1093

1094

1095

1096

1097