Full Title:

Genome wide identification of bacterial genes required for plant

infection by Tn-seq

Kevin Royet[1], Nicolas Parisot[2], Agnès Rodrigue[1], Erwan Gueguen[1]* and Guy Condemine[1]*

* Both authors share co-last authorship

Corresponding author: erwan.gueguen@univ-lyon1.fr

Short title: *Dickeya dadantii* virulence genes in chicory

1 Univ Lyon, Université Lyon 1, INSA de Lyon, CNRS UMR 5240 Microbiologie

Adaptation et Pathogénie, F-69622 Villeurbanne, France.

2 Univ Lyon, INSA-Lyon, INRA, BF2I, UMR0203, F-69621, Villeurbanne, France

ABSTRACT

Soft rot enterobacteria (*Dickeya* and *Pectobacterium*) are major pathogens that cause diseases

on plants of agricultural importance such as potato and ornamentals. Long term studies to

identify virulence factors of these bacteria focused mostly on plant cell wall degrading

enzymes secreted by the type II secretion system and the regulation of their expression. To

identify new virulence factors we performed a Tn-seq genome-wide screen of a transposon

mutant library during chicory infection followed by high-throughput sequencing. This

allowed the detection of mutants with reduced but also increased fitness in the plant.

Virulence factors identified differed from those previously known since diffusible ones

(secreted enzymes, siderophores or metabolites) were not detected by this screen. In addition

to genes encoding proteins of unknown function that could be new virulence factors, others

could be assigned to known biological functions. The central role of the FlhDC regulatory

cascade in the control of virulence was highlighted with the identification of new members of

this pathway. Scarcity of the plant in certain amino acids and nucleic acids required presence

of the corresponding biosynthetic genes in the bacteria. Their products could be targets for

the development of antibacterial compounds. Among the genes required for full development

in chicory we also identified six genes involved in the glycosylation of the flagellin FliC,

glycosylation, which in other plant pathogenic bacteria contributes to virulence.

Author summary

Identification of virulence factors of plant pathogenic bacteria has relied on the test of

individual mutants on plants, a time-consuming method. New methods like transcriptomic or

proteomic can now be used but they only allow the identification of genes induced during the

infection process and non-induced genes may be missed. Tn-seq is a very powerful method to

identify genes required for bacterial growth in their host. We used for the first time this

method in a plant pathogenic bacteria to identify genes required for the multiplication of

Dickeya dadantii in chicory. We identified about 100 genes with decreased or increased

fitness in the plant. Most of them had no previously described role in bacterial virulence. We

unveiled important metabolic genes and regulators of motility and virulence. We showed that

D. dadantii flagellin is glycosylated and that this modification confers fitness to the bacteria

during plant infection. Our work opens the way to the use of Tn-seq with bacterial

phytopathogens. Assay by this method of large collections of environmental pathogenic

strains now available will allow an easy and rapid identification of new virulence factors.

Introduction

Dickeya are broad-host range phytopathogenic bacteria belonging to the Pectobacteriaceae family [1] that provoke the soft rot disease on many plant species. They are the cause of important losses on economically important crops such as potato, chicory and ornamentals. Identification and studies on the virulence factors of these bacteria have been performed mostly on the model strain D. dadantii 3937 and focused mainly on three domains/aspects, known to be important for disease development: plant cell wall degrading enzymes, the type III secretion system and iron metabolism [2]. Secretion of plant cell wall degrading enzymes has long ago been identified as the bacteria main virulence factor. Many studies focused on the identification and characterization of these secreted enzymes, mostly pectinases [3], of the regulators controlling their production (kdgR, pecS, pecT, hns, gacA), [4-8] of the genes whose expression is coregulated with that of the secreted enzyme genes [9, 10], and of the mechanism of their secretion by the type II secretion system [11]. Although of a lesser importance for Dickeya virulence, the same type of approach has been used to identify type III secretion system regulators and effectors [12] [13] [14]. Moreover, struggling for iron within the plant is strong. D. dadantii acquires this metal through production of two siderophores, chrysobactin and achromobactin [15] [16] [17]. Omics approaches have also been used to identify genes induced during plant infection [18] [19] [20]. These studies now provide a clearer picture on a complex network of factors required for D. dadantii virulence [2, 21]. However, these approaches may have missed some important factors not targeted by these analyses. More global screens need to be performed to identify these factors. Libraries of transposon-induced mutants were tested on plants to find mutants showing reduced virulence with Pectobacterium carotovorum and atrosepticum, two

other soft rot enterobacteria [22-24]. These studies identified auxotrophs, mutants defective

in production or secretion of exoenzymes and in motility. Other mutants with a more

complex phenotype were not characterized at this time. Moreover, the number of tested

mutants was limited by the necessity to test individually each mutant on plant. This type of

work has never been performed on *Dickeya* strains. To have a more complete view of the

genes required for the virulence of *Dickeya*, we used a high-throughput sequencing of a

saturated transposon library (Tn-seq) to screen tens of thousands random insertion mutants of

D. dadantii in laboratory medium and during infection of chicory. Tn-Seq involves creating

large transposon libraries, growing the mutants in a control and a selective condition,

sequencing the transposon insertion sites with next-generation sequencing, mapping sequence

reads to a reference genome and comparing the number of read in each gene in the two

conditions. Tn-seq has been extensively used to uncover essential genes required for mouse

colonization by human pathogens *Vibrio cholerae* [25], *Pseudomonas aeruginosa* [26] and

Streptococcus pneumoniae [27] or plant root colonization by *Pseudomonas simiae* [28] or

multiplication of *Pantoea stewartii* in corn xylem [29]. This latter bacteria relies on the

massive production of exopolysaccharides to block water transport and cause wilting. Thus,

Tn-seq is a very powerful method to identify genes required for bacterial growth in their host.

By applying this technique to screen a *D. dadantii* mutant library in chicory, we identified

metabolic pathways and bacterial genes required by a necrotrophic bacteria for growth *in*

planta. Among them, we found a cluster of genes required for flagellin glycosylation, a

modification known to be important for several plant pathogenic bacteria virulence.

Results and discussion

Characterization of *D. dadantii* 3937 *Himar1* transposon library

Many tools are available to perform Tn-seq [30]. In order to perform a Tn-seq experiment with *D. dadantii* 3937, we used a *Himar9* mariner transposon derivative carrying MmeI restriction sites in the inverted repeats (IR) and a kanamycin resistance cassette between the IRs [31]. We carried out a biparental mating between *E. coli* and *D. dadantii* on M63 agar medium without carbon source and amino acids. We obtained approximately 300 000 colonies that were pooled. Subsequent DNA sequencing (see below) showed the presence of transposon insertions in amino acid, vitamin, purine or pyrimidine biosynthesis pathways, demonstrating that mating on M63 minimal medium does not prevent the obtention of auxotroph mutants. To identify essential genes, mutants were grown in LB medium for several generations. Two DNA libraries were prepared from two cultures and subjected to high-throughput sequencing. The mariner transposon inserts into TA dinucleotides. The TPP software [32] was used to determine the number of reads at each TA site for each biological replicate. *D. dadantii* genome has 171,791 TA sites that can be targeted by the *Himar9* transposase. Pairs of biological replicates were compared. 37,794 and 48,101 unique insertions in TAs were detected in each sample, which corresponds to 22 and 28% density of insertion respectively (Table 1). The average number of reads per TA is 88 and 75, respectively. The results were reproducible with a Pearson correlation coefficient of 72% (Fig. S1) The location of the unique insertions showed an even distribution around the chromosome (Fig. 1A). For each gene, we calculated a $\log_2$ fold change (FC) corresponding to a ratio between the measured number of reads and the expected number of reads. The density plot (Fig. 1B) indicates that essential and non-essential genes are easily distinguishable, confirming the good quality of our Tn-seq libraries.

Then, gene essentiality of the Tn-seq input libraries was determined by using the TRANSIT software [32]. We decided to use the Hidden Markov Model (HMM) which predicts

essentiality and non-essentiality for individual insertion sites since it has been shown to give

good prediction in datasets with density as low as 20% [32]. The HMM analysis led to the

identification of 665 genes essential for growth in LB (ES), representing 14% of the genes of

D. dadantii 3937, a number in the range of those found for this type of analysis with bacteria.

The transposon we used does not allow us to discriminate between the direct effect of the

insertion or a polar effect on downstream genes. Goodall et al [33] have shown that this

overestimates the number of essential genes. Thus 665 must be considered has an upper limit

of the number of essential genes.

552 genes were categorized as Growth Defect genes (GD, i.e. mutations in these genes lead

to loss of fitness), 125 as growth advantage genes (GA, i.e mutations in these genes lead to

gain of fitness) and 3320 as non-essential genes (NE) (Table S5 and Fig. 1B).

TABLE 1 Tn-Seq analysis of *Dickeya dadantii* 3937

Mutant pool	Total no. of reads	No. of reads containing Tn end	No. of reads normalized[a]	No. of mapped reads to unique TA sites	No. of mapped reads to unique TA sites after LOESS correction	Density (%)[b]	Mean read count per TA[c]
LB #1	23,152,186	22,647,343	18,748,028	13,166,770 (70 %)	12,904,900 (69 %)	28 %	75
LB #2	30,105,412	27,963,154	18,748,028	15,535,291 (83 %)	15,195,582 (81 %)	22 %	88
Chicory #1	18,925,029	18,748,028	18,748,028	17,535,146 (94 %)	14,906,888 (79 %)	24 %	87
Chicory #2	27,607,717	26,555,297	18,748,028	17,477,706 (93 %)	16,955,724 (90 %)	23 %	99

[a] The number of reads containing the sequence of a Tn end were normalized for each sample according to the Chicory #1

[b] *Dickeya dadantii* 3937 genome has 171,791 TA sites. The density is the % of TAs for which mapped reads has been assigned by the TPP software.

[c] The mean value of mapped reads per TA

Genes necessary for chicory leaf maceration. We used chicory leaf infection as a model to

identify *D. dadantii* genes required for growth in plant tissues. Biological duplicates were

performed to insure the reproducibility of the results. Each chicory was inoculated with 10^7

bacteria from the mutant pool and after 2 days more than 10^{10} bacteria were collected from

the rotten tissue. Sequencing transposon insertion sites in these bacteria followed by the TPP

analysis indicated a density of unique insertion in TAs comparable to that of the input

datasets (23-24%). Surprisingly, the results were more highly reproducible than in LB with a

very high Pearson correlation coefficient of 98% (Fig. S1).

In order to test the statistical significance of the identified genes conferring to *D. dadantii* a

loss or a gain of fitness *in planta*, we performed the RESAMPLING (permutation test)

analysis of the TRANSIT software. The RESAMPLING method is a variation of the classical

permutation test in statistics that sums the reads at all TA sites for each gene in each

condition. It then calculates the difference of the sum of read-counts between the input (LB)

and output (chicory) datasets. The advantage of this statistical method is to attribute for each

gene an adjusted p-value (q-value). Genes with a significant difference between total read-

counts in LB and chicory achieve a q-value ≤ 0.05. The method also calculates a $\log_2$ fold-

change ($\log_2$FC) for each gene based on the ratio of the sum of read counts in the output

datasets (chicory) versus the sum of read counts in the input (LB) datasets [32]. Applied to

our Tn-seq datasets and selecting only genes achieving a FDR adjusted p-value (q-value) $\leq$

0.05, we identified 122 genes out of 4666 required for fitness *in planta*, as shown with the

volcano plot of RESAMPLING results comparing replicates grown in LB versus *in planta*

(Fig. S2). For these 122 genes, we applied an additional cutoff by removing 20 genes with a

mean read count in LB <5 (less than 5 reads in average / TA). These genes were categorized

as ES or GD in LB. We also removed from the analysis 6 genes with a $\log_2$FC comprised

between -2 and 2. By applying all these criteria, we retained only 96 genes for a further

analysis (Table 2). 92 of them were identified as GD genes in the chicory ($\log_2$FC $\leq$2), the 4

left as GA genes in the chicory ($\log_2$FC $\geq$2). A possible polar effect for genes being part of an

operon is analysed in Table 2: if a GD gene is upstream of another GD gene in the same

operon, a polar effect of insertions in the first gene on the second one cannot be excluded.

Some of these genes, in bold in Table 2, were already known to play a role in *D. dadantii*

virulence, confirming the validity of the Tn-seq approach. Using the Kyoto Encyclopedia of

Genes and Genomes (KEGG) [34], we discovered that certain metabolic pathways and

biological functions are very important for growth in chicory (Table S4). We highlight some

of them in the next sections of the article.

TABLE 2 Genes identified by Tn-seq exhibiting a fitness change in chicory compared to LB.

| | | | HMM | RESAMPLING | | | | | | | |
| | | | | | Mean reads[d] | | | | | | |
Locus[a]	Gene[a]	Function	State in LB[b]	No. of TAs[c]	LB	Chicory	ΔSum	log₂FC[e]	q-value[f]	In operon[g]	genes in operon (state)[h]
Dda3937_00335	*glpD*	glycerol-3-phosphate dehydrogenase	GD	33	650	0	-11,706	-12.56	0.00	N	
Dda3937_03379	*purL*	phosphoribosylformyl-glycineamide synthetase	NE	73	378	0	-21,944	-11.91	0.00	N	
Dda3937_03564	***opgG***	**Glucans biosynthesis protein G precursor**	**GA**	**40**	**1976**	**1**	**-90,843**	**-11.41**	**0.00**	**Y**	*opgG (-11.41) opgH (-9.79)*
Dda3937_00244	*purH*	phosphoribosylaminoimidazolecarboxamide formyltransferase/IMP cyclohydrolase	NE	37	145	0	-2,896	-11.25	0.00	Y	*purD (-1.66)* **purH (-11.25)**
Dda3937_00432	*hflK*	FtsH protease regulator	GD	28	339	0	-4,060	-11.12	0.03	Y	**hflK (-11.12)** *hflC (+0.06) yjeT (-1.38)*
Dda3937_02515	*purM*	phosphoribosylaminoimidazole synthetase	NE	21	344	0	-6,188	-10.57	0.00	Y	**purM (-10.57)** *purN (0)*
Dda3937_02627		4-hydroxythreonine-4-phosphate dehydrogenase	NE	26	129	0	-2,065	-10.06	0.00	Y	**Dda3937_02627 (-10.06)** *Dda3937_02626 (-3.77)*
Dda3937_00004	*guaB*	IMP dehydrogenase	NE	33	151	0	-3,915	-9.97	0.00	N	
Dda3937_03563	***opgH***	**Glucans biosynthesis glucosyltransferase H**	**GA**	**62**	**1409**	**2**	**-90,073**	**-9.79**	**0.00**	**Y**	*opgG (-11.41) opgH (-9.79)*
Dda3937_01284	*pyrB*	aspartate carbamoyltransferase	NE	17	159	0	-1,910	-9.68	0.00	Y	**pyrB (-9.68)** *pyrI (+1.33)*
Dda3937_03924	***rffG***	**dTDP-glucose 4,6-dehydratase**	**NE**	**23**	**317**	**1**	**-3,167**	**-9.38**	**0.02**	**Y**	**rffG (-9.38)** *rffH (-3.49) rfbC (-0.53) rfbD (-0.91)*
Dda3937_01389	*carB*	carbamoyl-phosphate synthase large subunit	NE	48	249	0	-7,967	-9.23	0.00	N	
Dda3937_03299	***acrA***	**MexE family multidrug efflux RND transporter periplasmic adaptor subunit**	**NE**	**34**	**196**	**0**	**-5,860**	**-9.03**	**0.00**	**Y**	*acrA (-9.03) acrB(-8.9)*
Dda3937_03300	***acrB***	**multidrug efflux system protein**	**NE**	**89**	**422**	**1**	**-31,986**	**-8.90**	**0.00**	**Y**	*acrA (-9.03) acrB(-8.9)*
Dda3937_03258	*pyrE*	orotate phosphoribosyltransferase	NE	14	175	0	-2,788	-8.81	0.00	N	
Dda3937_02336	*nlpI*	lipoprotein	GD	33	27	0	-601,000	-8.69	0.00	N	
Dda3937_02506	*nlpB (bamC)*	outer membrane protein assembly factor BamC	NE	20	47	0	-841,000	-8.69	0.00	Y	*dapA (+2.02)* **bamC (-8.69)**
Dda3937_04018	*pta*	phosphate acetyltransferase	GD	36	579	2	-10,400	-8.59	0.02	N	
Dda3937_03554	*pyrC*	dihydro-orotase	NE	25	343	1	-7,534	-8.44	0.00	N	
Dda3937_04573	*lpxM*	acyl (myristate) transferase	NE	33	63	0	-1,764	-8.31	0.00	N	
Dda3937_01116	*glnG*	Nitrogen regulation protein NR(I), Two-component system	NE	26	39	0	-629,000	-8.22	0.00	Y	*glnL (-0.2)* **glnG (-8.22)**
Dda3937_02099	*purF*	amidophosphoribosyltransferase	NE	32	107	0	-2,779	-8.19	0.00	Y	**purF (-8.19)** *cvpA (-1.92)*
Dda3937_04019	*ackA*	acetate kinase A and propionate kinase 2	NE	29	45	0	-1,063	-8.16	0.00	Y	*Dda3937_04020 (-2.48)* **ackA (-8.16)**
Dda3937_02189	*yejM*	Membrane-anchored periplasmic protein, alkaline phosphatase superfamily	GA	34	4160	15	-99,478	-8.08	0.00	Y	*yejL (0)* **yejM (-8.08)**
Dda3937_01390	*carA*	carbamoyl-phosphate synthase small subunit	NE	21	69	0	-956,000	-8.05	0.00	N	
Dda3937_01426	*ptsI*	Phosphoenolpyruvate-protein phosphotransferase of PTS system	NE	33	45	0	-1,176	-7.85	0.00	Y	*crr (-2.66)* **ptsI (-7.85)** *ptsH (0)*
Dda3937_00161	*cysQ*	3'(2'),5'-bisphosphate nucleotidase	NE	16	44	0	-434,000	-7.81	0.02	N	
Dda3937_00210	*cysI*	sulfite reductase beta subunit	NE	40	252	1	-7,515	-7.65	0.00	Y	*cysH (-8.93)* **cysI (-7.65)** *cysJ (-6.25)*
Dda3937_04075	*lysR*	LysR family transcriptional regulator	NE	13	2385	13	-18,976	-7.51	0.00	N	
Dda3937_02526	*yidR*	conserved protein	NE	18	50	0	-591,000	-7.50	0.00	N	
Dda3937_03888	*metB*	Cystathionine gamma-synthase	NE	21	118	1	-1,881	-7.34	0.01	Y	**metB (-7.34)** *metL (-3.23)*
Dda3937_00195	*relA*	(p)ppGpp synthetase I/GTP pyrophosphokinase	NE	55	256	2	-11,683	-7.12	0.00	Y	**relA (-7.12)** *rumA (-1.33)*
Dda3937_02532	***lfcR***	**Fructose repressor FruR, LacI family**	**NE**	**15**	**399**	**3**	**-4,756**	**-7.04**	**0.00**	**N**	
Dda3937_02226	*fliF*	Flagellar M-ring protein fliF	NE	46	476	4	-18,898	-7.02	0.00	Y	**fliF (-7.02)** *fliG (-4.26) fliH (-3.92)* **fliI (-6.56)** *fliJ (-5.44) fliK (-4.71)*
Dda3937_02206	*flgE*	Flagellar hook protein flgE	NE	50	597	5	-29,608	-7.00	0.00	Y	**flgE (-7)** *flgF (-4.76)* **flgG (-5.91)**
Dda3937_04507	*gnd*	phosphogluconate dehydrogenase (NADP(+)-dependent, decarboxylating)	GD	36	7	0	-190,000	-6.91	0.00	N	
Dda3937_00697	*degQ*	Protease	NE	28	80	1	-956,000	-6.87	0.01	N	
Dda3937_03631	*trxB*	thioredoxin-disulfide reductase	GD	25	16	0	-257,000	-6.85	0.03	N	
Dda3937_00361	*yrfF (igaA)*	intracellular growth attenuator protein	GD	38	22	0	-430,000	-6.78	0.03	N	
Dda3937_00588	*cysB*	Transcriptional dual regulator, O-acetyl-L-serine-binding protein	NE	29	90	1	-2,504	-6.75	0.00	N	
Dda3937_03783	*prc*	carboxy-terminal protease for penicillin-binding protein 3	NE	46	243	2	-11,557	-6.71	0.00	Y	**prc (-6.71)** *proQ (-1.82)*
Dda3937_00433	*hflX*	predicted GTPase	GD	27	16	0	-187,000	-6.69	0.04	N	
Dda3937_03427	*fliC*	flagellar filament structural protein (flagellin)	NE	33	96	1	-1,520	-6.61	0.03	N	
Dda3937_02223	*fliI*	Flagellum-specific ATP synthase fliI	NE	42	236	3	-7,009	-6.56	0.00	Y	*fliF (-7.02) fliG (-4.26) fliH (-3.92)* **fliI (-6.56)** *fliJ (-5.44) fliK (-4.71)*
Dda3937_04419	*hdfR*	DNA-binding transcriptional regulator	NE	29	117	1	-3,241	-6.34	0.00	N	
Dda3937_00209	*cysJ*	sulfite reductase alpha subunit	NE	41	180	2	-6,746	-6.25	0.00	Y	*cysH (-8.93) cysI (-7.65)* **cysJ (-6.25)**
Dda3937_02209	*flgH*	Flagellar L-ring protein flgH	NE	23	586	8	-13,875	-6.22	0.01	Y	**flgH (-6.22)** *flgI (-5.49) flgJ (-7.16)*
Dda3937_02246	*fabF*	beta-ketoacyl-[acyl-carrier-protein] synthase II	GD	41	10	0	-273,000	-6.15	0.00	N	

Gene ID	gene	description	State	c1	c2	c3	Difference	log2 ratio	P-value	Operon	Operon structure
Dda3937_00301	uvrD	ATP-dependent DNA helicase UvrD/PcrA	NE	42	29	0	-678,000	-6.11	0.00	N	
Dda3937_02212	flgK	Flagellar hook-associated protein flgK	NE	63	116	2	-4,808	-6.07	0.00	Y	**flgK (-6.07)** flgL (-5.58)
Dda3937_04046	purU	Formyltetrahydrofolate deformylase	NE	28	51	1	-1,105	-5.84	0.00	N	
Dda3937_03965	flhA	predicted flagellar export pore protein	NE	49	106	2	-3,532	-5.80	0.00	Y	flhE (-0.89) **flhA (-5.8) flhB (-5.31)** Dda3937_04633 (-1) cheZ (-3.29) cheY (-4.52) **cheB (-5.14) cheR (-4.67)**
Dda3937_02205	flgD	Flagellar basal-body rod modification protein flgD	NE	22	227	4	-4,905	-5.73	0.01	Y	flgB (-3.45) **flgC (-6.38) flgD (-5.73)**
Dda3937_01352	leuC	3-isopropylmalate dehydratase large subunit	NE	21	139	3	-2,457	-5.73	0.01	Y	**leuA (-4.69) leuB (-4.63) leuC (-5.73)** leuD (-6.26)
Dda3937_02784	flhC	Flagellar transcriptional activator flhC	NE	20	477	9	-11,222	-5.66	0.01	Y	**flhC (-5.66)** flhD (-4.1)
Dda3937_02782	motB	Flagellar motor rotation protein motB	NE	40	109	2	-4,067	-5.55	0.01	Y	**motA (-5.06) motB (-5.55) cheA (-4.89)** cheW (-5.39)
Dda3937_02210	flgI	Flagellar P-ring protein flgI	NE	26	163	4	-3,191	-5.49	0.00	Y	**flgH (-6.22) flgI (-5.49)** flgJ (-7.16)
Dda3937_02222	fliJ	Flagellar protein fliJ	NE	14	182	4	-2,486	-5.44	0.03	Y	**fliF (-7.02)** fliG (-4.26) fliH (-3.92) **fliI (-6.56) fliJ (-5.44)** fliK (-4.71)
Dda3937_02219	fliM	Flagellar motor switch protein fliM	NE	27	143	3	-3,339	-5.40	0.00	Y	fliL (-4.17) **fliM (-5.4)** fliN (-4.78) fliO (-6.89) fliP (-4.78) fliQ (-3.12) **fliR (-4.56)**
Dda3937_02774	flhB	Flagellar biosynthesis protein flhB	NE	32	186	5	-4,712	-5.31	0.00	Y	flhE (-0.89) **flhA (-5.8) flhB (-5.31)** Dda3937_04633 (-1) cheZ (-3.29) cheY (-4.52) **cheB (-5.14) cheR (-4.67)**
Dda3937_02777	cheB	Chemotaxis response regulator protein-glutamate methylesterase CheB	NE	31	282	8	-7,682	-5.14	0.00	Y	flhE (-0.89) **flhA (-5.8) flhB (-5.31)** Dda3937_04633 (-1) cheZ (-3.29) cheY (-4.52) **cheB (-5.14) cheR (-4.67)**
Dda3937_02783	motA	Flagellar motor rotation protein motA	NE	24	39	1	-834,000	-5.06	0.00	Y	**motA (-5.06) motB (-5.55) cheA (-4.89)** cheW (-5.39)
Dda3937_00565	tonB	TonB protein	NE	14	106	3	-2,062	-5.00	0.05	N	
Dda3937_00427	fbp	fructose-bisphosphatase	GA	33	805	27	-28,026	-4.92	0.01	N	
Dda3937_02781	cheA	Chemotaxis protein CheA	NE	50	151	5	-5,838	-4.89	0.00	Y	**motA (-5.06) motB (-5.55) cheA (-4.89)** cheW (-5.39)
Dda3937_03422		Carbamoyl-phosphate synthase small subunit	NE	43	379	13	-11,713	-4.85	0.02	Y	Dda3937_03422 (-4.85) Dda3937_03421 (-0.71)
Dda3937_02577	lysA	diaminopimelate decarboxylase	NE	23	332	0	-3,989	-4.79	0.00	N	
Dda3937_02207	flgF	Flagellar basal-body rod protein flgF	NE	21	35	1	-671,000	-4.76	0.00	Y	**flgE (-7) flgF (-4.76) flgG (-5.91)**
Dda3937_02230	fliD	Flagellar hook-associated protein fliD	NE	47	93	3	-2,506	-4.75	0.00	N	
Dda3937_04301	leuA	2-isopropylmalate synthase	NE	36	35	1	-944,000	-4.69	0.02	Y	**leuA (-4.69) leuB (-4.63) leuC (-5.73)** leuD (-6.26)
Dda3937_02778	cheR	Chemotaxis protein methyltransferase CheR	NE	30	462	18	-8,882	-4.67	0.05	Y	flhE (-0.89) **flhA (-5.8) flhB (-5.31)** Dda3937_04633 (-1) cheZ (-3.29) cheY (-4.52) **cheB (-5.14) cheR (-4.67)**
Dda3937_02228	fliT	Flagellar biosynthesis protein fliT	GD	16	8	0	-95,000	-4.63	0.05	Y	fliS (-6.36) **fliT (-4.63)**
Dda3937_04404	leuB	3-isopropylmalate dehydrogenase	NE	16	285	12	-3,835	-4.63	0.05	Y	**leuA (-4.69) leuB (-4.63) leuC (-5.73)** leuD (-6.26)
Dda3937_02214	fliR	Flagellar biosynthesis protein fliR	NE	33	268	11	-5,653	-4.56	0.05	Y	fliL (-4.17) **fliM (-5.4)** fliN (-4.78) fliO (-6.89) fliP (-4.78) fliQ (-3.12) **fliR (-4.56)**
Dda3937_03727	**kduI**	**4-deoxy-L-threo-5-hexosulose-uronate ketol-isomerase**	**NE**	**26**	**70**	**3**	**-2,015**	**-4.54**	**0.03**	N	
Dda3937_03267		O-antigen, teichoic acid lipoteichoic acids export membrane protein	ES	107	89	4	-1,181	-4.33	0.05	Y	**Dda3937_03267 (-4.33)** Dda3937_03268 (-1.07)
Dda3937_00415	epd	D-erythrose 4-phosphate dehydrogenase	NE	26	316	16	-4,793	-4.27	0.02	N	
Dda3937_02337	pnp	polynucleotide phosphorylase/polyadenylase	GD	50	5	0	-105,000	-3.97	0.00	N	
Dda3937_01683	purK	N5-carboxyaminoimidazole ribonucleotide synthase	NE	16	90	0	-722,000	-3.49	0.01	Y	purE (-5.75) **purK (-3.49)**
Dda3937_00689	yrbF (mlaF)	predicted toluene transporter subunit	GA	9	1254	114	-15,962	-3.47	0.01	Y	**yrbF (-3.47)** yrbE (-1.48) **yrbD (-3.09) yrbC (-2.81)** yrbB (-0.24))
Dda3937_02829	helD	DNA helicase IV	NE	26	99	9	-1,803	-3.46	0.01	N	
Dda3937_02252	ptsG	PTS system glucose-specific IICB component	NE	37	81	8	-2,928	-3.38	0.03	N	
Dda3937_00726	**tolC**	**transport channel**	**NE**	**34**	**184**	**0**	**-3,304**	**-3.35**	**0.00**	N	
Dda3937_02363	clpA	ATP-dependent Clp protease ATP-binding subunit	NE	44	64	8	-1,793	-3.02	0.03	Y	clpS (-2.07) **clpA (-3.02)**
Dda3937_02470	corC	magnesium and cobalt ions transport	NE	13	159	21	-1,377	-2.90	0.02	Y	lnt (+3.02) **corC (-2.09)**
Dda3937_00692	yrbC (mlaC)	predicted ABC-type organic solvent transporter	GA	23	740	106	-16,493	-2.81	0.01	Y	**yrbF (-3.47)** yrbE (-1.48) **yrbD (-3.09) yrbC (-2.81)** yrbB (-0.24)
Dda3937_02045	envC	murein hydrolase activator	NE	17	71	12	-825,000	-2.59	0.00	N	
Dda3937_01807	nuoM	NADH-quinone oxidoreductase subunit M	NE	29	57	10	-1,130	-2.47	0.03	Y	nuoN (-2.01) **nuoM (-2.47)**
Dda3937_03668	sufB	Fe-S cluster assembly protein	NE	32	116	21	-3,581	-2.44	0.00	Y	**sufB (-2.44)** sufA (-1.47)
Dda3937_02080	trkH	Potassium uptake protein	NE	36	65	13	-1,047	-2.33	0.05	Y	pepQ (-0.21) ; yigZ (+0.1) **trkH (-2.33)** hemG (+1.15)
Dda3937_03042	**fct**	**ferrichrysobactin outer membrane receptor**	**NE**	**80**	**244**	**51**	**-14,622**	**-2.25**	**0.01**	N	
Dda3937_01287	argI	Ornithine carbamoyltransferase	NE	24	279	59	-4,383	-2.23	0.03	N	
Dda3937_02456	rsmC	global regulatory protein RsmC	NE	10	116	221,705	2,659,067	10.90	0.028	N	
Dda3937_03858	gcpA	hypothetical protein	GA	55	3728	140,136	9,002,975	5.23	0.00	N	
Dda3937_03971	mltD	outer membrane-bound lytic murein transglycosylase D	NE	46	276	10,885	445,590	5.30	0.00	N	
Dda3937_00363	mrcA	penicillin-binding protein 1A (PBP1A)	NE	53	85	468	16,879	2.47	0.021	N	

180   [a] Genes for which a role in *D. dadantii* virulence has been described before are in bold. Underlined genes have been deleted to study the mutants in further analysis.

181   [b] State of each gene in LB defined by the TRANSIT software using an Hidden Markov Model: NE, Non-Essential ; GD, Growth-Defect ; E, Essential ; GA, Growth-Advantage.

182   [c] Mean reads per TA site for a gene in each growth condition

183   [d] Difference of reads between chicory and LB growth condition

184   [e] Ratio of reads between chicory and LB condition expressed in $\log_2$

185   [f] P-values adjusted for multiple comparisons using the Benjamini-Hochberg procedure (See Transit manual)

186   [g] Presence of the gene in an operon (Yes or No)

187   [h] Operon structure determined by analysis of *D. dadantii* 3937 RNA-seq datasets from Jiang X *et al*, Environ Microbiol. 2016 Nov;18(11):3651-3672. $\log_2$FC for each gene in operon are indicated in brackets, genes considered to be essential in chicory are indicated in bold (q-value <0.05).

Analysis of the genes of *D. dadantii* required for plant colonization.

(i) Metabolism. Chicory appears as an environment in which amino acids, nucleic acids and

some vitamins (pyridoxal) are scarce. Of the 92 genes identified as GD genes *in planta*, 8 are

involved in purine and 7 in pyrimidine metabolisms (Table S4). In the purine metabolism

pathway, the inosine monophosphate (IMP) biosynthesis pathway that produces IMP from L-

glutamine and 5-phosphoribosyl diphosphate is particularly important for *D. dadantii in*

planta since 5 out of the 10 genes of this pathway are significantly GD genes *in planta* (Fig.

2). IMP is the precursor of adenine and guanine. Next, IMP can be converted in xanthosine

5'-phosphate (XMP) by the IMP dehydrogenase GuaB. *guaB* gene is also a GD gene *in*

planta, with a strong $\log_2$FC of -10.06 (Fig. 2). In the pyrimidine synthesis, the uridine

monophosphate (UMP) biosynthesis pathway that converts L-glutamine to UMP, a precursor

of uracyl, is very important *in planta* since *carAB*, *pyrB*, *pyrC* and *pyrE*, involved in this

enzymatic pathway, are all required for growth *in planta* (Fig. 2). This pyrimidine

biosynthesis pathway is specific to bacteria. It is noteworthy that in the human pathogen *S.*

pneumoniae, mutants of this pathway have a fitness defect in the nasopharynx of infected

mice [27]. Hence, it looks that the pyrimidine biosynthesis pathway is particularly important

for multiplication of some bacterial species in the host.

Mutants in genes involved in the synthesis of sulfur-containing amino acids (*cysIJQ, metB),*

lysine (*lysA*) and leucine (*leuABC)* are disadvantaged in chicory (Table 2 and Fig. 3A). These

amino acids are known to be present in low concentration in plant tissues. Other amino acids

seem to be present in quantity sufficient for growth of *D. dadantii* auxotrophs. Low level of

certain amino acids probably induces the stringent response in the bacteria. Reduced growth

in the plant of the *relA* mutant, unable to synthesize the alarmone ppGpp, supports this

hypothesis. Glucose is the main sugar present in plant tissue, present as a circulating sugar or

a cellulose degradation product. Mutants in the PTS glucose transport system genes *ptsI* and

ptsG have a reduced growth in bacteria (Table 2) showing its importance as a carbon source

in planta.

Degradation of cell wall pectin by a battery of extracellular enzymes is the main determinant

of *Dickeya* pathogenicity. Mutants unable to produce or to secrete these enzymes by the type

II secretion system were not disfavored in chicory since these mutants could use for their

growth the pectin degradation compounds produced by enzymes secreted by other bacteria.

The redundancy of oligogalacturonate specific porins (KdgM and KdgN) and inner

membrane transporters (TogT and TogMNABC) allow entry of these compounds into the

bacteria even in a mutant in one of these transport systems. However, *kduI* mutants, blocked

in the intracellular part of the pectin degradation pathway, have a limited growth *in planta*,

confirming the importance of the pectin degradation pathway in the disease progression.

(ii) Stress resistance. Plant is an hostile environment for the bacteria that has to cope with

antimicrobial peptides, ROS, toxic compounds and acidic pH [35]. We observed that the

pump AcrABTolC, that can efflux a wide range of compounds [36], is important for survival

in chicory (Fig. S3). Stress can lead to the accumulation of phospholipids in the outer

membrane. This accumulation makes the bacteria more sensitive to small toxic molecules

[37]. Such a phospholipid accumulation probably occurs when the bacteria infect chicory

since *mlaC* and *mlaF* mutants, which are unable to prevent phospholipid accumulation in the

outer membrane, have a reduced growth in plant. Production of exopolysaccharides (EPS)

was shown to protect the bacteria during the first steps of infection [9]. We observed that

rffG and *wzx* mutants unable to synthesize EPS have a growth defect in chicory. A set of

genes required to repair or degrade altered proteins (*clpA, degQ, trxB*) are also important for

survival *in planta*. No gene directly involved in detoxification of ROS was detected in our

analysis. However, ROS can create DNA damage. The two helicases involved in DNA repair,

UvrD and HelD, give growth advantage in plant. Osmoregulated periplasmic glycans (OPG)

are polymers of glucose found in the periplasm of α, β and γ-proteobacteria. Their exact role

is unknown but their absence leads to avirulence in certain bacteria such as *D. dadantii* [38].

This absence induces a membrane stress that is sensed and transduced by the Rcs envelope

stress response system. This system controls the expression of many genes, including those

involved in motility, and those encoding plant cell wall degrading enzymes through the

RsmA-RsmB system [39-41]. Thus, mutants defective in OPG synthesis are expected to have

a reduced virulence. Indeed, in our experiment, mutants in the two genes involved in OPG

synthesis, *opgG* and *opgH* were non competitive in chicory (Table 2).

(iii) Iron uptake. *D. dadantii* produces two types of siderophores, achromobactin and

chrysobactin, that are required for the development of maceration symptoms in the iron

limited environment of plant hosts [42]. Once iron loaded, the siderophores are imported into

the bacteria. Import through the outer membrane requires a specific outer membrane channel

and the energy transducing complex formed by TonB ExbB and ExbD. While the absence of

synthesis of one of the siderophores can be compensated by the presence of siderophore

secreted by other bacteria in the growth medium, mutants of the TonB complex are totally

unable to acquire iron and thus are unable to grow in the plant. In accordance, *tonB* was

essential in chicory while the genes coding for siderophore synthesis or secretion were not.

Similarly a mutant devoid of the iron-loaded chrysobactin transport gene (*fct)* is non-

competitive.

(iv) Regulation. Mutants in several genes controlling virulence factor production have a

growth defect in the plant. The master regulator FlhDC acts as a regulator of both flagella and

virulence factor synthesis in many bacteria such as *Yersinia ruckeri*, *Edwardsiella tarda* and

Ralstonia solanacearum [43-45]. In *D. dadantii* FlhDC has recently been shown to control, in

addition to flagellar motility, type III secretion system and virulence factor synthesis through

several pathways [46]. We observed that *flhC* gives a growth advantage in chicory. In

addition, we uncovered that some genes regulating *flhDC* in other bacteria regulate *D.*

dadantii virulence, probably by controlling *flhDC* expression. *rsmC* is a poorly characterized

gene in *D. dadantii* but that has been studied in *Pectobacterium carotovorum*. It negatively

controls motility and extracellular enzyme production through modulating transcriptional

activity of FlhCD [47]. HdfR is a poorly characterized LysR family regulator that controls the

std fimbrial operon in *S. enterica* and FlhDC expression in *E. coli* [48]. *rsmC* mutants were

overrepresented in the chicory (Fig. 3B), indicating a gain of virulence for these mutants.

hdfR conferred fitness benefits during growth in chicory and could also act in *D. dadantii* as

activator of *flhDC* expression.

The GGDEF proteins are c-di-GMP synthase. Their genes are often located next to their

cognate EAL diguanylate phosphodiesterase gene. *ecpC (yhjH)* encodes an EAL protein that

was shown to activate virulence factor production in *D. dadantii* [49]. *gcpA*, which is located

next to *ecpC* encodes a GGDEF protein. Role of *gcpA* in *D. dadantii* virulence has recently

been described [50]. We observed that *gcpA* mutants (Dda_03858) were overrepresented in

chicory (Table 2). This increased virulence, a phenotype opposite to that described for the

ecpC mutants, indicates that overproduction of c-di-GMP could reduce *D. dadantii* virulence.

Of the eighteen regulators of the LacI family present in *D. dadantii*, four of them were found

to be involved in plant infection [51]. One of those, LfcR, which has been found important

for infection of chicory, Saintpaulia and *Arabidopsis*, was identified as important for chicory

infection in our experiment. LfcR is a repressor of adjacent genes [51]. Surprisingly none of

these genes appeared to play a role for chicory infection suggesting that other targets of LfcR

probably remain to be discovered.

Finally, it is noteworthy to mention that the *ackA* and *pta* genes are GD *in planta*. These

genes constitute the reversible Pta-AckA pathway. The steady-state concentration of acetyl-

phosphate (acetyl-P), a signaling molecule in bacteria, depends upon the rate of its formation

catalyzed by Pta and of its degradation catalyzed by AckA [52]. The GD phenotype of *D.*

dadantii ackA and *pta* mutants during infection suggests that acetyl-P might play a crucial

signaling role in the adaptation of *D. dadantii* to the plant tissue.

(v) Motility. Motility is an essential virulence factor of *D. dadantii* required to move on the

surface of the leaf, enter the wounds and propagate into the plant tissue [53-55]. Accordingly,

all the genes required for flagella synthesis, the flagella motor and genes regulating their

synthesis (*flhC*, *flhD*, *fliA*) (see above) are necessary for fitness during chicory infection (Fig.

3C and 5A). All the genes responsible for the transduction of the chemotaxis signal (*cheA*, *B*,

R, *W*, *X*, *Y* and *Z*) also confer a benefit *in planta* (Table 2). No methyl-accepting

chemoreceptor gene mutant was found. Like other environmental bacteria, *D. dadantii*

encodes many such proteins (47). They probably present some redundancy in the recognized

signal which prevented their detection by our screen.

D. dadantii **flagellin is modified by glycosylation**

A group of six genes located between *fliA* and *fliC* retained our interest since insertions in

these genes led to a growth defect in chicory (Fig. 4A). This effect does not result from

insertions in the first gene of the group since they are not expressed in operon [56].

Dda3937_03424 encodes an O-linked N-acetylglucosamine transferase and Dda3937_03419

encodes a protein with a nucleotide diphospho sugar transferase predicted activity. The other

ones could be involved in the modification of sugars (predicted function of Dda3937_03423:

nucleotide   sugar   transaminase,   Dda3937_03422:   carbamoyl   phosphate   synthase,

Dda3937_03421: oxidoreductase; Dda3937_03420: methyltransferase). Their location let

suppose that this group of genes could be involved in flagellin glycosylation. Analysis by

SDS-PAGE of FliC produced by the wild type, and the Dda3937_03424 and Dda3937_03419

mutants, showed that in the two latter strains the molecular weight of the protein diminished

(Fig. 4B). The molecular weight determined by mass spectroscopy was 28,890 Da for

FliC$_{A4277}$, 31,034 Da for FliC$_{A3422}$ and 32170 Da for the WT FliC. Thus, in the wild type

strain FliC is modified by the products of the genes Dda_03424 to Dda_03419, probably by

multiple glycosylation with a disaccharide. Absence of modification did not modify *D.*

dadantii motility (data not shown). The flagellin of the plant pathogens *Pseudomonas*

syringae pv *tabaci* and *Burkholderia cenocepacia* are also glycosylated and absence of this

modification lowered the ability of these bacteria to cause disease on tobacco and

Arabidopsis, respectively [57, 58]. Accordingly, in *D. dadantii*, FliC modification appears

important for multiplication of the bacteria in plant.

Additional genes could be involved in virulence

Several genes have a log$_2$FC >4 or <-4 but do not satisfy the statistical permutation test

adjusted by the false discovery rate method (q-value) (table S6). However, most of them

belong to the categories described above and could be required for growth *in planta*. Among

those with a log$_2$FC< -4 can be found genes involved in amino acid and nucleic acid synthesis

(*cysH, ilvC, pyrF, pyrD, purC, thrC, metA, cysK, lysC*), flagella and motility *(flgJ, fliO, flgC,*

fliS, flgG, flgA, flgL, cheW, fliN, fliP, fliK, fliG, fliL), pectin and glucose metabolism (*kduD,*

pgi), EPS synthesis (*gmd*), flagella glycosylation (*vioA*) and regulation (*zur*, *ecpC* and the

general RNA chaperone *hfq*).

Among the genes with a log$_2$FC > 4, three regulators can be noticed: *pecS*, *pecT* and

Dda3937_00840. *pecS* and *pecT* are known regulators of *D. dadantii* controlling the

expression of many factors involved in virulence [5, 6]. Thus, their mutation could confer an

increased fitness of the bacteria in chicory. *D. dadantii* possesses a functional *expI-expR*

quorum sensing system which does not seem to control plant virulence factor production

[59]. However, several LuxR family regulator genes which are not associated with a *luxI*

339 gene are present in the genome of the bacteria. Mutants of one of them (*Dda3937_00840*) are

340 overrepresented in the chicory. Its product is probably a repressor of genes conferring an

341 increased fitness *in planta*.

343 **Validation of the Tn-seq results**.

344 To validate the Tn-seq results, we performed coinoculation experiments in chicory leaves

345 with the wild type strain and various mutants in GA genes (*gcpA* and *rsmC*) or GD genes

346 (*hdfR*, *clpSA*, *metB*, *flhDC*, *purF*, *cysJ*, *degQ*, *pyrE*, *carA*, *leuA*, *guaB*, *purL*, *lysA*) in a 1/1

347 ratio. We calculated a competitive index (CI) by counting the number of each type of bacteria

348 in the rotten tissue after 24 h. We confirmed the ability of the Δ*rsmC* and Δ*gcpA* to overgrow

349 the wild type strain. At the opposite, the wild type strain overgrew the in frame deletion

350 mutants that were tested. The lowest competitive index were observed with mutants in

351 biosynthetic pathways such as Δ*leuA*, Δ*guaB*, Δ*purL*, Δ*lysA*.

352 Amino acid auxotroph mutants (Cys⁻, Leu⁻, Met⁻ and Lys⁻) tested by coinoculation

353 experiments could be phenotypically complemented *in planta*. Addition of both the non-

354 synthetized amino acid and the auxotroph mutant within the wound totally or almost

355 completely suppressed the growth defect of the auxotroph mutant *in planta* (Fig. S4). This

356 confirmed the low availability of certain amino acids in chicory. This result confirmed that

357 Tn-seq is a reliable technique to identify genes involved in plant colonization and virulence.

359 **Conclusion**

360 This Tn-seq experiment highlights new factors required for *D. dadantii* successful rotting of

361 chicory. Many genes known to be important for pathogenesis were not found in this screen

362 because their products are secreted and can be shared with other strains in the community.

363 This includes all the proteins secreted by the type II secretion system and small molecules

such as siderophores and butanediol. Other categories of genes were not found: for example,

no genes involved in response to acidic or oxidative stresses were identified. Chicory has

been described as an inadequate model to study the response of *D. dadantii* to oxidative stress

[60]. Similarly, the type III *hrp* genes were not identified in our study. The Hrp system is not

necessary for *D. dadantii* virulence and in our experimental conditions (high inoculum on

isolated chicory leaves) the necrotrophic capacities of *D. dadantii* (production of plant cell

wall degrading enzymes) is probably sufficient to provoke the disease. Our results also

uncover some unknown aspects of the infection process. Struggle for iron availability

between plant and bacterial pathogens has been well described. However, a competition for

amino acids and nucleic acid seems also to take place in the plant. The amount of nucleic

acids and of the cysteine, leucine, methionine, threonine and isoleucine amino acids is too

low in chicory to allow an efficient multiplication of bacteria defective in their biosynthesis.

Some enzymatic steps involved in their synthesis are specific to bacteria and fungi. Thus,

they could be good targets for the development of specific inhibitors [61] to fight *D. dadantii*.

Regulation of *D. dadantii* virulence has been extensively studied [2, 21]. However, new

regulatory genes controlling virulence were also detected in this study. An orphan LuxR

family regulator seems to play an important role in virulence. New members of the FlhDC

regulation pathway were also detected. A few genes of unknown function remain to be

studied.

D. dadantii can infect dozens of plants. Besides chicory, *D. dadantii* virulence tests are

usually performed on potato plant, tuber or slices, *Arabidopsis thaliana*, saintpaulia or celery.

Metabolic status or reaction defenses of these model plants are all different and bacterial

genes required for a successful infection will probably differ in each model. Testing several

of them will allow to determine the full virulence repertoire of the bacteria.

While Tn-seq has been used to study genes required for the infection of animals, there has

been no genome-wide study of factors necessary for a necrotrophic plant pathogen to develop

and provoke disease on a plant. Besides the genes of known function described in the Result

section, this study allowed the identification of several genes of unknown function required

for chicory rotting. Repetition of this experiment with other strains or on other plants will tell

if these genes encode strain or host specific virulence factors.

Methods

Bacterial strains and growth conditions. Bacterial strains, phages, plasmids and

oligonucleotides used in this study are described in Table S1 to Table S3. *D. dadantii* and *E.*

coli cells were grown at 30 and 37°C respectively in LB medium or M63 minimal medium

supplemented with a carbon source (2 g/L). When required antibiotics were added at the

following concentration: ampicillin, 100 µg/L, kanamycin and chloramphenicol, 25 µg/L.

Media were solidified with 1.5 g/L agar. Transduction with phage PhiEC2 was performed

according to [62]. The motility of each mutant was compared with that of the wild-type strain

on semisolid (0.4%) LB agar plates as previously described [63].

Construction of the transposon library

Five mL of an overnight culture of *D. dadantii* strain A350 and of *E. coli* MFDpir/pSamEC

were mixed and centrifuged 2 min at 6000 g. The bacteria were resuspended in 1 mL of M63

medium and spread onto a 0.45 µm cellulose acetate filter placed on a M63 medium agar

plate. After 8h, bacteria were resuspended in 1 mL M63 medium. An aliquot was diluted and

spread onto LB agar + kanamycin plates to estimate the efficiency of mutagenesis. The other

part was inoculated in 100 mL of LB medium + kanamycin and grown for 24 h at 30°C. To

confirm that the bacteria that grew were *D. dadantii* strains with a transposon but without

plasmid pSamEC, we checked that all the grown bacteria were kan^R, amp^S and

diaminopimelate (DAP) prototrophs (MFDpir is DAP$^-$). The bacteria were frozen in 40%

glycerol at -80°C and represent a library of about 300 000 mutants.

DNA preparation for high-throughput sequencing

An aliquot of the mutant library was grown overnight in LB medium + kanamycin. To

identify essential genes in LB, the culture was diluted 100-fold in LB and grown for 6 h. To

infect chicory, the overnight culture was centrifuged and resuspended at OD_{600} = 1 in M63

medium. Chicories cut in half were inoculated with 10 µL of this bacterial suspension and

incubated at 30°C with maximum moist. After 60 h, the rotten tissue was collected and

filtered through a cheesecloth. The bacteria were collected by centrifugation and washed

twice in M63 medium. DNA was extracted from 1.5 mL aliquots of bacterial suspension

adjusted to OD_{600}1.5 with the Promega Wizard Genomic DNA purification kit. Next steps of

the DNA preparation methods were adapted from [26]. All DNA gel-extraction were

performed onto a blue-light transilluminator of DNA stained with gel-green (Biotium) to

avoid DNA mutation and double-stranded breaks. 50 µg of DNA samples were digested with

50 U MmeI in a total volume of 1.2 mL for one hour at 37°C according to manufacturer's

instructions, then heat-inactivated for 20 minutes at 80°C, purified (QIAquick, PCR

purification kit Qiagen) and concentrated using a vacuum concentrator to a final volume of

25 µL. Digested DNA samples were run on a 1% agarose gel, the 1.0–1.5 kb band containing

the transposon and adjacent DNA was cut out and DNA was extracted from the gel according

to manufacturer's instructions (Qiaquik Gel Extraction Kit, Qiagen). This allowed recovery

of all the fragments containing genomic DNA adjacent to transposons (1201 bp of

transposable element with 32-34 bp of genomic DNA). A pair of single-stranded

complementary oligonucleotides containing an unique 5-nt barcode sequence (LIB_AdaptT

and LIB_AdaptB) was mixed and heated to 100°C, then slowly cooled down in a water bath

to obtain double-stranded adaptors with two-nucleotide overhangs. 1 µg DNA of each sample

was ligated to the barcoded adaptors (0.44 mM) with 2000 U T4 DNA ligase in a final

volume of 50 µL at 16°C overnight. Five identical PCR reactions from the ligation product

were performed to amplify the transposon adjacent DNA. One reaction contained 100 ng of

DNA, 1 unit of Q5 DNA polymerase (Biolabs), 1X Q5 Buffer, 0.2 mM dNTPs, 0.4 µM of the

forward primer (LIB_PCR_5, which anneals to the P7 Illumina sequence of the transposon)

and the reverse primer (LIB_PCR_3, which anneals to the P5 adaptor). Only 18 cycles were

performed to keep a proportional amplification of the DNA. Samples were concentrated

using a vacuum concentrator to a final volume of 25 µL. Amplified DNA was run on a 1.8%

agarose gel and the 125 bp band was cut-out and gel extracted (QIAquick, PCR purification

kit Qiagen). DNA was finally dialysed (MF-Millipore™ Membrane Filters) for 4 hours.

Quality control of the Tn-seq DNA libraries (size of the fragments and concentration) and

High-throughput sequencing on HiSeq 2500 (Illumina) was performed by MGX (CNRS

sequencing service, Montpellier). 6 DNA libraries were multiplexed on one flow-cell. After

demultiplexing, the total number of reads was comprised between 18 and 31 millions (Table

1).

Bioinformatics analysis:

Raw reads from the fastQ files were first filtered using cutadapt v1.11 [64] and only reads

containing the *mariner* inverted left repeat (ACAGGTTGGATGATAAGTCCCCGGTCTT)

were trimmed and considered *bona fide* transposon-disrupted genes. Trimmed reads were

then analyzed using a modified version of the TPP script available from the TRANSIT

software v2.0.2 [32]. The mapping step was modified to select only reads mapping uniquely

and without mismatch in the *D. dadantii* 3937 genome (Genbank CP002038.1). Then, the

counting step was modified to accurately count the reads mapping to each TA site in the

reference genome according to the Tn-seq protocol used in this study. Read counts per

insertion were normalized using the LOESS method as described in [65]. We next used the

TRANSIT software (version 2.0) to compare the Tn-seq datasets.

Strain construction. To construct the A4277 strain, gene Dda3937_03424 was amplified

with the oligonucleotides 19732+ and 19732-. The resulting fragment was inserted into the

pGEM-T plasmid (Promega). A *uidA*-kan^R cassette [66] was inserted into the unique AgeI

site of the fragment. The construct was recombined into the *D. dadantii* chromosome

according to [67]. Recombination was checked by PCR. To construct the in-frame deletion

mutants, the counter-selection method using the *sacB* gene was used [68]. The suicide

pRE112 plasmid containing 500 bp of upstream and downstream DNA of the gene to delete

was transferred by conjugation from the *E. coli MFDpir* strain into *D. dadantii* 3937.

Selection of the first event of recombination was performed on LB agar supplemented with

chloramphenicol at 30 µg/L. Transconjugants were then spread on LB agar without NaCl

supplemented with 5 % sucrose to allow the second event of recombination. In-frame

deletions were then checked by auxotrophy analysis and/or by PCR (Dreamtaq polymerase,

Thermofisher). In order to discriminate mutants from the wild strain during coinoculation

experiments, a Gm^R derivative of the WT strain was constructed by insertion of the mini-

Tn*7*-Gm into the *att*Tn7 site (close to the *glmS* gene) [69]. A 3937 Gm^R strain was made by

coelectroporation of pTn7-M [69] and pTnS3 [70] plasmids into *D. dadantii* 3937 strain. The

mini-Tn*7*-Gm delivered by the pTn7-M vector (suicide plasmid in *D. dadantii*) is inserted

into the *att*Tn7 site (close to the *glmS* gene) of recipient strain thanks to pTnS3 plasmid

encoding the Tn*7* site-specific transposition pathway. The Gm^R strain obtained was then

checked by PCR using attTn7-Dickeya3937-verif and 3-Tn7L primers (Table S3).

Protein techniques. Flagella were prepared from overnight LB grown cells. Bacteria

were pelleted, resuspended in 1/10 volume of water and passed 20 fold through a needle

on a syringe. Cells and cells debris were removed by centrifugation 5 min at 20 000 x g

[63]. Proteins were analyzed by SDS-polyacrylamide gel electrophoresis (SDS-PAGE).

Celery inoculation experiments. Wild Type and A4277 (glycosylation) mutant were grown

overnight in M63 + glycerol medium. Bacteria were washed in M63 medium and the OD_{600}

was adjusted to 1.0. Bacteria were diluted 10-fold in the same medium. 10 µL of the bacterial

suspension were inoculated into leaves in a hole made with a pipet tip. The wound was

covered with mineral oil and the leaves were incubated at 30°C at high humidity for 2 days

(celery). Length or rotten tissue was measured.

Coinoculation experiments. To determine the competitive index of the mutants, the wild

type strain and the mutant to test were grown overnight in M63 + glycerol medium. Bacteria

were washed in M63 medium and the OD_{600} was adjusted to 1.0. Bacteria were mixed to a

1:1 ratio and diluted 10-fold. For complementation experiments *in planta*, the dilution was

performed in M63 medium with 1mM of required amino acid. 10 µL of the mixture were

inoculated into chicory leaves. The wound was covered with mineral oil and the leaves were

incubated at 30 °C at high humidity. After 24 h the rotten tissue was collected, homogenized,

diluted in M63 and spread onto LB and LB + antibiotic plates. After 48 h at 30°C, colonies

were counted. The competitive index is the ratio (number of mutant bacteria/number of WT

bacteria) in the rotten tissue / (number of mutant bacteria/number of WT bacteria) in the

inoculum. For the genes whose absence confers a growth advantage in chicory according to

the Tn-seq experiment, in frame deletions were realized in a WT strain. The other mutants

were realized in the 3937 Gm[R] strain.

Nucleotide sequence accession numbers. The transposon sequence reads we obtained have

been submitted to the ENA database under accession number PRJEB20574.

Acknowledgments

We thank Geraldine Effantin, Veronique Utzinger, Matthias Schulz, Andrea Flipo, Leana

Corneloup and Barbara Gbaguidi for technical assistance, the members of the MTSB team

and Xavier Charpentier for discussion, Nicole Cotte-Pattat and Sarah Bigot for critical read of

the manuscript and James Paslesvert for encouragements.

References

1.   Adeolu M, Alnajar S, Naushad S, R SG. Genome-based phylogeny and taxonomy of

the 'Enterobacteriales': proposal for Enterobacterales ord. nov. divided into the families

Enterobacteriaceae, Erwiniaceae fam. nov., Pectobacteriaceae fam. nov., Yersiniaceae fam.

nov., Hafniaceae fam. nov., Morganellaceae fam. nov., and Budviciaceae fam. nov. Int J Syst

Evol Microbiol. 2016;66(12):5575-99. Epub 2016/09/14. doi: 10.1099/ijsem.0.001485.

PubMed PMID: 27620848.

2.   Charkowski A, Blanco C, Condemine G, Expert D, Franza T, Hayes C, *et al*. The

Role of Secretion Systems and Small Molecules in Soft-Rot Enterobacteriaceae

Pathogenicity. Annu Rev Phytopathol. 2012;50:425-49. PubMed PMID: 22702350.

3.   Hugouvieux-Cotte-Pattat N, Condemine G, Nasser W, Reverchon S. Regulation of

pectinolysis in *Erwinia chrysanthemi.* Annu Rev Microbiol. 1996;50:213-57. PubMed PMID:

8905080.

4.   Condemine G, Robert-Baudouy J. Analysis of an *Erwinia chrysanthemi* gene cluster

involved in pectin degradation. Mol Microbiol. 1991;5(9):2191-202. PubMed PMID:

1766386.

5.   Reverchon S, Nasser W, Robert-Baudouy J. *pecS*: a locus controlling pectinase,

cellulase and blue pigment production in *Erwinia chrysanthemi*. Mol Microbiol.

1994;11(6):1127-39. Epub 1994/03/01. PubMed PMID: 8022282.

6.   Surgey N, Robert-Baudouy J, Condemine G. The *Erwinia chrysanthemi pecT* gene

regulates pectinase gene expression. J Bacteriol. 1996;178(6):1593-9. PubMed PMID:

8626286.

7.   Nasser W, Faelen M, Hugouvieux-Cotte-Pattat N, Reverchon S. Role of the nucleoid-

associated protein H-NS in the synthesis of virulence factors in the phytopathogenic

bacterium *Erwinia chrysanthemi*. Mol Plant Microbe Interact. 2001;14(1):10-20. Epub

2001/02/24. doi: 10.1094/mpmi.2001.14.1.10. PubMed PMID: 11194867.

8.   Lebeau A, Reverchon S, Gaubert S, Kraepiel Y, Simond-Cote E, Nasser W, *et al*. The

GacA global regulator is required for the appropriate expression of *Erwinia chrysanthemi*

3937 pathogenicity genes during plant infection. Environ Microbiol. 2008;10(3):545-59.

PubMed PMID: 18177376.

9.   Condemine G, Castillo A, Passeri F, Enard C. The PecT repressor coregulates

synthesis of exopolysaccharides and virulence factors in *Erwinia chrysanthemi*. Mol Plant

Microbe Interact. 1999;12(1):45-52. PubMed PMID: 9885192.

10.   Reverchon S, Rouanet C, Expert D, Nasser W. Characterization of indigoidine

biosynthetic genes in *Erwinia chrysanthemi* and role of this blue pigment in pathogenicity. J

Bacteriol. 2002;184(3):654-65. Epub 2002/01/16. PubMed PMID: 11790734; PubMed

Central PMCID: PMCPMC139515.

11.   Condemine G, Dorel C, Hugouvieux-Cotte-Pattat N, Robert-Baudouy J. Some of the

out genes involved in the secretion of pectate lyases in *Erwinia chrysanthemi* are regulated by

kdgR. Mol Microbiol. 1992;6(21):3199-211. PubMed PMID: 1453958.

12.   Yang CH, Gavilanes-Ruiz M, Okinaka Y, Vedel R, Berthuy I, Boccara M, *et al. hrp*

genes of *Erwinia chrysanthemi* 3937 are important virulence factors. Mol Plant Microbe

Interact. 2002;15(5):472-80. Epub 2002/05/31. doi: 10.1094/mpmi.2002.15.5.472. PubMed

PMID: 12036278.

13.   Li Y, Hutchins W, Wu X, Liang C, Zhang C, Yuan X, et al. Derivative of plant

phenolic compound inhibits the type III secretion system of *Dickeya dadantii* via HrpX/HrpY

two-component signal transduction and Rsm systems. Molecular plant pathology.

2015;16(2):150-63. Epub 2014/07/06. doi: 10.1111/mpp.12168. PubMed PMID: 24986378.

14.   Yang S, Peng Q, Zhang Q, Zou L, Li Y, Robert C, *et al.* Genome-wide identification

of HrpL-regulated genes in the necrotrophic phytopathogen *Dickeya dadantii* 3937. PLoS

One. 2010;5(10):e13472. Epub 2010/10/27. doi: 10.1371/journal.pone.0013472. PubMed

PMID: 20976052; PubMed Central PMCID: PMCPMC2957411.

15.   Franza T, Mahe B, Expert D. *Erwinia chrysanthemi* requires a second iron transport

route dependent of the siderophore achromobactin for extracellular growth and plant

infection. Mol Microbiol. 2005;55(1):261-75. Epub 2004/12/23. doi: 10.1111/j.1365-

2958.2004.04383.x. PubMed PMID: 15612933.

16.   Franza T, Sauvage C, Expert D. Iron regulation and pathogenicity in *Erwinia*

chrysanthemi 3937: role of the Fur repressor protein. Mol Plant Microbe Interact.

1999;12(2):119-28. Epub 1999/02/02. doi: 10.1094/mpmi.1999.12.2.119. PubMed PMID:

9926414.

17.   Franza T, Expert D. The virulence-associated chrysobactin iron uptake system of

Erwinia chrysanthemi 3937 involves an operon encoding transport and biosynthetic

functions. J Bacteriol. 1991;173(21):6874-81. Epub 1991/11/01. PubMed PMID: 1657869;

PubMed Central PMCID: PMCPMC209040.

18.   Okinaka Y, Yang CH, Perna NT, Keen NT. Microarray profiling of *Erwinia*

chrysanthemi 3937 genes that are regulated during plant infection. Mol Plant Microbe

Interact. 2002;15(7):619-29. Epub 2002/07/18. doi: 10.1094/mpmi.2002.15.7.619. PubMed

PMID: 12118877.

19.   Yang S, Perna NT, Cooksey DA, Okinaka Y, Lindow SE, Ibekwe AM, *et al.*

Genome-wide identification of plant-upregulated genes of *Erwinia chrysanthemi* 3937 using

a GFP-based IVET leaf array. Mol Plant Microbe Interact. 2004;17(9):999-1008. Epub

2004/09/24. doi: 10.1094/mpmi.2004.17.9.999. PubMed PMID: 15384490.

20.   Chapelle E, Alunni B, Malfatti P, Solier L, Pedron J, Kraepiel Y, *et al.* A

straightforward and reliable method for bacterial *in planta* transcriptomics: application to the

Dickeya dadantii/Arabidopsis thaliana pathosystem. Plant J. 2015;82(2):352-62. Epub

2015/03/06. doi: 10.1111/tpj.12812. PubMed PMID: 25740271.

21.   Reverchon S, Muskhelisvili G, Nasser W. Virulence Program of a Bacterial Plant

Pathogen: The *Dickeya* Model. Progress in molecular biology and translational science.

2016;142:51-92. Epub 2016/08/31. doi: 10.1016/bs.pmbts.2016.05.005. PubMed PMID:

27571692.

22.   Hinton JC, Sidebotham JM, Hyman LJ, Perombelon MC, Salmond GP. Isolation and

characterisation of transposon-induced mutants of *Erwinia carotovora* subsp. *atroseptica*

exhibiting reduced virulence. Mol Gen Genet. 1989;217(1):141-8. Epub 1989/05/01. PubMed

PMID: 2549365.

23.   Pirhonen M, Saarilahti H, Karlsson M-B, Palva ET. Identification of pathogenicity

determinants of *Erwinia carotovora* subsp. *carotovora* by transposon mutagenesis. Molecular

Plant-Microbe Interaction. 1991;4:276-83.

24.   Lee DH, Lim JA, Lee J, Roh E, Jung K, Choi M, *et al*. Characterization of genes

required for the pathogenicity of *Pectobacterium carotovorum* subsp. *carotovorum* Pcc21 in

Chinese cabbage. Microbiology. 2013;159(Pt 7):1487-96. Epub 2013/05/17. doi:

10.1099/mic.0.067280-0. PubMed PMID: 23676432; PubMed Central PMCID:

PMCPMC3749726.

25.   Fu Y, Waldor MK, Mekalanos JJ. Tn-Seq analysis of *Vibrio cholerae* intestinal

colonization reveals a role for T6SS-mediated antibacterial activity in the host. Cell host &

microbe. 2013;14(6):652-63. Epub 2013/12/18. doi: 10.1016/j.chom.2013.11.001. PubMed

PMID: 24331463; PubMed Central PMCID: PMCPMC3951154.

26.   Skurnik D, Roux D, Aschard H, Cattoir V, Yoder-Himes D, Lory S, *et al*. A

comprehensive analysis of *in vitro* and *in vivo* genetic fitness of *Pseudomonas aeruginosa*

using high-throughput sequencing of transposon libraries. PLoS Pathog. 2013;9(9):e1003582.

Epub 2013/09/17. doi: 10.1371/journal.ppat.1003582. PubMed PMID: 24039572; PubMed

Central PMCID: PMCPMC3764216.

27.   van Opijnen T, Camilli A. A fine scale phenotype-genotype virulence map of a

bacterial pathogen. Genome research. 2012;22(12):2541-51. Epub 2012/07/25. doi:

10.1101/gr.137430.112. PubMed PMID: 22826510; PubMed Central PMCID:

PMCPMC3514683.

28.   Cole BJ, Feltcher ME, Waters RJ, Wetmore KM, Mucyn TS, Ryan EM, *et al.*

Genome-wide identification of bacterial plant colonization genes. PLoS Biol.

2017;15(9):e2002860. Epub 2017/09/25. doi: 10.1371/journal.pbio.2002860. PubMed PMID:

28938018; PubMed Central PMCID: PMCPMC5627942.

29.   Duong DA, Jensen RV, Stevens AM. Discovery of *Pantoea stewartii* subsp. *stewartii*

Genes Important for Survival in Corn Xylem through a Tn-Seq Analysis. Molecular plant

pathology. 2018. Epub 2018/02/27. doi: 10.1111/mpp.12669. PubMed PMID: 29480976.

30.   van Opijnen T, Camilli A. Transposon insertion sequencing: a new tool for systems-

level analysis of microorganisms. Nat Rev Microbiol. 2013;11(7):435-42. Epub 2013/05/29.

doi: 10.1038/nrmicro3033. PubMed PMID: 23712350; PubMed Central PMCID:

PMCPMC3842022.

31.   Wiles TJ, Norton JP, Russell CW, Dalley BK, Fischer KF, Mulvey MA. Combining

quantitative genetic footprinting and trait enrichment analysis to identify fitness determinants

of a bacterial pathogen. PLoS Genet. 2013;9(8):e1003716. Epub 2013/08/31. doi:

10.1371/journal.pgen.1003716. PubMed PMID: 23990803; PubMed Central PMCID:

PMCPMC3749937.

32.   DeJesus MA, Ambadipudi C, Baker R, Sassetti C, Ioerger TR. TRANSIT--A

Software Tool for Himar1 TnSeq Analysis. PLoS computational biology.

2015;11(10):e1004401. Epub 2015/10/09. doi: 10.1371/journal.pcbi.1004401. PubMed

PMID: 26447887; PubMed Central PMCID: PMCPMC4598096.

33.   Goodall ECA, Robinson A, Johnston IG, Jabbari S, Turner KA, Cunningham AF, *et*

al. The Essential Genome of *Escherichia coli* K-12. MBio. 2018;9(1). Epub 2018/02/22. doi:

10.1128/mBio.02096-17. PubMed PMID: 29463657.

34.   Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M. KEGG: Kyoto

Encyclopedia of Genes and Genomes. Nucleic Acids Res. 1999;27(1):29-34. Epub

1998/12/10. PubMed PMID: 9847135; PubMed Central PMCID: PMCPMC148090.

35.   Reverchon S, Nasser W. *Dickeya* ecology, environment sensing and regulation of

virulence programme. Environ Microbiol Rep. 2013;5(5):622-36. PubMed PMID: 24115612.

36.   Ravirala RS, Barabote RD, Wheeler DM, Reverchon S, Tatum O, Malouf J, *et al*.

Efflux pump gene expression in *Erwinia chrysanthemi* is induced by exposure to phenolic

acids. Mol Plant Microbe Interact. 2007;20(3):313-20. Epub 2007/03/24. doi: 10.1094/mpmi-

20-3-0313. PubMed PMID: 17378434.

37.   Malinverni JC, Silhavy TJ. An ABC transport system that maintains lipid asymmetry

in the gram-negative outer membrane. Proc Natl Acad Sci U S A. 2009;106(19):8009-14.

Epub 2009/04/23. doi: 10.1073/pnas.0903229106. PubMed PMID: 19383799; PubMed

Central PMCID: PMCPMC2683108.

38.   Page F, Altabe S, Hugouvieux-Cotte-Pattat N, Lacroix JM, Robert-Baudouy J, Bohin

JP. Osmoregulated periplasmic glucan synthesis is required for *Erwinia chrysanthemi*

pathogenicity. J Bacteriol. 2001;183(10):3134-41. PubMed PMID: 11325942.

39.   Bouchart F, Boussemart G, Prouvost AF, Cogez V, Madec E, Vidal O, *et al*. The

virulence of a *Dickeya dadantii* 3937 mutant devoid of osmoregulated periplasmic glucans is

restored by inactivation of the RcsCD-RcsB phosphorelay. J Bacteriol. 2010;192(13):3484-

90. Epub 2010/04/27. doi: 10.1128/jb.00143-10. PubMed PMID: 20418397; PubMed Central

PMCID: PMCPMC2897653.

40.   Madec E, Bontemps-Gallo S, Lacroix JM. Increased phosphorylation of the RcsB

regulator of the RcsCDB phosphorelay in strains of *Dickeya dadantii* devoid of

osmoregulated periplasmic glucans revealed by Phos-tag gel analysis. Microbiology.

2014;160(Pt 12):2763-70. Epub 2014/10/17. doi: 10.1099/mic.0.081273-0. PubMed PMID:

25320363.

41.   Wu X, Zeng Q, Koestler BJ, Waters CM, Sundin GW, Hutchins W, *et al*. Deciphering

the components that coordinately regulate virulence factors of the soft rot pathogen *Dickeya*

dadantii. Mol Plant Microbe Interact. 2014;27(10):1119-31. Epub 2014/09/03. doi:

10.1094/mpmi-01-14-0026-r. PubMed PMID: 25180688.

42.   Franza T, Expert D. Role of iron homeostasis in the virulence of phytopathogenic

bacteria: an 'a la carte' menu. Molecular plant pathology. 2013;14(4):429-38. Epub

2012/11/23. doi: 10.1111/mpp.12007. PubMed PMID: 23171271.

43.   Tans-Kersten J, Brown D, Allen C. Swimming motility, a virulence trait of *Ralstonia*

solanacearum, is regulated by FlhDC and the plant host environment. Mol Plant Microbe

Interact. 2004;17(6):686-95. Epub 2004/06/16. doi: 10.1094/mpmi.2004.17.6.686. PubMed

PMID: 15195951.

44.   Jozwick AK, Graf J, Welch TJ. The flagellar master operon *flhDC* is a pleiotropic

regulator involved in motility and virulence of the fish pathogen *Yersinia ruckeri*. J Appl

Microbiol. 2016. Epub 2016/12/17. doi: 10.1111/jam.13374. PubMed PMID: 27981729.

45.   Xu T, Su Y, Xu Y, He Y, Wang B, Dong X, et al. Mutations of flagellar genes *fliC12*,

fliA and *flhDC* of *Edwardsiella tarda* attenuated bacterial motility, biofilm formation and

virulence to fish. J Appl Microbiol. 2014;116(2):236-44. Epub 2013/10/15. doi:

10.1111/jam.12357. PubMed PMID: 24118854.

46.   Yuan X, Khokhani D, Wu X, Yang F, Biener G, Koestler BJ, *et al.* Cross-talk

between a regulatory small RNA, cyclic-di-GMP signalling and flagellar regulator FlhDC for

virulence and bacterial behaviours. Environ Microbiol. 2015;17(11):4745-63. Epub

2015/10/16. doi: 10.1111/1462-2920.13029. PubMed PMID: 26462993.

47.   Chatterjee A, Cui Y, Chatterjee AK. RsmC of *Erwinia carotovora* subsp. *carotovora*

negatively controls motility, extracellular protein production, and virulence by binding FlhD

and modulating transcriptional activity of the master regulator, FlhDC. J Bacteriol.

2009;191(14):4582-93. Epub 2009/05/19. doi: 10.1128/jb.00154-09. PubMed PMID:

19447906; PubMed Central PMCID: PMCPMC2704716.

48.   Ko M, Park C. H-NS-Dependent regulation of flagellar synthesis is mediated by a

LysR family protein. J Bacteriol. 2000;182(16):4670-2. Epub 2000/07/27. PubMed PMID:

10913108; PubMed Central PMCID: PMCPMC94646.

49.   Yi X, Yamazaki A, Biddle E, Zeng Q, Yang CH. Genetic analysis of two

phosphodiesterases reveals cyclic diguanylate regulation of virulence factors in *Dickeya*

dadantii. Mol Microbiol. 2010;77(3):787-800. Epub 2010/06/30. doi: 10.1111/j.1365-

2958.2010.07246.x. PubMed PMID: 20584146.

50.   Yuan X, Tian F, He C, Severin GB, Waters CM, Zeng Q, et al. The diguanylate

cyclase GcpA inhibits the production of pectate lyases via the H-NS protein and RsmB

regulatory RNA in *Dickeya dadantii*. Molecular plant pathology. 2018. Epub 2018/02/02.

doi: 10.1111/mpp.12665. PubMed PMID: 29390166.

51.   Van Gijsegem F, Wlodarczyk A, Cornu A, Reverchon S, Hugouvieux-Cotte-Pattat N.

Analysis of the LacI family regulators of *Erwinia chrysanthemi* 3937, involvement in the

bacterial phytopathogenicity. Mol Plant Microbe Interact. 2008;21(11):1471-81. PubMed

PMID: 18842096.

52.   Wolfe AJ. The acetate switch. Microbiol Mol Biol Rev. 2005;69(1):12-50. Epub

2005/03/10. doi: 10.1128/mmbr.69.1.12-50.2005. PubMed PMID: 15755952; PubMed

Central PMCID: PMCPMC1082793.

53.   Antunez-Lamas M, Cabrera-Ordonez E, Lopez-Solanilla E, Raposo R, Trelles-Salazar

O, Rodriguez-Moreno A, *et al*. Role of motility and chemotaxis in the pathogenesis of

Dickeya dadantii 3937 (ex *Erwinia chrysanthemi* 3937). Microbiology. 2009;155(Pt 2):434-

42. Epub 2009/02/10. doi: 10.1099/mic.0.022244-0. PubMed PMID: 19202091.

54.   Rio-Alvarez I, Munoz-Gomez C, Navas-Vasquez M, Martinez-Garcia PM, Antunez-

Lamas M, Rodriguez-Palenzuela P, *et al*. Role of *Dickeya dadantii* 3937 chemoreceptors in

the entry to *Arabidopsis* leaves through wounds. Molecular plant pathology. 2015;16(7):685-

98. Epub 2014/12/10. doi: 10.1111/mpp.12227. PubMed PMID: 25487519.

55.   Jahn CE, Willis DK, Charkowski AO. The flagellar sigma factor *fliA* is required for

Dickeya dadantii virulence. Mol Plant Microbe Interact. 2008;21(11):1431-42. Epub

2008/10/10. doi: 10.1094/mpmi-21-11-1431. PubMed PMID: 18842093.

56.   Jiang X, Zghidi-Abouzid O, Oger-Desfeux C, Hommais F, Greliche N,

Muskhelishvili G, *et al.* Global transcriptional response of *Dickeya dadantii* to environmental

stimuli relevant to the plant infection. Environ Microbiol. 2016;18(11):3651-72. Epub

2016/03/05. doi: 10.1111/1462-2920.13267. PubMed PMID: 26940633.

57.   Taguchi F, Yamamoto M, Ohnishi-Kameyama M, Iwaki M, Yoshida M, Ishii T, *et al*.

Defects in flagellin glycosylation affect the virulence of *Pseudomonas syringae* pv. *tabaci*

6605. Microbiology. 2010;156(Pt 1):72-80. Epub 2009/10/10. doi: 10.1099/mic.0.030700-0.

PubMed PMID: 19815579.

58.   Khodai-Kalaki M, Andrade A, Fathy Mohamed Y, Valvano MA. *Burkholderia*

cenocepacia Lipopolysaccharide Modification and Flagellin Glycosylation Affect Virulence

but Not Innate Immune Recognition in Plants. MBio. 2015;6(3):e00679. Epub 2015/06/06.

doi: 10.1128/mBio.00679-15. PubMed PMID: 26045541; PubMed Central PMCID:

PMCPMC4462625.

59.   Reverchon S, Bouillant ML, Salmond G, Nasser W. Integration of the quorum-

sensing system in the regulatory networks controlling virulence factor synthesis in *Erwinia*

chrysanthemi. Mol Microbiol. 1998;29(6):1407-18. Epub 1998/10/22. PubMed PMID:

9781878.

60.   Santos R, Franza T, Laporte ML, Sauvage C, Touati D, Expert D. Essential role of

superoxide dismutase on the pathogenicity of *Erwinia chrysanthemi* strain 3937. Mol Plant

Microbe Interact. 2001;14(6):758-67. Epub 2001/06/02. doi: 10.1094/mpmi.2001.14.6.758.

PubMed PMID: 11386371.

61.   Thangavelu B, Bhansali P, Viola RE. Elaboration of a fragment library hit produces

potent and selective aspartate semialdehyde dehydrogenase inhibitors. Bioorg Med Chem.

2015;23(20):6622-31. Epub 2015/09/26. doi: 10.1016/j.bmc.2015.09.017. PubMed PMID:

26404410; PubMed Central PMCID: PMCPMC4601562.

62.   Résibois A, Colet M, Faelen M, Schoonejans E, Toussaint A. phiEC2, a new

generalized transducing phage of *Erwinia chrysanthemi*. Virology. 1984;137(1):102-12.

PubMed PMID: 18639822.

63.   Shevchik VE, Condemine G, Robert-Baudouy J. Characterization of DsbC, a

periplasmic protein of *Erwinia chrysanthemi* and *Escherichia coli* with disulfide isomerase

activity. Embo J. 1994;13(8):2007-12. PubMed PMID: 8168497.

64.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. EMBnetjournal. 2011;17:10-2.

65.   Zomer A, Burghout P, Bootsma HJ, Hermans PW, van Hijum SA. ESSENTIALS:

software for rapid analysis of high throughput transposon insertion sequencing data. PLoS

One. 2012;7(8):e43012. Epub 2012/08/18. doi: 10.1371/journal.pone.0043012. PubMed

PMID: 22900082; PubMed Central PMCID: PMCPMC3416827.

66.   Bardonnet N, Blanco C. Improved vectors for transcriptional signal screening in

corynebacteria. FEMS Microbiol Lett. 1991;68(1):97-102. PubMed PMID: 1722768.

67.   Roeder DL, Collmer A. Marker-exchange mutagenesis of a pectate lyase isozyme

gene in *Erwinia chrysanthemi*. J Bacteriol. 1985;164(1):51-6. PubMed PMID: 2995324.

68.   Link AJ, Phillips D, Church GM. Methods for generating precise deletions and

insertions in the genome of wild-type *Escherichia coli*: application to open reading frame

characterization. J Bacteriol. 1997;179(20):6228-37. Epub 1997/10/23. PubMed PMID:

9335267; PubMed Central PMCID: PMCPMC179534.

69.   Zobel S, Benedetti I, Eisenbach L, de Lorenzo V, Wierckx N, Blank LM. Tn*7*-Based

Device for Calibrated Heterologous Gene Expression in *Pseudomonas putida*. ACS synthetic

biology. 2015;4(12):1341-51. Epub 2015/07/03. doi: 10.1021/acssynbio.5b00058. PubMed

PMID: 26133359.

70.   Choi KH, Mima T, Casart Y, Rholl D, Kumar A, Beacham IR, et al. Genetic tools for

select-agent-compliant manipulation of *Burkholderia pseudomallei*. Appl Environ Microbiol.

2008;74(4):1064-75. Epub 2007/12/25. doi: 10.1128/aem.02430-07. PubMed PMID:

18156318; PubMed Central PMCID: PMCPMC2258562.

Legend of figures

Fig. 1. **Quality control of the Tn-seq *D. dadantii* 3937 libraries**. (A) Frequency and

distribution of transposon sequence reads across the entire *D. dadantii 3937* genome. The

localization of transposon insertions shows no bias throughout the genome of *D. dadantii*

3937. B) Density plot of $\log_2$FC (measured reads/expected reads per gene).

Fig 2. Scheme of the purine and pyrimidine biosynthesis pathways in *D. dadantii* that

produce XMP (purine metabolism) and UMP (pyrimidine metabolism) from L-

glutamine. In red are indicated the growth defect genes in chicory that pass the permutation

test (q-value ≤ 0.05). In bold are genes for which GD phenotype was tested and confirmed

with in frame deletion mutants. The $\log_2$FC of read numbers between chicory and LB for

each gene is indicated in bracket. Some genes do not pass the permutation test (in black) but

have a strong negative $\log_2$FC. PRPP: 5-phosphoribosyl-1-pyrophosphate ; GAR: 5'-

phosphoribosyl-1-glycinamide ; FGAM: 5'-phosphoribosyl-*N*-formylglycinamide ; AIR: 5'-

phosphoribosyl-5-aminoimidazole ; CAIR: 5'-phosphoribosyl-5-aminoimidazole carboxylic

acid ; SAICAR: 5'-phosphoribosyl-4-(*N*-succino-carboxamide)-5-aminoimidazole ; AICAR:

5-aminoimidazole-4-carboxamide ribonucleotide ; IMP: inosine monophosphate ; XMP:

xanthine monophosphate ; UMP: uridine monophosphate.

Fig 3. Examples of essential and important genes revealed by Tn-seq. Number of reads at each transposon location in the sample grown in LB or in chicory. Data are averaged from biological replicates and normalized as described in the method section. Three regions of the genome representative of Tn-seq results are shown, with the predicted genes represented at the bottom of each panel. Peaks represent read number at TA sites. Black arrows represent genes that pass the permutation test (q-value ≤ 0.05). Small arrows indicate the presence of promoter (A) Essentiality of leucine biosynthetic genes in chicory. (B) Insertions in the 5' region of *rsmC* generate growth advantage for the bacteria in chicory. (C) Importance of genes involved in motility for growth in chicory.

Fig 4. Modification of FliC revealed by Tn-seq analysis and SDS-PAGE. (A) Importance of 6 genes located between *fliA* and *fliC* for growth in chicory. Log_2FC are indicated in bracket. Dda3937_03425 and Dda3937_03426 are duplicated transposase genes that have been removed from the analysis. Black arrow: GD in chicory (q-value ≤ 0.05) ; white arrow: genes that do not pass the permutation test (q-value > 0.05). Small arrows indicate the presence of promoter. (B) Analysis by SDS-PAGE of FliC produced by the wild type (lane 1), the A3422 (lane 2) and A4277 (lane 3) strains. (C) Maceration of celery leaves by the Wild Type and A4277 (glycosylation) mutant. Length of rotten tissue was measured 48 h post infection. Boxplot were generated by BoxPlotR from 9 data points. The calculated median value is 109 for the WT strain, 40 for the A4277 strain. Center lines show the medians; box limits indicate the 25th and 75th percentiles as determined by R software; whiskers extend 1.5 times the interquartile range from the 25th and 75th percentiles.

Fig 5. Competitive Index (CI) of several mutant strains. CI values were determined in chicory leaves as described in Methods. Each value is the mean of 5 experiments. Center lines show the medians; box limits indicate the 25th and 75th percentiles as determined by R software; whiskers extend 1.5 times the interquartile range from the 25th and 75th percentiles, outliers are represented by dots. n = 5 sample points. Numbers above the boxes indicate the average competitive index in Log_{10}. * indicates a significant difference relative to the WT (p<0.05, Welch's t-test).

Supporting information legends

Fig S1. Biological reproducibility of the Tn-seq results. Pairs of Tn-seq assay results are compared, with the total number of reads per gene plotted. Analysis of DNA samples corresponding to two independent cultures of the mutant pool grown (A) in LB medium (correlation coefficient R = 0.72) and (B) in chicory (correlation coefficient R = 0.98). Values represent average numbers of reads per gene from the pairs of biological replicates.s

 Fig S2. Volcano plot of resampling results comparing replicates grown in chicory versus in LB. Significant hits have q < 0.05 or $-log_{10}$ q > 1.3. Growth defect (GD) or growth advantage (GA) genes are indicated by a red frame.

Fig S3. *acrAB* are essential in chicory. Number of Tn-seq reads at each insertion site in the *acrA acrB* region in samples grown in LB or in chicory. Data are averaged from biological replicates and normalized as described in Methods. *dnaX* which encodes both the tau and gamma subunits of DNA polymerase is represented by a grey arrow. *dnaX* is essential gene in LB. *acrAB* represented by grey arrows are GD in chicoy (q-value ≤ 0.05).

Fig S4. Complementation of auxotroph mutants *in planta*. Each leaf was inoculated with 10^6 bacteria. Length of rotten tissue was observed after 24h. Bacteria were injected into the

wounded leaf with or without amino acid. Center lines show the medians; box limits indicate

the 25th and 75th percentiles as determined by R software; whiskers extend 1.5 times the

interquartile range from the 25th and 75th percentiles, outliers are represented by dots. n = 5

sample points. Numbers above the boxes indicate the average competitive index in Log_{10}. *

indicates a significant difference relative to the WT ($p<0.05$, Welch's t-test). ** Indicates an

absence of significant difference relative to the WT ($p>0.05$, Welch's t-test).

Table S1: bacterial strains used in this study

Table S2: plasmids used in this study

Table S3: oligonucleotides used in this study

Table S4 : number of genes implicated in KEGG pathway

Table S5: raw data of the HMM and resampling analysis by transit

Table S6: List of genes with $log_2FC <-2$ or >2 but with q-value >0.05

Fig.1

Fig.2

PURINE METABOLISM

PYRIMIDINE METABOLISM

Fig.3

Fig. 4

Fig. 5