Defining the functional significance of intergenic transcribed regions based on heterogeneous features of phenotype genes and pseudogenes

John P. Lloyd[1], Zing Tsung-Yeh Tsai[2], Rosalie P. Sowers[3], Nicholas L. Panchy[4], Shin-Han Shiu[1,4,5]*

[1] Department of Plant Biology, Michigan State University, East Lansing, MI 48824, USA

[2] Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI 48109, USA

[3] Department of Biochemistry and Molecular Biology, Pennsylvania State University, University Park, PA 16802, USA

[4] Genetics Program, Michigan State University, East Lansing, MI 48824, USA

[5] Ecology, Evolutionary Biology, and Behavior Program, Michigan State University, East Lansing, MI 48824, USA

* To whom correspondence should be addressed.

Correspondence to:

Shin-Han Shiu

Michigan State University

E-mail: shius@msu.edu

Telephone number: 517-353-7196

Running title: Functionality of intergenic transcripts

Keywords: Intergenic transcription, ncRNAs, definition of function, molecular evolution, machine learning, data integration

ABSTRACT

With advances in transcript profiling, the presence of transcriptional activities in intergenic

regions has been well established in multiple model systems. However, whether intergenic

expression reflects transcriptional noise or the activity of novel genes remains unclear. We

identified intergenic transcribed regions (ITRs) in 15 diverse flowering plant species and found

that the amount of intergenic expression correlates with genome size, a pattern that could be

expected if intergenic expression is largely nonfunctional. To further assess the functionality of

ITRs, we first built machine learning classifiers using *Arabidopsis thaliana* as a model that can

accurately distinguish functional sequences (phenotype genes) and nonfunctional ones

(pseudogenes and unexpressed intergenic regions) by integrating 93 biochemical, evolutionary,

and sequence-structure features. Next, by applying the models genome-wide, we found that

4,427 ITRs (38%) and 796 annotated ncRNAs (44%) had features significantly similar to

benchmark protein-coding or RNA genes and thus were likely parts of functional genes.

However, ~60% of ITRs and ncRNAs were more similar to nonfunctional sequences and should

be considered transcriptional noise unless falsified with experiments. The predictive framework

established here provides not only a comprehensive look at how functional, genic sequences are

distinct from likely nonfunctional ones, but also a new way to differentiate novel genes from

genomic regions with noisy transcriptional activities.

INTRODUCTION

Advances in sequencing technology have helped to identify pervasive transcription in intergenic regions with no annotated genes. These intergenic transcripts have been found in metazoa and fungi, including *Homo sapiens* (human; ENCODE Project Consortium 2012), *Drosophila melanogaster* (Brown et al. 2014), *Caenorhabditis elegans* (Boeck et al. 2016), and *Saccharomyces cerevisiae* (Nagalakshmi et al. 2008). In plants, ~7,000 and ~15,000 intergenic transcripts have also been reported in *Arabidopsis thaliana* (Yamada et al. 2003; Stolc et al. 2005; Moghe et al. 2013; Krishnakumar et al. 2015) and *Oryza sativa* (Nobuta et al. 2007), respectively. The presence of intergenic transcripts indicates that there may be additional genes in genomes that have escaped gene finding efforts thus far. Knowledge of the complete suite of functional elements present in a genome is an important goal for large-scale functional genomics efforts and the quest to connect genotype to phenotype. Thus the identification of functional intergenic transcribed regions (ITRs) represents a fundamental task that is critical to our understanding of the gene space in a genome.

Loss-of-function study represents the gold standard by which the functional significance of genomic regions, including ITRs, can be confirmed (Niu and Jiang 2013). In *Mus musculus* (mouse), at least 25 ITRs with loss-of-function mutant phenotypes have been identified (Sauvageau et al. 2013; Lai et al. 2015), indicating that they are *bona fide* genes. In addition, loss-of-function mutants have been used to confirm ITR functionality in mouse embryonic stem cell proliferation (Ivanova et al. 2006; Guttman et al. 2009) and male reproductive development (Heinen et al. 2009), as well as brain and eye development in *Danio rerio* (Ulitsky et al. 2011). In human, 162 long intergenic non-coding RNAs harbor phenotype-associated SNPs, suggesting that these expressed intergenic regions may be functional (Ning et al. 2013). In addition to intergenic expression, most model organisms feature an abundance of annotated non-coding RNA (ncRNA) sequences (Zhao et al. 2016), which are mostly identified through the presence of expression occurring outside of annotated genes. Thus, the only difference between ITRs and most ncRNA sequences is whether or not they have been annotated. Similar to the ITR examples above, a small number of ncRNAs have been confirmed as functional through loss-of-function experimental characterization, including but not limited to *Xist* in mouse (Penny et al. 1996; Marahrens et al. 1997), *Malat1* in human (Bernard et al. 2010), *bereft* in *D. melanogaster* (Hardiman et al. 2002), and *At4* in *A. thaliana* (Shin et al. 2006). However, despite the presence

of a few notable examples, the number of ITRs and ncRNAs with well-established functions is

dwarfed by those with no known function.

While some ITRs and ncRNAs are likely novel genes, intergenic transcription may also

be the byproduct of noisy expression that can occur due to nonspecific landing of RNA

Polymerase II (RNA Pol II) or spurious regulatory signals that drive expression in random

genomic regions (Struhl 2007). Thus, whether an intergenic transcript is considered functional

cannot depend solely on the fact that it is expressed. In addition to being biochemically active,

the genomic region must be under selection. This line of logic has revived the classical ideas on

differentiating "causal role" and "selected effect" functionality (Doolittle et al. 2014). A "causal

role" definition requires a definable activity to consider a genomic region as functional

(Cummins 1975; Amundson and Lauder 1994), which is adopted by the ENCODE Consortium

(2012) to classify ~80% of the human genome as having biochemical functions. This finding has

been used as evidence to disprove the presence of junk DNA that is not under natural selection

(see Eddy 2013). This has drawn considerable critique because biochemical activity itself is not

an indication of selection (Graur et al., 2013; Niu and Jiang, 2013). Instead, selected effect

functionality is advocated to be a more suitable definition for a genomic region with discernible

activity (Amundson and Lauder 1994; Graur et al., 2013; Doolittle et al. 2014). Under the

selected effect functionality definition, ITRs and most annotated ncRNA genes remain

functionally ambiguous.

Functional ITRs represent genic sequences that have not been identified with

conventional gene finding programs. Such programs incorporate sequence characteristics,

transcriptional evidence, and conservation information to define genic regions that are expected

to be functional. Thus, genes that lack the features typically associated with genic regions remain

unidentified. Due to the debate on the definitions of function post-ENCODE, Kellis et al. (2014)

suggested that evolutionary, biochemical, and genetic evidences provide complementary

information to define functional genomic regions. Integration of chromatin accessibility,

transcriptome, and conservation evidence was successful in identifying regions in the human

genome that are under selection (Gulko et al. 2014). Moreover, a comprehensive integration of

biochemical, evolutionary, and genetic evidence resulted in highly accurate identification of

human disease genes and pseudogenes (Tsai et al. 2017). However, it is not known if such

predictions are possible or if the features that define functional genomic regions in human are

applicable in other species. In plants, even though many biochemical signatures are known to be

associated with genic regions, these signatures have not been incorporated to assist in identifying

the functional genomic regions.

112   To investigate the functionality of intergenic transcription, we first identified ITRs in 15

flowering plant species with 17-fold genome size differences and evaluated the relationship

between the prevalence of intergenic expression and genome size. Next, we determined whether

93 evolutionary, biochemical, and sequence-structure features could distinguish functional

sequences (phenotype genes) and nonfunctional ones (pseudogenes) using *A. thaliana* as a

model. We then jointly considered all 93 features to establish functional gene prediction models

using machine learning methods. Finally, we applied the models to ITRs and annotated ncRNAs

to determine whether these functionally ambiguous sequences are more similar to known

functional or likely nonfunctional sequences.

RESULTS & DISCUSSION

Relationship between genome size and intergenic expression indicates that intergenic

transcripts may generally be nonfunctional

Transcription of an unannotated, intergenic region could be due to nonfunctional transcriptional

noise or the activity of a novel gene. If noisy transcription occurs due to random landing of RNA

Pol II or spurious regulatory signals, a naïve expectation is that, as genome size increases, the

amount of intergenic expression would increase accordingly. By contrast, we expect that the

extent of expression for genic sequences will not be significantly correlated with genome size

because larger plant genomes do not necessarily have more genes (r^2=0.01; p=0.56; see

Methods). Thus, to gauge if intergenic transcribed regions (ITRs) generally behave more like

what we expect of noisy or genic transcription, we assessed the correlation between genome size

and the amount of intergenic expression occurring within 15 flowering plant species.

133   We first identified genic and intergenic transcribed regions using leaf transcriptome data

from 15 flowering plants with 17-fold differences in genome size (Supplemental Table 1).

Identical numbers of RNA-sequencing (RNA-seq) reads (30 million) and the same mapping

procedures were used in all species to facilitate cross-species comparisons (see Methods).

Transcribed regions were considered as ITRs if they did not overlap with any gene annotation

and had no significant translated sequence similarity to plant protein sequences. As expected, the

amount of expression originating from annotated genic regions had no significant correlation

with genomes size (r^2=0.03; p=0.53; **Fig. 1A**). In contrast, the amount of intergenic expression

occurring was significantly and positively correlated (r^2=0.30; p=0.04; **Fig. 1B**). Because more

intergenic expression is occurring in species with more genome space, this is consistent with the

interpretation that a significant proportion of intergenic expression represents transcriptional

noise. However, the correlation between genome size and intergenic expression explained ~30%

of the variation (**Fig. 1B**), suggesting that other factors also affect ITR content, including the

possibility that some ITRs are truly functional, novel genes. To further evaluate the functionality

of intergenic transcripts, we next identified the biochemical and evolutionary features of

functional genic regions and tested whether intergenic transcripts in *A. thaliana* were more

similar to functional or nonfunctional sequences.

Expression, conservation, and epigenetic features are significantly distinct between

benchmark functional and nonfunctional genomic sequences

To determine whether intergenic transcripts resemble functional sequences, we first asked what

features allow benchmark functional and nonfunctional genomic regions to be distinguished. For

benchmark functional sequences, we used genes with visible loss-of-function phenotypes when

mutated (referred to as phenotype genes, n=1,876; see Methods). These phenotype genes were

considered functional based on the selected effect functionality criterion (Neander 1991) because

their mutations have significant growth and/or developmental impact and likely contribute to

reduced fitness. For benchmark nonfunctional genomic regions, we utilized pseudogene

sequences (n=761; see Methods). These pseudogenes exhibit sequence similarity to known

genes, but harbor disabling mutations, including frame shifts and/or in-frame stop codons, that

result in the production of presumably nonfunctional protein products. Considering that only 2%

of pseudogenes are maintained over 90 million years of divergence between human and mouse

(Svensson et al. 2006), it is expected that the majority of pseudogenes are no longer under

selection (Li et al. 1981).

We evaluated 93 gene or gene product features for their ability to distinguish between

phenotype genes and pseudogenes. These features were grouped into seven categories, including

chromatin accessibility, DNA methylation, histone 3 (H3) marks, sequence conservation,

sequence-structure, transcription factor (TF) binding, and transcription activity. Feature values

(Supplemental Table 2) were calculated for a randomly-selected 500 base pair (bp) window

inside a phenotype gene or pseudogene. We used Area Under the Curve - Receiver Operating

Characteristic (AUC-ROC) as a metric to measure how well a feature distinguishes between

phenotype genes and pseudogenes. AUC-ROC values range between 0.5 (random guessing) and

1 (perfect separation of functional and nonfunctional sequences), with AUC-ROC values of 0.7,

0.8, and 0.9 considered fair, good, and excellent performance, respectively. Among the seven

feature categories, transcription activity features were highly informative (median AUC-

ROC=0.88; **Fig. 2A**). Sequence conservation, DNA methylation, TF binding, and H3 mark

features were also fairly distinct between phenotype genes and pseudogenes (median AUC-ROC

~ 0.7 for each category; **Fig. 2B-E**). By contrast, chromatin accessibility and sequence-structure

features were largely uninformative (median AUC-ROC=0.51 and 0.55, respectively; **Fig.**

2F,G). The poor performance of chromatin accessibility features is likely because the DNase I

hypersensitive site (DHS) datasets were sparse, as only 2-6% of phenotype gene and pseudogene

sequences overlapped a DHS peak. Further, median nucleosome occupancy of phenotype genes

(median normalized nucleosome occupancy = 1.22) is only slightly lower than that of

pseudogenes (median = 1.31; Mann Whitney U test, $p < 2e-4$). For sequence-structure features

based on dinucleotide structures (see Methods), we found that poor performance was likely due

to phenotype genes and pseudogenes sharing similar dinucleotide sequence compositions

(r^2=0.99, p<3e-16).

Error rates for functional region predictions are high when only single features are

considered

Within each feature category, there was a wide range of performance between features (**Fig. 2**,

Supplemental Table 3) and there were clear biological or technical explanations for features that

perform poorly. For the transcription activity category, 17 out of 24 features had an AUC-ROC

performance >0.8, including the best-performing feature, expression breadth (AUC-ROC=0.95;

Fig. 2A). However, five transcription activity-related features performed poorly (AUC-

ROC<0.65), including the presence of expression (transcript) evidence (AUC-ROC=0.58; **Fig.**

2A). This is because 80% of pseudogenes were considered expressed in ≥1 of 51 RNA-seq

datasets, demonstrating that presence of transcripts should not be used by itself as evidence of

functionality. For the sequence conservation category, maximum and average phastCons

conservation scores were highly distinct between phenotype genes and pseudogenes (AUC-

ROC=0.83 and 0.82, respectively; **Fig. 2B**). On the other hand, identity to best matching

nucleotide sequences found in *Brassicaceae* and algal species were not informative (AUC-

ROC=0.55 and 0.51, respectively; **Fig. 2B**). This was because 99.8% and 95% of phenotype

genes and pseudogenes, respectively, had a potentially homologous sequence within the

Brassicaceae family, and only 3% and 1%, respectively, in algal species. Thus, *Brassicaceae*

genomes were too similar and algal genomes too dissimilar to *A. thaliana* to provide meaningful

information. H3 mark features also displayed high variability. The most informative H3 mark

features were based on the number and coverage of activation-related marks (AUC-ROC=0.87

and 0.85, respectively; **Fig. 2E**), consistent with the notion that histone marks are often jointly

associated with active genomic sequences to provide a robust regulatory signal (Schreiber and

Bernstein 2002; Wang et al. 2008). By comparison, the coverage and intensity of H3 lysine 27

trimethylation (H3K27me3) and H3 threonine 3 phosphorylation (H3T3ph) were largely

indistinct between phenotype genes and pseudogenes (AUC-ROC range: 0.55-0.59; **Fig. 2E**).

Despite this high variability in performance, some features and feature categories had

high AUC-ROCs, suggesting that these features may individually provide sufficient information

for distinguishing between functional and nonfunctional genomic regions. To assess this

possibility, we next evaluated the error rates of function predictions based on single features. We

first considered expression breadth of a sequence, the best predicting feature of functionality.

Despite high AUC-ROC (0.95), the false positive rate (FPR; % of pseudogenes predicted as

phenotype genes) was 21% when only expression breadth was used, while the false negative rate

(FNR; % of phenotype genes predicted as pseudogenes) was 4%. Similarly, the best-performing

H3 mark- and sequence conservation-related features had FPRs of 26% and 32%, respectively,

and also incorrectly classified at least 10% of phenotype genes as pseudogenes. Thus, error rates

are high even when considering well-performing single features, indicating the need to jointly

consider multiple features for distinguishing phenotype genes and pseudogenes.

Consideration of multiple features in combination produces accurate predictions of

functional genomic regions

To consider multiple features in combination, we first conducted principle component (PC)

analysis to investigate how well phenotype genes and pseudogenes could be separated. Between

the first two PCs, which jointly explain 40% of the variance in the feature dataset, phenotype

genes (**Fig. 3A**) and pseudogenes (**Fig. 3B**) were distributed in largely distinct space. However,

there remains substantial overlap, indicating that standard parametric approaches are not well

suited to distinguishing between benchmark functional and nonfunctional sequences. Thus, we

instead considered all 93 features for phenotype gene and pseudogenes in combination using

random forest (referred to as the full model; see Methods). The phenotype gene and pseudogene

sequences and associated conservation, biochemical, and sequence-structure features were

separated into distinct training and testing sets and the full model was generated and validated

using independent data subsets (cross-validation). The full model provided more accurate

predictions (AUC-ROC=0.98; FNR=4%; FPR=10%; **Fig. 3C**) than any individual feature (**Fig.**

2; Supplemental Table 3). An alternative measure of performance based on the precision

(proportion of predicted functional sequences that are truly functional) and recall (proportion of

truly functional sequences predicted correctly) values among predictions generated by the full

model also indicated that the model was performing well (**Fig. 3D**). When compared to the best-

performing single feature (expression breadth), the full model had a similar FNR but half the

FPR (10% compared to 21%). Thus, the full model is highly capable of distinguishing between

phenotype genes and pseudogenes.

We next determined the relative contributions of different feature categories in predicting

phenotype genes and pseudogenes and whether models based on a subset of features would

perform similarly as the full model. Seven prediction models were established, each using only

the subset of features from a single category (**Fig. 2**). Although none of these category-specific

models had performance as high as the full model, the models based on transcription activity,

sequence conservation, and H3 mark features scored highly (AUC-ROC=0.97, 0.92, and 0.91,

respectively; **Fig. 3C**). Particularly, the transcription activity feature category model performed

almost as well as the full model (FNR=6%, FPR=12%). We emphasize that the breadth and level

of transcription are the causes of the strong performance of the transcription activity-only model,

not the presence of expression evidence.

To evaluate whether the strong performance of the full model is being driven solely by

transcription activity-related features, we also built a function prediction model did not consider

these features (full (-TX), **Fig. 3C,D**). We found that the model excluding transcription activity

features performed almost as well as the full model and similarly to the transcription activity-

feature-only model, but with an increased FPR (AUC-ROC=0.96; FNR=3%; FPR=20%). This

indicates that predictions of functional regions are not reliant solely on transcription data.

Instead, a diverse array of features can be considered to make highly accurate predictions of the

functionality of a genomic sequence. Meanwhile, our finding of the high performance of the

transcription activity-only model highlights the possibility of establishing an accurate model for

distinguishing functional genic and nonfunctional genomic sequences in plant species with only

a modest amount of transcriptome data.

Functional likelihood allows the prediction of functional and nonfunctional genomic

regions

To provide a measure of the potential functionality of any sequence in the *A. thaliana* genome,

including ITRs and ncRNAs, we utilized the confidence score from the full model as a

"functional likelihood" value (Tsai et al. 2017; see Methods). The functional likelihood (FL)

score ranges between 0 and 1, with high values indicating that a sequence is more similar to

phenotype genes (functional) and low values indicating a sequence more closely resembles

pseudogenes (nonfunctional). FL values for all genomic regions examined in this study are

available in Supplemental Table 4. As expected, phenotype genes had high FL values

(median=0.97; **Fig. 4A**) and pseudogenes had low values (median=0.01; **Fig. 4B**). To call

sequences as functional or not, we defined a threshold FL value of 0.35 (see Methods). Using

this threshold, 96% of phenotype genes (**Fig. 4A**) and 90% of pseudogenes (**Fig. 4B**) are

correctly classified as functional and nonfunctional, respectively, demonstrating that the full

model is highly capable of distinguishing functional and nonfunctional sequences.

We next applied our model to predict the functionality of annotated protein-coding genes,

transposable elements, and unexpressed intergenic regions. Most annotated protein-coding genes

not included in the phenotype gene dataset had high FL scores (median=0.86; **Fig. 4C**) and 80%

were predicted as functional. Of the 20% of protein-coding genes that were predicted as

nonfunctional, we expect that at least 4% represent false negatives based on the FNR of the full

model. The actual FNR among protein-coding genes may be higher, however, as phenotype

genes represent a highly active and well conserved subset of all genes. However, a subset of the

low-scoring protein-coding genes may also represent gene sequences undergoing functional

decay and *en route* to pseudogene status. To assess this possibility, we examined 1,940 *A.*

thaliana "decaying" genes that may be experiencing pseudogenization due to promoter

disablement (Yang et al. 2011) and found that, while these decaying genes represented only 7%

of all *A. thaliana* annotated protein-coding genes, they made up 45% of protein-coding genes

predicted as nonfunctional (Fisher's Exact Test (FET), $p < 1E\text{-}11$). In addition to protein-coding

genes, we evaluated the FLs of transposable elements (TEs) and randomly-selected, unexpressed

intergenic regions that are most likely nonfunctional. As expected, the FLs were low for both

TEs (median=0.03, **Fig. 4D**) and unexpressed intergenic regions (median=0.07; **Fig. 4E**), and

99% of TEs and all unexpressed intergenic sequences were predicted as nonfunctional, further

demonstrating the utility of the function prediction model. Overall, the FL measure provides a

useful metric to distinguish between phenotype genes and pseudogenes. In addition, the FLs of

annotated protein-coding genes, TEs, and unexpressed intergenic sequences agree with *a priori*

expectations regarding the functionality of these sequences.

302   Exclusion of features from multiple tissues increases prediction performance for narrowly-

303   expressed sequences

Although the full model performs exceedingly well, there remain false predictions. There are 76

phenotype genes (4%) predicted as nonfunctional (referred to as low-FL phenotype genes). We

assessed why these phenotype genes were not correctly identified by first asking what category

of features were particularly distinct between low-FL and the remaining phenotype genes. We

found that the major category that led to the misclassification of phenotype genes was

transcription activity, as only 7% of low-scoring phenotype genes were predicted as functional in

the transcription activity-only model, compared to 98% of high FL phenotype genes (**Fig. 5**). By

contrast, >65% of low-FL phenotype genes were predicted as functional when sequence

conservation, H3 mark, or DNA methylation features were used. This could suggest that the full

model is less effective in predicting functional sequences that are weakly or narrowly expressed.

While sequence conservation features are distinct between functional and nonfunctional

sequences when considered in combination, a significantly higher proportion of low-FL

phenotype genes were specific to the *Brassicaceae* family, with only 33% present in

dicotyledonous species outside of the *Brassicaceae*, compared to 78% of high-scoring phenotype

genes (FET, $p < 4e\text{-}12$), thus our model likely has reduced power in detecting lineage-specific

genes.

Given the association between transcription activity features and functional predictions,

we next investigated how functional predictions performed for conditionally-functional and

narrowly-expressed sequences. We found that genes with conditional phenotypes (see Methods)

had no significant differences in FLs (median=0.96) as those with phenotypes under standard

growth conditions (median=0.97; U test, p=0.38, Supplemental Fig. 1A). Thus, our model can

capture conditionally functional sequences. Next, we evaluated FL distributions among

sequences with different breadths of gene expression. For this comparison, we focused on non-

stress, single-tissue expression datasets (Supplemental Table 5), which was distinct from the

expression breadth feature in the prediction model that considered all datasets. While phenotype

genes were better predicted than pseudogenes among sequences with the same number of tissues

with expression evidence (U tests, all $p < 1.7E-06$; Supplemental Fig. 1B), 65% of the 62

phenotype genes expressed in ≤3 tissues were predicted as nonfunctional. Further, there was a

significant correlation between the number of tissues with expression evidence and FL values of

all sequences in our analysis (r^2=0.77; $p < 2E-16$). Thus, the function prediction model is biased

against narrowly-expressed phenotype genes.

We also found that 80 pseudogenes (10%) were defined as functional (high-FL

pseudogenes). Consistent with misclassifications among phenotype genes, a key difference

between high-FL pseudogenes and those that were correctly predicted as nonfunctional was that

high-FL pseudogenes were more highly and broadly expressed (**Fig. 5**). A significantly higher

proportion of high-FL pseudogenes came from existing genome annotation as 19% of annotated

pseudogenes were classified as functional, compared to 4% of pseudogenes identified through a

computational pipeline (Zou et al 2009) (FET, $p < 1.5E-10$). We found that high-FL pseudogenes

might be more recently pseudogenized and thus have not yet lost many genic signatures, as the

mean number of disabling mutations (premature stop or frameshift) per kb in high-scoring

pseudogenes (1.9) were significantly lower than that of low-scoring pseudogenes (4.0; U test, $p <$

0.02). Lastly, we cannot rule out the possibility that a small subset of high-scoring pseudogenes

represent truly functional sequences, rather than false positives (e.g. Poliseno et al. 2010; Karreth

et al. 2015). Overall, the misclassification of both narrowly-expressed phenotype genes and

broadly-expressed pseudogenes highlights the need for an updated prediction model that is less

influenced by expression breadth.

To tailor functional predictions to narrowly-expressed sequences, we generated a "tissue-agnostic" model that attempts to minimize the contribution of biochemical activities occurring in many tissues by excluding expression breadth and features that were available across multiple tissues (see Methods). The tissue-agnostic model performed similarly to the full model (AUC-ROC=0.97; FNR=4%; FPR=15%; Supplemental Fig. 2; Supplemental Table 4). Importantly, the proportion of phenotype genes expressed in ≤3 tissues predicted as functional increased by 23% (35% in the full model to 58% in the tissue-agnostic model, Supplemental Fig. 1C), indicating that the tissue-agnostic model is more suitable for predicting the functionality of narrowly-expressed sequences than the full model, although there was an increase in FPR (from 10% to 15%). We next sought to evaluate the FL of ITR and annotated ncRNA sequences utilizing both the full model and the tissue-agnostic model, as these sequences were often narrowly-expressed (Supplemental Fig. 3A).

Intergenic transcribed regions and annotated ncRNAs are mostly predicted as nonfunctional

A subset of ITRs and ncRNAs likely represent novel genes or unannotated exon extensions of known genes (Johnson et al. 2005; Moghe et al. 2013). Nevertheless, most ITRs and ncRNAs are functionally ambiguous, as they are predominantly identified by the presence of expression evidence and few have been characterized genetically. To evaluate the functionality of ITRs and ncRNAs, we applied both the full and tissue-agnostic models to 895 ITRs, 136 TAIR ncRNAs, and 252 Araport long ncRNAs (referred to as Araport ncRNAs; see Methods). The median FLs based on the full model were low (0.09) for both ITRs (**Fig. 4F**) and Araport ncRNAs (**Fig. 4G**), and only 15% and 9% of these sequences were predicted as functional, respectively. By contrast, TAIR ncRNAs had a significantly higher median FL value (0.53; U tests, both *p*<5e-31; **Fig. 4H**) and 68% were predicted as functional. The higher proportion of functional TAIR ncRNA predictions compared to ITRs and Araport ncRNAs could be best explained by differences in features from the transcription activity category (**Fig. 5**). We also note that a greater proportion of ITRs and Araport ncRNAs are predicted as functional when considering only DNA methylation or H3 mark features (**Fig. 5**). However, these two category-specific models also had higher false positive rates (unexpressed intergenic sequences and pseudogenes, **Fig. 5**). Thus,

single feature models do not provide additional support for the functionality of most Araport

ncRNAs and ITRs.

We next applied the tissue-agnostic model that is less biased against narrowly-expressed

sequences (Supplemental Fig. 1C) to ITRs and TAIR/Araport ncRNAs that were generally

narrowly-expressed (Supplemental Fig. 3A). Compared to the full model, around twice as many

ITRs (30%) and Araport ncRNAs (19%) but a similar number of TAIR ncRNA (67%) were

predicted as functional. Considering the union of the full and tissue-agnostic model predictions,

268 ITRs (32%), 57 Araport ncRNAs (23%), and 105 TAIR ncRNAs (77%) were likely

functional. ITRs and annotated ncRNAs closer to annotated genes tended to be predicted as

functional (Supplemental Fig. 4A). Using the 95[th] percentile of intron lengths for all genes as a

threshold to call ITRs and annotated ncRNAs as proximal or distal to neighboring genes, 57% of

likely functional and 35% of likely nonfunctional ITRs and ncRNAs were proximal to

neighboring genes, respectively (FET, $p < 2E-09$). To assess if a subset these likely functional,

proximal ITRs/ncRNAs may be unannotated exons of known genes, we assessed whether they

tended to have similar features with their neighbors. Compared to feature similarities between

neighboring and random gene pairs (Supplemental Fig. 4B-D), likely functional ITRs/ncRNAs

were less similar to their neighbors, regardless of proximity (Supplemental Fig. 4C,D). Thus,

despite their proximity to annotated genes, it remains unclear if some ITRs or annotated ncRNAs

represent unannotated exon extensions of known genes or not. In addition, for proximal

functional ITRs/annotated ncRNAs, we cannot rule out the possibility that they represent false-

positive functional predictions due to the accessible and active chromatin states of nearby genes.

Given the challenge in ascertaining the origin of likely functional, proximal ITRs/ncRNAs, we

instead conservatively estimate that 116 distal, functional ITRs and annotated ncRNAs may

represent fragments of novel genes.

Intergenic transcribed regions and annotated ncRNAs do not resemble benchmark RNA

genes

Thus far, we predicted the majority of ITR and annotated ncRNA sequences as nonfunctional.

We demonstrated that the full model was able to predict conditional phenotype genes

(Supplemental Fig. 1A) and the tissue-agnostic model was more effective than the full model in

predicting narrowly expressed phenotype genes (Supplemental Fig. 1B,C). Thus, conditional or

tissue-specific functionality do not fully explain why the majority of ITRs and ncRNAs are

predicted as nonfunctional. However, the function prediction models so far were built by

contrasting protein-coding genes with pseudogenes and it remains possible that these protein-

coding gene-based models can not accurately predict RNA genes. To evaluate this possibility,

we generated a tissue-agnostic model using features calculated from a randomly-selected 100 bp

sequence within a phenotype protein-coding gene or pseudogene body (for features, see

Supplemental Table 6). The reason for using 100 bp sequence is that most RNA genes are too

short to be considered by earlier models, which were based on 500 bp sequences. In addition,

features from the tissue agnostic model are more suitable for RNA gene prediction as annotated

RNA genes tend to be more narrowly expressed than phenotype genes (U tests, all $p < 2e-05$;

Supplemental Fig. 3B). The 100 bp tissue-agnostic model performed similarly to the full 500 bp

model in distinguishing between phenotype protein-coding genes and pseudogenes, except with

higher FNR (AUC-ROC=0.97; FNR=13%; FPR=5%; Supplemental Fig. 5), but only predicted

three out of six RNA genes with documented mutant phenotypes (phenotype RNA genes) as

functional (Supplemental Fig. 5I). Further, other RNA Pol II-transcribed RNA genes exhibited

mixed predictions from the 100 bp tissue-agnostic model, as 15% of microRNA (miRNA)

primary transcripts (Supplemental Fig. 5J), 73% of small nucleolar RNAs (snRNAs;

Supplemental Fig. 5K), and 50% of small nuclear RNAs (snRNAs; Supplemental Fig. 5L) were

predicted as functional. Although the proportion of phenotype RNA genes predicted as

functional (50%) is significantly higher than the proportion of pseudogenes predicted as

functional (5%, FET, $p < 0.004$), this finding suggests that a model built with protein-coding

genes has a substantial FNR for detecting RNA genes.

To determine whether the suboptimal predictions by the phenotype protein-coding gene-

based models are because RNA genes belong to a class of their own, we next built a multi-class

function prediction model aimed at distinguishing four classes of sequences: benchmark RNA

genes (n=46), phenotype protein-coding genes (1,882), pseudogenes (3,916), and randomly-

selected, unexpressed intergenic regions (4,000). Benchmark RNA genes include six phenotype

RNA genes and 40 high-confidence miRNA primary transcript sequences (see Methods).

Unexpressed intergenic sequences were included to provide another set of likely nonfunctional

sequences distinct from pseudogenes. Expression breadth and tissue-specific features were

excluded from the four-class model and 100 bp sequences were used. In the four-class model,

87% of benchmark RNA genes, including all six phenotype RNA genes, were predicted as

functional sequences (65% RNA gene-like and 22% phenotype protein-coding gene-like; **Fig.**

6A). In addition, 95% of phenotype genes were predicted as functional (**Fig. 6B**), including 80%

of narrowly expressed genes, an increase of 22% over the 500 bp tissue-agnostic model

(Supplemental Fig. 1B). For the remaining two sequence classes, 70% of pseudogenes (**Fig. 6C**)

and 100% of unexpressed intergenic regions (**Fig. 6D**) were predicted as nonfunctional (either

pseudogenes or unexpressed intergenic sequences). Thus, the four-class model improves

prediction accuracy of RNA genes and narrowly expressed genes. However, the inclusion of

RNA genes in the model has significantly increased the ambiguity in pseudogene classification.

Since the four-class model was able to distinguish benchmark RNA genes from

nonfunctional sequence classes, we next evaluated whether ITRs and annotated ncRNAs

resemble functional sequences with the four-class model. Note that the 100 bp model used here

allowed us to evaluate an additional 10,938 ITRs and 1,406 annotated ncRNAs. We find that

34% of ITR, 38% of Araport ncRNA, and of 65% TAIR ncRNAs were predicted as functional

sequences (**Fig. 6E-G**). To provide an overall estimate of the proportion of likely-functional

ITRs and annotated ncRNAs, we considered the predictions from the four-class model (**Fig. 6**),

the full model (**Fig. 3,4**), and the tissue-agnostic models (Supplemental Fig. 2,5). Based on

support from at least one of the four models, we classified 4,437 ITRs (38%) and 796 annotated

ncRNAs (44%) as functional, as they resembled either phenotype protein-coding or RNA genes.

Our findings lend support that they are likely parts of novel or annotated genes. Meanwhile, we

find that a substantial number of ITRs (62%) and annotated ncRNAs (56%) are predicted as

nonfunctional. Moreover, at least a third of ITRs (**Fig. 6E**) and Araport ncRNAs (**Fig. 6F**) most

closely resemble unexpressed intergenic regions. Thus, we show that the majority of ITRs and

annotated ncRNA regions resemble nonfunctional genomic regions, and therefore could

represent regions of noisy transcription.

CONCLUSION

Discerning the location of functional regions within a genome represents a key goal in genomic

biology. Despite advances in computational gene finding, it remains challenging to determine

whether intergenic transcribed regions (ITRs) represent functional or noisy biochemical activity.

We established robust function prediction models based on the evolutionary, biochemical, and

structural characteristics of phenotype genes and pseudogenes. The prediction models accurately

define functional and nonfunctional regions and are applicable genome-wide. These results echo

recent findings that human phenotype genes could be distinguished from pseudogenes (Tsai et al.

2017). Given that function predictions were successful in both plant and metazoan model

systems, integrating the evolutionary and biochemical features of known genes will likely be

applicable to any species. The next step will be to test whether function prediction models can be

applied across species, which could ultimately allow the phenotype data and omics resources

available in model systems to effectively guide the identification of functional regions in non-

models.

Expression data was highly informative to functional predictions. We found that the

prediction model based on only 24 transcription activity-related features performs nearly as well

as the full model that integrates additional information including conservation, H3 mark,

methylation, and TF binding data. In human, use of transcription data from cell lines also

produced highly accurate predictions of functional genomic regions (AUC-ROC=0.96; Tsai et al.

2017). Despite the importance of transcription data, we emphasize that the presence of

expression evidence is an extremely poor predictor. Taken together, these results indicate that

function prediction models can be established in any species, model or not, with a modest

number of transcriptome datasets (e.g. 51 in this study and 19 in human). One caveat of the

current model is that narrowly-expressed phenotype genes are frequently predicted as

pseudogene and broadly-expressed pseudogenes tend to be called functional. To improve the

function prediction model, it will be important to explore additional features unrelated to

transcription. Because few phenotype genes are narrowly-expressed (5%) in the *A. thaliana*

training data, more phenotyping data for narrowly expressed genes will be crucial as well.

Upon application of the function prediction models genome-wide, we found that 4,427

ITRs and 796 annotated ncRNAs in *A. thaliana* are likely functional. Assuming each entry

equals a novel gene, this estimate represents a 19% increase in annotated gene space (excluding

annotated ncRNAs) for the model plant. However, considering the high false positive rates (e.g.

10% for the full and 31% for the four-class model), this is most likely an overestimate of the

number of novel genes contributed by functional ITRs and annotated ncRNAs. In addition, we

emphasize that the majority of ITRs and ncRNAs resemble pseudogenes and random

unexpressed intergenic regions. Similarly, most human ncRNAs are more similar to

nonfunctional sequences than they are to protein coding and RNA genes (Tsai et al. 2017).

Furthermore, the significant relationship between the amount of intergenic expression occurring

in a species and the size of a genome is consistent with the interpretation that intergenic

transcripts are generally nonfunctional. Thus, instead of assuming any expressed sequence must

be functionally significant, we advocate that the null hypothesis should be that it is not,

particularly considering that most ITRs and annotated ncRNAs have not been experimentally

characterized. The machine learning framework we have described provides an approach for

distinguishing between functional and noisy biochemical activity, and will help defining the gene

space in a genome.

METHODS

Identification of transcribed regions in leaf tissue of 15 flowering plants

RNA-sequencing (RNA-seq) datasets were retrieved from the Sequence Read Archive (SRA) at

the National Center for Biotechnology Information (NCBI; www.ncbi.nlm.nih.gov/sra/) for 15

flowering plant species (Supplemental Table 1). All datasets were generated from leaf tissue and

sequenced on Illumina HiSeq 2000 or 2500 platforms. Genome sequences and gene annotation

files were downloaded from Phytozome v.11 (www.phytozome.net; Goodstein et al. 2011) or

Oropetium Base v.01 (www.sviridis.org; VanBuren et al. 2015). Genome sequences were repeat

masked using RepeatMasker v4.0.5 (www.repeatmasker.org) if a repeat-masked version was not

available. Only one end from paired-end read datasets were utilized in downstream processing.

Reads were trimmed to be rid of low scoring ends and residual adaptor sequences using

Trimmomatic v0.33 (LEADING:3 TRAILING:3 SLIDINGWINDOW:4:20 MINLEN:20, Bolger

et al. 2014) and mapped to genome sequences using TopHat v2.0.13 (default parameters except

as noted below; Kim et al. 2013). Reads ≥20 nucleotides in length that mapped uniquely within a

genome were used in further analysis.

For each species, thirty million mapped reads were randomly selected from among all

datasets and assembled into transcript fragments using Cufflinks v2.2.1 (default parameters

except as noted below, Trapnell et al. 2010), while correcting for sequence-specific biases during

the sequencing process by providing an associated genome sequence with the -b flag. The

expected mean fragment length for assembled transcript fragments in Cufflinks was set to 150

from the default of 200 so that expression levels in short fragments would not be overestimated.

The 1st and 99th percentile of intron lengths for each species were used as the minimum and

maximum intron lengths, respectively, for both the TopHat2 and Cufflinks steps. Intergenic

transcribed regions (ITRs) were defined by transcript fragments that did not overlap with gene

annotation and did not have significant six-frame translated similarity to plant protein sequences

in Phytozome v.10 (BLASTX E-value < 1E-05). The correlation between assembled genome

size and gene counts was determined with data from the first 50 published plant genomes

(Michael and Jackson, 2013).

Phenotype data sources

Mutant phenotype data for *A. thaliana* protein-coding genes was collected from a published

dataset (Lloyd and Meinke 2012), the Chloroplast 2010 database (Ajjawi et al. 2010; Savage et

al. 2013), and the RIKEN phenome database (Kuromori et al. 2006) as described by Lloyd et al.

(2015). Phenotype genes used in our analyses were those whose disruption resulted in lethal or

visible defects under standard laboratory growth conditions. Genes with documented mutant

phenotypes under standard conditions were considered as a distinct and non-overlapping

category from other annotated protein-coding genes. We identified six RNA genes with

documented loss-of-function phenotypes through literature searches (Supplemental Table 7): *At4*

(AT5G03545; Shin et al. 2006), *MIR164A* and *MIR164D* (AT2G47585 and AT5G01747,

respectively; Guo et al. 2005), *MIR168A* (AT4G19395; Li et al. 2012b), and *MIR828A* and *TAS4*

(AT4G27765 and AT3G25795, respectively; Hsieh et al. 2009). Conditional phenotype genes

were those belonging to the Conditional phenotype group as described by Lloyd and Meinke

(2012). Loss-of-function mutants of these genes exhibited phenotype only under stress

conditions.

Arabidopsis thaliana **genome annotation**

A. thaliana protein-coding gene, miRNA gene, snoRNA gene, snRNA gene, ncRNA region,

pseudogene, and transposable element annotations were retrieved from The Arabidopsis

Information Resource v.10 (TAIR10; www.arabidopsis.org; Berardini et al. 2015). Additional

miRNA gene and lncRNA region annotations were retrieved from Araport v.11

(www.araport.org; Krishnakumar et al. 2015). A pseudogene-finding pipeline (Zou et al. 2009)

was used to identify additional pseudogene fragments and count the number of disabling

mutations (premature stop or frameshift mutations). Genes, pseudogenes, and transposons with

overlapping annotation were excluded from further analysis. Overlapping lncRNA annotations

were merged for further analysis. When pseudogenes from TAIR10 and the pseudogene-finding

pipeline overlapped, the longer pseudogene annotation was used.

A. thaliana ITRs analyzed include: (1) the Set 2 ITRs in Moghe et al. (2013), (2) the

novel transcribed regions from Araport v.11, and (3) additional ITRs from 206 RNA-seq datasets

(Supplemental Table 5). Reads were trimmed, mapped, and assembled into transcript fragments

as described above, except that overlapping transcript fragments from across datasets were

merged. ITRs analyzed did not overlap with any TAIR10, Araport11, or pseudogene annotation.

Overlapping ITRs from different annotated subsets were kept based on a priority system:

Araport11 > Set 2 ITRs from Moghe et al. (2013) > ITRs identified in this study. For each

sequence entry (gene, ncRNA, pseudogene, transposable element, or ITR), a 100 and 500 base

pair (bp) window was randomly chosen for calculating feature values and subsequent model

building steps. Feature descriptions are provided in the following sections. The feature values for

randomly selected 500 and 100 bp windows are provided in Supplemental Table 2 and 6,

respectively. Additionally, non-expressed intergenic sequences were randomly-sampled from

genome regions that did not overlap with annotated genes, pseudogenes, transposable elements,

or regions with genic or intergenic transcript fragments (100 bp, n=4,000; 500 bp, n=3,716). All

100 and 500 bp windows described above are referred to as sequence windows throughout the

Methods section.

Sequence conservation and structure features

There were 10 sequence conservation features examined. The first two were derived from

comparisons between *A. thaliana* accessions including nucleotide diversity and Tajima's D

among 81 accessions (Cao et al. 2011) using a genome matrix file from the 1,001 genomes

database (www.1001genomes.org). The python scripts are available through GitHub

(https://github.com/ShiuLab/GenomeMatrixProcessing). The remaining eight features were

derived from cross-species comparisons, three based on multiple sequence and five based on

pairwise alignments. Three multiple sequence alignment-based features were established using

aligned genomic regions between *A. thaliana* and six other plant species (*Glycine max*,

Medicago truncatula, *Populus trichocarpa*, *Vitis vinifera*, *Sorghum bicolor*, and *Oryza sativa*)

(referred to as conserved blocks). For each conserved block, the first feature was the proportion

of a sequence window that overlapped a conserved block (referred to as coverage), and the two

other features were the maximum and average phastCons scores within each sequence window.

The phastCons score was determined for each nucleotide within conserved blocks (Li et al.

2012a). Nucleotides in a sequence window that did not overlap with a conserved block were

assigned a phastCons score of 0. For each sequence window, five pairwise alignment-based

cross-species conservation features were the percent identities to the most significant BLASTN

match (if E-value<1E-05) in each of five taxonomic groups. The five taxonomic groups included

the *Brassicaceae* family ($n_{species}$=7), other dicotyledonous plants (22), monocotyledonous plants

(7), other embryophytes (3), and green algae (5). If no sequence with significant similarity was

present, percent identity was scored as zero.

For sequence-structure features, we used 125 conformational and thermodynamic

dinucleotide properties collected from DiProDB database (Friedel et al. 2009). Because the

number of dinucleotide properties was high and dependent, we reduced the dimensionality by

utilizing principal component (PC) analysis as described previously (Tsai et al. 2015). Sequence-

structure values corresponding to the first five PCs were calculated for all dinucleotides in and

averaged across the length of a sequence window and used as features when building function

prediction models.

Transcription activity features

We generated four multi-dataset and 20 individual dataset transcription activity features. To

identify a set of RNA-seq datasets to calculate multi-dataset features, we focused on the 72 of

206 RNA-seq datasets each with ≥20 million reads (see above; Supplemental Table 5).

Transcribed regions were identified with TopHat2 and Cufflinks as described in the RNA-seq

analysis section except that the 72 *A. thaliana* RNA-seq datasets were used. Following transcript

assembly, we excluded 21 RNA-seq datasets because they had unusually high RPKM (Reads Per

Kilobase of transcript per Million mapped reads) values (median RPKM value

range=272~2,504,294) compared to the rest (2~252). The remaining 51 RNA-seq datasets were

used to generate four multi-dataset transcription activity features including: expression breadth,

95[th] percentile expression level, maximum transcript coverage, and presence of expression

evidence (for values see Supplemental Table 5). Expression breadth was the number of RNA-seq

datasets that have ≥1 transcribed region that overlapped with a sequence window. The 95^{th}

percentile expression level was the 95^{th} percentile of RPKM values across 51 RNA-seq datasets

where RPKM values were set to 0 if there was no transcribed region for a sequence window.

Maximum transcript coverage was the maximum proportion of a sequence window that

overlapped with a transcribed region across 51 RNA-seq datasets. Presence of expression

evidence was determined by overlap between a sequence window and any transcribed region in

the 51 RNA-seq datasets.

In addition to features based on multiple datasets, 20 individual dataset features were

derived from 10 datasets: seven tissue/organ-specific RNA-seq datasets including pollen

(SRR847501), seedling (SRR1020621), leaf (SRR953400), root (SRR578947), inflorescence

(SRR953399), flower, (SRR505745) and silique (SRR953401), and three datasets from non-

standard growth conditions, including dark-grown seedlings (SRR974751) and leaf tissue under

drought (SRR921316) and fungal infection (SRR391052). For each of these 10 RNA-seq

datasets, we defined two features for each sequence window: the maximum transcript coverage

(as described above) and the maximum RPKM value of overlapping transcribed regions (referred

to as Level in **Fig. 2**). If no transcribed regions overlapped a sequence window, the maximum

RPKM value was set as 0. For the analysis of narrowly- and broadly-expressed phenotype genes

and pseudogenes (Supplemental Fig. 1B,C), we used 28 out of 51 RNA-seq datasets generated

from a single tissue and in standard growth conditions to calculate the number of tissues with

evidence of expression (tissue expression breadth). In total, seven tissues were represented

among the 28 selected RNA-seq datasets (see above; Supplemental Table 5), and thus tissue

expression breadth ranges from 0 to 7 (note that only 1 through 7 are shown in Supplemental Fig.

1B,C due to low sample size of phenotype genes in the 0 bin). The tissue breadth value is distinct

from the expression breadth feature used in model building that was generated using all 51

datasets and considered multiple RNA-seq datasets from the same tissue separately (range: 0-

51).

Histone 3 mark features

Twenty histone 3 (H3) mark features were calculated based on eight H3 chromatin

immunoprecipitation sequencing (ChIP-seq) datasets from SRA. The H3 marks examined

include four associated with activation (H3K4me1: SRR2001269, H3K4me3: SRR1964977,

H3K9ac: SRR1964985, and H3K23ac: SRR1005405) and four associated with repression

(H3K9me1: SRR1005422, H3K9me2: SRR493052, H3K27me3: SRR3087685, and H3T3ph:

SRR2001289). Reads were trimmed as described in the RNA-seq section and mapped to the

TAIR10 genome with Bowtie v2.2.5 (default parameters; Langmead et al. 2009). Spatial

Clustering for Identification of ChIP-Enriched Regions (SICER) v.1.1 (Xu et al. 2014) was used

to identify ChIP-seq peaks with a false discover rate ≤ 0.05 with a non-overlapping window size

of 200, a gap parameter of 600, and an effective genome size of 0.92 according to Koehler et al.

(2011). For each H3 mark, two features were calculated for each sequence window: the

maximum intensity among overlapping peaks and peak coverage (proportion of overlap with the

peak that overlaps maximally with the sequence window). In addition, four multi-mark features

were generated. Two of the multi-mark features were the number of activating marks (0-4)

overlapping a sequence window and the proportion of a sequence window overlapping any peak

from any of the four activating marks (activating mark peak coverage). The remaining two multi-

mark features were the same as the two activating multi-mark features except focused on the four

repressive marks.

DNA methylation features

Twenty-one DNA methylation features were calculated from bisulfite-sequencing (BS-seq)

datasets from seven tissues (pollen: SRR516176, embryo: SRR1039895, endosperm:

SRR1039896, seedling: SRR520367, leaf: SRR1264996, root: SRR1188584, and inflorescence:

SRR2155684). BS-seq reads were trimmed as described above and processed with Bismark v.3

(default parameters; Krueger and Andrews 2011) to identify methylated and unmethylated

cytosines in CG, CHH, and CHG (H = A, C, or T). Methylated cytosines were defined as those

with ≥5 mapped reads and with >50% of mapped reads indicating that the position was

methylated. For each BS-seq dataset, the percentage of methylated cytosines in each sequence

window for CG, CHG, and CHH contexts were calculated if the sequence window had ≥5

cytosines with ≥5 reads mapping to the position. To determine whether the above parameters

where reasonable, we assessed the false positive rate of DNA methylation calls by evaluating the

proportion of cytosines in the chloroplast genome that are called as methylated, as the

chloroplast genome has few DNA methylation events (Ngernprasirtsiri et al. 1988; Zhang et al.

2006). Based on the above parameters, 0-1.5% of cytosines in CG, CHG, or CHH contexts in the

chloroplast genome were considered methylated in any of the seven BS-seq datasets. This

indicated that the false positive rates for DNA methylation calls were low and the parameters

were reasonable.

Chromatin accessibility and transcription factor binding features

Chromatin accessibility features consisted of ten DHS-related features and one micrococcal

nuclease sequencing (MNase-seq)-derived feature. DHS peaks from five tissues (seed coat,

seedling, root, unopened flowers, and opened flowers) were retrieved from the Gene Expression

Omnibus (GSE53322 and GSE53324; Sullivan et al. 2014). For each of the five tissues, the

maximum DHS peak intensity and DHS peak coverage were calculated for each sequence

window. Normalized nucleosome occupancy per bp based on MNase-seq was obtained from Liu

et al. (2015). The average nucleosome occupancy value was calculated across each sequence

window. Transcription factor (TF) binding site features were based on *in vitro* DNA affinity

purification sequencing data of 529 TFs (O'Malley et al. 2016). Two features were generated for

each sequence window: the total number of TF binding sites and the number of distinct TFs

bound.

Single-feature prediction performance

The ability for each single feature to distinguish between functional and nonfunctional regions

was evaluated by calculating AUC-ROC value with the Python scikit-learn package (Pedregosa

et al. 2011). AUC-ROC values range between 0.5 (equivalent to random guessing) and 1 (perfect

predictions) and values above 0.7, 0.8, and 0.9 are considered to be fair, good, and excellent,

respectively. Thresholds to predict sequences as functional or nonfunctional using a single

feature were defined by the feature value that produced the highest F-measure, the harmonic

mean of precision (proportion of sequences predicted as functional that are truly functional) and

recall (proportion of truly functional sequences predicted as functional). The F-measure allows

consideration of both false positives and false negatives at a given threshold. FPR were

calculated as the percentage of negative (nonfunctional) cases with values above or equal to the

threshold and thus falsely predicted as functional. FNR were calculated as the percentage of

positive (functional) cases with values below the threshold and thus falsely predicted as

nonfunctional.

Binary classification with machine learning

For binary classification (two-class) models that contrasted phenotype genes and pseudogenes,

the random forest (RF) implementation in the Waikato Environment for Knowledge Analysis

software (WEKA; Hall et al. 2009) was utilized. Three types of two-class models were

established, including the full model (500 bp sequence window, **Fig. 3C,D** and **Fig. 4**), tissue-

agnostic models (500 bp, Supplemental Fig. 2; 100 bp, Supplemental Fig. 5), and single feature

category models (**Fig. 3C,D**). For each model type, we first generated 100 balanced datasets by

randomly selecting equal numbers of phenotype genes (positive examples) and pseudogenes

(negative examples). For each of these 100 datasets, 10-fold stratified cross-validation was

utilized, where the model was trained using 90% of sequences and tested on the remaining 10%.

Thus, for each model type, a sequence window had 100 prediction scores, where each score was

the proportion of 500 random forest trees that predicted a sequence as a phenotype gene in a

balanced dataset. The median of 100 prediction scores was used as the functional likelihood (FL)

value (Supplemental Table 4). The FL threshold to predict a sequence as functional or

nonfunctional was defined based on maximum F-measure as described in the previous section.

We tested multiple -K parameters (2 to 25) in the WEKA-RF implementation, which alters the

number of randomly-selected features included in each RF tree (Supplemental Table 8), and

found that 15 randomly-selected features provided the highest performance based on AUC-ROC

(calculated and visualized using the ROCR package; Sing et al. 2005). Binary classification

models were also built using all features from 500 bp sequences (equivalent to the full model)

with the Sequential Minimal Optimization - Support Vector Machine (SMO-SVM)

implementation in WEKA (Hall et al. 2009). The results of SMO-SVM models were highly

similar to the full RF results: *PCC* between the FL values generated by RF and SMO-

SVM=0.97; AUC-ROC of SMO-SVM=0.97; FPR=12%; FNR=3%. By comparison, the full RF

model had AUC-ROC=0.98, FPR=10%, FNR=4%.

Tissue-agnostic models were generated by excluding the expression breadth feature and

95[th] percentile expression level and replacing all features from RNA-seq, BS-seq, and DHS

datasets that were available in multiple tissues. For multiple-tissue RNA-seq data, the maximum

expression level across 51 RNA-seq datasets (in RPKM) and maximum coverage (as described

in the transcription activity section) of a sequence window in any of 51 RNA-seq datasets were

used. For multi-tissue DNA methylation features, minimum proportions of methylated cytosines

in any tissue in CG, CHG, and CHH contexts were used. For DHS data, the maximum peak

intensity and peak coverage was used instead. In single feature category predictions, fewer total

features were used and therefore lower –K values (i.e. the number of random features selected

when building random forests) were considered in parameter searches (Supplemental Table 8).

Multi-class machine learning model

For the four-class model, benchmark RNA gene, phenotype protein-coding gene, pseudogene,

and random unexpressed intergenic sequences were used as the four training classes. Benchmark

RNA genes consisted of six RNA genes with documented loss-of-function phenotypes and 40

high-confidence miRNA genes from miRBase (www.mirbase.org; Kozomara and Griffiths-Jones

2014). We generated 250 datasets with equal proportions (larger classes randomly sampled) of

training sequences. Two-fold stratified cross-validation was utilized due to the low number of

benchmark RNA genes. The features included those described for the tissue-agnostic model and

focused on 100 bp sequence windows. The RF implementation, *cforest*, in the *party* package of

R (Strobl et al. 2008) was used to build the classifiers. The four-class predictions provide

prediction scores for each sequence type: an RNA gene, phenotype protein-coding gene,

pseudogene, and unexpressed intergenic score (Supplemental Table 4). The prediction scores

indicate the proportion of random forest trees that classify a sequence as a particular class.

Median prediction scores from across 100 balanced runs were used as final prediction scores.

Scores from a single balanced dataset models sum to 1, but not the median from 100 balanced

runs. Thus, the median scores were scaled to sum to 1. For each sequence window, the maximum

prediction score among the four classes was used to classify a sequence as phenotype gene,

pseudogene, unexpressed intergenic, or RNA gene.

FIGURE LEGENDS

Figure 1. Relationship between genome size and number of nucleotides covered by RNA-seq

reads (expression) in 15 flowering plant species. (*A*) annotated genic regions. (*B*) intergenic

regions excluding any annotated features. Mb: megabase. Gb: gigabase. Dotted lines: linear

model fits. r^2: square of Pearson's correlation coefficient.

Figure 2. Predictions of functional (phenotype gene) and non-functional (pseudogene) sequences

based on each individual feature. Prediction performance is measured using Area Under the

Curve - Receiver Operating Characteristic (AUC-ROC). Features include those in the categories

of (*A*) transcription activity, (*B*) sequence conservation, (*C*) DNA methylation, (*D*) transcription

factor (TF) binding, (*E*) histone 3 (H3) marks, (*F*) sequence structure, and (*G*) chromatin

accessibility. AUC-ROC ranges in value from 0.5 (equivalent to random guessing) to 1 (perfect

predictions). Dotted lines: median AUC-ROC of features in a category.

Figure 3. Predictions of functional and nonfunctional sequences based on multiple features.

Smoothed scatterplots of the first two principle components (PCs) of (*A*) phenotype gene and (*B*)

pseudogene features. The percentages on the axes in *(A)* indicate the feature value variation

explained by the associated PC. (*C*) AUC-ROC values of function prediction models built when

considering all features (Full), all except transcription activity (TX)-related features (Full (-TX)),

and all features from each category. The category abbreviations follow those in **Fig. 2**. (*C*)

Precision-recall curves of the models with matching colors from (*B*). The models were built

using feature values calculated from 500 bp sequence windows.

Figure 4. Functional likelihood distributions of various sequence classes based on the full

model. (*A*) Phenotype genes. (*B*) Pseudogenes. (*C*) Annotated protein-coding genes. (*D*)

Transposable elements. (*E*) Random unexpressed intergenic sequences. (*F*) Intergenic

transcribed regions (ITR). (*G*) Araport11 ncRNAs. (*H*) TAIR10 ncRNAs. The full model was

established using 500 bp sequence windows. Higher and lower functional likelihood values

indicate greater similarity to phenotype genes and pseudogenes, respectively. Vertical dashed

lines indicate the threshold for calling a sequence as functional or nonfunctional. The

percentages to the left and right of the dashed line indicate the percent of sequences predicted as

functional or nonfunctional, respectively.

Figure 5. Proportion of phenotype genes, pseudogenes, ITRs, and ncRNAs predicted as

functional in the full and single-category models. Percentages of sequence classes that are

predicted as functional in models based on all features and the single category models, each

using all features from a category (abbreviated according to **Figure 2**. The models are sorted

from left to right based on performance (AUC-ROC). The colors of and numbers within the

blocks indicate the proportion sequences predicted as functional by a given model. Phenotype

gene and pseudogene sequences are shown in three sub-groups: all sequences (All), and those

predicted as functional (high functional likelihood (FL)) and nonfunctional (low FL) in the full

model. ITR: intergenic transcribed regions.

Figure 6. Function predictions based on a four-class prediction model. (*A*) Stacked bar plots

indicate the prediction scores of benchmark RNA genes for each of the four classes: dark blue -

phenotype protein-coding gene (Ph), cyan - RNA gene (RNA), red - pseudogene (Ps), yellow –

random intergenic sequence (Ig). A benchmark RNA gene is classified as one of the four classes

according to the highest prediction score. The color bars below the chart indicate the predicted

class, with the same color scheme as the prediction score. Sequences classified as Ph or RNA

were considered functional, while those classified as Ps or Ig were considered nonfunctional.

Percentages below a classification region indicate the proportion of sequences classified as that

class. (*B*) Phenotype protein-coding gene prediction scores. (*C*) Pseudogene prediction scores.

(*D*) Random unexpressed intergenic region prediction scores. Note that no sequence was

predicted as functional. (*E*) Intergenic transcribed region (ITR), (*F*) Araport11 ncRNA regions.

(*G*) TAIR10 ncRNA regions.

SUPPLEMENTAL FIGURE LEGENDS

Supplemental Figure 1. Impacts of conditional phenotypes and expression breadth on the

function prediction model. (*A*) Functional likelihood distributions of phenotype genes with

mutant phenotypes under standard growth conditions (non-conditional) and non-standard growth

conditions such as stressful environments (conditional) based on the 500 bp full model. Feature

values were calculated from a random 500 bp region from within the sequence body. Higher and

lower functional likelihood values indicate a greater similarity to phenotype genes and

pseudogenes, respectively. (*B,C*) Distributions of functional likelihood scores for phenotype

genes (blue) and pseudogenes (red) for sequences with various breadths of expression for (*B*) the

500 bp full model and (*C*) the 500 bp tissue-agnostic model generated by excluding the

expression breadth and features available from multiple tissues. The tissue-agnostic model is

aimed toward minimizing the effects of biochemical activity occurring across multiple tissues

and predicts a greater proportion of narrowly-expressed phenotype genes as functional compared

to the full model.

Supplemental Figure 2. Distributions of functional likelihood scores based on the 500 bp tissue-
agnostic model. (*A*) Phenotype genes. (*B*) Pseudogenes. (*C*) Annotated protein-coding genes. (*D*)
Transposable elements. (*E*) Random unexpressed intergenic sequences. (*F*) Intergenic
transcribed regions (ITR). (*G*) Araport11 ncRNAs. (*H*) TAIR10 ncRNAs. Vertical dashed lines
display the threshold to define a sequence as functional or nonfunctional. The numbers to the left
and right of the dashed line show the percentage of sequences predicted as functional or
nonfunctional, respectively.

Supplemental Figure 3. Distributions of expression breadth of different sequence classes. (*A*)
Based on 500 bp feature regions. (*B*) Based on 100 bp feature regions.

Supplemental Figure 4. Distance of ITRs and annotated ncRNA regions to and feature
similarity with neighboring genes. (*A*) Distance from intergenic transcribed regions (ITRs) and
annotated ncRNAs to the closest neighboring gene. ITR and ncRNA sequences are separated by
whether they are predicted as functional (F) or nonfunctional (NF) by the 500 bp full model. (*B*)
Feature similarity based on Pearson's Correlation Coefficients (PCC) between random pairs of
ITRs, Araport11 ncRNAs, TAIR10 ncRNAs, or annotated genes. (*C*) Feature similarity between
proximal neighbors (within 95th percentile (456 bp) of intron lengths), and (*D*) Feature similarity
between distal neighbors (>456 bp). Pairs involving ITRs and annotated ncRNAs were divided
by whether the ITR or ncRNA sequence was predicted as functional (F) or nonfunctional (NF)
by the full model. Feature values were quantile normalized prior to calculating correlations.

Supplemental Figure 5. Distributions of functional likelihood scores based on the 100 bp tissue-
agnostic model. (*A*) Phenotype genes. (*B*) Pseudogenes. (*C*) Protein-coding gene. (*D*)
transposable elements. (*E*) Random unexpressed intergenic sequences. (*F*) Intergenic transcribed
regions (ITR). (*G*) Araport11 ncRNAs. (*H*) TAIR10 ncRNAs. (*I*) RNA genes with loss-of-
function mutant phenotypes. (J) MicroRNAs, (*K*) Small nucleolar RNAs, (*L*) Small nuclear
RNAs. The tissue-agnostic model was built with 100 bp features and while excluding the
expression breadth and tissue-specific features. Higher functional likelihood values indicate
greater similarity to phenotype genes while lower values indicate similarity to pseudogenes.
Vertical dashed lines display the threshold to define a sequence as functional or nonfunctional.

The numbers to the left and right of the dashed line show the percentage of sequences predicted

as functional or nonfunctional, respectively.

SUPPLEMENTAL TABLES

Supplemental Table 1. Leaf tissue RNA-sequencing datasets for 15 flowering plant species

Supplemental Table 2. Conservation, biochemical, and sequence-structure feature values

calculated from 500 bp sequences.

Supplemental Table 3. False positive and false negative rates for single feature classifications.

Supplemental Table 4. Function predictions for all models generated in this study.

Supplemental Table 5. RNA-sequencing datasets for identifying intergenic transcribed regions,

calculating transcription activity features, and assessing tissue-specific predictions.

Supplemental Table 6. Conservation, biochemical, and sequence-structure feature values

calculated from 100 bp sequences.

Supplemental Table 7. RNA genes with documented loss-of-function phenotypes.

Supplemental Table 8. K parameters tested for random forest runs.

DATA ACCESS

All data are available in the text of this article or in the supplemental material.

ACKNOWLEDGEMENTS

The authors wish to thank Christina Azodi, Ming-Jung Liu, Gaurav Moghe, Bethany Moore, and

Sahra Uygun and for providing processed data and discussion. This work was supported by

National Science Foundation grants (IOS-1126998, IOS-1546617, and DEB-1655386) to S.-H.S,

and Research Experience for Undergraduates support to R.P.S, as well as the Michigan State

University Dissertation Continuation Fellowship to J.P.L.

AUTHOR CONTRIBUTIONS

J.P.L., Z.T.-Y.T., and S.-H.S. designed the research. J.P.L., Z.T.-Y.T., R.P.S., and N.L.P.

performed the research. J.P.L., Z.T-Y.T., R.P.S., N.L.P., and S.-H.S. wrote the article.

DISCLOSURE DECLARATION

The authors have no conflicts of interest to disclose.

REFERENCES

Ajjawi I, Lu Y, Savage LJ, Bell SM, Last RL. 2010. Large-scale reverse genetics in Arabidopsis:

case studies from the Chloroplast 2010 Project. *Plant Physiol* **152**: 529–540.

Amundson R, Lauder GV. 1994. Function without purpose. *Biol Philos* **9**: 443–469

Bennetzen JL. 2005. Mechanisms of Recent Genome Size Variation in Flowering Plants. *Ann*

Bot **95**: 127–132.

Berardini TZ, Reiser L, Li D, Mezheritsky Y, Muller R, Strait E, Huala E. 2015. The

Arabidopsis information resource: Making and mining the "gold standard" annotated

reference plant genome. *Genesis* **53**: 474–485.

Bernard D, Prasanth KV, Tripathi V, Colasse S, Nakamura T, Xuan Z, Zhang MQ, Sedel F,

Jourdren L, Coulpier F, et al. 2010. A long nuclear-retained non-coding RNA regulates

synaptogenesis by modulating gene expression. *EMBO J* **29**: 3082–3093.

Boeck ME, Huynh C, Gevirtzman L, Thompson OA, Wang G, Kasper DM, Reinke V, Hillier

LW, Waterston RH. 2016. The time-resolved transcriptome of C. elegans. *Genome Res* **26**:

1441–1450.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics* **30**: 2114–2120.

Brenchley R, Spannagl M, Pfeifer M, Barker GLA, D'Amore R, Allen AM, McKenzie N,

Kramer M, Kerhornou A, Bolser D, et al. 2012. Analysis of the bread wheat genome using

whole-genome shotgun sequencing. *Nature* **491**: 705–710.

Brown JB, Boley N, Eisman R, May GE, Stoiber MH, Duff MO, Booth BW, Wen J, Park S,
Suzuki AM, et al. 2014. Diversity and dynamics of the Drosophila transcriptome. *Nature*
512: 393–399.

Cabili MN, Trapnell C, Goff L, Koziol M, Tazon-Vega B, Regev A, Rinn JL. 2011. Integrative
annotation of human large intergenic noncoding RNAs reveals global properties and
specific subclasses. *Genes Dev* **25**: 1915–1927.

Cao J, Schneeberger K, Ossowski S, Günther T, Bender S, Fitz J, Koenig D, Lanz C, Stegle O,
Lippert C, et al. 2011. Whole-genome sequencing of multiple Arabidopsis thaliana
populations. *Nat Genet* **43**: 956–963.

Chen Z-X, Sturgill D, Qu J, Jiang H, Park S, Boley N, Suzuki AM, Fletcher AR, Plachetzki DC,
FitzGerald PC, et al. 2014. Comparative validation of the D. melanogaster modENCODE
transcriptome annotation. *Genome Res* **24**: 1209–1223.

Cummins R. 1975. Functional Analysis. *J Philos* **72**: 741.

Doolittle WF, Brunet TDP, Linquist S, Gregory TR. 2014. Distinguishing between "function"
and "effect" in genome biology. *Genome Biol Evol* **6**: 1234–1237.

Eddy SR. 2013. The ENCODE project: missteps overshadowing a success. *Curr Biol* **23**: R259–
61

ENCODE Project Consortium. 2012. An integrated encyclopedia of DNA elements in the human
genome. *Nature* **489**: 57–74.

Friedel M, Nikolajewa S, Sühnel J, Wilhelm T. 2009. DiProDB: a database for dinucleotide
properties. *Nucleic Acids Res* **37**: D37–40.

Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten
U, Putnam N, et al. 2011. Phytozome: a comparative platform for green plant genomics.
Nucleic Acids Res **40**: D1178–D1186.

Graur D, Zheng Y, Price N, Azevedo RBR, Zufall RA, Elhaik E. 2013. On the immortality of
television sets: "function" in the human genome according to the evolution-free gospel of

ENCODE. *Genome Biol Evol* **5**: 578–590.

Gulko B, Gronau I, Hubisz MJ, Siepel A. 2014. *Probabilities of Fitness Consequences for Point*
Mutations Across the Human Genome. http://dx.doi.org/10.1101/006825.

Guo H-S, Xie Q, Fei J-F, Chua N-H. 2005. MicroRNA directs mRNA cleavage of the
transcription factor NAC1 to downregulate auxin signals for Arabidopsis lateral root
development. *Plant Cell* **17**: 1376–1386.

Guttman M, Amit I, Garber M, French C, Lin MF, Feldser D, Huarte M, Zuk O, Carey BW,
Cassady JP, et al. 2009. Chromatin signature reveals over a thousand highly conserved large
non-coding RNAs in mammals. *Nature* **458**: 223–227.

Hall M, Frank E, Holmes G, Pfahringer B, Reutemann P, Witten IH. 2009. The WEKA data
mining software. *ACM SIGKDD Explorations Newsletter* **11**: 10.

Hanada K, Zhang X, Borevitz JO, Li W-H, Shiu S-H. 2007. A large number of novel coding
small open reading frames in the intergenic regions of the Arabidopsis thaliana genome are
transcribed and/or under purifying selection. *Genome Res* **17**: 632–640.

Hanada K, Higuchi-Takeuchi M, Okamoto M, Yoshizumi T, Shimizu M, Nakaminami K, Nishi
R, Ohashi C, Iida K, Tanaka M, et al. 2013. Small open reading frames associated with
morphogenesis are hidden in plant genomes. *Proc Natl Acad Sci U S A* **110**: 2395–2400.

Hardiman KE, Brewster R, Khan SM, Deo M, Bodmer R. 2002. The bereft gene, a potential target of the
neural selector gene cut, contributes to bristle morphogenesis. *Genetics* **161**: 231–247.

Heinen TJAJ, Staubach F, Häming D, Tautz D. 2009. Emergence of a New Gene from an Intergenic
Region. *Curr Biol* **19**: 1527–1531.

Hsieh L-C, Lin S-I, Shih AC-C, Chen J-W, Lin W-Y, Tseng C-Y, Li W-H, Chiou T-J. 2009.
Uncovering small RNA-mediated responses to phosphate deficiency in Arabidopsis by deep
sequencing. *Plant Physiol* **151**: 2120–2132.

Ibarra-Laclette E, Lyons E, Hernández-Guzmán G, Pérez-Torres CA, Carretero-Paulet L, Chang T-H, Lan
T, Welch AJ, Juárez MJA, Simpson J, et al. 2013. Architecture and evolution of a minute plant
genome. *Nature* **498**: 94–98.

Ivanova N, Dobrin R, Lu R, Kotenko I, Levorse J, DeCoste C, Schafer X, Lun Y, Lemischka IR.
2006. Dissecting self-renewal in stem cells with RNA interference. *Nature* **442**: 533–538.

Johnson JM, Edwards S, Shoemaker D, Schadt EE. 2005. Dark matter in the genome: evidence
of widespread transcription detected by microarray tiling experiments. *Trends Genet* **21**: 93–
102.

Karreth FA, Reschke M, Ruocco A, Ng C, Chapuy B, Léopold V, Sjoberg M, Keane TM, Verma
A, Ala U, et al. 2015. The BRAF pseudogene functions as a competitive endogenous RNA
and induces lymphoma in vivo. *Cell* **161**: 319–332.

Kellis M, Wold B, Snyder MP, Bernstein BE, Kundaje A, Marinov GK, Ward LD, Birney E,
Crawford GE, Dekker J, et al. 2014. Defining functional DNA elements in the human
genome. *Proc Natl Acad Sci U S A* **111**: 6131–6138.

Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. 2013. TopHat2: accurate
alignment of transcriptomes in the presence of insertions, deletions and gene fusions.
Genome Biol **14**: R36.

Koehler R, Issac H, Cloonan N, Grimmond SM. 2011. The uniqueome: a mappability resource
for short-tag sequencing. *Bioinformatics* **27**: 272–274.

Kozomara A, Griffiths-Jones S. 2014. miRBase: annotating high confidence microRNAs using
deep sequencing data. *Nucleic Acids Res* **42**: D68–73.

Krishnakumar V, Hanlon MR, Contrino S, Ferlanti ES, Karamycheva S, Kim M, Rosen BD,
Cheng C-Y, Moreira W, Mock SA, et al. 2015. Araport: the Arabidopsis information portal.
Nucleic Acids Res **43**: D1003–9.

Krueger F, Andrews SR. 2011. Bismark: a flexible aligner and methylation caller for Bisulfite-
Seq applications. *Bioinformatics* **27**: 1571–1572.

Kuromori T, Wada T, Kamiya A, Yuguchi M, Yokouchi T, Imura Y, Takabe H, Sakurai T,

Akiyama K, Hirayama T, et al. 2006. A trial of phenome analysis using 4000 Ds-insertional

mutants in gene-coding regions of Arabidopsis. *Plant J* **47**: 640–651.

Lai K-MV, Gong G, Atanasio A, Rojas J, Quispe J, Posca J, White D, Huang M, Fedorova D,

Grant C, et al. 2015. Diverse Phenotypes and Specific Transcription Patterns in Twenty

Mouse Lines with Ablated LincRNAs. *PLoS One* **10**: e0125522.

Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment

of short DNA sequences to the human genome. *Genome Biol* **10**: R25.

Li F, Zheng Q, Vandivier LE, Willmann MR, Chen Y, Gregory BD. 2012a. Regulatory impact of

RNA secondary structure across the Arabidopsis transcriptome. *Plant Cell* **24**: 4346–4359.

Liu M-J, Seddon AE, Tsai ZT-Y, Major IT, Floer M, Howe GA, Shiu S-H. 2015. Determinants

of nucleosome positioning and their influence on plant gene expression. *Genome Res* **25**:

1182–1195.

Li W, Cui X, Meng Z, Huang X, Xie Q, Wu H, Jin H, Zhang D, Liang W. 2012b. Transcriptional

regulation of Arabidopsis MIR168a and argonaute1 homeostasis in abscisic acid and abiotic

stress responses. *Plant Physiol* **158**: 1279–1292.

Li WH, Gojobori T, Nei M. 1981. Pseudogenes as a paradigm of neutral evolution. *Nature* **292**:

237–239.

Lloyd J, Meinke D. 2012. A comprehensive dataset of genes with a loss-of-function mutant

phenotype in Arabidopsis. *Plant Physiol* **158**: 1115–1129.

Lloyd JP, Seddon AE, Moghe GD, Simenc MC, Shiu S-H. 2015. Characteristics of Plant

Essential Genes Allow for within- and between-Species Prediction of Lethal Mutant

Phenotypes. *Plant Cell* **27**: 2133–2147.

Marahrens Y, Panning B, Dausman J, Strauss W, Jaenisch R. 1997. Xist-deficient mice are defective in

dosage compensation but not spermatogenesis. *Genes Dev* **11**: 156–166.

Mattick JS. 2009. The Genetic Signatures of Noncoding RNAs. *PLoS Genet* **5**: e1000459.

Mercer TR, Dinger ME, Mattick JS. 2009. Long non-coding RNAs: insights into functions. *Nat Rev Genet* **10**: 155–159

Michael TP, Jackson S. 2013. The First 50 Plant Genomes. *Plant Genome* **6**: 0.

Moghe GD, Lehti-Shiu MD, Seddon AE, Yin S, Chen Y, Juntawong P, Brandizzi F, Bailey-Serres J, Shiu S-H. 2013. Characteristics and significance of intergenic polyadenylated RNA transcription in Arabidopsis. *Plant Physiol* **161**: 210–224.

Nagalakshmi U, Wang Z, Waern K, Shou C, Raha D, Gerstein M, Snyder M. 2008. The Transcriptional Landscape of the Yeast Genome Defined by RNA Sequencing. *Science* **320**: 1344–1349.

Neander K. 1991. Functions as Selected Effects: The Conceptual Analyst's Defense. *Philos Sci* **58**: 168–184.

Ngernprasirtsiri J, Kobayashi H, Akazawa T. 1988. DNA methylation as a mechanism of transcriptional regulation in nonphotosynthetic plastids in plant cells. *Proc Natl Acad Sci U S A* **85**: 4750–4754.

Ning S, Wang P, Ye J, Li X, Li R, Zhao Z, Huo X, Wang L, Li F, Li X. 2013. A global map for dissecting phenotypic variants in human lincRNAs. *Eur J Hum Genet* **21**: 1128–1133.

Niu D-K, Jiang L. 2013. Can ENCODE tell us how much junk DNA we carry in our genome? *Biochem Biophys Res Commun* **430**: 1340–1343.

Nobuta K, Venu RC, Lu C, Beló A, Vemaraju K, Kulkarni K, Wang W, Pillay M, Green PJ, Wang G-L, et al. 2007. An expression atlas of rice mRNAs and small RNAs. *Nat Biotechnol* **25**: 473–477.

O'Malley RC, Huang S-SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti A, Ecker JR. 2016. Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape. *Cell* **166**: 1598.

Pang KC, Frith MC, Mattick JS. 2006. Rapid evolution of noncoding RNAs: lack of conservation does not mean lack of function. *Trends Genet* **22**: 1–5.

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Prettenhofer P,
Weiss R, Dubourg V, et al. 2011. Scikit-learn: Machine Learning in Python. *J Mach Learn Res* **12**:
2825–2830.

Penny GD, Kay GF, Sheardown SA, Rastan S, Brockdorff N. 1996. Requirement for Xist in X
chromosome inactivation. *Nature* **379**: 131–137.

Poliseno L, Salmena L, Zhang J, Carver B, Haveman WJ, Pandolfi PP. 2010. A coding-
independent function of gene and pseudogene mRNAs regulates tumour biology. *Nature*
465: 1033–1038.

Ponting CP, Belgard TG. 2010. Transcribed dark matter: meaning or myth? *Hum Mol Genet* **19**:
R162–R168.

Sauvageau M, Goff LA, Lodato S, Bonev B, Groff AF, Gerhardinger C, Sanchez-Gomez DB,
Hacisuleyman E, Li E, Spence M, et al. 2013. Multiple knockout mouse models reveal
lincRNAs are required for life and brain development. *Elife* **2**: e01749.

Savage LJ, Imre KM, Hall DA, Last RL. 2013. Analysis of essential Arabidopsis nuclear genes
encoding plastid-targeted proteins. *PLoS One* **8**: e73291.

Schnable JC, Pedersen BS, Subramaniam S, Freeling M. 2011. Dose–Sensitivity, Conserved
Non-Coding Sequences, and Duplicate Gene Retention Through Multiple Tetraploidies in
the Grasses. *Front Plant Sci* **2**. http://dx.doi.org/10.3389/fpls.2011.00002.

Schnable JC, Wang X, Pires JC, Freeling M. 2012. Escape from preferential retention following
repeated whole genome duplications in plants. *Front Plant Sci* **3**: 94.

Schreiber SL, Bernstein BE. 2002. Signaling Network Model of Chromatin. *Cell* **111**: 771–778

Shin H, Shin H-S, Chen R, Harrison MJ. 2006. Loss of At4 function impacts phosphate
distribution between the roots and the shoots during phosphate starvation. *Plant J* **45**: 712–
726.

Sing T, Sander O, Beerenwinkel N, Lengauer T. 2005. ROCR: visualizing classifier performance
in R. *Bioinformatics* **21**: 3940–3941.

Slotkin RK, Keith Slotkin R, Martienssen R. 2007. Transposable elements and the epigenetic

regulation of the genome. *Nat Rev Genet* **8**: 272–285.

Soltis DE, Albert VA, Leebens-Mack J, Bell CD, Paterson AH, Zheng C, Sankoff D,

Depamphilis CW, Wall PK, Soltis PS. 2009. Polyploidy and angiosperm diversification. *Am*

J Bot **96**: 336–348.

Stergachis AB, Haugen E, Shafer A, Fu W, Vernot B, Reynolds A, Raubitschek A, Ziegler S,

LeProust EM, Akey JM, et al. 2013. Exonic Transcription Factor Binding Directs Codon

Choice and Affects Protein Evolution. *Science* **342**: 1367–1372.

Stolc V, Samanta MP, Tongprasit W, Sethi H, Liang S, Nelson DC, Hegeman A, Nelson C,

Rancour D, Bednarek S, et al. 2005. Identification of transcribed sequences in Arabidopsis

thaliana by using high-resolution genome tiling arrays. *Proc Natl Acad Sci U S A* **102**:

4453–4458.

Strobl C, Boulesteix A-L, Kneib T, Augustin T, Zeileis A. 2008. Conditional variable importance for

random forests. *BMC Bioinformatics* **9**: 307

Struhl K. 2007. Transcriptional noise and the fidelity of initiation by RNA polymerase II. *Nat*

Struct Mol Biol **14**: 103–105.

Sullivan AM, Arsovski AA, Lempe J, Bubb KL, Weirauch MT, Sabo PJ, Sandstrom R, Thurman

RE, Neph S, Reynolds AP, et al. 2014. Mapping and dynamics of regulatory DNA and

transcription factor networks in A. thaliana. *Cell Rep* **8**: 2015–2030.

Svensson O, Arvestad L, Lagergren J. 2006. Genome-wide survey for biologically functional

pseudogenes. *PLoS Comput Biol* **2**: e46.

Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold

BJ, Pachter L. 2010. Transcript assembly and quantification by RNA-Seq reveals

unannotated transcripts and isoform switching during cell differentiation. *Nat Biotechnol* **28**:

511–515.

Tsai ZT-Y, Shiu S-H, Tsai H-K. 2015. Contribution of Sequence Motif, Chromatin State, and

DNA Structure Features to Predictive Models of Transcription Factor Binding in Yeast.
PLoS Comput Biol **11**: e1004418.

Tsai ZT-Y, Lloyd JP, Shiu S-H. 2017. Defining functional, genic regions in the human genome
through integration of biochemical, evolutionary, and genetic evidence. *Mol Biol Evol* **In**
press

Ulitsky I, Shkumatava A, Jan CH, Sive H, Bartel DP. 2011. Conserved function of lincRNAs in
vertebrate embryonic development despite rapid sequence evolution. *Cell* **147**: 1537–1550.

VanBuren R, Bryant D, Edger PP, Tang H, Burgess D, Challabathula D, Spittle K, Hall R, Gu J,
Lyons E, et al. 2015. Single-molecule sequencing of the desiccation-tolerant grass
Oropetium thomaeum. *Nature* **527**: 508–511.

Wang Z, Zang C, Rosenfeld JA, Schones DE, Barski A, Cuddapah S, Cui K, Roh T-Y, Peng W,
Zhang MQ, et al. 2008. Combinatorial patterns of histone acetylations and methylations in
the human genome. *Nat Genet* **40**: 897–903.

Woodhouse MR, Schnable JC, Pedersen BS, Lyons E, Lisch D, Subramaniam S, Freeling M.
2010. Following Tetraploidy in Maize, a Short Deletion Mechanism Removed Genes
Preferentially from One of the Two Homeologs. *PLoS Biol* **8**: e1000409.

Xu S, Grullon S, Ge K, Peng W. 2014. Spatial clustering for identification of ChIP-enriched
regions (SICER) to map regions of histone methylation patterns in embryonic stem cells.
Methods Mol Biol **1150**: 97–111.

Yamada K, Lim J, Dale JM, Chen H, Shinn P, Palm CJ, Southwick AM, Wu HC, Kim C,
Nguyen M, et al. 2003. Empirical analysis of transcriptional activity in the Arabidopsis
genome. *Science* **302**: 842–846.

Yang L, Takuno S, Waters ER, Gaut BS. 2011. Lowly expressed genes in Arabidopsis thaliana
bear the signature of possible pseudogenization by promoter degradation. *Mol Biol Evol* **28**:
1193–1203.

Zhang X, Yazaki J, Sundaresan A, Cokus S, Chan SW-L, Chen H, Henderson IR, Shinn P,

Pellegrini M, Jacobsen SE, et al. 2006. Genome-wide high-resolution mapping and

functional analysis of DNA methylation in arabidopsis. *Cell* **126**: 1189–1201.

Zhao J, Sun BK, Erwin JA, Song J-J, Lee JT. 2008. Polycomb proteins targeted by a short repeat

RNA to the mouse X chromosome. *Science* **322**: 750–756.

Zhao Y, Li H, Fang S, Kang Y, Wu W, Hao Y, Li Z, Bu D, Sun N, Zhang MQ, et al. 2016.

NONCODE 2016: an informative and valuable data source of long non-coding RNAs.

Nucleic Acids Res **44**: D203–8.

Zou C, Lehti-Shiu MD, Thibaud-Nissen F, Prakash T, Buell CR, Shiu S-H. 2009. Evolutionary

and expression signatures of pseudogenes in Arabidopsis and rice. *Plant Physiol* **151**: 3–15.

1124

1125 **FIGURES**

Figure 1.

1126

1127

Figure 3.

Figure 4.

Figure 5.

1133

1134

Figure 6.

1135

1136

 SUPPLEMENTAL FIGURES

**Supplemental
Figure 1.**

1138

1139

**Supplemental
Figure 2.**

1140

1141

Supplemental Figure 3.

1142

1143

**Supplemental
Figure 4.**

1144

1145

**Supplemental
Figure 5.**

1146

1147