

***De novo* transcriptome sequencing and comparative analysis of midgut tissues of four non-model insects pertaining to Hemiptera, Coleoptera, Diptera and Lepidoptera**

Rajesh Kumar Gazara¹, Christiane Cardoso², Daniel Bellieny-Rabelo¹, Clélia Ferreira²,
Walter R. Terra^{2,*} and Thiago Motta Venancio^{1,*}

¹ Laboratório de Química e Função de Proteínas e Peptídeos, Centro de Bociências e Biotecnologia, Universidade Estadual do Norte Fluminense Darcy Ribeiro; Campos dos Goytacazes, Brazil. ² Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, São Paulo, Brazil.

* Corresponding authors

warterra@iq.usp.br

thiago.venancio@gmail.com

ABSTRACT

Despite the great morphological diversity of insects, there is a regularity in their digestive functions, which is apparently related to their physiology. In the present work we report the *de novo* midgut transcriptomes of four non-model insects from four distinct orders: *Spodoptera frugiperda* (Lepidoptera), *Musca domestica* (Diptera), *Tenebrio molitor* (Coleoptera) and *Dysdercus peruvianus* (Hemiptera). We employed a computational strategy to merge assemblies from two different algorithms, which substantially increased the quality of the assembled transcriptomes. Unigenes were annotated and analyzed using the eggNOG database, which allowed us to assign some level of functional and evolutionary information to 79.7% to 93.1% of the transcriptomes. We found interesting transcriptional patterns, such as: i) the intense use of lysozymes in digestive functions of *M. domestica* larvae, which are streamlined and adapted to feed on bacteria; ii) the up-regulation of orthologous UDP-glycosyl transferase and cytochrome P450 genes in the whole midguts different species, supporting the presence of an ancient defense frontline to counter xenobiotics; iii) evidence supporting roles for juvenile hormone binding proteins in the midgut physiology, probably as a way to activate genes that help fight anti-nutritional substances (e.g. protease inhibitors). The results presented here shed light on the digestive and structural properties of the digestive systems of these distantly related species. Furthermore, the produced datasets will also be useful for scientists studying these insects.

Keywords: insects; transcriptomes; whole midgut; anterior midgut; xenobiotics; juvenile hormone

INTRODUCTION

Although insects can be extremely variable in morphology, there is a clear regularity in their digestive functions, which is apparently related to their physiology (Terra and Ferreira, 2012, Terra, 1990). In this regard, the availability of midgut transcriptomes obtained by second-generation sequencing offers a new way to evaluate the differences and similarities of the midgut physiology across divergent groups of insects. With this aim, the midgut transcriptomes of 4 insect species pertaining to 4 different orders were compared here. We provide below a summary of the life styles and available physiological data of the chosen insects, as a basis for discussing results obtained from the transcriptome data.

The yellow meal worm, *Tenebrio molitor*, is a pest of stored products. The *T. molitor* midgut is buffered at pH 5.6 at the two anterior thirds and at pH 7.9 at the posterior midgut (PM). Carbohydrate digestion occurs largely at the anterior midgut (AM), whereas proteins are continuously digested by cysteine proteases at the AM and later by serine proteases located at the lumen and surface of PM cells. A counter current flux of water, powered by secretion of fluid at the posterior midgut and absorption at the anterior midgut, recovers digestive enzymes from inside the peritrophic membrane (i.e. an acellular chitin-protein membrane surrounding the food inside midgut) before excretion (Terra et al., 1985, Cristofolletti et al., 2001).

The fall armyworm, *Spodoptera frugiperda*, is a pest of several crops, including corn. Digestion in *S. frugiperda* initially occurs in a highly alkaline midgut, mainly by amylases and serine proteases inside the peritrophic membrane. Intermediate and final digestive steps occur via the action of enzymes located between the peritrophic membrane and midgut cells and at the surface of midgut cells. A counter current flux of fluid, similar to that described above in *T. molitor*, is also present in *S. frugiperda* (Ferreira et al., 1994a, Ferreira et al., 1994b).

The peculiarities displayed by *Musca domestica* larvae (as well by other *Cyclorrhaphous* flies) in regard to their digestive physiology are related to the process of killing and digesting bacteria (Espinoza-Fuentes and Terra, 1987). The midgut of *M. domestica* larvae can be divided in three parts. The middle midgut cells are thought to have an embryonic origin distinct from the AM and PM cells (Poulson and Waterhouse, 1960). The *M. domestica* middle midgut is very acidic, which is probably an adaptation to kill bacteria, the major food source of *M. domestica*

larvae. Further, AM amylases deplete the starch content of the meal, making bacteria more susceptible to low pH, digestive lysozymes and cathepsins D of the middle midgut. The nutrients from the killed bacteria pass to the PM, where most of the digestion takes place. Therefore, in a sense, the *M. domestica* PM corresponds to the entire midgut of the majority of the other insect larvae (Espinoza-Fuentes and Terra, 1987).

The midgut of the cotton-stainer, *Dysdercus peruvianus*, is divided into three sections (V1-V3), which are linked to the hindgut by V4. *D. peruvianus* lacks a chitin-protein peritrophic membrane and have instead a lipoprotein perimicrovillar membrane that ensheathes the midgut cells microvilli as glove fingers. The midgut is acidic, with a pH of 5.8. V1 is the site of carbohydrate digestion and absorption of water and glucose. V2 and V3 regions are responsible for protein digestion (mainly by cathepsin L, a cysteine protease) and amino acid absorption (Silva and Terra, 1994).

In the present work we report *de novo* transcriptome sequencing and analysis of midgut tissues of *S. frugiperda* (Lepidoptera), *M. domestica* (Diptera), *T. molitor* (Coleoptera) and *D. peruvianus* (Hemiptera). We used a combined assembly approach to recover longer transcripts, which were annotated with a step-wise pipeline using different databases. This strategy allowed the annotation of a large fraction of the contigs. The sequencing reads were further used to estimate the transcriptional levels of the contigs. The integration of functional annotations and transcriptional levels allowed us to find common and divergent aspects in the digestive processes of these four distantly related species, as well as a number of differentially expressed genes and pathways that are critical for midgut physiology and biochemistry.

RESULTS AND DISCUSSION

De novo transcriptome assembly and annotation

A total 609,407, 677,080, 608,637 and 747,904 reads from *D. peruvianus*, *T. molitor*, *M. domestica* and *S. frugiperda*, respectively, were assembled using a combined assembly approach (Table 1, see methods for details). Initial assemblies for all species were obtained by merging Newbler 2.5 and MIRA assemblies, as previously described (Kumar and Blaxter, 2010). Newbler assembly resulted in 8,093, 12,370, 5,329 and 10,315 contigs of *D. peruvianus*, *T.*

molitor, *M. domestica* and *S. frugiperda*, respectively (Table 2). MIRA assemblies resulted in 27,395, 37,703, 20,589, and 28,308 contigs for *D. peruvianus*, *T. molitor*, *M. domestica* and *S. frugiperda*, respectively (Table 2). After combining the assemblies for each species using CAP3, we obtained 6,774, 9,583, 4,248 and 7,264 longer and less redundant contigs, for *D. peruvianus*, *T. molitor*, *M. domestica* and *S. frugiperda*, respectively (Table 2; Supplementary figure 1). After filtering bacterial and rRNA contaminants, we obtained 6,395, 9,010, 4,005 and 6,833 of unigenes for *D. peruvianus*, *T. molitor*, *M. domestica* and *S. frugiperda* (Table S1). The final CAP3 transcriptome assemblies can be found on supplementary file 1.

EggNOG (Powell et al., 2014) is a database of orthologous groups (OGs) that have general functional annotation codes that are largely derived from the COG database (Tatusov et al., 1997). A stepwise function assignment strategy (see methods for details) allowed us to assign eggNOG function codes for 2970, 5764, 2506 and 3627 genes from *D. peruvianus*, *T. molitor*, *M. domestica* and *S. frugiperda*, respectively (Table 3; Table S2). Through BLASTX against uniref90, 1809, 2028, 716 and 1331 additional unigenes were annotated for *D. peruvianus*, *T. molitor*, *M. domestica* and *S. frugiperda*, respectively (Table S2). Finally, the remaining unigenes had their longest ORFs extracted and scanned for conserved PFAM domains, resulting in 317, 595, 299 and 533 additional annotated unigenes for *D. peruvianus*, *T. molitor*, *M. domestica* and *S. frugiperda*, respectively (Table S2). In total, we were able to assign some level of functional or evolutionary information to 79.7% to 93.1% of the *de novo* transcriptomes reported here (Table 4).

Differential gene expression and conservation across species

Next, we further explored the transcriptome datasets reported here to better understand the transcriptional programs of midgut tissues. We compared the expression levels of all genes between all possible tissue combinations in each species; PM vs CAR, PM vs WM, WM vs CAR for *D. peruvianus*; for the remaining species, the following comparisons were performed: AM vs CAR, AM vs WM and WM vs CAR (Figure 1; Table S4). We found a significantly lower number of DEGs between different midgut compartments than when these samples were compared with CAR, which reflect the physiological and anatomical similarities of

these midgut compartments. Notably, in *D. peruvianus* there is a smaller number of induced genes in midgut tissues (Figure 1), probably due to the simplification of the midgut of *D. peruvianus* ancestors (insects similar to extant cicads) resulting from their adaption to feed on plant sap, which is generally poor in macromolecules that demand digestion (Terra, 1990). Although other hemipterans regained a macromolecular feeding habit, they did this by deploying lysosomal enzymes instead of the serine proteases that were previously lost (Houseman et al., 1985). A similar trend has been observed in *M. domestica*, and is in accordance with the streamlined digestive apparatus of the larvae, which is adapted to digest bacterial cells.

By aggregating eggNOG family and function annotations to assembled unigenes, we were able to perform functional enrichment analyses to explore the global trends underlying the transcriptomes reported here. In *D. peruvianus*, there is a significant up-regulation of lipases and two endopeptidase families in WM when compared to CAR (Table S4; Table S5). These peptidase families encode C1 and D cathepsin family endopeptidases. Interestingly, this latter gene family is also significantly up-regulated in WM when compared to PM, suggesting that its expression is biased towards the anterior parts of the *D. peruvianus* midgut. It has been hypothesized that up-regulation of cathepsin L (cysteine proteases) and cathepsin D (aspartic proteases) genes are associated with their use as digestive enzymes, replacing lost digestive serine peptidases (Silva and Terra, 1994). Alternatively, it is also possible that those genes were actually not lost, but divergent in function. This is supported by the down-regulation of three families of serine peptidases (i.e. ENOG4103A6W, ENOG41032G6 and ENOG4103814) in WM (vs CAR). Further, the preferential up-regulation of cathepsin D in AM is probably an adaptation to remove cathepsin L inhibitors present in the cotton seeds, favoring cathepsin L as the main digestive enzyme in *D. peruvianus* PM (Pimentel et al., 2017). An opposite trend is observed in lipases, which were more expressed in PM than in WM samples (Table S4; Table S5), suggesting a more intense lipid digestion in the posterior parts of the *D. peruvianus* midgut. The up-regulation of lipases could be an adaptation to digest the large oil amounts that are sucked from the cotton seeds (Bewley et al., 2006). Finally, a family of genes encoding proteins with a domain of unknown function (DUF3421) is also enriched among genes up-regulated in *D.*

peruvianus WM (ENOG410382G). Although the functions of these genes are largely unclear, the two *Drosophila melanogaster* homologs from this family, CG13321 and CG32633, are highly expressed in embryos/larvae and digestive systems samples, respectively (modENCODE Anatomy and Development RNA-Seq Atlas, Flybase, accessed on Jan 2017). In addition, at least one member of this family appears as up-regulated in WM (vs CAR) in all four species analyzed here (Table 5), further emphasizing its importance in the insect midguts.

In *T. molitor* we found a total of 11 and 2 enriched families among up-regulated DEGs from WM vs CAR and AM vs WM comparisons, respectively (Table S5). These families comprise a number of up-regulated genes involved in digestive and structural functions, such as chitinases, beta-galactosidases, mannosidases, serine-proteases, chitin-binding proteins and peritrophins (Table S4; Table S5). Further, our results suggest that serine protease activity is more constrained to the PM, as these genes are down-regulated in the AM when compared to WM (Table S5). Importantly, the preferential expression of cysteine (e.g. cathepsin L) and serine proteases (e.g. trypsin and chymotrypsin) in the AM and PM of *T. molitor*, respectively, were previously described by direct enzymatic assays (Terra and Cristofolletti, 1996). This strong up-regulation allowed the isolation and characterization many enzymes, such as alpha-mannosidases (Moreira et al., 2015), chitinases (Genta et al., 2006), β -1,3-glucanases (Genta et al., 2009), β -glycosidases (Ferreira et al., 2001), β -galactosidases (Ferreira et al., 2003), serine proteases (Cristofolletti et al., 2001) and chymotrypsins (Sato et al., 2008). Furthermore, we found 5 chitin deacetylase-like genes (family ENOG41033EF) that are up-regulated in *T. molitor* WM when compared with CAR, which probably play key roles in determining the peritrophic membrane structure or permeability (Muthukrishnan et al., 2012). Interestingly, a *D. melanogaster* homolog of this gene (CDA9) is also highly expressed in larvae (modENCODE Anatomy and Development RNA-Seq Atlas, Flybase, accessed on Jan 2017).

In *M. domestica*, we also found a general trend for the up-regulation of genes involved in digestive functions in WM and AM. Further, some of these enzymes have been characterized, expressed as recombinant proteins and had their 3D structure resolved (Lemos et al., 1993, Cançado et al., 2007). This list comprises, for example, several genes from a lysozyme family (ENOG4103B29) associated with digestion (Table S5). Interestingly, the gene family encoding

DUF3421 proteins, up-regulated in *D. peruvianus* WM, is also up-regulated in *M. domestica* WM (vs CAR). This conservation in expression patterns between distant species indicates that this gene family is critical in midgut tissues, thus warranting further biochemical investigations.

Finally, we found a number of enriched gene families encoding endopeptidases, carbohydrate hydrolases and lipases, among others, as preferentially up-regulated in *S. frugiperda* WM when compared with CAR. Interestingly, three of these families (i.e. ENOG4103AQE, ENOG4103CXY and ENOG41036EN) are biased towards AM regions of the midgut (Table S5). ENOG4103AQE encodes peritrophin-A chitin binding proteins that have crucial roles in larvae of many insects, including the mosquito *A. aegypti* (Venancio et al., 2009). ENOG4103CXY comprises genes encoding esterases with the alpha/beta hydrolase fold, which are involved in lipid metabolism. Further, this family contains only genes from lepidopterans, including the silkworm *Bombyx mori* and the postman butterfly *Heliconius melpomene*. Finally, we found three genes encoding prostaglandin reductases (ENOG41036EN family) up-regulated in *S. frugiperda* AM, which might be involved in prostaglandin signaling at this compartment.

Regarding the conservation of up-regulated genes across different species, we also found some interesting trends, such as: 1) the absence of families with up-regulated members in AM (vs WM) of all species, which is probably due to different factors, such as the diversification of this tissue across species (*M. domestica*, see above) and the similarity between the transcriptional programs of WM and AM. Importantly, we were able to detect DEGs in AM, suggesting that this result is not merely due to inadequate tissue dissection or absence of differential expression throughout the midgut; 2) there are 5 gene families with at least one up-regulated member in the WM of each species (Table 5). These families encode some interesting functions, such as the haemolymph juvenile hormone (JH) binding protein (JHBP, family ENOG41039KJ, discussed in the next section). Other interesting family is ENOG410362J, which encodes orthologs of the *Drosophila* MESH gene (FBgn0051004). In *D. melanogaster*, MESH is highly expressed after 12hr of embryo development is essential for the development of 1st instar larvae, probably due to its crucial roles in septate junction formation (Izumi et al., 2012). Thus, the up-regulation of MESH genes in such distantly related species reveal the importance of this ancient structural component in the development of the larval

midgut; 3) the up-regulation of orthologous families involved in the catabolism of xenobiotics across species. One notable example is a family of UDP-glucuronosyl and UDP-glucosyl transferases [ENOG41033B3; PFAM PF00201 (UDPGT)], with at least one gene up-regulated in the WM of *D. peruvianus*, *S. frugiperda* and *T. molitor*. This domain is present on a relatively big family of enzymes involved in cuticle formation, pigmentation, olfaction and detoxification of xenobiotics. This latter function is of particular importance to WM, the primary site of digestion that has constant contact with deleterious substances (Heidel-Fischer and Vogel, 2015). Interestingly, in *D. melanogaster*, there are 10 UDPGT genes (8 from the ENOG41033B3 family) in a cluster of ~51.5Kb at chromosome 3R (from 11,126,480 to 11,178,000), strongly supporting the expansion of this family via tandem gene duplication. Furthermore, a family of cytochrome P450 genes (ENOG41035SH), known to be involved in xenobiotic detoxification (Heidel-Fischer and Vogel, 2015), is also up-regulated in WM of *D. peruvianus* and *S. frugiperda*, supporting the presence of an ancient defense program that is activated in the midgut of distantly related species.

Conservation of the top highly expressed genes

Next, we analyzed the conservation patterns of the most highly expressed genes in each tissue in the 4 species considered here. In spite of the evolutionary distance between these species, we found in WM and CAR, respectively, a total of 35 and 36 gene families with at least one conserved member in each species among the top 10% most highly expressed genes (Figure 2 and Tables 6 and S6). Among the 35 families with highly expressed genes in WM, 22 also have highly expressed members in CAR and are probably housekeeping genes. The remaining 13 could have more specific roles in the digestive tract. These families cover some interesting functions, such as peptidoglycan recognition proteins (ENOG41032AN), which is involved in innate immunity of the moth *Trichoplusia ni* (Kang et al., 1998). CG9380, a *D. melanogaster* homolog from this family, displays high expression levels in larval and adult digestive systems (modENCODE Anatomy and Development RNA-Seq Atlas, Flybase, accessed on Jan 2017). Interestingly, no genes from this family are among the top 10% expressed genes in AM, indicating that their functions could be more relevant in more posterior midgut sections.

Another interesting gene family showing biased transcription in midgut of several species is ENOG41039KJ, which encodes JHBPs (mentioned above), which protect the JH molecules from hydrolysis in the hemolymph. Furthermore, JH and JHBP can help to counter the effects of anti-nutritional substances (e.g. protease inhibitors) via up-regulation of midgut proteases in the migratory locust *Locusta migratoria* (Spit et al., 2016). JH/JHPB has also been demonstrated to induce the expression of proteases in other insects, such as *Helicoverpa armigera* (Sui et al., 2009), *A. aegypti* (Sui et al., 2009) and *Hodotermopsis sjostedti* (Cornette et al., 2013). Finally, we have also found a family of serine carboxypeptidases (i.e. ENOG410351F) in this dataset. This is interesting because most digestive carboxypeptidases in these insects (except for *D. peruvianus*, which does not have digestive metallo-carboxypeptidases) are metallo-carboxypeptidases, whereas the serine carboxypeptidases are mainly lysosomal (Ferreira et al., 2015). This result suggests that the recruitment of lysosomal carboxypeptidases predates the divergence of these insect groups, although more analyses are warranted to rule out the possibility of independent events after radiation.

There are 67 gene families with highly expressed members in the AM of *M. domestica*, *S. frugiperda* and *T. molitor* (AM of *D. peruvianus* was not studied) (Figure 2 and Table 7). Although most of these families encode genes that are also highly expressed in carcass, some of them are obviously related to digestive function such as: alpha-amylase, aminopeptidase, alkaline phosphatase, serine-type carboxypeptidase, cathepsin, glucosylceramidase, serine endopeptidase, and serpins (Table 7). This is probably due to the fact that carcass includes the Malpighian tubules, which are known to harbor several enzymes and transporters similar to those found in the midgut (Beyenbach et al., 2010). The same phenomenon occurs with kidneys and intestine in mammals (Boron and Boulpaep, 2005). Further, among the 10% most expressed genes, *T. molitor* and *S. frugiperda* share more OGs with one another than the latter with *M. domestica* (Figure 2). Although unexpected given the phylogenetic relationships of these species, this result can be explained by the adaptation of *M. domestica* to digest bacteria (see Introduction), which largely remodeled its digestive enzymatic apparatus. Most of these 67 gene families encode enzymes involved in primary metabolism and ribosomal proteins. However, important regulatory gene families are also present. A notable example is the

ENOG41038D7 family of bZIP transcription factors, which includes the *D. melanogaster* ATF4 (CRC, *cryptocephal*). CRC is involved in critical developmental processes, such as larval molting and pupariation (Hewes et al., 2000), probably in a pathway involving the transcriptional regulation of the Ecdysis triggering hormone (ETH) in endocrine cells (Gauthier and Hewes, 2006). Interestingly, ATF4 is also involved in endoplasmic reticulum stress response through the regulation of a metabolic shift involving the up-regulation of genes encoding glycolytic enzymes (Lee et al., 2015). Accordingly, other genes encoding glycolytic enzymes are also present in the set of 67 gene families mentioned above (i.e. ENOG41032XF: Triose-phosphate isomerase; ENOG41035ZK: Glyceraldehyde-3-phosphate dehydrogenase). Therefore, important parts of this system are present in high levels in the AM of different species, suggesting that the regulatory mechanism outlined above play important roles in the development of this tissue.

CONCLUSIONS

In the present work we described the *de novo* sequencing and assembly of transcriptomes of distantly-related, non-model insects of agronomical and medical importance. We used a computational approach to merge results from different algorithms and integrated our data with the eggNOG database, which helped us to annotate and analyze the evolution of several gene families of interest. We believe that the results provided here are not only important to shed light on fundamental biochemical aspects underlying the digestive process of the species in question, but also contribute with a large dataset that can be further used by the scientific community to design novel experiments to study genes of interest in various fields.

MATERIALS AND METHODS

Insects

S. frugiperda (Lepidoptera) were laboratory-reared, as previously described (Parra, 1986). *S. frugiperda* larvae were kept in glass vials with the diet at a temperature of 25°C and 14:10 photoperiod (photophase: scotophase). The diet for *S. frugiperda* was composed of kidney bean "carioca" (*Phaseolus vulgaris*), wheat germ, ascorbic acid, yeast, agar and microbial inhibitors. Adults were fed a 10% honey solution. Stock cultures of *D. peruvianus* (Hemiptera)

were reared in glass bottles covered with a piece of cloth, under natural photoperiod conditions at a relative humidity of 50-70% at $24 \pm 2^\circ\text{C}$. Insects had access to water and cotton (*Gossypium hirsutum*) seeds that were previously disinfested by freezing. Adults of *D. peruvianus* were used, as this is the stage of more intense feeding. Larvae of *M. domestica* (Diptera) were reared in a mixture of fermented commercial pig food and rice hull (1:2, v/v) (Targa and Peres, 1979). The larvae used in this study were actively feeding individuals at larval third instar. *T. molitor* (Coleoptera), were bred on wheat bran at 24-26 °C and relative humidity of 70-75%. Fully-grown larvae (each weighing about 0.12 g) of both sexes were used.

Sample preparation and sequencing

Larvae from *S. frugiperda*, *M. domestica*, and *T. molitor* and adults of *D. peruvianus* were immobilized on ice and dissected with gloves, sterile forceps and glassware treated with diethyl pyrocarbonate. Whole midgut (WM) and carcass (larval body without midgut, CAR) were separated. The anterior third of the midgut (i.e. AM) of *S. frugiperda*, *T. molitor*, and *M. domestica* and the PM (three last midgut sections, V2+V3+V4) of *D. peruvianus* were used. Dissected tissues were kept for a short period in an ethanol-ice bath and stored at -80°C . RNA was extracted using Trizol (Invitrogen). The mRNA was isolated from total RNA with Dynabeads mRNA purification kit (Life technologies). Total RNA and mRNA quality were checked with an Agilent 2100 Bioanalyzer instrument. cDNA libraries were constructed following Roche recommendations and sequenced on a 454 Genome Sequencer FLX (454 Life Sciences/Roche) at the sequencing facility of the Instituto de Química, University of São Paulo, Brazil.

De novo transcriptome assembly

The 454 reads were extracted from SFF files using the *sff_extract* program (https://bioinf.comav.upv.es/seq_crumbs/). The leading 15 bp of each read were trimmed. The 3' end of each read was also trimmed to keep reads at a maximum length of 500 bp. Reads with less than 100 bp were discarded. *De novo* assemblies were performed using Newbler (version 2.5) (Margulies et al., 2005), MIRA (version 4.0.2) (Chevreux, 2004) and CAP3 (Huang and Madan, 1999). It has been demonstrated that significantly improved *de novo* assemblies of 454

reads can be achieved by combining assemblies obtained with different tools. In particular, merging of MIRA and Newbler assemblies by CAP3 gave the best results (Kumar and Blaxter, 2010). Different 454 libraries of the same species were combined before the assembly step to improve coverage. In general, each CAP3 contig represents most (if not all) splicing isoforms of a gene; therefore, in the present work we use the term “unigenes” to refer to CAP3 contigs. We followed this strategy and assembled our data with Newbler and MIRA; the resulting assemblies were filtered to retain only contigs with at least 100 bp and combined with CAP3 (MIRA and Newbler contigs are treated pseudo-reads for the CAP3 assembly).

Transcriptome annotation

A stepwise strategy was used to functionally annotate genes as follows: 1) bi-directional BLASTX (Altschul et al., 1997) hit (BBH); 2) Best unidirectional BLASTX hit; steps 1 and 2 were performed using the eggNOG v4.0 arthropod database (Powell *et al.* 2014), with hit coverage of at least 50% of the shortest protein of the pair and e-value < 1e-3. Furthermore, contigs were assigned to eggNOG orthologous groups (OGs) using the results from steps 1 and 2; 3) unigenes not annotated in steps 1 and 2 were submitted to a BLASTX search against the Uniref90 database (e-value < 1e-3) (Suzek et al., 2007); 4) the remaining unigenes had their longest ORFs extracted with TransDecoder (<http://transdecoder.sf.net>) and analyzed with HMMER3 (e-value < 1e-3) (Eddy, 2011) searches against the PFAM database (Finn et al., 2014). Unigenes were filtered for bacterial contamination by using BLASTX (Altschul et al., 1997) against bacterial proteins from the eggNOG v4.0 database (Powell et al., 2014)(E-value < 1e-10). In addition, rRNAs contaminants were removed using BLASTN (E-value < 1e-10) against all Genbank (Clark et al., 2016) insect rRNA sequences (downloaded on June 26, 2015).

Differential expression analysis and functional enrichment

To estimate transcriptional levels, reads were mapped back to the respective transcriptome assemblies using BWA (Li and Durbin, 2009). Uniquely mapped reads were normalized by the “reads per kilobase per million mapped reads” method (RPKM). Differentially expressed genes (DEGs) between tissues of the same species (i.e. AM, PM, WM and CAR) were

inferred using edgeR (Robinson et al., 2010). Genes with p-value ≤ 0.001 and \log_2 (fold change) (\log_2FC) ≥ 2 or ≤ -2 were considered up- and down-regulated, respectively. Enriched (i.e. over-represented) eggNOG functions in the DEG sets were analyzed using the Fished Exact Test (p-value ≤ 0.05 ; Bonferroni-corrected) using R (www.r-project.org).

ACKNOWLEDGEMENTS

This work was supported by Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES).

REFERENCES

- ALTSCHUL, S. F., MADDEN, T. L., SCHÄFFER, A. A., ZHANG, J., ZHANG, Z., MILLER, W. & LIPMAN, D. J. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic acids research*, 25, 3389-402.
- BEWLEY, D. J., BLACK, M. & HALMER, P. 2006. Encyclopedia of seeds: science, technology and uses.
- BEYENBACH, K. W., SKAER, H. & DOW, J. A. T. 2010. The Developmental, Molecular, and Transport Biology of Malpighian Tubules. *Annu. Rev. Entomol*, 55, 351-74.
- BORON, W. F. & BOULPAEP, E. L. 2005. Medical Physiology. A cellular and molecular approach . Updated edition.
- CANÇADO, F. C., VALÉRIO, A. A., MARANA, S. R. & BARBOSA, J. A. R. G. 2007. The crystal structure of a lysozyme c from housefly *Musca domestica*, the first structure of a digestive lysozyme. *Journal of Structural Biology*, 160, 83-92.
- CHEVREUX, B. 2004. Using the miraEST Assembler for Reliable and Automated mRNA Transcript Assembly and SNP Detection in Sequenced ESTs. *Genome Research*, 14, 1147-1159.
- CLARK, K., KARSCH-MIZRACHI, I., LIPMAN, D. J., OSTELL, J. & SAYERS, E. W. 2016. GenBank. *Nucleic Acids Res*, 44, D67-72.
- CORNETTE, R., HAYASHI, Y., KOSHIKAWA, S. & MIURA, T. 2013. Differential gene expression in response to juvenile hormone analog treatment in the damp-wood termite *Hodotermopsis sjostedti* (Isoptera, Archotermopsidae). *Journal of Insect Physiology*, 59, 509-518.
- CRISTOFOLETTI, P. T., RIBEIRO, A. F. & TERRA, W. R. 2001. Apocrine secretion of amylase and exocytosis of trypsin along the midgut of *Tenebrio molitor* larvae. 47, 143-155.
- EDDY, S. R. 2011. Accelerated Profile HMM Searches. *PLoS Computational Biology*, 7, e1002195.
- ESPINOZA-FUENTES, F. P. & TERRA, W. R. 1987. Physiological adaptations for digesting bacteria. Water fluxes and distribution of digestive enzymes in *Musca domestica* larval midgut. *Insect Biochemistry*, 17, 809-817.
- FERREIRA, A. H. P., MARANA, S. R., TERRA, W. R. & FERREIRA, C. 2001. Purification, molecular cloning, and properties of a β -glycosidase isolated from midgut lumen of *Tenebrio molitor* (Coleoptera) larvae. *Insect Biochemistry and Molecular Biology*, 31, 1065-1076.
- FERREIRA, A. H. P., TERRA, W. R. & FERREIRA, C. 2003. Characterization of a β -glycosidase highly active on disaccharides and of a β -galactosidase from *Tenebrio molitor* midgut lumen. *Insect Biochemistry and Molecular Biology*, 33, 253-265.
- FERREIRA, C., CAPELLA, A. N., SITNIK, R. & TERRA, W. R. 1994a. Digestive Enzymes in Midgut Cells , Endo- and Ectoperitrophic Contents , and Peritrophic Membranes of *Spodoptera frugiperda* (Lepidoptera) Larvae. 313, 299-313.

- FERREIRA, C., CAPELLA, A. N., SITNIK, R. & TERRA, W. R. 1994b. Properties of the digestive enzymes and the permeability of the peritrophic membrane of *Spodoptera frugiperda* (Lepidoptera) larvae. 107, 631-640.
- FERREIRA, C., REBOLA, K. G., CARDOSO, C., BRAGATTO, I., RIBEIRO, A. F. & TERRA, W. R. 2015. Insect midgut carboxypeptidases with emphasis on S10 hemipteran and M14 lepidopteran carboxypeptidases. *Insect Mol Biol*, 24, 222-39.
- FINN, R. D., BATEMAN, A., CLEMENTS, J., COGGILL, P., EBERHARDT, R. Y., EDDY, S. R., HEGER, A., HETHERINGTON, K., HOLM, L., MISTRY, J., SONNHAMMER, E. L. L., TATE, J. & PUNTA, M. 2014. Pfam: the protein families database. *Nucleic Acids Research*, 42, D222-D230.
- GAUTHIER, S. A. & HEWES, R. S. 2006. Transcriptional regulation of neuropeptide and peptide hormone expression by the *Drosophila* dimmed and cryptocephal genes. *Journal of Experimental Biology*, 209, 1803-1815.
- GENTA, F. A., BLANES, L., CRISTOFOLETTI, P. T., DO LAGO, C. L., TERRA, W. R. & FERREIRA, C. 2006. Purification, characterization and molecular cloning of the major chitinase from *Tenebrio molitor* larval midgut. *Insect Biochemistry and Molecular Biology*, 36, 789-800.
- GENTA, F. A., BRAGATTO, I., TERRA, W. R. & FERREIRA, C. 2009. Purification, characterization and sequencing of the major β -1,3-glucanase from the midgut of *Tenebrio molitor* larvae. *Insect Biochemistry and Molecular Biology*, 39, 861-874.
- HEIDEL-FISCHER, H. M. & VOGEL, H. 2015. Molecular mechanisms of insect adaptation to plant secondary compounds. *Current Opinion in Insect Science*, 8, 8-14.
- HEWES, R. S., SCHAEFER, A. M. & TAGHERT, P. H. 2000. The cryptocephal gene (ATF4) encodes multiple basic-leucine zipper proteins controlling molting and metamorphosis in *Drosophila*. *Genetics*, 155, 1711-1723.
- HOUSEMAN, J. G., MORRISON, P. E. & DOWNE, A. E. R. 1985. Cathepsin B and aminopeptidase in the posterior midgut of *Euschistus euschistoides* (Hemiptera: Phymatidae). *Can J. Zool.*, 63, 1288-1291.
- HUANG, X. & MADAN, A. 1999. CAP3: A DNA sequence assembly program. *Genome research*, 9, 868-77.
- IZUMI, Y., YANAGIHASHI, Y. & FURUSE, M. 2012. A novel protein complex, Mesh-Ssk, is required for septate junction formation in the *Drosophila* midgut. *J Cell Sci*, 125, 4923-33.
- KANG, D., LIU, G., LUNDSTRÖM, A., GELIUS, E. & STEINER, H. 1998. A peptidoglycan recognition protein in innate immunity conserved from insects to humans. *Proceedings of the National Academy of Sciences*, 95, 10078-10082.
- KUMAR, S. & BLAXTER, M. L. 2010. Comparing de novo assemblers for 454 transcriptome data. *BMC genomics*, 11, 571.
- LEE, J. E., ONEY, M., FRIZZELL, K., PHADNIS, N. & HOLLIN, J. 2015. *Drosophila melanogaster* Activating Transcription Factor 4 Regulates Glycolysis During Endoplasmic Reticulum Stress. *Genes Genomes Genetics* 5, 667-675.
- LEMOES, F. J. A., RIBEIRO, A. F. & TERRA, W. R. 1993. A bacteria-digesting midgut-lysozyme from *Musca domestica* (diptera) larvae. Purification, properties and secretory mechanism. *Insect Biochemistry and Molecular Biology*, 23, 533-541.
- LI, H. & DURBIN, R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics (Oxford, England)*, 25, 1754-60.
- MARGULIES, M., EGHOLM, M., ALTMAN, W. E., ATTIIYA, S., BADER, J. S., BEMBEN, L. A., BERKA, J., BRAVERMAN, M. S., CHEN, Y.-J., CHEN, Z., DEWELL, S. B., DU, L., FIERRO, J. M., GOMES, X. V., GODWIN, B. C., HE, W., HELGESEN, S., HO, C. H., IRZYK, G. P., JANDO, S. C., ALENQUER, M. L. I., JARVIE, T. P., JIRAGE, K. B., KIM, J.-B., KNIGHT, J. R., LANZA, J. R., LEAMON, J. H., LEFKOWITZ, S. M., LEI, M., LI, J., LOHMAN, K. L., LU, H., MAKHIJANI, V. B., MCDADE, K. E., MCKENNA, M. P., MYERS, E. W., NICKERSON, E., NOBILE, J. R., PLANT, R., PUC, B. P., RONAN, M. T., ROTH, G. T., SARKIS, G. J., SIMONS, J. F., SIMPSON, J. W., SRINIVASAN, M., TARTARO, K. R., TOMASZ, A., VOGT, K. A., VOLKMER, G. A., WANG, S. H., WANG, Y., WEINER, M. P., YU, P., BEGLEY, R. F. & ROTHBERG, J. M. 2005. Genome sequencing in microfabricated high-density picolitre reactors. *Nature*, 437, 376-381.
- MOREIRA, N. R., CARDOSO, C., RIBEIRO, A. F., FERREIRA, C. & TERRA, W. R. 2015. Insect midgut α -mannosidases from family 38 and 47 with emphasis on those of *Tenebrio molitor*. *Insect Biochemistry and Molecular Biology*, 67, 94-104.
- MUTHUKRISHNAN, S., MERZENDORFER, H., ARAKANE, Y. & KRAMER, K. J. 2012. Chitin metabolism in insects. In: GILBERT, L. I. (ed.) *Insect Molecular Biology and Biochemistry*. London: Academic Press/Elsevier.

- PARRA, J. R. P. 1986. Criação de insetos para estudos com patógenos. *Controle Microbiano de Insetos*. São Paulo: Editora Manole.
- PIMENTEL, A. C., FUZITA, F. J., PALMISANO, G., FERREIRA, C. & TERRA, W. R. 2017. Role of cathepsins D in the midgut of *Dysdercus peruvianus*. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 204, 45-52.
- POULSON, D. & WATERHOUSE, D. 1960. Experimental Studies on Pole Cells and Midgut Differentiation in Diptera. *Australian Journal of Biological Sciences*, 13, 541.
- POWELL, S., FORSLUND, K., SZKLARCZYK, D., TRACHANA, K., ROTH, A., HUERTA-CEPAS, J., GABALDON, T., RATTEI, T., CREEVEY, C., KUHN, M., JENSEN, L. J., VON MERING, C. & BORK, P. 2014. eggNOG v4.0: nested orthology inference across 3686 organisms. *Nucleic Acids Research*, 42, D231-D239.
- ROBINSON, M. D., MCCARTHY, D. J. & SMYTH, G. K. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*, 26, 139-140.
- SATO, P. M., LOPES, A. R., JULIANO, L., JULIANO, M. A. & TERRA, W. R. 2008. Subsite substrate specificity of midgut insect chymotrypsins. *Insect Biochemistry and Molecular Biology*, 38, 628-633.
- SILVA, C. P. & TERRA, W. R. 1994. Digestive and Absorptive Sites Along the Midgut of the Cotton Seed Sucker Bug *Dysdercus peruvianus* (Hemiptera : Pyrrhocoridae). 24, 493-505.
- SPIT, J., HOLTOF, M., BADISCO, L., VERGAUWEN, L., VOGEL, E., KNAPEN, D. & VANDEN BROECK, J. 2016. Transcriptional Analysis of The Adaptive Digestive System of The Migratory Locust in Response to Plant Defensive Protease Inhibitors. *Sci Rep*, 6, 32460.
- SUI, Y. P., WANG, J. X. & ZHAO, X. F. 2009. The impacts of classical insect hormones on the expression profiles of a new digestive trypsin-like protease (tlp) from the cotton bollworm, *Helicoverpa armigera*. *Insect Molecular Biology*, 18, 443-452.
- SUZEK, B. E., HUANG, H., MCGARVEY, P., MAZUMDER, R. & WU, C. H. 2007. UniRef: comprehensive and non-redundant UniProt reference clusters. *Bioinformatics*, 23, 1282-1288.
- TARGA, H. J. & PERES, C. A. 1979. Radiation-induced dominant lethal mutations in oocytes of *Musca domestica*. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 63, 153-160.
- TATUSOV, R. L., KOONIN, E. V. & LIPMAN, D. J. 1997. A genomic perspective on protein families. *Science (New York, N.Y.)*, 278, 631-7.
- TERRA, W. R. 1990. Evolution of Digestive Systems of Insects. *Annual Review of Entomology*, 35, 181-200.
- TERRA, W. R. & CRISTOFOLETTI, P. T. 1996. Midgut proteinases in three divergent species of Coleoptera. *Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology*, 113, 725-730.
- TERRA, W. R. & FERREIRA, C. 2012. Biochemistry and molecular biology of digestion. In: GILBERT, L. I. (ed.) *Insect Molecular Biology and Biochemistry*. London: Academic Press/Elsevier.
- TERRA, W. R., FERREIRA, C. & BASTOS, F. 1985. Phylogenetic considerations of insect digestion. Disaccharidases and the spatial organization of digestion in the *Tenebrio molitor* larvae. *Insect Biochemistry*, 15, 443-449.
- VENANCIO, T. M., CRISTOFOLETTI, P. T., FERREIRA, C., VERJOVSKI-ALMEIDA, S. & TERRA, W. R. 2009. The *Aedes aegypti* larval transcriptome: A comparative perspective with emphasis on trypsins and the domain structure of peritrophins. *Insect Molecular Biology*, 18, 33-44.

FIGURES




Figure 1: Differentially expressed genes. a) summary of all up ($\log_{2}FC \geq 2$, $p\text{-value} \leq 0.001$) and down regulated genes ($\log_{2}FC \leq -2$, $p\text{-value} \leq 0.001$); b) Venn diagram representing eggNOG families with at least one gene from the respective species up-regulated in anterior midgut compared with whole midgut; c) Venn diagram of eggNOG families with at least one gene from the respective species up-regulated in whole midgut compared with carcass. Species abbreviations: Mdom: *M. domestica*, Dper: *D. peruvianus*; Tmol: *T. molitor* and; Sfru: *S. frugiperda*.




Figure 2: Venn diagram representing eggNOG families with at least one gene among the 10% most highly expressed genes in the respective species in: a) whole midgut; b) carcass ; c) anterior midgut. Species abbreviations: Mdom: *M. domestica*, Dper: *D. peruvianus*; Tmol: *T. molitor* and; Sfru: *S. frugiperda*.

TABLES

Table 1: Sequencing and mapping statistics

Species	Raw reads	Total mapped reads	Uniquely mapped reads
<i>D. peruvianus</i>	609,407	531,707	445,533
<i>M. domestica</i>	608,637	559,459	491,359
<i>S. frugiperda</i>	747,904	696,223	602,167
<i>T. molitor</i>	677,080	624,423	550,598

Note: The mapping statistics were derived from the remapping of the original reads against their respective assembled transcriptome.

Table 2: Assembly statistics

Species	Newbler	MIRA	CAP3	Total contigs after contaminant filtering
<i>D. peruvianus</i>	8,093	27,395	6,774	6,395
<i>M. domestica</i>	5,329	20,589	4,248	4,005
<i>S. frugiperda</i>	10,315	28,308	7,264	6,833
<i>T. molitor</i>	12,370	37,703	9,583	9,010

Table 3: Functional classification of eggNOG annotated transcripts

Classification (one letter eggNOG code)	Number of transcripts (in %)			
	<i>D. peruvianus</i>	<i>M. domestica</i>	<i>S. frugiperda</i>	<i>T. molitor</i>
Function unknown(S)	523(17.61)	494(19.71)	894(24.65)	1255(21.77)
Posttranslational modification, protein turnover, chaperones(O)	398(13.4)	316(12.61)	433(11.94)	690(11.97)
Signal transduction mechanisms(T)	331(11.14)	205(8.18)	291(8.02)	621(10.77)
Translation, ribosomal structure and biogenesis(J)	208(7)	142(5.67)	162(4.47)	225(3.9)
Intracellular trafficking, secretion, and vesicular transport(U)	198(6.67)	138(5.51)	154(4.25)	320(5.55)
Carbohydrate transport and metabolism(G)	169(5.59)	171(6.82)	245(6.75)	426(7.39)
Transcription(K)	160(5.39)	55(2.19)	146(4.03)	331(5.74)
Energy production and conversion(C)	149(5.02)	63(2.51)	207(5.71)	195(3.38)
Lipid transport and metabolism(I)	145(4.88)	172(6.86)	175(4.82)	326(5.66)
RNA processing and modification(A)	130(4.38)	39(1.56)	113(3.12)	207(3.59)
Amino acid transport and metabolism(E)	128(4.31)	26(1.04)	241(6.64)	205(3.56)
Cytoskeleton(Z)	122(4.11)	106(4.23)	145(4)	229(3.97)
Secondary metabolites biosynthesis, transport and catabolism(Q)	93(3.13)	108(4.31)	138(3.8)	197(3.42)
Inorganic ion transport and metabolism(P)	74(2.49)	65(2.59)	89(2.45)	150(2.6)
Cell cycle control, cell division, chromosome partitioning(D)	50(1.68)	25(1)	51(1.41)	116(2.01)
Cell wall/membrane/envelope biogenesis(M)	49(1.65)	24(0.96)	34(0.94)	88(1.53)
Nucleotide transport and metabolism(F)	43(1.45)	44(1.76)	63(1.74)	82(1.42)
Coenzyme transport and metabolism(H)	33(1.11)	32(1.28)	35(0.96)	61(1.06)
Replication, recombination and repair(L)	32(1.08)	21(0.84)	54(1.49)	111(1.93)

Chromatin structure and dynamics(B)	27(0.91)	10(0.4)	32(0.88)	60(1.04)
Extracellular structures(W)	26(0.88)	16(0.64)	32(0.88)	36(0.62)
Nuclear structure(Y)	17(0.57)	3(0.12)	8(0.22)	29(0.5)
Defense mechanisms(V)	16(0.54)	28(1.12)	37(1.02)	54(0.94)
Cell motility(N)	2(0.07)	2(0.08)	3(0.08)	4(0.07)

Note: A given unigene can be assigned to more than one function code.

Table 4: Annotation summary

Properties	<i>D. peruvianus</i>	<i>M. domestica</i>	<i>S. frugiperda</i>	<i>T. molitor</i>
Total contigs	6,395	4,005	6,833	9,010
BBH hit - eggNOG	2,402	1,976	2,823	4,410
Top BLAST hit - eggNOG	568	530	804	1,354
Top BLAST hit - uniref90	1,809	716	1,331	2,208
Hit with Pfam	317	299	533	595
Annotated contigs (total)	5,096	3,521	5,491	8,387
Annotated contigs (%)	79.69	87.92	80.36	93.09

Table 5: Orthologous groups with at least one member up-regulated in whole midgut when compared to carcass in all studied insects.

eggNOG ID	Description
ENOG410362J	AMOP (MESH genes)
ENOG41034JA	glycoside hydrolase family 31
ENOG410382G	Protein of unknown function (DUF3421)
ENOG41032RI	Aminopeptidase
ENOG41039KJ	Haemolymph juvenile hormone binding protein (JHBP)

Table 6: Orthologous groups with at least one member among the top 10% most highly expressed genes in whole midgut.

eggNOG ID	Description	Dper	Mdom	Sfru	Tmol
ENOG41031YA	ATPase subunit B	Y	Y	Y	N
ENOG4103206	ribosomal protein L19	Y	Y	Y	Y
ENOG4103269	ribosomal protein S9	Y	Y	Y	Y
ENOG41032AN	NA; possibly involved in peptidoglycan recognition	N	N	N	N
ENOG41032F0	ribosomal protein P0	Y	Y	Y	Y
ENOG41032JU	vacuolar ATP synthase subunit H	N	N	Y	N
ENOG41032Q3	Tubulin is the major constituent of microtubules.	Y	Y	Y	Y
ENOG41032RI	Aminopeptidase	N	N	N	N
ENOG4103369	activated protein kinase c	Y	Y	Y	Y
ENOG410336G	citrate synthase	Y	Y	Y	N
ENOG410338D	ribosomal protein L5	Y	Y	Y	Y

ENOG41033B8	Mitochondrial membrane ATP synthase	Y	Y	N	Y
ENOG41033DB	Protein disulfide isomerase	Y	Y	Y	Y
ENOG41033N4	Ribosomal protein L15	Y	Y	Y	Y
ENOG41033P7	ribosomal protein	Y	Y	Y	Y
ENOG41034RS	Sterol carrier protein	N	N	Y	Y
ENOG41034Y5	Involved in calcium binding and microtubule stabilization (By similarity)	Y	Y	Y	Y
ENOG410351T	40S ribosomal protein S8	Y	Y	Y	Y
ENOG4103520	Thiolase, C-terminal domain	N	N	N	N
ENOG4103523	polyadenylate-binding protein	Y	Y	Y	Y
ENOG410356S	Cathepsin	Y	N	Y	Y
ENOG4103598	40S ribosomal protein	Y	Y	Y	Y
ENOG410359F	This protein promotes the GTP-dependent binding of aminoacyl-tRNA to the A-site of ribosomes during protein biosynthesis (By similarity)	Y	Y	Y	Y
ENOG41035BZ	Produces ATP from ADP in the presence of a proton gradient across the membrane (By similarity)	Y	Y	Y	Y
ENOG41035IX	Tubulin is the major constituent of microtubules.	Y	Y	Y	Y
ENOG41035KN	Required for the assembly and or stability of the 40S ribosomal subunit.	Y	Y	Y	Y
ENOG4103648	Arginine kinase	Y	Y	Y	Y
ENOG41036EA	ribosomal protein S3	Y	Y	Y	Y
ENOG41036JR	Ribosomal protein S16	Y	Y	Y	Y
ENOG41036K1	Ribosomal protein S11	Y	Y	Y	Y
ENOG410373X	Tetraspanin family	N	N	N	N
ENOG410375W	SERine Proteinase Inhibitors	N	Y	Y	Y
ENOG41037D0	cytochrome C oxidase	N	N	Y	Y
ENOG410382G	Protein of unknown function (DUF3421)	N	N	N	Y
ENOG41039KJ	Haemolymph juvenile hormone binding protein (JHBP)	N	N	N	N

The last four columns indicate whether the gene family is also highly expressed in carcass, which is a strong feature of housekeeping genes. Species abbreviations: Mdom: *M. domestica*, Dper: *D. peruvianus*; Tmol: *T. molitor* and; Sper: *S. frugiperda*.

Table 7: Common OGs with at least one member in the top 10% most highly expressed genes in anterior midgut.

eggNOG_cID	Description	Mdom		Sfru		Tmol	
		WM	CAR	WM	CAR	WM	CAR
ENOG41031YA	ATPase subunit B	Y	Y	Y	Y	Y	N
ENOG4103206	ribosomal protein L19	Y	Y	Y	Y	Y	Y
ENOG4103269	ribosomal protein S9	Y	Y	Y	Y	Y	Y

ENOG41032DQ	Calreticulin	N	N	Y	Y	Y	Y
ENOG41032F0	ribosomal protein P0	Y	Y	Y	Y	Y	Y
ENOG41032FD	ribosomal protein	Y	Y	Y	Y	Y	Y
ENOG41032FU	K05692 actin beta gamma 1	Y	Y	Y	Y	Y	Y
ENOG41032Q3	Tubulin is the major constituent of microtubules.	Y	Y	Y	Y	Y	Y
ENOG41032QK	alpha-amylase	Y	Y	Y	N	Y	Y
ENOG41032QY	Peroxiredoxin (EC 1.11.1.15)	N	N	N	N	N	Y
ENOG41032RI	Aminopeptidase	Y	N	Y	N	Y	N
ENOG41032XF	Triose-phosphate isomerase	Y	N	Y	Y	Y	Y
ENOG41032ZC	Angiotensin-converting enzyme	Y	Y	Y	N	Y	Y
ENOG4103369	activated protein kinase c	Y	Y	Y	Y	Y	Y
ENOG410336G	citrate synthase	Y	Y	Y	Y	Y	N
ENOG4103370	Occurs in almost all aerobically respiring organisms and serves to protect cells from the toxic effects of hydrogen peroxide (By similarity)	Y	Y	Y	Y	Y	Y
ENOG410338D	ribosomal protein L5	Y	Y	Y	Y	Y	Y
ENOG41033AG	Phosphopyruvatehydratase(EC 4.2.1.11)	N	Y	Y	Y	N	Y
ENOG41033B3	Glucuronosyltransferase (EC 2.4.1.17)	N	N	Y	N	Y	N
ENOG41033B8	Mitochondrial membrane ATP synthase	Y	Y	Y	N	Y	Y
ENOG41033DB	Protein disulfide isomerase	Y	Y	Y	Y	Y	Y
ENOG41033DK	oxidoreductase activity	Y	N	Y	N	Y	N
ENOG41033E4	Ribosomal protein L12	Y	Y	Y	Y	Y	Y
ENOG41033N4	Ribosomal protein L15	Y	Y	Y	Y	Y	Y
ENOG41033P7	ribosomal protein	Y	Y	Y	Y	Y	Y
ENOG41033ZE	Actins	N	Y	Y	Y	Y	Y
ENOG410340Q	Malate dehydrogenase	Y	Y	Y	Y	Y	Y
ENOG4103418	Insulinase (Peptidase family M16)	N	N	Y	Y	Y	Y
ENOG410347N	Aldehyde dehydrogenase	Y	N	Y	Y	N	Y
ENOG41034BW	Phosphate carrier protein	N	Y	Y	Y	Y	Y
ENOG41034I7	Produces ATP from ADP in the presence of a proton gradient across the membrane (By similarity)	N	Y	Y	Y	Y	Y
ENOG41034JA	Glycosyl hydrolase family 31	N	N	Y	N	N	N
ENOG41034MU	5'-nucleotidase, C-terminal domain	Y	N	Y	N	Y	N
ENOG41034RS	Sterol carrier protein	Y	N	Y	Y	Y	Y
ENOG41034UM	Enoyl-CoA hydratase	N	Y	Y	Y	Y	Y
ENOG41034Y5	Involved in calcium binding and microtubule stabilization (By similarity)	Y	Y	Y	Y	Y	Y
ENOG41034Z5	fructose-bisphosphatealdolase	N	Y	Y	Y	Y	Y
ENOG410351F	serine-type carboxypeptidase activity	Y	N	Y	N	Y	N
ENOG410351T	40S ribosomal protein S8	Y	Y	Y	Y	Y	Y

ENOG410351Z	peroxisomal multifunctional enzyme type	Y	N	Y	Y	Y	N
ENOG4103520	Thiolase, C-terminal domain	Y	N	Y	Y	Y	N
ENOG4103523	polyadenylate-binding protein	Y	Y	Y	Y	Y	Y
ENOG410356S	Cathepsin	Y	N	Y	Y	Y	Y
ENOG4103598	40S ribosomal protein	Y	Y	Y	Y	Y	Y
ENOG410359F	This protein promotes the GTP-dependent binding of aminoacyl-tRNA to the A-site of ribosomes during protein biosynthesis (By similarity)	Y	Y	Y	Y	Y	Y
ENOG41035A9	Dehydrogenase reductase SDR family member	Y	N	Y	N	N	Y
ENOG41035BZ	Produces ATP from ADP in the presence of a proton gradient across the membrane (By similarity)	Y	Y	Y	Y	Y	Y
ENOG41035GM	Larval storage protein (LSP) which may serve as a store of amino acids for synthesis of adult proteins	Y	Y	N	Y	N	Y
ENOG41035IX	Tubulin is the major constituent of microtubules.	Y	Y	Y	Y	Y	Y
ENOG41035KN	Required for the assembly and or stability of the 40S ribosomal subunit.	Y	Y	Y	Y	Y	Y
ENOG41035N8	Glucosylceramidase (EC 3.2.1.45)	Y	Y	Y	N	Y	N
ENOG41035RN	alkaline phosphatase	Y	N	Y	N	Y	N
ENOG41035ZK	Glyceraldehyde-3-phosphate dehydrogenase	Y	Y	Y	Y	N	N
ENOG4103648	Arginine kinase	Y	Y	Y	Y	Y	Y
ENOG41036EA	ribosomal protein S3	Y	Y	Y	Y	Y	Y
ENOG41036JR	Ribosomal protein S16	Y	Y	Y	Y	Y	Y
ENOG41036PA	ATP synthase	N	Y	Y	Y	Y	Y
ENOG41036ZV	Endonuclease_NS	Y	Y	Y	N	N	N
ENOG410371B	NA	Y	N	N	N	Y	N
ENOG410373X	Tetraspanin family	Y	N	Y	N	Y	N
ENOG410375W	SERine Proteinase Inhibitors	Y	Y	Y	Y	Y	Y
ENOG41037D0	cytochrome C oxidase	Y	N	Y	Y	Y	Y
ENOG41037R1	serine-type endopeptidase activity	Y	Y	Y	N	Y	N
ENOG410382G	Protein of unknown function (DUF3421)	Y	N	Y	N	Y	Y
ENOG41038D7	bZIP transcription factor	N	N	Y	Y	Y	Y
ENOG41038K5	serine protease	Y	Y	Y	N	Y	N
ENOG41039KJ	Haemolymph juvenile hormone binding protein (JHBP)	Y	N	Y	N	Y	N

The last six columns indicate whether the gene family is also highly expressed in carcass or WM of each species, similarly to Table S5. Species abbreviations: Mdom: *M. domestica*, Dper: *D. peruvianus*; Tmol: *T. molitor* and; Sper: *S. frugiperda*.