Predation of stocked masu salmon (*Oncorhynchus masou*) by riparian wildlife

Journal:	*Canadian Journal of Fisheries and Aquatic Sciences*
Manuscript ID	Draft
Manuscript Type:	Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Miyamoto, Kouta; National Research Institute of Fisheries Science, Japan Fisheries Research and Education Agency Squires, Theodore; Graduate school of Agriculture, Hokkaido University Araki, Hitoshi; Hokkaido University,
Please Select from this Special Issues list if applicable:	N/A
Keyword:	PREDATOR-PREY INTERACTION < General, hatchery management, wildlife, camera trap, body size

SCHOLARONE™
Manuscripts

Predation of stocked masu salmon (*Oncorhynchus masou*) by riparian wildlife

Kouta Miyamoto, Theodore E. Squires, and Hitoshi Araki

K. Miyamoto. National Research Institute of Fisheries Science, Japan Fisheries Research and

Education Agency, Nikko, Tochigi 321-1661, Japan, and Graduate school of Agriculture,

Hokkaido University, N9W9 Kita, Sapporo 060-8589, Japan

E-mail: mkouta@affrc.go.jp

T. E. Squires. Graduate school of Agriculture, Hokkaido University, N9W9 Kita, Sapporo

060-8589, Japan

E-mail: squires@res.agr.hokudai.ac.jp

H. Araki. Research Faculty of Agriculture, Hokkaido University, N9W9 Kita, Sapporo 060-8589,

Japan

E-mail: arakih@res.agr.hokudai.ac.jp

Corresponding author: Kouta Miyamoto, National Research Institute of Fisheries Science,

Japan Fisheries Research and Education Agency, Nikko, Tochigi 321-1661, Tel.: +81 288 55

0055; Fax.: +81 288 55 0064; E-mail: mkouta@fra.affrc.go.jp

Predation of stocked masu salmon (*Oncorhynchus masou*) by riparian wildlife

K. Miyamoto, T. E. Squires and H. Araki

Abstract

Predation after release is one of the major concerns of hatchery fish conservation and

propagation. However, the relationship among the size of hatchery fish, the predator species, and

their behaviors in natural environments is largely unknown. To understand the relationship, we

conducted predation experiments in outdoor tanks and a semi-natural stream with exposure to

local predators. Two different ranges of fork lengths of masu salmon (*Oncorhynchus masou*)

were examined as prey sizes. Camera trap data showed that grey herons (*Ardea cinerea*) were the

primary predator animal in the system, and that most herons utilized shallow areas in the

morning or evening. Increasing the density of stocked salmon brought in more grey herons. More

importantly, predation by grey herons resulted in the survival rate of larger salmon being

significantly lower than that of the smaller salmon. Our results suggest that it is important to

understand local predators, adjust the optimum body size of hatchery fish at release, and choose

the appropriate stocking site and time of day for maximizing the effectiveness of fish stocking.

Key words: predator-prey interaction, hatchery management, wildlife, camera trap, body size

Introduction

Predation by riparian wildlife is widely recognized as a key factor influencing the

survival of fishes in stream ecosystems (Kruuk, 1995; Draulans 1987; Roby et al. 2003;

Steinmetz et al. 2003). Salmonid fish populations reduced by riparian wildlife have been

reported, not only for stocked populations, but also for wild populations (e.g. Osterback et al.

2013; Frechette et al. 2015). It is also noteworthy that stocked salmonids are often intensively

preyed upon by birds (Wood, 1987; Martel & Dill, 1995; Harvey & Nakamoto, 2013) and

mammals (Roberts & Garcia de Leaniz, 2011). Hatchery-reared salmonid fishes are routinely

stocked in natural waterways as part of conservation efforts or stock enhancement (Brown and

Laland 2003; Salvanes and Braithwaite 2006; Fraser 2008). Therefore, if we implement a

management strategy that decreases local predation pressures, then effective conservation or

stock enhancement methods should be improved.

Studies examining the predator-prey interaction between riparian animals and fish have

been generally based on either investigating predation pressure of riparian animals without

observing fish response (Draulans 1987; Post et al. 1998; Harvey & Nakamoto 2013) or

investigating the mortality rate of fish without observing the behavior of predators (Penaluna et

al. 2015). However, the predation effect is influenced by many factors: fish migration, predator

species diversity, the number of predators present, and the duration of their sojourn (Gawlik,

2002, Steinmetz et al. 2003). Thus, methodological improvements which decrease predation

pressure on stocked fish require a profound understanding of both predator and prey behaviors.

Additionally, as food size preference depends on predator species (Carss and Marquiss 1991) and

the ability of fish to avoid predators depends on fish size (Dill & Fraser 1984), we need to

consider the size effect of prey fish.

To evaluate the impacts of predation by riparian animals, it is necessary to evaluate the

relationships among the predator species, their behaviors, and their preferred size of prey in a

place that allows for free movement of all parties. However, the number of studies on this

relationship is limited due to the difficulty in identifying species of predatory riparian animals

and observing their predation behavior directly. These studies are further complicated due to

complex behaviors exhibited by many predator species, such as individual movement for feeding

and nesting needs (Collis et al., 2002), movement due to changes in prey densities (Kushlan,

1976; Gawlik, 2002), and movement resulting from human disturbance (Klein, 1993).

Recently, camera trapping that uses fixed cameras, triggered by infrared sensors, to

'trap' images of passing animals, provide the opportunity to collect information on animal

behavior (Silveira et al. 2003; Wegge et al. 2004). Because camera trapping is a non-invasive

technique, it causes minimal environmental disturbance (Henschel & Ray 2003; Silveira et al.

2003). In this study, we conducted a predation test using camera traps with outdoor experimental

tanks and a semi-natural stream to investigate the relationships between the different predator

species, their behavior, and the size of stocked salmon.

Materials and Methods

Test fish

The test fish used in the experiments were hatchery-reared masu salmon (*Oncorhynchus*

masou) from the Shobu-shimizu River in Tochigi Prefecture, Japan (approximately 36°45'N,

139°27'E). Eggs were obtained from mature adults that had migrated into the Shobu-shimizu

River, and were handled according to standard hatchery procedures directed by the National

Research Institute of Fisheries Science (NRIFS) facility at Nikko. Young of the year (YOY) ($n =$

305) and one year old (OYO) ($n = 305$) salmon were employed in this study. The YOY and OYO

salmon were selected from five rearing tanks (50 cm width × 120 cm length × 20 cm depth) and

a large tank (3 m width × 1.5 m length × 0.9 m depth) respectively. The YOY fish used were

75–100 mm fish (75–100 mm FL-group, hereafter) and the OYO used were 135–160 mm fish

(135–160 mm FL-group, hereafter). Before the start of the study, fish were fed daily food rations

(commercial trout pellets) equal to 1.5-2.0% of their estimated body weight.

Tank experiment

We conducted predation tests in four outdoor tanks for three days from August to

September, 2013. Four FRP circular tanks (120-cm diameter × 15-cm high) at NRIFS in Nikko

were used in a forested and grassy area and each had a thin layer of natural gravel (2–7-cm

gravel and about 13-cm cobble substrate) on the bottom (Fig. 1a). A 10-cm high and 8-cm roof

fence that was installed on each tank both provided the fish with cover and prevented them from

jumping out. A camera (Trophy Cam HD, Bushnell, Overland Park, KS, USA) was set in the

south side of each tank to monitor the whole tank. The presence of cover and gravel substrate

facilitated near natural behavior of the masu salmon, allowing hiding and escaping (Miyamoto

2016a), thus minimizing experimental stress. During the study period the stocked fish were fed

daily food rations (commercial trout pellets) equal to around 2.0% of their body weight. Spring

water, 10.2 ± 0.3°C (mean ± SD) was introduced into each aquarium at a rate of 6 L/min. Two

different fork lengths (FL) of masu salmon were used as the size of prey fish, 75–100 mm

FL-group (mean ± SD, 91.9 ± 7.0 mm) and 135–160 mm FL-group (145.8 ± 6.8 mm).

To evaluate the size-selective predation risk, 30 fish (15 from each FL-group) were

placed in each tank. The number of fish of each FL-group that survived was counted each day.

The survival rate of salmon was then compared between the two FL-groups. To identify the

predator animals, the number of photos taken by camera trap (described in the section below)

was counted.

Next, to investigate the relationship between the number of stocked salmon and the

frequency in which predator animals appeared, three outdoor tanks were used. The test fish were

divided into 10, 5, and 0 individuals and stocked randomly in each of the three tanks. Three days

later, the number of photos of potential predators taken by camera trap (described in the section

below) was counted for each tank. This trial was replicated three times for each FL-group.

Predation test in a semi-natural stream

The predation test was conducted for 20 days from August to September, 2013 using a

semi-natural stream (mean ± standard deviation [SD]: 120 m long, 95.8 ± 4.3-cm wide; 2%

gradient drain) in a forested and grassy area at NRIFS (Fig. 1b). The stream was constructed of

stone, wood, and soil. Spring water, at 10.2 ± 0.3°C (mean ± SD) was drained at 18 L/s. The

stream was forked to allow free passage into four 30 meter sections (Sec. 1-4 from upper to

lower). Each section contained three pools and three riffles. The pools were about 80-cm long

and about 60-cm wide with maximum depth of approximately 40-cm. Woody debris (c. 70-cm

long and c. 20-cm wide) were placed in the sides of each pool. The riffles were 9–12-cm deep

and contained a mixture of 2–7-cm gravel and about 13-cm cobble substrate. A 50-cm waterfall

was built in the top of the stream with two metal gates (1-cm × 2-cm mesh) at the top to prevent

fish from escaping. The water drain was separated by a metal mesh gate (1-cm × 2-cm mesh)

with the bottom quarter of the gate covered with plastic mesh (8-mm × 8-mm mesh). A total of

400 fish (200 from each FL-group) were stocked in Sec. 2 after measuring their fork length and

body weight. At the end of the study period, a backpack electrofishing unit (Smith-Root,

Vancouver, WA, USA) was used to remove and count the fish that had survived. The electric

shock was repeated until the fish count was zero twice in a row.

To investigate the distribution of salmon in the stream, the fish in each section were

closed in by fish block nets (8-mm × 8-mm mesh) at the end of the experiment to prevent them

from moving between the sections. When fish were caught by electrofishing, each fish was

lightly anesthetized with 2-phenoxyethanol, its fork length and body weight were measured, and

the section that the fish was caught in was recorded. To identify the predator animals, 3 cameras

(Trophy Cam HD, Bushnell, Overland Park, KS, USA) were arranged in each section and placed

at 10-m intervals along the riverside to monitor both sides of the stream, and the number of

photos that were taken by camera traps (described in the section below) was counted for each

section. In addition, to estimate the predation behavior of wild animals, the position of potential

predators (at pool or riffle) in photos was recorded. Then, to investigate the position that

predators utilized, the proportion of predators located around pools or riffles was calculated in

each section, then the average proportions were compared.

Camera trap

To assess predator encounters during day and night, we recorded potential predators

using motion and infrared sensor camera traps in the tank experiments and the semi-natural

stream experiment. Each camera was mounted on a wooden stake so that the camera was about

50 cm above the water's surface. Cameras were triggered with a passive infrared motion sensor;

the camera was set to wait 15 seconds after an initial trigger entered its sensor range before

attempting to detect additional triggers. To identify predators and estimate the frequency of their

visits to the study site, all the photos containing potential predators were checked by KM. For

some ambiguous species identifications, additional checking was performed by a local wildlife

expert Dr. T. Takeda from Nikko National Park. When more than one potential predator was

captured in a photo, the species and their number was recorded. In addition, the number of

photos that showed predators capturing or eating fish was counted, and the predator species was

recorded.

Statistical analyses

In the tank tests, Two-way ANOVA was used to determine the effects of study period

and fish fork length on the survival rate of salmon. Additionally, a second set of ANOVA tests

were used to determine how the number of salmon, and salmon size, effected the number of

photos containing riparian predators in the stream test. One-way ANOVA was used to compare

the average proportions of the predators located around pools and riffles in the stream. Data was

log (x + 0.5) transformed prior to analysis, or, when proportions were tested, data was arcsin $\sqrt{x}$

transformed. Tukey HSD test was used as a post-hoc test.

For the semi-natural stream experiment, Pearson's chi-square test was used to compare

the proportion of the number of fish captured at each section. However, the omnibus chi-square

value does not specify which combination of categories contributes to statistical significance;

thus, the adjusted standardized residuals (ASR) was used for each value to determine

discrepancies between the observed and expected value (Haberman, 1973). |ASR| > 1.96 and >

2.56 show $P < 0.05$ and $P < 0.01$, respectively. A $P < 0.05$ was considered significant in all

statistical analyses. Data analyses were generated using IBM SPSS software (version 21.0 of

IBM SPSS Statistics for Windows).

Results

Tank experiment

In the tank experiment, cameras captured grey heron (*Ardea cinerea*), Japanese marten

(*Martes melampus*), raccoon dog (*Nyctereutes procyonoides*), and large-billed crow (*Corvus*

macrorhynchos). The total number of photos containing potential predators was 186 (including

three photos with ambiguous species, which were identified by the local wildlife expert). A total

of 173 photos of grey heron were taken and there were six photos or fewer of each of the other

animals (Japanese marten: three times, raccoon dog: four times, large-billed crow: six times).

Twenty one photos showed grey herons capturing prey fish. Therefore, in tank experiments, grey

herons were regarded as the main predator.

With regard to the size-selective predation risk, both the length of the study period and

the fork length of the stocked salmon had significant effects on the fish survival (Period, $F_{2, 18} =$

10.56, $P < 0.010$; Salmon length, $F_{1, 18} = 36.67$, $P < 0.001$; Period × Salmon length, $F_{2, 18} = 0.18$,

$P = 0.833$). The fish survival rate was significantly higher in the 75-100 mm FL-group than in

the 135-160 mm FL-group during the study period (Tukey HSD test, all shows $P < 0.050$) (Fig.

2).

In the Two-way ANOVA test for frequency of predator appearance, only the number of

fish in the tank had a significant effect on the number of photos containing grey herons (Number

of fish, $F_{2, 12} = 149.26$, $P < 0.001$; Fork length, $F_{1, 12} = 2.48$, $P = 0.141$; Number of fish × Fork

length, $F_{2, 12} = 0.02$, $P = 0.981$) and the number of photos containing grey herons increased

significantly when the number of fish in the tanks was large (Tukey HSD test, all shows $P <$

0.001) (Fig. 3). At the end of the three-day experiment, there were no surviving salmon.

Stream experiment

The overall survival rate of the fish in this experiment was 18.0% ($n = 72$). The survival

rate was significantly higher for the 75–100 mm FL-group (33.0%, $n = 66$) than for the 135-160

mm FL-group (3.0%, $n = 6$) (Peason's χ^2 test, $\chi^2 = 60.98$, $df = 1$, $p < 0.001$).

There was no significant difference between the proportion of fish that survived in each

section between both fish groups (Peason's χ^2 test, $\chi^2 = 0.77$, $df = 3$, $p = 0.86$). Comparing the

actual proportion of surviving fish (75-100 and 135-160 mm) in each section to the expected

surviving proportion (we hypothesized a uniform 1:1:1:1 ratio) showed there was a significant

difference (Peason's χ^2 test, $\chi^2 = 15.43$, $df = 3$, $p < 0.001$). Accordingly, the number of fish

surviving in Sec. 2 was significantly higher than the average number of fish capture in each

section (ASR = 3.4, *p* < 0.010) (Fig. 4).

Cameras captured grey heron, brown dipper (*Cinclus pallasii*), Japanese red fox (*Vulpes*

vulpes japonica) and large-billed crow during the predation test in the semi-natural stream. The

total number of photos containing potential predators was 470. Among them, 455 contained grey

heron. There were fewer than 10 photos of other animals (brown dipper: one times, Japanese red

fox: seven times, large-billed crow: seven times). The most grey heron photos (*n* = 104) were

taken at Sec. 2 on day-5 (Fig. 5). 76.5 % of all pictures that contained grey heron were taken at

Sec.2. Five photos had two grey herons in them, and two photos had three grey herons in them,

all photos with multiple herons were taken in Sec. 2 between day-4 and day-7 after stocking.

Seventeen photos showed grey herons capturing prey fish. Grey herons were most frequently

photographed in the mornings and evenings.

In order to investigate the position that grey herons utilized, 412 photos were analyzed

(455 minus 43 photos for undistinguishable positioning). The average proportion of photos

showing grey heron around the pools (6.88 ± 10.68 %) (Fig. 1c) was significantly lower than

those around the riffles (93.13 ± 10.68 %) (Fig. 1d) ($F_{1, 6}$ = 53.13, *P* < 0.010) (Fig. 6). The

overall proportions of photos taken of grey herons in the morning (3:00-9:00) and in the evening

(15:00-21:00) were 49.3% (*n* = 224) and 47.9% (*n* = 218), respectively.

Discussion

In the 20-day stream experiment, we found that 82% of fish were taken and that grey

heron was the most frequently visiting predator. It has been previously reported that salmonid

populations can be seriously damaged by avian predation, (Feltham 1995; Stewart et al.2005),

but there have been no quantitative assessments of the impact of grey heron on river fisheries

(Harris et al. 2008). Our results suggest that at least in our experimental settings, grey herons can

significantly reduce salmonid populations by consuming juvenile fish. On the other hand, it is

important to note that water depth can strongly influence wading birds' selection of foraging

habitat (Master et al. 2005, Gawlik and Crozier 2007). The length of a grey heron's legs restricts

the maximum depth at which they can forage up to 16 cm deep (Ntiamoa-Baidu et al. 1998),

thereby limiting the habitat suitable for hunting. Our observation that grey herons showed a

significant preference for the riffle, rather than the pool, in the semi-natural stream is consistent

with previous studies. Thus, where local salmonids reside in shallow water, it is reasonable to

expect that grey herons exert a high predation pressure.

On the first day of the tank experiments, grey herons preyed upon large salmon more

often than upon small salmon. In both the tank and the stream experiments, the survival rate of

large salmon was less than that of small salmon. Birds of the heron family (Ardeidae) including

the grey heron are reported to show a preference for larger mosquito fish *Gambusia affinis*

(Britton and Moser 1982) and sailfin molly *Poecilia latipinna* (Trexler et al., 1994) as prey. It is

suggested that larger fish are easier for avian predators to detect than small ones (Eriksson 1985,

Magnhagen 1988). After the second day of the tank experiment, however, both sizes of salmon

showed similar levels of decline on their survival rate. One possible explanation for this result is

the decreased opportunities to prey upon larger fish. Our results suggest that size-selective

predation by grey herons depends on the density of preferred prey size, and it appears that

size-selective predation occurred in the stream experiment as well as in the tank experiments.

Therefore, if fish with different body sizes are stocked in rivers, it might be important to consider

how size composition will potentially influence the behavior of predators.

It is established that larger fish are safer from fish-on-fish predation than smaller fish

(Peterson and Wroblewski 1984, Houde 1987, Miller et al. 1988, Miyamoto and Araki 2017).

One potential cause for the higher survival rate of larger fish is improved swimming ability that

allows them to better avoid predators as they grow (Beamish 1978; Lundvall et al. 1999).

However, the grey heron is an ambush predator that usually stands upright and waits for a fish to

approach (Tojo, 1996), so the swimming ability that a fish possesses to avoid aquatic predators

has significantly limited benefits to them in this case (Miyamoto 2016b), and large size might

make fish more detectable. Therefore, to decrease predation risk by wild animals, we suggest that

hatchery managers choose an optimum body size of released fish based on the predator species

inhabiting the stocking area. Additionally, to block the vision of grey herons or prevent access, it

can be recommended to stock fish in a structurally complex place that has a lot of woody debris

and/or big rocks if many grey herons are around during the stocking season.

Generally, in salmonids, larger individuals avoid exposure to a predatory threat and

reduce growth rate, whereas smaller individuals are less cautious and maintain their growth rate

even in the presence of a threat (Reinhardt 1999). Based on this behavior, it is expected that the

OYO salmon are more difficult for grey heron to catch than the YOY salmon. In contrast to this,

it was also reported that the hatchery environment selects for bolder individuals that spend more

time in open areas and are more active than wild specimens (Sundström et al., 2004). So, OYO

salmon that spent a longer time than the YOY salmon in the hatchery may display a higher

tendency for bold behavior (Roberts et al 2014). Thus, to fully understand the predation risks for

stocked salmon, further studies that estimate the individual behavior of fish (e.g. bold and shy)

and improve the maladapted behavior of fish (Berejikian et al 1999, Roberts et al 2011) will be

needed.

In the tank experiment where different numbers of salmon were stocked, the number of

grey herons present or the duration of their sojourn at the tanks was positively correlated to the

number of stocked fish. In other studies, large numbers of fish have been observed to remain

close to where they were stocked (Cresswell 1981), and piscivorous water birds have displayed

similar density and temporal trends in response to stocking (Gawlik 2002; Draulans 1987). This

suggests that for the duration of our stream experiment many salmon stayed in Sec. 2, as multiple

grey herons simultaneously appeared frequently in that section. We also captured more fish in

Sec. 2 by electrofishing than in any other sections on the final day of the stream experiment.

These results indicate that most stocked fish gathered without migrating from the initial stocking

site and continued to be preyed upon there. This situation may have serious implications for the

conservation and propagation of salmonids and it is important to further investigate the spatial

and temporal relationships between salmonids and their predators.

291   The great cormorant (*Phalacrocorax carbo*) is a type of piscivorous water bird, whose

predation on stocked fish became a serious problem, just after the fish were released (Kumada et

al. 2013). In this study, the grey herons appeared only two days after the fish were released, then

the number of photos rapidly increased for a few days indicating increased predation, or higher

predator abundance. This suggests that grey herons exhorted the strongest predation pressure

soon after stocking. To solve this problem, it is important to stock the fish in a safe place and

allocate as much time as possible for them to acclimate to the surrounding environment. The

results of this study showed that the grey heron has a tendency to avoid deep water and foraging

in the middle of day and night. In contrast, it previously was reported that grey herons forage

constantly during the daytime (Richner 1986; Sawara et al 1990), so human activities (Klein,

1993) and other feeding sites (Richner 1986) in the surrounding environment could have had an

effect on the observed behavior. Grey herons infrequently forage at night, and it has been

suggested that such feeding behavior is inefficient (Sawara et al 1990). Therefore, to decrease the

predation pressure of grey herons, it might be wise to stock fish in a deep area, or at night

(Roberts et al 2009), or both. In this way, if we can develop techniques to mitigate predatory

damage to fisheries by investigating the relationship between stocked fish and predatory animals,

it will not only help the conservation and propagation of stocked fish, but also support the

conservation of wild animals as well.

Acknowledgements

We would like to thank Kouji Mutou, Hidefumi Nakamura and Masaharu Murakami of the

National Research Institute of Fisheries Science for their help with the care of the fish. We thank

Dr Tsutomu Takeda of Nikko Yumoto Visitor Center of Nikko National Park for his help with

the identification of wildlife. We are grateful to Predrag Davidovic and Petar Ilic for critically

reading the manuscript and providing valuable comments. This work was supported by JSPS

KAKENHI Grant Number JP26292102.

References

Beamish, F.W.H. 1978. Swimming capacity. *In* Fish physiology, Vol. VII. *Edited by* W. S.

Hoar and D.J. Randall. Academic Press, New York. pp. 101–187.

Brown, C., and Laland K.N. 2003. Social learning in fishes: a review. Fish Fish. **4**: 280-288.

Berejikian, B.A., Smith, R.J.F., Tezak, E.P., Schroder, S.L., and Knudsen, C.M. 1999.

Chemical alarm signals and complex hatchery rearing habitats affect antipredator behavior and

survival of chinook salmon (*Oncorhynchus tshawytscha*) juveniles. Can. J. Fish. Aquat. Sci. **56**:

830–838.

Carss, D.N., and Marquiss, M. 1991. Avian predation at farmed and natural fisheries. *In*

Interaction between fisheries and environment. *Edited by* M.C. Lucas, I. Diak, and L. Laird.

Proc. Institute of Fisheries Management, 22nd Annual Study Course. University of Aberdeen.

University of Aberdeen. UK. pp 179–196.

Collis, K., Roby, D.D., Craig, D.P., Adamany, S., Adkins, J.Y., and Lyons, D.E. 2002. Colony

size and diet composition of piscivorous waterbirds on the lower. Columbia River: Implications

for losses of juvenile salmonids to avian predation. T. Am. Fish. Soc. **131**: 537–550.

Cresswell, R.C. 1981. Post‑stocking movements and recapture of hatchery‑reared trout

released into flowing waters–a review. J. Fish Biol. **18**(4): 429–442.

Dill, L.M., and Fraser, A.H.G. 1984. Risk of predation and the feeding behavior of juvenile

coho salmon (*Oncorhynchus kisutch*). Behav. Ecol. Sociobiol. **16**: 65–71.

Draulans, D. 1987. The effect of prey density on foraging behaviour and success of adult and

first-year grey herons (*Ardea cinerea*). J. Anim. Ecol., 479–493.

Eriksson, M.O. 1985. Prey detectability for fish-eating birds in relation to fish density and

water transparency. Ornis Scand. **16**: 1–7.

Feltham, M.J. 1995. Consumption of Atlantic salmon smolts and parr by goosanders: estimates

from doubly‑labelled water measurements of captive birds released on two Scottish rivers. J.

Fish Biol. **46**: 273–281.

Fraser, D.J. 2008. How well can captive breeding programs conserve biodiversity? A review of

salmonids. Evol. Appl. **1**: 535–586.

Frechette, D., Osterback, A.M.K., Hayes, S.A., Moore, J.W., Shaffer, S.A., Pavelka, M.,

Winchell, C., and Harvey, J. T. 2015. Assessing the Relationship between Gulls Larus spp. and

Pacific Salmon in Central California Using Radiotelemetry. N. Am. J. Fish. Manage. **35**(4):

775–788.

Gawlik, D.E. 2002. The effects of prey availability on the numerical response of wading birds.

Ecol. Monogr. **72**(3): 329–346.

Harris, C.M., Calladine, J.R., Wernham, C.V., and Park, K.J. 2008. Impacts of piscivorous

birds on salmonid populations and game fisheries in Scotland: a review. Wildlife Biol. **14**(4):

395–411.

Henschel, P., and Ray, J. 2003 Leopards in African Rainforests: Survey and Monitoring

Techniques. Wildlife Conservation Society, New York, NY.

Harvey, B.C., and R. J. Nakamoto, 2013. Seasonal and Among-Stream Variation in Predator

Encounter Rates for Fish Prey. T. Am. Fish. Soc. **142**: 621–627.

Houde, E.D. 1987. Fish early life dynamics and recruitment variability. Am. Fish. Soc. Symp.

2: 17–29.

Klein, M.L. 1993. Waterbird behavioral responses to human disturbances. Wildlife Soc. B. **21**:

31–39.

Kruuk, H. 1995. Wild Otters. Predation and populations. Oxford University Press. New York.

Kumada, N., Arima, T., Tsuboi, J., Ashizawa, A., and Fujioka, M. 2013. The multi-scale

aggregative response of cormorants to the mass stocking of fish in rivers. Fish. Res. **137**: 81–87.

Kushlan, J.A. 1976. Wading bird predation in a seasonally fluctuating pond. Auk **93**: 464–476.

Lundvall, D., Svanbäck, R., Persson, L., and Byström, P. 1999. Size-dependent predation in

piscivores: interactions between predator foraging and prey avoidance abilities. Can. J. Fish.

Aquat. Sci. **56**: 1285–1292. Magnhagen, C. 1988. Predation risk and foraging in juvenile pink

(*Oncorhynchus gorbuscha*) and chum salmon (*O. keta*). Can. J. Fish. Aquat. Sci. **45**(4): 592–596.

Martel, G., and Dill, L.M. 1995. Influence of movement by coho salmon (*Oncorhynchus*

kisutch) parr on their detection by common mergansers (*Mergus merganser*). Ethology **99**:

139–149.

Master, T.l., Lesier, J.K. Bennett, K.A. Bretsch, J.K., and Wolfe H.J. 2005. Patch selection by

Snowy Egrets. Colon. Waterbird. **28**: 220–224.

Miller, T.J., Crowder, L.B., Rice, J.A., and Marschall, E.A. 1988. Larval size and recruitment

mechanism in fisheries: toward a conceptual framework. Can. J. Fish. Aquat. Sci. **45**:

1657–1670.

Miyamoto, K. 2016a. Effect of visual and chemical stimuli on predator avoidance behavior in

juvenile masu salmon *Oncorhynchus masou*. Aquacul. Sci. **64**: 43–51.

Miyamoto, K. 2016b. Effects of body color luminance and behavioral characteristics on

predation risk in salmonid fishes. Hydrobiologia, **783**: 249–256.

Miyamoto, K., and Araki, H. 2017. Differentiated predation risk on hatchery-reared juvenile

masu salmon by white-spotted charr with different body sizes. Fish. Sci. doi:

10.1007/s12562-016-1059-8

Ntiamoa-Baidu Y., Piersma, T., Wiersma, P., Poot, M., Battley, P., and Gordon, C. 1998. Water

depth selection, daily feeding routines and diets of waterbirds in coastal lagoons in Ghana. Ibis.

140(1): 89–103.

Osterback, A.M.K., Frechette, D. M., Shelton, A. O., Hayes, S. A., Bond, M. H., Shaffer S. A.,

and Moore, J. W. 2013. High predation on small populations: avian predation on imperiled

salmonids. Ecosphere **4**(9): 116.

Peterson, I., and Wroblewski, J.S. 1984. Mortality rate of fishes in the pelagic ecosystem. Can.

J. Fish. Aquat. Sci. **41**(7): 1117–1120.

Reinhardt, U.G. 1999. Predation risk breaks size-dependent dominance in juvenile coho

salmon (*Oncorhynchus kisutch*) and provides growth opportunities for risk-prone individuals.

Can. J. Fish. Aquat. Sci. **56**(7): 1206–1212.

Richner, H. 1986. Winter feeding strategies of individually marked herons. Anim. Behav. **34**:

881–886.Roberts, L.J., Taylor, J., Gough, P.J., Forman, D.W., and de Garcia de Leaniz, C., 2009.

Night stocking facilitates nocturnal migration of hatchery‐reared Atlantic salmon, *Salmo salar*,

smolts. Fisheries Manag. Ecol. **16**: 10–13.

Roberts, L.J., and Garcia de Leaniz, C. 2011. Something smells fishy: predator-naïve salmon

use diet cues, not kairomones, to recognize a sympatric mammalian predator. Anim. Behav. **82**:

619–625.

Roberts, L.J., Taylor J., and Garcia de Leaniz, C., 2011. Environmental enrichment reduces

maladaptive risk-taking behavior in salmon reared for conservation. Biol. Conserv. **144**:

1972–1979.

Roberts, L.J., Taylor, J., Gough, P.J., Forman, D.W., and Garcia de Leaniz, C. 2014. Silver

spoons in the rough: can environmental enrichment improve survival of hatchery Atlantic salmon

Salmo salar in the wild? J. Fish Biol. **85**: 1972–1991.

Roby, D.D., Lyons, D.E., Craig, D.P., Collis, K., and Visser, G.H. 2003. Quantifying the effect

of predators on endangered species using a bioenergetics approach: Caspian terns and juvenile

salmonids in the Columbia River estuary. Can. J. Zool. **81**: 250–265.

Penaluna, B.E., Jason, B.D., and Noakes, D.L. 2015. Instream cover and shade mediate avian

predation on trout in semi-natural streams. Ecol. Freshw. Fish **25**: 405–411.

Post, J.R., Parkinson, E.A., and Johnston, N.T. 1998. Spatial and temporal variation in risk to

piscivory of age-0 rainbow trout: patterns and population level consequences. T. Am. Fish. Soc.

127: 932–942.

Salvanes, A.G.V., and Braithwaite, V. 2006. The need to understand the behaviour of fish

reared for mariculture or restocking. ICES J. Mar. Sci. **63**: 346–354.

Sawara, Y., Azuma, N., Fukui, K., Demachi, G., and Sakuyama, M. 1990. Feeding activity of

the Grey Heron Ardea cinerea in tidal and non-tidal environments. Jap. J. Ornithol. **39**: 45–52.

Silveira, L., Jacomo, A.T.A., and Diniz-Filho, J.A.F. 2003. Camera trap, line transect census

and track surveys: a comparative evaluation. Biol. Conserv. **114**: 351–355.

Steinmetz, J., Kohler, S.L., and Soluk, D.A. 2003. Birds are overlooked top predators in

aquatic food webs. Ecology **84**: 1324–1328.

Sundström, L.F., Petersson, E., Höjesjö, J., Johnsson, J.I., and Järvi, T. 2004. Hatchery

selection promotes boldness in newly hatched brown trout (*Salmo trutta*): implications for

dominance. Behav. Ecol. **15**(2): 192–198.

Stewart, D.C., Middlemas, S.J., Gardiner, W.R., Mackay, S., and Armstrong, J.D. 2005. Diet

and prey selection of cormorants (*Phalacrocorax carbo*) at Loch Leven, a major stocked trout

fishery. J. Zool. **267**: 191–201.

Tojo, H. 1996. Habitat Selection, Foraging Behavior and Prey of Five Heron Species in Japan.

Jap. J. Ornithol. **45**: 141–158.

Wegge, P., Pokheral, G.P., and Jnawali, S.R. 2004. Effects of trapping effort and trap shyness

on estimates of tiger abundance from camera trap studies. Anim. Conserv. **7**: 251–256.

Wood, C.C. 1987. Predation on juvenile Pacific salmon by the common merganser (*Merges*

merganser) on eastern Vancouver Island I: Predation during the seaward migration. Can. J. Fish.

Aquat. Sci. **44**: 941–949.

Figure captions

Fig. 1. a and b: outdoor FRP circular tank (a) and semi-natural stream (b) at NRIFS in

Nikko. c and d: examples of photo of grey heron in the pool (c) and in the riffle (d) in

the semi-natural stream.

Fig. 2. Survival rate of masu salmon (*Oncorhynchus masou*) with different body sizes

in outdoor tanks.

Fig. 3. Correlation between the number of taken photos of grey heron (*Ardea cinerea*)

and the number of masu salmon (*Oncorhynchus masou*) in outdoor tanks.

Fig. 4. The sum of masu salmons (*Oncorhynchus masou*), 75–100 and 135–160 mm,

captured at the each section in the semi-natural stream. Broken line indicates the

expected surviving proportion (we hypothesized a uniform 1:1:1:1 ratio). Asterisk

denotes a significant deviation from the average value (** $p < 0.010$). Each section

(Sec.) was named from the upper most section in the stream, Sec. 1, Sec. 2, Sec. 3 and

Sec. 4, Sec. 2 was the stocking location.

Fig. 5. Number of photos of grey heron (*Ardea cinerea*) in each study section of

semi-natural stream.

Fig. 6. The proportion of the photos that showed grey heron located around the pool

or riffle in each study section of the semi-natural stream. Asterisk denotes a significant

difference (∗ $p < 0.050$, ∗∗ $p < 0.010$).

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5