Predation of stocked salmon by riparian wildlife in semi-natural stream

Journal:	*Marine and Freshwater Research*
Manuscript ID	Draft
Manuscript Type:	Research paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Miyamoto, Kouta; National Research Institute of Fisheries Science, Japan Fisheries Research and Education Agency, Squires, Theodore Araki, Hitoshi
Keyword:	fisheries, birds, ecology

SCHOLARONE™
Manuscripts

Predation of stocked salmon by riparian wildlife in semi-natural stream

Running head: Predation of stocked salmon by riparian wildlife

Kouta Miyamoto[A,B,D], Theodore E. Squires[B], and Hitoshi Araki[C]

[A]National Research Institute of Fisheries Science, Japan Fisheries Research and Education

Agency, Nikko, Tochigi 321-1661, Japan

[B]Graduate school of Agriculture, Hokkaido University, N9W9 Kita, Sapporo 060-8589, Japan

[C]Research Faculty of Agriculture, Hokkaido University, N9W9 Kita, Sapporo 060-8589, Japan

[D]Corresponding author. E-mail: mkouta@affrc.go.jp

Abstract.   Predation after release is one of the major concerns of hatchery fish

conservation and propagation. However, the relationship among the size of hatchery fish, the

predator species, and their behaviors in natural environments is largely unknown. To understand

the relationship, we conducted predation experiments in outdoor tanks and a semi-natural stream

with exposure to local predators. Two different ranges of fork lengths of masu salmon

(*Oncorhynchus masou*) were examined as prey sizes. Camera trap data showed that grey herons

(*Ardea cinerea*) were the primary predator animal in the system, and that most herons utilized

shallow areas in the morning or evening. Increasing the density of stocked salmon brought in

more grey herons. More importantly, predation by grey herons resulted in the survival rate of

larger salmon being significantly lower than that of the smaller salmon. Our results suggest that

it is important to understand local predators, adjust the optimum body size of hatchery fish at

release, and choose the appropriate stocking site and time of day for maximizing the

effectiveness of fish stocking.

Additional key words: predator-prey interaction, hatchery management, size-selective predation,

piscivorous water bird

Introduction

Predation by riparian wildlife is widely recognized as a key factor influencing the survival of

fishes in stream ecosystems (Kruuk, 1995; Draulans 1987; Roby *et al.* 2003; Steinmetz *et al.*

2003; Sahashi and Yoshiyama 2015). Salmonid fish populations reduced by riparian wildlife

have been reported, not only for stocked populations, but also for wild populations (e.g.

Osterback *et al.* 2013; Frechette *et al.* 2015). It is also noteworthy that stocked salmonids are

often intensively preyed upon by birds (Wood, 1987; Martel and Dill, 1995; Harvey and

Nakamoto, 2013) and mammals (Roberts and Garcia de Leaniz, 2011). Hatchery-reared

salmonid fishes are routinely stocked in natural waterways as part of conservation efforts or

stock enhancement (Brown and Laland 2003; Salvanes and Braithwaite 2006; Fraser 2008).

Therefore, if we implement a management strategy that decreases local predation pressures, then

effective conservation or stock enhancement methods should be improved.

40   Studies examining the predator-prey interaction between riparian animals and fish have

been generally based on either investigating predation pressure of riparian animals without

observing fish response (Draulans 1987; Post *et al*. 1998; Harvey and Nakamoto 2013) or

investigating the mortality rate of fish without observing the behavior of predators (Penaluna *et*

al. 2015). However, the predation effect is influenced by many factors: the condition of fish

(Hostetter *et al*. 2012), fish migration (Roberts *et al*. 2009), the number of predators present, and

the duration of their sojourn (Gawlik, 2002, Steinmetz *et al*. 2003). Thus, methodological

improvements which decrease predation pressure on stocked fish require a profound

understanding of both predator and prey behaviors. Additionally, as food size preference depends

on predator species (Carss and Marquiss 1991, Sogard 1997) and the ability of fish to avoid

predators depends on fish size (Dill and Fraser 1984), we need to consider the size effect of prey

fish.

52   To evaluate the impacts of predation by riparian animals, it is necessary to evaluate the

relationships among the predator species, their behaviors, and their preferred size of prey in a

place that allows for free movement of all parties. However, the number of studies on this

relationship is limited due to the difficulty in identifying species of predatory riparian animals

and observing their predation behavior directly. These studies are further complicated due to

complex behaviors exhibited by many predator species, such as individual movement for feeding

and nesting needs (Collis *et al*. 2002), movement due to changes in prey densities (Kushlan,

1976; Gawlik, 2002), and movement resulting from human disturbance (Klein, 1993).

Recently, camera trappings that use fixed cameras, triggered by infrared sensors, to

'trap' images of passing animals, provide the opportunity to collect information including

animals' species, time of appearance in each day, appearance frequency, and duration of sojourn

(Silveira *et al*. 2003; Wegge *et al*. 2004). Because camera trapping is a non-invasive technique, it

causes minimal environmental disturbance (Henschel and Ray 2003; Silveira *et al*. 2003). In this

study, we conducted a predation test using camera traps with outdoor experimental tanks and a

semi-natural stream to investigate the relationships between the different predator species, their

behavior, and the size of stocked salmon.

Materials and Methods

Test fish

The test fish used in the experiments were hatchery-reared masu salmon (*Oncorhynchus masou*)

from the Shobu-shimizu River and the Jigoku River in Tochigi Prefecture, Japan (approximately

36°45'N, 139°27'E). Eggs were obtained from mature adults that had migrated into the

Shobu-shimizu River and the Jigoku River, and were handled according to the standard hatchery

procedures directed by the National Research Institute of Fisheries Science (NRIFS) facility at

Nikko. Young of the year (YOY) ($n = 305$) and one year old (OYO) ($n = 305$) salmon were used

in this study. The YOY and OYO salmon were graded from five rearing tanks (50 cm width ×

120 cm length × 20 cm depth, the supply amount of water is set at 18 L/min) and a large tank (3

m width × 1.5 m length × 0.9 m depth, the supply amount of water is set at 150 L/min)

respectively. The YOY fish used were 75–100 mm fish (75–100 mm FL-group, hereafter) and

the OYO used were 135–160 mm fish (135–160 mm FL-group, hereafter). The body size range

of the YOY salmon is bigger than that of general stocked YOY salmon in Japan (Sahashi *et al*.

2015), however the body size ranges are within those of the masu salmon that are stocked in the

Shobu-shimizu River and the Jigoku River. In addition, the size range of YOY masu salmon are

recommended as an effective body size for preventing predation by piscivorous fish in rivers

around central Japan (Miyamoto and Araki 2017). Before the start of the study, fish were fed

daily food rations (commercial trout pellets) equal to 1.5–2.0% of their estimated body weight.

Tank experiment

We conducted predation tests in four outdoor tanks for three days from August to September,

2013. Four Fibre-Reinforced Plastic (FRP) circular tanks (120-cm diameter × 15-cm high) at

NRIFS in Nikko were used in a forested and grassy area and each had a thin layer of natural

gravel (2–7-cm gravel and about 13-cm cobble substrate) on the bottom (Fig. 1a). A 10-cm high

and 8-cm roof fence that was installed on each tank, both to provide the fish with cover and

prevented them from jumping out. A camera (Trophy Cam HD, Bushnell, Overland Park, KS,

USA) was set in the south side of each tank to monitor the whole tank. The presence of cover

and gravel substrate are set so that masu salmon can show near natural behavior, allowing hiding

and escaping (Miyamoto 2016a), thus minimizing experimental stress. During the study days the

stocked fish were fed daily food rations (commercial trout pellets) equal to around 2.0% of their

body weight. Spring water, $10.2 \pm 0.3°C$ (mean $\pm$ SD) was introduced into each tank at a rate of

6 L/min. Two different fork lengths (FL) of masu salmon were used as the size of prey fish,

75–100 mm FL-group (mean $\pm$ SD, 91.9 ± 7.0 mm) and 135–160 mm FL-group (145.8 ± 6.8

mm).

To evaluate the size-selective predation risk, 30 fish (15 from each FL-group) were

placed in each tank. The number of fish of each FL-group that survived was counted each day for

three days. The survival rate of salmon was then compared between the two FL-groups. To

identify the predator animals, the number of photos taken by camera trap (described in the

section below) was counted.

Next, to investigate the relationship between the number of stocked salmon and the

frequency in which predator animals appeared, three FRP circular tanks were used. The test fish

were divided into 10, 5, and 0 individuals and stocked randomly in each of the three tanks. Three

days later, the number of photos of potential predators taken by camera trap (described in the

section below) was counted for each tank. This trial was replicated three times for each

FL-group.

Predation test in a semi-natural stream

The predation test was conducted for 20 days from August to September, 2013 using a

semi-natural stream (mean ± standard deviation [SD]: 120 m long, 95.8 ± 4.3-cm wide; 2%

gradient drain) in a forested and grassy area at NRIFS (Fig. 1b). The stream was constructed of

stone, wood, and soil. Spring water, at 10.2 ± 0.3°C (mean ± SD) was drained at 18 L/s. The

stream was forked to allow free passage into four 30 meter sections (Sec. 1-4 from upper to

lower). Each section contained three pools and three riffles. The pools were about 80-cm long

and about 60-cm wide with maximum depth of approximately 40-cm. Woody debris (c. 70-cm

long and c. 20-cm wide) were placed in the sides of each pool. The riffles were 9–12-cm deep

and contained a mixture of 2–7-cm gravel and about 13-cm cobble substrate. A 50-cm waterfall

was built in the top of the stream with two metal gates (1-cm × 2-cm mesh) at the top to prevent

fish from escaping. The water drain was separated by a metal mesh gate (1-cm × 2-cm mesh)

with the bottom quarter of the gate covered with plastic mesh (8-mm × 8-mm mesh). A total of

400 fish (200 from each FL-group) were stocked in Sec. 2 after measuring their fork length and

body weight. At the end of the study period, a backpack electrofishing unit (Smith-Root,

Vancouver, WA, USA) was used to remove and count the fish that had survived. The electric

shock was repeated until the fish count was zero twice in a row.

To investigate the distribution of salmon in the stream, the fish in each section were

closed in by fish block nets (8-mm × 8-mm mesh) at the end of the experiment to prevent them

from moving between the sections. When fish were caught by electrofishing, each fish was

lightly anesthetized with 100 ppm 2-phenoxyethanol (Wako Pure Chemical Industries, Tokyo,

Japan), its fork length and body weight were measured, and the section that the fish was caught

in was recorded. To identify the predator animals, 3 cameras (Trophy Cam HD, Bushnell,

Overland Park, KS, USA) were arranged in each section and placed at 10-m intervals along the

riverside to monitor both sides of the stream, and the number of photos that were taken by

camera traps (described in the section below) was counted for each section. In addition, to

estimate the predation behavior of wild animals, the position of potential predators (at pool or

riffle) in photos was recorded. Then, to investigate the position that predators utilized, the

proportion of predators located around pools or riffles was calculated in each section, then the

average proportions were compared.

Camera trap

To assess predator encounters during day and night, potential predators were recoded using

motion and infrared sensor camera traps in the tank experiments and the semi-natural stream

experiment. Each camera was mounted on a wooden stake so that the camera was about 50 cm

above the water's surface. Cameras were triggered with a passive infrared motion sensor; the

camera was set to wait 15 seconds after an initial trigger entered its sensor range before

attempting to detect additional triggers. To identify predators and estimate the frequency of their

visits to the study site, all the photos containing potential predators were checked by KM. For

some ambiguous species identifications, additional checking was performed by a local wildlife

expert Dr. T. Takeda from Nikko National Park. When more than one potential predator was

captured in a photo, the species and their number was recorded. In addition, the number of

photos that showed predators capturing or eating fish was counted, and the predator species was

recorded.

Statistical analyses

In the tank tests, Two-way ANOVA was used to determine the effects of study days and fish fork

length on the survival of salmon. Additionally, a second set of ANOVA tests were used to

determine how the number of salmon, and salmon size, effected the number of photos containing

riparian predators in the stream test. One-way ANOVA was used to compare the average

proportions of the predators located around pools and riffles in the stream. Count data that

contain zeros was log $(x + 0.5)$ transformed prior to analysis (Yamamura 1999), or, when

proportions were tested, data were arcsin $\sqrt{x}$ transformed. Tukey HSD test was used as a

post-hoc test.

170   For the semi-natural stream experiment, Pearson's chi-square test was used to compare

the proportion of the number of fish captured at each section. However, the omnibus chi-square

value does not specify which combination of categories contributes to statistical significance;

thus, the adjusted standardized residuals (ASR) was used for each value to determine

discrepancies between the observed and expected value (Haberman, 1973). $|ASR| > 1.96$ and >

2.56 show $P < 0.05$ and $P < 0.01$, respectively. A $P < 0.05$ was considered significant in all

statistical analyses. Data analyses were generated using IBM SPSS software (version 21.0 of

IBM SPSS Statistics for Windows).

Results

Tank experiment

In the tank experiment, cameras captured grey heron (*Ardea cinerea*), Japanese marten (*Martes*

melampus), raccoon dog (*Nyctereutes procyonoides*), and large-billed crow (*Corvus*

macrorhynchos). The total number of photos containing potential predators was 186 (including

three photos with ambiguous species, which were identified by the local wildlife expert). A total

of 173 photos of grey heron were taken and there were six photos or fewer of each of the other

animals (Japanese marten: three times, raccoon dog: four times, large-billed crow: six times).

Twenty one photos showed grey herons capturing prey fish. Therefore, in tank experiments, grey

herons were regarded as the main predator.

With regard to the size-selective predation risk, both the length of the study days and the

fork length of the stocked salmon had significant effects on the fish survival (Period, $F_{2, 18}$ =

10.56, $P < 0.010$; Salmon length, $F_{1, 18}$ = 36.67, $P < 0.001$; Period × Salmon length, $F_{2, 18}$ = 0.18,

$P = 0.833$) (Table 1). The fish survival rate was significantly higher in the 75-100 mm FL-group

than in the 135-160 mm FL-group during the study days (Tukey HSD test, all shows $P < 0.050$)

(Fig. 2).

195   In the Two-way ANOVA test for frequency of predator appearance, only the number of

fish in the tank had a significant effect on the number of photos containing grey herons (Number

of fish, $F_{2, 12}$ = 149.26, $P < 0.001$; Fork length, $F_{1, 12}$ = 2.48, $P = 0.141$; Number of fish × Fork

length, $F_{2, 12}$ = 0.02, $P = 0.981$) (Table 2) and the number of photos containing grey herons

increased significantly when the number of fish in the tanks was large (Tukey HSD test, all

shows $P < 0.001$) (Fig. 3). At the end of the three-day experiment, there were no surviving

salmon.

Stream experiment

The overall survival of the fish in this experiment was 18.0% ($n = 72$). The survival of them was

significantly higher for the 75–100 mm FL-group (33.0%, $n = 66$) than for the 135-160 mm

FL-group (3.0%, $n = 6$) (Peason's χ^2 test, $\chi^2 = 60.98$, $df = 1$, $p < 0.001$).

207   There was no significant difference between the proportion of fish that survived in each

section between both fish groups (Peason's χ^2 test, $\chi^2 = 0.77$, $df = 3$, $p = 0.86$). Comparing the

actual proportion of surviving fish (75-100 and 135-160 mm) in each section to the expected

surviving proportion (we hypothesized a uniform 1:1:1:1 ratio) showed there was a significant

difference (Peason's χ^2 test, $\chi^2 = 15.43$, $df = 3$, $p < 0.001$). Accordingly, the number of fish

surviving in Sec. 2 was significantly higher than the average number of fish capture in each

section (ASR $= 3.4$, $p < 0.010$) (Fig. 4).

214   Cameras captured grey heron, brown dipper (*Cinclus pallasii*), Japanese red fox (*Vulpes*

vulpes japonica) and large-billed crow during the predation test in the semi-natural stream. The

total number of photos containing potential predators was 470. Among them, 455 contained grey

heron. There were fewer than 10 photos of other animals (brown dipper: one times, Japanese red

fox: seven times, large-billed crow: seven times). The most grey heron photos ($n = 104$) were

taken at Sec. 2 on day-5 (Fig. 5). Of all the pictures that contained grey herons, 76.5 % were

taken at Sec.2. Five photos had two grey herons in each picture, and two photos had three grey

herons in each picture, all photos with multiple herons were taken in Sec. 2 between day-4 and

day-7 after stocking. Seventeen photos showed grey herons capturing prey fish. Grey herons

were most frequently photographed in the mornings and evenings.

224   In order to investigate the position that grey herons utilized, 412 photos were analyzed

(455 minus 43 photos for undistinguishable positioning). The average proportion of photos

showing grey heron around the pools (6.88 ± 10.68 %) (Fig. 1c) was significantly lower than

those around the riffles (93.13 ± 10.68 %) (Fig. 1d) ($F_{1, 6} = 53.13$, $P < 0.010$) (Fig. 6). The

overall proportions of photos taken of grey herons in the morning (3:00-9:00) and in the evening

(15:00-21:00) were 49.3% (n = 224) and 47.9% (n = 218), respectively.

Discussion

In the 20-day stream experiment, 82% of fish were lost and that grey heron was the most

frequently visiting predator. It has been previously reported that salmonid populations can be

seriously damaged by avian predation (Feltham 1995; Stewart et al.2005; Evans et al. 2016), but

quantitative assessments of the impact of grey heron on river fisheries is rare (Hodgens et al.

2004). Our results suggest that at least in our experimental settings, grey herons can significantly

reduce salmonid populations by consuming juvenile fish. It is important to note that water depth

can strongly influence wading birds' selection of foraging habitat (Master et al. 2005, Gawlik

and Crozier 2007). The length of a grey heron's legs restricts the maximum depth at which they

can forage up to 16 cm deep (Ntiamoa-Baidu et al. 1998), thereby limiting the habitat suitable

for hunting. Our observation that grey herons showed a significant preference for the riffle,

rather than the pool, in the semi-natural stream is consistent with previous studies. Thus, where

local salmonids reside in shallow water, it is reasonable to expect that grey herons exert a high

predation pressure.

245   On the first day of the tank experiments, grey herons preyed upon large salmon more

often than upon small salmon. In both the tank and the stream experiments, the survival rate of

large salmon was less than that of small salmon. Birds of the heron family (Ardeidae) including

the grey heron are reported to show a preference for larger mosquito fish *Gambusia affinis*

(Britton and Moser 1982) and sailfin molly *Poecilia latipinna* (Trexler *et al*. 1994) as prey. It is

suggested that larger fish are easier for avian predators to detect than small ones (Eriksson 1985,

Magnhagen 1988). After the second day of the tank experiment, however, both sizes of salmon

showed similar levels of decline on their survival rate. One possible explanation for this result is

the decreased opportunities to prey upon larger fish. Our results suggest that size-selective

predation by grey herons depends on the density of preferred prey size, and it appears that

size-selective predation occurred in the stream experiment as well as in the tank experiments.

Therefore, if fish with different body sizes are stocked in rivers, it might be important to consider

how size composition will potentially influence the behavior of predators.

It is established that larger fish are safer from fish-on-fish predation than smaller fish

(Peterson and Wroblewski 1984, Houde 1987, Miller *et al*. 1988, Miyamoto and Araki 2017).

One potential cause for the higher survival rate of larger fish is improved swimming ability that

allows them to better avoid predators as they grow (Beamish 1978; Lundvall *et al*. 1999).

However, the grey heron is an ambush predator that usually stands upright and waits for a fish to

approach (Tojo, 1996), so the swimming ability that a fish possesses to avoid aquatic predators

has significantly limited benefits to them in this case (Miyamoto 2016b), and large size might

make fish more detectable. Therefore, to decrease predation risk by wild animals, we suggest that

hatchery managers choose an optimum body size of released fish based on the predator species

inhabiting the stocking area. Additionally, to block the vision of grey herons or prevent access, it

can be recommended to stock fish in a structurally complex place that has a lot of woody debris

and/or aquatic vegetation (Diehl 1988) if many grey herons are around during the stocking

season.

Generally, in salmonids, larger individuals avoid exposure to a predatory threat and

reduce growth rate, whereas smaller individuals are less cautious and maintain their growth rate

even in the presence of a threat (Reinhardt 1999). Based on this behavior, it is expected that the

OYO salmon are more difficult for grey heron to catch than the YOY salmon. In contrast to this,

it was also reported that the hatchery environment selects for bolder individuals that spend more

time in open areas and are more active than wild specimens (Sundström *et al*. 2004). So, OYO

salmon that spent a longer time than the YOY salmon in the hatchery may display a higher

tendency for bold behavior (Roberts *et al*. 2014). Thus, to fully understand the predation risks for

stocked salmon, further studies that estimate the individual behavior of fish (e.g. bold and shy)

and improve the maladapted behavior of fish (Berejikian *et al*. 1999, Roberts *et al*. 2011) will be

needed.

In the tank experiment where different numbers of salmon were stocked, the number of

grey herons present or the duration of their sojourn at the tanks was positively correlated to the

number of stocked fish. In other studies, large numbers of fish have been observed to remain

close to where they were stocked (Cresswell 1981), and piscivorous water birds have displayed

similar density and temporal trends in response to stocking (Draulans 1987; Gawlik 2002). This

suggests that for the duration of our stream experiment many salmon stayed in Sec. 2, as multiple

grey herons simultaneously appeared frequently in that section. We also captured more fish in

Sec. 2 by electrofishing than in any other sections on the final day of the stream experiment.

These results indicate that most stocked fish gathered without migrating from the initial stocking

site and continued to be preyed upon there. This situation may have serious implications for the

conservation and propagation of salmonids and it is important to further investigate the spatial

and temporal relationships between salmonids and their predators.

294   The results of this study showed that the grey heron has a tendency to avoid deep water

and foraging in the middle of day and night. In contrast, it previously was reported that grey

herons forage constantly during the daytime (Richner 1986; Sawara *et al*. 1990), so human

activities (Klein, 1993) and other feeding sites (Richner 1986) in the surrounding environment

could have had an effect on the observed behavior. Grey herons infrequently forage at night, and

it has been suggested that such feeding behavior is inefficient (Sawara *et al*. 1990). Therefore, to

decrease the predation pressure of grey herons, just after the fish were released, it might be wise

to stock fish in a deep area or at night (Roberts *et al*. 2009), or both. In this way, if we can

develop techniques to mitigate predatory damage to fisheries by investigating the relationship

between stocked fish and predatory animals, it will not only help the conservation and

propagation of stocked fish, but also support the conservation of wild animals as well.

Acknowledgements

We would like to thank Kouji Mutou, Hidefumi Nakamura and Masaharu Murakami of the

National Research Institute of Fisheries Science for their help with the care of the fish. We thank

Dr Tsutomu Takeda of Nikko Yumoto Visitor Center of Nikko National Park for his help with

the identification of wildlife. We are grateful to Predrag Davidovic and Petar Ilic for critically

reading the manuscript and providing valuable comments. Dr Tomoyuki Nakamura of the

NRIFS helped us with valuable advice. This work was supported by JSPS KAKENHI Grant

Number JP26292102.

References

Beamish, F. W. H. (1978). Swimming capacity. In ´Fish physiology, Vol. VII´. (Eds W. S. Hoar

and D.J. Randall.) pp. 101−187. (Academic Press, New York.)

Berejikian, B. A., Smith, R. J. F., Tezak, E. P., Schroder, S. L., and Knudsen, C. M. (1999).

Chemical alarm signals and complex hatchery rearing habitats affect antipredator behavior and

survival of chinook salmon (*Oncorhynchus tshawytscha*) juveniles. *Canadian Journal of*

Fisheries and Aquatic Sciences **56**, 830−838.

Brown, C., and Laland, K. N. (2003). Social learning in fishes: a review. Brown, C. and

Laland, K.N., 2003. Social learning in fishes: a review. *Fish and fisheries* **4**, 280–288.

Carss, D. N., and Marquiss, M. (1991). Avian predation at farmed and natural fisheries. In

´Interaction between fisheries and environment´. (Eds M. C. Lucas, I. Diak, and L. Laird. Proc.

Institute of Fisheries Management, 22nd Annual Study Course.) pp. 179–196. (University of

Aberdeen. UK.)

Collis, K., Roby, D. D., Craig, D. P., Adamany, S., Adkins, J. Y., and Lyons, D. E. (2002).

Colony size and diet composition of piscivorous waterbirds on the lower. Columbia River:

Implications for losses of juvenile salmonids to avian predation. *Transactions of the American*

Fisheries Society **131**, 537–550.

Cresswell, R. C. (1981). Post‐stocking movements and recapture of hatchery‐reared trout

released into flowing waters−a review. *Journal of Fish Biology* **18**, 429–442.

Diehl, S. (1988). Foraging efficiency of three freshwater fishes: effects of structural

complexity and light. *Oikos* **53**, 207–214.

Dill, L. M., and Fraser, A. H. G. (1984). Risk of predation and the feeding behavior of juvenile

coho salmon (*Oncorhynchus kisutch*). *Behavioral Ecology and Sociobiology* **16**, 65–71.

Draulans, D. (1987). The effect of prey density on foraging behaviour and success of adult and

first-year grey herons (*Ardea cinerea*). *The Journal of Animal Ecology*, 479–493.

Eriksson, M. O. (1985). Prey detectability for fish-eating birds in relation to fish density and

water transparency. *Ornis Scandinavica* **16**, 1–7.

Evans, A. F., Payton, Q., Turecek, A., Cramer, B., Collis, K., Roby, D. D., Loschl, P. J.,

Sullivan, L., Skalski, J., Weiland, M. and Dotson, C. (2016). Avian Predation on Juvenile

Salmonids: Spatial and Temporal Analysis Based on Acoustic and Passive Integrated

Transponder Tags. *Transactions of the American Fisheries Society* **145**, 860–877.

Feltham, M. J. (1995). Consumption of Atlantic salmon smolts and parr by goosanders:

estimates from doubly ‐ labelled water measurements of captive birds released on two Scottish

rivers. *Journal of Fish Biology* **46**, 273–281.

Fraser, D. J. (2008). How well can captive breeding programs conserve biodiversity? A review

of salmonids. *Evolutionary Applications* **1**, 535–586.

Frechette, D., Osterback, A. M. K., Hayes, S. A., Moore, J. W., Shaffer, S. A., Pavelka, M.,

Winchell, C., and Harvey, J. T. (2015). Assessing the Relationship between Gulls Larus spp. and

Pacific Salmon in Central California Using Radiotelemetry. *North American Journal of Fisheries*

Management **35**, 775–788.

Gawlik, D. E. (2002). The effects of prey availability on the numerical response of wading

birds. *Ecological monographs* **72**, 329–346.

Harvey, B. C., and Nakamoto, R. J. (2013). Seasonal and Among-Stream Variation in Predator

Encounter Rates for Fish Prey. *Transactions of the American Fisheries Society* **142**, 621–627.

Henschel, P., and Ray, J. (2003). ´Leopards in African Rainforests: Survey and Monitoring

Techniques.´ (Wildlife Conservation Society, New York)

Hodgens, L. S., Blumenshine, S. C. and Bednarz, J. C. (2004). Great blue heron predation on

stocked rainbow trout in an Arkansas tailwater fishery. *North American Journal of Fisheries*

Management **24**, 63–75.

Hostetter, N. J., Evans, A. F., Roby, D. D. and Collis, K. (2012). Susceptibility of juvenile

steelhead to avian predation: the influence of individual fish characteristics and river conditions.

Transactions of the American Fisheries Society **141**, 1586–1599.

Houde, E. D. (1987). Fish early life dynamics and recruitment variability. *Transactions of the*

American Fisheries Society **2**, 17–29.

Klein, M. L. (1993). Waterbird behavioral responses to human disturbances. *Wildlife Society*

Bulletin **21**, 31–39.

Kruuk, H. (1995). ´Wild Otters. Predation and populations.´ (Oxford University Press. New

York.)

Kushlan, J. A. (1976). Wading bird predation in a seasonally fluctuating pond. *The Auk* **93**,

464–476.

Lundvall, D., Svanbäck, R., Persson, L., and Byström, P. (1999). Size-dependent predation in

piscivores: interactions between predator foraging and prey avoidance abilities. *Canadian*

Journal of Fisheries and Aquatic Sciences **56**, 1285–1292.

Magnhagen, C. (1988). Predation risk and foraging in juvenile pink (*Oncorhynchus*

gorbuscha) and chum salmon (*O. keta*). *Canadian Journal of Fisheries and Aquatic Sciences* **45**,

592–596.

Martel, G., and Dill, L. M. (1995). Influence of movement by coho salmon (*Oncorhynchus*

kisutch) parr on their detection by common mergansers (*Mergus merganser*). *Ethology* **99**,

139–149.

Master, T. l., Lesier, J. K. Bennett, K. A. Bretsch, J. K., and Wolfe H. J. (2005). Patch selection

by Snowy Egrets. *Colonial Waterbirds* **28**, 220–224.

Miller, T. J., Crowder, L. B., Rice, J. A., and Marschall, E. A. (1988). Larval size and

recruitment mechanism in fisheries: toward a conceptual framework. *Canadian Journal of*

Fisheries and Aquatic Sciences **45**, 1657–1670.

Miyamoto, K. (2016a). Effect of visual and chemical stimuli on predator avoidance behavior

in juvenile masu salmon *Oncorhynchus masou*. *Aquaculture Science* **64**, 43–51.

Miyamoto, K. (2016b). Effects of body color luminance and behavioral characteristics on

predation risk in salmonid fishes. *Hydrobiologia* **783**, 249–256.

Miyamoto, K., and Araki, H. (2017). Differentiated predation risk on hatchery-reared juvenile

masu salmon by white-spotted charr with different body sizes. *Fisheries Science* **83**, 245–250.

Ntiamoa-Baidu Y., Piersma, T., Wiersma, P., Poot, M., Battley, P., and Gordon, C. (1998). Water

depth selection, daily feeding routines and diets of waterbirds in coastal lagoons in Ghana. *Ibis*

140, 89–103.

Osterback, A. M. K., Frechette, D. M., Shelton, A. O., Hayes, S. A., Bond, M. H., Shaffer S.

A., and Moore, J. W. (2013). High predation on small populations: avian predation on imperiled

salmonids. *Ecosphere* **4**, 116.

Penaluna, B. E., Jason, B. D., and Noakes, D. L. (2015). Instream cover and shade mediate

avian predation on trout in semi-natural streams. *Ecology of Freshwater Fish* **25**, 405–411.

Peterson, I., and Wroblewski, J. S. (1984). Mortality rate of fishes in the pelagic ecosystem.

Canadian Journal of Fisheries and Aquatic Sciences **41**, 1117–1120.

Post, J. R., Parkinson, E. A., and Johnston, N. T. (1998). Spatial and temporal variation in risk

to piscivory of age-0 rainbow trout: patterns and population level consequences. *Transaction of*

the American Fisheries Society **127**, 932–942.

Reinhardt, U. G. (1999). Predation risk breaks size-dependent dominance in juvenile coho

salmon (*Oncorhynchus kisutch*) and provides growth opportunities for risk-prone individuals.

Canadian Journal of Fisheries and Aquatic Sciences **56**, 1206–1212.

Richner, H. (1986). Winter feeding strategies of individually marked herons. *Animal*

Behaviour **34**, 881–886.

Roberts, L. J., and Garcia de Leaniz, C. (2011). Something smells fishy: predator-naïve

salmon use diet cues, not kairomones, to recognize a sympatric mammalian predator. *Animal*

Behaviour **82**, 619–625.

Roberts, L. J., Taylor J., and Garcia de Leaniz, C. (2011). Environmental enrichment reduces

maladaptive risk-taking behavior in salmon reared for conservation. Biological Conservation **144**,

1972–1979.

Roberts, L. J., Taylor, J., Gough, P. J., Forman, D. W., and de Garcia de Leaniz, C. (2009).

Night stocking facilitates nocturnal migration of hatchery‐reared Atlantic salmon, *Salmo salar*,

smolts. *Fisheries Management and Ecology* **16**, 10–13.

Roberts, L. J., Taylor, J., Gough, P. J., Forman, D. W., and Garcia de Leaniz, C. (2014). Silver

spoons in the rough: can environmental enrichment improve survival of hatchery Atlantic salmon

Salmo salar in the wild? *Journal of fish biology* **85**, 1972–1991.

Roby, D. D., Lyons, D. E., Craig, D. P., Collis, K., and Visser, G. H. (2003). Quantifying the

effect of predators on endangered species using a bioenergetics approach: Caspian terns and

juvenile salmonids in the Columbia River estuary. *Canadian Journal of Zoology* **81**, 250–265.

Sahashi, G. and Yoshiyama, T. (2015). A hump-shaped relationship between migration distance

and adult pink salmon morphology suggests interactive effects of migration costs and bear

predation. *Canadian Journal of Fisheries and Aquatic Sciences* **73**, 427–435.

Sahashi, G., Morita, K., Ohnuki, T. and Ohkuma, K. (2015). An evaluation of the contribution

of hatchery stocking on population density and biomass: a lesson from masu salmon juveniles

within a Japanese river system. *Fisheries Management and Ecology* **22,** 371–378.

Salvanes, A. G. V., and Braithwaite, V. (2006). The need to understand the behaviour of fish

reared for mariculture or restocking. *ICES Journal of Marine Science: Journal du Conseil* **63**,

346–354.

Sawara, Y., Azuma, N., Fukui, K., Demachi, G., and Sakuyama, M. (1990). Feeding activity of

the Grey Heron Ardea cinerea in tidal and non-tidal environments. *Japanese Journal of*

Ornithology **39**, 45–52.

Silveira, L., Jacomo, A. T. A., and Diniz-Filho, J. A. F. (2003). Camera trap, line transect

census and track surveys: a comparative evaluation. *Biological Conservation* **114**, 351–355.

Sogard, S. M. (1997). Size-selective mortality in the juvenile stage of teleost fishes: a review.

Bulletin of Marine Science **60**, 1129–1157.

Steinmetz, J., Kohler, S. L., and Soluk, D. A. (2003). Birds are overlooked top predators in

aquatic food webs. *Ecology* **84**, 1324–1328.

Stewart, D. C., Middlemas, S. J., Gardiner, W. R., Mackay, S., and Armstrong, J. D. (2005).

Diet and prey selection of cormorants (*Phalacrocorax carbo*) at Loch Leven, a major stocked

trout fishery. *Journal of Zoology* **267**, 191–201.

Sundström, L. F., Petersson, E., Höjesjö, J., Johnsson, J. I., and Järvi, T. (2004). Hatchery

selection promotes boldness in newly hatched brown trout (*Salmo trutta*): implications for

dominance. *Behavioral Ecology* **15**, 192–198.

Tojo, H. (1996). Habitat Selection, Foraging Behavior and Prey of Five Heron Species in

Japan. *Japanese Journal of Ornithology* **45**, 141–158.

Wegge, P., Pokheral, G. P., and Jnawali, S. R. (2004). Effects of trapping effort and trap

shyness on estimates of tiger abundance from camera trap studies. *Animal Conservation* **7**,

251–256.

Wood, C. C. (1987). Predation on juvenile Pacific salmon by the common merganser (*Merges*

merganser) on eastern Vancouver Island I: Predation during the seaward migration. *Canadian*

Journal of Fisheries and Aquatic Sciences **44**, 941–949.

Yamamura, K. (1999). Transformation using (x+ 0.5) to stabilize the variance of populations.

Researches on Population Ecology **41**, 229–234.

Figure captions

Fig. 1. a and b: outdoor fibre-reinforced plastic circular tank (a) and semi-natural stream (b) at

National Research Institute of Fisheries Science in Nikko. c and d: examples of photo of grey

heron in the pool (c) and in the riffle (d) in the semi-natural stream.

Fig. 2. Survival rate of masu salmon (*Oncorhynchus masou*) with different body sizes groups

in outdoor tanks. Circles and solid line: using 75−100 mm salmon; black circles and dashed line:

using 135−160 mm salmon. Data are means ± standard error.

Fig. 3. Relationship between the number of taken photos of grey heron (*Ardea cinerea*) and

the number of masu salmon (*Oncorhynchus masou*) in outdoor tanks. Circles and solid line:

using 75−100 mm salmon; black circles and dashed line: using 135−160 mm salmon. Data are

means ± standard error.

Fig. 4. The number of juvenile masu salmon (*Oncorhynchus masou*) recovered at each section

in the semi-natural stream. In each section of the stream, fish number was calculated as the sum

of fish remaining from each size category (75-100 and 135-160 mm). Broken line indicates the

expected surviving proportion (we hypothesized a uniform 1:1:1:1 ratio). Asterisk denotes a

significant deviation from the average value (∗∗ $p < 0.010$). Each section (Sec.) was named from

the upper most section in the stream, Sec. 1, Sec. 2, Sec. 3 and Sec. 4, Sec. 2 was the stocking

location.

Fig. 5. Number of photos of grey heron (*Ardea cinerea*) in each study section of semi-natural

stream. X-axis show the number of days elapsed from stocking (day 0) to completion at day 20.

Fig. 6. The percent of the photos that showed grey heron located around the pool or riffle

across each study section of the semi-natural stream. Asterisk denotes a significant difference (∗

$p < 0.050$, ∗∗ $p < 0.010$). Data are means ± standard error.

Table 1. Results from two-way ANOVA to determine the effect
of study days, salmon length and their interaction on the survival
of masu salmon (*Oncorhynchus masou*)

Variable	*df*	*MS*	*F*	*p*
Study days	2	0.38	10.56	**0.001**
Salmon length	1	1.31	36.67	**< 0.001**
Period × Salmon length	2	0.01	0.18	0.833
Error	18	0.04		

Table 2. Results from two-way ANOVA to determine the effect of number of salmon, salmon length and their interaction on the number of photos containing grey herons (*Ardea cinerea*)

Variable	df	MS	F	p
Number of salmon	2	1.11	149.26	**< 0.001**
Salmon length	1	0.02	2.48	0.141
Number of salmon × Salmon length	2	0.00	0.02	0.981
Error	12	0.01		

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5

Fig.6