Loss of synteny to conserved noncoding elements impact the fetus-specific

gene expression in mammalian brain

Meenakshi Bagadia[§], Keerthivasan Raanin Chandradoss[§], Yachna Jain, Harpreet Singh,

Mohan Lal, Kuljeet Singh Sandhu*

Department of Biological Sciences

Indian Institute of Science Education and Research (IISER) - Mohali

Knowledge City, Sector - 81, SAS Nagar 140306, India

[§]*Equal contribution*

To whom correspondence should be addressed

Kuljeet Singh Sandhu

Assistant Professor

Department of Biological Sciences

Indian Institute of Science Education and Research (IISER) - Mohali

E. mail: sandhuks@iisermohali.ac.in

Abstract

Conserved noncoding elements (CNEs) have significant regulatory influence on their neighbouring genes. Loss of synteny to CNEs through genomic rearrangements can, therefore, impact the transcriptional states of the cognate genes. Yet, the evolutionary implications of such chromosomal position effects have not been studied. Through genome-wide analysis of CNEs and the cognate genes of representative species from 5 different mammalian orders, we observed significant loss of synteny to CNEs in rat lineage. The CNEs and genes losing synteny had significant association with the fetal, but not the post-natal, brain development as assessed through ontology terms, developmental gene expression, chromatin marks and genetic mutations. The loss of synteny correlated with the independent evolutionary loss of fetus-specific upregulation of genes in rat brain. DNA-breakpoints implicated in brain abnormalities of germline origin had significant representation between CNE and the gene that exhibited loss of synteny, signifying the underlying developmental tolerance of genomic rearrangements that had allowed the evolutionary splits of CNEs and the cognate genes in rodent lineage. These observations highlighted the non-trivial impact of chromosomal position-effect in shaping the evolutionary dynamics of mammalian brain development and might explain loss of brain traits, like cerebral folding of cortex, in rodent lineage.

Author Summary

Expression of genes is regulated by proximally located non-coding regulatory elements. Loss of linear proximity between gene and its regulatory element thus can alter the expression of gene. Such a phenomenon can be tested at whole genome scale using evolutionary methods. We compared the positions of genes and regulatory elements in 5 different mammals and identified the significant loss of proximities between gene and their regulatory elements in rat during evolution. Brain development related function was selectively enriched among the genes and regulatory elements that had lost the proximity in rat. The observed separation of genes and their regulatory elements was strongly associated with the evolutionary loss of developmental gene expression pattern in rat brain, which coincided with the loss of brain traits in rodents. The study highlighted the importance of relative chromosomal positioning of genes and their gene regulatory elements in the evolution of phenotypes.

Keywords: conserved non-coding elements, enhancer, genome organization, synteny, evolution, position-effect.

Introduction

Around 4-8% of the human genome is evolutionary constrained, of which coding elements contribute only about 1.5%, while rest is non-coding [1-3]. Massive data produced by ENCODE and Epigenome Roadmap projects have confirmed that majority of the evolutionary constrained non-coding DNA serve as protein binding sites[4, 5]. These conserved noncoding elements (CNEs) are interwoven with the protein coding genes in a complex manner. Ample evidence converges to non-trivial regulatory impact of CNEs on proximal gene. Deletion of a non-coding region between sclerostin (SOST) gene, a negative regulator of bone formation, and MEOX1 impacts the expression of SOST and is strongly associated with Van Buchmen disease characterized by progressive overgrowth of bones [6]. Similarly, deletion of a 10kb non-coding region downstream to stature homeobox (SHOX) gene is associated with Leri Weill dyschondrosteosis syndrome, a skeletal dysplasias condition [7]. Mutations in CNEs downstream to PAX6 gene prevent its expression and are associated with Aniridia, a congenital eye malformation. Genetic errors in locus control region (LCR) at alpha and beta globin loci strongly associate with alpha/beta-thalassemia [8, 9]. Maternal deletion of Igf2/H19 ICR disrupts the Igf2 imprinting leading to bi-allelic expression of Igf2, which is strongly associated with Beckwith Weidman syndrome [10]. Loss of a CNE proximal to androgen receptor is strongly associated with evolutionary loss of penile spines and sensory vibrissae in human [11].

Around 200,000 human-anchored Conserved Non-coding Elements (CNEs) have been identified in mammals, which are likely to exhibit gene regulatory potential, as measured through enhancer-associated chromatin marks [12-14]. Most CNEs position around developmental genes [14-16]. However, establishing causal relationship between CNE and the phenotype remains a daunting task. Though genome wide association studies (GWAS) have uncovered a whole repertoire of non-coding variants with phenotypic associations [17], it is difficult to identify the causal variants. More recently, pooled CRISPR-Cas technique has been implemented to alter the non-coding elements to assess their function more precisely [18]. These methods are difficult to be scaled up for high throughput genotype-phenotype associations. With the availability of whole genome sequences of multiple species, evolutionary methods are instrumental in deciphering genotype-phenotype associations. Through comprehensive multi-species comparison, it has been inferred that most CNEs are syntenic to the nearest gene in linear proximity and are likely to regulate the same [14, 19]. Attempts have been made to link evolutionary loss and sequence divergence of CNEs to lineage specific traits, like auditory system in echo-locating mammals and adaptively morphed pectoral flippers in marine mammals [20, 21]. In this study, we asked the question whether the lineage-specific evolutionary alterations in relative chromosomal positions of CNEs are associated with lineage-specific changes in

gene expression. Through analysis of chromosomal positions of orthologous CNEs and genes from 5

different mammals, we observed that a significant number of genes had lost synteny to their

adjacent CNEs independently in rat lineage. This loss of synteny was significantly associated with the

down-regulation of genes involved in neuorogenesis and neuronal migration during fetal brain

development, which coincided with the evolutionary loss of several brain traits independently in rat

lineage. The study suggested significant contribution of chromosomal position-effect in the

evolutionary divergence of developmental gene expression trajectories in mammals.

Results

Loss of synteny between CNE and the proximal gene

Using chromosomal position data of CNEs and genes from representative primate (human), rodent

(rat), carnivore (dog), perrisodactyl (horse), and artiodactyls (cow), we obtained 51434 'syntenic'

CNE-gene pairs (4241 genes), wherein the CNE and the nearest gene-TSS were <1 Mb distance apart

in all 5 species. There were 3579 'non-syntenic' CNE-gene pairs (334 genes), wherein the CNE and

the gene-TSS were on different chromosomes or were >2Mb apart independently in one of the

species (Figure 1A, Figure S1, Methods). The rationale of 1Mb distance cutoff for the synteny was

based on the observation that the distribution of all CNE-gene distances saturated when approached

1Mb range (Figure 1B). Similar approach has been used earlier to infer enhancer-promoter linkage

based on evolutionary syntney between the two in 1Mb range[19]. The distance cut-off of 2Mb for

loss of synteny made sure that the minimal expansion in CNE-gene distance would at-least be 2 fold.

To test if the CNEs in syntenic and non-syntenic sets were comparable, we assessed their lengths

and degree of conservation in mammalian genomes. Figure 1C showed insignificant differences in

the degree of sequence conservation of syntenic and non-syntenic CNEs, suggesting that the

sequence of non-syntenic CNEs had not diverged among mammals as compared to that of syntenic

CNEs. The length distribution of syntenic and non-syntenic CNEs showed only marginal difference

towards slightly longer CNEs in non-sytenic set (Figure 1D). However, the syntenic and non-syntenic

CNEs were located in the genomic domains of distinct sequence properties. We analysed the

enrichment of SINE, LINE and LTR retrotransposons, which covers upto 50% of mammalian genome,

around syntenic and non-syntenic CNEs. Syntenic CNEs were enriched in the region of open

chromatin, as signified through greater enrichment SINE content, and might have more wide-spread

role across different cell-lineages as compared with the non-syntenic CNEs (Figure 1E). In contrast,

the non-syntneic CNEs were located in the domains enriched with long terminal repeats (LTR),

marking their susceptibility to genomic rearrangements through mechanisms like non-allelic

homologous recombination (NAHR) [22, 23] (Figure 1E). LINE elements, in general, did not exhibit

significant difference in the two sets. These observations were largely consistent across species,

marking an ancestral property (Figure 1E), except in the rat wherein the LINE elements were

enriched around non-syntenic CNEs. This exception can be explained by the fact that the rodents

retained least of the ancestral LINE elements (1%) as compared to most other mammals (7-8%) and

had accumulated new LINE elements[24].

Relatively large number of syntenic CNE-gene pairs (93.5%) confirmed the widespread conservation

of linear proximity between CNE and its adjacent gene [14]. Among the total 3579 non-syntenic

instances, 2711 (75%) were associated with the rat genome alone, coherent with the significantly

greater number of structural variations in rodent clade[25] (Figure 2A-B). Positive scaling between

number of non-syntenic instances and the break-point distances of species from the common

ancestor signified that CNE-gene synteny was an ancestral trait (Figure 2B). Due to significant loss of

synteny in rat lineage as compared to others, we focussed on rat instances in this study. By 'Loss of

synteny' or 'non-syntenic' set, we referred to loss of CNE-gene synteny in rat from figure 2C

onwards. To directly assess the proportion of non-syntenic CNE-gene pairs associated with structural

variations, we analysed the rodent-specific evolutionary break-points (Methods). We observed that

930 (34%) of all non-syntenic instances in rat had at-least one rodent-specific break-point in

between the gene-TSS and the CNE as compared to 319 (11%) on an average for the random null

prepared through distance-controlled bootstrap sampling of syntenic CNE-gene pairs (Figure 2C,

Methods). This suggested that the loss of CNE-gene synteny in rat could largely be explained through

rodent-specific genomic rearrangements (Figure 2C). We further argue that the sequence alignment

based annotations of evolutionary breakpoints might not represent the entire repertoire of genomic

rearrangements and, therefore, analysed the neighbouring genes on either side of non-syntenic

CNEs to map the various rearrangement scenario through which CNE-gene synteny was lost. We

found that that the translocation like scenario, as marked by (i) in Figure 2D, largely explained the

inter-chromosomal (*trans*) splits of CNE and the adjacent gene. The scenario, which reflected the

mapping artefacts, like in panel (iii), was under-represented (5%, Figure 2D). Analysis of intra-

chromosomal (*cis*) splits suggested inversion-like events separating the CNE-gene pairs. Scenario (iv)

and (v) showed events where region adjacent to CNE (on left side in scenario-iv and right side in

scenario-v) had undergone local rearrangements, of which 30% and 90% events respectively were

confirmed as inversion events by analysing the change in relative strand orientation of neighboring

genes. We illustrated examples of *trans* and *cis* splits of CNE and the genes in Figure 2E. Gene

POU3F2 on human chromosome 6 was syntenic to a CNE, which was 45Kb upstream. The

orthologous CNE and the gene in rat were on chromosome 8 and 5 respectively marking the *trans*

split of CNE and the gene through translocation (Figure 2E). Another CNE was 18kb upstream to

gene ADAM23 on human chromosome 2. The rat orthologues were separated by 2.4Mb on

chromosome 9 through an inversion (Figure 2E).

We concluded that the rodent-specific genomic rearrangements largely explained the loss of CNE-

gene synteny in rat.

Genes that had lost synteny to CNEs in rat were associated with the fetal brain development

Significant differences in the genomic attributes around syntenic and non-syntenic CNEs hinted at

their distinct functional roles. To assess their functions, syntenic and non-syntenic gene-lists were

subjected to Gene Ontology (GO) and Mammalian Phenotype Ontology (MPO) analyses. The analysis

of GO terms revealed enrichment of general as well as various tissue-specific development related

terms in the syntenic set, while non-syntenic set was specifically enriched with nervous system

development related terms (Figure 3A). In MPO analysis, syntenic set exhibited of enrichment of

neonatal lethality and skeletal phenotypes, while non-syntenic set was associated with brain

morphology related phenotypes (Figure S2A). Species, other than rat, did not exhibit enrichment of

any particular functional term, owing to smaller sample size.

We further followed the above observations through tissue-specific gene expression analysis in

human. The syntenic set had widespread representation of genes expressed in different cell-lineages

and, therefore, did not exhibit significant tissue-specificity, while genes in non-syntenic set were

specifically expressed in brain (Figures 3B). The brain-specific expression of genes in non-syntenic

set was also confirmed through enrichment analysis of anatomical terms from *bgee* database (Figure

S2B). Within brain, non-syntenic set was enriched with the genes specifically expressed in cerebral

cortex during fetal, but not post-natal development (Figure 3C). In contrast, the genes in syntenic set

did not exhibit any specificity for brain tissues and developmental stages (Figures 3C). These

observations highlighted fetal brain-specific roles of genes in the non-syntenic set.

Non-syntenic CNEs function as fetal brain-specific enhancers

To test whether the differences between syntenic and non-syntenic sets observed through

functional analysis of genes, were coherent with the associated CNEs, we tested the regulatory

potential of CNEs by analyzing their epigenomic properties across tissues. Through analysis of

enhancer associated chromatin state annotations from Epigenome Roadmap, ENCODE and Fantom

consortia (Methods), we observed that 74% of syntenic and 61% of non-syntenic CNEs overlapped

with the enhancer-associated regulatory sites in at-least one of the tissues or cell-types, marking the

enhancer potential of CNEs. Relatively less representation of enhancers in the non-syntenic set

might relate to their tissue or developmental stage specific functions, a hypothesis that we further

reconciled through detailed analysis of histone modification associated with enhancers, namely

Histone-3-Lysine-4-mono-methylation (H3K4me1). We chose this mark because of its strong

association with the enhancer potential and the availability of genome-wide datasets for all the cell-

lineages we were interested in. We observed that: 1) the CNEs in syntenic set exhibited consistent

H3K4me1 enrichment across several fetal and adult tissues like thymus (endodermal), muscle

(mesodermal), heart (mesodermal), intestine(endodermal) and brain (ectodermal) (Figure 4A, Figure

S3A-B); 2) H3K4me1 enrichment over CNEs in non-syntenic set was specifically higher (comparable

to that of syntenic CNEs) in fetal, but not in adult brain (Figure 4A). We further observed the

significant enrichment of binding sites of ectoderm-specific transcription factors, which were

specifically upregulated in fetal brain, in non-syntenic CNEs as compared to syntenic CNEs (Figure S4,

Methods). These observations were largely coherent with our proposal that non-syntenic CNE-gene

pairs were associated with fetal brain development.

To assess the physical enhancer-promoter association, we generated virtual 4C data by processing

available HiC datasets of fetal and adult human brains (Figure S3C). Figure 4B showed the significant

fetal-to-adult ratio of the proportion of non-syntenic CNE-gene pairs showing significant physical

interactions as compared to that of syntenic CNE-gene pairs. We illustrated the physical interactions

between CNE and the gene through examples (Figure 4C, Figure S3D). Transcription start sites of

GPR85 and FEZF2 genes, both associated with neurological phenotypes[26-28], showed significant

interaction frequency to their cognate CNEs in human fetal brain, but not in adult brain. The

H3K4me1 signals at TSSs and CNEs were also significant in fetal brain as compared to adult.

Epigenomic analyses thus suggested that the majority of the non-syntenic CNEs exhibited enhancer

associated hallmarks in fetal brain.

By mapping the trait/disease associated SNPs from Genome Wide Associated Studies (GWAS) and

the nearby SNPs (proxy) in the linkage disequilibrium based on 1000 genome data, we observed that

105 of the non-syntenic CNEs were having at-least one brain related SNP (Figure 4D). This

representation was statistically significant when compared with that of syntenic set (Figure 4D).

These observations represented genetic evidence of brain-specific roles of non-syntenic CNEs. We

highlighted the example of EPHA4 gene, which is required for radial neuron migration and is

involved in the pathways leading to lissencephaly and schizophrenia in human[29, 30]. Upstream

CNE to this gene had a schizophrenia associated SNP. Fetal brain specificity of CNE and gene
expression was illustrated using H3K4me1 and RNA-seq tracks of fetal and post-natal human brains
(Figure 4E).
240
241 Therefore, our observations through enhancer datasets, epigenomic marks, differential motif
242 enrichment analysis and brain associated SNPs concomitantly established that the non-syntenic
243 CNEs were specific to fetal brain development in human.
244

Developmental tolerance loss of synteny events
While we have shown that the genes and the CNEs that had lost synteny in rat were associated with
fetal brain development, whether or not CNE-gene proximity was causally linked with the brain-
specific expression of the cognate gene remained to be addressed. Towards this, we assessed the
representation of germ-line breakpoints associated with the congenital disorders exhibiting brain
abnormalities and the somatic cancer breakpoints, between CNE and gene-TSS in syntenic and non-
syntenic sets. Since the observed germ-line break-points are the ones that had survived through
germ-line and the embryonic development, their presence and absence between CNE and the
adjacent gene signifies developmental tolerance and intolerance respectively of loss of CNE-gene
synteny. On the contrary, the cancer breakpoints of somatic origin do not undergo such selection
and hence do not indicate the developmental tolerance or lack thereof. Figure 5A showed relative
proportion of non-syntenic CNE-gene pairs having at-least one DNA breakpoint between gene-TSS
and the CNE superimposed onto random null prepared from syntenic CNE-gene pairs of similar
distance distribution as that of non-syntenic. We observed significantly greater representation of
germ-line breakpoints in non-syntenic set as compared to syntenic set, while representation of
somatic breakpoints showed insignificant difference (Figure 5A). We interpreted that DNA
breakpoints between non-syntenic CNE and the genes were developmentally tolerable and genomic
rearrangements thereof in the ancestral genome might have served as an evolutionary substrate for
position effect. We further showed a few examples of germ-line chromosomal rearrangements that
had split the CNE and the adjacent gene in congenital disorders with brain abnormalities (Figure 5B).
Example (i) in Figure 5B showed a chromothripsis event wherein an inversion had split the CNE-gene
pair. The involved gene BCL11A regulates cortical neuron migration and mutations therein associate
with microcephaly and intellectual disability in human[31]. BCL11A gene also exhibited 3.6 fold loss
of expression in peripheral blood of the patient having genomic rearrangement as compared with
the normal mother of the patient. Example (ii) showed a translocation event splitting a CNE and
CCDC68 gene. Genetic mutations in CCDC68 are associated with schizophrenia, bipolar disorder and

autism[32]. In example (iii), an inversion had split the CNE from DNAJB6 gene, which has role in
neuritogenesis and neuroprotective functions[33].

Loss of synteny to CNEs coincided with the loss of fetus-specific upregulation of genes in rat brain
An important question was whether or not the evolutionary loss of CNE-gene synteny in rat was
associated with the loss of expression. To assess the functional fate of associated genes, we
compared their time-course gene expression trajectories for developing cerebral cortex of human,
rat and sheep (as an out-group). Sheep was inducted in the analysis due to the availability of gene
expression datasets for pre- and post-natal tissues. We found that 99.4% of CNE-gene pairs that had
lost synteny in rat were syntenic in sheep too, confirming the independent loss of synteny in rat
lineage. We observed relative loss of fetus-specific gene expression in rat brain as compared to that
of human and sheep, suggesting that the loss of synteny correlates with the loss of fetus-specific
gene expression in developing rat brain (Figure 6). Enrichment of neurogenesis related genes and
down-regulation thereof in fetal brain of rat has implications in understanding loss of brain traits in
rat lineage, as discussed in detail in the discussion section.

Taken together, our analysis suggested a strong association between evolutionary dynamics of
chromosomal positions of gene regulatory elements and the gain or loss of gene expression, aligning
to the notion of 'position effect'. Tissue and developmental stage specific impact of position effect
highlighted the possibility of its significant role in altering developmental dynamics towards
evolutionary gain or loss of lineage-specific traits.

Discussion
It is not always the change in number and the sequence of protein coding regions in the genome
that leads to the phenotype alternation in evolution, the dynamics of gene expression is equally
relevant in the context. One way the gene expression is altered is through position-effect, i.e.,
relative chromosomal position of the gene in the genome can alter its expression through regulatory
elements and chromatin states in the neighbourhood. Position effect was first discovered through
the observation that the chromosomal arrangement of duplicated copies of *bar* gene in *bar*-mutant
flies had influence on its expression and consequently causes the relative decrease in number of eye
facets [34, 35].   Similarly, *white* gene when localized near heterochromatin gives mottled eye
phenotype with red and white patches in drosophila eye [36]. Despite its significance, the role of
position effect in evolution of traits has not been investigated thoroughly. Through comparative
genomic analysis, we showed that the CNE-gene pairs that were syntenic in most mammals, but lost

the close linear proximity independently in rat were associated with the alteration in the
transcriptional program during fetal brain development, presenting evidence how the position-effect
might have impacted the evolution of lineage-specific phenotypes by modulating the developmental
trajectories in early stages.

Enhancers can function at distance longer than several Mbs and spatial synteny has been observed
among genomic regions that had been rearranged in the evolution[37, 38]. How might then the loss
of linear proximity to CNEs downregulate the expression of genes? Evidence suggests that such long-
range or trans enhancer promoter interactions occur at the cost of increased expression noise[39-
41]. As a result, the overall expression level in a tissue is expected to decline due to increased
stochastic fluctuations in gene expression across cells. Therefore, we hypothesized that the loss of
linear proximity between CNE and the gene would have compromised with the expression level of
the gene by allowing stochastic variations in enhancer-promoter interactions.

Enrichment of brain development related genes in the non-syntenic set might relate to
developmental plasticity of brain as compared to other tissues. Genomic alterations at the loci which
are important in the development of basic body plan and functioning would be embryonic lethal,
which largely explained the significant representation of skeletal/heart development and neonatal
lethality related genes in the syntenic set (Figure 3A, S2A). Brain, despite having neurodevelopment
plasticity, exhibits least genome-wide expression divergence across mammalian species[42, 43], but
within the space of small non-syntenic gene-set the expression divergence was observed. This
suggested that the least expression divergence observed for brain were due to cellular functions that
needed to be precisely regulated to maintain delicately shaped brain tissues of all the mammals in
general, while the ones that exhibited divergence would implicate in developmental functions
specific to fetal brain. Our data showed that one of the ways, such expression divergence was
modulated in the evolution was through alteration of genomic proximity between CNEs and the
neighbouring genes. Fetus brain-specific downregulation of neurogenesis related genes that had lost
synteny to CNEs in rat aligned to the hypothesis that observed genomic alterations might link to
brain traits that were lost in rodent lineage. We showed evidence that among the species taken in
this analyses, rat exhibited most number of independently modified brain traits, including the ones
directly associated with neurogenesis, like absence of cerebral folding of cortex, absence of
claustrum separation from cortex, absence of lateral geniculate nucleus magnocellular layer etc
(Figure S5). Of these, loss of cerebral folding of cortex, i.e., lissencephalic or smooth brain phenotype
is the largest visible alteration in the rodent brains. Folded or gyrencephalic brain, in general, is

considered as adaptation for the mammals with greater encephalization quotient, intelligence and

complex behavioural traits[44, 45]. It can, therefore, be contended that the CNE-gene proximity and

the associated fetal brain-specific expression was not lost in rat, but were rather gained in other

mammals that had bigger and gyrencephalic brains. However, we argue that significant non-

uniformity in cerebral cortex has been observed across several different mammalian species[46] and

the assumption that the common ancestor of placental mammals had a smaller and simpler brain

has been challenged recently[47, 48]. Evidence converges to gyrencephalic brain of eutherian

ancestor and the subsequent loss of cortical gyration in some lineages including rodents has been

supported[49]. The enrichment of genes associated with brain morphology phenotypes (Figure S2),

ECM-receptor interactions & actin cytoskeleton regulation (ACTN, ITGA1, ITGA11, ASAP1, LAMB4,

CD36 etc), and the ones implicated in human cortical malformations (MYCN, NRXN1, RASA1, DDX11,

FEZF2, EFNA5, GLI3 etc) [50]in the non-syntenic set further supported the loss of synteny in rat

rather than gain of synteny in other mammals (Figure S6). We also emphasized that the loss of

synteny in rat was inferred by filtering the CNE-gene pairs which were syntenic in all other species,

hence were evolutionarily constrained, except in rat. Assessing gain of synteny was difficult because

a CNE-gene pair that was non-syntenic in all species except one cannot be considered as

evolutionarily constrained CNE-gene pair. We suspected that gain of synteny inferred in this flawed

manner would not have shown any functional association. This indeed was observed through an

independent analysis (Figure S6).

It remains arguable whether or not the alterations in brain traits in rodent lineage represented the

adaptive selection or was a product of neutral drift. Some studies have suggested that smaller and

lissencephalic brain was adaptively selected among mammals with distinct life history traits, like

narrow habitat and smaller social groups, than that of gyrencephalic species[48]. Distinct neurogenic

potential of gyrencephalic and lissencephalic species has been attributed to the observed

difference[48]. Increased proliferative potential of basal progentior cells is necessary and sufficient

to explain the gyrencephalic brains[48].   The loss of such proliferative potential, which was likely an

ancestral trait, might have caused inefficient neurogenic program in lissencephalic species. Our

observation that the genes that had lost the synteny to CNEs in rat were involved in neuronal

differentiation and were downregulated in fetal rat brain is largely coherent with the above

proposal.

Altogether, our observations highlighted the link between genome order and the evolutionary

dynamics of temporal gene expression pattern associated with mammalian brain development. The

study also suggested that the genomic rearrangements, without any change in the genomic content,

might impact the developmental trajectories and shape the evolution of phenotypes.

Acknowledgement

Financial support to KSS from Department of Science and Technology (EMR/2015/001681) is duly

acknowledged. MB and ML were financially supported by IISER-Mohali. KRC, YJ and HS thank ICMR,

CSIR and SERB-DST agencies respectively for their fellowships.

Methods

Compilation of chromosomal position data

Human (hg19), rat (rn5), dog (camFam3), horse (equCab2) and cow (bosTau6) genome assemblies

were used in the analysis. Conserved Non-coding Elements (CNEs) were taken from Marcovitz et al

(Marcovitz et al.2016), which in turn were obtained by curating mammalian CNEs anchored to the

human genome (hg19). Our choice of the aforementioned species and the CNE dataset was

constrained by following considerations: i) We wanted sufficient evolutionary depth in the analysis

and Marcovitz el al had considered 20 sequenced mammalian genomes to identify CNEs; 2) Since our

analysis considered the chromosomal positions of CNEs and the genes, we only considered the

genomes for which complete chromosome assemblies were available. For example, chromosome

assemblies for the orders Cetacean, Chiroptera and Proboscidea etc. are not presently available; 3)

In order to obtain the sufficient number of orthologous genes across species, we restricted our

analysis to fewer mammalian lineages only. Considering multiple species would have compromised

the total number of orthologous genes to start with.

We obtained the orthologue positions of human CNEs in query species using standard approach of

mapping through LiftOver (https://genome-store.ucsc.edu/) chains at 0.95 mapping coverage[21].

Finally, we compiled 114219 CNEs that were having orthologous positions in all 5 species. We

independently obtained the table of orthologous genes across 5 mammals from Ensembl. Using CNE

and gene tables, the list of nearest genes that were within 1Mb to the CNEs was obtained for

human. The position of orthologous CNEs and genes in other mammals were assessed and CNE-gene

pairs were classified as syntenic if the distance between the two was less than 1Mb in all 5 species

and as non-syntenic if the CNE and the gene were >2Mb apart or were on different chromosomes in

one of the species and remained within 1Mb in rest of the species. If there were multiple

orthologues for the same gene, we took the nearest gene to the CNE on the same chromosome to

ensure that a syntenic pair should not have classified as non-syntenic because of mapping errors.

The distance cut-off of 1Mb was determined based on distribution of number of CNE-gene pairs at
different distance cut-offs. At around 1Mb, the overall distribution approaches a plateau and the
numbers do not increase significantly after that (Figure 1B). The lack of synteny cut-off of 2Mb
ensured that CNE and the gene were distant at-least by 2 fold in their non-syntenic form when
compared to their syntenic form. Larger distance cut-off was also likely to be robust against the
annotation artefacts of gene coordinates. A flow-chart illustrating the overall strategy is given in
Figure S1. All the data are available in supplementary data file.

To assess the genome assembly artefacts, we mapped the non-syntenic CNE-gene pairs of rn5 rat
assembly to rn6 assembly. Out of 2711 CNE-gene pairs, 2667 pairs (98.4%) were successfully lifted
over to rn6. All pairs maintained non-sytenic status in rn6 too (Figure S7). We further mapped the
non-syntenic CNE-pairs to known problematic regions of rat genome. Out of 2711 CNE, only 3 CNEs
(0.1%) and out of total 245 genes, only three genes (ABCC6, FOS, BNIP2; 1.2%) mapped to these
regions. Exclusion of these regions was unlikey to change our claims.

Analysis of genomic attributes
Chromosomal coordinates of repeat elements were downloaded from UCSC table browser. Repeat
elements were mapped +/- 50kb around syntenic and non-syntenic CNEs and average value of
enrichment in 2kb bins were plotted. For conservation analysis PhyloP scores of placental mammals
(http://ccg.vital-it.ch/mga/hg19/phylop/phylop.html) were mapped +/- 1kb to CNE.

Functional enrichment analysis
Gene Ontology and Mammalian Phenotype Ontology analysis was performed using GREAT
(http://bejerano.stanford.edu/great/public/html/). Tissue specificity analysis was done using TSEA
(http://genetics.wustl.edu/jdlab/tsea/), CSEA (http://genetics.wustl.edu/jdlab/csea-tool-2/). The
tissue specificity index (pSI) score of a gene i in a tissue k, over the given tissues $j=1,2,...m$ was
calculated as per Dougherty et al [51] using following equation:

$$pSI_{i,k} = \frac{\sum_{j=2}^{m} \left(rank(\frac{x_{i,k}}{x_{i,j}})\right)}{m-1}$$

Where $x_{i,1}$ is the expression level of gene i in tissue 1 and $x_{i,j}$ is the expression level of gene i in tissue
j. A stringent pSI cut-off of 0.05 was taken for the analysis. For syntenic gene set, we randomly
sampled 245 genes (the size of non-syntenic gene set) from 4241 syntenic genes 100 times and
plotted the mean and the standard error of the significance (-log10 of corrected p-value) of overlap
between the candidate gene-sets and the tissues specific genes in the genome. The random sample

of syntenic genes that exhibited most significant overlap with the brain-specific genes was taken for

the expression specificity analysis among brain tissues across developmental stages.

Normalized gene expression data for developing cerebral corte and heart of human, rat and sheep

were taken from BRAINSPAN (human cortex; http://www.brainspan.org/static/download.html),

GSE71148(human heart), Stead et al (rat cortex), GSE53512 (rat heart), Clark et al (sheep cortex) and

GSE66725 (sheep heart). Average gene expression with 90% confidence intervals were plotted across

development time course.

Enhancer analysis

Regulatory potential of CNEs was assessed by mapping ChromHMM data obtained from Epigenome

roadmap   (http://egg2.wustl.edu/roadmap/web_portal/imputed.html#chr_imp)   and   ENCODE

(http://hgdownload.soe.ucsc.edu/goldenPath/hg19/encodeDCC/wgEncodeBroadHmm/)   projects

onto CNEs. Enhancer coordinates from FANTOM (http://enhancer.binf.ku.dk/presets/) were also

mapped to CNEs. Cumulative overlap across aforementioned three resources was calculated.

Datasets for H3K4me1 methylation for fetal and post-natal/adult human tissues were obtained from

Epigenome Roadmap (http://www.roadmapepigenomics.org/data/) with following accession IDs and

age groups: fetal brain (E081, E082; 17GW), adult brain (E067, E068, E069, E071, E072, E073, E074;

pooled 73Yr/75Yr/81Yr), fetal muscle (E089, E090; 15GW), post-natal muscle (E107; pooled

54Yr/72Yr) and fetal thymus (E093; 15GW), post-natal thymus (E112; 3Yr), fetal heart (E083, 91

days), post-natal heart (E95, E104, E105, pooled 3Yr/34Yr), fetal small intestine(E085, 15GW) and

post-natal small intestine(E109, pooled 3Yr/30Yr). Fold-change over input DNA was used for

aggregation plots. WashU epigenome browser was used for visualization. Motif analysis was

performed through RSAT's 'peak-motif' package (http://rsat.sb-roscoff.fr/peak-motifs_form.cgi)

using JASPAR core matrices for vertebrate genomes. Syntenic CNEs were taken as background

control sequences.

Mapping of proxy GWAS SNPs

Total 251835 GWAS SNPs were obtained from GWASdb (http://jjwanglab.org/gwasdb). From this

data, 71990 brain related SNPs were obtained by analyzing the HPO terms associated with brain

associated phenotypes. We extended this data to total 533388 nearby SNPs (proxy) that were in

linkage disequilibrium to 71990 brain related GWAS SNPs based on 1000 genome data using SNAP

algorithm (https://personal.broadinstitute.org/plin/snap/index.php). Random null was prepared by

picking CNE samples, of same sample size and CNE-gene distance as of non-syntenic set, from the

syntenic set 1000 times and mapping SNPs to each of these samples. Number of CNEs with at-least

one SNP was counted for each sample. The distribution of these numbers was regarded as random

null. P-value was calculated as following:

$$—$$

476   Where B = number of re-sampling iterations (1000)

477   k_b = Number of sampled syntenic CNEs having at-least one SNP.

478   k = Number of observed non-syntenic CNE having atleast one SNP.

Virtual 4C data

SRA files of HiC datasets for fetal and adult brains were obtained from GSE77565 and GSE87112.

Datasets were processed using HiCUP (https://github.com/theaidenlab/juicer/wiki/HiCCUPS) and

contact maps were normalized using iterative correction and eigen vector decomposition (ICE)

method (https://github.com/hiclib/iced). TSS in each CNE-gene pair was taken as bait (reference

point) and its intra-chromosomal interactions were obtained from HiC matrices. Loess regression

line was fit to the HiC counts as a function of genomic distance from the bait. Signifcant interactions

with the bait were identified by applying cut-off of 3-standard deviation distance from the regression

line [52].

DNA breakpoint analysis

We obtained 552 germline breakpoints associated with congenital disorders having brain

abnormalities and 68018 somatic cancer breakpoints from van Heesch et al [53]. The matching RNA-

seq data of perpheral blood of patient and the mother were obtained from European Nucleotide

Archive(https://www.ebi.ac.uk/ena) with the accession IDs ERX358048 and ERX358046 respectively.

Total 2061 evolutionary DNA break-points for rodents were taken from Bourque etal (2004 & 2006),

Larkin etal and Lemaitre etal et al [54-57]. These breakpoints were then mapped onto inter-spacer

regions between CNE and the nearest gene-TSS. The random null was obtained by picking CNE-gene

pairs, of same sample size and CNE-gene distance as of non-syntenic set, from the syntenic set 1000

times and mapping the breakpoint in the inter-spacer regions. Number of CNE-gene pairs with at-

least one break-point in between was counted for each sample. The distribution of these numbers

was regarded as random null. P-values was calculated using equation as in GWAS SNP analysis. The

break-point distances from the ancestor were obtained from Luo et al[25].

Mammalian traits

Status of traits in 5 mammals were obtained from project ID P773 of Morphobank database (https://morphobank.org/). Traits that exhibited same status atleast in 3 of the mammals including rat, but showed a different status in human were classified as independently modified traits in human. Similarly the traits that had same status in atleast 3 species including human, but changed status in rat were denoted as independently modified traits in rat.

Availability of data. All datasets presented in this article are available as supplementary data:

References

1. Rands CM, Meader S, Ponting CP, Lunter G: **8.2% of the Human genome is constrained: variation in rates of turnover across functional element classes in the human lineage.** *PLoS Genet* 2014, **10**:e1004525.
2. Lindblad-Toh K, Garber M, Zuk O, Lin MF, Parker BJ, Washietl S, Kheradpour P, Ernst J, Jordan G, Mauceli E, et al: **A high-resolution map of human evolutionary constraint using 29 mammals.** *Nature* 2011, **478**:476-482.
3. Thurman RE, Rynes E, Humbert R, Vierstra J, Maurano MT, Haugen E, Sheffield NC, Stergachis AB, Wang H, Vernot B, et al: **The accessible chromatin landscape of the human genome.** *Nature* 2012, **489**:75-82.
4. Consortium EP: **An integrated encyclopedia of DNA elements in the human genome.** *Nature* 2012, **489**:57-74.
5. Skipper M, Eccleston A, Gray N, Heemels T, Le Bot N, Marte B, Weiss U: **Presenting the epigenome roadmap.** *Nature* 2015, **518**:313.
6. Loots GG, Kneissel M, Keller H, Baptist M, Chang J, Collette NM, Ovcharenko D, Plajzer-Frick I, Rubin EM: **Genomic deletion of a long-range bone enhancer misregulates sclerostin in Van Buchem disease.** *Genome Res* 2005, **15**:928-935.
7. Sabherwal N, Bangs F, Roth R, Weiss B, Jantz K, Tiecke E, Hinkel GK, Spaich C, Hauffa BP, van der Kamp H, et al: **Long-range conserved non-coding SHOX sequences regulate expression in developing chicken limb and are associated with short stature phenotypes in human patients.** *Hum Mol Genet* 2007, **16**:210-222.
8. Driscoll MC, Dobkin CS, Alter BP: **Gamma delta beta-thalassemia due to a de novo mutation deleting the 5' beta-globin gene activation-region hypersensitive sites.** *Proc Natl Acad Sci U S A* 1989, **86**:7470-7474.
9. Hatton CS, Wilkie AO, Drysdale HC, Wood WG, Vickers MA, Sharpe J, Ayyub H, Pretorius IM, Buckle VJ, Higgs DR: **Alpha-thalassemia caused by a large (62 kb) deletion upstream of the human alpha globin gene cluster.** *Blood* 1990, **76**:221-227.
10. Sparago A, Cerrato F, Vernucci M, Ferrero GB, Silengo MC, Riccio A: **Microdeletions in the human H19 DMR result in loss of IGF2 imprinting and Beckwith-Wiedemann syndrome.** *Nat Genet* 2004, **36**:958-960.
11. McLean CY, Reno PL, Pollen AA, Bassan AI, Capellini TD, Guenther C, Indjeian VB, Lim X, Menke DB, Schaar BT, et al: **Human-specific loss of regulatory DNA and the evolution of human-specific traits.** *Nature* 2011, **471**:216-219.
12. Seridi L, Ryu T, Ravasi T: **Dynamic epigenetic control of highly conserved noncoding elements.** *PLoS One* 2014, **9**:e109326.
13. Roh TY, Wei G, Farrell CM, Zhao K: **Genome-wide prediction of conserved and nonconserved enhancers by histone acetylation patterns.** *Genome Res* 2007, **17**:74-81.
14. Babarinde IA, Saitou N: **Genomic Locations of Conserved Noncoding Sequences and Their Proximal Protein-Coding Genes in Mammalian Expression Dynamics.** *Mol Biol Evol* 2016, **33**:1807-1817.

553   15.   Woolfe A, Goodson M, Goode DK, Snell P, McEwen GK, Vavouri T, Smith SF, North P,
554   Callaway H, Kelly K, et al: **Highly conserved non-coding sequences are associated with**
555   **vertebrate development.** *PLoS Biol* 2005, **3**:e7.
556   16.   Akalin A, Fredman D, Arner E, Dong X, Bryne JC, Suzuki H, Daub CO, Hayashizaki Y, Lenhard B:
557   **Transcriptional features of genomic regulatory blocks.** *Genome Biol* 2009, **10**:R38.
558   17.   Welter D, MacArthur J, Morales J, Burdett T, Hall P, Junkins H, Klemm A, Flicek P, Manolio T,
559   Hindorff L, Parkinson H: **The NHGRI GWAS Catalog, a curated resource of SNP-trait**
560   **associations.** *Nucleic Acids Res* 2014, **42**:D1001-1006.
561   18.   Rajagopal N, Srinivasan S, Kooshesh K, Guo Y, Edwards MD, Banerjee B, Syed T, Emons BJ,
562   Gifford DK, Sherwood RI: **High-throughput mapping of regulatory DNA.** *Nat Biotechnol*
563   2016, **34**:167-174.
564   19.   Naville M, Ishibashi M, Ferg M, Bengani H, Rinkwitz S, Krecsmarik M, Hawkins TA, Wilson
565   SW, Manning E, Chilamakuri CS, et al: **Long-range evolutionary constraints reveal cis-**
566   **regulatory interactions on the human X chromosome.** *Nat Commun* 2015, **6**:6904.
567   20.   Davies KT, Tsagkogeorga G, Rossiter SJ: **Divergent evolutionary rates in vertebrate and**
568   **mammalian specific conserved non-coding elements (CNEs) in echolocating mammals.**
569   *BMC Evol Biol* 2014, **14**:261.
570   21.   Marcovitz A, Jia R, Bejerano G: **"Reverse Genomics" Predicts Function of Human Conserved**
571   **Noncoding Elements.** *Mol Biol Evol* 2016, **33**:1358-1369.
572   22.   Chan JE, Kolodner RD: **A genetic and structural study of genome rearrangements mediated**
573   **by high copy repeat Ty1 elements.** *PLoS Genet* 2011, **7**:e1002089.
574   23.   Roeder GS, Fink GR: **DNA rearrangements associated with a transposable element in yeast.**
575   *Cell* 1980, **21**:239-249.
576   24.   Buckley RM, Kortschak RD, Raison JM, Adelson DL: **Similar Evolutionary Trajectories for**
577   **Retrotransposon Accumulation in Mammals.** *Genome Biol Evol* 2017, **9**:2336-2353.
578   25.   Luo H, Arndt W, Zhang Y, Shi G, Alekseyev MA, Tang J, Hughes AL, Friedman R: **Phylogenetic**
579   **analysis of genome rearrangements among five mammalian orders.** *Mol Phylogenet Evol*
580   2012, **65**:871-882.
581   26.   Matsumoto M, Straub RE, Marenco S, Nicodemus KK, Matsumoto S, Fujikawa A, Miyoshi S,
582   Shobo M, Takahashi S, Yarimizu J, et al: **The evolutionarily conserved G protein-coupled**
583   **receptor SREB2/GPR85 influences brain size, behavior, and vulnerability to schizophrenia.**
584   *Proc Natl Acad Sci U S A* 2008, **105**:6133-6138.
585   27.   Chen L, Zheng J, Yang N, Li H, Guo S: **Genomic selection identifies vertebrate transcription**
586   **factor Fezf2 binding sites and target genes.** *J Biol Chem* 2011, **286**:18641-18649.
587   28.   Eckler MJ, Larkin KA, McKenna WL, Katzman S, Guo C, Roque R, Visel A, Rubenstein JL, Chen
588   B: **Multiple conserved regulatory domains promote Fezf2 expression in the developing**
589   **cerebral cortex.** *Neural Dev* 2014, **9**:6.
590   29.   Steinecke A, Gampe C, Zimmer G, Rudolph J, Bolz J: **EphA/ephrin A reverse signaling**
591   **promotes the migration of cortical interneurons from the medial ganglionic eminence.**
592   *Development* 2014, **141**:460-471.
593   30.   Senturk A, Pfennig S, Weiss A, Burk K, Acker-Palmer A: **Ephrin Bs are essential components**
594   **of the Reelin pathway to regulate neuronal migration.** *Nature* 2011, **472**:356-360.
595   31.   Wiegreffe C, Simon R, Peschkes K, Kling C, Strehle M, Cheng J, Srivatsa S, Liu P, Jenkins NA,
596   Copeland NG, et al: **Bcl11a (Ctip1) Controls Migration of Cortical Projection Neurons**
597   **through Regulation of Sema3c.** *Neuron* 2015, **87**:311-325.
598   32.   Cross-Disorder Group of the Psychiatric Genomics C: **Identification of risk loci with shared**
599   **effects on five major psychiatric disorders: a genome-wide analysis.** *Lancet* 2013,
600   **381**:1371-1379.
601   33.   Calabrese V, Scapagnini G, Ravagna A, Giuffrida Stella AM, Butterfield DA: **Molecular**
602   **chaperones and their roles in neural cell differentiation.** *Dev Neurosci* 2002, **24**:1-13.

34.   Sturtevant AH: **A Further Study of the so-Called Mutation at the Bar Locus of Drosophila.**
Genetics 1928, **13**:401-409.

35.   Sturtevant AH: **The Effects of Unequal Crossing over at the Bar Locus in Drosophila.**
Genetics 1925, **10**:117-147.

36.   Martin-Morris LE, Csink AK, Dorer DR, Talbert PB, Henikoff S: **Heterochromatic trans-**
inactivation of Drosophila white transgenes. *Genetics* 1997, **147**:671-677.

37.   Veron AS, Lemaitre C, Gautier C, Lacroix V, Sagot MF: **Close 3D proximity of evolutionary**
breakpoints argues for the notion of spatial synteny. *BMC Genomics* 2011, **12**:303.

38.   Bagadia M, Singh A, Singh Sandhu K: **Three Dimensional Organization of Genome Might**
Have Guided the Dynamics of Gene Order Evolution in Eukaryotes. *Genome Biol Evol* 2016,
8:946-954.

39.   Singh A, Bagadia M, Sandhu KS: **Spatially coordinated replication and minimization of**
expression noise constrain three-dimensional organization of yeast genome. *DNA Res*
2016, **23**:155-169.

40.   Sandhu KS: **Did the modulation of expression noise shape the evolution of three**
dimensional genome organizations in eukaryotes? *Nucleus* 2012, **3**:286-289.

41.   Kustatscher G, Grabowski P, Rappsilber J: **Pervasive coexpression of spatially proximal**
genes is buffered at the protein level. *Mol Syst Biol* 2017, **13**:937.

42.   Khaitovich P, Enard W, Lachmann M, Paabo S: **Evolution of primate gene expression.** *Nat*
Rev Genet 2006, **7**:693-702.

43.   Strand AD, Aragaki AK, Baquet ZC, Hodges A, Cunningham P, Holmans P, Jones KR, Jones L,
Kooperberg C, Olson JM: **Conservation of regional gene expression in mouse and human**
brain. *PLoS Genet* 2007, **3**:e59.

44.   Prothero JW, Sundsten JW: **Folding of the cerebral cortex in mammals. A scaling model.**
Brain Behav Evol 1984, **24**:152-167.

45.   Toro R, Perron M, Pike B, Richer L, Veillette S, Pausova Z, Paus T: **Brain size and folding of**
the human cerebral cortex. *Cereb Cortex* 2008, **18**:2352-2357.

46.   Herculano-Houzel S, Collins CE, Wong P, Kaas JH, Lent R: **The basic nonuniformity of the**
cerebral cortex. *Proc Natl Acad Sci U S A* 2008, **105**:12593-12598.

47.   Rowe TB, Macrini TE, Luo ZX: **Fossil evidence on origin of the mammalian brain.** *Science*
2011, **332**:955-957.

48.   Lewitus E, Kelava I, Kalinka AT, Tomancak P, Huttner WB: **An adaptive threshold in**
mammalian neocortical evolution. *PLoS Biol* 2014, **12**:e1002000.

49.   Kelava I, Lewitus E, Huttner WB: **The secondary loss of gyrencephaly as an example of**
evolutionary phenotypical reversal. *Frontiers in Neuroanatomy* 2013, **7**:16.

50.   Pinero J, Bravo A, Queralt-Rosinach N, Gutierrez-Sacristan A, Deu-Pons J, Centeno E, Garcia-
Garcia J, Sanz F, Furlong LI: **DisGeNET: a comprehensive platform integrating information**
on human disease-associated genes and variants. *Nucleic Acids Res* 2017, **45**:D833-D839.

51.   Dougherty JD, Schmidt EF, Nakajima M, Heintz N: **Analytical approaches to RNA profiling**
data for the identification of genes enriched in specific cells. *Nucleic Acids Res* 2010,
38:4218-4230.

52.   Klein FA, Pakozdi T, Anders S, Ghavi-Helm Y, Furlong EE, Huber W: **FourCSeq: analysis of 4C**
sequencing data. *Bioinformatics* 2015, **31**:3085-3091.

53.   van Heesch S, Simonis M, van Roosmalen MJ, Pillalamarri V, Brand H, Kuijk EW, de Luca KL,
Lansu N, Braat AK, Menelaou A, et al: **Genomic and functional overlap between somatic**
and germline chromosomal rearrangements. *Cell Rep* 2014, **9**:2001-2010.

54.   Larkin DM, Pape G, Donthu R, Auvil L, Welge M, Lewin HA: **Breakpoint regions and**
homologous synteny blocks in chromosomes have different evolutionary histories.
Genome Res 2009, **19**:770-777.

55.   Bourque G, Zdobnov EM, Bork P, Pevzner PA, Tesler G: **Comparative architectures of**
mammalian and chicken genomes reveal highly variable rates of genomic rearrangements
across different lineages. *Genome Res* 2005, **15**:98-110.
56.   Bourque G, Pevzner PA, Tesler G: **Reconstructing the genomic architecture of ancestral**
mammals: lessons from human, mouse, and rat genomes. *Genome Res* 2004, **14**:507-516.
57.   Lemaitre C, Zaghloul L, Sagot MF, Gautier C, Arneodo A, Tannier E, Audit B: **Analysis of fine-**
scale mammalian evolutionary breakpoints provides new insight into their relation to
genome organisation. *BMC Genomics* 2009, **10**:335.

Figure legends

Figure 1. Synteny and lack thereof between CNE and the nearest gene. (a) Illustration of the strategy to infer

the synteny and lack thereof between CNE and the neighboring gene across 5 representative mammalian

orders. CNE-gene pairs were classified as syntenic if remained proximal (<1Mb) in all the 5 species and as non-

syntenic if departed by > 2mb or on different chromosomes in on of the species while maintaining synteny in

other 4 species. (b) Cumulative distribution of all CNE-gene distances in the human genome. Most CNE-gene

pairs were <1Mb distance and, therefore, cut-off of 1Mb was applied for CNE-gene synteny. (c) Sequence

conservation, as measured through mammalian PhyloP scores, and (d) length distribution of CNEs in syntenic

and non-syntenic sets. (e) Enrichment of retrotransposons +/- 50Kb around syntenic and non-syntenic CNEs.

Asterisk indicate significant p-values (<0.05) calculated using Mann Whitney U test of enrichment values +/-

10kb around CNEs.

Figure 2. Genomic rearrangements underlying the loss of synteny (a) Number of non-syntenic instances and

genes (in parentheses) in different mammals. P-value for the non-syntenic instances in rat and the next highest

value (in dog) was calculated using Fisher's exact test. (b) Scaling between number of non-syntenic cases and

the evolutionary break-point distance from the common mammalian ancestor. (c) Number of non-syntenic

CNE-gene pairs having atleast one rodent-specific break-point inbetween CNE and gene-TSS, overlaid onto null

distribution prepared from syntenic set. (d) Distinct *trans* and *cis* chromosomal rearrangements as inferred

from the analysis of genes flanking the non-syntenic CNEs. Shown are the neighboring genes around CNE. Red

color represents the target gene and orange color represents the nearest gene on the other side of the CNE.

Red color in the barplot marks the proportion for which inversion could be confirmed through analysis of gene

orientations. Shown are the two examples illustrating loss of CNE-gene synteny in rat. In first example, a CNE

was located 45kb upstream to the gene POU3F2 in human, but were split on different chromosomes in rat.

Second example shows that an inversion event had distanced the CNE and the proximal ADAM23 gene upto

2.4Mb in rat genome.

Figure 3. Functional characterization of genes in syntenic and non-syntenic sets. (a) Enrichment of Gene

Ontology (GO) terms among genes in syntenic and non-syntenic sets. P-values shown were corrected using

Benjamini-Hochberg method. (b) Tissue-specific expression analysis of genes in syntenic and non-syntenic set.

Relative significance was plotted as negative of log10 transformed corrected p-values of Fisher's exact test for

the overlap with the tissue-specific genes at stringency score (pSI) < 0.05. Horizontal grey colored dashed line

represents the p-value of 0.01. For syntenic set, mean values and standard errors of significance for 100

random samples of 245 genes (the size of non-syntenic set) from syntenic sets were plotted. (c) Expression

specificity of genes in syntenic and non-syntenic sets across brain regions and across developmental stages.

For syntenic set, the sample that exhibited maximum significance for brain specificity in panel-b was taken.

Size of the nested hexagons represents the proportion of all genes specifically expressed in particular tissue at

particular developmental stage. Hexagons are nested inwards based on relative stringency of tissue specificity

scores (pSI=0.05, 0.01, 0.001 & 0.0001 respectively). Color gradient represents the magnitude of corrected p-

values of Fisher's exact test.

Figure 4. Enhancer properties of syntenic and non-syntenic CNEs. (a) H3K4me1 ChIP enrichment (over input)

on and around CNEs in syntenic and non-syntenic sets in fetal and post-natal tissues. P-values for the

difference between syntenic and non-syntenic CNEs were calculated using Mann Whitney U test of H3K4me1

enrichment values in 1kb spanning windows on either side of the CNEs. (b) Virtual 4C analysis of CNE-gene

interactions in fetal and adult human brains. The barplot shows fetal-to-adult ratio of proportion of CNE-gene

pairs exhibiting significant physical chromatin interactions. (c) The examples of GPR85 and FEZF2 genes are

shown for illustration. Vertical grey and yellow bars represent the TSS and CNE positions respectively. Red and

grey curves show virtual 4C and h3K4me1 signals respectively. Black smooth line represents the loess fit of the

4C signal as function of genomic distance from the reference point. The dotted line represents 3σ distance

from the loess regression line. (d) Proportion of non-syntenic CNEs having at-least one brain associated SNP

superimposed onto the null distribution obtained from syntenic CNEs. P-value was calculated using boot-strap

method by randomly sampling 2711 CNEs (size of non-syntenic set) from syntenic set 1000 times. (e) An

example of EPHA4 gene and its proximal CNE having a Schizophrenia associated GWAS SNP is shown. The

tracks for PhyloP conservation score, H3K4me1 and RNA-seq data of fetal and post-natal brain aligned

accordingly. The orthologous CNE and gene were 7.2 Mb apart on chromosome 9 in rat.

Figure 5. Tolerance and intolerance of CNE-gene split. **(a)** Number of non-syntenic CNE-gene pairs flanking at-

least one germ-line breakpoint associated with the congenital disorders having brain abnormalities (upper

panel) and somatic cancer breakpoint (lower panel), superimposed onto null distributions obtained from the

syntenic set of same CNE-gene distance distribution as that of non-syntenic set. P-values were calculating

using boot-strap method with 1000 random samplings. (b) Examples illustrating chromothripsis, translocation

and inversion events breaking CNE-gene synteny in congenital disorders with brain abnormalities. Mutations in

BCL11A (i), CCDC68 (ii) and DNAJB6 (iii) genes are associated with abnormal brain phenotypes as elaborated in

the results section. The right most panel shows difference in expression level of BCL11A gene in the patient

having genomic rearrangement and the normal mother.

Figure 6. Evolutionary dynamics of developmental gene expression associated with the loss of CNE-gene
synteny. Red curves in the plots represent the mean expression of genes in the non-syntenic set and grey area
represents 95% confidence interval. Grey colored elongated triangles below the x-axis represent the fetal and
post-natal time course. Lack of triangles at some places denotes unavailability of multiple time points in the
data. Left panel represents cerebral cortex and right panel is for heart datasets (control).

Supplementary Figure Legends

Figure S1. Flowchart illustrating overall strategy to obtain syntenic and non-syntenic CNE-gene pairs.

Figure S2. Enrichment of (a) Mammalian Phenotype Ontology (MPO) terms , and (b) Bgee anatomical terms in
syntenic (left panel) and non-syntenic (right panel) gene-sets. P-values were corrected using Benjamini-
Hochberg method.

Figure S3. Extension of Epigenomic analyses of syntenic and non-syntenic CNEs. (A-B) H3K4me1 ChIP
enrichment (over inut DNA) around syntenic (grey) and non-syntenic (red) CNEs in (a) fetal, and (b) post-natal
tissues. (C) Strategy used to assess the CNE-TSS interactions using HiC datasets of human fetal and adult
brains. All-to-all HiC interactions were filtered for TSS-to-all interactions for the genes in syntenic and
nonsynteic sets. The resultant data was analogous to 4C and was analyzed using method atypical for 4C
analysis. Loess regression line was fit to 4C counts as a function of genomic distance from the refrence point
(TSS in this case). The distance of 3-standard deviation from this regression line was taken as signicance cut-off
for the interactions impinging onto TSS. (D) Example of CNE-TSS interactions identified in the non-syntenic set.
Upper and lower panels represent fetal and adult brain data. Red line: 4C signal; grey line: H3K4me1; Black
line: Loess fit; Dotted line: 3-standard deviation cut-off for signicance.

Figure S4. Sequence motif enrichment analysis for non-synteic CNEs. 'Peak-motifs' from RSAT was used to
identify over-represented sequence motifs in the non-synteic CNEs while taking syntenic CNEs as background
control. (a) Sequence motifs, their e-values and the matching transcription factors (TFs) from JASPAR. (b)
Tissue specificity analyses of TFs. Red bars represent the tissues, wherein the TFs exhibit significant specificity
(P<0.05). (c) Time course gene expression of TFs during human brain development. Red curve represents
average expression of TFs and grey colour denotes 90% confidence interval.

Figure S5. Independently modified traits in rat and human. (A) bar plot representing rat-to-human ratio for the
proportion of traits independently modified in each trait class. (B) List of traits that were independetly
modified in rat.

Figure S6. Analysis of gain of synteny instances in human. Gain of synteny instances were identified using the
strategy shown. No enrichment was found in the GO process terms. The most enriched GO term is shown.

Figure S7. Comparison of non-syntenic (rat) CNE-gene pairs in rn5 and rn6 genome assemblies. Shown is the

scatter plot of CNE-gene distances (log10 scale) the non-syntenic set in rn5 and rn6 assemblies. All the non-

syntenic CNE-gene pairs in rn5 remained non-syntenic in rn6.