TITLE

Full title: Whole genome sequencing of pharmacogenetic drug response in racially

and ethnically diverse children with asthma

Short title: Pharmacogenetic drug response in ethnically diverse children with

asthma

AUTHORS

Angel C.Y. Mak[1,¶,*], Marquitta J. White[1,¶], Zachary A. Szpiech[2,¶], Walter L. Eckalbar[1,¶],

Sam S. Oh[1], Maria Pino-Yanes[3,4], Donglei Hu[1], Pagé Goddard[1], Scott Huntsman[1],

Joshua Galanter[1], Dara G. Torgerson[1], Ann Chen Wu[5,6], Blanca E. Himes[7], Soren

Germer[8], Julia M. Vogel[8], Karen L. Bunting[8], Celeste Eng[1], Sandra Salazar[1], Kevin L.

Keys[1], Jennifer Liberto[1], Thomas J. Nuckton[1], Thomas A. Nguyen[1], Pui-Yan Kwok[9,10],

Albert M. Levin[11], Juan C. Celedón[12], Erick Forno[12], Hakon Hakonarson[13,14], Patrick

M. Sleiman[13,14], Amber Dahlin[5], Kelan G. Tantisira[5], Scott T. Weiss[5], Denise

Serebrisky[15], Emerita Brigino-Buenaventura[16], Harold J. Farber[17], Kelley Meade[18],

Michael A. Lenoir[19], Pedro C. Avila[20,#a], Saunak Sen[1,#b] Shannon M. Thyne[21], William

Rodriguez-Cintron[22], Cheryl A. Winkler[23], Andrés Moreno-Estrada[24], Karla

Sandoval[24], Jose R. Rodriguez-Santana[25], Rajesh Kumar[26,27], L. Keoki Williams[28,29],

Nadav Ahituv[1], Elad Ziv[1], Max A. Seibold[30], Robert B. Darnell[8,31,32], Noah Zaitlen[1],

Ryan D. Hernandez[2,10,33], Esteban G. Burchard[1,2] on behalf of the Trans-Omics for

Precision Medicine Whole Genome Sequencing Program (TOPMed)^

AFFILIATIONS

[1] Department of Medicine, University of California San Francisco, San Francisco,

California, USA.

[2] Department of Bioengineering and Therapeutic Sciences, University of California

San Francisco, San Francisco, California, USA.

[3] Research Unit, Hospital Universitario N.S. de Candelaria, Universidad de La Laguna,

Santa Cruz de Tenerife, Spain.

[4] CIBER de Enfermedades Respiratorias, Instituto de Salud Carlos III, Madrid, Spain.

[5] Channing Division of Network Medicine, Department of Medicine, Brigham and

Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA.

[6] Precision Medicine Translational Research (PRoMoTeR) Center, Department of

Population Medicine, Harvard Medical School and Pilgrim Health Care Institute,

Boston, Massachusetts, USA.

[7] Department of Biostatistics, Epidemiology and Informatics, University of

Pennsylvania, Philadelphia, Pennsylvania, USA.

[8] New York Genome Center, New York, New York, USA.

[9] Cardiovascular Research Institute, University of California San Francisco, San

Francisco, California, USA.

[10] Institute for Human Genetics, University of California San Francisco, San

Francisco, California, USA.

[11] Department of Public Health Sciences, Henry Ford Health System, Detroit

Michigan, USA.

[12] Division of Pediatric Pulmonary Medicine, Allergy and Immunology, University of

Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA.

[13] Center for Applied Genomics, The Children's Hospital of Philadelphia Research

Institute, Philadelphia, Pennsylvania, USA.

[14] Department of Pediatrics, Perelman School of Medicine, University of

Pennsylvania, Philadelphia, Pennsylvania, USA.

[15] Pediatric Pulmonary Division, Jacobi Medical Center, Bronx, New York, USA.

[16] Department of Allergy and Immunology, Kaiser Permanente Vallejo Medical

Center, Vallejo, California, USA.

[17] Department of Pediatrics, Baylor College of Medicine and Texas Children's

Hospital, Houston, Texas, USA.

[18] Children's Hospital and Research Center, Oakland, California, USA.

[19] Bay Area Pediatrics, Oakland, California, USA.

[20] Department of Medicine, Northwestern University, Chicago, Illinois, USA.

[21] Department of Pediatrics, David Geffen School of Medicine, University of

California, Los Angeles, Los Angeles, California, USA.

[22] Veterans Caribbean Health Care System, San Juan, Puerto Rico.

[23] Basic Science Laboratory, Center for Cancer Research, National Cancer Institute,

Leidos Biomedical Research, Frederick National Laboratory, Frederick, Maryland,

USA.

[24] National Laboratory of Genomics for Biodiversity (UGA-LANGEBIO), CINVESTAV,

Irapuato, Guanajuato, Mexico.

[25] Centro de Neumologia Pediatrica, San Juan, Puerto Rico.

[26] Feinberg School of Medicine's Division of Allergy and Immunology, Northwestern

University, Chicago, Illinois, USA.

[27] Ann & Robert H. Lurie Children's Hospital of Chicago, Chicago, Illinois, USA.

[28] Department of Internal Medicine, Henry Ford Health System, Detroit, Michigan,

USA.

[29] Center for Health Policy and Health Services Research, Henry Ford Health System,

Detroit, Michigan, USA.

[30] Center for Genes, Environment and Health, Department of Pediatrics, National

Jewish Health, Denver, Colorado, USA.

[31] Laboratory of Molecular Neuro-Oncology, The Rockefeller University, New York,

New York, USA.

[32] Howard Hughes Medical Institute, The Rockefeller University, New York, New

York, USA.

[33] Quantitative Biosciences Institute, University of California San Francisco, San

Francisco, California, USA.

[#a]Current address: Allergy & ENT Associates. The Woodlands, Texas, USA.

[#b]Current address: Division of Biostatistics, Department of Preventive Medicine,

University of Tennessee Health Science Center, Memphis, Tennessee, USA.

[¶] These authors contributed equally to this work

* Correspondence and material requests should be addressed to Angel C.Y. Mak

(angel.mak@ucsf.edu)

ˆ Membership of the TOPMed team is provided in the Acknowledgments.

ABSTRACT

Asthma is the most common chronic disease of children, with significant

racial/ethnic differences in prevalence, morbidity, mortality and therapeutic

response. Albuterol, a bronchodilator medication, is the first-line therapy for asthma

treatment worldwide. We performed the largest whole genome sequencing (WGS)

pharmacogenetics study to date using data from 1,441 minority children with

asthma who had extremely high or low bronchodilator drug response (BDR). We

identified population-specific and shared pharmacogenetic variants associated with

BDR, including genome-wide significant ($p < 3.53$ x 10^{-7}) and suggestive ($p < 7.06$ x

10^{-6}) loci near genes previously associated with lung capacity (*DNAH5*), immunity

(*NFKB1* and *PLCB1*), and β-adrenergic signaling pathways (*ADAMTS3* and *COX18*).

Functional analyses centered on *NFKB1* revealed potential regulatory function of

our BDR-associated SNPs in bronchial smooth muscle cells. Specifically, these

variants are in linkage disequilibrium with SNPs in a functionally active enhancer,

and are also expression quantitative trait loci (eQTL) for a neighboring gene,

SLC39A8. Given the lack of other asthma study populations with WGS data on

minority children, replication of our rare variant associations is infeasible. We

attempted to replicate our common variant findings in five independent studies

with GWAS data. The age-specific associations previously found in asthma and

asthma-related traits suggest that the over-representation of adults in our

replication populations may have contributed to our lack of statistical replication,

despite the functional relevance of the *NFKB1* variants demonstrated by our

functional assays. Our study expands the understanding of pharmacogenetic

analyses in racially/ethnically diverse populations and advances the foundation for

precision medicine in at-risk and understudied minority populations.

AUTHOR SUMMARY

Asthma is the most common chronic disease among children. Albuterol, a bronchodilator medication, is the first-line therapy for asthma treatment worldwide. In the U.S., asthma prevalence is the highest among Puerto Ricans, intermediate among African Americans and lowest in Whites and Mexicans. Asthma disparities extend to mortality, which is four- to five-fold higher in Puerto Ricans and African Americans compared to Mexicans [1]. Puerto Ricans and African Americans, the populations with the highest asthma prevalence and death rate, also have the lowest albuterol bronchodilator drug response (BDR). We conducted the largest pharmacogenetic study using whole genome sequencing data from 1,441 minority children with asthma who had extremely high or low albuterol bronchodilator drug response. We identified population-specific and shared pharmacogenetic variants associated with BDR. Our findings help inform the direction of future development of asthma medications and our study advances the foundation of precision medicine for at-risk, yet understudied, racially/ethnically diverse populations.

INTRODUCTION

Asthma is a chronic inflammatory disorder of the airways characterized by

recurrent respiratory symptoms and reversible airway obstruction. Asthma affects

5% of the world population [2] and is the most common chronic disease among

children [3, 4]. In the United States (U.S.), asthma is the most racially disparate

health condition among common diseases [5, 6]. Specifically, U.S. asthma prevalence

is highest among Puerto Ricans (36.5%), intermediate among African Americans

(13.0%) and European Americans (12.1%), and lowest among Mexican Americans

(7.5%) [7]. These disparities also extend to asthma mortality, which is four- to five-

fold higher in Puerto Ricans and African Americans compared to Whites and

Mexican Americans [1].

Current asthma guidelines recommend inhaled β_2-agonists (e.g., albuterol) for

treatment of acute asthma symptoms. Albuterol is a short-acting β_2-adrenergic

receptor (β_2AR) agonist, and it produces bronchodilation by causing rapid smooth

muscle relaxation in the airways. Albuterol is the most commonly prescribed asthma

medication in the world and is the mainstay of acute asthma management across all

ethnic groups [8, 9]. Among low income and minority populations in the U.S.,

albuterol is often the only medication used for asthma regardless of asthma severity

[10, 11]. Response to albuterol is quantified based on bronchodilator drug response

(BDR) using spirometry. We and others have demonstrated that there is significant

variability in BDR among individuals and between populations [12, 13]. Specifically,

the populations with the highest asthma prevalence and mortality also have the

lowest drug response to albuterol: Puerto Rican and African American children have

significantly lower BDR than Whites and Mexican American children [13, 14]. This

variation in drug response across racial/ethnic groups may contribute to the

observed disparities in asthma morbidity and mortality [15-19].

BDR is a complex trait, influenced by environmental and genetic factors, with

heritability estimates ranging from 47% to 92% [20-22]. Genome-wide association

studies (GWAS) have identified several common single nucleotide polymorphisms

(SNPs) associated with BDR in populations of European descent [23-25]. To date,

only one GWAS of BDR has been conducted among African Americans [26]. While

that study identified a novel BDR-associated locus, it did not replicate known

associations discovered in populations of European descent, suggesting that BDR

may be determined in part by population-specific variants. Our previous study of

genetic predictors of BDR in Latino populations identified a significant contribution

of population-specific rare variants to BDR [27].

GWAS were designed to identify common variants associated with disease

through the use of genotyping arrays that relied on linkage disequilibrium to

tag/represent variants not explicitly genotyped on the array itself. Early GWAS

arrays were optimized for performance in populations of European origin and

lacked the ability to capture race-/ethnic-specific genetic variation due to

differences in linkage disequilibrium (LD) across racially/ethnically diverse

populations [28]. Recent generations of arrays have attempted to tailor genotyping

panels for major HapMap populations (Affymetrix Axiom® World Arrays [29]), or to

include population-specific and trans-ethnic tag SNPs to statistically infer genotypes

not directly captured in diverse populations (Illumina Infinium® Multi-Ethnic

Genotyping Array [30]). However, imputation accuracy decreases significantly with

variant frequency [31, 32], making it difficult to use genotyping arrays to study rare

and/or population-specific variants.

The most striking weakness of GWAS is the inability to adequately capture rare

variation. Whole exome sequencing (WES) and other forms of targeted sequencing

were developed to address the inability of genotyping arrays to capture rare

variation. WES only allows for the capture of common and rare variants within

coding and flanking regions. Studies have shown that a large number of variants

associated with complex disease lie within non-coding regions of the genome

(reviewed in Zhang and Lupski, 2015 [33]). Additionally, the target capture

procedures result in uneven sequence coverage, limiting the reliability of SNP

calling for loci close to the boundary of targeted regions. WES also has limited usage

for the detection of structural variation, which depends heavily on uniform coverage

across the genome.

Whole genome sequencing (WGS) is the ideal technology for identifying disease-

causing variants that are rare and/or population-specific. Unlike GWAS genotyping

arrays or targeted sequencing technologies, WGS allows the detection of common

and rare variants in coding and non-coding regions. WGS is the only technology

capable of a truly comprehensive and agnostic evaluation of genetic sequence

variation in the context of complex disease. The persistent lack of large-scale genetic

studies conducted in populations of non-European descent further exacerbates

racial/ethnic disparities in clinical and biomedical research [34-36]. The application

of WGS to the evaluation of genetic factors within a racially/ethnically diverse study

population is a necessary step toward eliminating health disparities in BDR and

other complex phenotypes.

In this study, we performed WGS on 1,441 minority children with asthma from

the tails of the BDR distribution (**S1 Fig**). Our study included high and low drug

responders from three ethnic groups: Puerto Ricans (PR) (n=483), Mexicans (MX)

(n=483), and African Americans (AF) (n=475). An overview of the subject selection

process and main analyses performed in this study is presented in **Fig 1**. We

identified multiple BDR-associated common and rare variants that are population-

specific or shared among populations. This study is part of the National Heart, Lung,

and Blood Institute's Trans-Omics for Precision Medicine Whole Genome

Sequencing (TOPMed) program and represents the largest WGS study thus far to

investigate genetic variants important for bronchodilator drug response in racially

and ethnically diverse children with asthma.

RESULTS

Descriptive characteristics of study subjects

Descriptive characteristics for all study subjects (N=1,441, including 483 Puerto Ricans, 483 Mexicans and 475 African Americans) are summarized in **Table 1**. Covariates and demographic variables were assessed for significant differences between high and low drug responders for each racial/ethnic group. Significant differences were found for age (Mexicans, $p<0.001$), baseline lung function (pre-FEV_1 % predicted, $p<0.001$), total Immunoglobulin E (tIgE, $p < 0.001$), and atopy. Baseline lung function (pre-FEV_1 % predicted) was defined as the percentage of observed FEV_1 relative to the expected population average FEV_1 estimated using the Hankinson lung function prediction equations [37].

225 **Table 1.** Study Population Description (N=1,441).

Descriptive Statistics		Puerto Ricans (N=483)			Mexicans (N=483)			African Americans (N=475)		
		High BDR	Low BDR	*P*	High BDR	Low BDR	*P*	High BDR	Low BDR	*P*
Number of Subjects		239	244	-	243	240	-	233	242	-
Percent Male		53.6%	53.3%	1.0	60.1%	52.1%	0.08	55.4%	47.9%	0.12
Median Age, yr (IQR)		11.6 (9.7 - 14.8)	12.2 (10.1 - 15.2)	0.18	11.7 (9.6 - 14.0)	13.3 (10.6 - 16.0)	<0.001	13.8 (11.0 - 16.8)	13.8 (10.9 - 17.1)	0.48
Mean Global Ancestry Proportions	AFR	0.24	0.22	0.44	0.05	0.05	0.37	0.79	0.79	0.80
	EUR	0.63	0.64	0.27	0.37	0.36	0.84	0.19	0.20	0.70
	NAM	0.13	0.13	0.93	0.58	0.59	0.90	0.02	0.02	0.90
BMI Category, N	Obese	76	67	0.32	100	96	0.85	82	83	0.85
	Non-Obese	163	177		143	144		151	159	
Pre-FEV$_1$ % Predicted, N	< 80%	149	56	<0.001	43	7	<0.001	47	6	<0.001
	≥ 80%	90	188		200	233		186	236	
Median ΔFEV$_1$, % (IQR)		21.2 (18.2 - 25.7)	5.0 (2.9 - 6.3)	-	12.7 (10.3 - 16.8)	3.6 (2.0 - 4.9)	-	15.5 (13.3 - 20.3)	3.3 (2.0 - 4.4)	-
Median tIgE, mL (IQR)		407.5 (126.8-952.8)	191.9 (50.5-542.2)	<0.001	247.5 (64.2-817.0)	105.7 (35.4-332.0)	<0.001	281.6 (97.3-552.4)	128.8 (36.6-351.3)	<0.001
Atopy, N		177	118	<0.001	155	117	<0.001	139	111	<0.001

226
BDR: bronchodilator drug response. IQR: interquartile range. **Pre-FEV$_1$ % predicted**: percentage of measured FEV$_1$ relative to predicted FEV$_1$
estimated by the Hankinson lung function prediction equations prior to administration to albuterol. **ΔFEV$_1$**: a quantitative measure of BDR,
measured as the percent change in baseline FEV$_1$ after administration of albuterol. High and low drug responders were chosen from the
extremes of the BDR (ΔFEV$_1$) distribution. **tIgE**: measure of total Immunoglobulin E from serum in milliliters. **Atopy**: tIgE measurement greater
than or equal to 100.

We estimated genetic ancestry for all participants (see **Methods**) and found that

the major ancestry proportions in Puerto Ricans, Mexicans and African Americans

are European, Native American and African ancestries, respectively (**Table 1**, **S2**

Fig). Analysis of genetic substructure of the three admixed populations by principal

component analysis (PCA) demonstrated that the three populations displayed the

characteristic spectrum of ancestry found in admixed populations (**S3 Fig**).

Variant summary statistics

Genetic variant summary statistics revealed that the average number of variants by

population corresponded to the proportion of African ancestry: the most variants

were found among African Americans, followed by Puerto Ricans and Mexicans (**Fig**

2a, **Table 2**).

Table 2. Summary statistics of variants

	Puerto Ricans	Mexicans	African Americans
Range in variants, per individual (mean)			
All	3.7M – 4.4 M (4.0 M)	3.5M – 4.1M (3.7M)	4.0M – 4.5M (4.3M)
SNPs	3.4M – 4.0 M (3.6M)	3.2M - 3.7M (3.4M)	3.6M – 4.1M (4.0M)
Indels	284,067 – 344,493 (305,618)	272,635 – 321,778 (289,096)	311,997 – 354,646 (339,570)
Others*	21,233 – 29,177 (24,728)	19,703 – 26,772 (23,190)	22,014 – 30,993 (27,911)
No. of biallelic SNPs, union**	29.2M	28.1M	36.3M
By allele frequency of data			
Rare (< 1%)	18,169,292 (62%)	20,029,291 (71%)	22,847,938 (63%)
Common (≥ 1%)	11,007,125 (38%)	8,092,157 (29%)	13,459,004 (37%)
Population-specific***	6,680,909	9,687,651	14,114,142
Singleton	4,616,383 (69%)	7,983,950 (82%)	10,574,879 (75%)
< 5%	6,676,286 (> 99%)	9,675,271 (> 99%)	14,096,844 (> 99%)
≥ 5%	4,623 (< 1%)	12,380 (< 1%)	17,298 (< 1%)
By novelty (dbSNP150)			
Known	27,034,179 (93%)	25,911,801 (92%)	33,540,387 (92%)
Novel	2,142,238 (7%)	2,209,647 (8%)	2,766,555 (8%)
By protein impact****			
Coding	284,269 (1%)	289,055 (1%)	362,823 (1%)
Nonsynonymous	157,541	164,312	203,029
Stopgain / stoploss	3,194	3,531	4,158
Splicing	2,111	2,214	2,620
Other coding	121,423	118,998	153,016
Noncoding	28,833,461 (99%)	27,769,434 (99%)	35,878,723 (99%)
Not annotated^	58,687 (< 1%)	62,959 (< 1%)	65,396 (< 1%)

* Includes multi-nucleotide polymorphism (MNPs), complex, symbolic and mixed variants as
defined by GATK VariantEval.
** Biallelic SNPs with less than 10% genotype missingness per population were included.
*** Biallelic SNPs that are present in only one of the three studied populations.
**** Other coding variants include those annotated as exonic and synonymous in ANNOVAR.
^ Not annotated: biallelic SNPs not included in the annotation pipeline because they are not
present in TOPMed freeze 2 and 3 data releases.

The majority of observed variants (>90%) were SNPs. The union of biallelic SNPs

from all individuals in each population varied from 28.1M among Mexicans, 29.2M

among Puerto Ricans to 36.3M among African Americans. Approximately 65% of

biallelic SNPs were rare (non-reference allele frequency < 1%, **Fig 2b**, **Table 2**).

Biallelic SNPs that were population-specific (i.e., SNPs found in only one population)

accounted for 23% (6.68M / 29.2M in Puerto Ricans) to 39% (14.1M / 36.3M in

African Americans) of the biallelic SNPs observed in each population. Over 99% of

the population-specific SNPs had a non-reference allele frequency less than 5% and

the majority of these population-specific SNPs (69% to 82%) were also singletons

(**Fig 2c**, **Table 2**). Based on dbSNP build 150, an average of 8% of biallelic SNPs

were novel (**Fig 2d**, **Table 2**).

In all three populations, 99% of the biallelic SNPs were observed in noncoding

regions. Based on the Combined Annotation Dependent Depletion (CADD) score

[38], which estimates the deleteriousness of a variant, over 99% of highly

deleterious biallelic SNPs (CADD score ≥ 25) were observed in coding regions,

regardless of ethnicity (**Table 3**). This may be due to the relatively limited

availability of functional annotations as positive training data for CADD to estimate

deleteriousness in non-coding regions [39]. The percentage of singletons in these

highly deleterious biallelic SNPs varied from 51% (Puerto Ricans) to 70%

(Mexicans).

Table 3. CADD score summary statistics of biallelic SNPs*.

Population	CADD score	No. of SNPs*	% Coding	% Singletons
Puerto Ricans	0-9	26,936,285	0.5%	29%
	10-19	2,027,392	4%	32%
	20-24	123,481	32%	37%
	≥ 25	30,572	>99%	51%
Mexicans	0-9	25,932,006	0.5%	44%
	10-19	1,967,096	4%	49%
	20-24	124,173	34%	55%
	≥ 25	35,214	>99%	70%
African Americans	0-9	33,468,245	0.5%	36%
	10-19	2,573,523	4%	40%
	20-24	158,967	32%	46%
	≥ 25	40,811	>99%	64%

272   *Biallelic SNPs with less than 10% genotype missingness per population were included.

BDR association testing with common variants

We performed genome-wide association testing of common variants with BDR
(dichotomized as high/low drug responders from the extremes of the BDR
distribution) for each population, adjusting by age, sex, body mass index (BMI)
categories, and the first ten principal components (PCs) (see **Methods** section
"Single locus BDR association testing on common variants" for rationale on
including these covariates). We then performed a trans-ethnic meta-analysis on
these results across all three populations. For all three populations, 94% of all
common variants tested are known variants annotated in dbSNP Build 150 (**S1**
Table).

A universal p-value threshold of 5×10^{-8} is often used to determine significance in

GWAS. This statistical threshold was calculated based on Bonferroni correction

under the assumption of 1,000,000 independent tests using patterns of linkage

disequilibrium based primarily on individuals of European descent and has been

shown to be non-generalizable for WGS studies or genetic studies of non-European

populations in general [40-42].The number of independent tests varies by LD

patterns, which in turn vary by race/ethnicity [41]. We calculated the effective

number of independent tests for each population and for our trans-ethnic meta-

analysis, and generated racially/ethnically adjusted genome-wide significance

thresholds for each population (see **Methods**). Population-specific genome-wide

significance thresholds after correcting for the number of effective tests (adjusted

genome-wide significance level) were 1.57×10^{-7} for Puerto Ricans, 2.42×10^{-7} for

Mexicans, and 9.59×10^{-8} for African Americans (see **Methods**). These numbers are

highly concordant with WGS significance thresholds derived from the African

American (ASW), Mexican (MXL), and Puerto Rican (PUR) 1000 Genomes

sequencing data [41]. The adjusted genome-wide significance level for our trans-

ethnic meta-analysis was 3.53×10^{-7}. Significance thresholds for discovery analyses

in genome-wide association studies can often produce false negative results [43-46].

To minimize Type II error, suggestive associations are often included in replication

and functional validation studies. We identified suggestive associations based on the

following widely used formula: 1/(effective number of tests) [43-46].

While no significant associations were identified from the population-specific

analyses (**S8 Fig**), our trans-ethnic meta-analysis identified ten unique loci

(represented by 27 SNPs) significantly ($p < 3.53$ x 10^{-7}) or suggestively ($p < 7.06$ ×

10^{-6}) associated with BDR status (**Fig 3a, Table 4, S2 Table)**. We annotated all 27

SNPs by performing a thorough bioinformatics search in ENCODE, NHGRI-EBL

GWAS Catalog and PubMed databases. Their previously reported lung-related

phenotype associations and functional annotations are reported in **S4 Table** and **S9**

Table.

Two SNPs, rs17834628 and rs35661809, located on chromosome 5 were

significantly associated with BDR ($p = 1.18$ x 10^{-8} and 3.33 x 10^{-8}); additionally,

population-specific analyses show that the direction of effect for these two variants

is concordant across all three populations (**Fig 3b, S3 Table**). **Fig 3c** displays a

LocusZoom plot of rs17834628 with 400 kb flanking regions. Three of the 27

identified SNPs were located within genes. Specifically, two SNPs are located in the

third and fifth introns of *NFKB1* (rs28450894 and rs4648006), and a third SNP,

rs16995064, mapped to intron 7 of *PLCB1* **(Table 4)**. Among the *NFKB1* SNPs, the

low BDR-associated T allele of rs28450894 is found predominantly among African

populations (minor allele frequency [MAF] 8.8% – 28.7%), followed by European

populations (MAF 3.7% - 7.6%) and Puerto Ricans (MAF 6.2%), and is relatively

rare in Mexicans (MAF 1.5%) based on 1000 Genomes data (**S6 Fig**). Combined,

these 27 SNPs explain 23%, 16%, and 18% of the variation in BDR status in Puerto

Ricans, Mexicans, and African Americans, respectively **(S5 Table)**.

Table 4. Results from trans-ethnic BDR association tests for common variants.

Chr	Start	rsID	Effect allele	OR (95%CI)	p	Effect allele frequency			Nearest genes
						PR	MX	AA	
5	12978566	rs17834628	A	1.67 (1.29-2.16)	1.18E-08*	0.32	0.42	0.17	*LINC01194* (-173k), *MIR4454* (-311k), *CTNND2* (-1074k), *DNAH5* (712k)
5	12968341	rs35661809	G	1.59 (1.20-2.10)	3.33E-08*	0.34	0.43	0.24	*LINC01194* (-163k), *MIR4454* (-300k), *CTNND2* (-1064k), *DNAH5* (722k)
5	12975934	rs17237639	G	1.61 (1.30-2.00)	1.22E-07	0.31	0.43	0.16	*LINC01194* (-171k), *MIR4454* (-308k), *CTNND2* (-1072k), *DNAH5* (715k)
5	12975187	rs1017452	G	1.60 (1.31-1.96)	2.11E-07	0.31	0.43	0.16	*LINC01194* (-170k), *MIR4454* (-307k), *CTNND2* (-1071k), *DNAH5* (715k)
5	12975322	rs1017454	A	1.60 (1.31-1.96)	2.11E-07	0.31	0.43	0.16	*LINC01194* (-170k), *MIR4454* (-307k), *CTNND2* (-1071k), *DNAH5* (715k)
5	12975265	rs1017453	C	1.56 (1.25-1.95)	6.40E-07	0.31	0.42	0.16	*LINC01194* (-170k), *MIR4454* (-307k), *CTNND2* (-1071k), *DNAH5* (715k)
5	12972636	rs17237443	C	1.59 (1.28-1.97)	9.85E-07	0.29	0.42	0.11	*LINC01194* (-170k), *MIR4454* (-307k), *CTNND2* (-1071k), *DNAH5* (715k)
1	209324294	rs10746419	T	1.29 (0.75-2.25)	1.19E-06	0.49	0.54	0.53	*MIR205HG* (278k), *MIR205* (281k), *CAMK1G* (433k), *LAMB3* (464k)
5	12961545	rs17833938	A	1.56 (1.28-1.91)	1.45E-06	0.30	0.42	0.12	*LINC01194* (-156k), *MIR4454* (-294k), *CTNND2* (-1057k), *DNAH5* (729k)
6	104240500	rs13437006	C	1.56 (1.21-2.02)	1.61E-06	0.22	0.24	0.32	*HACE1* (935k), *LINC00577* (1144k), *LIN28B* (1164k)
15	101230457	rs1565749	A	1.66 (1.18-2.32)	1.64E-06	0.18	0.15	0.18	*ASB7* (-39k), *LINS1* (-88k), *PRKXP1* (-131k)
5	12948369	rs34845041	T	1.56 (1.26-1.92)	1.77E-06	0.30	0.42	0.12	*LINC01194* (-143k), *MIR4454* (-280k), *CTNND2* (-1044k), *DNAH5* (742k)
5	12975108	rs1017451	T	1.55 (1.24-1.93)	1.96E-06	0.30	0.42	0.13	*LINC01194* (-170k), *MIR4454* (-307k), *CTNND2* (-1071k), *DNAH5* (715k)
5	12950432	rs62347395	G	1.55 (1.26-1.92)	2.02E-06	0.30	0.42	0.12	*LINC01194* (-145k), *MIR4454* (-282k), *CTNND2* (-1046k), *DNAH5* (740k)
15	101231049	rs57924834	A	1.59 (1.25-2.03)	2.04E-06	0.23	0.20	0.20	*ASB7* (-39k), *LINS1* (-89k), *PRKXP1* (-132k)
4	137382142	rs17048684	A	1.8 (1.06-3.05)	2.20E-06	0.11	0.14	0.18	*LINC00613* (-547k), *PCDH18* (1058k)
5	12959598	rs1438293	G	1.55 (1.24-1.93)	2.73E-06	0.29	0.42	0.11	*LINC01194* (-154k), *MIR4454* (-292k), *CTNND2* (-1055k), *DNAH5* (731k)
20	8635168	rs16995064	G	1.96 (1.12-3.43)	3.30E-06	0.12	0.13	0.05	**PLCB1 (intron 7)**, *PLCB4* (415k)
12	19821401	rs66544720	T	0.66 (0.55-0.78)	3.66E-06	0.33	0.37	0.16	*AEBP2* (-146k), *PLEKHA5* (-292k)
6	104235591	rs6926020	C	1.57 (1.25-1.97)	3.68E-06	0.19	0.23	0.27	*HACE1* (940k), *LINC00577* (1149k), *LIN28B* (1169k)
4	103453535	rs28450894	T	0.47 (0.34-0.64)	3.75E-06	0.06	0.03	0.12	*SLC39A8* (-187k), **NFKB1 (intron 3)**, *MANBA* (99k)
4	103461559	rs4648006	T	0.47 (0.34-0.64)	3.75E-06	0.06	0.03	0.12	*SLC39A8* (-195k), **NFKB1 (intron 5)**, *MANBA* (91k)
22	27826429	rs60163793	G	2.01 (1.20-3.38)	4.30E-06	0.04	0.14	0.15	*MN1* (318k), *PITPNB* (421k)
12	19824386	rs7313907	C	0.66 (0.55-0.79)	4.35E-06	0.33	0.37	0.16	*AEBP2* (-149k), *PLEKHA5* (-295k)
12	19820677	rs11044754	A	0.66 (0.55-0.79)	4.54E-06	0.33	0.37	0.16	*AEBP2* (-146k), *PLEKHA5* (-291k)
15	101233236	rs55638658	A	1.61 (1.13-2.30)	5.08E-06	0.18	0.15	0.18	*ASB7* (-41k), *LINS1* (-91k), *PRKXP1* (-134k)
6	54581204	rs13200833	A	0.66 (0.48-0.90)	5.15E-06	0.32	0.24	0.22	*TINAG* (-326k), *MLIP* (-450k), *FAM83B* (130k)

The top ten unique loci (represented by 27 SNPs) significantly ($p < 3.53 \times 10^{-7}$) or suggestively ($p < 7.06 \times 10^{-6}$) associated with BDR status in our trans-ethnic meta-analysis. **Chr and Start**: chromosome locations of SNPs in GRCh37 coordinates. All significantly and suggestively associated common variants are presented above. **Nearest genes**: The four nearest transcripts from RefSeq were identified and genes with multiple transcripts were reported once only with the distance to the nearest transcripts indicated in parentheses. Negative distances indicate upstream genes. Genes that overlap with BDR-associated SNPs are bold. High drug responders were assigned as cases in analyses throughout this study.* p-values that achieve adjusted genome-wide significance for trans-ethnic meta-analysis ($p < 3.53 \times 10^{-7}$).

It has been shown that functionally relevant variants do not always display the

lowest *p*-values in association studies [47]. To avoid false negative results, it is

strongly suggested that replication analyses for two-stage genome-wide studies [43-

46]. Therefore, we included all 27 SNPs in replication analyses we performed

separately and via meta-analysis in five independent populations (GALA I, SAGE I,

HPR, SAPPHIRE and CHOP) (**S6-S7 Table**). None of the 27 SNPs were significantly

associated with BDR status in our replication analyses **(S6-S7 Table)**. It is

important to note that our largest replication cohort (SAPPHIRE) did not include

children (median age = 30 and 28 for high and low BDR groups, respectively, **S6**

Table). All other replication populations included less than 500 individuals per

study **(S6 Table)**. It is well known that there are age-specific associations with

asthma and asthma-related phenotypes [48, 49]. It is unclear whether the same is

true for BDR. It is possible that the overrepresentation of adult patients in our

available replication populations may explain, in part, our lack of replication.

In addition to performing WGS association analyses to identify genetic variants

associated with variation in BDR, we also performed H3K27ac chromatin

immunoprecipitation sequencing analysis (ChIP-seq) experiments in primary

bronchial smooth muscle cells (BSMCs) to identify potential regulatory regions

marked by H3K27ac peaks. Albuterol's mechanism of action involves binding with

the β_2-adrenergic receptor in bronchial smooth muscle cells causing rapid onset of

airway tissue relaxation and bronchodilation. BSMCs are therefore considered one

of the most relevant cell types for molecular studies of BDR [50]. We observed two

H3K27ac ChIP-seq signals that overlapped with variants in moderate to high LD (R^2

= 0.47 to 0.82) with two *NFKB1* SNPs (rs28450894 and rs4648006) we identified

through trans-ethnic meta-analysis, implying that these variants may have

regulatory functions in BSMC (**S5 Fig a, S8 Table**).

The gold standard for identifying true signals in genetic association studies is to

use *p*-values from a primary and/or replication study to prioritize variants for

further investigation. The use of *p*-values as the sole metric for prioritization is

problematic for three reasons: (1) the *p*-value statistic is dependent on sample size

and effect magnitude, (2) *p*-values do not incorporate biological knowledge, and (3)

one cannot use *p*-values to distinguish between true association signals and noise of

the same magnitude [47, 51, 52]. Instead of relying solely on *p*-values, we applied

the Diverse Convergent Evidence (DiCE) [53] approach to prioritize each of the 27

BDR-associated SNPs from our trans-ethnic meta-analysis for inclusion in further

function analyses **(S15 Table, S4 Fig)**. After integrating information from our WGS

analysis, publicly available bioinformatics data, and ChIP-Seq experiments in BSMCs,

the *NFKB1* locus had the highest DiCE evidence score, indicating that this locus had

the strongest evidence of functional relevance to BDR variation **(S4 Fig)**. Therefore,

all further functional experiments were focused on variants within this locus.

Functional assays on the NFKB1 Locus

Both H3K27ac ChIP-seq regions that overlapped with the BDR-associated *NFKB1*

locus were tested for enhancer activity using luciferase enhancer assays. The

sequences of these two *NFKB1* intronic regions were cloned into a pGL4.23

enhancer assay vector (Promega, Madison, WI, USA), which contains a minimal

promoter and a luciferase reporter gene. The pGL4.23 vector with the viral SV40

promoter was used as a positive control, and the pGL4.23 empty vector as a

negative control. All constructs were tested for their enhancer activity in BSMCs.

One enhancer, *NFKB1* Region 2, showed significantly increased enhancer activity

over empty vector (2.24-fold increase, p = 8.70 x 10^{-6}, unpaired t-test; **S5 Fig b**).

Given the relevance of *NFKB1* in immune pathways and asthma, we also

performed RNA sequencing (RNA-seq) experiments to verify whether the identified

intronic *NFKB1* SNPs regulate gene expression of neighboring genes. Among genes

within 1Mb of rs28450894 meeting expression reliability cutoffs (see **Methods**), we

found that the low BDR-associated T allele of rs28450894 is significantly associated

with decreased expression of *SLC39A8* in blood (**S7 Fig**, p = 0.0066, FDR-adjusted p

= 0.0856, $\log_2(\beta)$ = -0.327).

We observed that two known BDR candidate genes, *ADCY9* and *CRHR2*, which

achieved replication in a previous GWAS of BDR performed in the full GALA II

population but did not replicate in the current study (**S10 Table**) [27]. In the

previous study, GWAS array data, supplemented by imputation, were used to

evaluate genetic associations with BDR measured as a continuous trait. To

determine whether the discrepancy between findings was due to data type

(imputed array-based vs. WGS-based) or study design (continuous trait vs. extreme

phenotype), the common variant analysis in the current analysis was repeated

among the subset of samples that had array-based and WGS data (n = 1,414 out of

1,441). Based on the top 1,000 BDR-associated SNPs from the current common

variant analysis, there was high correlation between association p-values generated

from imputed array-based and WGS-based genotypes (Spearman correlation = 1.0),

suggesting that data type is not the cause of the observed discrepancy (**S9 Fig a**).

Nearly all SNPs with high imputation R^2 exhibited high genotype concordance

between array-based and WGS-based genotypes, confirming high imputation quality

for most common SNPs ($\geq$ 99.7%). (**S9 Fig b and c**). We also performed linear

regression to analyze BDR (ΔFEV_1) as a continuous trait using imputed array-based

data. The most significantly associated SNP identified in the trans-ethnic meta-

analysis using extreme phenotype analysis displayed the same direction of effect as

analyzing BDR as a continuous trait (OR=1.67 in extreme phenotype analysis and

β=0.51 in continuous analysis). These observations indicate that the discrepancy

between findings may be due to differences in statistical power afforded by the

different study designs (continuous trait vs. extreme phenotype). For common

variant analyses, dichotomization of a continuous outcome results in a loss of

statistical power [54-57]. For example, a population of 2,000 individuals has 80%

power to identify moderate genetic associations (β=0.3) for common variants with

minor allele frequencies $\geq$ 0.05 when the outcome is continuous. If this population

were re-analyzed after dichotomizing the continuous outcome at the median

(cases=1,000, controls=1,000), power would be reduced to 62%. The opposite effect

is observed in rare variant analyses. The extreme phenotype study design is a

specific type of dichotomous outcome study design that has been shown to increase

power and the probability of identifying functional rare variants [56-58]. It should

also be noted that the previously published results were discovered in one

population (Puerto Ricans), whereas the results from our trans-ethnic meta-analysis

describe associations that are conserved across three populations (Puerto Ricans,

Mexicans, and African Americans).

BDR association testing using common and rare variants

We tested the combined effects of common and rare variants on BDR using SKAT-O

[59] to examine variants in 1kb sliding windows, which moved across the genome in

500bp increments. The same covariates used for common variant association

testing were applied.

After identifying the effective number of tests and adjusting for multiple

comparisons on each population separately (see **Methods**), we identified three

population-specific loci associated with BDR at genome-wide significance levels;

two were found in Mexicans on chromosome 1 and chromosome 11, and one in

African Americans on chromosome 19 (**Fig 4a-c, Table 5, S11 Table**).

Table 5. Results from association testing on combined effects of common and rare variants on BDR

Chr	Start	Stop	*p*	Population	nCommon	nRare	Nearest genes
1	114177000	114178000	4.40E-09	MX	2	1	***MAGI3* (intron 9)**, *PHTF1* (62k), *RSBN1* (126k)
11	27507000	27508000	6.59E-09	MX	2	3	*LOC105376671* (-3k), *LGR4* (-13k), *LIN7C* (8k)
19	10424000	10425000	3.12E-11	AA	1	2	*ZGLP1* (-4k), *ICAM5* (-17k), ***FDX1L* (intron 3)**, *RAVER1* (2k)
4	73478000	73479000	6.25E-08	Combined	10	23	*ADAMTS3* (-43k), *COX18* (441k)
8	97926000	97927000	1.32E-08	Combined	3	13	*SDC2* (-302k), ***CPQ* (intron 4)**, *LOC101927066* (37k), *TSPYL5* (359k)

Chr, start and stop: GRCh37 chromosome coordinates; **AA**: African Americans; **MX**: Mexicans; **Combined**: all individuals in all three populations. **nCommon**
and **nRare**: number of common and rare variants, respectively. **Nearest genes**: The four nearest transcripts from RefSeq were identified and genes with
multiple transcripts were reported once only with the distance to the nearest transcripts indicated in parentheses. Negative distances indicate upstream
genes. Genes that overlap with BDR-associated SNPs are bold.

We also performed association testing across all three populations in a single

analysis. Pooling subjects increased the sample size and thereby maximized the

power of the SKAT-O association test. To minimize any potential effect of

confounding by population substructure, association testing also included local

genetic ancestry, defined as the proportions of Native American and African

ancestries for the window under testing. Two loci on chromosomes 4 and 8 were

found to be genome-wide significant ($p < 1.53 \times 10^{-7}$) (**Fig 4d, Table 5**). A total of 60

variants were identified from all SKAT-O regions reported in **Table 5**. Six of the 60

variants were located within predicted regulatory regions (**S12 Table**). Specifically,

three variants located on chromosome 11 identified in Mexicans overlap with a

CTCF (transcriptional repressor) binding site and comprise a chromatin insulator

region. The five regions identified in our combined and population- specific SKAT-O

analyses independently explained 4% to 8% of the variation in BDR in their

respective populations (**Table 5, S5 Table**).

We examined alternative grouping strategies for rare variants, including

grouping by (1) genes from transcription start to end sites with or without 50kb

flanking regions, (2) transcription start site with 20kb flanking regions, and (3)

H3K27ac ChIP-seq peaks in airway epithelial cells and airway smooth muscle cells.

Association tests with these alternate grouping strategies identified no further

significant associations.

DISCUSSION

We identified population-specific and shared common and rare variants associated

with bronchodilator drug response in three ethnically diverse populations of

children with asthma. WGS, unlike GWAS genotyping arrays and targeted

sequencing, provides comprehensive detection of common and rare variants in

coding and non-coding regions. African Americans, Latinos, and other minorities

have been dramatically underrepresented in GWAS [34-36]. Combined, the 27

variants identified from our common variant analyses (**Table 4**) explained 23%,

16%, and 18% of the variation in BDR in Puerto Ricans, Mexicans, and African

Americans, respectively, after adjusting for clinical covariates (**S5 Table**). The five

SKAT-O regions identified in our combined and population-specific analyses

independently explained 4% to 8% of the BDR variation in their respective

populations (**Table 5, S5 Table**). Our study represents an important investment

from the NIH/NHLBI to include underrepresented populations in large whole

genome sequencing efforts and to improve racial/ethnic diversity in clinical and

biomedical research.

Our trans-ethnic common variants meta-analysis identified one locus on

chromosome 5 that was associated with BDR at a genome-wide significance level (p

$< 5 \times 10^{-8}$). The proximity of this BDR-associated locus to *DNAH5* and *LINC01194* is

of particular interest. A SNP in *DNAH5* has been associated with total lung capacity

in White subjects with chronic obstructive pulmonary disease [60]. In a separate

GWAS, the *DNAH5/LINC01194* locus was reported among Europeans to be

associated with levels of IgE [61, 62], a biomarker associated with asthma

endotypes. Baseline lung function (FEV_1) and total IgE levels are associated with

asthma severity and can predispose an individual to lower bronchodilator drug

responsiveness [13, 14, 63]. We found two *NFKB1* intronic variants on chromosome

4 associated with BDR at a suggestive significance level. The NFκB protein has a

known role in allergic response, and various studies have demonstrated that the

NFκB pathway is activated in patients with asthma, as reviewed by Edwards *et al.*

[64].

ChIP-seq and functional enhancer assays in BSMCs suggest these *NFKB1* intronic

variants may regulate expression of nearby genes. This was in fact supported by our

RNA-seq data, which showed that individuals with the low BDR-associated T allele

genotype displayed reduced expression of the neighboring gene *SLC39A8*, which has

previously been found to be responsive to cytokine treatment in airway epithelial

cells [65] and had reduced expression in mice with allergic airway inflammation

[66]. Recent studies have also shown that *SLC39A8* is unique among zinc

transporters in that upregulation of *SLC39A8* is sufficient to protect lung epithelium

against TNF-α-induced cytotoxicity [67]. Additionally, the higher frequency of the

low BDR-associated allele (T allele of rs28450894 in *NFKB1*) in African populations

suggests that the low BDR-associated allele tracks with African ancestry. This may

explain why admixed populations with higher proportions of African ancestry, i.e.,

African Americans and Puerto Ricans, have lower bronchodilator drug

responsiveness [14], and by extension may shed light on the higher asthma

morbidity and mortality in these populations.

Another intronic variant (chromosome 20, rs16995064, *PLCB1* intron 7) was

associated with BDR at a suggestive significance level. *PLCB1* is highly relevant, as

this gene has been reported to be differentially expressed in therapy-resistant

childhood asthma compared to controlled persistent asthma or age-matched

healthy control subjects in a Swedish cohort [68]. Functional studies also reported

that silencing *PLCB1* inhibited the effect of lipopolysaccharide-induced endothelial

cell inflammation through inhibiting expression of proinflammatory cytokines [69].

Further functional studies are necessary to establish the role of *NFKB1* and *PLCB1*

on BDR.

Apart from assessing the individual effect of common variants on BDR, we also

identified various combined effects of rare variants that were population-specific or

shared across populations. While some of the nearest genes are uncharacterized or

have no known functional relationship to BDR (*MAGI3*, *LOC105376671*, *LIN7C* and

CPQ), there appears to be functional relevance for the locus between *ADAMTS3* and

COX18. The *ADAMTS3* and *COX18* locus were associated with β-adrenergic responses

in cardiovascular-related traits in mice [70]. This locus was significantly associated

with cardiac atrial weight in mice treated with the β blocker atenolol; the

association also replicated in mice treated with the β agonist isoproterenol. These

findings suggest that SNPs found in this locus may modify β adrenergic signaling

pathways in BDR. In the present study, we also identified BDR association with rare

variants within the *CPQ* gene, which encodes a protein from the carboxypeptidase

family. Although no previous BDR association has been identified for *CPQ*, another

member of the carboxypeptidase family, carboxypeptidase A3 (CPA3), is known to

be expressed at higher levels in the airway epithelium among subjects with T_H2-

high asthma [71, 72]. Further studies are necessary to determine the role of *CPQ* in

BDR.

GWAS-based BDR-associated common variants in GALA II have previously been

reported [27]. However, these variants did not replicate in the current study, likely

due to different study designs between the previous and current investigations. The

previous BDR GWAS used an array-based genotyping panel to examine children

with asthma across the entire BDR spectrum, i.e., BDR was used as a continuous

variable. In contrast, the current study sequenced the entire genome to investigate

only the extremes of the BDR distribution (i.e., high and low drug responders). By

repeating our current analysis using a subset of individuals who had array and WGS

data, we confirmed that the major discrepancy between the two studies is due to

study design instead of differences in data type. The contrast in results between

GWAS and WGS due to differences in study design implies that varied study designs

are necessary for a comprehensive understanding of variants associated with

asthma-related phenotypes and drug response. Studying samples from the extreme

tails of drug response distribution has been recognized as one of the success factors

in the study design of pharmacogenomic GWAS [73]. Furthermore, it was recently

demonstrated that the power gain from studying extreme phenotypes is much

greater in analyses of rare variants compared to common variant studies [55]. Since

cost is often a limiting factor for WGS studies, choosing an extreme phenotypic study

design may be beneficial for the study of rare variants and the discovery of common

variant associations that may otherwise be missed when sampling across the entire

phenotypic spectrum.

We did not identify BDR-associated variants from β_2AR signaling pathways.

Instead, most of the BDR-associated genes identified in this study are related to lung

function and allergic response, including total IgE levels and cytokine production in

mast cells. This suggests that at least part of BDR may be due to the predisposition

or intrinsic state of airway smooth muscle cells. Genetic variation may determine

individuals' intrinsic expression levels of candidate genes, which in turn determine

whether their response to albuterol is beneficial.

A higher percentage of African ancestry often implies a higher degree of genetic

variation [74]. Although Puerto Ricans have higher proportions of African ancestry

than Mexicans (**Table 1**), they have fewer population-specific SNPs, an observation

that is consistent with findings from our contributions to the 1000 Genomes Project

[75] and our independent work. This is likely due to the fact that Puerto Ricans have

gone through recent population bottlenecks [76]. We demonstrated that Puerto

Ricans may be more genetically related than expected [76], suggesting that our

current relatedness filters may be too conservative for Puerto Ricans.

Including admixed populations in whole genome sequencing studies has

important scientific implications. First, it allows for discovery of genetic variation of

multiple ancestral populations in a single study. Second, it is extremely useful to

study admixed populations with ancestries that are currently underrepresented in

existing genetic repositories. For example, the widely popular PCSK9 inhibitors used

to treat hypercholesterolemia were discovered by studying the genetics of African

Americans but the biology and final drug development have benefited all patients

regardless of race/ethnicity [77]. Finally, studying admixed populations such as

Mexicans will enhance the understanding of genetic variation in Native American

ancestry, an area that is currently lacking in all major sequencing efforts.

Although an extensive effort was made to replicate the top BDR-associated

variants, we were unable to replicate our results because few studies of non-

European populations exist, as we and others have documented [34-36, 78]. Our

efforts to perform replication of rare BDR-associated variants were further hindered

by the lack of studies with whole genome sequencing data. These challenges

highlight the need to include more racially/ethnically diverse populations in all

clinical and biomedical research.

In an era of precision medicine, addressing questions about the impact of genetic

factors on therapeutic drug response in globally diverse populations is essential for

making precision medicine socially and scientifically precise [5]. This study

advances our understanding of genetic analysis in admixed populations and may

play an important role in advancing the foundation of precision medicine for

understudied and racially and ethnically diverse populations.

METHODS

Data availability

TOPMed whole genome sequencing data are available to download by submitting a

data access request through dbGaP. The dbGaP study accession numbers for GALA II

and SAGE II are phs000920.v1.p1 and phs000921.v1.p1. WGS and array genotype

data for each study are available through dbGaP under the same accession numbers.

Study cohorts and sample details

This study examined a subset of subjects with asthma from the Study of African

Americans, Asthma, Genes & Environments (SAGE II) [49, 79-81] and the Genes-

Environments & Admixture in Latino Americans (GALA II) study [27]. SAGE II

recruited African American subjects from the San Francisco Bay area. GALA II

recruited Latino subjects from Puerto Rico and the mainland United States (Bronx,

NY; Chicago, IL; Houston, TX; San Francisco Bay Area, CA). Ethnicity of the subjects

was self-reported and all four of the participant's biological grandparents must have

reported the same ethnicity.

A total of 1,484 individuals from three ethnic groups (494 Puerto Ricans, 500

Mexicans and 490 African Americans), representing the extremes of the

bronchodilator response (BDR, see below) distribution were selected for whole

genome sequencing. Genomic DNA was extracted and purified from whole blood

using Wizard® Genomic DNA Purification Kits (Promega, Madison, WI, USA).

Bronchodilator response measurements

Spirometry was performed and BDR (i.e., ΔFEV_1, defined as the relative change in

FEV_1) was calculated as previously described [27]. In brief, BDR was calculated

according to American Thoracic Society/European Respiratory Society guidelines

[82] as the percent change in FEV_1 after 2 doses of albuterol: that is, BDR = (post-

FEV_1 – pre-FEV_1) / pre-FEV_1.

High and low drug responders were selected from the extremes of BDR

distribution from GALA II and SAGE II (**Fig 1**). **S1 Fig** highlighted the BDR

distribution of the 1,441 individuals who passed WGS data quality control (see

"WGS data processing and quality control"). The ΔFEV_1 cutoffs for high and low

responders are as follows: high responders ($\Delta FEV_1 > 16.29$ for Puerto Ricans, > 8.55

for Mexicans and > 11.81 for African Americans); low responders ($\Delta FEV_1 < 7.23$ for

Puerto Ricans, < 6.05 for Mexicans and < 5.53 for African Americans).

Analysis on descriptive data of study subjects

Dichotomous variables were tested for association with BDR using Fisher's exact

test. Continuous variables were tested for normality using the Shapiro-Wilk test.

Normally and non-normally distributed continuous variables were tested using

Student's t-test and the Wilcoxon rank-sum test, respectively.

Sample quality control and whole genome sequencing

DNA samples were quantified by fluorescence using the Quant-iT PicoGreen dsDNA

assay (ThermoFisher Scientific, Waltham, MA, USA) on a Spectramax fluorometer

(Molecular Devices, Sunnyvale, CA, USA). Sample integrity was ascertained using the

Fragment Analyzer™ (Advanced Analytical Technologies, Inc., Ankeny, IA, USA).

Samples passing QC were genotyped using the HumanCoreExome-24 array

(Illumina®, San Diego, CA, USA). Genotyping results were analyzed using

VerifyIDintensity [83] to flag sample contamination. Sequencing libraries were

constructed using the TruSeq PCR-free DNA HT Library Preparation Kit (Illumina®,

San Diego, CA, USA) with 500ng DNA input. Briefly, genomic DNA was sheared using

a Covaris sonicator (Covaris, Woburn, MA, USA), followed by end-repair and bead-

based size selection of fragmented molecules. Selected fragments were then A-tailed

and sequence adaptors were ligated onto the fragments, followed by a final bead

purification of the libraries. Final libraries were reviewed for size distribution using

Fragment Analyzer and quantified by qPCR (Kapa Biosystems, Wilmington, MA,

USA). Libraries were sequenced on a HiSeq X system (Illumina®, San Diego, CA, USA)

with v2 chemistry, using a paired-end read length of 150 bp, to a minimum of 30x

mean genome coverage.

WGS data processing and quality control

Sequencing data were demultiplexed using bcl2fastq version 2.16.0.10 (Illumina®,

San Diego, CA, USA) and aligned to human reference hs37d5 with decoy sequences

using BWA-MEM v0.7.8 [84]. Data were further processed using the GATK best-

practices v3.2-2 pipeline [85]. Quality control procedures included marking of

duplicate reads using Picard tools v1.83 (http://picard.sourceforge.net),

realignment around indels, and base quality recalibration using 1000 Genomes

Phase 1 high confidence SNPs, HapMap v3.3, dbSNP v137, 1000 Genomes omni2.5,

1000 Genomes Phase 1 indels, and both Mills and 1000 Genomes gold standard

indels. Single-sample genotypes were called using GATK HaplotypeCaller followed

by joint genotyping of all subjects. The resulting multi-sample Variant Call Format

(VCF) file was used for variant quality score recalibration (VQSR). A 99.8% truth

sensitivity tranche level was used for SNPs and 99.0% for indel variants. SNP calls

were used to check for sample contamination using VerifyBAMId [83], and sample

identity was confirmed by requiring > 99.5% concordance with SNP array

(HumanCoreExome-24 array) genotypes.

As part of NIH's Trans-Omics for Precision Medicine (TOPMed) Program, BAM

files were submitted to the Informatics Resource Center (IRC) at the University of

Michigan. All 1,484 samples sequenced passed TOPMed's IRC quality control metrics

(mean genome coverage >30X; >95% of genome covered at >10X; and <3%

contamination).

VCF-level variants were filtered by GATK version 3.4.46 and VCFtools version

0.1.14 [86]. Variants were filtered according to the following procedures: (1)

remove variants that were not indicated as "PASS" in the VCF FILTER column, (2)

remove variants in low complexity regions [87] (downloaded from

https://github.com /lh3/varcmp/tree/master/scripts/LCR-hs37d5.bed.gz), and (3)

keep sample genotypes that have minimum read depths of 10 and genotype

qualities of 20 (DP ≥ 10 and GQ ≥ 20). The ratio of homozygous to heterozygous

variants (hom/het), ratio of transitions to transversions (Ti/Tv), and other variant

summary statistics were generated using GATK VariantEval. VCF files were

converted into PLINK format using PLINK 1.9 software [88] according to

recommended best practices [89]. Genotype consistency between WGS data and

previously published Axiom® Genome-Wide LAT 1 array (Affymetrix, Santa Clara,

CA) genotype data (dbGaP phs000920.v1.p1 and phs000921.v1.p1) was assessed

using VCFtools [86]. Individuals with percentage consistency three S.D. below the

mean (< 96.3%) were removed (N=7, **S10 Fig**). Cryptic relatedness was detected

using REAP [90]. Global ancestry and allele frequency used by REAP were estimated

using ADMIXTURE in supervised mode [91]. Related individuals (kinship coefficient

> 0.044, corresponding to a third degree relationship [92]) were excluded from

further analysis (N=36), yielding a final sample size of 1,441 for downstream

analysis. Downstream analyses were only performed on biallelic SNPs that passed

all quality filters mentioned above and had less than 10% of genotype missingness.

The 10% genotype missingness filter was applied per population instead of across

all three populations except for the rare variant analysis performed with all three

populations combined (see **Methods** section "Multi-variant analyses of combined

effects of rare variants on BDR").

Principal component analysis

Principal component analysis (PCA) was performed to control for hidden population

substructure using EIGENSTRAT's smartpca program [93]. After using PLINK 1.9 to

remove biallelic SNPs with low minor allele frequency (MAF ≤ 0.05) and in linkage

disequilibrium (R^2 > 0.5 in a 50-SNP window with a shift size of 5 SNPs), 710,256

variants remained for input into smartpca.

Local ancestry estimation

Reference genotypes for European and African ancestries were obtained from the

Axiom® Genotype Data Set [94]. SNPs with less than a 95% call rate were removed.

Since no Native American reference samples are available in the HapMap database,

reference genotypes for Native American ancestry were generated from 71 Native

American individuals previously genotyped on the Axiom® Genome-Wide LAT 1

array [27].

To call local ancestry tracts, we first created a subset of our WGS data

corresponding to sites found on the Axiom® Genome-Wide LAT 1 array, leaving

765,321 markers. Using PLINK 1.9, we merged these data with our European (CEU),

African (YRI), and Native American (NAM) reference panels, which overlapped at

434,145 markers. After filtering multi-allelic SNPs and SNPs with > 10% missing

data, we obtained a final merged dataset of 428,644 markers. We phased all samples

using SHAPEIT2 [95] and called local ancestry tracts jointly with RFMix [96] under a

three-way admixture model based on the African, European, and Native American

reference genotypes described above.

Variant annotation

TOPMed freeze 2 and 3 variants were annotated using the WGSA annotation

pipeline [97]. Annotated VCF files were downloaded from the TOPMed Data

Coordinating Center SFTP sites. dbSNP150 annotation was added separately by

using VCF file downloaded from NCBI dbSNP ftp site [98]. ENCODE (v4) annotations

were downloaded as BED files from the UCSC Table Browser (Feb.2009

[GRCh37/hg19] assembly). The conversion from GRCh37 to GRCh38 coordinates

was performed using liftOver from the UCSC Genome Browser Utilities [99].

Single locus BDR association testing on common variants

An additive logistic regression model was used to evaluate the association of

biallelic common variants (MAF > 1%) with BDR using PLINK 1.9 separately for

each population. Throughout this study, high drug responders were assigned as

cases. Logistic regression models included the covariates age, sex and body mass

index (BMI) categories to account for previously reported confounders of asthma

and BDR [100-107], and the first ten principal components (PCs) to correct for

population substructure in admixed populations. BMI and age- and sex-specific BMI

percentiles (BMI-pct) were calculated as previously described [49] and used for

assignment to BMI categories. For subjects aged 20 years and over, BMI categories

were defined as follows: underweight (BMI < 18), normal (18 ≤ BMI < 25),

overweight (25 ≤ BMI < 30) and obese (BMI ≥ 30). For subjects under 20 years of

age, BMI categories were defined as follows: underweight (BMI-pct < 5), normal (5 ≤

BMI-pct < 85), overweight (85 ≤ BMI-pct < 95) and obese (BMI-pct ≥ 95). Baseline

lung function (pre-FEV_1) has a significant impact on variation in BDR drug response.

Pre-FEV_1 was not included as a covariate in association analyses, as variation in pre-

FEV_1 was indirectly captured by several of the ten principal components already

included in association models (**S13 Table**). A correlation matrix showing the

relationship between pre-FEV_1, age, sex, BMI status, mean global ancestry, and the

top ten principal components is presented in **S13 Table**. The correlation matrix was

constructed using Spearman correlation coefficients and the accompanying

association tests for the significance of each correlation. Population-specific

genome-wide significance thresholds for the single locus analyses were calculated

based on genotypes using the autocorrelation-based *effectiveSize()* function in the R

package *'coda'* as published by Sobota *et al.* [41]. Population-specific genome-wide

significance thresholds after adjusting for the effective number of tests (adjusted

genome-wide significance) were 1.57×10^{-7} for Puerto Ricans, 2.42×10^{-7} for

Mexicans, and 9.59×10^{-8} for African Americans. Suggestive significance thresholds

were calculated as one divided by the effective number of tests [43]. Linkage

disequilibrium patterns (Genome build: hg19/1000 Genomes Nov 2014 AMR) of the

flanking regions of BDR-associated SNPs were visualized using LocusZoom [108].

Quantile-quantile (q-q) plots were generated using a uniform distribution as the

expected p-value distribution (**S11 Fig a-c**). The genomic inflation factor (λ_{GC}) was

calculated using the R package *'gap'*.

Trans-ethnic meta-analysis of common variant effects on BDR

A meta-analysis of the effects of common variants on BDR across the three

populations was performed using METASOFT [109]. We used the Han and Eskin

random effects model optimized for detecting associations under heterogeneous

genetic effects from different study conditions [109]. The number of effective tests

was estimated using the R package *'coda'* as described above, yielding an adjusted

genome-wide significance threshold of 3.53×10^{-7} and a suggestive significance

threshold of 7.06×10^{-6}. Allele frequency variation in the world population was

visualized using the Geography of Genetic Variants Browser (GGV) beta v0.92 [110].

The q-q plot and λ_{GC} were generated in the same way as described above (**S11 Fig d**,

see **Methods** section, "Single locus BDR association testing on common variants").

Calculation of variation in BDR explained by common variants

Total variation in BDR explained was estimated by calculating McFadden's pseudo

R^2 [111] in each population separately, after first pruning significantly and

suggestively associated variants (**Table 4**) for LD using the LD prune function in

PLINK 1.9 (R^2 cut-off: 0.6, window size = 50 SNPs, shift = 5 SNPs). Calculation of

pseudo R^2 was adjusted for age, sex, BMI category, and principal components 1-10.

McFadden's Pseudo R^2 is defined as:

$$R^2 = 1 - \frac{\ln L\ (M_{FULL})}{\ln L\ (M_{INTERCEPT})}$$

L: Estimated likelihood of model

M_{FULL}: Model with all predictors

$M_{INTERCEPT}$: Model with no predictors

Multi-variant analyses of combined effects of common and rare variants on

BDR

Combined effects of common and rare variants on BDR were analyzed using SKAT-O

[59]. Common and rare variants were collapsed into 1kb windows sliding across the

GRCh37 genome in steps of 500 base pairs. A total of 5.3 million windows were

analyzed and the R package *'coda'* was used to determine the number of effective

tests based on autocorrelation of the association *p*-value, as described above.

Adjusted genome-wide significance thresholds for Puerto Ricans (8.15×10^{-8}),

Mexicans (8.60×10^{-8}), African Americans (6.94×10^{-8}) and for all three populations

combined (1.53×10^{-7}) were used to identify windows of variants significantly

associated with BDR. The same covariates used for common variant association

testing were used for analyses of individual populations. For analyses of individuals

combined across all three populations, we avoided confounding from population

substructure by including local ancestry as additional covariates, defined as the

proportions of Native American and African ancestries for the window under

testing. The q-q plots and λ_{GC} were generated in the same way as described above

(**S12 Fig**, see **Methods** section, "Single locus BDR association testing on common

variants").

Variation in BDR explained by associated SKAT-O regions

Variation in BDR explained by SKAT-O regions was calculated as described above

for common variant analyses using McFadden's pseudo R^2 [111], but with one

addition: variants were weighted using a weighted kernel as described in SKAT-O

[112].

Single locus BDR association and trans-ethnic meta-analysis of array data

To address the discrepancy between our current common variant analysis results

with previously published BDR GWAS results [27], we used 1,414 of the 1,441

individuals who had both Axiom® Genome-Wide LAT 1 array (see **Methods** section,

"WGS data processing and QC") and WGS data available to rerun the single locus

BDR association testing and trans-ethnic meta-analysis. Array data were imputed to

the Haplotype Reference Consortium [113] (HRC release 1) panel using the

Michigan Imputation Server [114]. We used the top 1,000 BDR-associated SNPs to

examine the relationship between array-based and WGS-based association p-values,

genotype discordance, and imputation R^2. Correlation between the array-based and

WGS-based association p-values was determined by Spearman correlation. We also

performed single locus BDR association testing and trans-ethnic meta-analysis by

applying linear regression on 1,122 Puerto Ricans, 662 Mexicans and 1,105 African

Americans using BDR (ΔFEV_1) as a continuous trait. HRC imputed array-based data

and the same covariates as described above were used for the analysis.

Replication of top BDR-associated common variants

Replication cohorts included the Genetics of Asthma in Latino Americans Study

(GALA I) [13, 115], the Study of African Americans, Asthma, Genes & Environments

(SAGE I) [80], a case-control study of childhood asthma in Puerto Ricans (HPR)

[116], the Study of Asthma Phenotypes and Pharmacogenomic Interactions by Race-

Ethnicity (SAPPHIRE) [117] and a cohort from the Children's Hospital of

Philadelphia (CHOP) [118]. Descriptive statistics for the replication cohorts are

shown in **S6 Table**.

Logistic regression in replication analyses was performed using the same

population-specific extreme BDR cut-offs applied to the discovery analyses in the

current study (see **Methods** section, "Bronchodilator response measurements"). All

27 significantly and suggestively associated SNPs from the discovery analysis

(**Table 4**) were assessed in each replication study population. An additive genetic

model was assumed for each SNP tested. Association models were adjusted for sex,

age, BMI categories, and the first ten principal components as in the discovery

analysis. Meta-analysis across all replication studies was performed using

METASOFT as described above (**Methods**, "Trans-ethnic meta-analysis of common

variant effects on BDR").

The GALA I and SAGE I replication cohorts included 108 Puerto Ricans, 202

Mexicans and 141 African Americans with BDR measurements and complete data

for all covariates (age, sex, BMI categories and the first ten PCs). Genotype data were

imputed to the HRC panel using the Michigan Imputation Server [114]. Replication

in the HPR cohort involved 414 Puerto Rican subjects. Spirometry data were

collected as previously described [119]. Genome-wide genotyping was performed

using the Illumina HumanOmni2.5 BeadChip platform (Illumina Inc., San Diego, CA)

and processed as previously described [120]. Genotype data were phased with

SHAPE-IT [121] and imputation was performed with IMPUTE2 [122] using all

populations from 1000 Genomes Project Phase 3 as reference [75]. The SAPPHIRE

replication cohort consisted of 1,022 African Americans with asthma. Genome-wide

genotyping was performed using the Axiom® Genome-Wide AFR 1 array (Affymetrix

Inc., Santa Clara, CA) as previously described [26]. Genotype data were imputed to

the cosmopolitan 1000 Genomes Phase 1 version haplotypes using the Michigan

Imputation Server [114]. The CHOP replication cohort included 280 African

Americans. Genotyping was performed as described [118], and genotype data were

imputed to the HRC panel using the Sanger Imputation server [113].

Identification of nearest genes for BDR-associated loci

The four nearest transcripts to BDR-associated loci were identified by using the

"closest" command in BEDTools with the parameters "-d -k 4" and the RefSeq gene

annotations (Feb.2009 [GRCh37/hg19] assembly) downloaded in refFlat format

from the UCSC Table Browser [123]. Genes with multiple transcripts were reported

only once. When reporting the nearest gene, the "closest" command in BEDTools

with the parameters "-D a" was applied.

858 **Primary bronchial smooth muscle cell culture**

859 Cryopreserved primary human bronchial smooth muscle from two donors (from

860 Lonza catalog number CC-2576, lot number 0000212076 and from ATCC catalog

861 number PCS-130-011, lot number 62326179) was thawed and expanded in Lonza

862 Smooth Muscle Growth Media (SmGM; catalog number CC-3182) on T75 flasks (E&K

863 Scientific Products, catalog number 658175).

864 **H3K27ac ChIP-seq assay**

865 Upon reaching 80% confluency, BSMCs were serum-starved by replacing SmGM

866 with smooth muscle basal media (SmBM) for 24 hours. BSMCs were then grown in

867 SmBM containing 5% FBS for 4 hours, then fixed in 1% formaldehyde for 10 min

868 and quenched with 0.125 M glycine for 5 minutes. Cells were removed from the T75

869 flasks by scraping in cold PBS containing sodium butyrate (20 mM, Diagenode,

870 catalog number C12020010). Chromatin sheering was carried out using a Covaris S2

871 sonicator. Sheared chromatin was used for immunoprecipitation with antibodies

872 against active chromatin marks (H3K27ac; Abcam, ab4729) using the Diagenode

873 LowCell# ChIP kit (CAT#C01010072), following the manufacturer's protocol.

874 Libraries were prepared using the Rubicon DNA-Seq kit (CAT#R400406) following

875 the manufacturer's protocol and sequenced on an Illumina HiSeq 4000 using single-

876 end 50-bp reads to a sequencing depth of at least 25 million reads (submitted under

877 BioProject PRJNA369271). Uniquely mapping raw reads were aligned using Bowtie

878 [124] under default settings. Peak regions for each individual were called using

MACS2 [125, 126] and reproducible peaks identified using the ENCODE IDR pipeline

[127].

Diverse Convergent Evidence approach for variant prioritization

The Diverse Convergent Evidence (DiCE) approach is a logical, heuristic framework

for integrating multiple types of observational, bioinformatics, and laboratory

evidence to prioritize variants discovered from high throughput genetic studies for

further evaluation in functional experiments [53]. Results from the trans-ethnic

meta-analysis and from the replication analyses were considered observational

data. Laboratory evidence was provided by the identified peaks in our ChIP-seq

analyses performed in BSMCs. Informatic evidence was compiled using Ensembl,

PubMed, the NHGRI-EBI GWAS Catalog, and ENCODE to identify previously reported

associations with BDR or asthma-related phenotypes, and predicted biological

functions associated with assessed loci. After compiling the observational,

informatic and laboratory evidence for each suggestively or significantly associated

variant, DiCE constructs an evidence matrix to estimate the strength of the

information supporting each association. DiCE scores > 6 were considered strong

evidence that a given locus was involved in the pathophysiology of BDR, and

variants with the highest DiCE score, after meeting this criterion, were prioritized

for downstream functional analyses.

Luciferase assays

NFKB1 candidate enhancer sequences were amplified from human genomic DNA

(Roche) using oligonucleotides designed in Primer3 with 18 and 20 bp overhangs

for forward and reverse primers, respectively (5⬛-GGCCTAACTGGCCGGTAC-3⬛ and

5⬛-CGCCGAGGCCAGATCTTGAT-3⬛), complementary to sequences flanking the KpnI

and EcoRV sites in the pGL4.23 Gate A vector (Promega, Madison, WI, USA) using

Phusion High-Fidelity PCR kit (NEB, catalog number M0531S). PCR primers were

designed around the edges of these ChIP-seq peaks and the most complete fragment

that was successfully amplified was used for luciferase assays. PCR products where

then cloned into the pGL4.23 vector using the Gibson Assembly method (NEB,

catalog number E2611S). Smooth muscle cells were plated at 50-70% confluency in

24-well cell culture plates (Falcon, catalog number 353047) and grown to 80%

confluency in SmGM. Transfections were carried out by combining

polyethyleneimine (PEI) with DNA vectors at a 1:1 ratio by weight in opti-MEM (Life

Technologies, catalog number 31985070). The transfection mixture consisted of

225 ng of enhancer assay vectors and 25 ng of pGL4.24 (Renilla transfection

efficiency control) with 250 ng of PEI in 50 μL of opti-MEM. After 15 minutes, 500

μL of SmBM was added to the transfection mixture and the combination added to

cell culture. Cells were incubated for 4 hours in SmBM plus the transfection mixture,

then media was replaced with SmGM for 24 hours. Cells were then washed with PBS

and enhancer assay cells were lysed with 100 μL of Passive Lysis Buffer (Promega,

Madison, WI, USA). Reporter activity was measured using the Dual-Luciferase

Reporter Assay System (Promega, Madison, WI, USA) and measured on the Glomax

96 well plate luminometer (Promega, Madison, WI, USA). The luciferase assay was

carried out in two separate experiments, with three independent replicates per

experiment (three wells of cells were transfected per construct per experiment).

Each well was then split into two technical replicates for luciferase activity

measurements with a luminometer.

RNA extraction, library preparation and sequencing

Among the African American subjects with WGS data in our study, 39 samples were

selected for RNA-seq based on BDR status and the number of copies of low-BDR

associated alleles at rs28450894. The number of samples in each category is shown

in **S14 Table**. Peripheral blood samples were collected into PAXgene Blood RNA

tubes (PreAnalytiX, Hombrechtikon, Switzerland). Total RNA was extracted from

PAXgene Blood RNA tubes using the MagMAX™ for Stabilized Blood Tubes RNA

Isolation Kit (CAT#4451894, Thermo Fisher Scientific, Waltham, MA, USA)

according to manufacturer's protocols. RNA integrity and yield were assessed using

an Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA). Globin

depletion was performed using GLOBINclear™ kit (CAT#AM1980, Thermo Fisher

Scientific, Waltham, MA, USA). Library preparation and ribosomal depletion were

performed using KAPA Stranded RNA-seq Kit with RiboErase (CAT#KK8483, Kapa

Biosystems, Wilmington, MA, USA) according to the manufacturer's protocols. Each

sample was uniquely barcoded with NEXTflex™ DNA Barcodes (CAT#514104, Bioo

Scientific®, Austin, TX, USA). Barcoded libraries were pooled and sequenced on 4

lanes on a HiSeq 4000 sequencing system (Illumina®, San Diego, CA, USA) with a

paired-end read length of 100 bp at the University of California, San Francisco's

Center for Advanced Technology.

RNA-seq data processing and analysis

Raw sequencing reads were aligned to the human reference genome (hg19) using

STAR (v2.4.2a) [128]. Gene read counts were obtained from uniquely mapped reads

based on Ensembl annotation (v75) [129]. DESeq2 [130] was used to analyze read

counts for differential gene expression changes between genotypes, including an

interaction term with genotype and sex (genotype * sex). We used a linear model to

account for sex, age and library prep batch, and a custom model matrix to correct for

GC content difference between genes. After normalization for sequencing depth and

GC percentage, genes with fewer than an average of five normalized read counts per

sample and fewer than 20 samples with at least one read count were removed. This

filtering process kept 19,592 Ensembl genes for analysis. Fold change, raw and FDR-

adjusted p-value for the genotype term was reported. Genes were then further

filtered to analyze the locus surrounding rs28450894 for differential gene

expression by including all genes with a transcriptional start site within 1Mbp of

rs28450894. P-values were then corrected using the false discovery rate method to

account for the 13 genes in this locus. Significant level FDR-adjusted p-value is ≤ 0.1.

ACKNOWLEDGEMENTS

Whole genome sequencing (WGS) for the Trans-Omics in Precision Medicine

(TOPMed) program was supported by the National Heart, Lung, and Blood Institute

(NHLBI). WGS for "NHLBI TOPMed: Genes-environments & Admixture in Latino

Americans (GALA II) Study" (phs000920) and "NHLBI TOPMed: Study of African

Americans, Asthma, Genes and Environments (SAGE II)" (phs000921) was

performed at the New York Genome Center (3R01HL117004-01S3). We

acknowledge New York Genome Center investigators and teams for whole genome

sequencing sample preparation, quality control, data generation, data processing

and initial joint genotyping. Centralized read mapping and genotype calling, along

with variant quality metrics and filtering were provided by the TOPMed Informatics

Research Center (3R01HL-117626-02S1). Phenotype harmonization, data

management, sample-identity QC, and general study coordination were provided by

the TOPMed Data Coordinating Center (3R01HL-120393-02S1). We gratefully

acknowledge the studies and participants who provided biological samples and data

for TOPMed. We also gratefully acknowledge the contributions of the investigators

of the NHLBI TOPMed Consortium (https://www.nhlbiwgs.org/topmed-banner-

authorship). C.A.W. would like to declare that the content of this publication does not

necessarily reflect the views or policies of the Department of Health and Human

Services, nor does mention of trade names, commercial products, or organizations imply

endorsement by the U.S. Government.

REFERENCES

1. Lara Akinbami. Centers for Disease Control and Prevention. (2015). Asthma

Prevalence, Health Care Use and Mortality: United States, 2003-05. [online]

Available at: http://www.cdc.gov/nchs/data/hestat/asthma03-05/asthma03-

05.htm [Accessed 9/12 2017].

2. Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, et al. (2012) Years lived

with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-

2010: A systematic analysis for the global burden of disease study 2010. Lancet

380(9859): 2163-2196.

3. World   Health   Organization.   (2017).   Asthma.   [online]   Available   at:

http://www.who.int/mediacentre/factsheets/fs307/en/   [Accessed   9/12

2017].

4. World Health Organization. (2007). Global surveillance, prevention and control

of chronic respiratory diseases: a comprehensive approach. [online] Available

at: http://www.who.int/gard/publications/GARD_Manual/en/ [Accessed 9/12

2017].

5. Oh SS, White MJ, Gignoux CR, Burchard EG. (2016) Making precision medicine

socially precise. take a deep breath. Am J Respir Crit Care Med 193(4): 348-350.

6. Burchard EG. (2014) Medical research: Missing patients. Nature 513(7518):

301-302.

7.   Barr RG, Aviles-Santa L, Davis SM, Aldrich TK, Gonzalez F, II, et al. (2016)

Pulmonary disease and age at immigration among hispanics. Results from the

Hispanic Community Health Study/Study of Latinos. Am J Respir Crit Care Med

193(4): 386-395.

8.   Palmer LJ, Silverman ES, Weiss ST, Drazen JM. (2002) Pharmacogenetics of

asthma. Am J Respir Crit Care Med 165(7): 861-866.

9.   Nelson HS. (1995) Beta-adrenergic bronchodilators. N Engl J Med 333(8): 499-

506.

10.   Eggleston PA, Malveaux FJ, Butz AM, Huss K, Thompson L, et al. (1998)

Medications used by children with asthma living in the inner city. Pediatrics

101(3 Pt 1): 349-354.

11.   Finkelstein JA, Lozano P, Farber HJ, Miroshnik I, Lieu TA. (2002) Underuse of

controller medications among medicaid-insured children with asthma. Arch

Pediatr Adolesc Med 156(6): 562-567.

12.   Drazen JM, Silverman EK, Lee TH. (2000) Heterogeneity of therapeutic

responses in asthma. Br Med Bull 56(4): 1054-1070.

13.   Burchard EG, Avila PC, Nazario S, Casal J, Torres A, et al. (2004) Lower

bronchodilator responsiveness in puerto rican than in mexican subjects with

asthma. Am J Respir Crit Care Med 169(3): 386-392.

14.   Naqvi M, Thyne S, Choudhry S, Tsai HJ, Navarro D, et al. (2007) Ethnic-specific

differences in bronchodilator responsiveness among african americans, puerto

ricans, and mexicans with asthma. J Asthma 44(8): 639-648.

15.   Wechsler ME, Castro M, Lehman E, Chinchilli VM, Sutherland ER, et al. (2011)

Impact of race on asthma treatment failures in the asthma clinical research

network. Am J Respir Crit Care Med 184(11): 1247-1253.

16.   Martinez FD. (2005) Safety of long-acting beta-agonists--an urgent need to

clear the air. N Engl J Med 353(25): 2637-2639.

17.   Dixon AE. (2011) Long-acting beta-agonists and asthma: The saga continues.

Am J Respir Crit Care Med 184(11): 1220-1221.

18.   Nelson HS, Weiss ST, Bleecker ER, Yancey SW, Dorinsky PM, et al. (2006) The

salmeterol   multicenter   asthma   research   trial: A   comparison   of   usual

pharmacotherapy for asthma or usual pharmacotherapy plus salmeterol. Chest

129(1): 15-26.

19.   Kramer JM. (2009) Balancing the benefits and risks of inhaled long-acting

beta-agonists--the influence of values. N Engl J Med 360(16): 1592-1595.

20.   McGeachie MJ, Stahl EA, Himes BE, Pendergrass SA, Lima JJ, et al. (2013)

Polygenic   heritability   estimates   in   pharmacogenetics: Focus   on   asthma   and

related phenotypes. Pharmacogenet Genomics 23(6): 324-328.

21.   Nieminen MM, Kaprio J, Koskenvuo M. (1991) A population-based study of

1041   bronchial asthma in adult twin pairs. Chest 100(1): 70-75.

1042   22.   Fagnani C, Annesi-Maesano I, Brescianini S, D'Ippolito C, Medda E, et al. (2008)

1043   Heritability and shared genetic effects of asthma and hay fever: An italian study

1044   of young twins. Twin Res Hum Genet 11(2): 121-131.

1045   23.   Himes BE, Jiang X, Hu R, Wu AC, Lasky-Su JA, et al. (2012) Genome-wide

1046   association analysis in asthma subjects identifies SPATS2L as a novel

1047   bronchodilator response gene. PLoS Genet 8(7): e1002824.

1048   24.   Duan QL, Lasky-Su J, Himes BE, Qiu W, Litonjua AA, et al. (2014) A genome-

1049   wide association study of bronchodilator response in asthmatics.

1050   Pharmacogenomics J 14(1): 41-47.

1051   25.   Israel E, Lasky-Su J, Markezich A, Damask A, Szefler SJ, et al. (2015) Genome-

1052   wide association study of short-acting beta2-agonists. A novel genome-wide

1053   significant locus on chromosome 2 near ASB3. Am J Respir Crit Care Med

1054   191(5): 530-537.

1055   26.   Padhukasahasram B, Yang JJ, Levin AM, Yang M, Burchard EG, et al. (2014)

1056   Gene-based association identifies SPATA13-AS1 as a pharmacogenomic

1057   predictor of inhaled short-acting beta-agonist response in multiple population

1058   groups. Pharmacogenomics J 14(4): 365-371.

27.   Drake KA, Torgerson DG, Gignoux CR, Galanter JM, Roth LA, et al. (2014) A

genome-wide association study of bronchodilator response in latinos implicates

rare variants. J Allergy Clin Immunol 133(2): 370-378.

28.   Conrad DF, Jakobsson M, Coop G, Wen X, Wall JD, et al. (2006) A worldwide

survey of haplotype variation and linkage disequilibrium in the human genome.

Nat Genet 38(11): 1251-1260.

29.   Hoffmann TJ, Zhan Y, Kvale MN, Hesselson SE, Gollub J, et al. (2011) Design

and coverage of high throughput genotyping arrays optimized for individuals of

east asian, african american, and latino race/ethnicity using imputation and a

novel hybrid SNP selection algorithm. Genomics 98(6): 422-430.

30.   Illumina. (2016). Infinium® Multi-Ethnic Global BeadChip. [online] Available

at: https://www.illumina.com/content/dam/illumina-

marketing/documents/products/datasheets/multi-ethnic-global-data-sheet-

370-2016-001.pdf [Accessed 9/12 2017].

31.   Zheng HF, Rong JJ, Liu M, Han F, Zhang XW, et al. (2015) Performance of

genotype imputation for low frequency and rare variants from the 1000

genomes. PLoS One 10(1): e0116487.

32.   Huang J, Howie B, McCarthy S, Memari Y, Walter K, et al. (2015) Improved

imputation of low-frequency and rare variants using the UK10K haplotype

reference panel. Nat Commun 6: 8111.

33.   Zhang F, Lupski JR. (2015) Non-coding genetic variants in human disease. Hum
1080   Mol Genet 24(R1): R102-10.

1081   34.   Bustamante CD, Burchard EG, De la Vega FM. (2011) Genomics for the world.
1082   Nature 475(7355): 163-165.

1083   35.   Popejoy AB, Fullerton SM. (2016) Genomics is failing on diversity. Nature
1084   538(7624): 161-164.

1085   36.   Oh SS, Galanter J, Thakur N, Pino-Yanes M, Barcelo NE, et al. (2015) Diversity
1086   in clinical and biomedical research: A promise yet to be fulfilled. PLoS Med
1087   12(12): e1001918.

1088   37.   Hankinson JL, Odencrantz JR, Fedan KB. (1999) Spirometric reference values
1089   from a sample of the general U.S. population. Am J Respir Crit Care Med 159(1):
1090   179-187.

1091   38.   Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, et al. (2014) A general
1092   framework for estimating the relative pathogenicity of human genetic variants.
1093   Nat Genet 46(3): 310-315.

1094   39.   Gulko B, Hubisz MJ, Gronau I, Siepel A. (2015) A method for calculating
1095   probabilities of fitness consequences for point mutations across the human
1096   genome. Nat Genet 47(3): 276-283.

40.   Altshuler D, Daly MJ, Lander ES. (2008) Genetic mapping in human disease.

Science 322(5903): 881-888.

41.   Sobota RS, Shriner D, Kodaman N, Goodloe R, Zheng W, et al. (2015)

Addressing population-specific multiple testing burdens in genetic association

studies. Ann Hum Genet 79(2): 136-147.

42.   Pe'er I, Yelensky R, Altshuler D, Daly MJ. (2008) Estimation of the multiple

testing burden for genomewide association studies of nearly all common

variants. Genet Epidemiol 32(4): 381-385.

43.   Duggal P, Gillanders EM, Holmes TN, Bailey-Wilson JE. (2008) Establishing an

adjusted p-value threshold to control the family-wide type 1 error in genome

wide association studies. BMC Genomics 9: 516-2164-9-516.

44.   Wang H, Thomas DC, Pe'er I, Stram DO. (2006) Optimal two-stage genotyping

designs for genome-wide association scans. Genet Epidemiol 30(4): 356-368.

45.   Skol AD, Scott LJ, Abecasis GR, Boehnke M. (2007) Optimal designs for two-

stage genome-wide association studies. Genet Epidemiol 31(7): 776-788.

46.   Reed E, Nunez S, Kulp D, Qian J, Reilly MP, et al. (2015) A guide to genome-

wide association analysis and post-analytic interrogation. Stat Med 34(28):

3769-3792.

47.   Zaykin DV, Zhivotovsky LA. (2005) Ranks of genuine associations in whole-

genome scans. Genetics 171(2): 813-823.

48.   Castro-Giner F, de Cid R, Gonzalez JR, Jarvis D, Heinrich J, et al. (2010)

Positionally cloned genes and age-specific effects in asthma and atopy: An

international population-based cohort study (ECRHS). Thorax 65(2): 124-131.

49.   White MJ, Risse-Adams O, Goddard P, Contreras MG, Adams J, et al. (2016)

Novel genetic risk factors for asthma in african american children: Precision

medicine and the SAGE II study. Immunogenetics 68(6-7): 391-400.

50.   Johnson M. (1998) The beta-adrenoceptor. Am J Respir Crit Care Med 158(5 Pt

3): S146-53.

51.   Nuzzo R. (2014) Scientific method: Statistical errors. Nature 506(7487): 150-

152.

52.   Malley JD, Dasgupta A, Moore JH. (2013) The limits of p-values for biological

data mining. BioData Min 6(1): 10-0381-6-10.

53.   Ciesielski TH, Pendergrass SA, White MJ, Kodaman N, Sobota RS, et al. (2014)

Diverse convergent evidence in the genetic analysis of complex disease:

Coordinating omic, informatic, and experimental evidence to better identify and

validate risk factors. BioData Min 7: 10-0381-7-10. eCollection 2014.

54.   Cumberland PM, Czanner G, Bunce C, Dore CJ, Freemantle N, et al. (2014)

Ophthalmic statistics note: The perils of dichotomising continuous variables. Br

J Ophthalmol 98(6): 841-843.

55.   Peloso GM, Rader DJ, Gabriel S, Kathiresan S, Daly MJ, et al. (2016) Phenotypic

extremes in rare variant study designs. Eur J Hum Genet 24(6): 924-930.

56.   Li D, Lewinger JP, Gauderman WJ, Murcray CE, Conti D. (2011) Using extreme

phenotype sampling to identify the rare causal variants of quantitative traits in

association studies. Genet Epidemiol 35(8): 790-799.

57.   Guey LT, Kravic J, Melander O, Burtt NP, Laramie JM, et al. (2011) Power in the

phenotypic extremes: A simulation study of power in discovery and replication

of rare variants. Genet Epidemiol 35(4): 236-246.

58.   Lamina C. (2011) Digging into the extremes: A useful approach for the analysis

of rare variants with continuous traits? BMC Proc 5 Suppl 9: S105-6561-5-S9-

S105.

59.   Lee S, Wu MC, Lin X. (2012) Optimal tests for rare variant effects in sequencing

association studies. Biostatistics 13(4): 762-775.

60.   Lee JH, McDonald ML, Cho MH, Wan ES, Castaldi PJ, et al. (2014) DNAH5 is

associated with total lung capacity in chronic obstructive pulmonary disease.

Respir Res 15: 97-014-0097-y.

61.   Ortiz RA, Barnes KC. (2015) Genetics of allergic diseases. Immunol Allergy Clin

North Am 35(1): 19-44.

62.   Ramasamy A, Curjuric I, Coin LJ, Kumar A, McArdle WL, et al. (2011) A

genome-wide meta-analysis of genetic variants associated with allergic rhinitis

and grass sensitization and their interaction with birth order. J Allergy Clin

Immunol 128(5): 996-1005.

63.   Naqvi M, Choudhry S, Tsai HJ, Thyne S, Navarro D, et al. (2007) Association

between IgE levels and asthma severity among african american, mexican, and

puerto rican patients with asthma. J Allergy Clin Immunol 120(1): 137-143.

64.   Edwards MR, Bartlett NW, Clarke D, Birrell M, Belvisi M, et al. (2009) Targeting

the NF-kappaB pathway in asthma and chronic obstructive pulmonary disease.

Pharmacol Ther 121(1): 1-13.

65.   Zhen G, Park SW, Nguyenvu LT, Rodriguez MW, Barbeau R, et al. (2007) IL-13

and epidermal growth factor receptor have critical but distinct roles in

epithelial cell mucin production. Am J Respir Cell Mol Biol 36(2): 244-253.

66.   Murgia C, Grosser D, Truong-Tran AQ, Roscioli E, Michalczyk A, et al. (2011)

Apical localization of zinc transporter ZnT4 in human airway epithelial cells and

its loss in a murine model of allergic airway inflammation. Nutrients 3(11): 910-

928.

67.   Besecker B, Bao S, Bohacova B, Papp A, Sadee W, et al. (2008) The human zinc

transporter SLC39A8 (Zip8) is critical in zinc-mediated cytoprotection in lung

epithelia. Am J Physiol Lung Cell Mol Physiol 294(6): L1127-36.

68.   Persson H, Kwon AT, Ramilowski JA, Silberberg G, Soderhall C, et al. (2015)

Transcriptome analysis of controlled and therapy-resistant childhood asthma

reveals distinct gene expression profiles. J Allergy Clin Immunol 136(3): 638-

648.

69.   Lin YJ, Chang JS, Liu X, Tsang H, Chien WK, et al. (2015) Genetic variants in

PLCB4/PLCB1 as susceptibility loci for coronary artery aneurysm formation in

kawasaki disease in han chinese in taiwan. Sci Rep 5: 14762.

70.   Hersch M, Peter B, Kang HM, Schupfer F, Abriel H, et al. (2012) Mapping

genetic variants associated with beta-adrenergic responses in inbred mice.

PLoS One 7(7): e41032.

71.   Dougherty RH, Sidhu SS, Raman K, Solon M, Solberg OD, et al. (2010)

Accumulation of intraepithelial mast cells with a unique protease phenotype in

T(H)2-high asthma. J Allergy Clin Immunol 125(5): 1046-1053.e8.

72.   Woodruff PG, Boushey HA, Dolganov GM, Barker CS, Yang YH, et al. (2007)

Genome-wide profiling identifies epithelial cell genes associated with asthma

and with treatment response to corticosteroids. Proc Natl Acad Sci U S A

104(40): 15858-15863.

73.   Motsinger-Reif AA, Jorgenson E, Relling MV, Kroetz DL, Weinshilboum R, et al.

1192   (2013) Genome-wide association studies in pharmacogenomics: Successes and

1193   lessons. Pharmacogenet Genomics 23(8): 383-394.

1194   74.   Tishkoff SA, Reed FA, Friedlaender FR, Ehret C, Ranciaro A, et al. (2009) The

1195   genetic structure and history of africans and african americans. Science

1196   324(5930): 1035-1044.

1197   75.   1000 Genomes Project Consortium, Auton A, Brooks LD, Durbin RM, Garrison

1198   EP, et al. (2015) A global reference for human genetic variation. Nature

1199   526(7571): 68-74.

1200   76.   Zou JY, Park DS, Burchard EG, Torgerson DG, Pino-Yanes M, et al. (2015)

1201   Genetic and socioeconomic study of mate choice in latinos reveals novel

1202   assortment patterns. Proc Natl Acad Sci U S A 112(44): 13621-13626.

1203   77.   Hall SS. (2013) Genetics: A gene of rare effect. Nature 496(7444): 152-155.

1204   78.   Burchard EG, Oh SS, Foreman MG, Celedon JC. (2015) Moving toward true

1205   inclusion of racial/ethnic minorities in federally funded studies. A key step for

1206   achieving respiratory health equality in the united states. Am J Respir Crit Care

1207   Med 191(5): 514-521.

1208   79.   Borrell LN, Nguyen EA, Roth LA, Oh SS, Tcheurekdjian H, et al. (2013)

1209   Childhood obesity and asthma control in the GALA II and SAGE II studies. Am J

1210   Respir Crit Care Med 187(7): 697-702.

80.   Nishimura KK, Galanter JM, Roth LA, Oh SS, Thakur N, et al. (2013) Early-life

air pollution and asthma risk in minority children. the GALA II and SAGE II

studies. Am J Respir Crit Care Med 188(3): 309-318.

81.   Thakur N, Oh SS, Nguyen EA, Martin M, Roth LA, et al. (2013) Socioeconomic

status and childhood asthma in urban minority youths. the GALA II and SAGE II

studies. Am J Respir Crit Care Med 188(10): 1202-1209.

82.   Pellegrino R, Viegi G, Brusasco V, Crapo RO, Burgos F, et al. (2005)

Interpretative strategies for lung function tests. Eur Respir J 26(5): 948-968.

83.   Jun G, Flickinger M, Hetrick KN, Romm JM, Doheny KF, et al. (2012) Detecting

and estimating contamination of human DNA samples in sequencing and array-

based genotype data. Am J Hum Genet 91(5): 839-848.

84.   Li H, Durbin R. (2009) Fast and accurate short read alignment with burrows-

wheeler transform. Bioinformatics 25(14): 1754-1760.

85.   DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, et al. (2011) A

framework for variation discovery and genotyping using next-generation DNA

sequencing data. Nat Genet 43(5): 491-498.

86.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, et al. (2011) The variant

call format and VCFtools. Bioinformatics 27(15): 2156-2158.

87.   Li H. (2014) Toward better understanding of artifacts in variant calling from

high-coverage samples. Bioinformatics 30(20): 2843-2851.

88.   Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, et al. (2015) Second-

generation PLINK: Rising to the challenge of larger and richer datasets.

Gigascience 4: 7-015-0047-8. eCollection 2015.

89.   Genetics For Fun. Best practice for converting VCF files to plink format.

[online]   Available   at: http://apol1.blogspot.nl/2014/11/best-practice-for-

converting-vcf-files.html.

90.   Thornton T, Tang H, Hoffmann TJ, Ochs-Balcom HM, Caan BJ, et al. (2012)

Estimating kinship in admixed populations. Am J Hum Genet 91(1): 122-138.

91.   Alexander DH, Novembre J, Lange K. (2009) Fast model-based estimation of

ancestry in unrelated individuals. Genome Res 19(9): 1655-1664.

92.   Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, et al. (2010) Robust

relationship inference in genome-wide association studies. Bioinformatics

26(22): 2867-2873.

93.   Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, et al. (2006)

Principal components analysis corrects for stratification in genome-wide

association studies. Nat Genet 38(8): 904-909.

94.   Affymetrix. (2014). Axiom genotype data set. [online] Available at:

1248   http://www.affymetrix.com/support/technical/

1249   sample_data/axiom_db/axiomdb_data.affx [Accessed 4/1 2014].

1250   95.   Delaneau O, Zagury JF, Marchini J. (2013) Improved whole-chromosome

1251   phasing for disease and population genetic studies. Nat Methods 10(1): 5-6.

1252   96.   Maples BK, Gravel S, Kenny EE, Bustamante CD. (2013) RFMix: A

1253   discriminative modeling approach for rapid and robust local-ancestry inference.

1254   Am J Hum Genet 93(2): 278-288.

1255   97.   Liu X, White S, Peng B, Johnson AD, Brody JA, et al. (2016) WGSA: An

1256   annotation pipeline for human genome sequencing studies. J Med Genet 53(2):

1257   111-112.

1258   98.   Sherry ST, Ward MH, Kholodov M, Baker J, Phan L, et al. (2001) dbSNP: The

1259   NCBI database of genetic variation. Nucleic Acids Res 29(1): 308-311.

1260   99.   Kuhn RM, Haussler D, Kent WJ. (2013) The UCSC genome browser and

1261   associated tools. Brief Bioinform 14(2): 144-161.

1262   100.   Yao TC, Ou LS, Yeh KW, Lee WI, Chen LC, et al. (2011) Associations of age,

1263   gender, and BMI with prevalence of allergic diseases in children: PATCH study. J

1264   Asthma 48(5): 503-510.

101.　Nicolai T, Pereszlenyiova-Bliznakova L, Illi S, Reinhardt D, von Mutius E.

(2003) Longitudinal follow-up of the changing gender ratio in asthma from

childhood to adulthood: Role of delayed manifestation in girls. Pediatr Allergy

Immunol 14(4): 280-283.

102.　Joseph M, Elliott M, Zelicoff A, Qian Z, Trevathan E, et al. (2016) Racial

disparity in the association between body mass index and self-reported asthma

in children: A population-based study. J Asthma 53(5): 492-497.

103.　Dixon AE, Shade DM, Cohen RI, Skloot GS, Holbrook JT, et al. (2006) Effect of

obesity on clinical presentation and response to treatment in asthma. J Asthma

43(7): 553-558.

104.　Ullah MI, Newman GB, Saunders KB. (1981) Influence of age on response to

ipratropium and salbutamol in asthma. Thorax 36(7): 523-529.

105.　Mohamed MH, Lima JJ, Eberle LV, Self TH, Johnson JA. (1999) Effects of

gender and race on albuterol pharmacokinetics. Pharmacotherapy 19(2): 157-

161.

106.　Carroll CL, Bhandari A, Zucker AR, Schramm CM. (2006) Childhood obesity

increases duration of therapy during severe asthma exacerbations. Pediatr Crit

Care Med 7(6): 527-531.

107.   Forno E, Lescher R, Strunk R, Weiss S, Fuhlbrigge A, et al. (2011) Decreased

1284   response to inhaled steroids in overweight and obese asthmatic children. J

1285   Allergy Clin Immunol 127(3): 741-749.

1286   108.   Pruim RJ, Welch RP, Sanna S, Teslovich TM, Chines PS, et al. (2010)

1287   LocusZoom: Regional visualization of genome-wide association scan results.

1288   Bioinformatics 26(18): 2336-2337.

1289   109.   Han B, Eskin E. (2011) Random-effects model aimed at discovering

1290   associations in meta-analysis of genome-wide association studies. Am J Hum

1291   Genet 88(5): 586-598.

1292   110.   Marcus JH, Novembre J. (2017) Visualizing the geography of genetic variants.

1293   Bioinformatics 33(4): 594-595.

1294   111.   McFadden D. (1979) Quantitative methods for analysing travel behavior of

1295   individuals: Some recent developments. In: Hensher DA, Stopher PR, editors.

1296   Behavioural travel modelling. London: Croom Helm. pp. 279-318.

1297   112.   Lee S, Emond MJ, Bamshad MJ, Barnes KC, Rieder MJ, et al. (2012) Optimal

1298   unified approach for rare-variant association testing with application to small-

1299   sample case-control whole-exome sequencing studies. Am J Hum Genet 91(2):

1300   224-237.

113.   McCarthy S, Das S, Kretzschmar W, Delaneau O, Wood AR, et al. (2016) A

reference panel of 64,976 haplotypes for genotype imputation. Nat Genet

48(10): 1279-1283.

114.   Das S, Forer L, Schonherr S, Sidore C, Locke AE, et al. (2016) Next-generation

genotype imputation service and methods. Nat Genet 48(10): 1284-1287.

115.   Torgerson DG, Gignoux CR, Galanter JM, Drake KA, Roth LA, et al. (2012)

Case-control admixture mapping in latino populations enriches for known

asthma-associated genes. J Allergy Clin Immunol 130(1): 76-82.e12.

116.   Han YY, Forno E, Brehm JM, Acosta-Perez E, Alvarez M, et al. (2015) Diet,

interleukin-17, and childhood asthma in puerto ricans. Ann Allergy Asthma

Immunol 115(4): 288-293.e1.

117.   Levin AM, Wang Y, Wells KE, Padhukasahasram B, Yang JJ, et al. (2014)

Nocturnal asthma and the importance of race/ethnicity and genetic ancestry.

Am J Respir Crit Care Med 190(3): 266-273.

118.   Ong BA, Li J, McDonough JM, Wei Z, Kim C, et al. (2013) Gene network

analysis in a pediatric cohort identifies novel lung function genes. PLoS One

8(9): e72899.

119.   Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, et al. (2010)

Standardisation of lung function testing: The authors' replies to readers'

comments. Eur Respir J 36(6): 1496-1498.

120.   Chen W, Brehm JM, Lin J, Wang T, Forno E, et al. (2015) Expression
quantitative trait loci (eQTL) mapping in puerto rican children. PLoS One 10(3):
e0122464.

121.   Delaneau O, Marchini J, Zagury JF. (2011) A linear complexity phasing
method for thousands of genomes. Nat Methods 9(2): 179-181.

122.   Howie BN, Donnelly P, Marchini J. (2009) A flexible and accurate genotype
imputation method for the next generation of genome-wide association studies.
PLoS Genet 5(6): e1000529.

123.   Karolchik D, Hinrichs AS, Furey TS, Roskin KM, Sugnet CW, et al. (2004) The
UCSC table browser data retrieval tool. Nucleic Acids Res 32(Database issue):
D493-6.

124.   Langmead B, Trapnell C, Pop M, Salzberg SL. (2009) Ultrafast and memory-
efficient alignment of short DNA sequences to the human genome. Genome Biol
10(3): R25-2009-10-3-r25. Epub 2009 Mar 4.

125.   Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, et al. (2008) Model-based
analysis of ChIP-seq (MACS). Genome Biol 9(9): R137-2008-9-9-r137. Epub
2008 Sep 17.

126.   Feng J, Liu T, Qin B, Zhang Y, Liu XS. (2012) Identifying ChIP-seq enrichment
using MACS. Nat Protoc 7(9): 1728-1740.

127.   Landt SG, Marinov GK, Kundaje A, Kheradpour P, Pauli F, et al. (2012) ChIP-

seq guidelines and practices of the ENCODE and modENCODE consortia.

Genome Res 22(9): 1813-1831.

128.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, et al. (2013) STAR:

Ultrafast universal RNA-seq aligner. Bioinformatics 29(1): 15-21.

129.   Flicek P, Amode MR, Barrell D, Beal K, Billis K, et al. (2014) Ensembl 2014.

Nucleic Acids Res 42(Database issue): D749-55.

130.   Love MI, Huber W, Anders S. (2014) Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol 15(12): 550.

1349

1350

1351 **FIGURE CAPTIONS**

1352 **Fig 1.** An overview of the main analyses performed in the current study. More

1353 detailed descriptions of the discovery and replication cohort demographics and

1354 analyses performed for common and rare variant analysis can be found in Methods.

1355

1356 **Fig 2.** (**a**) Number of variants per sample. The bin size is 0.025M variants. (**b**) Allele

1357 frequency of biallelic SNPs (relative to GRCh37). (**c**) Allele frequency of population-

1358 specific biallelic SNPs. (d) Novel biallelic SNPs based on dbSNP build150.

1359

Fig 3. (**a**) Manhattan plot of the trans-ethnic meta-analysis of single locus BDR

association testing. Top ten BDR-associated loci are circled. Black line indicates

universal genome-wide significance threshold (5.00×10^{-8}), red line indicates

adjusted genome-wide significance threshold (3.53×10^{-7}), and blue line indicates

suggestive significance threshold (7.06×10^{-6}). (**b**) Forest plot of the population-

specific and joint effect of the two most significantly associated SNPs, rs17834628

and rs35661809. The R^2 between these two SNPs is 0.93 in Puerto Ricans, 0.96 in

Mexicans and 0.66 in African Americans. (**c**) The most significantly associated SNP

(rs17834628) is plotted together with 400kb flanking regions on either side. Color

of the dots shows the LD of each SNP with rs17834628 based on the 1000 Genomes

Nov 2014 AMR population. Multiple SNPs in high LD ($R^2 > 0.8$, red) reached a

suggestive significance level.

1372

1373   **Fig 4.** Manhattan plot of SKAT-O analysis of biallelic common and rare SNPs

1374   grouped by 1kb windows sliding across chromosome 1 to 22 in **(a)** Puerto Ricans

1375   **(b)** African Americans, **(c)** Mexicans, and **(d)** all populations combined. Bonferroni-

1376   corrected genome-wide and suggestive significance levels are marked by red and

1377   blue lines, respectively.

1378

 SUPPORTING INFORMATION CAPTIONS

 S1 Fig. Distribution of bronchodilator drug response (BDR) in Puerto Ricans,

 Mexican (GALA II) and African Americans (SAGE). The 1,441 subjects selected from

 the extreme of the BDR distribution for this study were highlighted.

 S2 Fig. Global ancestry composition for **(a)** Puerto Ricans, **(b)** Mexicans and **(c)**

 African Americans. Each individual is represented by a vertical line and the ancestry

 composition is colored based on the percentage composition of African (red),

 European (blue) and Native American (green) ancestries.

 S3 Fig. Plot of the first two principal components of variation based on WGS

 genotypes of 1,441 individuals.

 S4 Fig. Diverse Convergent Evidence (DiCE) prioritization of 27 common variants

 (**Table 4**). Y-axis indicates the points allotted per SNP for each form of evidence

 according to **S15 Table**: statistical evidence (grey), informatic evidence (blue), and

 experimental evidence (orange). Evidence-specific scores for each SNP are provided

 in the table. SNPs with a DiCE score ≥ 4 are labeled with the nearest gene.

 S5 Fig. Regions overlapped with SNPs in LD with BDR-associated SNPs show

 enhancer activity in BSMCs. **(a)** H3K27ac ChIP-seq peaks in BSMCs overlap with

 SNPs in LD with rs28450894 (marked red). The GRCh37 coordinates are

 chr4:103486504-103491377 for region 1 and chr4:103527184-103531814 for

 region 2. In addition to BSMCs, H3K27ac data is shown for Roadmap Epigenomic

peripheral blood mononuclear cells (PBMSs) and lung tissue, as well as ENCODE

project data for GM12878, H1 human embryonic stem cell (H1-ESC) line, human

skeletal muscle cells and myoblasts (HSMM) and normal human epidermal

keratinocytes (NHEK). **(b)** Luciferase assay results for regions tested for enhancer

activity in BSMCs. *NFKB1* region 2 significantly increased the expression of

luciferase over empty vector control (Fold change = 2.24, $p < 0.01$). A red line marks

a fold change of one compared to the empty vector.

S6 Fig. A GGV plot showing the allele frequency of rs2845894 for different

populations based on 1000 Genomes Project. European populations include CEU

(Utah residents [CEPH] with northern and western ancestry), FIN (Finnish in

Finland), GBR (British in England and Scotland), TSI (Toscani in Italia) and IBS

(Iberian population in Spain). American populations include MXL (Mexican ancestry

from Los Angeles USA), PUR (Puerto Ricans from Puerto Rico), CLM (Colombians

from Medellin, Colombia) and PEL (Peruvians from Lima, Peru). African populations

include ASW (Americans of African ancestry in SW USA), ACB (African Caribbeans in

Barbados), GWD (Gambian in western divisions in the Gambia), MSL (Mende in

Sierra Leone), YRI (Yoruba in Ibadan, Nigeria), ESN (Esan in Nigeria) and LWK

(Luhya in Webuye, Kenya).

S7 Fig. Boxplot showing increasing number of copies of low BDR-associated T allele

of rs28450894 is associated with decreased expression of *SLC39A8* in blood

regardless of sex ($p = 0.0066$, FDR adjusted $p = 0.0856$, $\log_2(\beta) = -0.327$).

S8 Figure. Manhattan plot of the single locus BDR association testing for **(a)** Puerto

Ricans, **(b)** Mexicans and **(c)** African Americans. The horizontal lines are colored

blue for the suggestive significance thresholds and red for the Bonferroni-adjusted

genome-wide significance thresholds. Since no associations were close to the

Bonferroni-adjusted genome-wide significance thresholds in Mexicans and African

Americans, the red line is only marked in Puerto Ricans.

S9 Fig. A comparison of the top 1000 BDR associations between the array-based

and WGS-based genotype data. **(a)** A plot of association p-values of array-based and

WGS-based data. The p-values from the two data types showed high correlation,

especially for SNPs with more significant p-values. Genotype discordance of SAGE II

(b) and GALA II **(c)** WGS SNPs. The corresponding imputation R^2 of the same SNPs

in the HRC imputed array data is indicated by red ($R^2 \geq 0.8$) or blue ($R^2 < 0.8$).

S10 Fig. Percentage genotype concordance between Axiom LAT1 array and WGS

genotypes. Grey horizontal lines mark one, two and three standard deviations (S.D.)

from the mean percentage genotype concordance.

S11 Fig. Quantile-quantile (q-q) plots of the single locus BDR association for **(a)**

Puerto Ricans, **(b)** Mexicans and **(c)** African Americans and **(d)** trans-ethnic meta-

analysis. The genomic inflator factors (λ_{GC}) are shown on the q-q plots.

S12 Fig. Quantile-quantile (q-q) plots of SKAT-O analysis of biallelic common and

rare SNPs grouped by 1kb windows for **(a)** Puerto Ricans, **(b)** Mexicans and **(c)**

African Americans and **(d)** all individuals in all three populations. The genomic

inflator factors (λ_{GC}) are shown on the q-q plots.

S1 Table. Novel common variants in discovery cohort by population.

S2 Table. Chromosomal location of BDR-associated common variants.

S3 Table. Common variant BDR association results by population.

S4 Table. Lung-related phenotypes previously reported for BDR-associated
common variants and their nearest genes.

S5 Table. Variation in BDR Status explained by significant and suggestively
associated variants identified by trans-ethnic meta-analysis and SKAT-O.

S6 Table. Descriptive statistics for replication cohorts.

S7 Table. Replication of common variant associations.

S8 Table. Size of DNA inserts used in luciferase assay and key variants included
within those sequences.

S9 Table. Functional annotations for BDR-associated common variants in Table 4.

S10 Table. Replication of previously reported BDR-associated SNPs in the current
study by population and trans-ethnic meta-analysis.

S11 Table. Chromosomal location of identified SKAT-O regions by genome build.

S12 Table. Functional annotations for variants within SKAT-O regions.

S13 Table. Correlation between baseline lung function, top ten principle

components, other covariates included in association analyses in the discovery

study population (N =1,441).

S14 Table. Number of African American samples selected for RNA-Seq based on

BDR status and number of copies of low BDR-associated allele.

S15 Table. Diverse Convergent Evidence (DiCE) approach scoring rubric.

Total Available Subjects

GALA II / SAGE

6,172 Subjects
- 2,319 PR
- 1,556 MX
- 2,297 AA
- 830 Others

WGS Study Population Selection
(Extremes of BDR distribution)

1,484 Subjects
- 494 PR
- 500 MX
- 490 AA

WGS Quality Control

1,441 Subjects
- 483 PR
- 483 MX
- 475 AA

Discovery Analysis

Common variant	Common and rare variant
Logistic regression	*SKAT-O, 1kb, 500bp increment*
PR	PR
MX	MX
AA	AA
Meta-analysis	Combined

Replication of 27 Associated Common Variants

BDR (high/low)
GALA I (18/90 PR, 105/97 MX), SAGE I (57/84 AA), HPR (42/372 PR),
SAPPHIRE (465/557 AA), CHOP (158/122 AA)

Perform ChIP-seq to Identify H3K27ac Peaks in BSMC

Prioritize Associated Loci for Further Analyses

DiCE Priority Score

Identify NFKB1 Loci as Highest Priority for Further Analyses

Functional analyses

Luciferase assay, RNA-seq (39 AA)