Title

A standardized framework for representation of ancestry data in genomics

studies, with application to the NHGRI-EBI GWAS Catalog

Authors

1. Joannella Morales[1*], jmorales@ebi.ac.uk

2. Emily H. Bowler[1], ehbowler13@gmail.com

3. Annalisa Buniello[1], buniello@ebi.ac.uk

4. Maria Cerezo[1], mcerezo@ebi.ac.uk

5. Peggy Hall[2], Peggy.Hall@nih.gov

6. Laura W. Harris[1], ljwh@ebi.ac.uk

7. Emma Hastings[1], emmakhastings@gmail.com

8. Heather A. Junkins[2], junkinsh@mail.nih.gov

9. Cinzia Malangone[1], cinzia@ebi.ac.uk

10. Aoife C. McMahon[1], aoifem@ebi.ac.uk

11. Annalisa Milano[1], annalisa@ebi.ac.uk

12. Danielle Welter[1], dwelter@ebi.ac.uk

13. Tony Burdett[1], tburdett@ebi.ac.uk

14. Fiona Cunningham[1], fiona@ebi.ac.uk

15. Paul Flicek[1], flicek@ebi.ac.uk

16. Helen Parkinson[1], parkinson@ebi.ac.uk

17. Lucia A. Hindorff[2], hindorffl@mail.nih.gov

18. Jacqueline A. L. MacArthur[1*], jalm@ebi.ac.uk

[1]*European Molecular Biology Laboratory, European Bioinformatics Institute,*

Wellcome Genome Campus, Hinxton, Cambridge, CB10 1SD, UK

[2]*Division of Genomic Medicine, National Human Genome Research Institute,*

National Institutes of Health, Bethesda, MD 20892, USA

Author information

31 Joannella Morales and Jacqueline MacArthur are corresponding authors.

32 Lucia A. Hindorff and Jacqueline A.L. MacArthur share joint last authorship of

33 this manuscript.

Abstract

Background

The accurate description of ancestry is essential to interpret and integrate

human genomics data, and to ensure that advances in the field of genomics

benefit individuals from all ancestral backgrounds. However, there are no

established guidelines for the consistent, unambiguous and standardized

description of ancestry. To fill this gap, we provide a framework, designed for

the representation of ancestry in GWAS data, but with wider application to

studies and resources involving human subjects.

Result

Here we describe our framework and its application to the representation of

ancestry data in a widely-used publically available genomics resource, the

NHGRI-EBI GWAS Catalog. We present the first analyses of GWAS data

using our ancestry categories, demonstrating the validity of the framework to

facilitate the tracking of ancestry in big data sets. We exhibit the broader

relevance and integration potential of our method by its usage to describe the

well-established HapMap and 1000 Genomes reference populations. Finally,

to encourage adoption, we outline recommendations for authors to implement

when describing samples.

Conclusions

While the known bias towards inclusion of European ancestry individuals in

GWA studies persists, African and Hispanic or Latin American ancestry

populations contribute a disproportionately high number of associations,

suggesting that analyses including these groups may be more effective at

identifying new associations. We believe the widespread adoption of our

framework will increase standardization of ancestry data, thus enabling

improved analysis, interpretation and integration of human genomics data and

furthering our understanding of disease.

Keywords

Genomics - Genome wide association studies – GWAS Catalog – Ancestry –

diversity – Population Genetics

Background

The past 15 years have seen a dramatic growth in the field of genomics, with

numerous efforts focused on understanding the etiology of common human

disease and translating this to advances in the clinic. Genome-wide

association studies (GWAS), in particular, are now a well-established

mechanism to identify links between genetic variation and human disease[1–

3]. The NHGRI-EBI GWAS Catalog[2,4], one of the largest repositories of

summary GWAS data, contains over 3,000 publications and 43,000 SNP-trait

associations as of July 2017. The Catalog is indispensable for researching

existing findings on common diseases, enabling further investigations to

identify causal variants, understand disease mechanisms and establish

targets for treatment[5–8].

Essential to the interpretation and application of genomic data is the accurate

description of the ancestry of the samples studied. Levels of genetic diversity

and patterns of linkage disequilibrium (LD) vary by ancestry, with important

implications for the experimental design, the generalizability of results and the

identification of causal variants. Standardized ancestry representation is

necessary to integrate data from different sources for further analysis and to

enable robust search functionalities in bioinformatics resources. There are

currently no established guidelines for the characterization and classification

of ancestral background information. This has led to ambiguity and

inconsistency, along with challenges in accessing and integrating data.

The need for genetic studies in more ancestrally diverse populations has been

repeatedly articulated[9], most recently by Popejoy and Fullerton[10] and by

Manolio, et al.[11]. The benefit of including diverse populations extends

throughout the translational research spectrum, from GWAS discovery efforts

to genomic medicine, for which variant interpretation can be greatly aided by

ancestrally diverse sequence information[12,13]. Although inclusion efforts

are improving over time, it is challenging to assess the status of such efforts,

or to implement improved approaches, without a standardized way of

representing ancestry data.

To fill these gaps, we here provide a framework to systematically describe

and represent detailed ancestry information. Our method is driven by an

analysis of data we manually curated from over 3,000 GWAS publications.

We developed it for immediate application to the GWAS Catalog to make

curated data accessible, searchable and compatible with other genomics

data. However, the framework is broadly applicable to studies and resources

involving human subjects, and its widespread adoption will enable improved

analysis, interpretation and integration of data, ultimately furthering our

understanding of the genetic architecture of disease in different human

populations.

Results

Standardized ancestry framework

We developed the framework to enable the generation of comprehensive and

standardized representation of ancestry information for samples included in

GWAS. We had several motivations: 1) our observation, from curating

thousands of publications, that ancestry data is often poorly represented,

inconsistent or even completely absent, 2) the requirement to describe,

represent and store ancestry data in a manner that allows robust searching

and visualization of data in the GWAS Catalog, 3) the increased interest in

ancestry from the scientific community and 4) the need to analyze the

ancestry of samples, assess diversity and generate metrics that would allow

the community to identify and address gaps in this area.

Our framework involves representing ancestry data in two forms: (1) a

detailed sample description and (2) an ancestry category from a controlled list

(Table 1). Detailed descriptions aim to capture accurate, informative and

comprehensive information regarding the ancestry or genealogy of the

samples. Ancestry categories are used to establish hierarchical relationships

between groups and populations. We believe this dual representation of

ancestry is both informative and useful. The two types of descriptions

complement each other. The detailed description is granular and

heterogeneous, whereas the use of a limited number of categories reduces

complexity, facilitating data representation, integration, searchability and

further analyses. The framework also allows for country information to be

recorded providing additional detail on sample demographics.

<u>Application of the framework to the GWAS Catalog and other resources</u>

Application of this framework by curators to samples reported in publications

relies on manual interpretation and extraction of author-reported data. To

ensure consistent application by GWAS Catalog curators, we created a set of

comprehensive data extraction guidelines (Supplementary Note). When the

information provided by authors is limited or ambiguous, we consider country

of recruitment demographics and peer-reviewed population genetics

publications.

We have now generated detailed descriptions and assigned ancestry

categories to 3,000 publications, representing 4,000 separate GWA studies

and 83 million individuals, as of July 2017. A full list of all detailed descriptions

currently included in the Catalog is provided in Supplementary Table 1.

Examples that illustrate how the framework was applied to Catalog samples

can be found in Supplementary Table 2. All curated ancestry data is available

from the GWAS Catalog website[4] (Figure 1) and via download[14].

Detailed description

The detailed description aims to accurately represent the ancestry or

genealogy of each distinct group analyzed in a specific study in detail, as

reported by the author. Information about the homogeneity of the samples,

including whether the cohort is admixed or taken from a founder or isolated

population, is included. In the GWAS Catalog, the majority of the detailed

descriptions include terms that describe the location of participants' ancestors

over the past few generations ("French", "Japanese"), while admixed

populations are primarily described using ethnic descriptors ("Hispanic").

Isolated populations are described using either location or ethnicity terms in

addition to being described explicitly as genetically isolated ("Old Order Amish

(founder or genetic isolate) population", "Norfolk Island (founder or genetic

isolate) population".

Ancestry categories

Ancestry category assignment from the list presented in Table 1 requires

careful consideration.   When clearly stated, author-reported categories are

extracted, with precedence given to genetically-inferred data. If a category is

not stated, curators infer the category based on the detailed description for

the sample, which, as noted above, represents author-provided information.

In the absence of any ancestry data, the category "Not Reported" is assigned,

unless geographical location of sample recruitment is stated. In such

instances, curators infer ancestry from external sources, such as the United

Nations[15] and The World Factbook[16]. Selecting a category for samples

that derive from a country with a homogenous demographic composition,

such as Japan, is straightforward. However, for samples from populations with

limited known genetic genealogy, such as Azerbaijan, or for samples recruited

in countries with ancestral diversity, such as Singapore, assigning a category

is   more   challenging.   These   sources   are   particularly   useful   to   obtain

geographical and country-specific population information. The World Factbook

is   a   regularly   updated,   comprehensive   compendium   of   worldwide

demographic data, covering all countries and territories of the world. However,

since it does not necessarily provide ancestry data, the World Factbook is

consulted when the only known information is the country of recruitment of

samples. We expect that as increased care is taken to accurately report

ancestry   data,   reliance   on   this   resource   will   decrease.   Peer-reviewed

population genetic studies that characterize the genetic background of a given

population may also be consulted. This is particularly helpful in cases where

the sample cohort self-reported or is described using geographical or ethno-

cultural terms, such as "Scandinavian" or "Punjabi Sikh". Supplementary

Table 3 provides a list of countries for which external sources were consulted.

If the ancestry data provided in publications does not allow the resolution of

samples into ancestrally distinct sets, more than one category may be

selected from the list in Table 1, such as in the Catalog entry for Jiang R *et.*

al. [17][18].

207   Country information

Country of recruitment (Figure 1) and country of origin provides additional

demographic information and is extracted for each distinct sample set.

Country of origin or recruitment is author-reported and not inferred from

ancestry data. An exception is made for occasions when authors combine

country of recruitment with an ancestry description ("Singaporean Chinese").

In these cases, we infer the country of recruitment ("Singapore") although it is

not explicitly stated. Country of origin is defined as the country of origin of the

study participant's grandparents or as the genealogy of the participants dating

several generations.

Wider application of the framework

The HapMap[19] and 1000 Genomes[20] projects have delivered a

comprehensive survey of human genetic variation across worldwide

populations. The application of our method to the ancestry representation of

these reference populations therefore provides huge integration potential and

demonstrates the relevance of our framework beyond GWAS publications.

For all populations, we assigned ancestry category, country of recruitment,

country of origin and a detailed description, if provided by each project

(Supplementary table 4). We are developing an integrated tool to provide

access to population genetic and linkage disequilibrium data from these

reference populations. This will be available in the near future.

To facilitate the application of our approach to databases and resources, we

have developed an ancestry-specific ontology based on our framework. We

have defined terms, identified synonyms and established hierarchical

relationships between all curated terms and categories. Integration of the

ontology into any search interface will enable users to perform more powerful

and precise ancestry-related queries[21]. We aim to integrate it into the

GWAS Catalog website in the near future. The ancestry ontology[22] can be

browsed and downloaded (manuscript in preparation).

<u>Application of framework to assess changes in diversity</u>

Several members of the community have called for greater efforts to increase

diversity in genomics studies[10]. However, it is difficult to assess progress or

identify concrete areas for improvement without a method to easily track

changes in ancestry data. Our framework addresses this challenge by

requiring the use of specific categories when describing ancestry. This

process establishes hierarchical relationships between populations, thus

facilitating reproducible tracking of changes in standardized categories over

time. A number of authors[9,10] have reviewed the ancestry distribution in the

Catalog, but focused exclusively on the detailed descriptions, which are

heterogeneous as they are based on the authors' language. Here we present

the first analyses using our ancestry categories and demonstrate the validity

of our framework to facilitate the tracking of ancestry in big data sets.

As expected, similar to previous reports[10], we found that the majority (78%)

of individuals in the Catalog are exclusively of European ancestry (Figure 2a).

The second largest group includes individuals of Asian descent (11%), with

East Asians comprising 9% of the Catalog's samples. The disproportionate

focus on Europeans was more prevalent in the earlier years of the Catalog

(86% of individuals in studies published between 2005 and 2010; 76%

between 2011 and 2016, Figure 3). The reduced proportion of European

ancestry individuals added to the Catalog in the last 5 years reflects an

increase in Asian (7.5% to 11.4%, 1.5-fold increase), African (0.8% to 2.8%,

3.5-fold increase), Hispanic or Latin American (0.1% to 1.2%, 9-fold increase)

and Middle Eastern (0.01% to 0.08%, 7-fold increase) samples. Though the

proportion of Hispanic or Latin Americans exhibited the largest increase, when

considering the absolute number of individuals, the largest increase, by far,

came from Asian populations; Asian ancestry individuals increased from

almost 900,000 in the first 5 years to over 7 million added to the Catalog in the

last 5 years, compared to an increase of 721,000 Hispanic or Latin

Americans.

A similar trend is observed when analyzing the ancestry distribution in

independent GWA studies. Approximately 50% of all studies are performed on

exclusively European ancestry individuals, with an additional 24% of studies

including some samples of European descent (Figure 2b). A more granular

analysis of the traits with the largest number of GWAS in the Catalog

presented the same European bias, with 57-80% of studies, depending on the

trait, carried out in European ancestry individuals, followed by East Asians (7-

28% of studies) (Supplementary Fig. 3). In studies that analyzed multiple

ancestries, the vast majority (> 90%) include some European ancestry

individuals, regardless of the trait. The traits that display the largest proportion

of ancestral diversity are anthropometric traits, such as body mass index

(BMI) and height, and common diseases, including type 2 diabetes and

cardiovascular disease (Supplementary Fig. 3).

Interestingly, when we focused our analysis on the number of associations

identified in each ancestry category, we noted a different distribution to the

ancestry distribution of individuals (Figure 2c). This disparity is particularly

pronounced for studies including African or Hispanic or Latin American

samples; African ancestries contribute 2.4% of individuals but 7% of

associations, while Hispanic/Latin Americans contribute 1.3% of individuals

compared to 4.3% of associations. The opposite effect was seen in

Europeans, with 54% of associations compared to 78% of individuals. In

addition, we also observed a disproportionate number of associations

contributed by the "Multiple ancestries" category, likely reflecting the Catalog's

inclusion of trans-ethnic meta-analyses and replication efforts in diverse

ancestries.

Recommendations to authors

The analysis of the over 3,000 GWAS publications revealed inconsistent and

ambiguous reporting of ancestry data, with a significant percentage of studies

(~ 4%) not reporting any ancestry information at all. Given that there are no

established guidelines for the description of ancestry, and in an effort to assist

the community as it seeks to improve in this area, we here provide a set of

specific recommendations for authors, also summarized in Box 1. We believe

implementation of these recommendations will improve the quality of reporting

and have a positive impact on the interpretation of published results, data re-

use and reproducibility.

We recommend that authors make every effort to generate a detailed

description for each distinct set of individuals included in their studies. Authors

should also assess whether the genetic diversity of each distinct set is

representative of one of the known populations listed and defined in Table 1,

and indicate the corresponding category in the publication. If authors have no

knowledge about the ancestry of the participants, are not able to determine it

or cannot share it due to confidentiality concerns, we suggest noting this

explicitly in the publication.

Where possible authors should provide genetically inferred, in addition to self-

reported, ancestry information as the latter is often not an accurate

representation of the underlying genetic background. Software to assess and

control for ancestry is readily available and computationally feasible[23–28].

Encouragingly the proportion of studies that assess ancestry by genetic

methods has increased, from 25% in the early days of GWAS (the first 100

eligible publications, 2005-2008) to 57% in 2016 (the first 100 eligible

publications; Supplementary Fig. 4). Thus, we suggest authors consider using

the methods listed in Supplementary Box 1 when inferring the ancestry of

samples.

In general, terms that pertain to an individual's ethno-cultural background

should be avoided, unless this provides additional information regarding the

genealogy of the samples. In such cases a descriptor that accurately reflects

the underlying genetics should also be provided. For example, when

describing "Punjabi Sikh" participants, every effort should be made to assess

the genetic background and to indicate this in the publication, for example by

stating "Punjabi Sikh South Asian ancestry individuals" rather than simply

"Punjabi Sikh" or "Sikh".

Particular care should be taken to note if a sample derives from founder or

genetically isolated population; given their homogeneity and reduced genetic

variation, these populations are especially well-suited for GWAS[29] and are

increasingly used as sample sources. To reduce ambiguity, when describing

isolates, the broader genetic background within which the population clusters

should also be indicated. For example, Old Order Amish participants should

be described as "Old Order Amish population isolate individuals of European

ancestry". While describing admixed populations can be challenging due to

varying levels of admixture, every effort should be made to explicitly note

whether the population is admixed and the ancestral backgrounds that

contribute to admixture. For example "Hispanics/Latinos are ethnically

heterogeneous, with admixture of European, West African, and Amerindian

ancestral populations", as stated in Hodonsky 2017[30].

Discussion

Summary

In this article we describe a framework for the standardized representation of

ancestry data from genomics studies. Our method provides structure to

unstructured data, enabling robust searching across large datasets and

integration across resources. We have established validity by application to

the over 3,000 GWAS publications currently in the GWAS Catalog and

performed a detailed analysis. These data represent over 83 million

individuals from diverse ancestral backgrounds, and, as such, it is likely one

of the largest and most widely-used repositories of curated ancestry data. We

have demonstrated relevance to, and integration potential with, other data

types and studies by using our method to describe well-characterized

reference populations, such as the HapMap and 1000 Genomes project

populations. This will greatly facilitate integration of studies involving these

populations with data included in the Catalog, and, indeed, with any other

resource that implements our framework. We display the utility of the ancestry

categories to simplify the tracking of efforts towards diversity, allowing the

identification of gaps and highlighting specific areas for improvement.

Interestingly, in addition to confirming known biases, our category-based

analyses revealed that African and Hispanic or Latin American ancestry

populations contribute a disproportionately high number of associations,

suggesting that analyses including these groups may be more effective at

identifying new associations. Finally, stemming from our extensive manual

review of publications, we note a lack of current standards with regard to

ancestry reporting and offer recommendations to authors to implement when

describing their samples. This, we believe, will increase consistency and

reduce ambiguity, facilitating the interpretation of results.

<u>Limitations to the framework</u>

There are challenges inherent to both the design of the framework and its

application. We recognize the sensitivities surrounding the concepts of race,

ethnicity and ancestry, and that these terms are often used interchangeably

without making a distinction between physical appearance, cultural traditions

and genetic variation. This conflation can often be observed in censuses and

other demographic tools, influencing how individuals and communities

describe their background. The United States Census, for example, defines

"White" as "a person having origins in any of the original peoples of Europe,

the Middle East, or North Africa"[31], even though Middle Eastern and North

African populations are known to cluster, in genetic analyses, independently

from European ancestry populations. We thus recommend that authors

continue to move away from relying solely on self-reported information and,

as much as possible, also use genomic mechanisms to infer and describe the

ancestry of participants.

We are aware that classifying the global human population is a challenging

endeavor and the subject of numerous publications by expert population

geneticists. Due to human evolution and patterns of migration, the ancestry of

a particular population is complex and its definition is dependent on time. It is

generally accepted that all modern human populations originated in Africa,

and, due to the relatively small amount of genetic variation between

populations from disparate geographical locations[20], genetic diversity

among modern populations may be more suitably described as a continuum.

However, it is both possible and useful to generate informative groupings.

Reference populations, such as those included in the HapMap and 1000

Genomes Projects, or ancestry informative markers[32] that allow populations

to be distinguished, have been characterized, and methods have been

developed to adjust for population stratification and separate samples into

clusters. In fact, these analyses between and within populations have

demonstrated that clusters identified through genomic methods generally

align with geographical and regional groupings [33]. Taking this into account,

we designed our framework such that our categories closely resemble

classifications currently in use by the community and defined by experts.

We do not view our categories as exhaustive or static. We envision that as

more cohorts from diverse populations are characterized, there might arise a

need to create additional categories or sub-categories. In addition,

anticipating that admixture is likely to increase in the future, due to migration,

for example, we also created categories to represent known (for example,

"Hispanic or Latin American") and emerging (for example, "Other admixed

ancestries") admixed groups. We recognize that classification of admixed

samples is particularly challenging. The degree and type of admixture may

vary within the population, and the accuracy of classification requires well-

defined reference samples, which are lacking for some groups. As the

community moves towards genetically-inferred ancestry descriptions, our

categories are likely to become more precise and granular over time.

We believe there are benefits to utilizing categories. Practically, categories

facilitate data integration and allow robust searches, which is an essential

component of databases such as the GWAS Catalog. Also, the use of

categories can be useful when carrying out further analyses. For example,

querying the generalizability of results, identifying ancestry-specific

associations or utilizing linkage disequilibrium information from a reference

population to identify independent signals. We did not set out to define novel

or authoritative global ancestral classifications. Rather, we aim to formalize

and encourage the use of existing classifications to increase standardization

and improve resource functionality, ultimately enabling more robust scientific

analyses.

<u>Assessing diversity in genomics</u>

Several reports have been published urging the scientific community to

ensure that individuals from all ancestry backgrounds benefit from advances

in the field of genomics[9,10]. Our proposed framework lays out a mechanism

for the generation of consistent and comprehensive ancestry descriptions and

this, in turn, facilitates the establishment of metrics and ultimately, the tracking

of ancestry data over time.

Our analysis of individuals in GWA studies, using ancestry categories rather

than Catalog detailed descriptions, as carried out in previous studies, confirms

the persistent bias towards inclusion of European ancestry samples and the

modest trend towards increased diversity.   Our more robust analysis of

approximately 43,000 associations found a disproportionately larger

proportion of associations derived from African and Hispanic or Latin

American populations, many of which have significant African admixture[34],

than is expected based on the proportion of individuals. We suggest that the

higher degree of genetic diversity and reduced linkage disequilibrium (LD) in

African populations[35] offers an explanation for this result. Shorter LD blocks

in African populations facilitate the separation of nearby but independent

signals in a way that is more challenging in European populations, in which

LD blocks tend to be longer. Additionally, the inclusion of larger numbers of

individuals from African or Hispanic or Latin American populations allows for

the identification of variants specific to these ancestries. Together, these

observations suggest that utilizing samples from diverse populations for

genomic studies may be advantageous and yield increased and more

comprehensive results.

There are limitations to our analysis. First, considering that some cohorts

have been included in numerous GWAS, it is highly likely that some

individuals are represented multiple times in the Catalog. The impact of this is

the skewing of results towards commonly-used or publicly available cohorts,

which are perhaps likely to be of European or Asian ancestry. Another

limitation stems from our criteria for inclusion of associations. Since we only

include variants with a p-value $< 1 \times 10^{-5}$ and only the "index" variant at each

locus, our analysis does not take into account all associations. To address

this and make the Catalog more comprehensive, we now also include in the

Catalog published summary statistics[36]. Finally, we were unable to assign a

category to associations identified in studies that include multiple ancestries.

This may be a factor contributing to the reduced number of associations

derived from European populations, since the vast majority of multiple

ancestry studies include Europeans (Figure 2b).

The analysis of diverse ancestries is advantageous from a scientific

perspective. No one population contains all human variants[37], and alleles

that are rare in one population may be common in a different population and

thus easier to detect. Studies of diverse populations may also aid in fine

mapping of existing signals or in identifying population-specific functional

variation[37,38]. Variant interpretation for genomic medicine in ancestrally

diverse or admixed populations relies on the availability of non-European

allele frequencies, with potentially serious clinical consequences if such data

are not available[12]. Finally, disease burden of common or complex diseases

(e.g., cardiovascular disease or cancer) disproportionately impacts non-

European populations. Of the commonly studied traits, the largest diversity of

backgrounds was found for common anthropometric traits, heart disease, and

type 2 diabetes. This is perhaps not surprising considering that metrics for

these traits are easy to obtain, and the two diseases are among the top ten

causes of death around the world, according to the World Health

Organization[39]. It is also consistent with the observation that diseases for

which global disease burden is substantial tend to lead to increased funding

and research infrastructure. While we are encouraged by the trend we have

seen in recent years towards increased diversity, we note that there are still

very clear gaps as some groups continue to be underserved or ignored. We

strongly urge the scientific community to expand their efforts to assemble and

analyze cohorts, including especially underrepresented communities.

Recommendations to authors

Our analysis also validates the need for a framework to improve the

description of ancestry. Approximately 5.8% of individuals in the Catalog

(2005 - 2016) are currently labeled with the category "Not reported" due to a

lack of adequate information in the publication. Although confidentiality

concerns certainly contribute to this, this large proportion of uncharacterized

samples supports the notion that guidelines for the reporting of ancestry data

are an absolute necessity. For this reason, we offer recommendations to

increase standardization of ancestry reporting, with an emphasis on

genetically-inferred ancestry, in publications (Box 1). We encourage

implementation by authors reporting ancestry data and by editors reviewing

publications that include human subjects.

Conclusions

Genome-wide association studies have been enormously successful.

However, the lack of clarity regarding the ancestry of samples and the lack of

studies including diverse ancestral backgrounds raises questions about the

interpretation and generalizability of results across populations. The
framework we provide aims to address these challenges. It improves
standardization of ancestry, increases integration of data, supports the
assessment of diversity in large sets and facilitates further analyses. Its
widespread adoption will enable the scientific community to investigate the
generalizability of trait-associations across diverse populations, to identify
associations unique to specific ancestries, to identify novel variants with
clinical implications, and to help pinpoint causative variants, thus increasing
our understanding of common diseases.

Methods

GWAS Catalog data curation
Details of GWAS publication identification, GWAS Catalog eligibility criteria
and curation methods can be found on the GWAS Catalog website[40].
Extracted information encompasses publication information, study cohort
information, including ancestry, and SNP-trait association results. Curation of
ancestry data from the literature was performed according to Ancestry
Extraction Guidelines outlined in the Supplementary Note.

1000 Genomes and HapMap Project population ancestry assignment
Information describing the 1000 Genomes[20] phase 3 and HapMap
Project[19] phase 3 populations was taken from the Coriell Institute
website[41]. Ancestry information, including ancestry category, country of
recruitment, country of origin and additional information, was assigned to each

population following the GWAS Catalog ancestry extraction guidelines

(Supplementary Note).

<u>GWAS Catalog ancestry analysis</u>

To determine the distribution of individuals, associations and traits by ancestry

category, we first downloaded all Catalog data in tabular form[14]. All data

(gwas-catalog-associations_ontology-annotated.tsv,   gwas-catalog-

ancestry.tsv,   gwas-catalog-studies_ontology-associated.tsv,   gwas-efo-trait-

mappings.tsv) included in these analyses were curated from GWA studies

published between 2005 and the end of 2015, with a release date of July 18

2017. The data can be found on the Catalog's FTP site[42].

<u>Analysis of ancestry assessment methods in a subset of the GWAS Catalog</u>

We selected the first 100 publications included in the Catalog (approximately

covering the period between March 2005 to January 2008), and for

comparison, the first 100 publications from 2016. For each publication, the

method was assessed and classified into one of the following: 1. Self-

reported, 2. Genetically assessed, 3. Ancestry stated without method, 4.

Inferred from limited ancestry-related information (e.g. country information), 5.

No ancestry information reported and 6. Mixed method (when a combination

of methods was utilized to describe the study samples). Publications classified

as "Genetically assessed" includes those where the author had clearly

identified the genetic ancestry or admixture of the population, for example by

using methods such as those described in Supplementary Box 1. It also

includes those that confirmed self-reported information or defined samples

based on self-reports but then excluded genetic outliers. Publications where

no ancestry was stated, but curators inferred an ancestry based on country

information are included in the fourth classification. In many cases authors

used a statistical method to assess or control for ancestry or population

stratification, without assigning individuals to a particular category, for

example using a continuous axis of genetic variation from PCA to compute

the association statistic. However, since this did not add any information that

curators could use to assign a population ancestry to the study, it was not

included under category 2.

Declarations.

Ethics approval and consent to participate

Not Applicable

Consent for publication

Not Applicable

Availability of data and materials

The datasets generated and/or analyzed during the current study are

available on the NHGRI-EBI GWAS Catalog search interface[4] and in

spreadsheet form[14].

Competing interests

PF is a member of the Scientific Advisory Board of Omicia, Inc.

Funding

Research reported in this publication was supported by the National

Human Genome Research Institute and the National Institute of General

Medical Sciences of the National Institutes of Health under Award Numbers

U41-HG007823 and U41-HG006104. The content is solely the responsibility

of the authors and does not necessarily represent the official views of the

National Institutes of Health. This research was also supported by the

European Molecular Biology Laboratory. L.A.H., P.H. and H.J. are employees

of the National Human Genome Research Institute.

Authors' contributions

J.M., J.A.L.M., P.H. H.A.J. and L.A.H. conceived this study and

developed the ancestry framework. J.M., J.A.L.M., E.H.B., A.B., M.C., P.H.,

L.W.H., H.A.J., A.C.M., A.M. and L.A.H. performed curation of ancestry data

of GWAS Catalog publications. J.M., J.A.L.M., M.C., T.B. and L.A.H. analyzed

the distribution of ancestry categories in the Catalog and interpreted the data.

L.W.H, J.A.L.M., L.H. and J.M. assessed the methods of ancestry

determination utilized in GWAS Catalog studies and interpreted the data.

A.C.M. and J.M. generated the figures. J.M., J.A.L.M and L.W.H. generated

the Tables. E.H., D.W., C.M. and T.B. developed the GWAS Catalog curation

and search interfaces. D.W. created the ancestry ontology, with contributions

from J.M., J.A.L.M. and E.H.B. All authors contributed to the final manuscript,

with J.M., J.A.L.M. and L.A.H. playing the key roles.

Acknowledgements

The authors wish to thank all GWAS Catalog users and authors of studies included in the Catalog. We also thank Chris Gignoux for his expert review of the genomic methods of ancestry determination discussed in this manuscript and to Teri Manolio for valuable discussion.

References

1. Visscher PM, Wray NR, Zhang Q, Sklar P, McCarthy MI, Brown MA, et al. 10 Years of GWAS Discovery: Biology, Function, and Translation. Am. J. Hum. Genet. 2017;101:5–22.

2. MacArthur J, Bowler E, Cerezo M, Gil L, Hall P, Hastings E, et al. The new NHGRI-EBI Catalog of published genome-wide association studies (GWAS Catalog). Nucleic Acids Res. 2017;45:D896–901.

3. Welter D, MacArthur J, Morales J, Burdett T, Hall P, Junkins H, et al. The NHGRI GWAS Catalog, a curated resource of SNP-trait associations. Nucleic Acids Res. 2014;42:D1001-1006.

4. GWAS Catalog [Internet]. [cited 2017 Aug 4]. Available from: http://www.ebi.ac.uk/gwas/

5. Onengut-Gumuscu S, Chen W-M, Burren O, Cooper NJ, Quinlan AR, Mychaleckyj JC, et al. Fine mapping of type 1 diabetes susceptibility loci and evidence for colocalization of causal variants with lymphoid gene enhancers. Nat. Genet. 2015;47:381–6.

6. Sivakumaran S, Agakov F, Theodoratou E, Prendergast JG, Zgaga L,

Manolio T, et al. Abundant pleiotropy in human complex diseases and traits.

Am. J. Hum. Genet. 2011;89:607–18.

7. Pal LR, Moult J. Genetic Basis of Common Human Disease: Insight into the

Role of Missense SNPs from Genome-Wide Association Studies. J. Mol. Biol.

2015;427:2271–89.

8. Mullen J, Cockell SJ, Woollard P, Wipat A. An Integrated Data Driven

Approach to Drug Repositioning Using Gene-Disease Associations. PloS

One. 2016;11:e0155811.

9. Need AC, Goldstein DB. Next generation disparities in human genomics:

concerns and remedies. Trends Genet. TIG. 2009;25:489–94.

10. Popejoy AB, Fullerton SM. Genomics is failing on diversity. Nature.

2016;538:161–4.

11. Manolio TA. In Retrospect: A decade of shared genomic associations.

Nature. 2017;546:360–1.

12. Manrai AK, Funke BH, Rehm HL, Olesen MS, Maron BA, Szolovits P, et

al. Genetic Misdiagnoses and the Potential for Health Disparities. N. Engl. J.

Med. 2016;375:655–65.

13. Li YR, Keating BJ. Trans-ethnic genome-wide association studies:

advantages and challenges of mapping in diverse populations. Genome Med.

2014;6:91.

14. GWAS Catalog [Internet]. [cited 2017 Aug 4]. Available from:

http://www.ebi.ac.uk/gwas/docs/file-downloads

15. UNSD — Methodology [Internet]. [cited 2017 Aug 4]. Available from:

https://unstats.un.org/unsd/methodology/m49/

16. The World Factbook — Central Intelligence Agency [Internet]. [cited 2017

Aug 4]. Available from: https://www.cia.gov/library/publications/resources/the-

world-factbook/index.html

17. GWAS Catalog [Internet]. [cited 2017 Aug 14]. Available from:

http://www.ebi.ac.uk/gwas/search?query=22391508

18. Jiang R, French JE, Stober VP, Kang-Sickel J-CC, Zou F, Nylander-

French LA. Single-Nucleotide Polymorphisms Associated with Skin Naphthyl–

Keratin Adduct Levels in Workers Exposed to Naphthalene. Environ. Health

Perspect. 2012;120:857–64.

19. International HapMap 3 Consortium, Altshuler DM, Gibbs RA, Peltonen L,

Altshuler DM, Gibbs RA, et al. Integrating common and rare genetic variation

in diverse human populations. Nature. 2010;467:52–8.

20. 1000 Genomes Project Consortium, Auton A, Brooks LD, Durbin RM,

Garrison EP, Kang HM, et al. A global reference for human genetic variation.

Nature. 2015;526:68–74.

21. Smith B, Ashburner M, Rosse C, Bard J, Bug W, Ceusters W, et al. The

OBO Foundry: coordinated evolution of ontologies to support biomedical data

integration. Nat. Biotechnol. 2007;25:1251–5.

22. Ancestry Ontology [Internet]. [cited 2017 Aug 4]. Available from:

http://www.ebi.ac.uk/ols/ontologies/ancestro

23. Porras-Hurtado L, Ruiz Y, Santos C, Phillips C, Carracedo A, Lareu MV.

An overview of STRUCTURE: applications, parameter settings, and

supporting software. Front. Genet. 2013;4:98.

24. Alexander DH, Novembre J, Lange K. Fast model-based estimation of

ancestry in unrelated individuals. Genome Res. 2009;19:1655–64.

25. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D.

Principal components analysis corrects for stratification in genome-wide

association studies. Nat. Genet. 2006;38:904–9.

26. Kang HM, Sul JH, Service SK, Zaitlen NA, Kong S-Y, Freimer NB, et al.

Variance component model to account for sample structure in genome-wide

association studies. Nat. Genet. 2010;42:348–54.

27. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et

al. PLINK: a tool set for whole-genome association and population-based

linkage analyses. Am. J. Hum. Genet. 2007;81:559–75.

28. Price AL, Zaitlen NA, Reich D, Patterson N. New approaches to

population stratification in genome-wide association studies. Nat. Rev. Genet.

2010;11:459–63.

29. Cronin S, Berger S, Ding J, Schymick JC, Washecka N, Hernandez DG,

et al. A genome-wide association study of sporadic ALS in a homogenous

Irish population. Hum. Mol. Genet. 2008;17:768–74.

30. Hodonsky CJ, Jain D, Schick UM, Morrison JV, Brown L, McHugh CP, et

al. Genome-wide association study of red blood cell traits in

Hispanics/Latinos: The Hispanic Community Health Study/Study of Latinos.

PLoS Genet. 2017;13:e1006760.

31. Bureau UC. About [Internet]. Available from:

https://www.census.gov/topics/population/race/about.html

32. Paschou P, Lewis J, Javed A, Drineas P. Ancestry informative markers for

fine-scale individual assignment to worldwide populations. J. Med. Genet.

2010;47:835–47.

33. Novembre J, Johnson T, Bryc K, Kutalik Z, Boyko AR, Auton A, et al.

Genes mirror geography within Europe. Nature. 2008;456:98–101.

34. Adhikari K, Mendoza-Revilla J, Chacón-Duque JC, Fuentes-Guajardo M,

Ruiz-Linares A. Admixture in Latin America. Curr. Opin. Genet. Dev.

2016;41:106–14.

35. Campbell MC, Hirbo JB, Townsend JP, Tishkoff SA. The peopling of the

African continent and the diaspora into the new world. Curr. Opin. Genet. Dev.

2014;29:120–32.

36. GWAS Catalog [Internet]. [cited 2017 Aug 4]. Available from:

http://www.ebi.ac.uk/gwas/downloads/summary-statistics

37. 1000 Genomes Project Consortium, Auton A, Brooks LD, Durbin RM,

Garrison EP, Kang HM, et al. A global reference for human genetic variation.

Nature. 2015;526:68–74.

38. Asimit JL, Hatzikotoulas K, McCarthy M, Morris AP, Zeggini E. Trans-

ethnic study design approaches for fine-mapping. Eur. J. Hum. Genet. EJHG.

2016;24:1330–6.

39. WHO | The top 10 causes of death [Internet]. WHO. Available from:

http://www.who.int/mediacentre/factsheets/fs310/en/

40. GWAS Catalog [Internet]. [cited 2017 Aug 4]. Available from:

https://www.ebi.ac.uk/gwas/docs/methods

41. Coriell Biorepository [Internet]. [cited 2017 Aug 4]. Available from:

https://catalog.coriell.org/

42. Index of /pub/databases/gwas/releases/2017/07/18/ [Internet]. [cited 2017

Aug 4]. Available from:

ftp://ftp.ebi.ac.uk/pub/databases/gwas/releases/2017/07/18/

43. Huoponen K, Schurr TG, Chen Y, Wallace DC. Mitochondrial DNA

variation in an aboriginal Australian population: evidence for genetic isolation

and regional differentiation. Hum. Immunol. 2001;62:954–69.

44. Nagle N, Ballantyne KN, van Oven M, Tyler-Smith C, Xue Y, Taylor D, et

al. Antiquity and diversity of aboriginal Australian Y-chromosomes. Am. J.

Phys. Anthropol. 2016;159:367–81.

45. Martínez-Cruz B, Vitalis R, Ségurel L, Austerlitz F, Georges M, Théry S, et

al. In the heartland of Eurasia: the multilocus genetic landscape of Central

Asian populations. Eur. J. Hum. Genet. EJHG. 2011;19:216–23.

46. Scott EM, Halees A, Itan Y, Spencer EG, He Y, Azab MA, et al.

Characterization of Greater Middle Eastern genetic variation for enhanced

disease gene discovery. Nat. Genet. 2016;48:1071–6.

47. Homburger JR, Moreno-Estrada A, Gignoux CR, Nelson D, Sanchez E,

Ortiz-Tello P, et al. Genomic Insights into the Ancestry and Demographic

History of South America. PLoS Genet. 2015;11:e1005602.

48. Reich D, Patterson N, Campbell D, Tandon A, Mazieres S, Ray N, et al.

Reconstructing Native American population history. Nature. 2012;488:370–4.

49. Kayser M. The human genetic history of Oceania: near and remote views

of dispersal. Curr. Biol. CB. 2010;20:R194-201.

Acknowledgements.

Research reported in this publication was supported by the National Human

Genome Research Institute and the National Institute of General Medical

Sciences of the National Institutes of Health under Award Numbers U41-

HG007823 and U41-HG006104. The content is solely the responsibility of the

authors and does not necessarily represent the official views of the National

Institutes of Health. This research was also supported by the European

Molecular Biology Laboratory. L.A.H., P.H. and H.J. are employees of the

National Human Genome Research Institute. The authors wish to thank all

GWAS Catalog users and authors of studies included in the Catalog. We also

thank Chris Gignoux for his expert review of the genomic methods of ancestry

determination discussed in this manuscript and to Teri Manolio for valuable

discussion.

Figures, tables and additional files

Table 1. Ancestry categories. Distinct regional population groupings used in this

framework. They are assigned to cohorts with distinct and well-defined patterns of

genetic variation, in addition to individuals with inferred relatedness to these cohorts.

A full list of GWAS Catalog sample descriptions assigned to each category can be

found in supplementary table 2.

Ancestry category	Definition	Examples of detailed descriptions for samples included in the category
Aboriginal Australian	Includes individuals who either self-report or have been described by authors as Australian Aboriginal. These are expected to be descendants of early human migration into Australia from Eastern Asia and can be distinguished from other Asian populations by mtDNA and Y chromosome variation[43,44].	Martu Australian Aboriginal

African American or Afro-Caribbean	Includes individuals who either self-report or have been described by authors as African American or Afro-Caribbean. This category also includes individuals who genetically cluster with reference populations from this region, for example 1000 Genomes and/or HapMap ACB or ASW populations. We note that there is likely to be significant admixture with European ancestry populations.	African American, African Caribbean
African unspecified	Includes individuals that either self-report or have been described as African, but there was not sufficient information to allow classification as African American, Afro-Caribbean or Sub-Saharan African.	African, non-Hispanic black
Asian unspecified	Includes individuals that either self-report or have been described as Asian but there was not sufficient information to allow classification as East Asian, Central Asian, South Asian or South-East Asian.	Asian, Asian American
Central Asian	Includes individuals who either self-report or have been described by authors as Central Asian[45]. We note that there does not appear to be a suitable reference population for this population and efforts are required to fill this gap.	Silk Road (founder/genetic isolate)

East Asian	Includes individuals who either self-report or have been described by authors as East Asian or one of the sub-populations from this region (e.g Chinese). This category also includes individuals who genetically cluster with reference populations from this region, for example 1000 Genomes and/or HapMap CDX, CHB, CHS and JPT populations.	Chinese, Japanese, Korean
European	Includes individuals who either self-report or have been described by authors as European, Caucasian, White or one of the sub-populations from this region (e.g Dutch). This category also includes individuals who genetically cluster with reference populations from this region, for example 1000 Genomes and/or HapMap CEU, FIN, GBR, IBS and TSI populations.	Spanish, Swedish
Greater Middle Eastern (Middle Eastern, North African or Persian)	Includes individuals who self-report or were described by authors as Middle Eastern, North African, Persian or one of the sub-populations from this region (e.g. Saudi Arabian)[46]. We note there is heterogeneity in this category with different degrees of admixture as well as levels of genetic isolation. We note that there does not appear to be a suitable reference population for this	Tunisian, Arab, Iranian

	category and efforts are required to fill this gap.	
Hispanic or Latin American	Includes individuals who either self-report or are described by authors as Hispanic, Latino, Latin American or one of the sub-populations from this region. This category includes individuals with known admixture of primarily European, African and Native American ancestries, though some may have also a degree of Asian (e.g. Peru). We also note that the levels of admixture vary depending on the country, with Caribbean countries carrying higher levels of African admixture when compared to South American countries, for example. This category also includes individuals who genetically cluster with reference populations from this region, for example 1000 Genomes and/or HapMap CLM, MXL, PEL and PUR populations[34,47].	Brazilian, Mexican
Native American	Includes indigenous individuals of North, Central and South America, descended from the original human migration into the Americas from Siberia[48]. We note that there does not appear to be a suitable reference population for this category and	Pima Indian, Plains American Indian

	efforts are required to fill this gap.	
Not Reported	Includes individuals for which no ancestry or country of recruitment information is available.	
Oceanian	Includes individuals that either self-report or have been described by authors as Oceanian or one of the sub-populations from this region (e.g. Native Hawaiian)[49]. We note that there does not appear to be a suitable reference population for this category and efforts are required to fill this gap.	Solomon Islander, Micronesian
Other	Includes individuals where an ancestry descriptor is known but insufficient information is available to allow assignment to one of the other categories.	Surinamese, Russian
Other admixed ancestry	Includes individuals who either self-report or have been described by authors as admixed and do not fit the definition of the other admixed categories already defined ("African American or Afro-Caribbean" or "Hispanic or Latin American").	
South Asian	Includes individuals who either self-report or have been described by authors as South	Bangladeshi, Sri Lankan

	Asian or one of the sub-populations from this region (e.g Asian Indian). This category also includes individuals who genetically cluster with reference populations from this region, for example 1000 Genomes and/or HapMap BEB, GIH, ITU, PJL and STU populations.	Sinhalese
South East Asian	Includes individuals who either self-report or have been described by authors as South East Asian or one of the sub-populations from this region (e.g Vietnamese). This category also includes includes individuals who genetically cluster with reference populations from this region, for example 1000 Genomes KHV population. We note that East Asian and South East Asian populations are often conflated. However, recent studies indicate a unique genetic background for South East Asian populations.	Thai, Malay
Sub-Saharan African	Includes individuals who either self-report or have been described by authors as Sub-Saharan African or one of the sub-populations from this region (e.g. Yoruban). This category also includes individuals who genetically cluster with reference populations from this region for example 1000 Genomes	Yoruban, Gambian

| | and/or HapMap ESN, LWK, GWD, MSL, MKK and YRI populations. | |

Box 1. **Recommendations for authors reporting ancestry data in**

publications. These recommendations were generated by expert curators

following a detailed review of the over 3,000 GWAS publications included in

the Catalog.

1. Preferentially use genomic methods to assess the ancestry of samples

   included in the GWAS Catalog. See Box 1 for a description of

   commonly used methods.

2. Indicate whether the background of participants was self-reported,

   inferred by genomic methods or a combination of both. If genetically

   inferred, indicate the analytical procedure utilized.

3. Provide detailed information for each distinct group of samples,

   a. Ancestry descriptors should be as granular as possible (e.g.

      Yoruban instead of Sub-Saharan African, Japanese instead of

      Asian).

   b. Avoid using country or citizenship as a substitute for ancestry.

   c. Avoid using geographic descriptors that are part of a cohort

      name as a substitute for ancestry (e.g. TwinsUK cannot be

      assumed to be European ancestry).

   d. If a population self-identifies using sociocultural descriptors (e.g.

      Old Order Amish), clearly state the genetic ancestry within which

      this sub-population falls.

   e. If samples were derived from an isolated or founder population

      with limited genetic heterogeneity, clearly state the genetic

ancestry within which this sub-population falls.

f.   If available, genetic genealogy or ancestry of grandparents or

parents should be included.

4. Assign an ancestry category for each distinct group of samples. See

Table 1 for a list of ancestry categories. Refer to Supplementary Table

1 for a list of descriptors in use in the Catalog with their category

assignments.

5. Provide the sample size for each distinct group of samples included in

the analysis.

6. Provide country of recruitment.

7. If ancestry information is not available due to confidentiality, or any

other concerns, note this in the publication.

Figures

1. Figure 1 – Representation of ancestry data in the GWAS Catalog
search interface
2. Figure 2 – Ancestry category distribution in the GWAS Catalog
a. Figure 2a - Distribution of individuals by ancestry category
b. Figure 2b - Distribution of studies by ancestry category
c. Figure 2c - Distribution of associations by ancestry category
3. Figure 3 – Distribution of individuals in GWAS Catalog studies
published between 2005 – 2010 compared to 2011 – 2016.

Figure 1. Representation of ancestry data in the GWAS Catalog search interface (www.ebi.ac.uk/gwas). Ancestry-related data is found in the Studies and Associations tables (underlined in black) when searching the Catalog. This figure shows the results of a search for PubMed Identifier 27145994. The sample description can be found in the Studies table, either by pressing "Expand all Studies" or the "+" on the study of interest (highlighted in red). Sample ancestry is captured in 2 forms: (1) Detailed description (highlighted in blue) and (2) Ancestry category (highlighted in green). The latter follows the format: sample size, category, (country of recruitment). In cases where multiple ancestries are included in a study, the ancestry associated with a particular association is found as an annotation in the p-value column in the Associations table (highlighted in pink).

Figure 2. Ancestry category distribution in the GWAS Catalog. This figure summarizes the distribution of ancestry categories in percentages, of individuals (N=83,200,468; panel a), studies (N= 4,100; panel b) and associations (N=43,919; panel c). The largest category in all panels is European (aqua). At the level of individuals (a), the largest non-European category is Asian (bright pink), with East Asian (light pink) accounting for the majority. Non-European, Non-Asian categories together (yellow) comprise 4% of individuals, and there are 6% (white) of samples for which an ancestry category could not be specified. Panel c demonstrates the disproportionate contribution of associations from African (blue) and Hispanic/Latin American (purple) categories, when compared to the percentage of individuals (a, blue, purple, respectively) and studies (b, blue, purple, respectively).

1 **Figure 3. Distribution of individuals in GWAS Catalog studies published**

2 **between 2005 – 2010 compared to 2011 – 2016.** This figure displays the

3 distribution of individuals in percentages, included in the 915 studies

4 published between 2005 – 2010 compared to the distribution of individuals

5 included in the 2,905 studies published between 2011 – 2016.