Boreal river impoundments caused little change in fish diversity but clear

community assemblage shifts: A multi-scale analysis

Katrine Turgeon*[1,2], Christian Turpin[2] & Irene Gregory-Eaves[1]

[1]Department of Biology, McGill University, 1205 Docteur Penfield Avenue, Montréal, QC,

CANADA, H3A 1B1

[2]Hydro-Québec, Environment and Corporate Affairs, 75 René-Lévesque, Montréal, QC,

CANADA, H2Z 1A4

*Corresponding author: katrine.turgeon@mail.mcgill.ca

Keywords: Beta-diversity; boreal reservoirs; fish; hydroelectricity; multi-scale approach;

Québec; species assemblages; upstream

Abstract

1. Hydroelectricity is often presented as a clean and renewable energy source, but river flow regulation and fragmentation caused by dams are recognized to impact aquatic biodiversity in temperate and tropical ecosystems. However, the effects of boreal river impoundment are not clear as the few studies that exist have not been able to separate the hydrological changes brought about by dams from other factors (*e.g.* fish stocking, and species introduction).

2. We adopted a multi-scale analysis to examine changes in nearshore fish communities over 20 years (spanning before and after impoundment) using a network of 24 sampling stations spread across from four reservoirs and two hydroelectricity complexes located in the boreal region (Northern Québec, Canada). Given the remote location, confounding factors were minimal.

3. We found no strong temporal trends in alpha- and gamma-diversity in impacted stations (upstream and downstream of the dam) relative to reference sites across the three spatial scales. Using beta-diversity analyses, we also detected a high stability in fish composition over time and space at the complex and reservoir scales.

4. At the scale of the sampling stations, we observed higher rates of species turnover (beta-diversity) coincident with the time of reservoir filling and shortly after. Likewise, we detected species assemblage shifts that correlated with time since impoundment only at the sampling station scale. This pattern was masked at the complex and reservoir scales.

5. *Synthesis and applications*. Overall, the isolated effect of impoundment in these remote boreal ecosystems caused no loss of species and little change in fish diversity over 20 years, but resulted in substantial species assemblage shifts. Our work shows that examining

42   community data at different scales is key to understand the anthropogenic impacts on fish

43   biodiversity.

1. Introduction

In response to increased demand for energy, many large dams are currently in operation,

or are being constructed to provide hydroelectricity (Grill *et al.* 2015; Winemiller *et al.* 2016).

Dams transform large rivers into large reservoirs, affecting numerous important physical,

chemical and biological processes (Ward & Stanford 1995; Friedl & Wüest 2002). Dams also

fragment rivers by creating barriers to movement (Nilsson *et al.* 2005; Pelicice, Pompeu &

Agostinho 2015), and alter the natural hydrological regime of the ecosystem (*i.e.*, discharge and

water levels) upstream and downstream of the dam (Kroger 1973; Poff *et al.* 2007). These

modifications are susceptible to affect the overall biodiversity and ecosystem functions

(Rosenberg, McCully & Pringle 2000; Vörösmarty *et al.* 2010; Liermann *et al.* 2012).

The effects of impoundment on fish communities have been extensively studied in

temperate (Martinez *et al.* 1994; Bonner & Wilde 2000; Gido, Matthews & Wolfinbarger 2000;

Taylor, Knouft & Hiland 2001; Gehrke, Gilligan & Barwick 2002; Quinn & Kwak 2003), and

more recently in tropical ecosystems where many new dams have been constructed (de Mérona,

Vigouroux & Tejerina-Garro 2005; Agostinho, Pelicice & Gomes 2008; Li, Madden & Xu 2012;

Lima *et al.* 2016). However, very little is known about the effects of impoundment in boreal

ecosystems (but see Tereshchenko & Strel'nikov 1997; Sutela & Vehanen 2008). This deficiency

is surprising considering that hydroelectricity is a major source of energy in some Nordic

countries (*e.g.*, Norway: 96% of domestic electricity generation, Iceland: 70%, Canada: 58% and

province of Québec in Canada: 95%; IEA 2016).

Long-term monitoring of fish assemblages in reservoirs is critical (Elliott 1990; Gido,

Matthews & Wolfinbarger 2000) because reservoirs are young (average of < 60 years), novel

ecosystems, and they are highly dynamic in the first decades following impoundment (*i.e.*, non-

trophic equilibrium phase, Grimard & Jones 1982; Turgeon *et al.* 2016). Moreover, the time

needed for the fish community to adapt (or not) to the new reservoir conditions, will depend on

several factors such as geographic location, reservoirs characteristics (*i.e.*, reservoir area, water

quality), dam operation and management (*i.e.*, drawdown), complexity of the food web, and

species life history traits. These potential sources of variability stress for the importance of

replication. To extract generalities about the effects of impoundment on fish community across

all latitudes, we need to improve our approach by having an exhaustive examination of the

following elements: 1) observations on fish communities spanning before to after impoundment,

collected routinely for many years, 2) data collected from multiple sampling stations downstream

and upstream of the dam and 3) parallel measurements made in reference sites to identify

climatic or other regional environmental drivers of change.

In this study, we took a multi-scale approach to examine how the impoundment of rivers

affects fish communities in four boreal reservoirs using a long-term dataset collected by Hydro-

Québec (Fig. 1). This dataset consists of a large network of 24 sampling stations (including

upstream and downstream stations as well as reference sites) and spans from before the

construction to 10 or 20 years after the start of its operation, allowing for one of the most

thorough and robust evaluations of how impoundment affect fish communities. Moreover,

because of its remoteness, this dataset provides a rare opportunity to isolate the effect of

impoundment on fish communities from other anthropogenic confounding factors.

2. Materials and Methods

2.1 Study sites

LG complex - The La Grande Rivière hydroelectric complex (hereafter called LG

complex) is located on the eastern side of James Bay (Québec, Canada), on the Canadian Shield.

The LG complex resulted in the creation of seven large reservoirs (Table 1, Fig. 1), and in the

diversion of three large rivers, the Caniapiscau (water flow at its mouth reduced by 43%), the

Eastmain (reduction of 86%) and the Opinaca (reduction of 86%; Roy & Messier 1989).

Through these hydrological changes, the average annual discharge in La Grande Rivière has

increased from 1700 $m^3 \cdot s^{-1}$ to 3400 $m^3 \cdot s^{-1}$ (Roy & Messier 1989). Data have been routinely

collected from three reservoirs in the LG complex: Robert-Bourassa (RB; impounded in 1979),

Opinaca (OP; impounded in 1980) and Caniapiscau (CA; impounded in 1982; Table 1, Fig. 1).

Reservoirs LG3, LG4 and LaForge II have been impounded in 1981, 1983 and 1983

respectively. Laforge I have been impounded later in 1993 (Fig. 1). The territory is free of other

industrial activities and sparsely occupied by the Indigenous Cree peoples. Some mitigation

measures over the years included fish habitat improvement (new spawning area, vegetation

control, creation of shelters, containment) and the maintenance of fish movement (*i.e.*, migratory

pass in Robert-Bourassa was created in 1980). Each of the three reservoirs was paired with a

natural lake in proximity to the reservoir ("REF" stations; Fig. 1). Fish community data were

collected in stations downstream of the dam ("DO" stations; Fig. 1) and upstream of the dam

("UR" if the station was a river or stream before and "UL" if the station was a lake before

impoundment; Fig. 1). We expected that UR stations would demonstrate a more pronounced

change in diversity and fish assemblages than UL stations because of the drastic change in

habitat from lotic to lentic conditions. One downstream station (DORB) had an increased flow

after impoundment, and the three others (DO1OP, DO2OP and DO1CA; Fig. 1) had decreased

flow because sampling stations were in rivers that were diverted to create the reservoirs.

Sainte-Marguerite complex - The Sainte-Marguerite complex (SM) is located on the

Moyenne-Côte Nord portion of the Canadian Shield (Eastern Québec, Canada; Fig. 1). The

Sainte-Marguerite 3 reservoir (SM3) is deeper than the LG reservoirs (Table 1) and is located

within a canyon shape valley. The Sainte-Marguerite river was impounded by Hydro-Québec in

1998 and took 4 years to fill. A smaller downstream reservoir (SM2) was created in 1954 by

Gulf Pulp and Paper (pulp industry) and is now managed by Gulf Power Co. The Sainte-

Marguerite watershed is also relatively free of anthropogenic perturbation. Fish community data

were collected in two stations upstream of the SM3 dam (two UR), in one UR station in SM2

reservoir that is downstream of SM3 but cannot be classified as a "true" downstream station, and

in one reference station (Table 1, Fig. 1).

2.2 Field sampling

In RB and OP reservoirs, nearshore fish communities were sampled annually from 1978

to 1984, and then in 1988, 1992, 1996 and 2000. In RB, the pre-impoundment period

corresponds to 1978 whereas in OP this period corresponds to the years 1978 and 1979. In CA

reservoir, fish communities were sampled annually from 1980 to 1982, and in the 1987, 1991,

1993, 1997 and 1999. In CA, pre-impoundment data correspond to the years 1980 and 1981. In

SM3 and SM2, the fish community was sampled in 1992, 1996, 2005 and 2011, with the former

two years corresponding to pre-impoundment period.

In LG complex, fish sampling occurred monthly from June to September-October, with a

total of five sampling times per season in RB and OP reservoirs, and four times per season in CA

until 1995. After 1995 in the LG complex, and for the whole period in SM complex, the fishing

protocol was optimized to concentrate the sampling effort in July and August (Deslandes &

Fortin 1994). To standardize the time series, we only used the data for the months of July and

August. Four gill nets were used, set in pairs. In each pair, there was an experimental

multifilament gill net (45.7 m in length x 2.4 m in depth; mesh sizes ranged between 2.5 to 10.2

cm). The second net in the pair was a gill net of uniform mesh size (either with a stretch mesh

size of 7.6 cm or 10.2 cm). Each net pair was set perpendicular to shore. In one of the net pairs,

the gill net with uniform mesh size was directed onshore while the other pair had the gill net with

uniform mesh size directed offshore. Sampling periods lasted 48h (nets visited every 24 h) in LG

complex until 1982 and 24h from 1983, and lasted 48h in the SM complex. All fish caught were

counted, measured and weighed. No seine net or minnow traps were used in this sampling

program and no gill nets were set in the pelagic zone. Thus, the presence and abundance of small

species from the nearshore area and pelagic species were underestimated.

In LG complex, changes in water quality in the photic zone (0 - 10 m) were monitored at

the same sampling stations. Water quality variables measured were average water temperature

measured at every m in the photic zone, water transparency (measured as secchi disk depth),

dissolved oxygen concentration, pH and specific conductivity (all measured with a Hydrolab

multiprobe). Details of the methodology used in the collection and analysis of these data were

presented by Fréchette (1980).

2.3 Statistical analyses

Alpha- and gamma-diversity analysis

Diversity was assessed with extrapolated species richness, Pielou's J Evenness index, and

Shannon's H′ diversity index. The extrapolated species richness represents the number of species

for a given standardized number of net lifts and we used the second-order jackknife index

(Jack2; function specpool in the vegan R package v. 2.4-1; Oksanen *et al.* 2016). Shannon's H′

diversity index takes evenness and species richness into account and quantifies the uncertainty in

predicting the species identity of an individual that is taken at random from the dataset. Pielou's

J′ Evenness index ranges from near 0 (indicating pronounced dominance) to near 1 (indicating an

almost equal abundance of all species).

To examine changes in diversity metrics over time in impacted stations in relation to

reference sites, we used General Linear Mixed Effects Models (glmm; applying the lme function

from the nlme package v. 3.1-128). Here, we were interested to compare the slopes (*i.e.*,

interaction term between time since impoundment [TSI] and impacted vs. reference sites [RI],

Table 2). We examined the effect of river impoundment on diversity metrics at three spatial

scales: at the hydroelectric complex scale (gamma-diversity; pooling data for all impacted

stations in each complex), at the reservoir scale (gamma-diversity; pooling data from impacted

sampling stations in each reservoir) and at the sampling station scale (alpha-diversity). To

control for spatio-temporal dependence, we used random factors where sampling stations were

nested within reservoirs: ~1 + TSI |STATION/RES (where RES stands for reservoir identity).

We also used an autoregressive correlation structure (corAR1) to control for temporal

autocorrelation. We determined the autoregressive process in each time series by plotting each

time series and by observing the autocorrelation function (ACF) and the partial autocorrelation

function (PACF) on detrended data using an autoregressive integrated moving average model

diagnostic (astsa package v. 1.3 in R; Stoffer 2014). Errors followed an autoregressive process of

degree 1. Our glmms with the autocorrelated structure did not perform better than those without

based on AICc scores (Burnham & Anderson 2002). We present only the glmms without the

autocorrelated structure.

Beta-diversity analysis

To test species turnover rate over time and space, we computed Local Contributions to

Beta-Diversity (LCBD; using the *beta.div* function in R available at

http://adn.biol.umontreal.ca/~numericalecology/FonctionsR/) and Species Contributions to Beta-

Diversity (SCBD) indices on Hellinger-transformed species abundance community matrices

(Legendre and De Cáceres 2013). LCBD values indicate how unique is any fish composition of a

site relative to other comparable sites by assessing its contribution to the total variation in fish

composition in space and/or time. SCBD indicates how large of a contribution is a species has to

overall beta diversity in the dataset (Legendre & De Cáceres 2013; Legendre & Gauthier 2014).

For details about the calculation of LCBD and SCBD and the. We computed LCBD at the

complex scale, at the reservoir scale, and at the sampling station scale and SCBD at the sampling

station scale. At the sampling station scale, each station was evaluated separately, so the turnover

rate was in relation to time only.

2.4 Variation partitioning

We examined fish species assemblages at the three spatial scales using unbiased variation

partitioning based on RDAs (Redundancy Analysis) and adjusted R^2 (Peres-Neto *et al.* 2006)

with the *varpart* function in the vegan package (v. 2.4-1). We used the forward selection

procedure of Blanchet, Legendre & Borcard (2008). With variation partitioning analyses, the

overall variation in species assemblages can be divided into "fractions" attributable to different

data matrices as well as combinations of these matrices (*i.e.*, shared variation). Here, we used

four matrices: time since impoundment [TSI; *i.e.,* including years before and after

impoundment], spatial heterogeneity [SH; latitude, longitude and identity of each sampling

station and reservoir], water quality variables [WQV; water transparency, dissolved oxygen, pH,

conductivity and temperature] and fishing gear [G]. The total variation of species assemblages

was decomposed into 15 fractions at the complex and reservoir scales, and eight fractions at the

sampling station scale because the [SH] matrix is irrelevant at the sampling station scale. We

used the Hellinger-transformed abundance values of species. To produce the most parsimonious

model in RDAs, we performed forward selection using the double stopping criteria (*ordiR2step*

function in the vegan R package v. 2.4-1; Blanchet, Legendre & Borcard 2008). Because of a

small sample size, these analyses were not possible in the SM complex (only 3 to 5 observations

per sampling station).

3. Results

3.1 Changes in alpha-, beta- and gamma-diversity

Overall, OP reservoir had a higher mean extrapolated richness and diversity, and SM3

had a lower richness and diversity than RB and CA reservoirs (Table 1). Downstream stations

generally had higher extrapolated richness than upstream stations, but did not differ in diversity

and evenness (Fig. 2). Across all scales and categories of impacted stations (U vs D and UL, UR

vs. D), the temporal trends in richness, diversity and evenness in impacted stations were weak

and comparable to those observed in reference sites for both complexes (complex scale; Fig. 2,

Table 2, reservoir scale; Table S1, sampling station scale; Tables S2-S5).

For completeness, we also examined the temporal trends in impacted stations only,

without comparison with reference sites. At the complex scale, we did not detect any temporal

trend in diversity metrics when categories of impacted stations were combined (Model 1; Table

S6). When station categories were added in the model (*i.e.*, U vs. D, or UR, UL vs. D), richness

marginally decreased over time (Models 2 and 3; Table S6). This trend was strongly driven by

the low richness values observed in 2000 in RB (lower fishing effort in this one year). When this

data point was excluded from the analysis, the trend was not significant anymore. At the

reservoir scale, we found some decreasing temporal trends in RB (Table S7, Models 1, 2, and 3)

but found no temporal trends in the other reservoirs (Table S7).

The lack of strong temporal trends in alpha- and gamma-diversity was echoed by an

absence of clear beta-diversity patterns across space and time at either the complex (Fig. 3 a and

Fig. S1) and reservoirs scale (and Fig. S2). At both scales, relatively few Local Contribution to

Beta-Diversity (LCBD) values were significant, and the weight of LCBD values did not relate to

impoundment, nor to the impacted stations. However, when beta-diversity analyses were

conducted at the sampling station scale (*i.e.,* only comparing any one site to itself through time),

many of the significant LCBD values were apparent in upstream stations during and shortly after

filling (Fig. 3 b), showing a higher species turnover rate during this period.

3.2 Drivers of the shift in species assemblage

At the complex and reservoirs scales, spatial heterogeneity (SH) among sampling stations

was the main driver structuring fish assemblages (Fig. 4 a-b, Table S8). The effect of

impoundment only became a dominant predictor at the sampling station scale (Fig. 4 c, Table

S9). At the scale of the LG complex, SH explained 45% of the variation across all shared

fractions (25% explained by SH alone) and a significant proportion of the variation was shared

with water quality variables (WQV; 15%; Fig. 4 a). A similar pattern was observed at the

reservoir scale (Fig. 4 b; Tables S8 and S9). At the scale of the sampling stations, most of the

variation was explained by the shared effect of TSI and WQV (24%; Fig. 4 c and Table S3),

which suggests that fish responded very locally to impoundment and in a large extent to changes

in water quality associated with impoundment.

3.3 Species affected by impoundment

The effect of impoundment on species differed among reservoirs, and among the

categories of sampling stations (Fig. 5 and Fig. 6). In several upstream stations, we observed a

shift from a catastomids-dominated community (longnose sucker, *Catostomus catostomus* and

white sucker, *C. commersonii*) toward a pike-coregonids (northern pike, *Esox lucius*, whitefish

Coregonus clupeaformis and cisco, *C. artedi*) community after impoundment (Fig. 6). This shift

was supported by high contribution to beta-diversity (SCBD) for these species in upstream

stations (Fig. 5 b, c, d and e). Changes in species assemblages in upstream stations appear to

have mostly occurred within the first 5 years of impoundment (Fig. 6). In downstream stations,

no consistent pattern was observed but the marked changes were a decrease of the lake sturgeon

(*Acipenser fulvescens*) and an increase in walleye (*Sander vitreus*) in OP (Fig. 6 f), and a

decrease in abundance of the brook trout (*Salvelinus fontinalis*) in CA. These patterns were

echoed by the SCBD values (Fig. 5 d, d and e). Reference sites were more stable but also

experienced some changes in fish community structure, with a fluctuating dominance between

two predators in RB (*i.e.,* walleye and burbot, *Lota lota;* Fig. 6 g), and between the lake trout

(*Salvelinus namaycush*) and the two catastomids in CA (Fig. 6 i).

4. Discussion

4.1 Fish community response to impoundment

There is an extensive literature on fish community responses to impoundment in

temperate and tropical reservoirs, but little was known about boreal reservoirs (but see

Tereshchenko & Strel'nikov 1997; Sutela & Vehanen 2008). Our analyses showed that in four

remote large boreal reservoirs, there were no significant temporal trends in fish alpha- and

gamma-diversity at three spatial scales. No native species were lost, and non-native fish did not

colonise our boreal study systems. Our work is in marked contrast to the tropic, where there is

evidence of a net loss of species (Liew, Tan & Yeo 2016). As such, there appears to be an

important heterogeneity in fish diversity responses to impoundment. However, almost all studies

to date (including ours) have shown a general change in fish assemblage after impoundment.

From the literature, four main mechanisms have been suggested to cause a change in fish

assemblage in reservoirs: 1) shift from a lotic to a lentic environment upstream of the dam, 2)

dams as barriers to free movement, 3) alteration of the natural hydrological regime, and 4) higher

susceptibility of reservoirs to be invaded by non-native species. Are these mechanisms

comparable across latitudes? The shift from lotic to lentic conditions upstream of the dam

represents an extreme transformation to fish habitats and can exert a suite of selective pressures

not experienced by fish during their evolutionary history. This is especially true in the tropics

where fish have evolved in flowing waters (Gomes & Miranda 2001) and may lack the

morphological and behavioral characteristics, or the reproductive strategy and plasticity to

successfully occupy the new lentic habitats (Gomes & Miranda 2001; Agostinho, Pelicice &

Gomes 2008). Given the predominance of large rivers and streams in the tropics and temperate

environments, significant losses in richness in these regions have been attributed to the

transformation of ecosystems in lentic ones (Martinez *et al.* 1994; de Mérona, Vigouroux &

Tejerina-Garro 2005; Sá-Oliveira *et al.* 2015; Lima *et al.* 2016). In boreal regions, both large

lakes and rivers are common (Messager *et al.* 2016) and the evolutionary young fish species

found in this region appear to be somewhat resilient to river impoundment. The creation of new

lentic habitats upstream of the dam, captured by the TSI variable in our study, is the most

plausible driver of the shift in assemblages, but did not wipe out any species.

Dams can also block migratory route of diadromous species and alter seasonal migration

of potamodromous species. Local losses or reduction in abundance of migratory species has been

attributed to river fragmentation by dams in tropical and temperate regions (Reyes-Gavilán *et al.*

1996; Galat *et al.* 1998; Gehrke, Gilligan & Barwick 2002; Okada, Agostinho & Gomes 2005;

Sá-Oliveira *et al.* 2015; Pelicice, Pompeu & Agostinho 2015; Lima *et al.* 2016). In our boreal

systems, dams did not appear to be a major barrier to migration or movement for most species

because focal fishes were not diadromous and do not undertake long spawning migration (Table

S10). Moreover, the barrier effect might also have been minimized as most dams were built on

pre-existing obstacles that were already impassable for fish (*i.e.,* high waterfall). However,

studies of this nature should be pursued in the future because of the high occurrence of anadromy

in some boreal regions (McDowall 2008).

The effects associated with altering the natural hydrological regimes on fish communities depend

in part on reservoir morphometry and on the magnitude of the alteration that is related to

reservoir management (*e.g.*, magnitude of drawdown, discharge and hypo- vs. epilimnitic water

release). As noted in previous works, the magnitude of change in discharge and drawdown can

have divergent effects. For example, a 76% decrease in discharge in the Canadian River strongly

affected fish assemblages downstream of the Ute dam (Ute reservoir, New Mexico, USA), but a

36% decrease in discharge did not have significant effects downstream of Sanford dam along the

same river (Lake Meredith reservoir, Texas, USA; Bonner & Wilde 2000). In our boreal

ecosystems, despite the diversion of some rivers (decrease in discharge of up to 90%), only the

lake sturgeon in Opinaca appear to be really strongly affected (Figs 5 and 6).

Intentional (*e.g.*, fishing bait) or unintentional introduction (*e.g.*, flooding creates new

connection between water bodies) of non-native species in reservoirs can promote a shift away

from native-dominated fish communities (Rodriguez Ruiz 1998; Gido *et al.* 2002; Johnson,

Olden & Vander Zanden 2008; Clavero & Hermoso 2010). As an extreme example, the

introduction of a voracious non-native predator (the peacock-bass; *Cichla kelberi*) in Rosana

reservoir (Paraná river basin), decreased fish richness by 80% after only three years (Pelicice and

Agostinho 2008). River basins where endemic species are abundant might be particularly

vulnerable (Dudgeon *et al.* 2006). In our boreal reservoirs, no non-native species has been

observed, and no endemic species were present in either the LG or SM complexes. The remote

location of our focal reservoirs also likely contributed to the lack of establishment of non-native

fishes.

The time it takes for fish communities to stabilize after impoundment is highly variable

among studies. It has been reported to be either quick (*i.e.,* within five years; Martinez *et al.*

1994), or much longer (more than 10 years; Quinn & Kwak 2003; Říha *et al.* 2009), highlighting

the need for data span decades after impoundment. Some states or phases can also be transient.

Říha *et al.* (2009) documented a five-phase succession in fish species with European reservoirs

aging. The time needed for the fish assemblages to stabilize will depend on fish behavior, life

history trait and adaptability, the stability of the food web, the strength of trophic interactions,

and on the management and operation of the dam and reservoir. If the dominant mechanisms are

related to reproduction and recruitment through the strength of year classes, the effect may take

years to be detectable. Species with some specific life history traits (*e.g.*, late age at first

reproduction), or positioned at higher trophic levels may have delayed responses to

impoundment. If the dominant mechanisms are through movement and redistribution due to river

fragmentation and change in habitat quality, then shifts can be detected quickly.

4.2 Multi-scale approach and study design

Equipped with fish assemblage data collected over decades after impoundment, and

across a large spatial network of sites in a remote boreal region, this study is unique in providing

the most data-rich analysis to date, and in its ability to isolate the effect of impoundment from

other factors that co-occur with hydroelectricity projects. Great insights are achieved when

multiple scales are considered because patterns observed in communities at a given scale are

often the consequence of a complex interplay between various processes occurring at multiple

scales (Wiens 1989; Whittaker, Willis & Field 2001). In this study, changes in fish assemblages

in response to impoundment were only detectable at the sampling station scale (Figs. 3 and 4). At

the complex and reservoirs scales, fish assemblage shifts were largely masked by some other

larger scale ecological processes (*i.e.,* diverse habitat types and natural barriers to movement

leading to different fish communities), which highlights the importance of a multi-scale approach

to evaluate the potential of anthropogenic impacts on aquatic ecosystems.

Scale matters, but having different categories of stations, reference sites, and time series

covering the periods before and after impoundment are equally important considerations to

understand the effects of impoundment on fish communities. We found the strongest shifts in

species assemblages in upstream stations (Fig 6), relative to references and downstream stations,

which clearly points to the impact of impoundment vs. regional environmental change. Finally,

time series should cover the period before and after impoundment, and preferably time series

should be long enough to cover the non-equilibrium trophic surge and reach the new ecosystem

equilibrium (Grimard & Jones 1982; Turgeon *et al.* 2016).

5. Conclusions

By using a network of sites with minimal confounding factors, and by conducting our

analyses at three spatial scales, we have provided strong empirical evidence that impounding

large rivers in these boreal ecosystems did not affect diversity, but resulted in a clear shift in fish

assemblages. Changes in fish assemblages to impoundment were most clearly detected with our

ordination and beta-diversity analyses conducted at the scale of the sampling station. Given the

strength of our multi-scale approach in providing a complete perspective on the scale at which

river impoundment affect fish community, we caution against large scale extrapolations and

correlation studies that may underestimate or mask anthropogenic effects on aquatic ecosystems.

Reservoirs are now dominant features of many landscapes, and they will become even more

common in the coming years, especially in tropical regions (Zarfl *et al.* 2014; Winemiller *et al.*

2016). Identifying which mechanisms related to impoundment and river regulation affect

species, evaluating the strength of their effects, and how they vary across regions can assist in

implementing mitigation measures and in minimizing biodiversity loss.

Authors' Contributions

KT, CT and IGE conceived the idea, KT analysed the data; KT led the writing of the

manuscript. All authors contributed critically to the drafts and gave final approval for

publication.

Literature Cited

Agostinho, A.A., Julio Jr, H.F. & Petrere Jr., M. (1994) Itaipu Reservoir (Brazil) : Impacts of the

Impoundment on the Fish Fauna and Fisheries. *Rehabilitation of Freshwater Fisheries, by*

I.G. Cowx., pp. 171–184. Fishing news Books, Blackwell Scientific Publications, Osney

Mead, Oxford, UK.

Agostinho, A.A., Pelicice, F.M. & Gomes, L.C. (2008) Dams and the fish fauna of the

Neotropical region: impacts and management related to diversity and fisheries. *Brazilian*

Journal of Biology, **68**, 1119–1132.

Blanchet, F.G., Legendre, P. & Borcard, D. (2008) Forward selection of explanatory variables.

Ecology, **89**, 2623–2632.

Bonner, T.H. & Wilde, G.R. (2000) Changes in the Canadian River Fish Assemblage Associated

with Reservoir Construction. *Journal of Freshwater Ecology*, **15**, 189–198.

Burnham, K.P. & Anderson, D.R. (2002) *Model Selection and Multimodel Inference: A Practical*

Information-Theoretic Approach. Second Edition. Springer-Verlag, New York.

Carmignani, J.R. & Roy, A.H. (2017) Ecological impacts of winter water level drawdowns on

lake littoral zones: a review. *Aquatic Sciences*, 1–22.

Clavero, M. & Hermoso, V. (2010) Reservoirs promote the taxonomic homogenization of fish

communities within river basins. *Biodiversity and Conservation*, **20**, 41–57.

Deslandes, J.-C. & Fortin, R. (1994) *Optimisation Du Réseau de Suivi Environnemental Des*

Populations de Poissons Des Milieux Affectés Par l'aménagement Du Complexe La Grande

(Phase I).

Dudgeon, D., Arthington, A.H., Gessner, M.O., Kawabata, Z.-I., Knowler, D.J., Lévêque, C.,

Naiman, R.J., Prieur-Richard, A.-H., Soto, D., Stiassny, M.L.J. & Sullivan, C.A. (2006)

Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological*

Reviews, **81**, 163–182.

Elliott, J.M. (1990) The need for long-term investigations in ecology and the contribution of the

Freshwater Biological Association. *Freshwater Biology*, **23**, 1–5.

Fréchette, J.-L. (1980) *Réseau Surveillance Écologique. Vol. 1*. Société d'énergie de la Baie

James. Environnement.

Friedl, G. & Wüest, A. (2002) Disrupting biogeochemical cycles - Consequences of damming.

Aquatic Sciences, **64**, 55–65.

Galat, D.L., Fredrickson, L.H., Humburg, D.D., Bataille, K.J., Bodie, J.R., Dohrenwend, J.,

Gelwicks, G.T., Havel, J.E., Helmers, D.L., Hooker, J.B., Jones, J.R., Knowlton, M.F.,

Kubisiak, J., Mazourek, J., McColpin, A.C., Renken, R.B. & Semlitsch, R.D. (1998)

Flooding to Restore Connectivity of Regulated, Large-River Wetlands. *BioScience*, **48**, 721–

733.

Gehrke, P.C., Gilligan, D.M. & Barwick, M. (2002) Changes in fish communities of the

Shoalhaven River 20 years after construction of Tallowa Dam, Australia. *River Research and*

Applications, **18**, 265–286.

Gido, K.B., Guy, C.S., Strakosh, T.R., Bernot, R.J., Hase, K.J. & Shaw, M.A. (2002) Long-Term

Changes in the Fish Assemblages of the Big Blue River Basin 40 Years after the

Construction of Tuttle Creek Reservoir. *Transactions of the Kansas Academy of Science*,

105, 193–208.

Gido, K.B., Matthews, W.J. & Wolfinbarger, W.C. (2000) Long-term changes in a reservoir fish

assemblage: stability in an unpredictable environment. *Ecological Applications*, **10**, 1517–

1529.

Gomes, L. c. & Miranda, L. e. (2001) Riverine characteristics dictate composition of fish

assemblages and limit fisheries in reservoirs of the upper Paraná River basin. *Regulated*

Rivers: Research & Management, **17**, 67–76.

Grill, G., Lehner, B., Lumsdon, A.E., MacDonald, G.K., Zarfl, C. & Liermann, C.R. (2015) An

index-based framework for assessing patterns and trends in river fragmentation and flow

regulation by global dams at multiple scales. *Environmental Research Letters*, **10**, 015001.

Grimard, Y. & Jones, H.G. (1982) Trophic Upsurge in New Reservoirs: A Model for Total

Phosphorus Concentrations. *Canadian Journal of Fisheries and Aquatic Sciences*, **39**, 1473–

1483.

Gubiani, É.A., Gomes, L.C., Agostinho, A.A. & Baumgartner, G. (2010) Variations in fish

assemblages in a tributary of the upper Paraná River, Brazil: A comparison between pre and

post-closure phases of dams. *River Research and Applications*, **26**, 848–865.

Haxton, T.J. & Findlay, C.S. (2009) Variation in large-bodied fish-community structure and

abundance in relation to water-management regime in a large regulated river. *Journal of Fish*

Biology, **74**, 2216–2238.

International Energy Agency (IEA). (2016) *Key World Energy Statistics*. Paris, Cedex, France.

Johnson, P.T., Olden, J.D. & Vander Zanden, M.J. (2008) Dam invaders: impoundments

facilitate biological invasions into freshwaters. *Frontiers in Ecology and the Environment*, **6**,

357–363.

Kroger, R.L. (1973) Biological Effects of Fluctuating Water Levels in the Snake River, Grand

Teton National Park, Wyoming. *American Midland Naturalist*, **89**, 478–481.

Legendre, P. & De Cáceres, M. (2013) Beta diversity as the variance of community data:

dissimilarity coefficients and partitioning. *Ecology Letters*, **16**, 951–963.

Legendre, P. & Gauthier, O. (2014) Statistical methods for temporal and space–time analysis of

community composition data. *Proceedings of the Royal Society of London B: Biological*

Sciences, **281**, 20132728.

Li, B., Madden, L.V. & Xu, X. (2012) Spatial analysis by distance indices: an alternative local

clustering index for studying spatial patterns. *Methods in Ecology and Evolution*, **3**, 368–377.

Liermann, C.R., Nilsson, C., Robertson, J. & Ng, R.Y. (2012) Implications of Dam Obstruction

for Global Freshwater Fish Diversity. *BioScience*, **62**, 539–548.

Liew, J.H., Tan, H.H. & Yeo, D.C.J. (2016) Dammed rivers: impoundments facilitate fish

invasions. *Freshwater Biology*, **61**, 1421–1429.

Lima, A.C., Agostinho, C.S., Sayanda, D., Pelicice, F.M., Soares, A.M.V.M. & Monaghan, K.A.

(2016) The rise and fall of fish diversity in a neotropical river after impoundment.

Hydrobiologia, **763**, 207–221.

Martinez, P.J., Chart, T.E., Trammell, M.A., Wullschleger, J.G. & Bergersen, E.P. (1994) Fish

species composition before and after construction of a main stem reservoir on the White

River, Colorado. *Environmental Biology of Fishes*, **40**, 227–239.

McDowall, R.M. (2008) Why are so many boreal freshwater fishes anadromous? Confronting

'conventional wisdom'. *Fish and Fisheries*, **9**, 208–213.

de Mérona, B. de, Vigouroux, R. & Tejerina-Garro, F.L. (2005) Alteration of Fish Diversity

Downstream from Petit-Saut Dam in French Guiana. Implication of Ecological Strategies of

Fish Species. *Hydrobiologia*, **551**, 33–47.

Messager, M.L., Lehner, B., Grill, G., Nedeva, I. & Schmitt, O. (2016) Estimating the volume

and age of water stored in global lakes using a geo-statistical approach. *Nature*

Communications, **7**, 13603.

Nilsson, C., Reidy, C.A., Dynesius, M. & Revenga, C. (2005) Fragmentation and Flow

Regulation of the World's Large River Systems. *Science*, **308**, 405–408.

Okada, E.K., Agostinho, A.A. & Gomes, L.C. (2005) Spatial and temporal gradients in artisanal

fisheries of a large Neotropical reservoir, the Itaipu Reservoir, Brazil. *Canadian Journal of*

Fisheries and Aquatic Sciences, **62**, 714–724.

Oksanen, J., Blanchet, G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R.,

O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, H.H., Szoecs, E. & Wagner, H. (2016)

Vegan: Community Ecology Package. R Package Version 2.4-1.

Pelicice, F.M., Pompeu, P.S. & Agostinho, A.A. (2015) Large reservoirs as ecological barriers to

downstream movements of Neotropical migratory fish. *Fish and Fisheries*, **16**, 697–715.

Peres-Neto, P.R., Legendre, P., Dray, S. & Borcard, D. (2006) Variation Partitioning of Species

Data Matrices: Estimation and Comparison of Fractions. *Ecology*, **87**, 2614–2625.

Poff, N.L., Olden, J.D., Merritt, D.M. & Pepin, D.M. (2007) Homogenization of regional river

dynamics by dams and global biodiversity implications. *Proceedings of the National*

Academy of Sciences, **104**, 5732–5737.

Quinn, J.W. & Kwak, T.J. (2003) Fish Assemblage Changes in an Ozark River after

Impoundment: A Long-Term Perspective. *Transactions of the American Fisheries Society*,

132, 110–119.

Reyes-Gavilán, F.G., Garrido, R., Nicieza, A.G., Toledo, M.M. & Braña, F. (1996) Fish

community variation along physical gradients in short streams of northern Spain and the

disruptive effect of dams. *Hydrobiologia*, **321**, 155–163.

Říha, M., Kubečka, J., Vašek, M., Seda, J., Mrkvička, T., Prchalová, M., Matěna, J., Hladík, M.,

Čech, M., Draštík, V., Frouzová, J., Hohausová, E., Jarolím, O., Jůza, T., Kratochvíl, M.,

Peterka, J. & Tušer, M. (2009) Long-term development of fish populations in the Římov

Reservoir. *Fisheries Management and Ecology*, **16**, 121–129.

Rodriguez Ruiz, A. (1998) Fish species composition before and after construction of a reservoir

on the Guadalete River (SW Spain). *Archiv für Hydrobiologie*, **142**, 353–369.

Rosenberg, D.M., McCully, P. & Pringle, C.M. (2000) Global-Scale Environmental Effects of

Hydrological Alterations: Introduction. *BioScience*, **50**, 746–751.

Roy, D. & Messier, D. (1989) A review of the effects of water transfers in the La Grande

Hydroelectric Complex (Québec, Canada). *Regulated Rivers: Research & Management*, **4**,

299–316.

Sá-Oliveira, J.C., Hawes, J.E., Isaac-Nahum, V.J. & Peres, C.A. (2015) Upstream and

downstream responses of fish assemblages to an eastern Amazonian hydroelectric dam.

Freshwater Biology, **60**, 2037–2050.

Stoffer, D. (2014) astsa: Applied Statistical Time Series Analysis. R package version 1.3.

Sutela, T. & Vehanen, T. (2008) Effects of water-level regulation on the nearshore fish

community in boreal lakes. *Hydrobiologia*, **613**, 13–20.

Taylor, C.A., Knouft, J.H. & Hiland, T.M. (2001) Consequences of stream impoundment on fish

communities in a small North American drainage. *Regulated Rivers: Research &*

Management, **17**, 687–698.

Tereshchenko, V.G. & Strel'nikov, A.S. (1997) Analysis of long-term changes in the fish

community in Rybinsk reservoir. *Journal of Ichthyology*, **37**, 590–598.

Turgeon, K., Solomon, C.T., Nozais, C. & Gregory-Eaves, I. (2016) Do novel ecosystems follow

predictable trajectories? Testing the trophic surge hypothesis in reservoirs using fish.

Ecosphere, **7**, n/a-n/a.

Turgeon, K., Turpin, C. & Gregory-Eaves, I. (in prep.) Do dams affect fish biodiversity in

similar ways across biomes? A global quantitative synthesis. *in prep.*

Vörösmarty, C.J., McIntyre, P.B., Gessner, M.O., Dudgeon, D., Prusevich, A., Green, P.,

Glidden, S., Bunn, S.E., Sullivan, C.A., Liermann, C.R. & Davies, P.M. (2010) Global

threats to human water security and river biodiversity. *Nature*, **467**, 555–561.

Ward, J.V. & Stanford, J.A. (1995) The serial discontinuity concept: Extending the model to

floodplain rivers. *Regulated Rivers: Research & Management*, **10**, 159–168.

Whittaker, R.J., Willis, K.J. & Field, R. (2001) Scale and species richness: towards a general,

hierarchical theory of species diversity. *Journal of Biogeography*, **28**, 453–470.

Wiens, J.A. (1989) Spatial Scaling in Ecology. *Functional Ecology*, **3**, 385–397.

Winemiller, K.O., McIntyre, P.B., Castello, L., Fluet-Chouinard, E., Giarrizzo, T., Nam, S.,

Baird, I.G., Darwall, W., Lujan, N.K., Harrison, I., Stiassny, M.L.J., Silvano, R. a. M.,

Fitzgerald, D.B., Pelicice, F.M., Agostinho, A.A., Gomes, L.C., Albert, J.S., Baran, E.,

Petrere, M., Zarfl, C., Mulligan, M., Sullivan, J.P., Arantes, C.C., Sousa, L.M., Koning, A.A.,

Hoeinghaus, D.J., Sabaj, M., Lundberg, J.G., Armbruster, J., Thieme, M.L., Petry, P.,

Zuanon, J., Vilara, G.T., Snoeks, J., Ou, C., Rainboth, W., Pavanelli, C.S., Akama, A.,

Soesbergen, A. van & Sáenz, L. (2016) Balancing hydropower and biodiversity in the

Amazon, Congo, and Mekong. *Science*, **351**, 128–129.

Zarfl, C., Lumsdon, A.E., Berlekamp, J., Tydecks, L. & Tockner, K. (2014) A global boom in

hydropower dam construction. *Aquatic Sciences*, **77**, 161–170.

Table 1. Reservoirs characteristics in the La Grande and Sainte-Marguerite 3 hydroelectric

complexes. Reservoir area represents the surface covered with water at maximum pool. The

area flooded represents the surface of terrestrial land flooded following impoundment and, in

brackets, the percentage of the reservoir that was terrestrial land before impoundment. The

extrapolated richness, diversity and evenness values are averages calculated across all

upstream and downstream stations for all time points.

Variables	Reservoirs			
	Opinaca	Robert-Bourassa	Caniapiscau	Ste-Marguerite 3
Latitude	52°38′58″N	53°45'00"N	54°31'46"N	50°42′52″N
Longitude	76°19′54″W	77°00'00"W	69°51'18"W	66°46′54″W
Trophic status	Oligotrophic	Oligotrophic	Oligotrophic	Oligotrophic
Area (km^2)	1040	2835	4275	262
Area flooded (km^2)	740 (71%)	2630 (92%)	3430 (80%)	230.5 (88%)
Volume (km^3)	8.4	61.7	53.8	12.5
Year of impoundment	1980	1979	1982	1998
Filling time (y)	0.5	1	2	4
Residency time (days)	124	183	803	366
Mean depth (m)	8	22	12	22.4
Max depth (m)	51	137	49	145
Annual drawdown (m)	3.6	3.3	2.1	14
Watershed area (km^2)	30 000	97 643	36 800	9 000
Extrapolated richness	11.1 ± 3.7	9.9 ± 3.1	9.3 ± 2.7	7.5 ± 3.1
Diversity	1.53 ± 0.28	1.48 ± 0.27	1.40 ± 0.28	1.23 ± 0.38
Evenness	0.76 ± 0.13	0.78 ± 0.12	0.72 ± 0.13	0.87 ± 0.11

Table 2. Estimate ± Standard error (SE), 95% Confidence intervals, t-values and degrees of freedom (DF) of model parameters used to predict change in extrapolated richness, diversity and evenness in La Grande hydroelectric complex. General linear mixed effects models were used to evaluate the effect of time since impoundment, stations categories (Impacted stations vs. reference sites) and their interaction on diversity metrics. Predictors not including 0 within their 95% CI are in bold. Reference sites are used as contrasts in the models.

Model parameter	Extrapolated richness		Diversity (H')		Evenness (J')	
	Estimate ± SE (95% CI)	t-value (DF)	Estimate ± SE (95% CI)	t-value (DF)	Estimate ± SE (95% CI)	t-value (DF)
Model 1: All impacted stations combined						
Intercept	-0.073 ± 0.309 (-0.678 to 0.532)	-0.236 (171)	0.485 ± 0.248 (-0.001 to 0.972)	1.954 (171)	0.421 ± 0.292 (-0.151 to 0.993)	1.443 (171)
Time since impoundment (TSI)	0.042 ± 0.179 (-0.309 to 0.393)	0.233 (171)	-0.184 ± 0.218 (-0.610 to 0.243)	-0.844 (171)	-0.145 ± 0.191 (-0.520 to 0.231)	-0.755 (171)
Ref vs. Impacted (RI)	0.052 ± 0.331 (-0.596 to 0.701)	0.159 (21)	**-0.562 ± 0.266 (-1.085 to -0.040)**	**-2.111 (21)**	-0.505 ± 0.312 (-1.117 to 0.107)	-1.616 (21)
TSI*RI	-0.186 ± 0.194 (-0.566 to 0.193)	-0.962 (171)	0.108 ± 0.235 (-0.352 to 0.569)	0.461 (171)	0.206 ± 0.206 (-0.199 to 0.611)	0.998 (171)
Model 2: Upstream (Up) and downstream (D) stations separately						
Intercept	-0.069 ± 0.270 (-0.598 to 0.461)	-0.254 (170)	0.480 ± 0.250 (-0.010 to 0.969)	1.920 (170)	0.424 ± 0.286 (-0.136 to 0.984)	1.483 (170)
TSI	0.036 ± 0.175 (-0.308 to 0.380)	0.207 (170)	-0.176 ± 0.227 (-0.620 to 0.269)	-0.776 (170)	-0.147 ± 0.190 (-0.518 to 0.225)	-0.775 (170)
Ref vs. D	0.592 ± 0.350 (-0.094 to 1.277)	1.692 (20)	-0.388 ± 0.323 (-1.020 to 0.244)	-1.203 (20)	**-0.778 ± 0.370 (-1.502 to -0.053)**	**-2.104 (20)**
Ref vs. Up	-0.098 ± 0.295 (-0.676 to 0.480)	-0.333 (20)	**-0.609 ± 0.273 (-1.144 to -0.074)**	**-2.233 (20)**	-0.432 ± 0.311 (-1.043 to 0.178)	-1.388 (20)
TSI*(Ref. vs D)	-0.309 ± 0.226 (-0.752 to 0.134)	-1.366 (170)	0.239 ± 0.295 (-0.339 to 0.817)	0.811 (170)	0.316 ± 0.244 (-0.163 to 0.794)	1.294 (170)
TSI*(Ref. vs Up)	-0.149 ± 0.194 (-0.530 to 0.232)	-0.767 (170)	0.055 ± 0.250 (-0.435 to 0.545)	0.220 (170)	0.177 ± 0.209 (-0.233 to 0.588)	0.848 (170)

Model 3: Upstream river (UR), upstream lake (UL) and downstream (D) stations separately

Intercept	-0.071 ± 0.271	-0.261	0.488 ± 0.248	1.971	0.441 ± 0.247	1.783
	(-0.603 to 0.461)	(169)	(0.003 to 0.974)	(169)	(-0.044 to 0.926)	(169)
TSI	0.035 ± 0.178	0.200	-0.192 ± 0.206	-0.934	-0.164 ± 0.175	-0.936
	(-0.312 to 0.383)	(169)	(-0.595 to 0.211)	(169)	(-0.508 to 0.180)	(169)
Ref vs. D	0.593 ± 0.351	1.690	-0.401 ± 0.320	-1.254	**-0.792 ± 0.319**	**-2.484**
	(-0.095 to 1.282)	(19)	(-1.029 to 0.226)	(19)	**(-1.418 to -0.167)**	(19)
Ref vs. UL	-0.024 ± 0.308	-0.078	**-0.654 ± 0.282**	**-2.316**	**-0.655 ± 0.281**	**-2.329**
	(-0.628 to 0.580)	(19)	**(-1.208 to -0.101)**	(19)	**(-1.206 to -0.104)**	(19)
Ref vs. UR	-0.233 ± 0.338	-0.689	-0.544 ± 0.309	-1.762	-0.044 ± 0.307	-0.142
	(-0.895 to 0.429)	(19)	(-1.149 to 0.061)	(19)	(-0.646 to 0.559)	(19)
TSI*(Ref. vs D)	-0.309 ± 0.229	-1.351	0.249 ± 0.267	0.934	0.336 ± 0.226	1.483
	(-0.757 to 0.139)	(169)	(-0.274 to 0.773)	(169)	(-0.108 to 0.779)	(169)
TSI*(Ref. vs UL)	-0.120 ± 0.206	-0.581	0.233 ± 0.237	0.981	0.311 ± 0.204	1.524
	(-0.524 to 0.284)	(169)	(-0.232 to 0.698)	(169)	(-0.089 to 0.711)	(169)
TSI*(Ref. vs UR)	-0.202 ± 0.228	-0.884	-0.244 ± 0.265	-0.918	-0.030 ± 0.226	-0.133
	(-0.650 to 0.246)	(169)	(-0.764 to 0.276)	(169)	(-0.473 to 0.413)	(169)

Figure 1. Map representing the before (light blue) and after (dark blue) impoundment hydrological conditions, and the location of the sampling stations in the La Grande hydroelectricity complex from three reservoirs; RB, OP and CA and 4 sampling stations in Sainte-Marguerite complex, Northern Québec. Stations located upstream of the dams that were in a river before impoundment are represented by yellow circles, and the ones that were in lakes before impoundment are represented by red circles. Sampling stations that were located downstream of the dams are represented by a blue triangle, and reference sites paired with each reservoir are represented by green squares. Dams are represented by a black line and power station by a turbine symbol. Reservoirs that are not the focus of our study but in the region, (LG3, LG4, LaForge I & II) are presented and were impounded at the following dates: 1981, 1983, 1993 and 1983 respectively.

Figure 2. Variation in extrapolated richness, diversity (H') and evenness (J') over time in impacted and reference stations at the LG complex level. Changes in diversity metrics over time in references sites (green) were compared with impacted stations (all categories combined from all reservoirs) in a, d, g panels, with impacted stations upstream and downstream of the dams in b, e, h panels, and with upstream stations that were lakes before being a reservoir (UL) and those that were a river before (UR) in c, f, i panels.

Figure 3. Local contribution to beta-diversity (LCBD) per station and year at: a) the LG complex level and b) the sampling station level. Circle areas are proportional to the LCBD values. Circles filled in white indicate significant LCBD at p<0.05. The lower panels represent mean values of LCBD per year with a distance weights least square (DWLS) curve fit. The right panel represents mean values of LCBD per station for the analysis at the complex level. Stations with a label starting with "UL" represent stations that were lakes before being a reservoir and those with a label starting with "UR" were rivers or stream before being a reservoir. Reference sites are in green, downstream stations in blue and upstream stations in yellow UR) and red (UL).

Figure 4. a) Variation partitioning analysis showing the contribution of four matrices (Time since impoundment [TSI], Spatial heterogeneity [SH], Water quality variables [WQ] and fishing gear [G]) to explain the variation in fish species assemblages at the a) LG complex level, b) at the reservoir level (average across reservoirs, see Table S7 for the breakdown per reservoir), and c) at the sampling stations (average across sampling stations, see Table S8 for the breakdown). All analyses included reference sites.

Figure 5. Radar charts of species contributions to beta-diversity (SCBD) computed for each sampling stations, and pooled: a) per reservoirs, b) per categories of sampling stations, for c) Robert-Bourassa, d) Opinaca and e) Caniapiscau and f) Sainte-Marguerite. Only the ten most common species are pictured.

Figure 6. Changes in relative abundance over time of the most common species (>5% of total catch) in sampling stations of LG reservoirs and SM3 reservoir. Relative abundance in upstream stations for each reservoir are in a, b, c and d panels, in downstream stations in e, f and g panels and in reference sites in h, i, j k panels. The white dashed line represents the start of reservoir filling. See Table S9 for the name of the species related to the four letters code.

Table S1. **Analysis at the reservoir scale**. Estimate ± Standard error (SE), 95% Confidence intervals and t-values, and degrees of

freedom (DF) of model parameters used to predict change in extrapolated richness, diversity and evenness in La Grande complex

reservoirs and Ste-Marguerite 3 reservoir. Generalized mixed effects models were used to evaluate the effect of time since

impoundment, stations categories and their interaction on diversity metrics. Reference sites are used as contrasts.

Model parameter	Extrapolated richness		Diversity (H')		Evenness (J')	
	Estimate ± SE (95% CI)	t-value (DF)	Estimate ± SE (95% CI)	t-value (DF)	Estimate ± SE (95% CI)	t-value (DF)
Robert-Bourassa						
Intercept	-0.325 ± 0.500 (-1.305 to 0.655)	-0.650 (70)	0.380 ± 0.357 (-0.301 to 1.061)	1.066 (70)	0.609 ± 0.375 (-0.111 to 1.328)	1.622 (70)
TSI	-0.003 ± 0.270 (-0.531 to 0.526)	-0.011 (70)	-0.450 ± 0.245 (-0.929 to 0.032)	-1.835 (70)	-0.457 ± 0.254 (-0.955 to 0.041)	-1.798 (70)
Ref vs. Impacted (RI)	0.361 ± 0.535 (-0.689 to 1.410)	0.674 (6)	-0.430 ± 0.383 (-1.022 to 0.144)	-1.122 (6)	-0.683 ± 0.403 (-1.450 to 0.094)	-1.696 (6)
TSI*RI	-0.232 ± 0.296 (-0.812 to 0.348)	-0.784 (70)	0.099 ± 0.270 (-0.451 to 0.615)	0.365 (70)	0.351 ± 0.281 (-0.211 to 0.890)	1.250 (70)
Opinaca						
Intercept	0.058 ± 0.604 (-1.125 to 1.241)	0.096 (62)	0.539 ± 0.527 (-0.468 to 1.545)	1.023 (62)	0.362 ± 0.587 (-0.761 to 1.484)	0.616 (62)
TSI	0.011 ± 0.299 (-0.576 to 0.598)	0.037 (62)	-0.143 ± 0.293 (-0.716 to 0.430)	-0.490 (62)	0.022 ± 0.282 (-0.531 to 0.574)	0.076 (62)
Ref vs. Impacted (RI)	-0.061 ± 0.652 (-1.339 to 1.217)	-0.094 (5)	-0.628 ± 0.569 (-1.715 to 0.459)	-1.104 (5)	-0.431 ± 0.634 (-1.645 to 0.780)	-0.681 (5)
TSI*RI	-0.118 ± 0.323 (-0.751 to 0.515)	-0.366 (62)	0.200 ± 0.316 (-0.419 to 0.818)	0.632 (62)	0.156 ± 0.305 (-0.439 to 0.753)	0.511 (62)
Caniapiscau						
Intercept	-0.061 ± 0.514 (-1.068 to 0.945)	-0.120 (35)	0.346 ± 0.501 (-0.612 to 1.303)	0.690 (35)	0.214 ± 0.579 (-0.612 to 1.303)	0.369 (35)
TSI	0.559 ± 0.528 (-0.475 to 1.593)	1.060 (35)	0.532 ± 0.514 (-0.451 to 1.515)	1.034 (35)	0.347 ± 0.505 (-0.451 to 1.515)	0.688 (35)
Ref vs. Impacted (RI)	0.031 ± 0.538 (-1.024 to 1.086)	0.057 (6)	-0.410 ± 0.525 (-1.413 to 0.593)	-0.782 (6)	-0.244 ± 0.612 (-1.413 to 0.593)	-0.400 (6)
TSI*RI	-0.672 ± 0.552 (-1.753 to 0.410)	-1.217 (35)	-0.376 ± 0.538 (-1.405 to 0.652)	-0.700 (35)	-0.194 ± 0.529 (-1.405 to 0.652)	-0.366 (35)

Ste-Marguerite 3

Intercept	0.162 ± 0.659	0.876	-0.627 ± 0.585	-1.072	**-1.457 ± 0.398**	-3.660
	(-1.130 to 1.454)	(7)	(-1.676 to 0.422)	(7)	**(-2.164 to -0.751)**	(7)
TSI	-0.297 ± 0.569	0.117	0.063 ± 0.504	0.125	0.056 ± 0.343	0.164
	(-1.412 to 0.817)	(7)	(-0.840 to 0.967)	(7)	(-0.553 to 0.666)	(7)
Ref vs. Impacted (RI)	-0.199 ± 0.751	-0.979	0.785 ± 0.654	1.199	**1.863 ± 0.450**	4.144
	(-1.670 to 1.273)	(2)	(-0.388 to 1.957)	(2)	**(1.062 to 2.661)**	(2)
TSI*RI	0.302 ± 0.697	0.488	0.286 ± 0.598	0.478	-0.174 ± 0.411	-0.423
	(-1.063 to 1.667)	(7)	(-0.786 to 1.357)	(7)	(-0.905 to 0.557)	(7)

Table S2. **Sampling stations in Robert-Bourassa**. Estimate ± Standard error (SE), 95% Confidence intervals (95% CI) and t-values of model parameters used to predict change in extrapolated richness, diversity (Shannon's H') and evenness (Pielou J') in impacted stations in Robert-Bourassa reservoir when compared to reference site. General linear models were used to evaluate the effect of time, stations and their interaction on diversity metrics. The reference site is used as a contrast in the model.

Model parameter	Extrapolated richness		Diversity (H')		Evenness (J')	
	Estimate ± SE (95% CI)	t-value (DF)	Estimate ± SE (95% CI)	t-value (DF)	Estimate ± SE (95% CI)	t-value (DF)
Intercept	-0.325 ± 0.254 (-0.823 to 0.173)	-1.280 (1)	0.380 ± 0.282 (-0.172 to 0.932)	1.350 (1)	**0.609 ± 0.294** **(0.032 to 1.185)**	**2.069** **(1)**
Year	-0.003 ± 0.225 (-0.443 to 0.437)	-0.013 (1)	-0.450 ± 0.249 (-0.938 to 0.039)	-1.805 (1)	-0.457 ± 0.260 (-0.967 to 0.053)	-1.757 (1)
DORB	**0.844 ± 0.354** **(0.149 to 1.538)**	**2.381** **(7)**	-0.683 ± 0.393 (-1.454 to 0.087)	-1.738 (7)	**-1.381 ± 0.411** **(-2.186 to -0.576)**	**-3.364** **(7)**
UR1RB	-0.275 ± 0.372 (-1.004 to 0.455)	-0.738 (7)	**-0.886 ± 0.413** **(-1.695 to -0.077)**	**-2.148** **(7)**	-0.439 ± 0.431 (-1.283 to 0.406)	-1.018 (7)
UR3RB	0.631 ± 0.367 (-0.089 to 1.350)	1.718 (7)	-0.025 ± 0.407 (-0.823 to 0.773)	-0.061 (7)	-0.466 ± 0.425 (-1.299 to 0.367)	-1.096 (7)
UL1RB	**1.051 ± 0.367** **(0.331 to 1.770)**	**2.863** **(7)**	-0.115 ± 0.407 (-0.913 to 0.683)	-0.283 (7)	-0.778 ± 0.425 (-1.611 to 0.055)	-1.830 (7)
UL2RB	0.008 ± 0.367 (-0.712 to 0.727)	0.021 (7)	-0.467 ± 0.407 (-1.265 to 0.331)	-1.148 (7)	**-0.885 ± 0.425** **(-1.718 to -0.052)**	**-2.081** **(7)**
UL3RB	-0.048 ± 0.367 (-0.767 to 0.672)	-0.130 (7)	**-0.839 ± 0.407** **(-1.637 to -0.041)**	**-2.061** **(7)**	-0.524 ± 0.425 (-1.357 to 0.310)	-1.231 (7)
Year*DORB	-0.392 ± 0.309 (-0.998 to 0.214)	-1.268 (7)	0.246 ± 0.343 (-0.426 to 0.918)	0.717 (7)	0.576 ± 0.358 (-0.126 to 1.278)	1.607 (7)
Year*UR1RB	-0.089 ± 0.328 (-0.733 to 0.555)	-0.272 (7)	0.267 ± 0.364 (-0.447 to 0.981)	0.733 (7)	0.352 ± 0.380 (-0.394 to 1.098)	0.926 (7)
Year*UR3RB	-0.278 ± 0.370 (-1.003 to 0.447)	-0.752 (7)	-0.205 ± 0.410 (-1.009 to 0.600)	-0.498 (7)	0.192 ± 0.429 (-0.648 to 1.031)	0.447 (7)
Year*UL1RB	-0.366 ± 0.370 (-1.091 to 0.359)	-0.989 (7)	0.346 ± 0.410 (-0.458 to 1.151)	0.844 (7)	0.523 ± 0.429 (-0.317 to 1.363)	1.220 (7)
Year*UL2RB	0.034 ± 0.370 (-1.783 to 1.851)	0.092 (7)	0.460 ± 0.410 (-0.344 to 1.264)	1.121 (7)	0.647 ± 0.429 (-0.193 to 1.487)	1.510 (7)
Year*UL3RB	-0.060 ± 0.370 (-1.770 to 1.650)	-0.161 (7)	-0.297 ± 0.410 (-1.102 to 0.507)	-0.725 (7)	-0.015 ± 0.429 (-0.855 to 0.825)	-0.036 (7)

Table S3. **Sampling stations in Opinaca**. Estimate ± Standard error (SE), 95% Confidence intervals (95% CI) and t-values of model parameters used to predict change in extrapolated richness, diversity (Shannon's H') and evenness (Pielou J') in impacted stations in Opinaca reservoir when compared to reference site. General linear models were used to evaluate the effect of time, stations and their interaction on diversity metrics. The reference site is used as a contrast in the model.

Model parameter	Extrapolated richness		Diversity (H')		Evenness (J')	
	Estimate ± SE (95% CI)	t-value	Estimate ± SE (95% CI)	t-value	Estimate ± SE (95% CI)	t-value
Intercept	0.058 ± 0.292 (-0.515 to 0.631)	0.198 (1)	0.539 ± 0.286 (-0.021 to 1.098)	1.887 (1)	0.362 ± 0.280 (-0.188 to 0.911)	1.289 (1)
Year	0.011 ± 0.295 (-0.566 to 0.588)	0.037 (1)	-0.143 ± 0.288 (-0.707 to 0.420)	-0.498 (1)	0.022 ± 0.282 (-0.532 to 0.575)	0.076 (1)
DO1OP	0.291 ± 0.414 (-0.520 to 1.101)	0.703 (6)	0.237 ± 0.404 (-0.554 to 1.029)	0.587 (6)	0.131 ± 0.397 (-0.647 to 0.908)	0.329 (6)
DO2OP	0.650 ± 0.414 (-0.160 to 1.461)	1.572 (6)	-0.614 ± 0.404 (-1.406 to 0.177)	-1.522 (6)	**-0.962 ± 0.397 (-1.739 to -0.185)**	**-2.426 (6)**
UR1OP	-0.204 ± 0.414 (-1.015 to 0.607)	-0.493 (6)	-0.552 ± 0.404 (-1.343 to 0.239)	-1.367 (6)	-0.340 ± 0.397 (-1.117 to 0.437)	-0.857 (6)
UR2OP	**-0.846 ± 0.414 (-1.657 to -0.036)**	**-2.046 (6)**	**-1.398 ± 0.404 (-2.189 to -0.606)**	**-3.462 (6)**	**-0.806 ± 0.397 (-1.583 to -0.028)**	**-2.031 (6)**
UL1OP	0.394 ± 0.414 (-0.417 to 1.204)	0.952 (6)	**-0.857 ± 0.404 (-1.649 to -0.066)**	**-2.123 (6)**	**-0.992 ± 0.397 (-1.769 to -0.214)**	**-2.500 (6)**
UL2OP	-0.634 ± 0.405 (-1.427 to 0.160)	-1.566 (6)	-0.552 ± 0.395 (-1.327 to 0.222)	-1.398 (6)	0.377 ± 0.388 (-0.383 to 1.138)	0.973 (6)
Year*DO1OP	0.166 ± 0.417 (-0.650 to 0.983)	0.399 (6)	0.431 ± 0.407 (-0.366 to 1.228)	1.060 (57)	0.196 ± 0.399 (-0.587 to 0.979)	0.492 (6)
Year*DO2OP	-0.454 ± 0.417 (-1.271 to 0.362)	-1.091 (6)	0.744 ± 0.407 (-0.054 to 1.541)	1.828 (6)	0.629 ± 0.399 (-0.154 to 1.412)	1.576 (6)
Year*UR1OP	-0.120 ± 0.417 (-0.936 to 0.697)	-0.288 (6)	-0.064 ± 0.407 (-0.862 to 0.733)	-0.158 (6)	-0.211 ± 0.399 (-0.994 to 0.572)	-0.528 (6)
Year*UR2OP	-0.015 ± 0.417 (-0.831 to 0.801)	-0.036 (6)	0.035 ± 0.407 (-0.762 to 0.832)	0.086 (6)	0.101 ± 0.399 (-0.682 to 0.883)	0.252 (6)
Year*UL1OP	-0.180 ± 0.417 (-0.996 to 0.636)	-0.432 (6)	0.242 ± 0.407 (-0.555 to 1.039)	0.595 (6)	0.220 ± 0.399 (-0.563 to 1.003)	0.550 (6)
Year*UL2OP	-0.090 ± 0.407 (-0.889 to 0.708)	-0.221 (6)	-0.158 ± 0.398 (-0.938 to 0.621)	-0.398 (6)	-0.003 ± 0.391 (-0.769 to 0.763)	-0.008 (6)

Table S4. **Sampling stations in Caniapiscau**. Estimate ± Standard error (SE), 95% Confidence intervals (95% CI) and t-values of model parameters used to predict change in extrapolated richness, diversity (Shannon's H') and evenness (Pielou J') in impacted stations in Caniapiscau reservoir when compared to reference site. General linear models were used to evaluate the effect of time, stations and their interaction on diversity metrics. The reference site is used as a contrast in the model.

Model parameter	Extrapolated richness		Diversity (H')		Evenness (J')	
	Estimate ± SE (95% CI)	t-value	Estimate ± SE (95% CI)	t-value	Estimate ± SE (95% CI)	t-value
Intercept	-0.061 ± 0.666 (-1.366 to 1.243)	-0.092 (1)	0.425 ± 0.659 (-0.866 to 1.716)	0.646 (1)	0.234 ± 0.485 (-0.716 to 1.185)	0.483 (1)
Year	0.559 ± 0.679 (-0.773 to 1.891)	0.823 (1)	0.503 ± 0.691 (-0.852 to 1.858)	0.728 (1)	0.327 ± 0.522 (-0.696 to 1.351)	0.627 (1)
UL6CA	-0.222 ± 0.638 (-1.473 to 1.029)	-0.348 (7)	-0.447 ± 0.631 (-1.684 to 0.789)	-0.709 (7)	-0.373 ± 0.618 (-1.585 to 0.839)	-0.603 (7)
UL5CA	-0.357 ± 0.638 (-1.608 to 0.894)	-0.559 (7)	-0.352 ± 0.631 (-1.589 to 0.884)	-0.559 (7)	0.581 ± 0.618 (-0.631 to 1.793)	0.939 (7)
UL2CA	0.189 ± 0.638 (-1.062 to 1.441)	0.296 (7)	-0.634 ± 0.631 (-1.870 to 0.603)	-1.004 (7)	-0.606 ± 0.618 (-1.818 to 0.606)	-0.980 (7)
UL1CA	-0.522 ± 0.721 (-1.935 to 0.891)	-0.724 (7)	-0.517 ± 0.693 (-1.876 to 0.843)	-0.745 (7)	-0.220 ± 0.680 (-1.552 to 1.112)	-0.323 (7)
DOCA	0.332 ± 0.630 (-0.903 to 1.567)	0.527 (7)	-0.640 ± 0.630 (-1.876 to 0.595)	-1.016 (7)	-0.742 ± 0.618 (-1.953 to 0.469)	-1.201 (7)
Year*UL6CA	-0.314 ± 0.651 (-1.591 to 0.963)	-0.482 (7)	-0.262 ± 0.661 (-1.557 to 1.033)	-0.396 (7)	-0.003 ± 0.647 (-1.272 to 1.266)	-0.005 (7)
Year*UL5CA	-0.564 ± 0.651 (-1.841 to 0.713)	-0.865 (7)	0.321 ± 0.661 (-0.973 to 1.616)	0.487 (7)	-0.092 ± 0.647 (-1.361 to 1.177)	-0.142 (7)
Year*UL2CA	-0.678 ± 0.651 (-1.954 to 0.599)	-1.040 (7)	-0.314 ± 0.661 (-1.609 to 0.981)	-0.475 (7)	-0.001 ± 0.647 (-1.271 to 1.268)	-0.002 (7)
Year*UL1CA	-0.180 ± 0.708 (-1.568 to 1.209)	-0.253 (7)	-0.383 ± 0.718 (-1.791 to 1.025)	-0.533 (7)	-0.327 ± 0.704 (-1.707 to 1.053)	-0.465 (7)
Year*DOCA	-0.923 ± 0.659 (-2.215 to 0.368)	-1.401 (7)	-0.805 ± 0.668 (-2.114 to 0.505)	-1.204 (7)	-0.571 ± 0.655 (-1.855 to 0.713)	-0.872 (7)

Table S5. **Sampling stations in Sainte-Marguerite**. Estimate ± Standard error (SE), 95% Confidence intervals (95% CI) and t-values of model parameters used to predict change in extrapolated richness, diversity (Shannon's H') and evenness (Pielou J') in impacted stations in Sainte-Marguerite reservoir when compared to reference site. Generalized mixed effects models were used to evaluate the effect of time, stations and their interaction on diversity metrics. The reference site is used as a contrast in the model.

Model parameter	Extrapolated richness		Diversity (H')		Evenness (J')	
	Estimate ± SE (95% CI)	t-value	Estimate ± SE (95% CI)	t-value	Estimate ± SE (95% CI)	t-value
Intercept	0.162 ± 0.843 (-0.215 to 0.538)	0.192 (1)	-0.627 ± 0.696 (-1.991 to 0.737)	-0.901 (1)	-1.457 ± 0.425 (-2.289 to -0.625)	-3.433 (1)
Year	-0.297 ± 0.727 (0.504 to -1.099)	-0.409 (1)	0.063 ± 0.600 (-1.112 to 1.238)	0.105 (1)	0.056 ± 0.366 (-0.660 to 0.773)	0.154 (1)
UR3SM	-0.443 ± 1.661 (0.081 to -0.966)	-0.267 (3)	0.746 ± 0.888 (-0.994 to 2.486)	0.840 (3)	1.767 ± 0.562 (0.666 to 2.868)	3.145 (3)
UR2SM	-0.224 ± 1.226 (0.135 to -0.582)	-0.183 (3)	0.698 ± 0.997 (-1.256 to 2.652)	0.700 (3)	1.565 ± 0.608 (0.373 to 2.757)	2.574 (3)
UR1SM	0.155 ± 1.129 (-0.114 to 0.423)	0.137 (3)	0.998 ± 0.923 (-0.812 to 2.807)	1.081 (3)	2.177 ± 0.563 (1.073 to 3.281)	3.864 (3)
Year* UR3SM	0.417 ± 1.726 (-0.057 to 0.891)	0.242 (3)	0.522 ± 0.824 (-1.092 to 2.136)	0.634 (3)	-0.080 ± 0.516 (-1.091 to 0.931)	-0.155 (3)
Year* UR2SM	0.320 ± 1.184 (-0.209 to 0.849)	0.270 (3)	0.355 ± 0.977 (-1.560 to 2.269)	0.363 (3)	0.207 ± 0.596 (-0.961 to 1.375)	0.348 (3)
Year* UR1SM	-0.122 ± 1.163 (0.084 to -0.328)	-0.105 (3)	-0.237 ± 0.960 (-2.118 to 1.644)	-0.247 (3)	-0.677 ± 0.585 (-1.824 to 0.471)	-1.156 (3)

Table S6. Estimate ± Standard error (SE), 95% Confidence intervals (95% CI), t-values and Degrees of Freedom (DF) of model

parameters used to predict change in extrapolated richness (Double jackknife estimation method), diversity (Shannon's H') and

evenness (Pielou's J') in La Grande mega-hydroelectricity complex (3 reservoirs, 22 stations). General linear mixed effects models

were used to evaluate the additive effect of time since impoundment (TSI) and categories of impacted stations (D vs. Up, or D vs. UL,

vs. UR) on diversity metrics. Predictors that did not include 0 within their 95% CI (*i.e.*, statistically "significant") are in bold.

Downstream stations are used as contrasts in the models.

Model parameter	Extrapolated richness		Diversity (Shannon's H')		Evenness (Pielou's J')	
	Estimate ± SE (95% CI)	t-value (DF)	Estimate ± SE (95% CI)	t-value (DF)	Estimate ± SE (95% CI)	t-value (DF)
Model 1: All impacted stations combined						
Intercept	-0.020 ± 0.120 (-0.256 to 0.215)	-0.170 (149)	-0.076 ± 0.100 (-0.272 to 0.120)	-0.764 (149)	-0.083 ± 0.114 (-0.307 to 0.141)	-0.729 (149)
TSI	-0.145 ± 0.075 (-0.293 to 0.003)	-1.920 (149)	-0.075 ± 0.089 (-0.248 to 0.099)	-0.841 (149)	0.062 ± 0.078 (-0.090 to 0.215)	0.801 (149)
Model 2: Upstream (U) and downstream (D) stations separately						
Intercept	**0.493 ± 0.218 (0.065 to 0.921)**	**2.259 (149)**	0.089 ± 0.212 (-0.327 to 0.504)	0.418 (149)	-0.311 ± 0.233 (-0.767 to 0.145)	-1.336 (149)
TSI	**-0.155 ± 0.074 (-0.300 to -0.010)**	**-2.093 (149)**	-0.077 ± 0.090 (-0.252 to 0.099)	-0.853 (149)	0.066 ± 0.077 (-0.086 to 0.217)	0.850 (149)
D vs. U	**-0.653 ± 0.247 (-1.136 to -0.169)**	**-2.645 (18)**	-0.216 ± 0.241 (-0.688 to 0.256)	-0.896 (18)	0.293 ± 0.262 (-0.220 to 0.807)	1.120 (18)
Model 3: Upstream river (UR), upstream lake (UL) and downstream (D) stations separately						
Intercept	**0.489 ± 0.218 (0.061 to 0.917)**	**2.240 (149)**	0.073 ± 0.218 (-0.355 to 0.501)	0.335 (149)	-0.341 ± 0.205 (-0.742 to 0.060)	-1.665 (149)
TSI	**-0.155 ± 0.074 (-0.301 to -0.009)**	**-2.081 (149)**	-0.071 ± 0.090 (-0.247 to 0.104)	-0.795 (149)	0.066 ± 0.074 (-0.080 to 0.212)	0.886 (149)
D vs. UL	**-0.571 ± 0.261 (-1.082 to -0.059)**	**-2.187 (17)**	-0.253 ± 0.262 (-0.766 to 0.260)	-0.965 (17)	0.131 ± 0.246 (-0.352 to 0.613)	0.532 (17)
D vs. UR	**-0.796 ± 0.294 (-1.372 to -0.219)**	**-2.707 (17)**	-0.076 ± 0.294 (-0.651 to 0.500)	-0.257 (17)	**0.728 ± 0.276 (0.187 to 1.269)**	**2.636 (17)**

Table S7. Estimate ± Standard error (SE), 95% Confidence intervals (95% CI) and t-values of model parameters used to predict change in extrapolated richness (Double jackknife estimation), diversity (Shannon-Weaver H') and evenness (Pielou J') in La Grande complex reservoirs and Ste-Marguerite 3 reservoirs. General linear mixed effects models were used to evaluate the additive effect of time since impoundment (TSI) and categories of impacted stations (D vs. Up, or D vs. UL, vs. UR) on diversity metrics. Predictors that did not include 0 within their 95% CI (*i.e.*, statistically "significant") are in bold. Downstream stations are used as contrasts in the models.

Model parameter	Extrapolated richness		Diversity (H')		Evenness (J')	
	Estimate ± SE (95% CI)	t-value (DF)	Estimate ± SE (95% CI)	t-value (DF)	Estimate ± SE (95% CI)	t-value (DF)
Robert-Bourassa						
Model 1: Effect of time (upstream and downstream stations combined)						
Intercept	0.039 ± 0.224 (-0.399 to 0.478)	0.176 (55)	-0.114 ± 0.143 (-0.395 to 0.167)	-0.796 (55)	-0.142 ± 0.160 (-0.456 to 0.171)	-0.888 (55)
Year	-0.213 ± 0.122 (-0.453 to 0.027)	-1.737 (55)	**-0.316 ± 0.120 (-0.551 to -0.080)**	-2.626 (55)	-0.078 ± 0.130 (-0.333 to 0.177)	-0.600 (55)
Model 2: Upstream (U) and downstream (D) stations separately						
Intercept	0.327 ± 0.487 (-0.627 to 1.280)	0.671 (55)	-0.274 ± 0.368 (-0.995 to 0.447)	-0.746 (55)	**-0.730 ± 0.276 (-1.271 to -0.188)**	-2.641 (55)
Year	-0.223 ± 0.120 (-0.457 to 0.011)	-1.865 (55)	**-0.305 ± 0.120 (-0.540 to -0.070)**	-2.543 (55)	-0.046 ± 0.122 (-0.285 to 0.193)	-0.378 (55)
D. vs. U	-0.350 ± 0.527 (-1.382 to 0.682)	-0.664 (4)	0.198 ± 0.406 (-0.598 to 0.995)	0.488 (4)	**0.722 ± 0.307 (0.119 to 1.324)**	**2.348** (4)
Model 3: Upstream river (UR), upstream lake (UL) and downstream (D) stations separately						
Intercept	0.367 ± 0.397 (-0.411 to 1.145)	0.925 (55)	-0.274 ± 0.271 (-0.804 to 0.257)	-1.011 (55)	-0.727 ± 0.277 (-1.270 to -0.185)	-2.628 (55)
Year	**-0.213 ± 0.118 (-0.444 to -0.018)**	**-1.806** (55)	**-0.294 ± 0.124 (-0.536 to -0.052)**	**-2.378** (55)	-0.055 ± 0.123 (-0.295 to 0.186)	-0.445 (55)
D. vs. UL	-0.127 ± 0.464	-0.275	0.451 ± 0.321	1.402	0.622 ± 0.328	1.897

	(-1.037 to 0.782)	(3)	(-0.179 to 1.080)	(3)	(-0.021 to 1.265)	(3)
D. vs. UR	-0.795 ± 0.487	-1.632	-0.187 ± 0.340	-0.550	**0.865 ± 0.348**	**2.482**
	(-1.749 to 0.159)	(3)	(-0.854 to 0.480)	(3)	**(0.182 to 1.548)**	(3)

Opinaca

Model 4: Effect of time (upstream and downstream stations combined)

Intercept	-0.003 ± 0.246	-0.012	-0.088 ± 0.215	-0.407	-0.067 ± 0.241	-0.278
	(-0.485 to 0.479)	(54)	(-0.509 to 0.334)	(54)	(-0.540 to 0.405)	(54)
Year	-0.107 ± 0.121	-0.884	0.058 ± 0.139	0.417	0.180 ± 0.131	1.373
	(-0.345 to 0.131)	(54)	(-0.214 to 0.330)	(54)	(-0.077 to 0.437)	(54)

Model 5: Upstream (U) and downstream (D) stations separately

Intercept	0.519 ± 0.341	1.519	0.594 ± 0.324	1.837	0.181 ± 0.442	0.408
	(-0.150 to 1.188)	(53)	(-0.040 to 1.229)	(53)	(-0.687 to 1.048)	(53)
Year	-0.107 ± 0.121	-0.877	0.060 ± 0.151	0.396	0.181 ± 0.137	1.318
	(-0.345 to 0.131)	(53)	(-0.237 to 0.357)	(53)	(-0.088 to 0.449)	(53)
D. vs. U	-0.781 ± 0.413	-1.890	**-1.020 ± 0.376**	**-2.709**	-0.371 ± 0.517	-0.717
	(-1.592 to 0.029)	(4)	**(-1.758 to -0.282)**	**(4)**	(-1.385 to 0.643)	(4)

Model 6: Upstream river (UR), upstream lake (UL) and downstream (D) stations separately

Intercept	0.517 ± 0.382	1.355	0.532 ± 0.295	1.806	0.092 ± 0.373	0.247
	(-0.231 to 1.265)	(52)	(-0.046 to 1.110)	(52)	(-0.639 to 0.824)	(52)
Year	-0.105 ± 0.121	-0.873	0.057 ± 0.144	0.396	0.178 ± 0.129	1.379
	(-0.342 to 0.131)	(52)	(-0.226 to 0.340)	(52)	(-0.075 to 0.431)	(52)
D. vs. UL	-0.678 ± 0.535	-1.266	**-1.150 ± 0.406**	**-2.835**	-0.627 ± 0.517	-1.213
	(-1.727 to 0.371)	(3)	**(-1.945 to -0.355)**	**(3)**	(-1.641 to 0.387)	(3)
D. vs. UR	-0.878 ± 0.532	-1.650	-0.712 ± 0.400	-1.780	0.143 ± 0.513	0.279
	(-1.920 to 0.165)	(3)	(-1.496 to 0.072)	(3)	(-0.862 to 1.148)	(3)

Caniapiscau

Model 7: Effect of time (upstream and downstream stations combined)

Intercept	-0.057 ± 0.188	-0.304	-0.076 ± 0.171	-0.445	-0.038 ± 0.224	-0.171
	(-0.427 to 0.312)	(32)	(-0.412 to 0.260)	(32)	(-0.478 to 0.401)	(32)
Year	-0.032 ± 0.185	-0.175	0.219 ± 0.180	1.217	0.153 ± 0.176	0.871
	(-0.394 to 0.330)	(32)	(-0.134 to 0.571)	(32)	(-0.191 to 0.497)	(32)

Model 8: Upstream (U) and downstream (D) stations separately

Intercept	0.427 ± 0.363	1.176	-0.039 ± 0.369	-0.106	-0.423 ± 0.523	-0.808
	(-0.285 to 1.139)	(31)	(-0.762 to 0.684)	(31)	(-1.448 to 0.603)	(31)
Year	-0.007 ± 0.174	-0.043	0.222 ± 0.181	1.223	0.135 ± 0.179	0.754
	(-0.349 to 0.335)	(31)	(-0.134 to 0.578)	(31)	(-0.215 to 0.485)	(31)
D. vs. U	-0.603 ± 0.413	-1.459	-0.047 ± 0.419	-0.111	0.488 ± 0.589	0.827

	(-1.413 to 0.207)	(5)	(-0.869 to 0.775)	(5)	(-0.668 to 1.643)	(5)

Ste-Marguerite 3
Model 9: Effect of time (upstream and downstream stations combined)

Intercept	-0.037 ± 0.379	-0.097	0.158 ± 0.306	0.514	0.397 ± 0.228	1.740
	(-0.780 to 0.706)		(-0.443 to 0.758)		(-0.050 to 0.845)	
Year	0.005 ± 0.425	0.011	0.349 ± 0.336	1.037	-0.125 ± 0.267	-0.468
	(-0.828 to 0.837)		(-0.310 to 1.008)		(-0.648 to 0.398)	

Figure S1. Local contribution to beta-diversity (LCBD) at the reservoir level in SM complex.
LCBD values indicate the extent to which each local community is unique in terms of its species composition. Circle surface areas are proportional to the LCBD values. Circles filled in white indicate significant LCBD indices at p<0.05. The upper panel represents mean values of LCBD per year and the right panel represents mean values of LCBD per station. Reference sites are labelled in green, and upstream stations in orange.

Figure S2. Local contribution to beta-diversity (LCBD) at the reservoir level in LG complex.
LCBD values indicate the extent to which each local community is unique in terms of its species composition. Circle surface areas are proportional to the LCBD values. Circles filled in white indicate significant LCBD indices at p<0.05. The upper panel represents mean values of LCBD per year and the right panel represents mean values of LCBD per station. Reference sites are labelled in green, downstream stations in blue and upstream stations in orange and red. Upstream stations are separated in two categories. Stations with a label starting with "UL" represent stations that were lakes before being a reservoir (orange) and the stations with a label starting with "UR" were rivers or stream before being a reservoir (red).

Table S8. Variation partitioning results at the reservoir level. Variation explained (adjusted R^2 statistics) by the Spatial heterogeneity [SH] Time since impoundment [TSI], Water quality variables [WQV], Fishing gear [G] matrices, their shared fractions and residual variation. Results are presented for analyses including and excluding reference sites (With; Without). For the significant variables in each matrix, see Fig. 4.

Partitions	Robert-Bourassa	Opinaca	Caniapiscau	Mean
SH	**0.12; 0.15**	**0.15; 0.19**	**0.15; 0.08**	**0.14; 0.14**
TSI	0.03; 0.01	0.05; 0.04	0.12; 0.15	0.07; 0.07
WQV	0.03; 0.01	0.03; 0.04	0.07; 0.15	0.04; 0.07
G	0.04; 0.07	0.06; 0.08	0.07; 0.07	0.06; 0.07
SH+TSI	0.00; 0.00	0.04; 0.03	0.03; 0.03	0.02; 0.02
SH+WQV	**0.19; 0.13**	**0.27; 0.27**	**0.16; 0.12**	**0.21; 0.17**
SH+G	0.00; 0.00	0.00; 0.00	0.00; 0.00	0.00; 0.00
TSI+WQV	0.01; 0.06	0.00; 0.02	0.00; 0.00	0.00; 0.03
TSI+G	0.00; 0.00	0.00; 0.00	0.00; 0.00	0.00; 0.00
SH+TSI+WQV	0.00; 0.00	0.00; 0.00	0.00; 0.00	0.00; 0.00
Residuals	0.59; 0.54	0.48; 0.41	0.48; 0.49	0.52; 0.48
Total SH	**0.31; 0.28**	**0.46; 0.49**	**0.34; 0.23**	**0.37; 0.33**
Total TSI	0.04; 0.07	0.09; 0.09	0.15; 0.18	0.09; 0.11
Total WQV	0.23; 0.25	0.30; 0.31	0.23; 0.22	0.25; 0.26
Total G	0.04; 0.07	0.06; 0.08	0.07; 0.07	0.06; 0.07

Table S9. Variation partitioning results at the station level in the LG complex. Variation explained (adjusted R^2 statistics) by the Spatial heterogeneity [SH] Time since impoundment [TSI], Water quality variables [WQV], Fishing gear [G] matrices, their shared contributions and residual variation. Station are listed by their type (D = downstream, R = reference stations, UR = upstream station that was a river or a stream before impoundment and UL = upstream station that was a lake before impoundment).

Reservoirs	Stations	Type	TSI	WQV	G	TSI+WQV	TSI+G	WQV+G	TSI+WQV+G	Residuals
RB	DORB	D	0.06	0.00	0.11	**0.13**	0.00	0.00	0.00	0.76
RB	REFRB	R	0.00	0.00	0.00	**0.02**	0.00	0.00	0.01	0.99
RB	UR1RB	UR	0.03	0.00	0.09	**0.30**	0.00	0.00	0.00	0.62
RB	UR2RB	UR	0.02	0.00	0.12	**0.23**	0.00	0.00	0.00	0.69
RB	UL1RB	UL	0.00	0.00	0.02	**0.17**	0.00	0.00	0.00	0.83
RB	UL2RB	UL	0.00	0.00	0.02	**0.50**	0.00	0.00	0.00	0.52
RB	UL3RB	UL	0.00	0.00	0.00	**0.25**	0.00	0.00	0.00	0.77
RB	UR3RB	UR	0.00	0.00	0.02	**0.42**	0.00	0.00	0.00	0.60
OP	REFOP	R	0.00	0.00	**0.02**	0.00	0.00	0.00	0.00	0.98
OP	DO2OP	D	0.03	0.00	0.03	**0.17**	0.00	0.00	0.00	0.80
OP	DO1OP	D	0.00	0.00	0.00	**0.00**	0.01	0.00	0.01	0.99
OP	UR2OP	UR	0.00	0.00	0.00	**0.17**	0.01	0.00	0.00	0.90
OP	UL1OP	UL	0.00	0.00	0.00	**0.18**	0.00	0.00	0.00	0.83
OP	UL2OP	UL	0.20	0.00	0.00	**0.42**	0.00	0.00	0.00	0.46
OP	UR1OP	UR	**0.29**	0.00	0.00	0.10	0.00	0.00	0.00	0.73
CA	DOCA	D	0.01	0.00	0.23	**0.34**	0.00	0.00	0.00	0.54
CA	REFCA	R	0.00	0.00	**0.37**	0.25	0.00	0.00	0.00	0.52
CA	UL6CA	UL	0.01	0.00	0.09	**0.31**	0.00	0.00	0.00	0.65
CA	UL5CA	UL	0.00	0.00	0.18	**0.29**	0.00	0.00	0.00	0.61
CA	UL2CA	UL	0.00	0.00	0.03	**0.51**	0.00	0.00	0.00	0.52
CA	UL1CA	UL	0.00	0.00	0.08	**0.24**	0.00	0.00	0.00	0.74
Mean for reference sites			0.00 ± 0.00	0.00 ± 0.00	**0.13 ± 0.21**	0.09 ± 0.14	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.01	0.83 ± 0.27
Mean for upstream stations			0.04 ± 0.09	0.00 ± 0.00	0.05 ± 0.06	**0.29 ± 0.13**	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.68 ± 0.13
Mean for downstream stations			0.03 ± 0.03	0.00 ± 0.00	0.09 ± 0.10	**0.16 ± 0.14**	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.01	0.77 ± 0.18
Mean for impacted stations			0.04 ± 0.08	0.00 ± 0.00	0.06 ± 0.07	**0.26 ± 0.14**	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.70 ± 0.14
Total mean			0.03 ± 0.07	0.00 ± 0.00	0.07 ± 0.09	**0.24 ± 0.16**	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.72 ± 0.26

Table S10. Fish species information. Fish species name (latin name (CODE) and vernacular), feeding habits, trophic level, migratory behavior and life history trait characteristics. Information have been mostly extracted from Fishbase (trophic level and feeding), Olden et al. (2006) and Mims & Olden (2013) (life history trait) and from Desroches and Picard (2013; feeding and migratory behavior).

Species latin name (CODE)	Vernacular name	Feeding	Trophic level	Migratory behavior	Life history trait*
Catostomus catostomus (CACA)	Longnose sucker	Benthivore	2.5 ± 0.3	Potadromous	Periodic
Catostomus commersonii (CACO)	White sucker	Benthivore	2.8 ± 0.2	Potadromous	Periodic
Acipenser fulvescens (ACFU)	Lake sturgeon	Benthivore	3.3 ± 0.5	Potadromous	Periodic
Coregonus artedi (COAR)	Cisco	Planktivore	3.4 ± 0.4	Potadromous	Periodic
Coregonus clupeaformis (COCL)	Whitefish	Planktivore	3.2 ± 0.2	Potadromous	Periodic
Prosopium cylindraceum (PRCY)	Round whitefish	Benthivore	3.3 ± 0.4	Potadromous	Periodic
Esox lucius (ESLU)	Northern pike	Piscivore	4.1 ± 0.4	Resident	Periodic
Lota lota (LOLO)	Burbot	Piscivore	3.8 ± 0.2	Potadromous	Periodic
Sander vitreus (SAVI)	Walleye	Piscivore	4.5 ± 0.0	Potadromous	Periodic
Perca flavescens (PEFL)	Yellow perch	Omnivore	3.7 ± 0.2	Resident	Periodic
Salvelinus fontinalis (SAFO)	Brook trout	Benthivore	3.3 ± 0.0	Resident/anadromous	Equilibrium
Salvelinus namaycush (SANA)	Lake trout	Piscivore	4.3 ± 0.5	Potadromous	Equilibrium
Semotilus atromaculatus (SEMA)	Creek chub	Benthivore	4.0 ± 0.5	Potadromous	Opportunistic

Percopsis omiscomaycus (PEOM)	Trout-perch	Omnivore	3.4 ± 0.5	Potadromous	Periodic
Notropis hudsonis (NOHU)	Spottail shiner	Benthivore	2.1 ± 0.1	Potadromous	Opportunistic
Couesius plumbeus (COPL)	Lake chub	Benthivore	3.4 ± 0.4	Potadromous	Opportunistic

* The three life history strategies in fishes represent the trade-offs between demographic parameters of survival, fecundity and duration of reproduction. Opportunistic strategists are small-bodied species with early maturation and low juvenile survivorship and are predicted to be associated with habitats defined by frequent and intense disturbance. Periodic strategists are characterized by large body size, late maturation, high fecundity and low juvenile survivorship and are likely to be favored in highly periodic (seasonal) environments. Equilibrium strategists are typically small to medium in body size with intermediate times to maturity, low fecundity per spawning event and high juvenile survivorship largely due to high parental care and small clutch size. Equilibrium strategists are predicted to be favored in more stable habitats with low environmental variation.

Froese, R. and D. Pauly. Editors. 2016. FishBase. World Wide Web electronic publication. www.fishbase.org, version 10.2016

Desroches, J.-F. and Picard, I. (2013). Poissons d'eau douce du Québec et des Maritimes. Éditions Michel Quintin. Québec. Canada

Mims M.C. & Olden J.D. (2013) Fish assemblages respond to altered flow regimes via ecological filtering of life history strategies. Freshwater Biology 58, 50–62.

Olden J.D., Poff, N.L. & Bestgen K.R. (2006) Life-history strategies predict fish invasions and extirpations in the Colorado River basin. Ecological Monographs 76(1), 25-40.