1

Disabling Cas9 by an anti-CRISPR DNA mimic

Jiyung Shin[1,2]*, Fuguo Jiang[2]*, Jun-Jie Liu[2,5]*, Nicolas L. Bray[1,2], Benjamin J. Rauch[3],

Seung Hyun Baik[1,2] , Eva Nogales[2,5], Joseph Bondy-Denomy[3], Jacob E. Corn[1,2]† and

Jennifer A. Doudna[1,2,4,5]†

[1]Innovative Genomics Institute, [2]Department of Molecular and Cell Biology, University of

California, Berkeley, California 94720; [3]Department of Microbiology and Immunology,

and Quantitative Biosciences Institute, University of California, San Francisco, CA

94158; [4]Howard Hughes Medical Institute; [5]Department of Chemistry, University of

California, Berkeley, California 94720; [5]Molecular Biophysics and Integrated Bioimaging

Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720

*These authors contributed equally to the project.

†Correspondence should be addressed to J.E.C. (jcorn@berkeley.edu) and J.A.D.

(doudna@berkeley.edu)

CRISPR-Cas9 gene editing technology is derived from a microbial adaptive

immune system, where bacteriophages are often the intended target. Natural

inhibitors of CRISPR-Cas9 enable phages to evade immunity and show promise

in controlling Cas9-mediated gene editing in human cells. However, the

mechanism of CRISPR-Cas9 inhibition is not known and the potential

applications for Cas9 inhibitor proteins in mammalian cells has not fully been

established. We show here that the anti-CRISPR protein AcrIIA4 binds only to

assembled Cas9-single guide RNA (sgRNA) complexes and not to Cas9 protein

alone. A 3.9 Å resolution cryo-EM structure of the Cas9-sgRNA-AcrIIA4 complex

revealed that the surface of AcrIIA4 is highly acidic and binds with 1:1

stoichiometry to a region of Cas9 that normally engages the DNA protospacer

adjacent motif (PAM). Consistent with this binding mode, order-of-addition

experiments showed that AcrIIA4 interferes with DNA recognition but has no

effect on pre-formed Cas9-sgRNA-DNA complexes. Timed delivery of AcrIIA4 into

human cells as either protein or expression plasmid allows on-target Cas9-

mediated gene editing while reducing off-target edits. These results provide a

mechanistic understanding of AcrIIA4 function and demonstrate that inhibitors

can modulate the extent and outcomes of Cas9-mediated gene editing.

Phage-encoded inhibitors of CRISPR-Cas bacterial immune systems evolved to enable

phage escape from destruction in bacterial cells (*1*) and have the potential to control

CRISPR-Cas enzymes that are deployed for gene editing applications in various cell

types (*2, 3*). Determination of the molecular basis for Cas9 inhibition could shed light on

the evolutionary "arms race" between phage and bacteria, as well as suggest new

approaches to regulate genome editing in eukaryotic cells. The 87 amino acid anti-

CRISPR-Cas9 protein AcrIIA4 is notable in both respects: it inhibits multiple Cas9

proteins including the widely used Cas9 ortholog from *Streptococcus pyogenes* (Spy),

and it blocks Cas9-mediated gene editing in human cells (*3*). To investigate the

molecular basis for AcrIIA4-mediated Cas9 inhibition, we first tested whether

recombinant AcrIIA4 protein interacts directly with SpyCas9 (Fig. 1A). Purified AcrIIA4

was incubated with *Spy*Cas9 in the presence or absence of a single-guide RNA

(sgRNA) that assembles with Cas9 to provide sequence-specific DNA recognition (*4*).

Size exclusion chromatography showed that AcrIIA4 binds to SpyCas9 only in the

presence of sgRNA (Fig. 1B and C, fig. S1), implying that AcrIIA4 recognizes a protein

surface created upon sgRNA-triggered conformational rearrangement (*5, 6*).

Furthermore, the AcrIIA4-bound Cas9-sgRNA complex is more resistant to proteolytic

digestion than the Cas9-sgRNA complex alone, implying that AcrIIA4 stabilizes a

particular Cas9 conformation (fig. S2).

To elucidate the molecular basis of AcrIIA4-mediated inhibition of Cas9 activity,

we performed cryo-electron microscopy (cryo-EM) single-particle analysis on a

SpyCas9-sgRNA complex bound to AcrIIA4. Cryo-EM images were collected on a Krios

microscope using zero-loss energy-filtered imaging and a K2 direct electron detector.

After unsupervised 3D classification of 840,000 particle images, refinement of a class

containing 285,600 particles resulted in an EM reconstruction of the SpyCas9-sgRNA-

AcrIIA4 complex with an overall resolution of 3.9 Å (fig. S3-S4). Subsequent local 3D

classification that was focused on Cas9's HNH nuclease domain ultimately yielded two

EM reconstructions, one at 3.9 Å resolution where the HNH was poorly resolved due to

flexibility (reconstruction 1, obtained as a combination of two 3D classes) (Fig. 2a and

fig. S5a) and one with better defined density for the HNH domain at an average

resolution of 4.5 Å (reconstruction 2) (fig. S4d and S5a; see Methods section and fig.

S3). In both cryo-EM reconstructions, the density for the HNH domain of Cas9 was

weaker than for the rest of the structure, consistent with the previously observed

conformational plasticity of the HNH nuclease domain in the pre-targeting state. An

atomic model for AcrIIA4 was built from reconstruction 1. The EM density map displays

excellent main-chain connectivity and side-chain densities for almost all residues of

AcrIIA4 (fig. S5). We then built an atomic model for the Cas9-sgRNA complex based on

the crystal structure of SpyCas9-sgRNA (PDB ID 4ZT0) and refined the entire SpyCas9-

sgRNA-AcrIIA4 model in real space to good stereochemistry (Fig. 2B and table S1).

Overall, Cas9 bound to AcrIIA4 resembles the pre-target state rather than the DNA-

bound state (fig. S6).

De novo model building demonstrated that AcrIIA4 binds to Cas9 with 1:1

stoichiometry and comprises a three-stranded antiparallel β sheet flanked by one alpha-

helix at the amino-terminal end and two α helices on the carboxy-terminal end

$(\alpha_1\beta_1\beta_2\beta_3\alpha_2\alpha_3)$. Remarkably, superposition of the AcrIIA4-bound Cas9 structure with

DNA-bound Cas9 revealed that AcrIIA4 sits exactly in the PAM-interacting cleft formed

between the alpha-helical recognition (REC) lobe and the nuclease lobe (Fig. 2B and

fig. S6). AcrIIA4 completely occupies the PAM binding pocket and thus blocks DNA

recognition through contacts between the β3 strand of AcrIIA4 and Cas9 PAM-binding

residues (R1333 and R1335) (Fig. 2D-E). In addition, AcrIIA4 wedges into the DNA

melting region immediately upstream of the PAM sequence and sits on top of +1

phosphate on the target strand and the flipped nucleotides on the nontarget strand,

indicating AcrIIA4 could also prevent DNA binding/unwinding. Consistent with this DNA-

mimicing binding mode, AcrIIA4 is extremely acidic (Fig. 2E). Interestingly, AcrIIA4 also

occupies the same space as the DNA-bound HNH domain and the linker connecting the

HNH and RuvC domains (Fig. 2D), suggesting AcrIIA4 could block the HNH movement

required for catalysis (Fig. 2B, C).

Collectively, our structural studies show that AcrIIA4 is a highly acidic DNA mimic

that blocks target DNA recognition through multiple mechanisms: 1) competitive

inhibition of PAM binding; 2) inhibition of DNA unwinding upstream of the PAM

sequence; and 3) inactivation of HNH domain movement from the inactive to active

conformation. These structural findings help explain the effectiveness of AcrIIA4 as an

inhibitor of Cas9-mediated DNA cleavage and cell-based genome editing. Previous

studies revealed the importance of Cas9's interactions with targeted DNA, mediated via

the PAM in an interaction preceding base pairing between the sgRNA and its target (7).

Transient Cas9-sgRNA association with PAM sequences in DNA is thought to enable a

rapid target sequence search. By mimicking the structure and electrostatic properties of

the DNA PAM sequence, AcrIIA4 might compete for initial DNA binding and thereby

prevent target recognition and cleavage (Fig. 2E-G).

To determine the functional effects of AcrIIA4 binding to Cas9-sgRNA

complexes, we subjected DNA substrates possessing a target sequence to Cas9-

sgRNA-catalyzed DNA cleavage assays. A linearized plasmid possessing a target

sequence and PAM was incubated with combinations of Cas9, sgRNA and AcrIIA4, and

the products were resolved by gel electrophoresis. Cas9-sgRNA alone completely cut

the target DNA within five minutes, while AcrIIA4 limited cleavage at even later time

points (Fig. 3A). To confirm that AcrIIA4 would also inactivate Cas9 at low enzyme and

inhibitor concentrations, we performed similar experiments with a radiolabeled DNA

target for increased sensitivity (Fig. 3B). Near-stoichiometric concentrations of AcrIIA4

inhibited DNA cleavage when titrated against 10 nM Cas9-gRNA complex, indicating an

apparent dissociation constant of less than 10 nM. These data show that AcrIIA4

functions as a robust Cas9 "off-switch" that can inhibit most Cas9 activity at low

concentrations.

The structure of Cas9 bound to AcrIIA4 suggested that the inhibitor competes for

the initial DNA recognition event of PAM binding. Yet we previously found that Cas9

binds so tightly to target DNA that its off-rate is negligible (*7, 8*). This implies an unusual

non-equilibrium mode of inhibition, in which AcrIIA4 requires access to Cas9-sgRNA

prior to formation of the Cas9-sgRNA-DNA complex. To test this prediction, we used

bilayer interferometry to measure the binding of catalytically inactivated Cas9 (dCas9) to

a DNA target in the presence of inhibitor. These experiments were performed under

stoichiometric binding conditions in which Cas9 was present at concentrations greater

than the dissociation constant of the Cas9-AcrIIA4 interaction. Pre-incubation of Cas9

with AcrIIA4 markedly inhibited Cas9 binding to DNA (on-rate) in a dose-dependent

fashion, including complete prevention of target engagement (Fig. 3C, fig. S7). An

electrophoretic mobility shift assay (EMSA) further confirmed that AcrIIA4 does not

impact sgRNA loading into Cas9 protein alone (fig. S8), but does inhibit Cas9-sgRNA

complex binding to a target DNA (fig. S9). Strikingly, allowing the Cas9-sgRNA-DNA

complex to form and then adding AcrIIA4 had no effect on Cas9's release of target DNA

(off-rate). These order-of-addition results are consistent with both the structure of

AcrIIA4 bound to Cas9 and Cas9's extremely slow dissociation from DNA. Together,

these data support a mechanism for AcrIIA4 inhibition in which the inhibitor blocks

Cas9's ability to bind and cut target DNA by obscuring the PAM-interacting domain.

To determine the ability of AcrIIA4 to regulate gene editing in human cells, we

first utilized human K562 erythroleukemia cells stably expressing a chromosomally-

integrated blue fluorescent protein (BFP) reporter (*8*). Nucleofection of Cas9-sgRNA

(Cas9 RNP) complexes targeting BFP resulted in loss of BFP fluorescence in almost all

cells as measured by flow cytometry (fig. S10). Simultaneous delivery of Cas9 RNP and

AcrIIA4 protein inhibited Cas9-mediated gene targeting by up to 80% (Fig. 4A blue

symbols; fig. S10). Simultaneous delivery of AcrIIA4 encoded in a plasmid inhibited

gene editing to a lesser extent, possibly due to a delay in expression of the inhibitor

from a plasmid relative to immediate nucleofection of Cas9 RNP (Fig. 4A green

symbols; fig. S10, S11). This implied that the order-of-addition effects we observed in

biophysical experiments may play a role during gene editing. To test this idea, we

nucleofected AcrIIA4 plasmid 24 hours prior to introducing Cas9 RNP into cells. The

presence of the AcrIIA4-encoding plasmid 24 hours prior to Cas9 RNP introduction

inhibited Cas9-mediated gene targeting to a far greater extent than was observed

during co-introduction, and to a similar extent as observed during co-introduction of

AcrIIA4 protein and Cas9 RNP (Fig. 4B, fig. S12). Remarkably, we found that addition of

AcrIIA4 six hours after Cas9-RNP reduced editing by ~50%, demonstrating the utility of

inhibitors for revealing *in vivo* gene editing kinetics (Fig. 4C, fig. S13). In sum,

controlling the timing of AcrIIA4 inhibition in human cells strongly affects the frequency

of gene editing at a given locus.

Using a given guide RNA, Cas9 may target both an on-target site and off-target

loci (*9-12*). But several lines of evidence suggest that off-target sites may be bound

without being immediately cleaved (*13-15*). Cas9 displaced from uncleaved sites (e.g.

by cellular factors) would be available for inhibition by AcrIIA4. Since inhibitor timing

experiments suggested that at least 50% of on-target Cas9 gene editing takes place

within the first 6 hours (Fig. 4C), we asked whether off-target editing could be reduced

by properly timed addition of inhibitor.

We examined on- and off-target editing using guide RNAs targeting the EMX1

and HBB loci. The off-target sites for both the EMX1 and HBB guides have been

previously described (*10, 16, 17*). Notably, the HBB guide is of therapeutic interest to

edit the causative mutation of sickle cell disease, but has an off-target that is nearly

identical to the on-target site. We edited the EMX1 and HBB loci using Cas9 RNPs with

and without pre-introduction of AcrIIA4 protein or plasmid. Consistent with experiments

at the BFP locus, we found that timed addition of AcrIIA4 retained substantial levels of

on-target editing. Strikingly, both T7E1 and TIDE analysis showed that timed addition of

AcrIIA4 almost completely abolished off-target editing at both HBB and EMX1 (Fig. 4D).

This suggests that Cas9 inhibitors have the potential to be useful in reducing off-target

events during research and therapeutic applications.

The recent and rapid expansion of the Cas9 toolkit for gene editing applications

has lacked an inducible "off-switch" to prevent undesired gene editing. Newly

discovered protein inhibitors, encoded by bacteriophages, provide an attractive solution

to this problem as these proteins are small and function well in human cells (*3*). Here,

we demonstrate that AcrIIA4, the most potent SpyCas9 inhibitor in human cells, acts as

a DNA mimic to block PAM recognition. The deployment of DNA mimics to inhibit a

DNA-binding protein is an elegant solution that is often deployed in the phage-host arms

race. A recent structural study revealed that a Class 1 anti-CRISPR (AcrF2) uses a

similar strategy (*18*). Further, phage-encoded restriction enzyme inhibitor proteins have

long been known to mimic the DNA target (*19*). Our results further show that Cas9

inhibitors delivered as either protein or expressed from a plasmid can modulate the

efficacy of gene editing at multiple loci in human cells. Although pre-addition of inhibitor

almost completely abolishes overall gene editing, timed addition of inhibitor after

initiating Cas9-sgRNA-based gene editing can adjust the amount of time that Cas9 is

active in the nucleus, thereby selectively limiting off-target editing. We anticipate that

Cas9 inhibitors could be broadly useful in situations where precise control of either on-

or off-target gene editing is desirable, such as during allele-specific therapeutic editing.

Figures

Fig 1. AcrIIA4 binds to the *S. pyogenes* Cas9-sgRNA complex.

A) A cartoon depiction of Cas9 protein loaded with the sgRNA binding to AcrIIA4

(pink). Cas9-sgRNA complexed with AcrIIA4 is unable to bind to the target DNA.

B) Size exclusion chromatogram of SpyCas9-sgRNA after preincubation with

AcrIIA4. Relevant peaks are indicated with arrowheads.

C) Coomassie and Ethidium Bromide (EB) stained polyacrylamide gel showing the

commigration of AcrIIA4 with Cas9 in the presence of guide RNA.

Fig. 2. Architecture of the *S. pyogenes* Cas9-sgRNA in complex with AcrIIA4

A) Cryo-EM reconstruction of the AcrII4-bound SpyCas9. The electron density map

was contoured at high threshold levels showing distinct features for each subunit.

B) The atomic model of SpyCas9-sgRNA-ArcrIIA4. AcrIIA4 (red) and sgRNA

(orange) are shown in ribbon diagram and SpyCas9 is displayed as surface

representation.

C) Representative region (hydrophobic core) of the cryo-EM density for AcrIIA4 with

the refined model superimposed.

D) Superposition with Cas9-guideRNA-dsDNA structure (PDB ID 5F9R). For clarity,

Cas9 is omitted except the HNH domain. Target and nontarget DNA strand is is

colored purple and beige, respectively.

E) Electrostatic surface potential of AcrIIA4. The inset shows the PAM recognition

residues (R1333 and R1334) are largely buried in an acidic pocket within

AcrIIA4.

F) and G) Close-up of AcrIIA4 binding pocket at different views showing AcrIIA4 is a

dsDNA mimic.

Fig. 3. AcrIIA4 inhibits DNA cleavage *in vitro*

A) A linearized plasmid was incubated with Cas9 alone, AcrIIA4 alone, Cas9 + sgRNA, or Cas9 + sgRNA + AcrIIA4 for 5, 15, or 30 minutes. Reactions were resolved via agarose gel electrophoresis and visualized by ethidium bromide staining.

B) AcrIIA4 inhibits SpyCas9 cleavage of radiolabeled target DNA *in vitro*. 10 nM SpyCas9-crRNA-tracrRNA complex were pre-incubated with increasing concentrations (0-100 nM) of AcrIIA4. Sub-stoichiometric synthetic oligonucleotide duplexes (2.5 nM) bearing a radiolabel at the 5' end of the complementary strand, were introduced for 6-minute cleavage reactions. Reactions were resolved by denaturing polyacrylamide gel electrophoresis and visualized by

phosphorimaging.

C) AcrIIA4 inhibits dCas9-sgRNA binding to a DNA target but does not affect target

release, as measured by biolayer interferometry. Pre-incubating increasing

concentrations of AcrIIA4 with dCas9-sgRNA reduces the on-rate of association

with a DNA target relative to no inhibitor (blue). Maximal inhibition is identical to

dCas9-sgRNA added to target DNA with a PAM mutation (red).

D) Addition of increasing concentrations of AcrIIA4 with pre-formed dCas9-sgRNA-

DNA complex has no effect upon the off-rate of dissociation.

Fig. 4. Timed delivery of AcrIIA4 differentially inhibits on- and off-target genome

editing in human cells.

A) Simultaneous delivery of Cas9 RNP and AcrIIA4 inhibits Cas9-mediated gene

targeting in human cells. K562 cells with a chromosomally integrated BFP (BFP-

K562) were nucleofected with Cas9 RNP and AcrIIA4 protein (Cas9:AcrIIA4

(molar ratio), 1:0.5, 1:1, 1:2, 1:3, 1:5) or plasmid (0.7µg and 2.8µg). NHEJ

frequencies are quantified by loss of BFP expression in BFP-K562 cells 96 hours

post nucleofection via flow cytometry. Data presented as mean ± SEM from at

least two biological replicates.

B) Administration of AcrIIA4 prior to Cas9 RNP completely inhibits Cas9-mediated

gene targeting. BFP-K562 cells were nucleofected with AcrIIA4 plasmid (0.7µg)

24 hours prior to Cas9 RNP delivery. Data presented as mean ± SEM from at

least two biological replicates.

C) Delivery of AcrIIA4 after introduction of Cas9 RNP yields intermediate inhibition

of Cas9 activity. BFP-K562 cells were nucleofected with AcrIIA4 protein

(Cas9:AcrIIA4 (molar ratio), 1:5) or plasmid (0.7µg) 6 hours post Cas9 RNP

delivery. Data presented as mean ± SEM from at least two biological replicates.

D) Proper timing of AcrIIA4 delivery diminishes off-target editing events while largely

retaining on-target editing. K562 cells were nucleofected with either HBB or

EMX1 targeting Cas9 RNP 6 hours prior to AcrIIA4 protein (Cas9:AcrIIA4 (molar

ratio), 1:5) or plasmid (0.7µg) delivery. (Top) Representative T7 endonuclease I

assay for visualization of HBB on- and off- target editing. (Bottom) Quantification

of on- and off-target editing at HBB and EMX1, as measured by TIDE analysis.

References

274   1.   J. Bondy-Denomy, A. Pawluk, K. L. Maxwell, A. R. Davidson, Bacteriophage genes that

275   inactivate the CRISPR/Cas bacterial immune system. *Nature* **493**, 429-432 (2013).

276   2.   A. Pawluk *et al.*, Naturally Occurring Off-Switches for CRISPR-Cas9. *Cell* **167**, 1829-

277   1838 e1829 (2016).

278   3.   B. J. Rauch *et al.*, Inhibition of CRISPR-Cas9 with Bacteriophage Proteins. *Cell* **168**,

279   150-158 e110 (2017).

280   4.   M. Jinek *et al.*, A programmable dual-RNA-guided DNA endonuclease in adaptive

281   bacterial immunity. *Science* **337**, 816-821 (2012).

282   5.   M. Jinek *et al.*, Structures of Cas9 endonucleases reveal RNA-mediated

283   conformational activation. *Science* **343**, 1247997 (2014).

284   6.   H. Nishimasu *et al.*, Crystal structure of Cas9 in complex with guide RNA and target

285   DNA. *Cell* **156**, 935-949 (2014).

286   7.   S. H. Sternberg, S. Redding, M. Jinek, E. C. Greene, J. A. Doudna, DNA interrogation by

287   the CRISPR RNA-guided endonuclease Cas9. *Nature* **507**, 62-67 (2014).

288   8.   C. D. Richardson, G. J. Ray, M. A. DeWitt, G. L. Curie, J. E. Corn, Enhancing homology-

289   directed genome editing by catalytically active and inactive CRISPR-Cas9 using

290   asymmetric donor DNA. *Nat Biotechnol* **34**, 339-344 (2016).

291   9.   Y. Fu *et al.*, High-frequency off-target mutagenesis induced by CRISPR-Cas nucleases

292   in human cells. *Nat Biotechnol* **31**, 822-826 (2013).

293   10.   P. D. Hsu *et al.*, DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat*

294   *Biotechnol* **31**, 827-832 (2013).

11.   P. Mali *et al.*, CAS9 transcriptional activators for target specificity screening and

paired nickases for cooperative genome engineering. *Nat Biotechnol* **31**, 833-838

(2013).

12.   V. Pattanayak *et al.*, High-throughput profiling of off-target DNA cleavage reveals

RNA-programmed Cas9 nuclease specificity. *Nat Biotechnol* **31**, 839-843 (2013).

13.   S. Kiani *et al.*, Cas9 gRNA engineering for genome editing, activation and repression.

Nat Methods **12**, 1051-1054 (2015).

14.   S. H. Sternberg, B. LaFrance, M. Kaplan, J. A. Doudna, Conformational control of DNA

target cleavage by CRISPR-Cas9. *Nature* **527**, 110-113 (2015).

15.   X. Wu *et al.*, Genome-wide binding of the CRISPR endonuclease Cas9 in mammalian

cells. *Nat Biotechnol* **32**, 670-676 (2014).

16.   T. J. Cradick, E. J. Fine, C. J. Antico, G. Bao, CRISPR/Cas9 systems targeting beta-

globin and CCR5 genes have substantial off-target activity. *Nucleic Acids Res* **41**,

9584-9592 (2013).

17.   M. A. DeWitt *et al.*, Selection-free genome editing of the sickle mutation in human

adult hematopoietic stem/progenitor cells. *Sci Transl Med* **8**, 360ra134 (2016).

18.   S. Chowdhury *et al.*, Structure Reveals Mechanisms of Viral Suppressors that

Intercept a CRISPR RNA-Guided Surveillance Complex. *Cell* **169**, 47-57 e11 (2017).

19.   M. D. Walkinshaw *et al.*, Structure of Ocr from bacteriophage T7, a protein that

mimics B-form DNA. *Mol Cell* **9**, 187-194 (2002).

Acknowledgments

EM-derived maps and atomic coordinates of the Cas9-sgRNA-AcrIIA4 structure have

been deposited in the EM Databank (EMDB) and the Protein Data Bank (PDB) with

accession codes **EMD-XXXX and YYYY, respectively.** The EM data was collected in

the EM facility of HHMI Janelia Research Campus. We thank Rick Huang and Zhiheng

Yu for expert electron microscopy assistance; and members of the Bondy-Denomy,

Corn, Doudna and Nogales labs for helpful discussions and critical reading of the

manuscript. F.J. is a Merck Fellow of the Damon Runyon Cancer Research. J.E.C. is

supported by the Li Ka Shing Foundation and the Heritage Medical Research Institute.

J.S. is supported by the National Institute on Aging of the National Institutes of Health

under Award Number T32 AG000266. J.B.-D. and B.R. are supported by the University

of California San Francisco Program for Breakthrough in Biomedical Research, funded

in part by the Sandler Foundation, and an NIH Office of the Director Early

Independence Award (DP5-OD021344). J.A.D. and E.N. are Investigators of the

Howard Hughes Medical Institute. This work was supported in part by HHMI. The

content is solely the responsibility of the authors and does not necessarily represent the

official views of the National Institutes of Health.