1   *Title:*
2   **The evolutionary dynamics of the odorant receptor gene family in corbiculate**
3   **bees**
4

5   *Authors:*
6   Philipp Brand[1,2] and Santiago R. Ramírez[1]
7
8   *Author affiliations:*
9   [1]Department for Evolution and Ecology, Center for Population Biology, University of
10   California Davis, One Shields Avenue, 95616 Davis, USA
11   [2]Population Biology Graduate Group, University of California Davis, One Shields
12   Avenue, 95616 Davis, USA
13
Corresponding author:
Philipp Brand, Department for Evolution and Ecology, Center for Population Biology,
University of California Davis, One Shields Avenue, 95616 Davis, USA, +1 530-752-
7614, pbrand@ucdavis.edu

Keywords:
Odorant receptors, Chemosensory genes, birth-death process, Genomic architecture,
Positive selection, Corbiculate bees, Honey bees, Bumble bees, Stingless bees, Orchid
bees

Abstract

Insects rely on chemical information to locate food, choose mates, and detect potential predators. It has been hypothesized that adaptive changes of the olfactory system facilitated the specialization and diversification of numerous insect lineages. In fact, modifications of life history strategies often occur in parallel with evolutionary changes in Odorant Receptor (OR) genes. However, the molecular mechanisms underlying the evolution of insect OR genes remain poorly understood. Corbiculate bees include economically and ecologically important bee pollinators that exhibit varying degrees of social organization, diverse sex pheromone systems, and manifold associations with floral resources. We investigated the molecular mechanisms governing the evolution of the OR gene family in corbiculate bees. Our results indicate that the genomic organization of the OR gene family has remained highly conserved for approximately 80 million years, despite exhibiting major changes in repertoire size among bee lineages. Moreover, the evolution of OR genes appears to be driven by lineage-specific gene duplications in few genomic regions harboring large numbers of OR genes. A selection analysis revealed that OR genes evolve under positive selection, with the strongest signals detected in recently duplicated copies. Our results indicate that chromosomal translocations had a minimal impact on OR evolution, and instead show that local molecular mechanisms drive OR repertoire size. Our results provide empirical support to the longstanding hypothesis that positive selection shaped the diversification of the OR gene family. Our results shed new light on the molecular mechanisms underlying the evolution of olfaction in insects and gene families in general.

Significance Statement

Despite playing a central role in the diversification of insect lineages, the evolution of insect olfactory systems remains poorly understood. Here, we analyze the Odorant Receptor (OR) repertoire across the corbiculate bees—a lineage of important insect pollinators—to test the hypothesis that genomic re-organization and positive selection shaped the evolution of the OR gene family. We show that locally acting molecular mechanisms are the driving force of OR gene family evolution in corbiculate bees. Moreover, we provide empirical support for the longstanding hypothesis that positive selection impacts the evolution of recently duplicated insect OR genes. Our results augment our understanding of the molecular mechanisms that govern the evolution of olfactory systems in insects and gene families in general.

Introduction

Animals have evolved sophisticated sensory systems that can detect and discriminate airborne volatile chemicals and provide precise information about food, enemies, and mating partners (1). Insects detect olfactory signals and cues via olfactory sensory neurons located in the antenna. Each olfactory sensory neuron expresses Odorant Receptor (OR) proteins on the cell membrane (2-4). It is the interaction of volatile odorant molecules with these OR proteins that initiates the signal transduction and transmission toward the olfactory centers in the insect brain (2, 4). Insect genomes are endowed with a diverse array of functional OR genes, each of which encodes a unique odorant receptor protein tuned to a specific set of odors (5-7). The entire repertoire of OR genes expressed in the olfactory organs determines the spectrum of chemical volatiles that an insect species can detect.

Comparative analyses of insect genomes have revealed that the number of OR genes and divergence among them can vary widely between species. Insect genomes may contain as few as 10 OR genes (head lice, (8)) and as many as 300 OR genes (ants, (9, 10)). In fact, the expansion and contraction of the OR gene family has been linked to shifts in the sensory ability of some insect lineages. This supports the idea that adaptation to novel food resources (11-13) or modifications in the pheromone communication system (14) can be mediated by peripheral changes in the OR gene repertoire. In addition, changes in the amino acid sequence of existing OR genes have been shown to correspond to adaptive shifts in sensory tuning (15, 16). However, we lack a clear understanding of the specific genetic and molecular mechanisms that drive the diversification of the insect OR gene family (17, 18).

Gene family evolution is determined by multiple molecular mechanisms including genomic drift, natural selection, and chromosomal rearrangements. Birth-death processes can directly impact the number of genes within a gene family (19). For example, genomic drift may generate new gene copies through gene duplication (gene 'birth') while existing copies can be purged via pseudogenization or deletion (gene 'death'). Novel mutations resulting in gene duplication or gene loss may be subsequently fixed through neutral genetic drift or positive selection (17, 19, 20). Although OR gene repertoire sizes have been analyzed extensively in various insect lineages, the relative contribution of positive selection and neutral processes in a birth-death evolutionary framework remains uncertain (17, 18).

The genomic organization and architecture of gene families may contain signatures of past evolutionary forces that shaped their diversification. For instance birth-death processes often produce tandem arrays of OR genes, which are groups of ancestrally duplicated genes located in close physical proximity along the genome (19, 21). Chromosomal rearrangements and transposition, on the other hand, can lead to inter-chromosomal translocations of OR genes (22, 23), and it has been hypothesized that changes in the location of OR genes along the genome can influence their evolution (18, 23, 24). Although both tandem duplications and

139 chromosomal rearrangements may influence the evolution of the OR gene family,
140 the relative contribution of each process remains unknown.
141
Like most insects, bees rely on olfactory information to regulate a wide array of
behaviors, including the location of food sources and nesting materials, the
identification of mating partners, and social interactions with other colony members
in social species. Corbiculate bees encompass a group of ecologically and
economically important bee lineages, including honey bees, bumble bees, stingless
bees, and orchid bees. Honey bees are estimated to pollinate one-third of
agricultural crops (25), and stingless bees and orchid bees are the major pollinators
of numerous tropical flowering plant species (26, 27). Corbiculate bees have evolved
a variety of phenotypic and behavioral traits that require specialized olfactory
functions. These include the evolution of an obligate cooperative social lifestyle
(eusociality), which has resulted in the evolution of specialized pheromone
communication systems in honey bees, bumble bees and stingless bees (28-31). In
addition, orchid bees evolved a unique pheromone communication system in which
male bees concoct species-specific perfume bouquets from scents collected from
flowers and other sources to subsequently use during courtship display (32-34).
However, despite the central role that olfactory encoding plays in the biology of
corbiculate bees, we lack a clear understanding of the evolution of the OR gene
family in this diverse group of bees.
160
Here we analyzed the evolution of the OR gene family in a set of species including
the four corbiculate bee clades, spanning 80 million years (my) of evolution. We
tested whether the evolution of the OR gene family has been influenced by changes
in the genomic organization of genes, as hypothesized based on patterns observed
in *Drosophila* and mammalian genomes (18, 23, 24). Second, we tested the
hypothesis that birth-death processes, in combination with positive selective
pressures, promoted the origin of novel OR genes. Finally, we estimated the relative
importance of positive selection in shaping OR sequence divergence, a mechanism
that has been hypothesized to drive sensory adaptation in insect lineages (15, 35,
36).
171
172 **Results**
173
The Evolution of Corbiculate Bee Odorant Receptors is Highly Dynamic. To
analyze the evolutionary dynamics of odorant receptor (OR) gene repertoires in
corbiculate bees, we reconstructed the phylogenetic relationships among the entire
OR gene family based on both whole-genome sequences and transcriptome data
obtained from ten bee species (Fig. 1). An initial comparison of OR annotation based
on genomic and transcriptomic data indicated no differences in annotation
accuracy, but as expected, the number of genes tended to be lower for
transcriptome data (Supplemental Text). Our analysis included published whole-
genome sequence data from at least one species per corbiculate bee tribe (21, 37-
39). In addition, we sequenced the genome of one orchid bee (*Euglossa dilemma;*
(40)) and the antennal transcriptomes of five additional orchid bee species. The

resulting dataset, which included three orchid bee species pairs, allowed us to
analyze the evolution of the OR gene family at multiple taxonomic levels over a
broad range of divergence times (0.15 to 80mya; Fig. 1). We compared the evolution
of the OR gene family, the largest and most dynamic chemosensory gene family in
insects, to the evolution of four additional chemosensory gene families, including
the Gustatory Receptors (GRs), Ionotropic Receptors (IRs), Odorant Binding
Proteins (OBPs), and Chemosensory Proteins (CSPs). We used genomic data only to
derive evolutionary patterns of gene family size since transcriptome data often yield
incomplete gene sets (Fig. 1).
194
Gene repertoire size varied substantially among bee lineages, ranging from 142 OR
genes in the orchid bee *Eufriesea mexicana* to 196 OR genes in the stingless bee
Melipona quadrifasciata (Fig. 1). The identified ORs were organized in 115
homologous OR ortho-groups consisting of orthologs and/or inparalogs (*i.e.* lineage-
specific expansions, Fig. 2, Fig. S1). Of these ortho-groups, 24 did not contain
duplications but instead consisted of simple 1:1 orthologous genes that were
present in three or more bee species (Fig. 2b). The remaining 91 ortho-groups
included species-specific, genus-specific, and tribe-specific duplications or larger
expansions (Fig. 2c). In contrast to the other four gene families analyzed, the
repertoire size of ORs is highly variable among corbiculate bee lineages. The
majority of genes in the non-OR gene families revealed simple 1:1 orthology with
few lineage-specific duplications resulting in similar repertoire sizes among bee
lineages (Fig. 1; Fig. S2). An exception to this was an observed increase in gene
family size in honey bee OBPs as well as bumble bee GRs (Supplemental Text).
Among all chemosensory genes, we identified 32 gene duplications and large
expansions in the honey bees, 36 in the bumble bees, 26 in the stingless bees, and 18
in the orchid bees, most of which were observed in the OR gene family (89%, Tab.
S1). These results show that the evolution of the corbiculate bee OR gene family
repertoire is characterized by highly dynamic lineage-specific gene expansions.
214
The Genomic Organization of the OR Gene Family is Highly Conserved in
Corbiculate Bees. Next, we investigated the genomic organization of OR genes
across the corbiculate bees. Our analysis revealed that OR genes are widely
distributed throughout the five bee genomes we analyzed. This supports previous
studies where OR genes were detected on scaffolds assigned to almost all
chromosomes in those species with available linkage groups (*i.e.* honey bees and
bumble bees; (21, 37, 38)). Congruent with a birth-death process for OR gene
evolution, the majority of genes were clustered in large tandem arrays and only few
ORs were found in isolated genomic regions as singletons. We tested whether the
chromosomal location of OR genes is conserved among corbiculate bees by
comparing the 200 kb flanking regions of all OR singletons and tandem arrays
among species using an "all-against-all" reciprocal blastn approach.
227
With the exception of a small number of ancestral translocations (Supplementary
Text), we found that the flanking regions and thus the physical chromosomal
locations of most OR genes were strictly homologous in all corbiculate bee species

(Tab. S2). Furthermore, all orthologous OR genes found in two or more species were
located in homologous chromosomal regions. Paralogous genes within species-
specific, genus-specific, and tribe-specific duplications and larger OR sublineage
expansions were always located in the same tandem array. Similarly, ORs located
within the same genomic region were generally more closely related to each other
than ORs located in different genomic regions that formed highly supported
monophyletic gene clades (Fig. 2a). On the basis of this strong correlation between
phylogenetic relationship and genomic location, we defined a total of 25 well-
supported subfamilies in the OR gene family that largely corresponded to groups
previously identified in ants (Fig. 2a; (41, 42)). In addition to the high conservation
of genomic locations of OR genes and tandem arrays, we found that the organization
of genes within tandem arrays (*i.e.* gene synteny) is also highly conserved
(Supplementary Text; Fig. S3). Our analysis identified similar patterns of
conservation in the genomic organization of other chemosensory gene families,
including Gustatory Receptors, Ionotropic Receptors, Odorant Binding Proteins, and
Chemosensory Genes (Dataset S1).
247
The Evolution of OR Repertoire Sizes is Governed by Dynamic Evolutionary
Changes in Large Subfamilies. Our analysis revealed that although the genomic
organization of OR tandem arrays is highly conserved across corbiculate bee
lineages, OR evolution is highly dynamic, with multiple taxon-specific OR subfamily
expansions. In order to investigate potential localized molecular mechanisms that
shape the evolution of the OR gene family, we analyzed the divergence of the 25 OR
subfamilies (see above). We found that variation in subfamily size among species
was positively correlated with mean subfamily size (Fig. 3a, Pearson coefficient
r=0.85, p=0; Orchid bees only: r=0.6, p=0.0013), a common pattern in gene family
evolution (Dennis et al. 2017). This result suggests that larger OR subfamilies have a
higher gene turnover than smaller gene subfamilies. Accordingly, all subfamilies
with one or two genes per species consisted of simple 1:1 orthologous genes across
the corbiculate bee genomes and transcriptomes (Fig. S1; Tab. S3).
261
Three of the OR subfamilies that exhibited high interspecific variation in OR
repertoire size had an expansion in a single bee species. The G04A subfamily was
expanded in the honey bee *A. mellifera* (Fig. 2c), the G09A subfamily was expanded
in the orchid bee *Ef. mexicana,* and the G09C subfamily was expanded in the
stingless bee *M. quadrifasciata* (Fig. 3b, Tab. S3). Several subfamily expansions were
observed in two or more species. Although most expansions are not shared by
multiple bee species with similar life history traits, the G02A subfamily was
expanded in all obligate eusocial bee species we included in our analysis (honey bee,
bumble bee and stingless bee; Fig. 3b). Conversely, ORs in the subfamily GUnA and
G12A were expanded in orchid bees (Fig. 3b, Tab. S3), which are known for their
unique perfume collecting behavior. The expansion we observed in the subfamilies
G02A and GUnA correspond to well-supported lineage-specific clusters of paralogs
in our phylogenetic analysis (Fig. S1), suggesting that these subfamilies are
independent expansions leading to elevated OR numbers. In contrast, more than
half of the orchid bee OR expansion in subfamily G12A was shared between the

orchid bee species included in our analysis (*i.e.* orthologous copies were present),
suggesting that the expansion was already present in their common ancestor.
279
Positive Selective Pressures Drive the Evolution of Both Paralogous and
Orthologous OR Genes. To investigate the selective pressures that shaped the
evolution of the corbiculate bee OR gene family, we directly compared the evolution
of ORs with the evolutionary patterns of the remaining major chemosensory gene
families, including the GRs, IRs, OBPs, and CSPs. We classified all annotated
chemosensory genes into orthologous groups consisting of at least four orthologs
and/or inparalogs based on our phylogenetic analysis (see Fig. 2b-c for examples of
orthologous groups). The resulting 159 orthologous groups (115 OR, 13 GR, 9 IR, 15
OBP, and 7 CSP orthologous groups) contained a total of 1588 chemosensory genes
(Dataset S1). We tested each of the orthologous groups for the presence of episodic
positive selection using branch-site model based dN/dS estimates (Smith et al.
2015). We identified 54 chemosensory genes (2.1% of all terminal branches) and 57
deeper branches (6.1% of 941 deep branches) with signatures of positive selection
under a 5% false discovery rate (FDR). This corresponds to 4.4% of all branches
tested. We found no correlation between divergence time and positive selective
pressures (Supplemental Text; Tab. S4). Furthermore, there was no difference in the
number of branches under positive selection between the four corbiculate bee
tribes (Tab. 1). However, a comparison across gene families revealed a higher
proportion of branches under positive selection in the GRs relative to any other
gene family in all bee lineages, supporting the hypothesis that GRs exhibit elevated
signatures of positive selection across many insect groups ($p<0.0001$; Fisher's Exact
Test; Supplementary Text) (12).
302
Those branches that exhibited signatures of positive selection were distributed
throughout the entire OR gene tree (Fig. S1). Interestingly, only 32% of all branches
that we identified to evolve under positive selection belonged to lineage-specific
duplications or larger gene expansion events (Tab. S5). Thus, the majority of
branches under positive selection correspond to orthologous rather than
paralogous gene copies. This pattern held true when we considered only those five
species with genomic data. However, when normalized by the total number of the
two branch classes (duplication and non-duplication branches, respectively), we
found that a higher proportion of all tested duplication branches were under
positive selection in comparison to non-duplicated branches (6% vs. 3.9%, Fisher's
Exact Test, $p< 0.05$, Tab. S5). Together, these results suggest that positive selective
pressures play an important role in the evolution and divergence of duplicated OR
genes.
316
317 In addition, we observed that 10 of the 11 OR subfamilies without any branches
318 under positive selection included simple (single-copy) 1:1 orthologs only
319 (Subfamilies G01A, G01B, G02B, G05A, G09D, G14B, G15A, G15B, GUnB, ORco; Fig.
320 S1). This observation suggests the existence of a core set of OR genes with
321 conserved chemosensory functions across all corbiculate bees.
322

Discussion

Our analysis focused on documenting signatures of positive selection as well as changes in the genomic organization of genes in the OR gene family analyzing 1394 ORs spanning multiple evolutionary time scales across the corbiculate bees. Our analysis revealed that the corbiculate bee OR gene family is highly dynamic (Fig. 1, Fig. 2, Fig. S1), a pattern consistent with that observed in other insect lineages. However, the genomic organization we identified in corbiculate bees differs from that previously reported in other insect groups. First, we found that the genomic organization of OR genes is highly conserved between corbiculate bee lineages, despite sharing a common ancestor 80 million years ago (Tab. S2). Second, OR gene repertoire sizes vary significantly between bee species due to lineage-specific expansions in a few large tandem arrays, some of which correlate with the evolution of obligate eusociality and pheromone communication systems (Fig. 3; Fig. S2). These findings suggest that the highly dynamic evolutionary history of the OR gene family is independent of changes in genomic organization and instead more likely influenced by locally acting molecular mechanisms. Furthermore, we found evidence for elevated rates of positive selection on duplicate ORs (Tab. 1; Tab S6), supporting the main prediction from the birth-death model of gene family evolution. Finally, we identified a fraction of bee ORs under positive selection that have not undergone previous duplication, highlighting the importance of positive selective pressures in OR sequence divergence, possibly in response to ecological adaptation.

Locally Acting Molecular Mechanisms are the Driving Force of Chemosensory Evolution. Our phylogenetic and genome annotation analysis revealed a high degree of genomic landscape conservation in the OR gene family across the corbiculate bees. This pattern was accompanied by a fast diversification rate of OR genes, indicating that genomic rearrangements had a relatively weak influence relatively to the stronger influence of local molecular mechanisms. The high degree of conservation in the overall genomic organization holds true even between corbiculate bee lineages that shared a common ancestor 80 million years ago. This is an unprecedented level of genomic conservatism in OR organization, and is particularly surprising given the diverse factors known to affect genome composition and genome architecture. In fact, the bee lineages we examined exhibit more than two-fold variation in chromosome number (9 in stingless bees to 20 in orchid bees; (43)), an order of magnitude of variation in genome size (ranging from 0.25GB in honey bees to 3GB in orchid bees; (37-40)), as well as exceptionally high recombination rates (44-46). Moreover, the high level of conservation stands in sharp contrast to known evolutionary mechanisms that affect genomic organization of OR genes in drosophilid flies, the only other insect group for which the evolution of chemosensory genes has been examined in comparable detail. The genomes of drosophilid flies exhibit complex genome rearrangements leading to a high degree of ortholog OR gene translocation within evolutionary time scales similar to those spanning the bee species we analyzed (22, 47, 48). A similarly dynamic pattern of OR tandem array architecture was previously observed in mammalian genomes (49). These observations lead to the hypothesis that genomic rearrangements are

369   important in the evolution of chemosensory receptors across vertebrate and
370   invertebrate lineages (18, 23, 24). In contrast, our results show that the evolution of
371   the OR gene family in corbiculate bees has resulted in highly divergent gene
372   repertoires without undergoing genome rearrangements.
373
Our results revealed no correlation between the evolution of genomic organization
and genome size. In fact, genome size of most corbiculate bees resembles those
reported for drosophilid flies (37, 50). Instead, a recent comparison of the genomes
of honey bees and bumble bees revealed high conservation of genome synteny
between these species (44), indicating that conserved genomic architecture of OR
tandem arrays may result from a low precedence of genome rearrangements in
corbiculate bees. Indeed, our analysis revealed that synteny among OR genes in
orchid bees is almost perfectly conserved even within highly dynamic genomic
areas that are dominated by gene duplication (Supplementary Text). Despite the
high level of conservation in the genomic organization reported here, we find a high
degree of OR gene divergence and turnover among corbiculate bees. This result
suggests that chemosensory gene evolution in corbiculate bees, and perhaps insects
in general, is shaped by local rather than global mechanisms of genome evolution.
387
Large Tandem Arrays are Hotspots for the Evolution of Novel ORs. The high rate
of variation we observed in OR repertoire size among lineages, along with the
prevalence of lineage-specific gene expansions in large OR tandem arrays, suggests
that large gene subfamilies play an important role in the evolution of novel OR genes
in corbiculate bees. Insect OR repertoires are highly dynamic between lineages in
comparison to other chemosensory gene families (18, 51). Our results confirm this
pattern for corbiculate bees. In contrast to previous studies, however, we find that
these expansions are driven by lineage-specific expansions within large tandem
arrays harboring entire OR subfamilies. This result is congruent with a recent
phylogenetic analysis that showed that gene duplication and pseudogenization
events of OR genes in Hymenoptera are often found in large subfamilies (52). The
increased gene turnover rate could be influenced by elevated rates of genome
replication errors resulting from non-allelic homology of a large number of closely
related paralogous loci in close proximity within large tandem arrays (53-55).
Nevertheless, other factors such as transposable element activity and the timing of
DNA replication can influence segmental duplication rates (53, 56). Independent of
the underlying genomic mechanism, our results suggest that large OR tandem arrays
are a major source of novelty in chemosensory receptors in corbiculate bees and
potentially other Hymenoptera. Accordingly, large tandem arrays are likely an
important source of functional novelty associated with the detection of novel
molecules or blends of chemical compounds. Hence, it is possible that the gene
expansions we identified coincide with key evolutionary innovations such as
complex social chemical communication in the obligate eusocial honey bees, bumble
bees and stingless bees or perfume-collecting behavior in orchid bees.
412
It has been hypothesized that OR gene family size is correlated with the degree of
complexity in the chemical communication system displayed by eusocial

Hymenoptera (57). Although our data do not support this general trend, we find a
positive correlation between OR gene subfamily size and the presence of obligate
eusociality. For instance, the OR subfamily G02A is expanded in all three obligate
eusocial bee species analyzed, relatively to the smaller size exhibited by the solitary
to weakly social orchid bees. Moreover, the OR subfamily G02A is expanded in ants
but not in the facultative social halictid bee *Lasioglossum albipes*, which exhibits
subfamily G02A gene numbers comparable to orchid bees (Subfamily L in (42, 52)).
Thus, it is possible that ORs in subfamily G02A play a central role in the olfactory
detection and processing of pheromones involved in social behavior. Similarly, an
expansion of the '9-exon' subfamily (here subfamilies G04A, G05B, G11A, G11B, and
G12A; Fig. 2a; Fig. 3b; Fig S1) was hypothesized to be important in the evolution of
eusociality due to a putative role of '9-exon' ORs in the detection of social
pheromones (namely cuticular hydrocarbons, (9, 10, 58)). We find an expansion of
the '9-exon' subfamily in the eusocial honey bees and bumble bees but not eusocial
stingless bees. This observation contrasts with the previous suggestion that the
expansion of this OR subfamily was concomitant with the evolution of eusociality in
Hymenoptera. However, it is possible that ORs involved in social pheromone
communication evolved independently in different Hymenoptera lineages. Future
studies should investigate the possibility that olfactory adaptations to social
behavior evolved through multiple analogous routes in social Hymenoptera (39,
59).
436
In addition, we found two OR subfamilies that exhibit an expansion in the orchid bee
lineage (namely GUnA and G12A). While large expansions were in general species-
specific clusters of paralogous ORs (*e.g.* Fig. 2c; Fig. S1), more than half of the ORs in
the orchid bee expansion of subfamily G12A were orthologous ORs found in all the
seven species included in our analysis. This shared expansion may correspond to an
orchid bee specific olfactory specialization. Since the only known olfactory behavior
clearly distinguishing orchid bees from all other corbiculate bees is male perfume-
collecting behavior (34, 60), it is possible that the G12A subfamily is involved in the
detection and encoding of these exogenous compounds. Functional analyses are
required to investigate the olfactory coding mediated by members of this OR
subfamily.
448
Selective Patterns of Corbiculate Bee Chemosensory Genes Support the Birth-
Death Model of Gene Family Evolution. It has been proposed that chemosensory
gene families evolve under a birth-death process where duplications of existing
genes give rise to new gene copies (gene 'birth') whereas pseudogenization purges
genes from the genome (gene 'death'; (19)). The birth-death model predicts a
relaxation of purifying selective pressures after duplication allowing subsequent
diversification of one or both copies through positive selection for newly
accumulated beneficial mutations. Accordingly, the evolution of highly dynamic
chemosensory gene families is likely driven by the action of positive selection.
Nevertheless, while previous studies detected a high impact of gene duplication and
loss on chemosensory gene family evolution, there is very little evidence for

diversification through positive selection (12, 22, 52, 61-63), even in large lineage-
specific gene family expansions (42).
462
Our analysis provides empirical evidence congruent with the selective pressures
predicted by the birth-death model of chemosensory gene family evolution. We
show that positive selective pressures are stronger on duplicated chemosensory
genes than in orthologous chemosensory genes across corbiculate bees. The
observed higher proportion of branches under positive selection is in contrast to
previous analyses of insect chemosensory genes and might be partly explained by
the former use of less powerful algorithms, reliance on a priori expectations,
incomplete testing, or differences in taxon sampling (64). It is thus unclear whether
the disparity between studies reflects technical differences or differences in
chemosensory gene evolution between different groups of insects. Regardless, our
findings suggest a non-negligible impact of positive selective pressures on
chemosensory gene evolution in corbiculate bees.
475
Diversification of Orthologous ORs through Positive Selection Drives Divergence
in Chemosensory Systems. Our selection analysis indicates that positive selection is
involved in the divergence of orthologous ORs, providing an evolutionary
mechanism for the diversification of chemosensory genes independent of gene
duplication. We found 54 ortholog chemosensory genes and deeper branches that
exhibit signatures of positive selection. Moreover, our analysis using three pairs of
sympatric species revealed the presence of orthologous chemosensory genes
evolving under positive selection irrespective of divergence times (0.15 - 13 my).
This result suggests that positive selection is an important evolutionary force
driving the diversification of orthologous chemosensory genes in corbiculate bees.
Similarly, Leary et al. (2012) (15) showed that positive selection of specific amino
acid substitutions in OR orthologs led to novel receptor functions through changes
in ligand binding affinities in a lineage of moths. Under this model, orthologous OR
genes undergo functional divergence driven by positive selection in the absence of a
duplication event. While OR gene duplications are more common in large
subfamilies in corbiculate bees, this mechanism might be important for the
evolution of interspecific functional divergence in smaller subfamilies. Thus, our
results suggest that positive selection is not only important in driving divergence
between OR duplicates within taxa, but also orthologous ORs between taxa. It is also
possible that a combination of these two mechanisms drives the evolution of
chemosensory genes in other insect lineages.
497
Evidence for ORs with Core Chemosensory Function in Corbiculate Bees. We
detected the existence of multiple highly conserved OR lineages that appear to be
subject to strong purifying selection. This observation supports the existence of a
core set of OR genes with conserved functions across all corbiculate bees and
perhaps other more distantly related bee lineages. We detected 10 OR subfamilies
and 14 orthologous groups within larger subfamilies that solely consisted of 1:1
orthologs. While most OR subfamilies and orthologous groups were characterized
by taxon-specific duplications and divergence driven by positive selective pressures,

our results suggest that these OR lineages are highly conserved throughout at least
80 million years of evolution. This observation implies that corbiculate bee genomes
contain a core set of olfactory genes with conserved chemosensory functions. In fact,
high conservation of OR genes is typical for some species with highly similar natural
history (12, 47, 65). Detailed functional analysis of these core OR genes may
elucidate the role of conserved olfactory functions in corbiculate bees of ecological
importance.
513
514   **Conclusion**
515
Odorant receptors (ORs) have been hypothesized to play an important role in
chemosensory adaptation, for example by enabling the evolution of novel
pheromone detection systems associated with social communication or mate
attraction (1, 66). However, little is known about the molecular mechanisms driving
the evolution of insect OR gene families. Our analysis of the OR gene family in an
important lineage of insect pollinators, the corbiculate bees, elucidated the role of
positive selective pressures and genomic organization on the evolution of the OR
gene family.
524
We show that OR gene family size is highly dynamic between corbiculate bee
species. In contrast, the underlying genomic organization of OR genes is highly
conserved among lineages over at least 80 million years. Consistent with these
findings, ORs are arranged in tandem arrays located in close proximity along the
genome. Moreover, the genomic location correlates with the phylogenetic
relationships among genes that fall in 25 well-supported OR subfamilies. We find
that a few large subfamilies contain a considerable amount of lineage-specific gene
duplications, explaining most of the variation in gene family size. Further, we show
that ORs evolve under strong positive selective pressures, with a higher impact on
duplicated genes. Our study suggests that locally acting molecular mechanisms are
the driving force of OR gene family evolution in corbiculate bees. Moreover, we
provide empirical support for the longstanding hypothesis that positive selection is
an important mechanism in the evolution of duplicated OR genes in insects. These
results elucidate the OR gene family evolution in corbiculate bees and extend the
current understanding of OR gene family evolution to other insect groups.
540
541   **Methods**
542
Sampling and Sequencing
Sampling and sequencing procedures followed Brand et al. 2015. Briefly, males of
the four orchid bee species *Euglossa flammea*, *Eg. imperialis*, *Eulaema bombiformis*
and *El. meriana* were sampled at the La Gamba Field Station in Costa Rica using
different chemical baits (67). Antennal transcriptomes for each species were based
on pools of the antennae of 20 individuals for the *Euglossa* species and 5 individuals
for the *Eulaema* species. Bees were chilled on ice and the antennae of each torpid
male were dissected using sterile forceps and transported either on liquid nitrogen

or RNA-later (Thermo Fisher Scientific, Waltham, MA) and kept frozen until RNA-
extraction.
553
Total-RNA was extracted and quantified as described (35). Barcoded cDNA libraries
were prepared using the NEBnext Ultra RNA Library Prep kit for Illumina (New
England Biolabs, Ipswich, MA) following the manufacturers protocol and sequenced
in two pools of two species on one HiSeq 2500 (*Euglossa*) and one HiSeq 4000
(*Eulaema*) lane for 2x100bp cycles. (Raw sequence reads are available at the NCBI
Sequence Read Archive XXX).
560
De novo *transcriptome assembly and transcript recovery.*
Identical raw reads were merged using fastuniq v1.1 (68) and subsequently
trimmed by trim_galore v0.3.7 (Babraham Bioinformatics) if sequencing primers or
low-quality bases (Phred-score <20) were detected. The pre-processed reads of
each species were used for *de novo* transcriptome assembly following the meta-
assembly approach developed in Brand et al. 2015 using the Trinity (69, 70) and
SOAPdenovo-trans v2 (71) assemblers. Briefly, each of the two assemblers was run
with multiple parameter sets controlling assembly stringency resulting in 25
assemblies per species. Trinity settings were set as described (35), whereas
SOAPdenovo-trans parameters –e and –d were set to 0, 2, 3, or 5 and 1, 3, 5, or 7,
respectively and run in all possible combinations. While Trinity was run with the
default single-k-mer mode, SOAPdenovo-trans was run in multi-k-mer mode with
nine different k-values (25, 31, 37, 43, 49, 59, 69, 79 and 89).
574
Annotation of chemosensory genes.
Chemosensory genes of five gene families (ORs, IRs, GRs, OBPs, and CSPs) were
annotated in the newly sequenced antennal transcriptomes of the orchid bee
species *Eg. flammea*, *Eg. imperialis*, *El. bombiformis*, and *El. meriana*, as well as the
publicly available genomes of the orchid bee species *Eg. dilemma* (40) and *Eufriesea*
mexicana (39), and the stingless bee *Melipona quadrifasciata* (39). Additionally, the
CSP gene family was annotated for the bumble bee *Bombus terrestris* (38). All four
de novo transcriptomes were annotated using a homology-based iterative tBLASTn
approach as previously described (35). In order to annotate the genomes, tBLASTn
was used to detect scaffolds harboring chemosensory genes with an e-value cutoff of
10e-6 based on gene family specific query libraries comprised of published bee,
wasp, ant and dipteran OR, IR, GR, CSP and OBP sequences (21, 35, 38, 41, 51, 72,
73). Subsequently, the queries were used to predict exon-intron boundaries with
exonerate (74) in a scaffold specific manner. All annotations were curated manually
and corrected if needed. All annotations can be found in Dataset S1.
590
Phylogenetic reconstruction of chemosensory gene families.
For each chemosensory gene family we estimated a maximum likelihood (ML) gene-
tree using RaxML (75) to infer the genealogical histories of the candidate
corbiculate gene family members. On that account, gene family specific protein
sequence alignments were constructed using MAFFT (76, 77) applying the L-INS-I
algorithm with the --maxiterate option set to 1000 (78). Based on the resulting

alignments ML trees were inferred under a JTT + G substitution model for each
family with 20 independent ML searches and 100 bootstrap replicates.
599
Analysis of the genomic organization of chemosensory gene families.
Based on genomic annotations of *Eg. dilemma*, *Ef. mexicana*, *M. quadrifasciata* (this
study), *A. mellifera* (21), and *B. terrestris* (38), we analyzed the evolutionary history
of genomic structure of chemosensory gene families within the corbiculate bees.
Therefore, we applied a reciprocal blast approach that compares the genomic
location of all chemosensory gene family tandem arrays and singletons to the
genomes of all other species. We required both 200kb flanking regions to be the
reciprocal best blast hit (79) in order to call homology of the genomic regions. In
case annotations were located less than 200kb away from the end of a scaffold, we
used the maximum flanking region length possible, which never corresponded to
less than 100kb. The location of ORs in the resulting genomic regions in
combination with the gene family phylogeny was used to classify all ORs into
subfamilies. Subfamilies were named after the linkage group they were located on in
the honey bee genome. Letters were used to allow for the unique indication of
subfamilies located on the same linkage group (*e.g.* G12B is the second subfamily
found to be located on linkage group 12 of the honey bee). In case a gene family was
located on a honey bee scaffold that could not be assigned to a linkage group, it was
named 'Un', in short for 'unassigned'.
618
In a second approach, we used the OR gene family phylogeny to infer the
evolutionary histories of micro-genomic synteny within tandem arrays. Orthologs
and paralogs were directly inferred from the ML phylogenies and subsequently
mapped to respective genomic coordinates. Since more distantly related species
share less orthologous than paralogous gene copies (*e.g.* (21, 52)), we focused on
the two orchid bees *Eg. dilemma* and *Ef. mexicana* for this analysis.
625
Selection analysis
Based on the phylogenetic reconstructions, we defined monophyletic orthologous
groups for each gene family consisting of four or more ortholog copies including in-
paralogs that were used for all subsequent selection analyses. In order to identify
genes under positive selection, we applied the adaptive branch-site relative effects-
likelihood (aBSREL) algorithm as implemented in HYPHY (80) to each subgroup to
identify branches with dN/dS ratios significantly larger than 1, indicating positive
selection along these branches (64). P-values for each aBSREL run were corrected
for multiple testing using a false-discovery-rate (FDR) of 5%. In order to analyze
patterns of recent gene family divergence in the orchid bees, we conducted pairwise
dN/dS analyses for the *Eulaema* species pair and the two *Euglossa* species pairs.
Therefore, we extracted orthologous copies from the gene family phylogenies and
performed pairwise nucleotide sequence alignments for each orthologous pair.
Each alignment was subsequently used for ML estimation of the pairwise dN/dS
ratio (model M1) in codeml (81, 82). dN/dS ratios for each pairwise comparison
were tested for significant deviations from the neutral expectation (dN/dS=1; model
M0) using likelihood-ratio tests.

Acknowledgements
645
We thank Michelle Stitzer, Judy Wexler, Mascha Koenen and the Ramírez lab for
valuable comments on the manuscript. We thank Hugh Robertson for sharing
chemosensory gene sequences of the honey bee and bumble bee. P.B. was supported
by a fellowship of the German academic exchange service (Deutscher Akademischer
Austauschdienst, DAAD) for parts of the project. SRR received support from the
David & Lucile Packard Foundation and the National Science Foundation (DEB-
1457753). This work used the Vincent J. Coates Genomics Sequencing Laboratory at
UC Berkeley, supported by NIH S10 Instrumentation Grants S10RR029668,
S10RR027303, and OD018174.

655
656 **References**
657

1.   Hansson BS, Stensmyr MC (2011) Evolution of insect olfaction. *Neuron*
72(5):698–711.

2.   Vosshall LB, Wong AM, Axel R (2000) An olfactory sensory map in the fly brain.
Cell 102(2):147–159.

3.   Elmore T, Smith DP (2001) Putative Drosophila odor receptor OR43b localizes to
dendrites of olfactory neurons. *Insect Biochem Mol Biol* 31(8):791–798.

4.   Dobritsa AA, van der Goes van Naters W, Warr CG, Steinbrecht RA, Carlson JR
(2003) Integrating the Molecular and Cellular Basis of Odor Coding in the
Drosophila Antenna. *Neuron* 37(5):827–841.

5.   Hallem EA, Ho MG, Carlson JR (2004) The molecular basis of odor coding in the
Drosophila antenna. *Cell* 117(7):965–979.

6.   Hallem EA, Carlson JR (2006) Coding of odors by a receptor repertoire. *Cell*
125(1):143–160.

7.   Wang G, Carey AF, Carlson JR, Zwiebel LJ (2010) Molecular basis of odor
coding in the malaria vector mosquito Anopheles gambiae. *Proc Natl Acad Sci*
USA 107(9):4418–4423.

8.   Kirkness EF, et al. (2010) Genome sequences of the human body louse and its
primary endosymbiont provide insights into the permanent parasitic lifestyle. *Proc*
Natl Acad Sci USA 107(27):12168–12173.

9.   Smith CD, et al. (2011) Draft genome of the globally widespread and invasive
Argentine ant (Linepithema humile). *Proc Natl Acad Sci USA* 108(14):5673–5678.

10.   Smith CR, et al. (2011) Draft genome of the red harvester ant Pogonomyrmex
barbatus. *Proc Natl Acad Sci USA* 108(14):5667–5672.

11. McBride CS (2007) Rapid evolution of smell and taste receptor genes during host
specialization in Drosophila sechellia. *Proc Natl Acad Sci USA* 104(12):4996–
5001.

12. McBride CS, Arguello JR (2007) Five Drosophila genomes reveal nonneutral
evolution and the signature of host specialization in the chemoreceptor
superfamily. *Genetics* 177(3):1395–1416.

13. Goldman-Huertas B, et al. (2015) Evolution of herbivory in Drosophilidae linked
to loss of behaviors, antennal responses, odorant receptors, and ancestral diet. *Proc
Natl Acad Sci USA* 112(10):3026–3031.

14. Gould F, et al. (2010) Sexual isolation of male moths explained by a single
pheromone response QTL containing four receptor genes. *Proc Natl Acad Sci USA*
107(19):8660–8665.

15. Leary GP, et al. (2012) Single mutation to a sex pheromone receptor provides
adaptive specificity between closely related moth species. *Proc Natl Acad Sci USA*
109(35):14081–14086.

16. Pellegrino M, Steinbach N, Stensmyr MC, Hansson BS, Vosshall LB (2011) A
natural polymorphism alters odour and DEET sensitivity in an insect odorant
receptor. *Nature* 478(7370):511–514.

17. Nei M, Niimura Y, Nozawa M (2008) The evolution of animal chemosensory
receptor gene repertoires: roles of chance and necessity. *Nat Rev Genet* 9(12):951–
963.

18. Sánchez-Gracia A, Vieira FG, Rozas J (2009) Molecular evolution of the major
chemosensory gene families in insects. *Heredity* 103(3):208–216.

19. Nei M, Rooney AP (2005) Concerted and birth-and-death evolution of multigene
families. *Annu Rev Genet* 39(1):121–152.

20. Innan H, Kondrashov F (2010) The evolution of gene duplications: classifying and
distinguishing between models. *Nat Rev Genet* 11(2):97–108.

21. Robertson HM, Wanner KW (2006) The chemoreceptor superfamily in the honey
bee, Apis mellifera: expansion of the odorant, but not gustatory, receptor family.
Genome Res 16(11):1395–1403.

22. Guo S, Kim J (2007) Molecular evolution of Drosophila odorant receptor genes.
Mol Biol Evol 24(5):1198–1207.

23. Conceição IC, Aguadé M (2008) High Incidence of Interchromosomal
Transpositions in the Evolutionary History of a Subset of Or Genes in Drosophila.
J Mol Evol 66(4):325–332.

24.   Kratz E, Dugas JC, Ngai J (2002) Odorant receptor gene regulation: implications
from genomic organization. *Trends in Genetics* 18(1):29–34.

25.   Klein A-M, et al. (2007) Importance of pollinators in changing landscapes for
world crops. *Proceedings of the Royal Society B: Biological Sciences*
274(1608):303–313.

26.   Slaa EJ, Chaves LAS, Malagodi-Braga KS, Hofstede FE (2006) Stingless bees in
applied pollination: practice and perspectives. *Apidologie* 37(2):293–315.

27.   Ramírez S, Dressler RL, Ospina M (2002) Abejas euglosinas (Hymenoptera:
Apidae) de la Región Neotropical: Listado de especies con notas sobre su biología.
3:7.

28.   Grüter C, Keller L (2016) Inter-caste communication in social insects. *Current
Opinion in Neurobiology* 38:6–11.

29.   Kocher SD, Grozinger CM (2011) Cooperation, Conflict, and the Evolution of
Queen Pheromones. *J Chem Ecol* 37(11):1263–1275.

30.   Leonhardt SD, Menzel F, Nehring V, Schmitt T (2016) Ecology and Evolution of
Communication in Social Insects. *Cell* 164(6):1277–1287.

31.   Leonhardt SD (2017) Chemical Ecology of Stingless Bees. *J Chem Ecol* 2(2):1–
18.

32.   Vogel S (1966) Parfümsammelnde Bienen als Bestäuber von Orchidaceen und
Gloxinia. *Österr bot Z* 113(3-4):302–361.

33.   Eltz T, Whitten WM, Roubik DW, Linsenmair KE (1999) Fragrance Collection,
Storage, and Accumulation by Individual Male Orchid Bees. *J Chem Ecol*
25(1):157–176.

34.   Roubik DW, Hanson PE (2004) *Orchid bees of tropical America: biology and field
guide* (Instituto Nacional de Biodiversidad (INBio), Santo Domingo De Heredia).

35.   Brand P, et al. (2015) Rapid evolution of chemosensory receptor genes in a pair of
sibling species of orchid bees (Apidae: Euglossini). *BMC Evol Biol* 15(1):176.

36.   McBride CS, et al. (2014) Evolution of mosquito preference for humans linked to
an odorant receptor. *Nature* 515(7526):222–227.

37.   Weinstock GM, et al. (2006) Insights into social insects from the genome of the
honeybee Apis mellifera. *Nature* 443(7114):931–949.

38.   Sadd BM, et al. (2015) The genomes of two key bumblebee species with primitive
eusocial organization. *Genome Biology* 16(1):76.

39. Kapheim KM, et al. (2015) Genomic signatures of evolutionary transitions from
solitary to group living. *Science*:aaa4788.

40. Brand P, et al. (2017) The Nuclear And Mitochondrial Genomes Of The
Facultatively Eusocial Orchid Bee Euglossa dilemma. *bioRxiv*:123687.

41. Zhou X, et al. (2012) Phylogenetic and transcriptomic analysis of chemosensory
receptors in a pair of divergent ant species reveals sex-specific signatures of odor
coding. *PLoS Genet* 8(8):e1002930.

42. Engsontia P, Sangket U, Robertson HM, Satasook C (2015) Diversification of the
ant odorant receptor gene family and positive selection on candidate cuticular
hydrocarbon receptors. *BMC Research Notes* 8(1):380.

43. Ross L, Blackmon H, Lorite P, Gokhman VE, Hardy NB (2015) Recombination,
chromosome number and eusociality in the Hymenoptera. *Journal of Evolutionary*
Biology 28(1):105–116.

44. Stolle E, et al. (2011) A second generation genetic map of the bumblebee Bombus
terrestris (Linnaeus, 1758) reveals slow genome and chromosome evolution in the
Apidae. *BMC Genomics* 12(1):48.

45. Rueppell O, et al. (2016) A New Metazoan Recombination Rate Record and
Consistently High Recombination Rates in the Honey Bee Genus Apis
Accompanied by Frequent Inversions but not Translocations. *Genome Biol*
Evol:evw269.

46. Wilfert L, Gadau J, Schmid-Hempel P (2007) Variation in genomic recombination
rates among animal taxa and the case of social insects. *Heredity* 98(4):189–197.

47. Nozawa M, Nei M (2007) Evolutionary dynamics of olfactory receptor genes in
Drosophila species. *Proc Natl Acad Sci USA* 104(17):7122–7127.

48. Obbard DJ, et al. (2012) Estimating divergence dates and substitution rates in the
Drosophila phylogeny. *Mol Biol Evol* 29(11):3459–3473.

49. Niimura Y, Nei M (2005) Comparative evolutionary analysis of olfactory receptor
gene clusters between humans and mice. *Gene* 346:13–21.

50. Bosco G, Campbell P, Leiva-Neto JT, Markow TA (2007) Analysis of drosophila
species genome size and satellite DNA content reveals significant differences
among strains as well as between species. *Genetics* 177(3):1277–1290.

51. Croset V, et al. (2010) Ancient protostome origin of chemosensory ionotropic
glutamate receptors and the evolution of insect taste and olfaction. *PLoS Genet*
6(8):e1001064.

52. Zhou X, et al. (2015) Chemoreceptor evolution in Hymenoptera and its

implications for the evolution of eusociality. *Genome Biol Evol*:evv149.

53.   Hastings PJ, Lupski JR, Rosenberg SM, Ira G (2009) Mechanisms of change in
gene copy number. *Nat Rev Genet* 10(8):551–564.

54.   Perry GH, et al. (2008) Copy number variation and evolution in humans and
chimpanzees. *Genome Res* 18(11):1698–1710.

55.   Perry GH, et al. (2006) Hotspots for copy number variation in chimpanzees and
humans. *Proc Natl Acad Sci USA* 103(21):8006–8011.

56.   Cardoso-Moreira M, Emerson JJ, Clark AG, Long M (2011) Drosophila
Duplication Hotspots Are Associated with Late-Replicating Regions of the
Genome. *PLoS Genet* 7(11):e1002340.

57.   LeBoeuf AC, Benton R, Keller L (2013) The molecular basis of social behavior:
models, methods and advances. *Current Opinion in Neurobiology* 23(1):3–10.

58.   McKenzie SK, Fetter-Pruneda I, Ruta V, Kronauer DJC (2016) Transcriptomics
and neuroanatomy of the clonal raider ant implicate an expanded clade of odorant
receptors in chemical communication. *Proc Natl Acad Sci USA*:201610800.

59.   Woodard SH, et al. (2011) Genes involved in convergent evolution of eusociality
in bees. *Proc Natl Acad Sci USA* 108(18):7472–7477.

60.   Michener CD (2007) *The Bees of the World. 2nd* (Ed. Johns Hopkins).

61.   Tunstall NE, Sirey T, Newcomb RD, Warr CG (2007) Selective pressures on
Drosophila chemosensory receptor genes. *J Mol Evol* 64(6):628–636.

62.   Gardiner A, Barker D, Butlin RK, Jordan WC, Ritchie MG (2008) Drosophila
chemoreceptor gene evolution: selection, specialization and genome size.
Molecular Ecology 17(7):1648–1657.

63.   Engsontia P, Sangket U, Chotigeat W, Satasook C (2014) Molecular evolution of
the odorant and gustatory receptor genes in lepidopteran insects: implications for
their adaptation and speciation. *J Mol Evol* 79(1-2):21–39.

64.   Smith MD, et al. (2015) Less is more: an adaptive branch-site random effects
model for efficient detection of episodic diversifying selection. *Mol Biol Evol*
32(5):1342–1353.

65.   Neafsey DE, et al. (2015) Highly evolvable malaria vectors: The genomes of 16
Anopheles mosquitoes. *Science* 347(6217):1258522–1258522.

66.   Smadja C, Butlin RK (2009) On the scent of speciation: the chemosensory system
and its role in premating isolation. *Heredity* 102(1):77–97.

67.   Pokorny T, et al. (2013) Acquisition of species-specific perfume blends: influence
of habitat-dependent compound availability on odour choices of male orchid bees
(Euglossa spp.). *Oecologia* 172(2):417–425.

68.   Xu H, et al. (2012) FastUniq: a fast de novo duplicates removal tool for paired
short reads. *PLoS ONE* 7(12):e52249.

69.   Grabherr MG, et al. (2011) Full-length transcriptome assembly from RNA-Seq
data without a reference genome. *Nat Biotechnol* 29(7):644–652.

70.   Haas BJ, et al. (2013) De novo transcript sequence reconstruction from RNA-seq
using the Trinity platform for reference generation and analysis. *Nat Protoc*
8(8):1494–1512.

71.   Xie Y, et al. (2014) SOAPdenovo-Trans: de novo transcriptome assembly with
short RNA-Seq reads. *Bioinformatics* 30(12):1660–1666.

72.   Forêt S, Maleszka R (2006) Function and evolution of a gene family encoding
odorant binding-like proteins in a social insect, the honey bee (Apis mellifera).
Genome Res 16(11):1404–1413.

73.   Vieira FG, Rozas J (2011) Comparative genomics of the odorant-binding and
chemosensory protein gene families across the Arthropoda: origin and
evolutionary history of the chemosensory system. *Genome Biol Evol* 3(0):476–
490.

74.   Slater GSC, Birney E (2005) Automated generation of heuristics for biological
sequence comparison. *BMC Bioinformatics* 6(1):31.

75.   Stamatakis A, Ludwig T, Meier H (2005) RAxML-III: a fast program for
maximum likelihood-based inference of large phylogenetic trees. *Bioinformatics*
21(4):456–463.

76.   Katoh K, Misawa K, Kuma K-I, Miyata T (2002) MAFFT: a novel method for
rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids*
Res 30(14):3059–3066.

77.   Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software
version 7: improvements in performance and usability. *Mol Biol Evol* 30(4):772–
780.

78.   Katoh K, Kuma K-I, Toh H, Miyata T (2005) MAFFT version 5: improvement in
accuracy of multiple sequence alignment. *Nucleic Acids Res* 33(2):511–518.

79.   Rivera MC, Jain R, Moore JE, Lake JA (1998) Genomic evidence for two
functionally distinct gene classes. *Proc Natl Acad Sci USA* 95(11):6239–6244.

80.   Pond SLK, Frost SDW, Muse SV (2005) HyPhy: hypothesis testing using

phylogenies. *Bioinformatics* 21(5):676–679.

81.   Bielawski JP, Yang Z (2004) A Maximum Likelihood Method for Detecting
854   Functional Divergence at Individual Codon Sites, with Application to Gene Family
855   Evolution. *J Mol Evol* 59(1):121–132.

856   82.   Yang Z (2007) PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol*
857   *Evol* 24(8):1586–1591.

859   **Figures**
860
Fig. 1 Gene family size dynamics in corbiculate bees. Numbers of previously (*A.*
mellifera, B. terrestris, Eg. viridissima) and newly annotated (all others) genes for
each gene family are indicated. Annotations derived from whole genomes and
antennal transcriptomes are labeled. Phylogenetic relationships and divergence
times of the ten corbiculate bee species following Ramirez et al. 2010 and Romiguier
et al. 2015 are shown. Annotated nodes indicate MRCA of bumble bees and stingless
bees (A), honey bees + A (B) and all corbiculate bees (C).

868
Fig. 2 Odorant receptor subfamily dynamics in corbiculate bees. Relationships
between subfamilies (**A**) are based on all annotated ORs of 10 corbiculate bee
species. Observed differences of within subfamily dynamics are shown with
subfamily G13A and subfamily G04A as examples (**B, C**). (**A**) OR subfamilies have
high bootstrap support while deeper nodes are of low support. All genes within a
subfamily are located in a single tandem array with highly conserved genomic
landscapes throughout corbiculate bees. The few exceptions are indicated with *. (**B,**
C) ORs in subfamily G13A are mainly simple 1:1 single copy orthologs found in all or
a subset of the analyzed species (marked '1:1') in contrast to subfamily G04A, where
lineage specific duplications and larger expansions in *A. mellifera*, *B. terrestris*, *Eg.*
dilemma and *Ef. mexicana* dominate (marked 'E'). Branches evolving under positive
selective pressures are indicated by a red star. Both, duplicate branches as well as
branches in 1:1 ortholog groups were under positive selection. The branch indicated
with a * in (**B**) indicates the sistergroup to all other ORs in subfamily G13A which is
found at a genomic location different from the rest of the subfamily in all corbiculate
bee species representing an ancient translocation event (see text for details).

885
Fig. 3 Evolution of OR subfamily size. (**A**) OR subfamily size variation among
corbiculates correlates with mean subfamily size. The pattern holds true for all
corbiculates (white dots, black dotted line) and the two orchid bee species alone
(black dots, grey dotted line). **(B)** OR subfamily repertoire numbers among
corbiculate bees. Numbers of ORs in all 25 subfamilies for the five species with
whole-genome data are indicated. Pseudogenes are excluded. *Eg. dilemma* and *Ef.*
mexicana represent the two orchid bee species. OR subfamily sizes are highly
variable between species, while increased subfamily sizes are shared by multiple

lineages or occur in single lineages only. 9Exon contains the combined repertoire of
subfamilies G04A, G05B, G11A, G11B, and G12A.
896
897
898 **Fig. 1**

	ORs	GRs	IRs	OBPs	CSPs	
A. mellifera	177	14	10	21	6	Genomic
B. terrestris	166	25	11	16	5	
M. quadrifasciata	196	16	10	14	6	
Ef. mexicana	142	16	9	16	7	
Eg. dilemma	183	13	9	15	7	
Eg. viridissima	85	4	5	11	6	Transcriptomic
Eg. flammea	115	6	9	13	6	
Eg. imperialis	108	11	9	14	7	
El. bombiformis	115	7	9	15	6	
El. meriana	110	10	10	18	6	

Honey bees Bumble bees
Stingless bees Orchid bees

899
900
901

 Fig. 2

A)

B) Subfamily G13A

C) Subfamily G04A

905   **Fig. 3**

906
907
908
909   **Tables**

910
911   **Tab. 1** Numbers of branches under positive selection for each chemosensory gene
912   family among the four corbiculate bee tribes.

913
914   Tab. 1

		Orchid bees	Honey bees	Bumble bees	Stingless bees	Total
ORs	Tested	1234	216	194	237	1881
	P ≤ 0.05	42 (3.4%)	5 (2.3%)	3 (1.6%)	14 (5.9%)	64 (3.4%)
GRs	Tested	116	11	32	15	174
	P ≤ 0.05	17 (14.7%)	2 (18.2%)	5 (15.6%)	4 (26.7%)	28 (16.1%)
IRs	Tested	101	9	9	9	128
	P ≤ 0.05	5 (5.0%)	0 (0%)	0 (0%)	0 (0%)	5 (3.9%)
OBPs	Tested	178	29	18	12	237
	P ≤ 0.05	3 (1.7%)	4 (13.8%)	0 (0%)	0 (0%)	7 (3.0%)
CSPs	Tested	76	6	5	6	93
	P ≤ 0.05	1 (1.3%)	0 (0%)	0 (0%)	0 (0%)	1 (1.1%)

915   All branches tested as well as the numbers and percentages of branches under positive selection after 5% FDR
916   correction are represented.

24